



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

A Multi-scale Spin-Glass Mean-Field Model

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Contucci, P., Mingione, E. (2019). A Multi-scale Spin-Glass Mean-Field Model. COMMUNICATIONS IN MATHEMATICAL PHYSICS, 368(3), 1323-1344 [10.1007/s00220-019-03308-8].

Availability:

This version is available at: <https://hdl.handle.net/11585/732135> since: 2020-02-29

Published:

DOI: <http://doi.org/10.1007/s00220-019-03308-8>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Contucci, P., Mingione, E. A Multi-scale Spin-Glass Mean-Field Model. *Commun. Math. Phys.* 368, 1323–1344 (2019).

The final published version is available online at : <https://doi.org/10.1007/s00220-019-03308-8>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

A Multi-Scale Spin-Glass Mean-Field Model

Pierluigi Contucci, Emanuele Mingione

Dipartimento di Matematica

Università di Bologna, 40127 Bologna, Italy

Abstract

In this paper a multi-scale version of the Sherrington and Kirkpatrick model is introduced and studied. The pressure per particle in the thermodynamical limit is proved to obey a variational principle of Parisi type. The result is achieved by means of lower and upper bounds. The lower bound is obtained with a Ruelle cascade using the interpolation technique, while the upper bound exploits factorisation properties of the equilibrium measure and the synchronisation technique.

Keywords: Spin glasses, Sherrington-Kirkpatrick model, multi-scale decomposition

1 The Multiscale SK model

The equilibrium statistical mechanics of a general disordered system can be described in between two prescriptions known, in the literature, as quenched and annealed. The spin-glass phase, for instance, is described by the quenched measure where the random coupling disorder is kept fixed while the spins are thermalised according to the Boltzmann distribution. This perspective is considered physically relevant because the relaxation time of the disorder interaction variables is much slower than the one for the spin variables. Conversely in the annealed prescription the disorder variables thermalise together with the field ones. In a paper by Talagrand [35] on mean field spin glasses it was shown how to define a generalised equilibrium measure depending on a real positive number ζ with thermodynamic pressure

$$P = \frac{1}{\zeta} \log \mathbb{E} Z_J^\zeta, \tag{1}$$

where Z_J is the partition function, a random variable depending on the disorder J obtained integrating on the spins. The origins of this description are to be found on the replica approach to spin glasses [24] where ζ is an integer. In [6] ζ was treated as a *scale* parameter in the unit interval to interpolate a general disordered system from the quenched case, obtained when $\zeta \rightarrow 0$, to the annealed one reached at $\zeta \rightarrow 1$.

In this paper we generalise the idea of (1) and consider a multi-scale equilibrium measure obtained by successive independent integration on suitably defined Gaussian couplings. The idea to study a system at different energy scales is common in mathematical-physics at least since the early days of the Euclidian approach to renormalisation group in quantum field theory (see [17, 34]). Recalling the basic concepts, a single scale model is defined as in (1). For two scales ζ_0 and ζ_1 the model is defined in terms of an interaction $J = (J_0, J_1)$ with independent components:

$$e^{\zeta_1 P^{(0)}} = \mathbb{E}_1 Z_J^{\zeta_1}, \quad (2)$$

and

$$e^{\zeta_0 P} = \mathbb{E}_0 e^{\zeta_0 P^{(0)}}. \quad (3)$$

For r scales $\zeta_0 < \zeta_1 < \dots < \zeta_{r-1} < \zeta_r = 1$ the recursion relations are

$$e^{\zeta_l P^{(l-1)}} = \mathbb{E}_l e^{\zeta_l P^{(l)}}, \quad (4)$$

where $0 \leq l \leq r$, $\mathbb{E}_r e^{P^{(r)}} = Z_J$ and $P = P^{(-1)}$.

The use of a multi-scale decomposition structure in the spin-glass problem made its first appearance with the celebrated work by Guerra on the Sherrington-Kirkpatrick model [22] where the covariance of a one-body exactly solvable system is split in many layers. The same method was later used by Talagrand in his complete proof, the upper bound, of the Parisi formula for the free energy density of the model [36]. The idea to use the recursive structure introduced above appeared also in the theoretical physics literature. In [25] the author uses it to investigate the properties of metastable states in a glassy system. In [14, 15] the authors introduce a multi-bath equilibrium showing that it can be used to describe the correlations and response functions for a class of dynamical systems in the limit of small entropy production.

What we propose here is a generalised mean-field model where a multi-scale structure is part of the model itself and involves the interacting covariance.

A fundamental tool throughout this work, that we will use to study the multi-scale equilibrium measure defined by (4), are the Ruelle Probability Cascades (RPC) [16, 27] whose use is consolidated in the spin-glass literature [7, 8, 10, 28]. A short appendix on RPC is provided at the end to make this work self-contained.

The main definitions follow.

Given $N \geq 1$ let us consider a system of N spins $\sigma = (\sigma_i)_{i \leq N} \in \Sigma_N = \{-1, 1\}^N$. Fix an integer $r \geq 1$ and denote by $\alpha \in \mathbb{N}^r$ an additional degree of freedom. A configuration of the system is

$$\sigma = (\sigma, \alpha) \in \Sigma_N \times \mathbb{N}^r \equiv \Sigma_{N,r} \quad (5)$$

Consider a sequence $\zeta = (\zeta_l)_{l \leq r}$ such that

$$0 = \zeta_{-1} < \zeta_0 < \zeta_1 < \dots < \zeta_{r-1} < \zeta_r = 1 \quad (6)$$

and let $(\nu_\alpha)_{\alpha \in \mathbb{N}^r}$ be the random weights of Ruelle Probability Cascade associated to the sequence ζ (see Appendix 5). For $\alpha, \beta \in \mathbb{N}^r$ we denote

$$\alpha \wedge \beta = \min \{0 \leq l \leq r \mid \alpha_1 = \beta_1, \dots, \alpha_l = \beta_l, \alpha_{l+1} \neq \beta_{l+1}\} \quad (7)$$

where $\alpha \wedge \beta = r$ if $\alpha = \beta$. It's useful to think \mathbb{N}^r as the set of leaves of an infinite tree $\mathcal{A} = \mathbb{N}^0 \cup \mathbb{N} \cup \mathbb{N}^2 \dots \cup \mathbb{N}^r$ of depth r and root $\mathbb{N}^0 = \{\emptyset\}$. Then $\alpha \wedge \beta$ denotes the level of their common ancestor, see (119).

Fix a sequence $\gamma = (\gamma_l)_{l \leq r}$ such that

$$0 = \gamma_0 < \gamma_1 < \dots < \gamma_r < \infty \quad (8)$$

and let $(g(\alpha))_{\alpha \in \mathbb{N}^r}$ be a family of centered gaussian random variables with covariance

$$\mathbb{E} g(\alpha^1) g(\alpha^2) = (\gamma_{\alpha^1 \wedge \alpha^2})^2 \quad (9)$$

Consider a gaussian process H_N on $\Sigma_{N,r}$ defined by

$$H_N(\sigma) = \frac{1}{\sqrt{N}} \sum_{i,j=1}^N g_{ij}(\alpha) \sigma_i \sigma_j \quad (10)$$

where $\sigma = (\sigma, \alpha) \in \Sigma_N \times \mathbb{N}^r$ and $(g_{ij}(\alpha))_{i,j=1,\dots,N}$ is a family of i.i.d. copies of $g(\alpha)$.

Given two configurations $\sigma^1 = (\sigma^1, \alpha^1)$, $\sigma^2 = (\sigma^2, \alpha^2) \in \Sigma_{N,r}$ the covariance of the process H_N is

$$\mathbb{E} H_N(\sigma^1) H_N(\sigma^2) = N (c_{N,\gamma}(\sigma^1, \sigma^2))^2 \quad (11)$$

where

$$c_{N,\gamma}(\sigma^1, \sigma^2) = \gamma_{\alpha^1 \wedge \alpha^2} q_N(\sigma^1, \sigma^2) \quad (12)$$

and

$$q_N(\sigma^1, \sigma^2) = \frac{1}{N} \sum_{i=1}^N \sigma_i^1 \sigma_i^2 \quad (13)$$

is the usual *overlap* between two configurations $\sigma^1, \sigma^2 \in \Sigma_N$. Notice that $q_N(\sigma^1, \sigma^2) \in [-1, 1]$ and $\alpha \wedge \alpha = r$ imply that $c_{N,\gamma}(\sigma^1, \sigma^2) \in [-\gamma_r, \gamma_r]$.

We denote by $\beta = (\zeta, \gamma)$ the couple of sequences in (6) and (8). Given β we by $p_N(\beta)$ the *quenched pressure density* of the Multiscale SK model, defined as

$$p_N(\beta) = \frac{1}{N} \mathbb{E} \log Z_N(\beta) \quad (14)$$

where

$$Z_N(\beta) = \sum_{\sigma \in \Sigma_{N,r}} \nu_{\alpha} e^{H_N(\sigma)} \quad (15)$$

We notice that $p_N(\beta)$ in (14) can be also defined recursively. Let $H_N(\sigma, l)$ be a gaussian process on $(\sigma, l) \in \Sigma_N \times \{1, \dots, r\}$ with covariance

$$\mathbb{E} H_N(\sigma^1, l) H_N(\sigma^2, l') = N \delta_{l,l'} \left(\sqrt{\gamma_l^2 - \gamma_{l-1}^2} q_N(\sigma^1, \sigma^2) \right)^2 \quad (16)$$

Then, by the property (122) of the RPC, it holds

$$p_N(\beta) = \frac{1}{N} \log Z_{0,N}(\beta) \quad (17)$$

where $Z_{0,N}$ is obtained recursively in the following way. We denote by \mathbb{E}_l denotes the average w.r.t. the randomness in $H_N(\sigma, l+1)$ and starting from

$$Z_{r,N}(\beta) = \sum_{\sigma} \prod_{1 \leq l \leq r} e^{H_N(\sigma, l)} \quad (18)$$

we define

$$Z_{l-1,N}^{\zeta_{l-1}} = \mathbb{E}_{l-1} Z_{l,N}^{\zeta_{l-1}} \quad (19)$$

for any $0 \leq l \leq r-1$.

For $r=1$ and a generic ζ_0 the model was studied and solved by Talagrand in [35]. If $\zeta_0 \rightarrow 0$ we recover the SK model at inverse temperature γ_1 .

2 Main result

The quenched pressure density p_N in (14) is completely determined by the choice of $\beta = (\zeta, \gamma)$. From now on r denotes the integer that defines the sequences ζ and γ in (6) and (8).

Consider an arbitrary integer $k \geq r$ and a sequence $\xi = (\xi_j)_{j \leq k}$ such that

$$0 = \xi_{-1} < \xi_0 < \xi_1 < \dots < \xi_k = 1 \quad (20)$$

Moreover we assume that

$$\zeta \subseteq \xi \quad (21)$$

It's useful to think ξ as a the image of some discrete distribution function. In other words given an arbitrary sequence $c = (c_j)_{j \leq k}$ such that

$$0 < c_0 < c_1 < \dots < c_k < \infty \quad (22)$$

we say that a random variable C taking values on the set c has distribution ξ if

$$\mathbb{P}(C = c_j) = \xi_j - \xi_{j-1} \quad (23)$$

for any $j \leq k$. Any couple of sequences (ξ, c) satisfying (20) and (22) combined with the relation (23) determines an element of $\mathcal{M}[0, c_k]$ where $\mathcal{M}[0, c_k]$ denotes the set of all distribution functions on $[0, c_k]$. However in our case the additional condition (21) implies that we look at a particular subset of $\mathcal{M}[0, c_k]$.

Definition 2.1. We denotes by $\mathcal{M}_\zeta[0, c_k]$ the set of all distribution function F on $[0, c_k]$ such that the sequence ζ is contained in the image of F .

Notice that if F is a discrete distribution on $[0, c_k]$ then it can be identified with a couple (ξ, c) satisfying (20) and (22) and the above definition implies that

$$F \in \mathcal{M}_\zeta[0, c_k] \Leftrightarrow \zeta \subseteq \xi \quad (24)$$

Now given the sequence ξ in (20) satisfying (21) consider the following subset of $\{0, \dots, k\}$

$$K_l = \{j : \zeta_{l-1} < \xi_j \leq \zeta_l, 0 \leq j \leq k\} \quad (25)$$

for any $l \leq r$. Given the sequence γ in (8) we construct a new sequence $\tilde{\gamma} = (\tilde{\gamma}_j)_{j \leq k}$ defining for any $j \leq k$

$$\tilde{\gamma}_j = \gamma_l \text{ if } j \in K_l \quad (26)$$

We also introduce an arbitrary sequence $q = (q_j)_{j \leq k}$ such that

$$0 = q_0 \leq q_1 \leq \dots \leq q_k = 1 \quad (27)$$

Definition 2.2. We denote X_β the set of all $x = (\xi, \tilde{\gamma}, q)$ such that ξ satisfies (20) and (21) while $\tilde{\gamma}$ and q are defined in (26) and (27) respectively.

Given $x = (\xi, \tilde{\gamma}, q) \in X_\beta$, consider the sequence $c = (c_j)_{j \leq k}$ where $c_j = \tilde{\gamma}_j q_j$ for any $j \leq k$. Then, from a physical point of view, the couple (ξ, c) associated to a suitable $x \in X_\beta$, represents the distribution of the overlap c_N in (12) w.r.t. the Gibbs measure in the thermodynamic limit.

Let $(J_j)_{1 \leq j \leq k}$ be a collection of i.i.d. standard gaussian random variables and define

$$Z_k = 2 \cosh \left(\sqrt{2} \sum_{1 \leq j \leq k} J_j \left(\tilde{\gamma}_j^2 q_j - \tilde{\gamma}_{j-1}^2 q_{j-1} \right)^{1/2} \right) \quad (28)$$

and recursively for $0 \leq j \leq k-1$

$$Z_{j-1}^{\xi_{j-1}} = \mathbb{E}_{j-1} Z_j^{\xi_{j-1}} \quad (29)$$

where \mathbb{E}_j denotes the average w.r.t. J_{j+1} .

For any $x \in X_\beta$ we define the Parisi functional for the Multiscale SK model the quantity

$$\mathcal{P}_\beta(x) = \log Z_0 - \frac{1}{2} \sum_{0 \leq j \leq k-1} \xi_j \left((\tilde{\gamma}_{j+1} q_{j+1})^2 - (\tilde{\gamma}_j q_j)^2 \right) \quad (30)$$

Using (122) one can prove that the Parisi functional (30) has another useful representation. Let $(\nu_\alpha)_{\alpha \in \mathbb{N}^k}$ the random weights of the RPC with parameter ξ . Consider two independent gaussian process z, y indexed by $\alpha \in \mathbb{N}^k$ with covariances

$$\mathbb{E} z(\alpha^1) z(\alpha^2) = 2 (\tilde{\gamma}_{\alpha^1 \wedge \alpha^2})^2 q_{\alpha^1 \wedge \alpha^2} \quad (31)$$

$$\mathbb{E} y(\alpha^1) y(\alpha^2) = (\tilde{\gamma}_{\alpha^1 \wedge \alpha^2} q_{\alpha^1 \wedge \alpha^2})^2 \quad (32)$$

Hence it holds

$$\mathcal{P}_\beta(x) = \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^k} \nu_\alpha 2 \cosh z(\alpha) - \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^k} \nu_\alpha 2 \exp y(\alpha) \quad (33)$$

The main result of this work is the following

Theorem 2.1. *The thermodynamic limit of the quenched pressure density of the Multiscale SK model $p_N(\beta)$ in (14) exists and is given by*

$$\lim_{N \rightarrow \infty} p_N(\beta) = \inf_{x \in X_\beta} \mathcal{P}_\beta(x) \quad (34)$$

where $\mathcal{P}_\beta(x)$ is the Parisi-like functional defined in (30) and the set X_β is defined in (2.2).

The existence of the thermodynamic limit of $p_N(\beta)$ can be proved regardless of (34) using a Guerra-Toninelli argument [21]. Indeed the covariance $c_{N,\gamma}$ in (12) depends on N only through the overlap q_N in (13), namely the covariance of an SK model.

It would be interesting to see if the functional \mathcal{P}_β is convex as it has been proved in the case of the SK model [2].

Notice also that in Talagrand's paper [35] where the case $r = 1, \zeta_0 \in (0, 1)$ is considered, the trial RPC starts from $\xi_0 = \zeta_0$. Even if this requirement is not present explicitly in the definition (30) for the trial functional \mathcal{P}_β , it's possible to show that condition (26) implies it.

3 Upper bound, Guerra's interpolation

In this section we give an upper bound for the quenched pressure of the Multiscale SK model p_N defined in (14). In the proof given here we use RPC formalism. The same result can be obtained working with the recursive definition (17) for $p_N(\beta)$ and applying Guerra's methods [19, 22].

Proposition 3.1. *The quenched pressure density of the Multiscale SK model $p_N(\beta)$ satisfies*

$$\limsup_{N \rightarrow \infty} p_N(\beta) \leq \inf_{x \in X_\beta} \mathcal{P}_\beta(x) \quad (35)$$

where the functional $\mathcal{P}_\beta(x)$ and the set X_β are defined in (30) and (2.2) respectively.

Proof. Let $(\nu_\alpha)_{\alpha \in \mathbb{N}^k}$ the random weights of the RPC with parameter $\xi = (\xi_j)_{j \leq k}$ in (20) and consider two independent gaussian process \tilde{g}, z indexed by $\alpha \in \mathbb{N}^k$ with covariances

$$\mathbb{E} \tilde{g}(\alpha^1) \tilde{g}(\alpha^2) = (\tilde{\gamma}_{\alpha^1 \wedge \alpha^2})^2 \quad (36)$$

$$\mathbb{E} z(\alpha^1) z(\alpha^2) = 2(\tilde{\gamma}_{\alpha^1 \wedge \alpha^2})^2 q_{\alpha^1 \wedge \alpha^2} \quad (37)$$

where $q = (q_j)_{j \leq k}$ and $\tilde{\gamma} = (\tilde{\gamma}_j)_{j \leq k}$ are defined in (27) and (26). Consider a gaussian process \tilde{H}_N on $\Sigma_{N,k}$ defined by

$$\tilde{H}_N(\sigma) = \frac{1}{\sqrt{N}} \sum_{i,j=1}^N \tilde{g}_{ij}(\alpha) \sigma_i \sigma_j \quad (38)$$

where $\tilde{g}_{ij}(\alpha)$ for $i, j = 1, \dots, N$ are i.i.d. copies of $\tilde{g}(\alpha)$ in (36).

Consider also a gaussian process G_N on $\Sigma_{N,k}$ independent from \tilde{H}_N defined by

$$G_N(\sigma) = \sum_{i=1}^N z_i(\alpha) \sigma_i \quad (39)$$

where $z_i(\alpha)$ for $i = 1, \dots, N$ are i.i.d. copies of $z(\alpha)$ in (37). Given two configurations $\sigma^1 = (\sigma^1, \alpha^1)$, $\sigma^2 = (\sigma^2, \alpha^2) \in \Sigma_{N,k}$ it's easy to check that the covariances of the process G_N and H_N are

$$\mathbb{E} G_N(\sigma^1) G_N(\sigma^2) = 2N c_{N,\tilde{\gamma}}(\sigma^1, \sigma^2) \tilde{\gamma}_{\alpha^1 \wedge \alpha^2} q_{\alpha^1 \wedge \alpha^2} \quad (40)$$

$$\mathbb{E} \tilde{H}_N(\sigma^1) \tilde{H}_N(\sigma^2) = N (c_{N,\tilde{\gamma}}(\sigma^1, \sigma^2))^2 \quad (41)$$

where

$$c_{N,\tilde{\gamma}}(\sigma^1, \sigma^2) = \tilde{\gamma}_{\alpha^1 \wedge \alpha^2} q_N(\sigma^1, \sigma^2) \quad (42)$$

For $t \in (0, 1)$ we define the interpolating Hamiltonian as

$$H_{N,t}(\boldsymbol{\sigma}) = \sqrt{t} \tilde{H}_N(\boldsymbol{\sigma}) + \sqrt{1-t} G_N(\boldsymbol{\sigma}) \quad (43)$$

and the interpolating pressure as

$$\varphi_N(t) = \frac{1}{N} \mathbb{E} \log Z_{N,t} \quad (44)$$

where

$$Z_{N,t} = \sum_{\boldsymbol{\sigma} \in \Sigma_{N,k}} \nu_{\boldsymbol{\alpha}} e^{H_{N,t}(\boldsymbol{\sigma})} \quad (45)$$

The Gibbs measure on $\Sigma_{N,k}$ associated to the Hamiltonian (43) is

$$\mu_{N,t}(\boldsymbol{\sigma}) = \frac{\nu_{\boldsymbol{\alpha}} e^{H_{N,t}(\boldsymbol{\sigma})}}{Z_{N,t}} \quad (46)$$

We denote by $\Omega_{N,t}(\cdot)$ the average w.r.t. $\mu_{N,t}^{\otimes \infty}$ and by $\langle \cdot \rangle_{N,t}$ the quenched expectation $\mathbb{E} \Omega_{N,t}(\cdot)$.

Keeping in mind that $q_N(\boldsymbol{\sigma}, \boldsymbol{\sigma}) = 1$ and $\tilde{\gamma}_{\boldsymbol{\alpha} \wedge \boldsymbol{\alpha}} = \tilde{\gamma}_k = \gamma_r$, then using integration by parts formula one obtains

$$2 \frac{d}{dt} \varphi_N = \tilde{\gamma}_k^2 - 2 (\tilde{\gamma}_k)^2 q_k + \langle (\tilde{\gamma}_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2} q_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2})^2 \rangle_{N,t} - \left\langle \left(c_{N,\tilde{\gamma}}(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2) - \tilde{\gamma}_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2} q_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2} \right)^2 \right\rangle_{N,t} \quad (47)$$

Now using the property (122) of RPC it's possible to show that

$$\langle (\tilde{\gamma}_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2} q_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2})^2 \rangle_{N,t} = \sum_{j \leq k} (\xi_j - \xi_{j-1}) (\tilde{\gamma}_j q_j)^2 = \tilde{\gamma}_k^2 - \sum_{0 \leq j \leq k-1} \xi_j \left((\tilde{\gamma}_{j+1} q_{j+1})^2 - (\tilde{\gamma}_j q_j)^2 \right) \quad (48)$$

In particular (47) implies that

$$\varphi_N(1) \leq \varphi_N(0) - \frac{1}{2} \sum_{0 \leq j \leq k-1} \xi_j \left((\tilde{\gamma}_{j+1} q_{j+1})^2 - (\tilde{\gamma}_j q_j)^2 \right) \quad (49)$$

Now since $H_{N,0}(\boldsymbol{\sigma}) \equiv G_N(\boldsymbol{\sigma})$ it holds

$$\varphi_N(0) = \frac{1}{N} \mathbb{E} \log \sum_{\boldsymbol{\alpha} \in \mathbb{N}^k} \nu_{\boldsymbol{\alpha}} \sum_{\boldsymbol{\sigma} \in \Sigma_N} e^{\sum_{i=1}^N z_i(\boldsymbol{\alpha}) \sigma_i} \quad (50)$$

and using again (122) one obtains

$$\varphi_N(0) = \mathbb{E} \log \sum_{\boldsymbol{\alpha} \in \mathbb{N}^k} \nu_{\boldsymbol{\alpha}} 2 \cosh(z(\boldsymbol{\alpha})) \quad (51)$$

Hence

$$\varphi(0) - \frac{1}{2} \sum_{0 \leq j \leq k-1} \xi_j \left((\tilde{\gamma}_{j+1} q_{j+1})^2 - (\tilde{\gamma}_j q_j)^2 \right) = \mathcal{P}_{\beta}(x) \quad (52)$$

On the other hand using the recursion in the property (122) one can represent $\varphi_N(1)$ in the following way. Let $\tilde{H}_N(\sigma, j)$ be a gaussian process on $(\sigma, j) \in \Sigma_N \times \{1, \dots, k\}$ with covariance

$$\mathbb{E} \tilde{H}_N(\sigma^1, j) \tilde{H}_N(\sigma^2, j') = N \delta_{jj'} \left(\sqrt{\tilde{\gamma}_j^2 - \tilde{\gamma}_{j-1}^2} q_N(\sigma^1, \sigma^2) \right)^2 \quad (53)$$

Then it holds

$$\varphi_N(1) = \frac{1}{N} \mathbb{E} \log \tilde{Z}_{0,N} \quad (54)$$

where $\tilde{Z}_{0,N}$ is obtained recursively starting from

$$\tilde{Z}_{k,N} = \sum_{\sigma} \prod_{1 \leq j \leq k} e^{H_N(\sigma, j)} \quad (55)$$

and for $0 \leq j \leq k-1$

$$\tilde{Z}_{j-1,N}^{\xi_{j-1}} = \mathbb{E}_{j-1} \tilde{Z}_{j,N}^{\xi_{j-1}} \quad (56)$$

where \mathbb{E}_j averages the randomness in $H_N(\sigma, j+1)$.

Now the key observation is that by definition the sequence $\tilde{\gamma}$ satisfies

$$\tilde{\gamma}_j = \gamma_l \text{ if } j \in K_l \quad (57)$$

If $\tilde{\gamma}_j = \tilde{\gamma}_{j-1}$ then by (53) the random variable $\tilde{H}_N(\sigma, j)$ is actually a centered gaussian with zero variance, namely its distribution is Dirac delta centered at the origin and it doesn't play any role. By (148) $\tilde{Z}_{0,N}$ can be represented using a new Ruelle Probability Cascade $(\tilde{\nu}_\alpha)_{\alpha \in \mathbb{N}^{k-1}}$ that is obtained from $(\nu_\alpha)_{\alpha \in \mathbb{N}^k}$ dropping the point process associated to the intensity ξ_{j-1} . A repeated use of the above argument implies that

$$\varphi(1) = p_N(\beta) \quad (58)$$

and then we get

$$p_N(\beta) \leq \mathcal{P}_\beta(x) \quad (59)$$

for every choice of the trial parameter $x \in X_\beta$ and then (35) follows.

□

4 The multi-scale Ghirlanda-Guerra identities

Consider quenched pressure density $p_N(\beta)$ in (14). It's standard to show that

$$\liminf_{N \rightarrow \infty} p_N(\beta) \geq \liminf_{N \rightarrow \infty} A_N \quad (60)$$

where

$$A_N = \mathbb{E} \log Z_{N+1} - \mathbb{E} \log Z_N \quad (61)$$

Now the strategy is to compare Z_{N+1} with Z_N . This procedure is known in mathematical-physics as Aizenman-Sims-Starr representation [3, 9]. Consider $\rho = (\sigma, \varepsilon) \in \Sigma_{N+1}$ with $(\sigma, \varepsilon) \in \Sigma_N \times \{-1, 1\}$ then

$$H_{N+1}(\rho, \alpha) = H'_N(\sigma, \alpha) + \varepsilon z_N(\sigma, \alpha) + O\left(\frac{1}{N}\right) \quad (62)$$

where

$$H'_N(\sigma, \alpha) = \frac{1}{\sqrt{N+1}} \sum_{i,j=1}^N g_{ij}(\alpha) \sigma_i \sigma_j \quad (63)$$

and

$$z_N(\sigma, \alpha) = \frac{1}{\sqrt{N+1}} \sum_{i=1}^N (g_{i,N+1}(\alpha) + g_{N+1,i}(\alpha)) \sigma_i \quad (64)$$

On the other hand

$$H_N(\sigma, \alpha) \stackrel{d}{=} H'_N(\sigma, \alpha) + y_N(\sigma, \alpha) \quad (65)$$

where

$$y_N(\sigma, \alpha) = \frac{1}{\sqrt{N(N+1)}} \sum_{i,j=1}^N g'_{ij}(\alpha) \sigma_i \sigma_j \quad (66)$$

for some array g' independent copy of g . Given two configurations $\sigma^1, \sigma^2 \in \Sigma_{N,r}$ the gaussian processes z_N and y_N defined in (64) and (66) respectively, have covariances

$$\mathbb{E} z_N(\sigma^1) z_N(\sigma^2) = 2 \frac{N}{N+1} \gamma_{\alpha^1 \wedge \alpha^2} c_{N,\gamma}(\sigma^1, \sigma^2) \quad (67)$$

$$\mathbb{E} y_N(\sigma, \alpha) y_N(\sigma', \beta) = \frac{N}{N+1} \left(c_{N,\gamma}(\sigma^1, \sigma^2) \right)^2 \quad (68)$$

The above relations implies that

$$A_N = \mathbb{E} \log \Omega'_N \left(2 \cosh(z_N(\sigma, \alpha)) \right) - \mathbb{E} \log \Omega'_N \left(\exp(y_N(\sigma, \alpha)) \right) \quad (69)$$

where $\Omega'_N = (\omega'_N)^{\otimes \infty}$ and ω'_N is the Gibbs measure on $\Sigma_{N,r}$ induced by the Hamiltonian H'_N in (63).

The Aizenmann-Sims-Starr representation A_N in (69) for the quenched pressure density has the same structure of the Parisi functional (33). Hence the strategy is to show that in the thermodynamic limit the distribution of $c_{N,\gamma}(\sigma^1, \sigma^2)$ under the random measure Ω'_N can be well approximated by a suitable RPC. We have two obstacles to overcome.

The first problem is to understand the joint probability distribution w.r.t. the limiting Gibbs measure of the two covariances $q_N(\sigma^1, \sigma^2)$ and $\gamma_{\alpha^1 \wedge \alpha^2}$. This situation is very similar to the case of the Multispecies SK model [5, 32] where it turns out that the Hamiltonian can be suitably perturbed in order to satisfy a *synchronization property* that allows to generate the joint probability of different overlaps functions using the same RPC. In addition since the parameter ξ associated to the RPC that express the Parisi functional (33) satisfies the condition $\zeta \subseteq \xi$, then the same must be true for the one that generates the limiting distribution of the above overlaps.

In this section we show that the Multiscale SK model can be suitably perturbed in order to satisfy the *synchronization property* that actually implies the condition $\zeta \subseteq \xi$.

Let H_N be the Hamiltonian function in (73) with parameters $\beta = (\zeta, \gamma)$ and $(\nu_\alpha)_{\alpha \in \mathbb{N}^r}$ the random weights of the RPC associated to the sequence ζ .

Let us consider a countable dense subset \mathcal{W} of $[0, 1]^2$ and a vector

$$w = (w_s)_{s=0,1} \in \mathcal{W} \quad (70)$$

For any $i \in \{0, \dots, N\}$, $w \in \mathcal{W}$ let us define

$$s_i(w) = \begin{cases} \sqrt{N} w_0 & \text{if } i = 0 \\ \sqrt{w_1} & \text{otherwise} \end{cases} \quad (71)$$

Let $(g^0(\alpha))_{\alpha \in \mathbb{N}^r}$ be a family of centered gaussian random variables with covariance

$$\mathbb{E} g^0(\alpha^1) g^0(\alpha^2) = \gamma_{\alpha^1 \wedge \alpha^2}, \quad (72)$$

Consider a gaussian process $h_{N,w,p}$ on $\Sigma_{N,r}$ defined by

$$h_{N,w,p}(\sigma) = \frac{1}{N^{p/2}} \sum_{i_1, \dots, i_p=0}^N g_{i_1, \dots, i_p}^{w,p}(\alpha) \sigma_{i_1} s_{i_1}(w) \cdots \sigma_{i_p} s_{i_p}(w) \quad (73)$$

where $\sigma_0 = 1$ while $g_{i_1, \dots, i_p}^{w,p}(\alpha)$ for $i_1, \dots, i_p = 1, \dots, N$, $p \geq 1$ and $w \in \mathcal{W}$ are i.i.d. standard gaussian random variables while if $i_l = 0$ for some $1 \leq l \leq p$ then $g_{i_1, \dots, i_p}^{w,p}(\alpha)$ is a family of i.i.d copies of $g^0(\alpha)$ in (72).

Then covariance of this process is

$$\mathbb{E} h_{N,w,p}(\boldsymbol{\sigma}^1) h_{N,w,p}(\boldsymbol{\sigma}^2) = (R_{N,w}(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2))^p \quad (74)$$

where

$$R_{N,w}(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2) = w_0 \gamma_{\alpha^1 \wedge \alpha^2} + w_1 q_N(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2) \quad (75)$$

We consider a weighted direct sum of the two previous overlaps because in the synchronization mechanism that we are going to exploit we need to control all the terms $\gamma^m q^n$ for generic integers m and n .

Since the set \mathcal{W} is countable, we can consider some one-to-one function $j : \mathcal{W} \rightarrow \mathbb{N}$. Consider now the following gaussian process

$$h'_N(\boldsymbol{\sigma}) = \sum_{w \in \mathcal{W}} \sum_{p \geq 1} 2^{-j(w)-p} (\sqrt{\gamma_r + 1})^{-p} x_{w,p} h_{N,w,p}(\boldsymbol{\sigma}) \quad (76)$$

where $X = (x_{w,p})_{w \in \mathcal{W}, p \geq 1}$ is a family of i.i.d. uniform random variables on $[1, 2]$.

Notice that the variance of the process h'_N is bounded uniformly on X , namely

$$\mathbb{E} h'_N(\boldsymbol{\sigma})^2 \leq 4 \quad (77)$$

For any $\boldsymbol{\sigma} \in \Sigma_{N,k}$ we define a perturbed Hamiltonian H_N^{pert} by

$$H_N^{\text{pert}}(\boldsymbol{\sigma}) = H_N(\boldsymbol{\sigma}) + s_N h'_N(\boldsymbol{\sigma}) \quad (78)$$

where s_N is a sequence of positive real numbers. We start observing that (77) implies that H_N^{pert} satisfies a *thermodynamic stability* condition

$$\mathbb{E} (H_N^{\text{pert}}(\boldsymbol{\sigma}))^2 \leq N \gamma_r^2 + 4 s_N^2 \quad (79)$$

uniformly on X . Consider the random function

$$\phi_{r,N} = \log \sum_{\boldsymbol{\sigma} \in \Sigma_{N,r}} \nu_{\boldsymbol{\alpha}} e^{H_N^{\text{pert}}(\boldsymbol{\sigma})} \quad (80)$$

Then $N^{-1} \mathbb{E} \phi_{r,N}$ is must be think as a small perturbation and the quantity $p_N(\beta)$ in (14). Indeed, it holds

$$p_N(\beta) \leq \frac{1}{N} \mathbb{E} \phi_{r,N} \leq p_N(\beta) + \frac{2s_N^2}{N} \quad (81)$$

Then if s_N satisfies

$$\lim_{N \rightarrow \infty} N^{-1} s_N^2 = 0 \quad (82)$$

the thermodynamic limits of $N^{-1} \mathbb{E} \phi_{r,N}$ and p_N coincide. Moreover RPC concentration inequality given in Proposition 5.1 implies that

$$\sup \{ \mathbb{E} |\phi_{r,N} - \mathbb{E} \phi_{r,N}| : 1 \leq x_p \leq 2, p \geq 1 \} \leq 4c(\zeta_0) \quad (83)$$

for some constant $c(\zeta_0)$ independent of N . Hence Theorem 3.2 in [29] and inequality (83) implies that if $s_N = N^\delta$ for $0 < \delta < 1/2$ we get the *Multispecies Ghirlanda-Guerra Identities* (Theorem 2 of [32]) that in our setting reads as follows.

Given two configurations $\boldsymbol{\sigma}^l = (\sigma^l, \boldsymbol{\alpha}^l), \boldsymbol{\sigma}^{l'} = (\sigma^{l'}, \boldsymbol{\alpha}^{l'}) \in \Sigma_{N,r}$ we set

$$R_{l,l'}(w) = R_{N,w}(\boldsymbol{\sigma}^l, \boldsymbol{\sigma}^{l'}) \quad (84)$$

and

$$R_{l,l'} = \begin{pmatrix} \gamma_{\boldsymbol{\alpha}^l \wedge \boldsymbol{\alpha}^{l'}} \\ q_N(\sigma^l, \sigma^{l'}) \end{pmatrix} \quad (85)$$

Given $n \geq 2$, let

$$R^n = (R_{l,l'})_{l,l' \leq n} \quad (86)$$

and for any bounded measurable function $f = f(R^n)$ we set

$$\langle f \rangle_N = \mathbb{E} \Omega_N(f) \quad (87)$$

where $\Omega_N = \mu_N^{\otimes \infty}$ while μ_N is the random Gibbs measure induced by H_N^{per} in (78).

For $p \geq 1$ and $w \in \mathcal{W}$ and conditionally on the i.i.d. uniform sequence $X = (x_{w,p})_{w \in \mathcal{W}, p \geq 1}$ let

$$\Delta_N(f, n, w, p, X) = \left| \langle f(R_{1,n+1}(w))^p \rangle_N - \frac{1}{n} \langle f \rangle_N \langle (R(w)_{1,2})^p \rangle_N - \frac{1}{n} \sum_{l=2}^n \langle f(R_{1,l}(w))^p \rangle_N \right| \quad (88)$$

By Theorem 2 in [32] we have that

$$\lim_{N \rightarrow \infty} \mathbb{E}_X \Delta_N(f, n, w, p, X) = 0 \quad (89)$$

where \mathbb{E}_X averages the random sequence X .

4.1 The Panchenko's synchronisation property

The synchronisation property is a powerful tool introduced by Panchenko [32] in his derivation of the lower bound for the multi-specie SK model [5]. It is moreover used in other mean-field settings [23,30,31].

By Lemma 3.3 in [29] there exists a non random sequence $X_N = (x_{w,p}^N)_{w \in \mathcal{W}, p \geq 1}$ such that (89) holds

$$\lim_{N \rightarrow \infty} \Delta_N(f, n, w, p, X_N) = 0 \quad (90)$$

In the rest of the work we assume to have such a sequence X_N . Consider the overlap function

$$Q_{l,l'} = \gamma_{\alpha^l \wedge \alpha^{l'}} + q_N(\sigma^l, \sigma^{l'}) \quad (91)$$

and the following overlap vector

$$\begin{pmatrix} R_{l,l'}^0 \\ R_{l,l'}^1 \end{pmatrix} = \begin{pmatrix} \gamma_{\alpha^l \wedge \alpha^{l'}} \\ q_N(\sigma^l, \sigma^{l'}) \end{pmatrix} \quad (92)$$

Consider also the arrays of the above overlap functions, namely

$$Q = (Q_{l,l'})_{l,l' \geq 1} \quad (93)$$

$$\begin{pmatrix} R^0 \\ R^1 \end{pmatrix} = \begin{pmatrix} R_{l,l'}^0 \\ R_{l,l'}^1 \end{pmatrix}_{l,l' \geq 1} \quad (94)$$

Let $(N_k)_{k \geq 1}$ be any subsequence along which the all above overlap arrays converges in distribution under the measure $\langle \cdot \rangle_N$. Since (90) holds, Theorem 3 in [32] implies that the arrays Q, R^0, R^1 satisfies the Ghirlanda-Guerra Identities [18,28], a factorisation property of the quenched equilibrium state (see also [1,13] for a related factorisation property).

Moreover Theorem 4 in [32] implies that the overlaps R^0 an R^1 are *synchronized*

Proposition 4.1. *For for any $s = 0, 1$ there exists a nondecreasing Lipschitz function*

$$L_0 : [0, \gamma_r + 1] \longrightarrow [0, \gamma_r], \quad L_1 : [0, \gamma_r + 1] \longrightarrow [0, 1]$$

such that

$$R_{l,l'}^s = L_s(Q_{l,l'}) \quad (95)$$

almost surely for all $l, l' \geq 1$

Notice that we can consider the domain and the range of L_s restricted to the positive real line because each of the overlap arrays Q, R^0, R^1 satisfies the Ghirlanda-Guerra identities and then the Talagrand's Positivity Principle holds (Theorem 2.16 in [29]).

The *synchronization property* of the previous Proposition is already a strong constraint on the limiting overlap distributions. Moreover by construction the overlap of the Multiscale SK model has an *a priori* hierarchical structure encoded in RPC with parameters ζ . The combination of these properties implies the following

Proposition 4.2. *Let $F_{Q_{12}}$ be the any weak limit of the distribution of one element of the array Q , then*

$$F_{Q_{12}} \in \mathcal{M}_\zeta[0, 2\gamma_r] \quad (96)$$

where $\mathcal{M}_\zeta[0, 2\gamma_r]$ is defined in 2.1.

Proof. The key observation is that the distribution of $R_{1,2}^0$ w.r.t. the perturbed Gibbs measure $\langle \cdot \rangle_N$ can be exactly computed for any N . Indeed by Theorem 3 of [33] it holds

$$\left\langle \mathbf{1}(\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2 = l) \right\rangle_N = \zeta_l - \zeta_{l-1} \quad (97)$$

for any $l \leq r$ and N integers.

Remark 1. *The quantity $\frac{1}{N} \mathbb{E} \phi_N$ has a recursive representation analogous to (17). In particular working with this representation (97) follows easily using the methods in [22].*

By definition $R_{1,2}^0 = \gamma_{\boldsymbol{\alpha}^1 \wedge \boldsymbol{\alpha}^2}$ then

$$\left\langle \mathbf{1}(R_{1,2}^0 \in A) \right\rangle_N = \sum_{l=0}^r \mathbf{1}(\{\gamma_l\} \in A) (\zeta_l - \zeta_{l-1}) \quad (98)$$

for any N and measurable set A . Since (98) doesn't depends on N the limit along any subsequence of the distribution of $R_{1,2}^0$ w.r.t. $\langle \cdot \rangle_N$ is given by (98). We denote by $\langle \cdot \rangle$ any of the above limiting measure that satisfies the synchronization property (95). Hence there exists a function L_0 such that

$$R_{1,2}^0 = L_0(Q_{1,2}) \text{ a.s.} \quad (99)$$

For any $l \leq r$ consider the set

$$A_l^0 = L_0^{-1}(\{\gamma_l\}) \quad (100)$$

Since L_0 is nondecreasing Lipschitz then A_l^0 is a closed interval or a single point and

$$\bigcup_{l \leq r} A_l^0 \equiv \text{supp}(Q_{1,2}) \quad (101)$$

Combining (98) and (99) we obtain

$$\left\langle \mathbf{1}(Q_{1,2} \in A_l^0) \right\rangle = \left\langle \mathbf{1}(R_{1,2}^0 = \gamma_l) \right\rangle = \zeta_l - \zeta_{l-1} \quad (102)$$

If we denote by Q_l^- the left extrema of A_l^0 then (102) and (101) implies that

$$F_{Q_{12}}(Q_l^-) = \zeta_{l-1} \quad (103)$$

for any $l \leq r$ and this proves the thesis. \square

5 Lower bound

Let $p_N(\beta)$ be the quenched pressure density of the Multiscale SK model (14) and replace the original Hamiltonian H_N with the perturbation H_N^{pert} in (78). We already know that this substitution doesn't affect the thermodynamic limit of $p_N(\beta)$. Moreover it entails a small change in the Aizenmann-Simms-Starr representation given in section 4. Indeed by Theorem 3.6 of [29] we have that

$$\liminf_{N \rightarrow \infty} p_N(\beta) \geq \liminf_{N \rightarrow \infty} \mathbb{E}_X A_N + o(1) \quad (104)$$

where $X = (x_{p,w})_{w \in \mathcal{W}, p \geq 1}$ is the family of random variables in (76) and

$$A_N = \mathbb{E} \log \Omega_N \left(2 \cosh(z_N(\sigma, \boldsymbol{\alpha})) \right) - \mathbb{E} \log \Omega_N \left(\exp(y_N(\sigma, \boldsymbol{\alpha})) \right) \quad (105)$$

Notice that A_N is the same functional appearing in (69) but now Ω_N is the infinite product of the random Gibbs measure induced by the Hamiltonian H_N^{pert} in (78).

Let us start observing that even if (105) is written in average over X , Lemma 3.3 of [29] ensures that one can choose a non random sequence $X_N = (x_p^{(N)})_{p \geq 1}$ such that

$$\liminf_{N \rightarrow \infty} p_N(\beta) \geq \liminf_{N \rightarrow \infty} A_N(X_N) + o(1) \quad (106)$$

and at the same time the *multi-scale Ghirlanda-Guerra Identities* (89) holds.

By (67) and Theorem 1.3 in [29] $A_N(X_N)$ is a continuous functional of the overlap array

$$\begin{pmatrix} R_{l,l'}^0 \\ R_{l,l'}^1 \end{pmatrix}_{l,l' \geq 1} = \begin{pmatrix} \gamma_{\boldsymbol{\alpha}^l \wedge \boldsymbol{\alpha}^{l'}} \\ q_N(\sigma^l, \sigma^{l'}) \end{pmatrix}_{l,l' \geq 1} \quad (107)$$

under the measure \mathbb{E}_{Ω_N} . Consider also the array

$$(Q_{l,l'})_{l,l' \geq 1} = (\gamma_{\alpha^l \wedge \alpha^{l'}} + q_{N,\gamma}(\sigma^l, \sigma^{l'}))_{l,l' \geq 1} \quad (108)$$

and a subsequence $(N_k)_{k \geq 1}$ along which all the above arrays converges in distribution to some arrays Q, R^0, R^1 w.r.t the measure induce by H_N^{per} . By construction the above arrays satisfy *multi-scale Ghirlanda-Guerra* (89) and then we can apply the results of section 4.1.

In particular by the *synchronization property* (Proposition 4.1) for any $s = 0, 1$ it holds

$$R_{l,l'}^s = L_s(Q_{l,l'}) \quad (109)$$

for some nondecreasing Lipschitz function L_s .

We denote by $\mu_{Q_{12}}$ the distribution of one element of the array Q . Let $k \geq 1$ be an integer and consider two sequences $\xi = (\xi_j)_{j \leq k}$ and $c = (c_j)_{j \leq k}$ such that

$$0 = \xi_{-1} < \xi_0 < \xi_1 < \dots < \xi_{r-1} < \xi_k = 1 \quad (110)$$

and

$$0 = c_0 < c_1 < \dots < c_k = 1 + \gamma_r \quad (111)$$

We choose the above couple (ξ, c) such that its associated discrete distribution ξ_c defined by (23) is close to $\mu_{Q_{12}}$ in some metric that metrizes weak convergence of distributions.

Moreover by Proposition 4.2 we know that the $F_{Q_{12}} \in \mathcal{M}_\zeta[0, 1 + \gamma_r]$ then we can assume without loss that the above ξ satisfies the key property

$$\zeta \subseteq \xi \quad (112)$$

Notice that (112) implies that $k \geq r$. Let $(\nu_\alpha)_{\alpha \in \mathbb{N}^k}$ be the random weights of the RPC associated to ξ in (110). By (89) the array Q satisfies the Ghirlanda-Guerra identities and then Theorems 2.13 and 2.17 in [32] imply that its distribution can be well approximated by the RPC associated to the above sequences ξ and q . This means that if we consider a family $(\alpha_l)_{l \geq 1}$ of i.i.d. samples from \mathbb{N}^k with distribution given by this RPC we have that the distribution of the array

$$(c_{\alpha^l \wedge \alpha^{l'}})_{l,l'} \quad (113)$$

will be close to the distribution of the array Q . For any $s = 0, 1$ we define a sequence

$$q_j^s = L_s(c_j) \quad 0 \leq j \leq k \quad (114)$$

then (109) implies that for any $s = 0, 1$ the distribution of the array

$$Q^s = (q_{\alpha^l \wedge \alpha^{l'}}^s)_{l, l'} \quad (115)$$

will be close to the distribution of the array R^s for any $s = 0, 1$.

We claim that the triple $(\xi, q^0, q^1) \in X_\beta$ where the set X_β is defined in (2.2). In other words we can set $q^0 \equiv \tilde{\gamma}$ and $q^1 \equiv q$ for some sequence $\tilde{\gamma}$ and q in (26) and (27) respectively.

Since we already know that ξ satisfies (112) it's enough to check that q_0 in (114) satisfies the condition (26). For a given $l \leq r$ consider the sets K_l and A_l^0 defined in (25) and (100) respectively. Then with probability one

$$c_j \in A_l^0 \Leftrightarrow j \in K_l \quad (116)$$

for any $j \leq k$ and any $l \leq r$. Hence, combining (114) and (116) we obtain that with probability one if $j \in K_l$ then $q_j^0 = \gamma_l$ which coincides with (26).

Given the above triple $(\xi, q^0, q^1) \in X_\beta$ consider the Parisi functional $\mathcal{P}(x)$ in (33). Notice that the quantity A_N (105) and are $\mathcal{P}(x)$ represented by the same continuous functional of the distribution of the arrays (R^0, R^1) in (107) and (Q^0, Q^1) in (115). Since by construction these arrays are close in some metric that metrizes weak convergence of distributions that one can use $\mathcal{P}_\beta(x)$ to approximate $A_N(X_N)$ as N goes to infinity (see Section 3.6 in [29]). Hence by (106)

$$\liminf_{N \rightarrow \infty} p_N(\beta) \geq \inf_{x \in X_\beta} \mathcal{P}_\beta(x) \quad (117)$$

and this conclude the proof of Theorem 2.1.

In this work we have analysed a multi-scale spin-glass mean-field model and obtained a variational principle that provides the solution for the free energy density. As a bypass result we obtained a full factorisation scheme of ultrametric nature. We plan to investigate how the multi-scale setting works with other mean-field cases, with hierarchical disordered models [11, 12] as well as to extend its use to finite dimensional models where alternative notions of equilibrium state, like for instance the *metastate* [26], have been advanced.

Acknowledgements

We want to thank several useful discussions with Diego Alberici, Francesco Guerra, Jorge Kurchan and especially Dmitry Panchenko whose observation led to a valuable improvement of proposition 5.1. P.C. was partially supported by PRIN project Statistical Mechanics and Complexity (2015K7KK8L), E.M. was partially supported by Progetto Almaidea 2018.

Appendix

For the benefit of the reader we summarise the main properties of Ruelle probability cascades used in the work. Here we follow Panchenko's monograph on the SK model [29]. For the interested reader we also mention the following works [4, 8, 27] on RPC and its applications to spin glasses theory.

Given an integer $r \geq 1$ let $\mathcal{A} = \mathbb{N}^0 \cup \mathbb{N} \cup \mathbb{N}^2 \dots \cup \mathbb{N}^r$ be a tree of depth r and root $\mathbb{N}^0 = \{\emptyset\}$. A vertex $\alpha = (n_1, \dots, n_p) \in \mathbb{N}^p$ for $1 < p < r$ has children $\alpha n = (n_1, \dots, n_p, n) \in \mathbb{N}^{p+1}$. Therefore each vertex $\alpha = (n_1, \dots, n_p)$ is connected to the root by the path

$$p(\alpha) = \{n_1, (n_1, n_2), \dots, (n_1, \dots, n_p)\} \quad (118)$$

We denote by $|\alpha|$ the distance between α and the root, namely the number of coordinates of α , thus by definition $\alpha \in \mathbb{N}^{|\alpha|}$. We also use the notation

$$\alpha \wedge \beta = |p(\alpha) \cap p(\beta)| \quad (119)$$

Let $\zeta = (\zeta_l)_{l=0, \dots, r-1}$ be a sequence such that

$$0 < \zeta_0 < \zeta_1 < \dots < \zeta_{r-1} < 1 \quad (120)$$

We denote by $(\nu_\alpha)_{\alpha \in \mathbb{N}^r}$ the random weights of the Ruelle probability cascade associated to the sequence ζ (Section 2.3 in [29]).

Consider a family of i.i.d. random variables $\omega = (\omega_p)_{1 \leq p \leq r}$ that have the uniform distribution on $[0, 1]$ and some function $X_r = X_r(\omega)$ which satisfies $\mathbb{E} \exp \zeta_{r-1} X_r < \infty$. Let us define recursively for $0 \leq l \leq r-1$

$$X_l = X_l(\omega_1, \dots, \omega_l) = \frac{1}{\zeta_l} \log \mathbb{E}_l \exp \zeta_l X_{l+1} \quad (121)$$

where \mathbb{E}_l denotes the expectation with respect to ω_{l+1} .

By definition X_0 is not random, moreover it can be represented through Ruelle Probability Cascades. Let $\omega_\alpha \in \mathcal{A} \setminus \mathbb{N}^0$ be a family of i.i.d. uniform $[0, 1]$ and set $\Omega_\alpha = (\omega_\beta)_{\beta \in p(\alpha)}$. Theorem 2.9 in [29] reads as follow

$$X_0 = \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^r} \nu_\alpha \exp X_r(\Omega_\alpha) \quad (122)$$

Actually the same argument used in [29] to prove (122) leads to a remarkable concentration result for Ruelle probability cascades.

Proposition 5.1. For any $r \geq 1$ the random variable

$$\phi_r = \log \sum_{\alpha \in \mathbb{N}^r} \nu_\alpha \exp X_r(\Omega_\alpha) \quad (123)$$

satisfies

$$\mathbb{E}(\phi_r - \mathbb{E}\phi_r)^2 \leq 4c(\zeta_0) \quad (124)$$

for some $c(\zeta_0)$ which doesn't depend on the distribution of X_r .

Proof. Let $(\nu_\alpha)_{\alpha \in \mathbb{N}^r}$ be the random weights of the Ruelle Probability Cascade associated to the sequence ζ in (120) that we rewrite as

$$\nu_\alpha = \frac{w_\alpha}{\sum_{\alpha \in \mathbb{N}^r} w_\alpha} \quad (125)$$

where the weights w_α are defined in section 2.3 of [29]. Let us start with the following lemma.

Lemma 1. Let $Z > 0$ be a random variable such that $\mathbb{E}Z^{\zeta_0} < \infty$ and let $(Z_\alpha)_{\alpha \in \mathbb{N}^r}$ be a sequence of i.i.d. copies of Z independent of all other random variables. For any $r \geq 1$ let

$$Y_r = \log \sum_{\alpha \in \mathbb{N}^r} w_\alpha Z_\alpha \exp X_r(\Omega_\alpha) \quad (126)$$

Then the following holds

$$\mathbb{E}(Y_r - \mathbb{E}Y_r)^2 = c(\zeta_0) < \infty \quad (127)$$

for some $c(\zeta_0)$ which doesn't depend on the distribution of X_r and Z .

Proof. The proof is by induction on r . Consider the case $r = 1$ then

$$Y_1 = \log \sum_{n \geq 1} w_n Z_n \exp(X_1(\Omega_n)) \quad (128)$$

The invariance property of the Poisson Dirichlet process (Theorem 2.6 in [29]) implies that

$$\sum_{n \geq 1} w_n Z_n \exp(X_1(\Omega_n)) \stackrel{d}{=} C \sum_{n \geq 1} w_n \quad (129)$$

where $C = \left(\mathbb{E}(Z \exp(X_1))^{\zeta_0}\right)^{1/\zeta_0}$. Since

$$\mathbb{E}(Y_1 - \mathbb{E}Y_1)^2 = \mathbb{E} \left(\log \sum_{n \geq 1} w_n Z_n \exp(X_1(\Omega_n)) - \mathbb{E} \log \sum_{n \geq 1} w_n Z_n \exp(X_1(\Omega_n)) \right)^2 \quad (130)$$

one can use the invariance property (129) in the r.h.s of the above line obtaining

$$\mathbb{E} (Y_1 - \mathbb{E} Y_1)^2 = \mathbb{E} \left(\log(C \sum_{n \geq 1} w_n) - \mathbb{E} \log(C \sum_{n \geq 1} w_n) \right)^2 = \mathbb{E} \left(\log \sum_{n \geq 1} w_n - \mathbb{E} \log \sum_{n \geq 1} w_n \right)^2 \quad (131)$$

Finally the same argument of Lemma 2.2 in [29] implies that

$$\mathbb{E} \log \sum_{n \geq 1} w_n < \infty, \quad \mathbb{E} \left(\log \sum_{n \geq 1} w_n \right)^2 < \infty$$

Therefore we can set

$$\mathbb{E} \left(\log \sum_{n \geq 1} w_n - \mathbb{E} \log \sum_{n \geq 1} w_n \right)^2 = c(\zeta_0) < \infty \quad (132)$$

for some $c(\zeta_0)$ that doesn't depends on the distribution of X_1 and Z .

Now for an arbitrary integer $r > 1$ consider the quantity

$$\mathbb{E} (Y_r - \mathbb{E} Y_r)^2 = \mathbb{E} \left(\log \sum_{\alpha \in \mathbb{N}^r} w_\alpha Z_\alpha \exp X_r(\Omega_\alpha) - \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^r} w_\alpha Z_\alpha \exp X_r(\Omega_\alpha) \right)^2 \quad (133)$$

The invariance property (2.57) in [29] implies that

$$\sum_{\alpha \in \mathbb{N}^r} w_\alpha Z_\alpha \exp(X_r(\Omega_\alpha)) \stackrel{d}{=} C \sum_{\alpha \in \mathbb{N}^{r-1}} w_\alpha U_\alpha \exp(X_{r-1}(\Omega_\alpha)) \quad (134)$$

where $C = (\mathbb{E} Z^{\zeta_{r-1}})^{1/\zeta_{r-1}}$ and $U_\alpha = \sum_{n \geq 1} u_{\alpha n}$ with $\mathbb{E} U_\alpha^{\zeta_{r-2}} < \infty$.

Then using (134) in the r.h.s. of (133) we obtain

$$\mathbb{E} (Y_r - \mathbb{E} Y_r)^2 = \mathbb{E} \left(\log \sum_{\alpha \in \mathbb{N}^{r-1}} w_\alpha U_\alpha \exp X_{r-1}(\Omega_\alpha) - \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^{r-1}} w_\alpha U_\alpha \exp X_{r-1}(\Omega_\alpha) \right)^2 \quad (135)$$

Finally notice that the above equation is of the same type of (127) with r replaced by $r - 1$ and Z_α by U_α and this conclude the proof by induction. □

Let's go back to the proof of Proposition 5.1. By (125) we can rewrite ϕ_r as

$$\phi_r = \tilde{\phi}_r + \log \sum_{\alpha \in \mathbb{N}^r} w_\alpha \quad (136)$$

where

$$\tilde{\phi}_r = \log \sum_{\alpha \in \mathbb{N}^r} w_\alpha \exp(X_r(\Omega_\alpha)) \quad (137)$$

Then we can write

$$\mathbb{E}(\phi_r - \mathbb{E}\phi_r)^2 \leq 2\mathbb{E}\left(\tilde{\phi}_r - \mathbb{E}\tilde{\phi}_r\right)^2 + 2\mathbb{E}\left(\log \sum_{\alpha \in \mathbb{N}^r} w_\alpha - \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^r} w_\alpha\right)^2 \quad (138)$$

Notice that we can apply Lemma 1 to compute the two terms in the r.h.s of (138) and this concludes the proof. \square

In this work we will use (122) also in the following particular setting. Let $q = (q_l)_{l=0, \dots, r}$ be a sequence such that

$$0 = q_0 < q_1 < \dots < q_r < \infty \quad (139)$$

and let $(J_l)_{1 \leq l \leq r}$ be a family of i.i.d. standard gaussian.

Consider a gaussian random variable

$$H_r = \sum_{1 \leq l \leq r} G_l \quad (140)$$

$$G_l = J_l (q_l - q_{l-1})^{1/2} \quad (141)$$

The covariance of G is given by

$$\mathbb{E} G_l G_l' = \delta_{l,l'} (q_l - q_{l-1}) \quad (142)$$

Consider the recursive construction (147) starting from

$$X_r = F(H_r) \quad (143)$$

for some function F that satisfies $\mathbb{E} \exp \zeta_{r-1} X_r < \infty$. Consider gaussian process g on \mathbb{N}^r defined by

$$g(\alpha) = \sum_{\beta \in p(\alpha)} J_\beta (q_{|\beta|} - q_{|\beta|-1})^{1/2} \quad (144)$$

where $(J_\alpha)_{\alpha \in \mathcal{A} \setminus \mathbb{N}^0}$ is a family of i.i.d. standard gaussian random variables. The covariance of the process g is

$$\mathbb{E} g(\alpha) g(\beta) = q_{\alpha \wedge \beta} \quad (145)$$

Then (122) implies that

$$X_0 = \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^r} \nu_\alpha \exp F(g(\alpha)) \quad (146)$$

Suppose that instead of (139) we have that $q_l = q_{l-1}$ for some $l \in \{1, \dots, r\}$. Then the random variable G_l in (141) is actually a centered gaussian with zero variance, namely its distribution is a Dirac delta at the origin. This implies that one can set $G_l \equiv 0$ and forget the average E_{l-1} getting $X_{l-1} = X_l$. In other words X_0 can be represented using a new Ruelle Probability Cascade $(\tilde{\nu}_\alpha)_{\alpha \in \mathbb{N}^{r-1}}$ that is obtained from $(\nu_\alpha)_{\alpha \in \mathbb{N}^r}$ dropping the point process associated to the intensity ζ_{l-1} .

Formally we consider the sequence $\tilde{\zeta} = \zeta \setminus \{\zeta_{l-1}\}$ and denote by $(\tilde{\nu}_\alpha)_{\alpha \in \mathbb{N}^{r-1}}$ the random weights of the Ruelle Probability Cascade associated to the sequence $\tilde{\zeta}$. Let ϕ be the one-to-one map between the sets $\{0, \dots, r\} \setminus \{l-1\}$ and $\{0, \dots, r-1\}$ and replace H_r in (140) with $\tilde{H}_{r-1} = \sum_{1 \leq l' \neq l \leq r} G_{\phi(l')}$ and starting from $\tilde{X}_{r-1} = F(\tilde{H}_{r-1})$ we recursively define

$$\tilde{X}_{\phi(l')} = \frac{1}{\zeta_{l'}} \log \mathbb{E}_l \exp \zeta_{l'} \tilde{X}_{\phi(l'+1)} \quad (147)$$

for any $0 < l' \neq l < r-1$. Then it holds

$$X_0 = \tilde{X}_0 = \mathbb{E} \log \sum_{\alpha \in \mathbb{N}^{r-1}} \tilde{\nu}_\alpha \exp F(\tilde{g}(\alpha)) \quad (148)$$

where $\tilde{g}(\alpha)$ is defined as in (144).

References

- [1] M.Aizenman, P.Contucci, On the stability of the quenched state in mean-field spin-glass models. *Journal of Statistical Physics*, Vol. 92, N. 5/6, 765-783, (1998).
- [2] A. Auffinger, A. W. Chen, The Parisi Formula has a Unique Minimzer. *Communications in Mathematical Physics*, **335**, Issue 3, pp 1429-1444 (2015)
- [3] M. Aizenman M., R. Sims , S. Starr S., An Extended Variational Principle for the SK Spin-Glass Model. *Phys. Rev. B*, 68:214403, (2003)
- [4] L.-P. Arguin, Spin glass computations and Ruelle's probability cascades. *J. Stat. Phys.*, 126(4-5):951-976, 2007
- [5] A. Barra, P. Contucci, E. Mingione, D. Tantari, Multi-species mean-field spin-glasses. Rigorous results. *Ann. Henri Poincaré*, **16**, 691-708 (2015)
- [6] A. Barra, F. Guerra and E. Mingione, Interpolating the Sherrington-Kirkpatrick replica trick. *Philosophical Magazine*, **92**, Issue 1-3, 78-97 (2012)

- [7] E. Bolthausen, N. Kistler, On a nonhierarchical version of the Generalized Random Energy Model, II: Ultrametricity. *Stochastic Processes and their Applications* **119**, Issue 7, 2357-2386, (2009).
- [8] E. Bolthausen, A.-S. Sznitman, On Ruelle's probability cascades and an abstract cavity method. *Comm. Math. Phys.* **197**(2), 247-276, (1998).
- [9] A. Bovier, A. Klimovsky, The Aizenman-Sims-Starr and Guerra schemes for the SK model with multidimensional spins *Electron. J. Probab.*, **14**, Nr. 8, 161-241, (2009)
- [10] A. Bovier, I. Kurkova, Derridas Generalized Random Energy models I-II *Annals de l' Institut Henri Poincaré*, **40**, 4 (2004)
- [11] M. Castellana, A. Barra, F. Guerra, Free-energy bounds for hierarchical spin models *Jou. Stat. Phys.*, 155- 2, pp 211222, (2014)
- [12] M. Castellana, G. Parisi, Non Perturbative effects in spin glasses. *Scirep, Nature* **5**, 8697 (2015)
- [13] P. Contucci, C. Giardinà, C. Giberti, Stability of the Spin Glass Phase under Perturbations *Europhysics Letters*, Vol. 96, N. 1, 17003-17006, (2011)
- [14] L. Cugliandolo, J. Kurchan, Thermal properties of slow dynamics, *Physica A: Statistical Mechanics and its Applications*, **263**, Issues 14, 242-251, 1999
- [15] L. Cugliandolo, J. Kurchan, A Scenario for the Dynamics in the Small Entropy Production Limit *Journal of the Physical Society of Japan* **69** (Suppl.A), 247-256, 2000
- [16] B. Derrida, E. Gardner, Solution of the generalized random energy model, *J. Phys. C* **19**, 2253 (1986).
- [17] G. Gallavotti, Renormalization theory and ultraviolet stability for scalar fields via renormalization group methods. *Rev. Mod. Phys.*, **57** 471 (1985)
- [18] S. Ghirlanda, F. Guerra, General properties of overlap probability distributions in disordered spin systems. Towards Parisi ultrametricity. *J. Phys. A: Math. Gen.* **31**, 9149-9155 (1998).
- [19] F. Guerra, Mathematical aspects of mean field spin glass theory. *Proceedings of the "4th European Congress of Mathematics"*, Stockholm, 2004
- [20] F. Guerra, Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model. *Comm. Math. Phys.*, **233**, 1-12 (2003).
- [21] F. Guerra, F.L. Toninelli, The thermodynamical limit in mean field spin glass model. *Comm. Math. Phys.*, **230**, 71-79 (2002)

- [22] F. Guerra, Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model. *Comm. Math. Phys.*, **233**, 1-12, 2003.
- [23] A. Jagannath, J. Ko, S. Sen, A connection between MAX k -CUT and the inhomogeneous Potts spin glass in the large degree limit. <https://arxiv.org/abs/1703.03455>
- [24] M. Mezard, G. Parisi and M. A. Virasoro, Spin Glass Theory and Beyond. *World Scientific*, 1987
- [25] R. Monasson, Structural glass transition and the entropy of the metastable states *Phys Rev Lett.*, 75(15):2847-2850, 1995
- [26] D. Stein, C. Newman, Spin Glasses and Complexity *Oxford University Press*, 2013
- [27] D. Ruelle, A mathematical reformulation of Derrida's REM and GREM. *Commun. Math. Phys.* **108**, 225 (1987).
- [28] D. Panchenko, The Parisi ultrametricity conjecture. *Annals of Mathematics* **177**, Issue 1, 383-393, 2013.
- [29] D. Panchenko, The Sherrington-Kirkpatrick Model. *Springer, New York* (2013)
- [30] D. Panchenko, Free energy in the mixed p -spin models with vector spins. *Annals of Probability* **46**, Nr. 2, 865-896, 2018
- [31] D. Panchenko, Free energy in the Potts spin glass *Annals of Probability* **46**, Nr. 2, 829-864, 2018
- [32] D. Panchenko, The free energy in a multispecies Sherrington Kirkpatrick model. *Annals of Probability*, **46**, No. 6, 3494-3513, 2015
- [33] D. Panchenko, M. Talagrand, <https://arxiv.org/abs/0708.3641>
- [34] J. Polchinski, Renormalization and Effective Lagrangians *Nucl.Phys. B*, **231**, 269-295, 1984.
- [35] M. Talagrand, Large Deviations, Guerra's and A.S.S. Schemes, and the Parisi Hypothesis *Journal of Statistical Physics*, **126**, Issue 4-5, 837-894, 2007
- [36] M. Talagrand, The Parisi formula. *Annals of Mathematics*, **163**, N. 1, 221-263, 2006