

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

In vivo and in vitro effects of selected antioxidants on rabbit meat microbiota

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

In vivo and in vitro effects of selected antioxidants on rabbit meat microbiota / Albonetti, Sabrina; Minardi, Paola; Trombetti, Fabiana; Savigni, Fabiana; Mordenti, Attilio Luigi; Baranzoni, Gian Marco; Trivisano, Carlo; Greco, Fedele Pasquale; Badiani, Anna. - In: MEAT SCIENCE. - ISSN 0309-1740. - ELETTRONICO. - 123:(2017), pp. 88-96. [10.1016/j.meatsci.2016.09.004]

Availability:

This version is available at: <https://hdl.handle.net/11585/566319> since: 2016-11-03

Published:

DOI: <http://doi.org/10.1016/j.meatsci.2016.09.004>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Albonetti, Sabrina, Paola Minardi, Fabiana Trombetti, Fabiana Savigni, Attilio Luigi Mordenti, Gian Marco Baranzoni, Carlo Trivisano, Fedele Pasquale Greco, e Anna Badiani. 2017. «In Vivo and in Vitro Effects of Selected Antioxidants on Rabbit Meat Microbiota». *Meat Science* 123 (gennaio): 88–96.

The final published version is available online at:
<https://doi.org/10.1016/j.meatsci.2016.09.004>.

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

***In vivo and in vitro* effects of selected antioxidants on rabbit meat microbiota**

Sabrina Albonetti^{a*}, Paola Minardi^c, Fabiana Trombetti^b, Fabiana Savigni^a, Attilio Luigi Mordenti^c,
Gian Marco Baranzoni^a, Carlo Trivisano^d, Fedele Pasquale Greco^d, Anna Badiani^c

^aLaboratorio di Igiene e Tecnologia Alimentare, ^bLaboratorio di Biochimica, ^cLaboratorio di Zootechnia, Nutrizione ed Alimenti, Dipartimento di Scienze Mediche Veterinarie, Via Tolara di Sopra 50, 40064 Ozzano Emilia (BO), Italy.

^dDipartimento di Scienze Statistiche “Paolo Fortunati”, Via Belle Arti 41, 40126 Bologna, Italy.

* Corresponding author: Sabrina Albonetti, Dipartimento di Scienze Mediche Veterinarie, Via Tolara di Sopra 50, 40064 Ozzano Emilia (BO), Italy. Tel: + 39.0547.338929; Fax: +39.0547.338941; E-mail address: sabrina.albonetti@unibo.it

Abstract

The purpose of this study was to investigate the effect of dietary vitamin E or EconomasETM supplementation on the growth of several background/pathogenic bacteria on rabbit carcasses and hamburgers during refrigerated storage. For 51 days, 270 New Zealand rabbits received either a basal diet, or experimental diets enriched with 100 or 200 mg/kg of vitamin E or EconomasETM. The bacteria studied were *Salmonella*, *Listeria monocytogenes*, *Pseudomonas*, *Enterobacteriaceae*, *Escherichia coli*, coagulase-positive staphylococci, plus both mesophilic and psychrotrophic aerobes. The growth of *Listeria monocytogenes* on contaminated patties was evaluated through a challenge test. The potential protective or antimicrobial effect of vitamin E or EconomasETM on *Listeria monocytogenes* or *Pseudomonas aeruginosa* were assessed *in vitro*. Diet did not influence the concentrations of bacteria found on rabbit carcasses and developing on hamburgers. Vitamin E (*in vivo* and *in vitro*) and EconomasETM *in vivo* had a protective antioxidant role, while EconomasETM *in vitro* had strong antibacterial activity against *Listeria monocytogenes*, but not against *Pseudomonas aeruginosa*.

Keywords: Meat safety; Antioxidants; Rabbit meat microbiota; Challenge test; *Listeria monocytogenes*; *Pseudomonas aeruginosa*

Chemical compounds studied in this article

35 DL α tocopherol acetate (PubChem CID: 86472); ethanol (PubChem CID: 702).

36

37

1. Introduction

Rabbit meat presents excellent nutritional and dietetic properties and meets the current demand for low-fat meat (Dalle Zotte & Szendrő, 2011). However, rabbit meat is expensive, time-consuming to prepare and rather perishable because it is prone to oxidative damage due to its high level of polyunsaturated fatty acids (Abdel-Khalek, 2013). Consequently, in order to expand the market, in addition to retail fresh cuts, many rabbit meat industries have tried to approach the consumer through the production of meat preparations, such as hamburgers or patties, which may benefit from antioxidants. Given the above, a relevant question might be raised, as already suggested by Sofos, Cabedo, Zerby, Belk, & Smith (2000): are those antioxidants able to protect bacterial cellular membranes as well, when present on the very same meat or meat preparation? There is plenty of literature regarding the effects of antioxidants on rabbit meat and meat preparations, but special attention has been paid to vitamin E, i.e. DL- α -tocopherol (VE) (Castellini, Dal Bosco, & Bernardini, 2001; Castellini, Dal Bosco, Bernardini, & Cyril, 1998; Dal Bosco, Castellini, Bianchi, & Mugnai, 2004; Lo Fiego et al., 2004). Due to its high antioxidant activity, VE, especially as α -tocopherol acetate, is commonly used in animal feed to promote growth and to improve meat quality; VE is deposited in muscle cell membranes and lipid depots, thus reducing lipid oxidation, which is one of the most significant causes of meat deterioration during refrigeration (**Hu et al., 2015**). Moreover, among the antioxidants it is possible to include the EconomasETM (EcoE), a patented commercial premixture of nutritional additives consisting mainly of L-ascorbic acid (50 000 mg/kg) and organic selenium produced by *Saccharomyces cerevisiae* CNCMI-3060 (750 000 mg/kg). Selenium in yeast is incorporated into organic compounds, mainly selenomethionine, and low molecular weight seleno-components. Selenium is an essential trace element involved in various physiological functions; as an integral part of selenoproteins, it plays an important role in the antioxidant defense system against reactive molecules and free radicals (**Ahmad et al., 2012**; Mehdi, Hornick, Istasse, & Dufrasne, 2013). Despite its importance, to our knowledge, information about the effect of dietary supplementation of antioxidant on microbial growth in rabbit carcasses and meat preparations is almost non-existent. In detail, carcasses and meat preparations from rabbits fed additional levels of VE or EcoE have not been studied so far in terms of their microbial status.

At each step of the food chain, meat and meat preparations might be contaminated and cold storage does not always inhibit the growth of bacteria. In particular, *Listeria monocytogenes* is a ubiquitous pathogen, and is especially dangerous because it is able to grow also at refrigeration temperatures, unlike most other foodborne pathogens (EFSA, 2014; Swaminathan, Cabanes, Zhang,

& Cossart, 2007). A wide variety of meats and processed products have been associated with *L. monocytogenes* contamination at a prevalence which can be high because of various conditions of storage, distribution and handling in addition to inadequate bacterial inactivation (Swaminathan et al., 2007). Furthermore, *L. monocytogenes* survives in foods for a long time, even under adverse conditions (Ramaswamy et al., 2007; Rocourt, BenEmbarek, Toyofuku, & Schlundt, 2003) and it causes severe symptoms and diseases (meningitis, septicemia and abortion) (**Ramaswamy et al., 2007**). It must also be remembered that human listeriosis cases in Europe have been increasing in recent years (EFSA & ECDC, 2014). In contrast, the *Pseudomonas* genus represents the dominant contaminant on rabbit carcasses and other packed meat (Bobbitt, 2002; Rodríguez-Calleja, Santos, Otero, & García-López, 2004). In particular, *Pseudomonas aeruginosa* is a food spoilage agent included in the list of bacteria carrying a biological risk, unlike all other species of *Pseudomonas*. In fact, the public health interest in these two microorganisms stems from the fact that they are both human pathogens and, according to regulations in Europe and the United States, these two bacteria are classified in risk group 2 on the basis of biohazard (EC, 2000; HHS, 2013).

The purpose of this work was to investigate the effect of dietary VE or EcoE supplementation on the growth of eight types of background or pathogenic bacteria on rabbit carcasses and rabbit meat preparations (hamburgers) during refrigerated storage. The growth of *Listeria monocytogenes* on contaminated patties was also evaluated through a challenge test. The potential protective or antimicrobial effect of VE or EcoE on *L. monocytogenes* or *Pseudomonas aeruginosa* were also assessed *in vitro*.

2. Materials and methods

This work represents the microbiological part of a multidisciplinary research project designed to evaluate the shelf-life of rabbit meat, including the study of carcass quality and the technological, nutritional and sensory quality of rabbit meat.

2.1. Animals and diets

Two hundred and seventy commercial New Zealand white rabbits (*Oryctolagus cuniculis*) provided by the Rabbit Genetic Centre of the Martini Group were selected for this study. Thirty-five-day-old males from a single breeding were randomly divided into five experimental **units (e.u.)** of 54 animals each. Every **e.u.** was housed under controlled temperature and light conditions (12 h light/12 h dark photoperiod cycle), equally and randomly divided into three cages (= **replicates**) having provision of *ad libitum* feeding and watering. A starter complete basal diet for growing rabbits and a subsequent finisher diet for fattening rabbits were formulated to meet the nutrient

106 requirements of the animals during the experimental period (Table 1).

107 Two antioxidants in two different concentrations were tested in this work. The basal diets of
108 two *e.u.* were supplemented with 100 or 200 mg/kg of DL- α -tocopherol acetate (Sigma-Aldrich, St.
109 Louis, MO, USA) (indicated as VE 100 and VE 200, respectively) while the diets of other two *e.u.*
110 were supplemented with 100 or 200 mg/kg of EcoE (Alltech Ireland Ltd., Dunboyne, Ireland)
111 (indicated as EcoE 100 and EcoE 200, respectively), as suggested by the producer. The remaining
112 *e.u.* was fed a normal diet and used as a control (CTRL). After 51 days, 256 animals (mortality
113 5.2%) were slaughtered in the Ma.Ge.Ma abattoir (Savignano sul Rubicone, FC, Italy); rabbits
114 underwent electrical stunning followed by cutting of the carotid arteries and jugular veins. Two
115 carcasses were discarded due to abscesses. Slaughter weights (g) \pm standard errors (SE) were:
116 2991 ± 35.09 (CTRL); 2934 ± 29.18 (VE 100); 2867 ± 38.85 (VE 200); 2905 ± 37.15 (EcoE 100);
117 2981 ± 30.31 (EcoE 200). The abattoir structure, layout and hygiene procedures were in compliance
118 with European Union requirements (EC, 2004). All handling procedures followed the
119 recommendations of the European Council Directive 86/609/EEC for the protection of animals used
120 for experimental and other scientific purposes (EEC, 1986).

121 Ten carcasses were randomly selected out of each *e.u.* (total number = 50). The selected
122 carcasses were transported to the DIMEVET laboratory of Food Hygiene and Technology in
123 accordance with traceability and cold chain. After 24 h at 4°C, carcass hygiene was tested and then
124 carcasses were used to produce hamburgers and patties.

125

126 2.2. Microbiological analyses

127 Microbiological assays on rabbit carcasses and meat preparations were performed using
128 international standard methods. Samples were prepared according to the ISO standard 6887-1 (ISO,
129 1999) and 6887-2 (ISO, 2003a) and were diluted with a solution of 0.1% tryptone (Oxoid Ltd.,
130 Basingstoke, England) and 0.85% NaCl (Oxoid Ltd.) in distilled water. ISO standard 6579 (ISO,
131 2007) and ISO 11290-1 (ISO, 2004a) were used respectively to detect *Salmonella* spp. and *L.*
132 *monocytogenes*, while ISO standard 4833-2 (ISO, 2013), 17410 (ISO, 2001b), 21528-2 (ISO,
133 2004c), 16649-2 (ISO, 2001a), 13720 (ISO, 2010), 6888-1 (ISO, 2003b), and 11290-2 (ISO, 2004b)
134 were used respectively to enumerate aerobic mesophilic bacteria, aerobic psychrotrophic bacteria,
135 *Enterobacteriaceae*, *Escherichia coli*, *Pseudomonas* spp., coagulase-positive staphylococci, and *L.*
136 *monocytogenes*.

137

138 2.3. Feed and carcass hygiene

139 The feeds for growing and fattening rabbits were preliminarily sampled and examined for *L.*

140 *monocytogenes* and *Salmonella* spp., as described in the specific ISO standards (subsection number
141 2.2).

142 After 24 h at 4°C, 15 random post-chill rabbit carcasses (three from each *e.u.*), according to
143 European Regulation No. 2073/2005 which rules carcass sampling on the slaughter line (EC, 2005),
144 were tested for the detection of *Salmonella* spp. and *L. monocytogenes*, and for the enumeration of
145 aerobic mesophilic bacteria, aerobic psychrotrophic bacteria, *Enterobacteriaceae*, *E. coli*,
146 *Pseudomonas* spp. and coagulase-positive staphylococci, as described in the specific ISO standards
147 (subsection number 2.2). The whole carcasses were sampled according to standards 17604 (ISO,
148 2003c) and 6887-2 (ISO, 2003a), using the excision method. In particular, 50 g of surface tissue (~
149 2 mm deep) from the neck, both external scapular regions, thorax, brisket, flanks, fore rib, and hind
150 limbs were obtained with sterile scalpels and forceps. Then, 10 g of tissue were cut using sterile
151 scissors, placed in sterile stomacher bags, diluted ten-fold, and blended for two minutes in a
152 stomacher (Lab blender 400, Abbot Park, USA) for enumeration procedures; 25 g was used for
153 detection procedures.

154

155 2.4. Trend of natural background bacteria on rabbit hamburgers

156 ***Ten post-chill carcasses from each e.u. were boned and minced using a refrigerated***
157 ***mincer (TC 32 Frozen, Sirman Spa, Padua, Italy) to generate a single batch of minced meat from***
158 ***which all the necessary hamburgers and patties were produced, without any seasoning or***
159 ***additive to avoid interferences with the tested antioxidants. The “single batch approach” was***
160 ***regarded as mandatory within the e.u., in that it was necessary to prepare a pabulum as***
161 ***homogeneous as possible and therefore not affected by intra e.u. differences, in order both to***
162 ***study the growth of natural background bacteria and to perform the Listeria challenge testing,***
163 ***each determination being repeated as many times as requested by the relevant ISO standard.***

164 Ninety hamburgers (100 ± 3 g) and 135 patties (30 ± 3 g) (18 and 27 respectively for each
165 *e.u.*) were produced, aerobically packaged, two by two (hamburgers) or three by three (patties) in
166 polystyrene trays, and wrapped with food plastic film.

167 All the trays were stored at 0-2°C in a cabinet (Quartet 200, Costan, Belluno, Italy) which
168 was closed with a lid every night to reproduce retail storage conditions. On the day of production
169 (Time 0) and after 1, 2, 4, 6, 8, 10, 12, and 16 days, one tray with two hamburgers per *e.u.* was
170 tested for the detection of *Salmonella* spp. and *L. monocytogenes*, and for the enumeration of
171 aerobic mesophilic bacteria, aerobic psychrotrophic bacteria, *Enterobacteriaceae*, *E. coli*,
172 *Pseudomonas* spp., and coagulase-positive staphylococci as described in the specific ISO standards
173 (subsection number 2.2). After each day of analysis, these ten hamburgers were discarded.

Patties were used for the *Listeria* challenge testing described in subsection 2.5.

2.5. *Listeria* challenge testing

Challenge testing was performed to assess the growth of *L. monocytogenes* in artificially contaminated patties of rabbit meat during and beyond the commercial shelf-life period. Four *L. monocytogenes* strains were used for the artificial contamination: a reference strain ATCC 7644 (clinical isolate) and three strains from the internal collection (87-1771 and 115-1924 from pork meat; 88-1777 from frozen chicken). All strains were stored in vials containing trypticase soy broth (TSB, Becton, Dickinson and company, Le Pont de Claix, France) with 0.6% yeast extract (YE, Becton Dickinson France SA, Le Pont de Claix, France) and 20% glycerol (Carlo Erba Reagents, Milan, Italy) at - 20°C. The inocula for each strain were prepared adding 100 µL of stock solution to 10 mL TSB-YE. The tube was incubated overnight at 37°C then 100 µL from every grown culture were added in 10 mL TSB-YE and incubated at 7°C for 96 h. *L. monocytogenes* strains cultures were grown at low temperatures to reduce the lag time period after inoculation in food samples (Uyttendaele et al., 2004). Appropriate volumes of diluted bacterial strains were mixed together and used to prepare an inoculum of 50 CFU/g. Contamination was performed by spreading the mixed bacterial cultures on the surface of patties using sterile glass rods. Contaminated patties were kept at 20°C for five minutes to allow adhesion of the bacterial cells on the product surface (Pal, Labuza, & Diez-Gonzalez, 2008) and then they were stored in the cabinet. The inoculum concentration was verified by plate counting on trypticase soy agar supplemented with 0.6% YE (TSA-YE). A tray with three patties for each *e.u.* was tested for the enumeration of *L. monocytogenes* on the same day as the artificial contamination (Time 0) and after 1, 2, 4, 6, 8, 10, 12, and 16 days. The patties were discarded after analysis.

2.6. In vitro effect of vitamin E and EconomasE™ on the growth of *L. monocytogenes* and *P. aeruginosa*

The effect of VE or EcoE supplementation on microbial growth in rabbit meat was further studied by evaluating the growth of two bacteria in TSB-YE with different concentrations of the two antioxidants. The strains *L. monocytogenes* ATCC 7644 (stored in vials containing TSB-YE supplemented with 20% glycerol at - 20°C) and *P. aeruginosa* ATCC 27853 (stored in dry pellet disk at 4°C) were used for these experiments. The inocula were prepared adding 100 µL of the stock solution or dissolving one pellet disk in 10 mL of TSB-YE followed by overnight incubation at 37°C. Then, 100 µL of each grown culture were transferred to a new tube containing 10 mL of TSB-YE and incubated at 37°C for 24 h. The culture was appropriately diluted and a volume of

approximately 100-200 µL added to a tube containing 10 mL of TSB-YE supplemented with one of the antioxidants to obtain a final concentration of 10 CFU/mL or 100 CFU/mL. VE (Sigma-Aldrich, St. Louis, MO, USA) was dissolved in ethanol 96% (solubility of 100 µL/mL) and diluted in TSB-YE to obtain concentrations of 10⁴, 10³, and 10² ppm (Table 2). The effect of ethanol at 10%, 1%, and 0.1% on bacterial growth and without VE was checked as well. For both bacterial strains, the three concentrations of VE along with the three controls were contaminated with both 10 CFU/mL and 100 CFU/mL. All the tubes were incubated at 37°C for *L. monocytogenes* and 25°C for *P. aeruginosa*, and after 24 h and 48 h the bacteria were quantified, as described in subsection 2.2. The experiment was repeated three times (Table 2).

EcoE was dissolved in TSB-YE through sonication for 45 minutes (Ultrasonic UTA Falc Instruments Srl, Treviglio, Italy); 4×10³ ppm was the highest concentration achieved. Differential thermal analyses were also performed before and after sonication to verify this process did not lead to physical and chemical changes in the additive. After appropriate dilutions in TSB-YE, the concentrations 4×10³, 3×10³, 2×10³, 10³, and 10² ppm were contaminated with 10 CFU/mL or 100 CFU/mL for both bacteria. All the tubes were incubated at 37°C for *L. monocytogenes* and 25°C for *P. aeruginosa*, and after 24 h and 48 h the bacteria were quantified as explained in subsection 2.2. The experiment was repeated three times.

225

226 2.7. Statistical analysis

Bacterial concentrations were transformed to a log₁₀ scale. The analysis of data concerning natural background bacteria on rabbit hamburgers and *Listeria* challenge testing (see sections 2.4 and 2.5) was performed by comparing growth curves estimated on the basis of microbial concentrations observed at times t=0, 1, 2, 4, 6, 8, 10, 12, 16. For each microorganism and for each diet, three different growth models were compared, namely the logistic model, the Gompertz model with the parameterizations proposed in Zwietering, Jongenburger, Rombouts, & Van't Riet (1990), and the Baranyi model with the parameterization proposed in Baranyi and Roberts (1994). These growth models are described in equations (1) - (3).

235 (1) Logistic model

$$236 \log_{10} N_t = \log_{10} N_0 + \frac{\log_{10}(N_{\infty} / N_0)}{1 + \exp\left(4\mu_m(\lambda - t)\ln(10)\left(\log_{10}(N_{\infty} / N_0)\right)^{-1} + 2\right)}$$

237 (2) Gompertz model

$$238 \log_{10} N_t = \log_{10} N_0 + \log_{10}(N_{\infty}/N_0) \exp\left(-\exp\left(\frac{e\mu_m(\lambda - t)}{\log_{10}(N_{\infty} / N_0)\ln(10)} + 1\right)\right)$$

239

240 (3) Baranyi model

$$241 \quad \log_{10} N_t = \log_{10} N_0 + \log_{10} \left(\frac{-1 + \exp(\mu_m \lambda) + \exp(\mu_m t)}{\exp(\mu_m t) - 1 + \exp(\mu_m \lambda) 10^{\log_{10}(N_\infty/N_0)}} \right)$$

242 Estimates of model parameters were obtained by means of the function *nls* (non-linear least
243 squares) implemented in the statistical software R version 3.1.0 (Copyright© 2011 The R
244 Foundation for Statistical Computing).

245 For each diet and microorganism, the best fitting model was selected on the basis of the
246 residual sum of squares (RSS) statistic. Since different growth models could be selected, direct
247 comparison in terms of parameters estimates is not meaningful; indeed, parameter interpretation is
248 different across models. The rationale for this approach is to favour model fitting over the
249 comparability of parameters.

250 In order to compare estimated curves for each microorganism, Confidence Intervals (CI) for
251 growth curves were obtained at each observational time following a bootstrap approach (Efron &
252 Tibshirani, 1993), as implemented in the R package *nlstool*. This is a distribution-free procedure
253 which is preferable to the usual approximation based on the normality assumption that holds in non-
254 linear regression only for large samples.

255 With regard to the *in vitro* effect of VE or EcoE on the growth of *L. monocytogenes* ATCC
256 7644 and *P. aeruginosa* ATCC 27853 (subsection number 2.6), only two observational times were
257 available: for this reason, growth curve estimation was not feasible and experimental results were
258 analyzed by means of an ANOVA model where observational time was considered as a
259 dichotomous experimental factor. Experimental vials were assigned to two experimental groups
260 defined by two different bacterial inoculum values and observed after 24 h and 48 h.

261 Let Y_{ijkt} be the \log_{10} concentration, at time t , of *L. monocytogenes* or *P. aeruginosa* in the i -
262 the experimental vial ($i=1,2,3$) assigned to the k -th ($k=1,2$) level of inoculum (I), and the j -the
263 treatment defined as follows:

- 264 $j=1$: Ethanol percentage=0.1, VE=0;
- 265 $j=2$: Ethanol percentage=0.1, VE=100;
- 266 $j=3$: Ethanol percentage=1, VE=0;
- 267 $j=4$: Ethanol percentage=1, VE=1000;
- 268 $j=5$: Ethanol percentage=10, VE=0;
- 269 $j=6$: Ethanol percentage=10, VE=10000.

270 The ANOVA model is specified as follows:

$$271 \quad Y_{ijkl} = \alpha + \mu_j^V + \mu_k^I + \mu_t^T + \mu_{jk}^{V:I} + \mu_{jt}^{V:T} + \mu_{kt}^{I:T} + \varepsilon_{ijkl} \quad \varepsilon_{ijkl} \sim N(0, \sigma^2) \quad (1)$$

272 where α denotes the general intercept and captures the \log_{10} concentration at the baseline,
273 parameters μ denote the main effects, parameters γ denote the second-order interactions.
274 Superscripts I , V and T refer respectively to experimental variables inoculum, vitamin E and time.
275 The model was parameterized such that the baseline represents the \log_{10} concentration at the first
276 level of each variable; i.e. all the considered effects are equal to 0 when $j = k = t = 1$ (baseline). A
277 stepwise selection procedure (not shown) suggested to ignore third-order interaction term.

278 Regarding EcoE, it was possible to manage the experimental variable EcoE (E) as a
279 continuous variable. As a consequence the specification of the model, selected using a stepwise
280 procedure, is:

$$281 \quad Y_{ijkl} = \alpha + \beta^E E_j + \mu_k^I + \mu_t^T + \beta_k^{E:I} E_{jk} + \beta_t^{E:T} E_{jt} + \gamma_{kt}^{I:T} + \varepsilon_{ijkl} \quad \varepsilon_{ijkl} \sim N(0, \sigma^2) \quad (2)$$

282 Parameter α denotes the general intercept: in order to maintain the interpretation of this intercept as
283 the \log_{10} concentration at baseline, variable E was shifted by subtracting its minimum value. The
284 main effects of variables I and T were captured by parameter μ , while γ denotes their second-order
285 interaction. Parameter β^E is the slope of the linear relationship between EcoE and bacterial
286 concentration when $k = 1$ and $t = 1$, while parameters $\beta_t^{E:T}$ and $\beta_k^{E:I}$ are the effect modifiers
287 referring to time and inoculum, respectively.

288

289 3. Results and discussion

290

291 3.1. Feed and carcass hygiene

292 The feeds for growing and fattening rabbits were negative for *L. monocytogenes* and
293 *Salmonella* spp.. The microbiota on 24 h post-chilled rabbit carcasses is summarized in Table 3; *L.*
294 *monocytogenes*, *Salmonella* spp., and coagulase-positive staphylococci were not reported since they
295 were absent in all carcasses, indicating good slaughtering practice.

296 Rabbit meat is not mentioned in the European Regulation No. 2073/2005 on the
297 microbiological criteria for the acceptability of carcasses (EC, 2005); for this reason, dressing of
298 rabbit carcasses was compared with that of beef carcasses. The \log_{10} means mesophilic aerobes and
299 *Enterobacteriaceae* were in agreement with other reports and, even if they were slightly high for
300 hygienically processed meat, values fell within the European limits for red meat. *Pseudomonas*
301 spp., most of them fluorescent strains, mesophilic aerobes and psychrotrophic aerobes, were the
302 main microorganisms of all groups after slaughter and revealed similar mean counts regardless of

the dietary treatment. *E. coli* counts were low, in accordance with the data of Bobbitt (2003). Relative variability among *e.u.* was observed for *E. coli* and *Enterobacteriaceae* with a variation within 1.5 log CFUs. *Pseudomonas* genus predominated on rabbit meat (Table 3), as already observed by Bobbitt (2002) and Rodríguez-Calleja et al. (2004). *Pseudomonas* is commonly the dominant meat spoilage bacteria at refrigerated temperatures, driven by enhanced catabolism of glucose and lactate (García-López, Prieto, & Otero, 1998).

3.2. Evolution of microbial growth on rabbit hamburgers

The hamburger background bacteria trend was determined using the counts of several microbiological indicators plotted as a function of time to monitor microbial population dynamics throughout refrigerated storage. The results of the microbiological analyses are shown in Fig. 1. Regarding the growth curves obtained, the choice of the best fitting model implies the selection of different growth curve families for different *e.u.*. As a consequence, direct comparison in terms of parameter estimates is not feasible, since parameter interpretation differs across models; the rationale for this approach is to favour model fitting over the comparability of parameter estimates. Comparison between growth curves is based on graphical examination of the estimated curves, along with the CIs obtained at the observational times.

First, it is important to note the absence of bacteria of public health significance: *Salmonella* spp., *L. monocytogenes*, and coagulase-positive staphylococci were absent in all samples, indicating good handling and meat processing practices during hamburger preparation. Initial concentrations of the other microbial populations investigated at time 0 fell within the acceptability limits set out in EC Regulation No. 2073/2005 (EC, 2005), although referring to other animal species. As expected, all microbial counts of all *e.u.* considerably increased throughout refrigerated storage. On average, growth rates for mesophiles, psychrotrophic bacteria, and *Pseudomonas* spp. were similar, reaching the plateau phase at day 4, independent from the tested diets. The storage flora of rabbit hamburgers was dominated by the genus *Pseudomonas*, in agreement with data reported by Soutos, Tzikas, Christaki, Papageorgiou, & Steris (2009). Regarding *E. coli*, hamburgers derived from the control *e.u.* (CTRL) showed the highest initial bacterial concentrations but the lowest and slowest growth. The most rapid and abundant growth was sustained by VE 200, followed by EcoE 200 and EcoE 100. Therefore, except for *E. coli*, there were no significant differences among the dietary treatments: the curves and the corresponding CIs overlapped considerably. This result was primarily influenced by the high initial concentrations of *E. coli* commonly present on rabbit meat.

3.3. *Listeria monocytogenes* challenge test

Fig. 2 shows the growth of *L. monocytogenes* in contaminated rabbit patties during and beyond the commercial shelf-life period. On the first day of analysis (Time 0), CTRL presented the highest amount of bacteria, significantly different from VE 100 and EcoE 200, probably because of slight contamination during patty manipulation, but the bacterial growth was the slowest throughout the subsequent period of analysis. EcoE 200 significantly allowed the highest and fastest growth rate, followed by EcoE 100 and VE 200, which overlapped. The relatively low initial level of *L. monocytogenes* suggested differences among dietary fortifications, confirming, as for *E. coli*, that EcoE (both amounts) and VE 200 supported the highest and fastest bacterial growth.

3.4. In vitro effect of vitamin E and EconomasE™ on the growth of *L. monocytogenes* and *P. aeruginosa*

The absence of information on the biological role of antioxidants on microbial growth in foods required *in vitro* assays investigating contact between VE or EcoE and the two chosen bacterial strains.

The interpretation of the results regarding VE was complex because the effect of the ethanol used as solvent must also be considered. VE, in fact, is insoluble in water, while it is soluble in organic solvents such as ethanol which at concentrations up to 1.25% does not inhibit bacteria growth, but bacteria are strongly inhibited in the presence of 5% ethanol (Oh & Marshall, 1993). Referring to the experimental design explained above and summarized in Table 2, analyses concerning *P. aeruginosa* ATCC 27853 refer to data collected at grid cells enclosed in the dashed line, while analyses concerning *L. monocytogenes* ATCC 7644 refer to the whole table. In fact, *P. aeruginosa* growth was inhibited at 10% ethanol, regardless of the starter inoculum. The results obtained for *L. monocytogenes* are reported in Table 4. The main effects capture the difference between the \log_{10} counts at baseline and the \log_{10} counts in experimental blocks, where all but one experimental variable was kept constant. The estimate of μ_2^I highlights a statistically significant increase corresponding to the increase in starter inoculum while the estimate of μ_2^T denotes a statistically significant reduction after 48 h. As regards the main effects μ_j^V , $j > 1$, results show a highly significant reduction in \log_{10} counts in experimental blocks where Ethanol=10. Second-order interactions measure the variation of \log_{10} counts with respect to the sum of the baseline and the main effects involved; as an example, an estimate of $\gamma_{62}^{V:I}$ equal to 1.628 means that the expected \log_{10} count where Ethanol=10, VE=10000 and inoculum=100 is equal to $9.432+0.954-9.868+1.628=2.146$. Estimates concerning second-order interactions were statistically significant only in blocks where Ethanol=10. VE seemed to protect bacterial cells since it allowed for the

370 survival and growth of *L. monocytogenes* in the presence of toxic levels of ethanol. The results
 371 obtained for *P. aeruginosa* are summarized in Table 5. Parameter interpretation was analogous to
 372 the previous model. It is worth noting that all parameters were statistically significant, except μ_2^T
 373 and $\gamma_{22}^{I:T}$; while time had a non-significant main effect, estimates of interactions between time and
 374 experimental factor V ($\gamma_{22}^{V:T}$, $\gamma_{32}^{V:T}$ and $\gamma_{42}^{V:T}$) demonstrated an increase in log₁₀ counts after 48 h at
 375 non-baseline values of V . Inoculum showed a positive main effect, but interactions with
 376 experimental factor V were all negative.

377 Regarding EcoE assays, it is important to emphasize that the contents of selenium were
 378 equal (for EcoE 100 ppm) or one log unit higher (for EcoE 1000 ppm) than the average level of
 379 selenium reported in rabbit muscle (0.18 ppm) by Puls (1988). Results concerning the *in vitro* effect
 380 of EcoE on the growth of *L. monocytogenes* were not analyzed by means of a statistical model since
 381 no appreciable variations were observed. As can be seen from Fig. 3, bacterial growth was
 382 completely inhibited in the presence of 1000 ppm and subsequent doses of EcoE for the inoculum
 383 10 CFU/mL (a), while with 100 CFU/mL there was a slight growth only at 24 h (b). On the
 384 contrary, *P. aeruginosa* showed growth at every dose of EcoE. Table 6 summarizes parameter
 385 estimates concerning the linear model (2).

386 Marginal effects of inoculum and time were significantly greater than zero. The estimate of
 387 the interaction term $\gamma_{22}^{I:T}$ highlights a lower growth when inoculum =100 and time =48 with respect
 388 to the growth expected, considering only marginal effects (note that the sum of the two main effects
 389 and the interaction term is positive). The negative sign of the estimate of β^E shows that, when
 390 inoculum =10 CFU/mL and time =24 h, the commercial additive had an inhibitory effect on
 391 microbial growth. However, when time =48 h, the estimate of $\beta_2^{E:T}$ shows that EcoE had a positive
 392 effect on bacterial growth. It can be assumed that the bacterium adapted over time.

393 According to the reported results, the microbiota found on rabbit carcasses and developing
 394 on rabbit hamburgers did not diversify on the basis of the different antioxidants added to the diet.
 395 However, dietary treatment with EcoE (both amounts) and VE 200 corresponded to the highest
 396 amount and the fastest growth rate of *E. coli* and *L. monocytogenes* on rabbit hamburgers and
 397 patties, respectively. VE is a powerful chain-breaking antioxidant with an essential role in
 398 maintaining the structural integrity of biological membranes, in which it primarily resides (Sun et
 399 al., 2012). It can therefore be assumed that antioxidant protection of tissue against oxidative damage
 400 also promotes microorganism growth. There is no clear evidence of the role of vitamins in
 401 improving the survival of bacteria, as these supplements are used to enhance the quality of the final
 402 product and expand the shelf-life period, but Shan, Ding, Fallourd, & Leyer (2010) reported that the

403 addition of ascorbic acid (vitamin C) can protect probiotic cells. According to Murata, Tanaka,
404 Kubo, & Fujita (2013), VE protects *S. aureus* cells from oxidative stress via free radical generation
405 induced by cardol (C_{15:3}). The role and effects of selenium as an antioxidant in most rabbit studies is
406 unclear (Abdel-Khalek, 2013), but it is important to emphasize that selenium has been investigated
407 for medical applications: Yang et al. (2009) reported a strong inhibitory activity of selenium-
408 enriched probiotics against pathogenic *E. coli* *in vivo* and *in vitro*; a series of organoselenium
409 compounds were successfully tested as antibacterial agents.

410 In the present study EcoE completely inhibited the Gram-positive *L. monocytogenes* *in vitro*
411 with doses ≥ 1000 ppm (corresponding to a toxic level of selenium of 1.5 ppm), but it had no effect
412 against the Gram negative *P. aeruginosa*. The resistance of *P. aeruginosa* can be explained by the
413 inability of selenium to cross the protective outer membrane of Gram-negative bacteria and
414 accumulate at the cell membrane or within the cytoplasm. However, this may not be the only
415 explanation, since low molecular weight components are able to reach the periplasm through the
416 porin proteins on the outer membranes (Helander et al., 1998). Another reason could be that
417 selenium is pumped out from the periplasm exceeding its penetration rate, but it is clear that further
418 studies are needed to better elucidate the effects of selenium on bacterial metabolism. Instead, EcoE
419 showed only a protective antioxidant role *in vivo*.

420

421 **4. Conclusions**

422

423 VE (*in vivo* and *in vitro*, since it allowed for the survival of bacteria at a toxic level of
424 ethanol) and EcoE *in vivo* had a protective antioxidant effect on bacteria. EcoE *in vitro* showed
425 strong antibacterial activity against Gram positive *L. monocytogenes* but not against Gram negative
426 *P. aeruginosa*. Diet did not influence the concentrations of bacteria found on rabbit carcasses and
427 developing on hamburgers.

428 This work is the first to study the effect of these selected antioxidant dietary supplements on
429 the microbiological status of rabbit carcasses and rabbit meat preparations.

430

431

432 **Acknowledgments**

433

434 This work was funded by the Department of Veterinary Medical Science (University of
435 Bologna). The authors are grateful to Martini Group (Martini Spa, Budrio di Longiano, FC, Italy)
436 who provided both animals and feeds and to Alltech (Alltech Ireland Ltd., Dunboyne, Ireland) who
437 provided EcoE. Thanks are extended to Isidoro Giorgio Lesci and Loretta Antoni for their skilful
438 technical assistance, Andrea Capitani (Alltech Italy) for his helpful attitude, and Proof-Reading-
439 Service.com for English language editing.

440

441

442 **References**

443

444 Abdel-Khalek, A. M. (2013). Supplemental antioxidants in rabbit nutrition: a review. *Livestock*
445 *Science*, 158, 95–105.

446 Ahmad, H., Tian, J., Wang, J., Khan, M. A., Wang, Y., Zhang, L., & Wang, T. (2012). Effects of
447 dietary sodium selenite and selenium yeast on antioxidant enzyme activities and oxidative
448 stability of chicken breast meat. *Journal of Agricultural and Food Chemistry*, 60,
449 7111–7120.

450 Baranyi, J., & Roberts, T. A. (1994). A dynamic approach to predicting bacterial growth in food.
451 *International Journal of Food Microbiology*, 23, 277–294.

452 Bobbitt, J. (2002). Shelf life and microbiological safety of selected new and emerging meats
453 destined for export markets. Rural Industries Research and Development Corporation,
454 RIRDC Publication No. 02–038. Available at: [https://rirdc.infoservices.com.au/items/02-](https://rirdc.infoservices.com.au/items/02-038)
455 [038](https://rirdc.infoservices.com.au/items/02-038). Accessed November 15, 2015.

456 Bobbitt, J. (2003). Buffalo, Camel, Crocodile, Emu, Kangaroo, Ostrich and Rabbit Meat. New
457 value added products. Rural Industries Research and Development Corporation, RIRDC
458 Publication No. 03–036. Available at:
459 [http://australiancamelindustry.com.au/cjamel/images/pdfs/camelmeat/RIRDC.Buffalo.Came](http://australiancamelindustry.com.au/cjamel/images/pdfs/camelmeat/RIRDC.Buffalo.Camel.Crocodile.Emu.Kangaroo.Ostrich.and.Rabbit.Meat.May.2003.pdf)
460 [l.Crocodile.Emu.Kangaroo.Ostrich.and.Rabbit.Meat.May.2003.pdf](http://australiancamelindustry.com.au/cjamel/images/pdfs/camelmeat/RIRDC.Buffalo.Camel.Crocodile.Emu.Kangaroo.Ostrich.and.Rabbit.Meat.May.2003.pdf). Accessed November 22,
461 2015.

462 Castellini, C., Dal Bosco, A., & Bernardini, M. (2001). Improvement of lipid stability of rabbit
463 meat by vitamin E and C administration. *Journal of the Science of Food and Agriculture*, 81,
464 46–53.

465 Castellini, C., Dal Bosco, A., Bernardini, M., & Cyril, H. W. (1998). Effect of dietary vitamin E on
466 the oxidative stability of raw and cooked rabbit meat. *Meat Science*, 50, 153–161.

467 Dal Bosco, A., Castellini, C., Bianchi, L., & Mugnai, C. (2004). Effect of dietary α -linolenic acid
468 and vitamin E on the fatty acid composition, storage stability and sensory traits of rabbit
469 meat. *Meat Science*, 66, 407–413.

470 Dalle Zotte, A. & Szendrő, Z. (2011). The role of rabbit meat as functional food. *Meat Science*, 88,
471 319–331.

472 EC (2000). Directive 2000/54/EC of the European Parliament and of the Council of 18 September
473 2000 on the protection of workers from risks related to exposure to biological agents at work
474 (seventh individual directive within the meaning of Article 16(1) of Directive 89/391/EEC).
475 *Official Journal of the European Communities*, L 262/, 17/10/2000, 21–45.

476 EC (2004). Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29
 477 April 2004 laying down specific hygiene rules for on the hygiene of foodstuffs. *Official*
 478 *Journal of the European Union*, L 139, 30/04/2004, available at:
 479 <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=OJ:L:2004:139:TOC>, accessed
 480 September 29, 2015.

481 EC (2005). Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological
 482 criteria for food-stuffs. *Official Journal of the European Union*, L 338, 22/12/2005, 1–26.

483 EEC (1986). Council Directive No 609/1986 of 24 November 1986 on the approximation of laws,
 484 regulations and administrative provisions of the Member States regarding the protection of
 485 animals used for experimental and other scientific purposes. *Official Journal of the*
 486 *European Communities*, L 358, 18/12/1986, 1–28.

487 Efron, B., & Tibshirani, R. J. (1993). *An introduction to the bootstrap*. New York: Chapman &
 488 Hall, (Chapter 13).

489 EFSA (2014). European Food Safety Authority (EFSA). Scientific Opinion on the public health risks
 490 related to the maintenance of the cold chain during storage and transport of meat. Part 2
 491 (minced meat from all species). *EFSA Journal*, 12(7):3783, 1–30.

492 EFSA & ECDC (2014). European Food Safety Authority (EFSA) & European Centre for Disease
 493 Prevention and Control (ECDC). The European Union summary report on trends and
 494 sources of zoonoses, zoonotic agents and food-borne outbreaks in 2012. *EFSA Journal*,
 495 12(2):3547, 1–312.

496 García-López, M. L., Prieto, M., & Otero, A. (1998). The physiological attributes of Gram-negative
 497 bacteria associated with spoilage of meat and meat products. In A. R. Davies, & R. G.
 498 Board. (Eds), *The microbiology of meat and poultry* (pp. 1–34). London: Blackie Academic
 499 and Professional.

500 Helander, I. K., Alakomi, H. L., Latva-Kala, K., Mattila-Sandholm, T., Pol, I., Smid, E. J., Gorris,
 501 L. G. M., & von Wright, A. (1998). Characterization of the action of selected essential
 502 components on Gram negative bacteria. *Journal of Agricultural and Food Chemistry*, 46,
 503 3590–3595.

504 HHS (2013). U.S. Department of Health and Human Services. NIH Guidelines for research
 505 involving recombinant or synthetic nucleic acid molecules (NIH Guidelines), November
 506 2013. 1–136. Available at http://osp.od.nih.gov/sites/default/files/NIH_Guidelines_0.pdf.
 507 Accessed October 07, 2015.

508 Hu, Z. P., Wang, T., Ahmad, H., Zhang, J. F., Zhang, L. L., & Zhong, X. (2015) Effects of different
509 formulations of α -tocopherol acetate (vitamin E) on growth performance, meat quality and
510 antioxidant capacity in broiler chickens. *British Poultry Science*, 56, 687–695.

511 ISO (1999). International Organization for Standardization. *Microbiology of food and animal*
512 *feeding stuffs - Preparation of test samples, initial suspension and decimal dilutions for*
513 *microbiological examination. Part 1: General rules for the preparation of the initial*
514 *suspension and decimal dilutions*. ISO, 6887-1. Geneva, Switzerland.

515 ISO (2001a). International Organization for Standardization. *Microbiology of food and animal*
516 *feeding stuffs - Horizontal method for the enumeration of β -glucuronidase-positive*
517 *Escherichia coli. Part 2: Colony-count technique at 44°C using 5-bromo-4-chloro-3-indolyl*
518 *β -D-glucuronide*. ISO, 16649-2. Geneva, Switzerland.

519 ISO (2001b). International Organization for Standardization. *Microbiology of food and animal*
520 *feeding stuffs - Horizontal method for the enumeration of psychrotrophic microorganisms*.
521 ISO, 17410. Geneva, Switzerland.

522 ISO (2003a). International Organization for Standardization. *Microbiology of food and animal*
523 *feeding stuffs - Preparation of test samples, initial suspension and decimal dilutions for*
524 *microbiological examination. Part 2: Specific rules for the preparation of meat and meat*
525 *products*. ISO, 6887-2. Geneva, Switzerland.

526 ISO (2003b). International Organization for Standardization. *Microbiology of food and animal*
527 *feeding stuffs - Horizontal method for the enumeration of coagulase-positive staphylococci*
528 *(Staphylococcus aureus and other species) - Technique using Baird-Parker agar medium*.
529 ISO, 6888-1:1999/Amd.1:2003. Geneva, Switzerland.

530 ISO (2003c). International Organization for Standardization. *Microbiology of food and animal*
531 *feeding stuffs - Carcass sampling for microbiological analysis*. ISO, 17604. Geneva,
532 Switzerland.

533 ISO (2004a). International Organization for Standardization. *Microbiology of food and animal*
534 *feeding stuffs - Horizontal method for the detection and enumeration of Listeria*
535 *monocytogenes. Part 1: Detection method*. ISO, 11290-1:1996/Amd.1:2004. Geneva,
536 Switzerland.

537 ISO (2004b). International Organization for Standardization. *Microbiology of food and animal*
538 *feeding stuffs - Horizontal method for the detection and enumeration of Listeria*
539 *monocytogenes. Part 2: Enumeration method*. ISO, 11290-2:1998/Amd.1:2004. Geneva,
540 Switzerland.

541 ISO (2004c). International Organization for Standardization. *Microbiology of food and animal*
542 *feeding stuffs - Horizontal methods for the detection and enumeration of*
543 *Enterobacteriaceae. Part 2: Colony-count method*. ISO, 21528-2. Geneva, Switzerland.

544 ISO (2007). International Organization for Standardization. *Microbiology of food and animal*
545 *feeding stuffs - Horizontal method for the detection of Salmonella spp.*. ISO,
546 6579:2002/Amd.1:2007. Geneva, Switzerland.

547 ISO (2010). International Organization for Standardization. *Meat and meat products - Enumeration*
548 *of presumptive Pseudomonas spp.*. ISO, 13720. Geneva, Switzerland.

549 ISO (2013). International Organization for Standardization. *Microbiology of the food chain -*
550 *Horizontal method for the enumeration of microorganisms. Part 2: Colony count at 30*
551 *degrees C by the surface plating technique*. ISO, 4833-2. Geneva, Switzerland.

552 Lo Fiego, D. P., Santoro, P., Macchioni, P., Mazzoni, D., Piattoni, F., Tassone, F., & De Leonibus,
553 E. (2004). The effect of dietary supplementation of vitamins C and E on the α -tocopherol
554 content of muscles, liver and kidney, on the stability of lipids, and on certain meat quality
555 parameters of the longissimus dorsi of rabbits. *Meat Science*, 67, 319–327.

556 Mehdi, Y., Hornick, J., Istasse, L., & Dufrasne I. (2013). Selenium in the environment, metabolism
557 and involvement in body functions. *Molecules*, 18, 3292–3311.

558 Murata, W., Tanaka, T., Kubo, I., & Fujita, K. (2013). Protective effects of α -tocopherol and
559 ascorbic acid against cardol-induced cell death and reactive oxygen species generation in
560 *Staphylococcus aureus*. *Journal of Medicinal Plant and Natural Product Research*, 79,
561 768–774.

562 Oh, D. - H., & Marshall, D. L. (1993). Antimicrobial activity of ethanol, glycerol monolaurate or
563 lactic acid against *Listeria monocytogenes*. *International Journal of Food Microbiology*, 20,
564 239–246.

565 Pal, A., Labuza, T. P., & Diez-Gonzalez, F. (2008). Shelf life evaluation for ready-to-eat sliced
566 uncured turkey breast and cured ham under probable storage conditions based on *Listeria*
567 *monocytogenes* and psychrotroph growth. *International Journal of Food Microbiology*, 126,
568 49–56.

569 Puls, R. (1988). *Mineral levels in animal health: diagnostic data*. Clearbrook, British Columbia:
570 Sherpa International, (Chapter 28).

571 Ramaswamy, V., Cresence, V. M., Rejitha, J. S., Lekshmi, M. U., Dharsana, K. S., Prasad, S. P., &
572 Vijila, H. M. (2007). *Listeria* – review of epidemiology and pathogenesis. *Journal of*
573 *Microbiology, Immunology and Infection*, 40, 4–13.

574 Rocourt, J., BenEmbarek, P., Toyofuku, H., & Schlundt, J. (2003). Quantitative risk assessment of
575 *Listeria monocytogenes* in ready-to-eat foods: the FAO/WHO approach. *FEMS Immunology*
576 *and Medical Microbiology*, 35, 263–267.

577 Rodríguez-Calleja, J. M., Santos, J. A., Otero, A., & García-López, M. (2004). Microbiological
578 quality of rabbit meat. *Journal of Food Protection*, 67, 966–971.

579 Shan, N. P., Ding, W. K., Fallourd, M. J., & Leyer, G. (2010). Improving the stability of probiotic
580 bacteria in model fruit juices using vitamins and antioxidants. *Journal of Food Science*,
581 75(5), M 278–282.

582 Sofos, J. N., Cabedo, L., Zerby, H., Belk, K. E., & Smith, G. C. (2000). Potential interaction
583 between antioxidants and microbial meat quality. In E.A., Decker, C., Faustman, & C.J.,
584 Lopez-Bote (Eds), *Antioxidants in muscle foods - Nutritional strategies to improve quality*
585 (pp. 427–453). New York (NY), USA: Wiley & Sons, Inc.

586 Soultos, N., Tzikas, Z., Christaki, E., Papageorgiou, K., & Steris, V. (2009). The effect of dietary
587 oregano essential oil on microbial growth of rabbit carcasses during refrigerated storage.
588 *Meat Science*, 81, 474–478.

589 Sun, Y., Ma, A., Li, Y., Han, X., Wang, Q., & Liang, H. (2012). Vitamin E supplementation
590 protects erythrocyte membranes from oxidative stress in healthy Chinese middle-aged and
591 elderly people. *Nutrition Research*, 32, 328–334.

592 Swaminathan, B., Cabanes, D., Zhang, W., & Cossart, P. (2007). *Listeria monocytogenes*. In M.P.,
593 Doyle, & L.R., Beuchat, (Eds.), *Food microbiology: fundamentals and frontiers* (pp.
594 457–492). 3rd ed. Washington (DC), USA: ASM Press.

595 Uyttendaele, M., Rajkovic, A., Benos, G., François, K., Devlieghere, F., & Debevere, J. (2004).
596 Evaluation of a challenge testing protocol to assess the stability of ready-to-eat cooked meat
597 products against growth of *Listeria monocytogenes*. *International Journal of Food*
598 *Microbiology*, 90, 219–236.

599 Yang, J., Huang, K., Qin, S., Wu, X., Zhao, Z., & Chen, F. (2009). Antibacterial action of selenium-
600 enriched probiotics against pathogenic *Escherichia coli*. *Digestive Diseases and Sciences*,
601 54, 246–254.

602 Zwietering, M. H., Jongenburger, I., Rombouts, F. M., & Van't Riet, K. (1990). Modeling of the
603 bacterial growth curve. *Applied and Environmental Microbiology*, 56, 1875–1881.