



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Soilless system on peat reduce trace metals in urban-grown food: unexpected evidence for a soil origin of plant contamination

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Pennisi, G., Orsini, F., Gasperi, D., Mancarella, S., Sanoubar, R., VITTORI ANTISARI, L., et al. (2016). Soilless system on peat reduce trace metals in urban-grown food: unexpected evidence for a soil origin of plant contamination. *AGRONOMY FOR SUSTAINABLE DEVELOPMENT*, 36(4), 1-11 [10.1007/s13593-016-0391-9].

Availability:

This version is available at: <https://hdl.handle.net/11585/588409> since: 2017-05-17

Published:

DOI: <http://doi.org/10.1007/s13593-016-0391-9>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1 **Soilless system on peat reduce trace metals in urban grown food:**
2 **unexpected evidence for a soil origin of plant contamination.**

3
4 Authors:

5 **GIUSEPPINA PENNISI**

6
7 **FRANCESCO ORSINI ***

8
9 **DANIELA GASPERI**

10
11 **SILVIA MANCARELLA**

12
13 **RABAB SANOUBAR**

14
15 **LIVIA VITTORI ANTISARI**

16
17 **GILMO VIANELLO**

18
19 **GIORGIO GIANQUINTO**

20
21 Research Centre for the Urban Environment on Agriculture and Biodiversity, Department of Agricultural
22 Sciences, University of Bologna, Bologna, Italy

23
24 *Corresponding author:

25

This is the accepted manuscript of an article published in *Agronomy for Sustainable Development*.
The final authenticated version is available online at: <https://doi.org/10.1007/s13593-016-0391-9>

This version is subjected to Springer Nature terms for reuse that can be found at:
<https://www.springer.com/gp/open-access/authors-rights/aam-terms-v1>

26 Abstract

27 **Urban horticulture is increasingly popular for social and economic benefits. However, edible urban**
28 **crops may be contaminated by airborne pollutants, thus leading to serious health risks. Therefore, a**
29 **better understanding of contamination risks of urban cultivation is needed in order to define safe**
30 **practices. In particular, whereas it is commonly accepted that the contamination of urban grown food**
31 **comes from airborne pollutants, little is known on a possible contamination by soils. Here we studied**
32 **trace metal risk in horticultural crops grown in an experimental urban allotment garden in Bologna,**
33 **Italy. Seven experiments were conducted between June and November 2015 on tomato, sweet basil,**
34 **onion, lettuce, kale, bulb fennel and radish. Treatments included two growing systems, soil and**
35 **soilless, and two fertilization managements: mineral and organic. Trace metal concentrations were**
36 **measured in soils, substrates and edible plant tissues. We identified preferentially translocated metals**
37 **by partitioning analysis of tomato, sweet basil and kale. Results showed that crops grown on soilless**
38 **system have a lower metal content, of -70% for Cr, -61% for Cu, -45% for Cd and -81% for Ni,**
39 **compared with those grown on soil. This finding demonstrates that the major contamination risk is**
40 **urban area is unexpectedly related to soil pollution.**

41

42 **Keywords: urban gardening; soilless cultivation; food safety; heavy metals; plant nutrition.**

43

44 1 Introduction

45

46 Urban agriculture activities are commonly found in Europe and all over the world. These include the
47 production of crop and livestock goods within cities and towns ([Zeza and Tasciotti 2010](#)), but also peri-urban
48 agricultural areas which may provide product to the local population, such as vegetables, medicinal plants,
49 fruit trees, ornamental plants, milk, meat and wool ([Lin et al. 2015](#)). Urban allotment gardens are one element
50 of the urban green infrastructure that is becoming increasingly important in urban landscape planning. As
51 reported in [Breuste and Artmann \(2014\)](#), they combine utility, social meaning, beauty, and several ecosystem
52 services such as food supply ([Drescher 2004](#)), air filtering ([Davies et al. 2011](#)), urban temperature and climate
53 regulation ([Phelan et al. 2015](#)), noise reduction ([Aylor 1972](#)), runoff mitigation ([Zhang et al. 2012](#)), and
54 biodiversity development ([Lin et al. 2015](#)). However, plant cultivation within cities may present environmental
55 risks associated to both air and soil pollution ([Alloway 2004](#)). In urban areas, air pollutants generally derive
56 from artificial sources e.g. vehicular emissions and fossil fuel burning ([Agrawal et al. 2003](#)). Urban air
57 pollution has increased rapidly in the past decades with fast industrialization, rapid growth of urban population,
58 increase in vehicular traffic, badly maintained roads and human activities. Consequently, agricultural land
59 adjacent to urban areas may be exposed to air pollutants of urban origin, which may result in contamination of
60 horticultural products beyond precautionary values. When this happens, a dietary exposure to trace metals can
61 result in significant human health risk ([Massaquoi et al. 2015](#)). An interesting study ([Säumel et al. 2012](#)),
62 explored the relationship between local traffic burden and the trace metal concentration in the edible biomass

63 of different horticultural crops cultivated by gardeners in the inner city of Berlin, Germany, analysing the
64 influence of traffic burden and of the existence of barriers between cultivation sites and nearby streets. The
65 study shows that a higher overall traffic burden increases trace metal content in the crop biomass while the
66 presence of barriers between cultivation site and roads strongly reduces trace metal content. As reported in
67 [Vittori Antisari et al. \(2015\)](#), the concentration of trace metals in urban grown vegetables is strictly related to
68 the site in the city where plants are grown, with vegetables grown nearby main roads generally presenting
69 greatest pollution levels. Urban soils can also be contaminated as they are often located on old urban sites,
70 impacted by human activities, such as industrial activities, road traffic, waste dumps and demolition sites ([Jean-
71 Soro et al. 2015](#)). They may contain hazardous substances that can be assimilated by plants grown in these
72 spaces and become a danger for human health. Among major pollutants in urban soils, potential toxic elements
73 (As, Hg, Cd, Co, Cr, Cu, Mn, Mo, Ni, Pb, Sn, Zn and Se), can be accumulated in foodstuffs and enter into the
74 food chain, causing serious problems to human health. Most important sources of trace metals in urban soils
75 are atmospheric depositions, agricultural input (e.g. fertilizers, pesticides and manure), use of low quality
76 compost, industrial activities, refineries. Exposure to metals can occur through a variety of processes. For As
77 and Cd, intake of plant-derived food represents a major fraction of potentially health-threatening human
78 exposure ([Clemens and Ma 2016](#)).

79 In Europe, urban gardens are primarily managed by the elders using traditional management practices often
80 related to the wide use of synthetic products, mainly fertilizers, herbicides and pesticides ([Szolnoki et al.,
81 2013](#)). One of the main problems that are generally encountered is the contamination risk associated with weed
82 and pest control, generally managed by application of chemicals. In 2001, a study showed that about 90% of
83 the German allotment holders use pesticides ([UM 2014](#)). Pesticides kill pests, but they may have a negative
84 effect on the useful garden fauna (e.g. pollinators, pest predators), may cause water, air and soil pollution, as
85 well as contamination of the edible products. Consistently, in recent years, the use of pesticides is being
86 reduced or even completely banned in allotment gardens, and their application has been dramatically reduced
87 as gardener awareness has grown ([Barthel et al. 2010](#)). Soil fertilization, on the other hand, is generally applied
88 by mineral inorganic fertilizers, which are easy to apply and allow the supply of the right dose of nutrients.
89 However, cases of over-application and soil and water contamination by nitrates and phosphates have been
90 repeatedly reported in urban gardens, due to the low agricultural skills of hobby farmers ([Tixier and de Bon
91 2006](#)). However, the potentially high environmental impact related to the considerable amount of inputs needed
92 to support the production may be avoided when sustainable practices are adopted. Weed management may be
93 addressed by using organic mulching, therein also improving moisture retention and preservation of soil
94 fertility. Pest control may be addressed in different ways ([Voigt et al. 2016](#)), including selection of disease and
95 pest resistant plants, adoption of companion plantings, using trap crop plants that attract pest insects saving
96 the nearby main crop, application of natural insecticides (e.g. Neem oil and Pyrethrum), application of pests'
97 diseases (e.g. *Bacillus thuringiensis*), and integration in the garden of flowering plants that attract beneficial
98 insects (both pollinators and natural pest enemies). Organic fertilization (e.g. by using manure from livestock
99 or poultry, or compost from vegetable waste) has the function of preserving soil characteristics allowing the

100 proper crop development (Tixier and de Bon 2006). Organic fertilizers are characterized by slow and long-
101 lasting action, and have beneficial effects on soil microflora, soil texture and water holding capacity.
102 With the increase of world urban population, which in 2007 has surpassed the rural population, the surface of
103 the city grows and evolves often uncontrollably. This leads to a continuous increase in impervious surfaces
104 and the consequent consumption of arable soils. In such conditions, urban horticulture development follows a
105 growing trend, representing not only a source of food, but also becoming a possible source of income
106 (Caldeyro-Stajano 2004). The two main constraints to the spread of urban horticulture are the shortage of soil
107 and water. Soil in the cities may be contaminated by air and soil pollutants, therefore cultivation of edible
108 products may not be feasible and, in general, in the cities often availability of land is scarce (Orsini et al. 2013).
109 In addition, water availability constitutes a further problem in the urban environment, since in some areas there
110 can be difficult access to drinking water. The adoption of simplified soilless cultivation systems allows
111 overcoming these constraints (Tixier and de Bon 2006). Among the different soilless systems, most commonly
112 adopted in urban gardens are those that make use of solid substrates in containers generally built on wood and
113 waterproofed with a plastic film (Orsini et al. 2014). The aim of the hereby illustrated research is to propose
114 alternative/innovative agricultural techniques instead of those typically used by local gardeners (on soil
115 cultivation with mineral or organic fertilizers purchased in local shops). Consistently, the risk of trace metals
116 contamination in vegetables cultivated in an experimental urban allotment garden was investigated by
117 comparing different growing systems (soil and soilless) and fertilization managements (mineral and organic).

118

119 **2 Materials and Methods**

120

121 The experimental allotment garden was previously identified (Vittori Antisari et al. 2015) as the most
122 contaminated by trace metals in terms of both tissue accumulation and leaf deposition in comparison with other
123 allotment gardens situated in different areas of the city and nearby different sources of contamination. The
124 experimental allotment garden (Fig. 1, with a surface of 40 m²) is part of a large area of gardens (so-called
125 Orti Salgari), built in the eighties, which occupies a surface of 27,000 m² and is composed by 398 allotment
126 gardens. It is situated nearby a main road of the city of Bologna (via San Donato, 300 m from the garden, 10³–
127 10⁴ vehicles d⁻¹), the main motorway (A14, 1,200 m from the garden, 10⁵ vehicles d⁻¹), the railway (1,600 m
128 from the garden, 700 trains d⁻¹) and it is placed 10 m from the local street (coordinates 44° 30' 54" N, 11° 23'
129 29" E). Moreover, the city incinerator (Frullo, 600 t d⁻¹ of urban waste processed) is located 3,000 m away,
130 north-east of the garden, with prevailing winds in direction East.

131

132 **2.1 Plant material and experimental design**

133

134 Seven experiments were conducted between June and November 2015. Plant species used were tomato
135 (*Lycopersicon esculentum*, cv San Marzano), sweet basil (*Ocimum basilicum*, cv. Napoletano), onion (*Allium*
136 *cepa*, cv. Gialla di Stoccarda), lettuce (*Lactuca sativa*, cv. Four Seasons), kale (*Brassica oleracea* var.

137 *Acephala* or tuscan kale or black cabbage), Florence/Bulb fennel (*Foeniculum vulgare*, var. *Azoricum* cv.
138 Carmo F1), and radish (*Raphanus sativus*, cv. Cherry Bell). Plants density was 5 plants m⁻² for kale and tomato,
139 10 plants m⁻² for fennel, lettuce and basil, 25 plants m⁻² for radish, 35 plants m⁻² for onion.
140 Treatments included growing media (soil vs peat) and fertilization management (mineral vs organic).
141 Experimental design was a strip plot with 2 replicates with growing media on the main plot and fertilization
142 management in the elemental plot. Each elemental plot hosted at least 6 plants.

143

144 2.2 Growing conditions

145

146 Some species (basil, tomato and radish) were sown in plastic containers filled with peat at the experimental
147 greenhouse facilities of the Agricultural Sciences Department of the University of Bologna under controlled
148 environmental conditions (25°C and 50% relative humidity). As they reached adequate growth (June 5, for
149 basil and tomato and September 28 for radish, at respectively 50, 28 and 10 days after sowing), seedlings were
150 transplanted at the experimental allotment garden. Other species (lettuce, onion, kale, fennel) were purchased
151 from a local retailer of products for agriculture (normally used by local gardeners, Garden Cap, Cadriano, BO,
152 Italy) and transplanted at the experimental allotment garden (respectively on June 5 for onion, and on
153 September 18, for lettuce, kale and fennel). Irrigation was provided daily with a drip irrigation system,
154 supplying 9 and 6 L m⁻² d⁻¹, respectively in summer (June to September), and early fall (October). Later on in
155 the season (November), irrigation was not supplied.

156 Plants were grown either on the allotment soil or onto elevated beds filled with commercial soil (Geotec, RO,
157 Italy). Growing containers were built according to [Orsini et al. \(2014\)](#), using recycled pallets and black plastic
158 films. Declared physical and biochemical features of commercial soil were Organic Carbon 25%, Organic
159 Nitrogen 1%, peat 60%, salinity 0.6 dS m⁻¹.

160 No pest control treatments were applied during the crop cycle. Consistently, organic and conventional
161 management protocols were diversified only according to the plant nutrient management. Plants grown under
162 mineral fertilization management were supplied with an N-P-K (5-7-14) chemical fertilizer (Tris, produced by
163 ALFE, MN, Italy), in measure of 50 g m⁻². Organic crop management was obtained by adding 125 g m⁻² of
164 mature horse manure (Fertistall pellet, Agrilinea, FE, Italy), with the following declared features Nitrogen 2%,
165 Organic Carbon 24%. The different doses of fertilizer were calculated to give the same amount of Nitrogen in
166 the two treatments.

167

168 2.3 Chemical analyses

169

170 *Soil and peat analysis.* Prior to experimentation, after the applications of the different fertilizers, soil and peat
171 samples were collected and analysed as reported below. Samples were air-dried and sieved (<2 mm). Measures
172 of pH (pHmeter, Crison, Barcelona, Spain) were performed with distilled water on 1:2.5 w/v. Total organic
173 carbon was measured by Dumas combustion with a EA 1110 CHN elemental analyser (Thermo Fisher

174 Scientific, Waltham, MA, USA) after dissolution of carbonates with 2 M HCl, and the organic matter was
175 obtained using 1.72 factors. Soil particle size distribution was determined by the pipette method (Gee and
176 Bauder 1986) and total carbonates (CaCO₃) were quantified by a volumetric method, according to Dietrich-
177 Fruehing.

178 The metal contents was determined according to Vittori Antisari et al. (2013). Briefly, the soil (0.25 g) was
179 treated with aqua regia (2 ml HNO₃ 65% plus 6 ml suprapur HCl 37%, suprapur grade Carlo Erba, MI, Italy)
180 in a microwave oven and the metal concentrations were determined by ICP-OES. The analysis of each sample
181 was replicated three times and compared with analyses of the International Reference Materials (BCR 141)
182 and laboratory internal standards (MO and ML), which was run after every 10 samples to check changes in
183 sensitivity. Controls with only reagents were also determined.

184 *Vegetable samples collection and analysis.* At full maturity (at respectively 38, 76, 76, 69, 69, 69 and 60 days
185 after transplanting, for basil, tomato, onion, lettuce, kale, fennel and radish), samples of the different plant
186 organs (roots, bulbs, stems, leaves, fruits) were collected for biochemical determinations. At harvest, fresh
187 weights were recorded and dry weights were determined after washing in tap water and drying at 58°C per 72
188 hours. Samples were dried in ventilated oven (T<40 °C) and ground in a blender with blades made of pure
189 titanium, carefully avoiding to introduce any further metal contamination to the samples (Vittori Antisari et al.
190 2012). Briefly, approximately 0.25 g of leaf sub-sample, weighted in Teflon bombs, was dissolved in 8 mL of
191 HNO₃ (suprapure, Merck, Kenilworth, New Jersey, USA) plus 2 mL of H₂O₂ (Carlo Erba, MI, Italy) using a
192 microwave oven (Milestone 2100, Sorisone, BG, Italy). After cooling, solutions were made up to 20 mL with
193 Milli-Q water and then filtered with Whatmann 42 filter paper. The major and trace elements were determined
194 by inductive coupled plasma optical emission spectrometry (ICP-OES, Spectro Ametek, MI, Italy). The ICP-
195 OES setting followed multi-standard solutions (CPI International, Amsterdam, The Netherland) that reproduce
196 the matrix effect present in samples and allow the lowering of detection limits. Instrument response was
197 assessed by measuring a standard sample (CRM 482 - Community Bureau of Reference, BCR).

198 *Statistical analysis.* The experimental data were statistically analysed using two ways analysis of variance
199 (ANOVA) and means of field replicates (n=6) were compared using Least Significant Difference (LSD) test.

200

201 2.4 Contamination indexes

202

203 *BioAccumulation Factor.* For the different trace metals, the BioAccumulation Factor (BAF) was calculated by
204 (eq. 1):

$$205 \text{BAF} = C_{\text{shoot}}/C_{\text{soil}} \quad (\text{eq. 1})$$

206 where C_{shoot} and C_{soil} are metals concentration in the plant edible portion of the shoot (mg kg⁻¹ dry weight) and
207 soil (mg kg⁻¹ dry weight), respectively (Ma et al. 2001; Cluis 2004).

208 *Translocation Factor.* Trace metals translocation from root to shoot was measured by Translocation Factor (TF)
209 calculated as described below (eq. 2).

$$210 \text{TF} = C_{\text{shoot}}/C_{\text{root}} \quad (\text{eq. 2})$$

211 where C_{shoot} and C_{root} are metals concentration in the edible part of the shoot ($mg\ kg^{-1}$ dry weight) and the one
212 of the root of plant ($mg\ kg^{-1}$ dry weight), respectively. Whenever C_{shoot} is equal or greater than C_{root} ($TF>1$),
213 metals are effectively translocated from root to the shoot, whereas root compartmentation occurs whenever
214 C_{root} is greater than C_{shoot} (Baker and Brooks 1989; Zhang et al. 2002; Fayiga and Ma 2006).

215 *Daily Metal Intake and Health Risk Index.* The Daily Metal Intake and Health Risk Index are indexes relating
216 to daily estimated consumption as well as health risks from the consumption of contaminated food. The US
217 Environmental Protection Agency's reference doses (US-EPA IRIS 2006) were used as reference points. The
218 Daily Metal Intake (DMI) was estimated using the eq. 3:

$$219 \quad DMI = C_{shoot} * 0.085 * DPC / BW \quad (eq. 3)$$

220 where C_{shoot} is the concentration of metals in the edible part of the shoot of the plant (as $mg\ kg^{-1}$ dry weight),
221 multiplied by a conversion factor of 0.085 to convert dry weight vegetable metal content to fresh weight,
222 according to Rattan et al. (2005). Daily plant consumption (DPC) and body weight (BW) were estimated based
223 on Leclercq et al. (2009).

224 Based on the Daily Metal Intake, it was possible to calculate the Health Risk Index, based on eq. 4:

$$225 \quad HRI = DMI/RfD \quad (eq.4)$$

226 where RfD is the Reference Dose, which, for the studied elements is respectively 1.5 (Cr), 0.04 (Cu), 0.004
227 (Pb) and 0.02 (Ni) ($mg\ kg^{-1}\ BW\ d^{-1}$) (US-EPA IRIS 2006; Jan et al. 2010). HRI values >1 are considered to
228 pose health risks (Cui et al. 2004; Rattan et al. 2005).

229

230 **3 Results and discussion**

231

232 **3.1 Growing media characteristics and metal concentrations**

233

234 Soil pH measured in the top 20 cm layer was significantly ($p<0.001$) higher than that of peat (**Table 1**).
235 Fertilization management did not influence pH of soil samples, whereas pH of peat samples were statistically
236 different in the two diverse fertilization methods with higher values on samples treated with mineral fertilizer
237 as compared to the organic ones (**Table 1**). The importance of this parameter is related to the influence that it
238 has on all the reactions that occur in nature. The pH values of the soil encountered in the present study were in
239 line with the regional average values, that, in Emilia Romagna soils, present pH values between 7 and 8.5,
240 classified as sub-alkaline soils (ARPA-RER 2004). Average $CaCO_3$ total content in soil samples was
241 significantly ($p<0.01$) higher as compared to peat samples (+53%), which was possibly correlated with the
242 differences in pH (Rowell 2014). On the other hand, organic matter was ten times higher in peat samples as
243 compared to soil samples (**Table 1**). Both the values of total carbonates and organic carbon (**Table 1**) measured
244 for soil samples were in line with the average values of the regional soils. No statistical differences in macro-
245 element content was observed between organic and mineral samples (**Table 1**). Significantly ($p<0.001$) higher
246 concentrations of Al, B, Ba, Fe, K, Mn were observed in soil samples; on the contrary, concentrations of Ca,
247 Na, S, and Si were significantly higher in peat samples (**Table 1**). No significant differences in Mg, P, Sr and

248 Ti content were observed between soil and peat samples. Trace metal contents, however, widely varied
249 between soil and peat samples. For every sample, the concentration of Cd was below the detection limit, and,
250 only for soil samples Mo concentration was below the detection limit. Significant ($p < 0.001$) differences
251 between soil and peat samples were revealed for every of the other microelements (As, Be, Co, Cr, Cu, Li,
252 Mo, Ni, Pb, Sb, Sn, V, Zn), with greater values in soil samples (**Table 1**). Comparing average soil
253 microelements concentrations with results of other researches in both urban and rural environments, the first
254 consideration is that trace metal concentration in soils may vary substantially according to both geogenic and
255 anthropic elements. As compared with other studies (Luo et al. 2011; Szolnoki et al. 2013; Izquierdo et al.
256 2015), the soil Cr content hereby measured (91 mg kg^{-1} , **Table 1**) turned out to be higher than previous
257 evidences in urban gardens where it ranged 12 to 31 mg kg^{-1} . Alternatively, Cu, Pb and Ni (respectively
258 accounting for 56, 44 and 46 mg kg^{-1} , **Table 1**) were within previously observed concentrations (Kabala et al.
259 2009; Luo et al. 2011; Szolnoki et al. 2013; Jean-Soro et al. 2014; Izquierdo et al. 2015) in both urban and
260 rural environments.

261

262 3.2 Trace metals content in vegetables' edible portions

263

264 Although potential health risks are generally associated with the assimilation of trace metals, there is a lack in
265 legislation concerning threshold limits for most trace metals in marketed crops. The only available EU
266 legislation (EU 2009) regulates upper limits of Pb and Cd in edible products. Consistently, Pb shall always be
267 under 0.1, 0.3, and 0.2 mg kg^{-1} of fresh weight (respectively in legumes, brassica, and all other vegetables),
268 which are about 1.2, 2.4 and 3.6 mg kg^{-1} on a dry weight basis. Furthermore, Cd threshold limits are set at
269 0.05, 0.1, and 0.2 mg kg^{-1} of fresh weight, respectively in vegetables whose edible part is the fruit, the
270 stem/root, or the leaf (EU 2009). Most dangerous elements beyond Pb and Cd are Cr, Cu, As, and Ni (Vittori
271 Antisari et al. 2015). Within the experiments hereby described, some of these elements (namely Cd and As)
272 were not detectable in vegetable samples. On the other hand, significant accumulation levels of Cr, Cu, Pb,
273 and Ni were found, as shown in **Fig. 2** and **3**. The recovery intervals for these elements, based on the CRM
274 482, were respectively 3.99 ± 0.51 (Cr), 7.03 ± 0.19 (Cu), 2.19 ± 0.49 (Ni) and 37.8 ± 1.47 (Pb). Given that no
275 significant interaction between growing system and fertilization management on the edible product content of
276 those trace metal was observed, the effects of these two factors will be separately described for each element.
277 *Chromium*. Cr content in edible organs of radish, fennel, lettuce and basil was significantly higher in plants
278 cultivated on soil as compared to those grown on peat, with greatest values in radish and lettuce (respectively
279 85 and 29 mg kg^{-1}) (**Fig. 2a**). Contrariwise, Cr accumulation in onion bulbs was higher in plants cultivated on
280 peat. Finally, kale leaves and tomato fruit did not present significant differences between the two growing
281 systems (**Fig. 2a**). Based on the fertilization managements (**Fig. 3a**), Cr accumulation was higher in samples
282 of radish, onion, lettuce and tomato supplied with organic fertilizer. Highest concentrations were observed in
283 radish and lettuce, respectively reaching 76 and 26 mg kg^{-1} . No significant differences could be associated
284 with fertilization management in fennel, basil or kale (**Fig. 3a**).

285 *Copper.* Cu content in edible organs of radish, onion, fennel, lettuce, basil and tomato was significantly higher
286 in plants cultivated on soil as compared to those grown on peat, with greatest values in radish and basil
287 (respectively 19 and 17 mg kg⁻¹) (**Fig. 2b**). No statistically significant differences could be observed in leaves
288 of soil- and peat- grown plants of kale, which also showed the lowest concentration (5 mg kg⁻¹) among studied
289 species grown on soil (**Fig. 2b**). Considering fertilization management, significant differences in Cu
290 accumulation were detectable only in onion bulbs and tomato fruits, with a slightly higher (respectively +4
291 and +1 %) concentration in plants grown with mineral fertilizer as compared to same products supplied with
292 organic fertilization (**Fig. 3b**). Highest concentrations were observed in radish and lettuce (14 mg kg⁻¹) (**Fig.**
293 **3b**).

294 *Lead.* Pb content in edible parts of radish, fennel, lettuce and basil was significantly higher in plants cultivated
295 on soil than those cultivated on peat (**Fig. 2c**), with values above EU safety thresholds only observed in radish.
296 Contrariwise, onion showed opposite results, with Pb accumulation being greater in onion cultivated on peat
297 as compared to same plants grown on soil (**Fig. 2c**). In general, radish and lettuce had the higher accumulation
298 level (respectively 7 and 3 mg kg⁻¹). Kale and tomato fruit did not present significant differences between the
299 two growing systems. Fertilization management did not affect Pb content in edible tissues of the studied
300 species, with exclusion of lettuce, where a significant 2-fold increase in Pb content was associated with organic
301 fertilization regime (**Fig. 3c**).

302 *Nickel.* Ni content in edible organs of radish, fennel, lettuce, basil and kale was significantly higher in plants
303 cultivated on soil as compared to those grown on peat, with greatest values in radish and lettuce (respectively
304 7 and 3 mg kg⁻¹) (**Fig. 2d**). Independently from the growing system, lowest accumulations were observed in
305 onion (0.10 mg kg⁻¹), and tomato (0 mg kg⁻¹) (**Fig. 2d**). Fertilization management did not affect Ni content in
306 edible tissues of the studied species, with exclusion of radish, where a 2.5-fold increase in Ni content was
307 associated with mineral fertilization regime (**Fig. 3d**).

308 In order to understand the potential health risk of the vegetables grown in the allotment garden, a comparative
309 analysis against values obtained in previous similar studies ([Douay et al. 2013](#); [Säumel et al. 2012](#); [Warming
310 et al. 2015](#)) was performed. Accordingly, and despite the apparently high content in soils, Cr values were
311 relatively low (with exclusion of radish that most accumulated Cr, with values of 85 mg kg⁻¹, **Fig. 2**). On the
312 other hand, Cu and Ni values were generally higher than previous experiences and Pb values varied
313 dramatically across species (with greatest values again on radish, about 7 mg kg⁻¹, **Fig. 2**). A previous study
314 ([Säumel et al. 2012](#)) addressed the quantification of trace metal contaminants in allotment gardens and food
315 purchased at local markets in the city of Berlin, showing lower values of Cu and Ni as compared to the hereby
316 presented results. This suggests that crops able to better exclude or compartmentalize those two elements
317 should be preferred at the experimental site.

318 When mean values of all studied vegetable species were compared, the adoption of simplified soilless systems
319 filled with non-contaminated growing substrate enabled to reduce trace metal contamination risk associated
320 with Cr (-70%, 6 vs 20 mg kg⁻¹, in peat vs soil, respectively), Cu (-61%, 7 vs 13 mg kg⁻¹), Pb (-45%, 0.8 vs 2.0
321 mg kg⁻¹) and Ni (-81%, 0.3 vs 2 mg kg⁻¹). Apparently, the greater values of contaminants associated with on-

322 soil cultivation should be related to the growing media content (**Table 1**). Furthermore, the observed
323 differences found in the pH values of soil and peat (**Table 1**) may have influenced the toxic mineral uptake,
324 given that substrate pH affect differently the solubility and availability of the elements, as commonly known
325 (e.g. [Kumpiene et al. 2008](#); [Zeng et al. 2011](#)).

326 Organic fertilization strategies did not affect substantially biochemical features and trace metal concentrations
327 of the growing media, as illustrated in **Table 1**. In the present study, however, some crop-specific differences
328 in trace metal accumulation in edible tissue were evidenced, with a general raise in Cr in radish and in Cr and
329 Pb in lettuce grown under organic fertilization. On the other hand, mineral fertilization resulted in greater Ni
330 in radish and Cu in onion and tomato. Comparative studies of organic versus mineral fertilization have rarely
331 addressed the effect on trace metal accumulation ([Chu and Wong 1987](#); [Singh et al. 2010](#); [Zaccone et al. 2010](#);
332 [Christensen and Elsgaard 2014](#)), often with controversial results ([Liñero et al. 2015](#)). Consistently, a priority
333 of the research is to address the understanding of how organic matter may affect (e.g. by chelating minerals)
334 plant uptake of contaminants.

335

336 3.3 Trace metal contamination indexes

337

338 In order to identify the crop capacity to exclude dangerous trace metals, the BioAccumulation Factor was used
339 to understand which species were most resilient to ion distribution in the growing media ([Rezvani and
340 Zaefarian 2011](#)). Consistently, crops may be classified as hyperaccumulators, accumulators and excluders,
341 based on the value of their BioAccumulation Factor being above, equal or below 1 unit. Accordingly, all
342 studied species acted as excluders, although to a different extent (**Table 2**). While highest values were
343 associated with radish roots, the lowest ones were observed for Cr and Ni in edible organs of tomato,
344 confirming the plant capability to avoid toxic elements loading into fruits ([Ali and Al-Qahtani 2012](#)). Once a
345 toxic element enters in plant by the root, it can either be translocated to shoots or compartmentalized within
346 roots. An indicator of the plant ability to avoid translocation of such elements into shoot is provided by the
347 Translocation Factor ([Rezvani and Zaefarian 2011](#)). The higher its value, the greater the rate of element that is
348 transferred to shoots. Among studied crops, three species (kale, sweet basil and tomato) representing three
349 different plant families (respectively Brassicaceae, Lamiaceae and Solanaceae) were studied for their trace
350 metal compartmentation in different plant organs (**Table 2**). For the sake of this publication, Translocation
351 Factor was calculated by relating the element concentration in the edible plant organ (leaf or fruit) to the
352 concentration of the same element in roots. Accordingly, no differences were observed on Cr, whereas greatest
353 values of Translocation Factor for Cu were observed in sweet basil and for Pb and Ni in kale, confirming the
354 hypothesis that lowest translocation occurs to fruits as compared to leaves in plants ([Tiwari et al. 2011](#)).
355 Consistently, a study over 15 plant native species grown on trace metal contaminated sites of Florida, showed
356 that both Pb and Cu absorbed at root level were only partially translocated to shoots ([Yoon et al. 2006](#)).

357 In order to better address the comprehension of the potential health risks associated with the consumption of
358 allotment garden vegetables, the application of Daily Metal Intake and Health Risk Index was performed on

359 the vegetables grown on the garden soil. Accordingly, values of Daily Metal Intake were extremely low
360 (ranging 0 to 20 $\mu\text{g kg}^{-1}$ Body Weight d^{-1} , data not shown) and this resulted in limited Health Risk Index values.
361 Health Risk Index was greatest in radish for Cr, Cu and Pb (**Table 2**), reaching respectively 0.015, 0.13 and
362 0.45 in adults (70 kg body weight and 0.22 kg d^{-1} Daily Plant Consumption, [Leclercq et al. 2009](#)) and 0.032,
363 0.270 and 0.951 in children (26 kg Body Weight and 0.13 kg d^{-1} Daily Plant Consumption, [Leclercq et al.](#)
364 [2009](#)) (**Table 2**). The same index was not affected by the crop species in neither adults (0.026) nor children
365 (0.055) when Ni was considered (**Table 2**). Health Risk Index estimated on both adults and children resulted
366 to be below the toxicity level of 1 ([US-EPA IRIS 2006](#)), although index value associated with Pb was greatest
367 in radish, confirming its associated health risks when grown in contaminated environments.

368

369 **4 Conclusions**

370

371 The experimental garden presented moderately high levels of Cr, whereas Cu, Pb and Ni were within ranges
372 observed in both urban and rural vegetable gardens assessed in previous studies. Indeed, while Cr content in
373 edible organs were lower than values observed in previous studies, Ni and Cu were higher than previous
374 evidences. Furthermore, Pb concentration in edible radish roots resulted to be above risk thresholds indicated
375 by the EU. Nonetheless, for all studied trace metals, Daily Metal Intake and Health Risk Index of plants grown
376 on soil were below the threshold value for health risks. Among studied plants, radish most accumulated toxic
377 elements, which resulted in significantly greater risk from all other studied crops, reaching values close to
378 toxicity for Pb when Health Risk Index was associated with children consumption. All studied crops presented
379 BioAccumulation Factor values always below 1, although with highest values for Cr, Cu, Pb and Ni in radish.
380 Translocation Factor of Cu from root to shoot was greater in leaves of sweet basil, while of Pb and Ni in kale,
381 as compared to the translocation to fruits in tomato. Plants grown on peat in a simplified soilless systems
382 showed reduced content of Cr, Cu, Pb and Ni as compared to those grown on soil, due to differences in the
383 growing media in terms of concentration of the element and biochemical properties (pH and CaCO_3). Organic
384 fertilization resulted in increased Cr and Pb and reduce Cu and Ni in the edible organs of some of the studied
385 crops. According to the results of the present study, the adoption of soilless growing systems where plants are
386 grown on peat may be a feasible strategy to cope with trace metal contamination risk in urban allotment garden.
387 The absence of clear differences in trace metal accumulation as affected by either mineral or organic fertilizers,
388 on the other hand, suggests that deeper studies should be conducted in order to define proper fertilization
389 strategies in potentially polluted environments.

390 **Acknowledgments**

391 The research was partially funded by the Erasmus+ project Urban Green Train (Urban Green Education for
392 Enterprising Agricultural Innovation). English language was checked by Dr. Agata Pennisi Sacoor.

393

394 **References**

- 395 Agrawal M, Singh B, Rajput M, Marshall F, Bell JNB (2003) Effect of air pollution on peri-urban agriculture:
396 a case study. *Environ Pollut* 126:323-329. doi: 10.1016/S0269-7491(03)00245-8
- 397 Ali MH, Al-Qahtani KM (2012) Assessment of some heavy metals in vegetables, cereals and fruits in Saudi
398 Arabian markets. *Egypt J Aquat Res* 38:31-37. doi:10.1016/j.ejar.2012.08.002
- 399 Alloway BJ (2004) Contamination of soils in domestic gardens and allotments: a brief overview. *Land Contam*
400 *Reclam* 12:179-187. doi:10.2462/09670513.658
- 401 ARPA-RER Regione Emilia Romagna (2004) Relazione sullo stato dell'ambiente della regione Emilia
402 Romagna.
- 403 Aylor DE (1972) Noise reduction by vegetation and ground. *J Acoust Soc Am* 51:197-205.
404 doi:10.1121/1.1912830
- 405 Baker AJM, Brooks RR (1989) Terrestrial higher plants which hyperaccumulate metallic elements. A review
406 of their distribution, ecology and phytochemistry. *Biorecov* 1:81-126
- 407 Barthel S, Folke C, Colding J (2010) Social-ecological memory in urban gardens - Retaining the capacity for
408 management of ecosystem services. *Glob Environ Change* 20:255-265.
409 doi:10.1016/j.gloenvcha.2010.01.001
- 410 Breuste JH, Artmann M (2014) Allotment gardens contribute to urban ecosystem service: case study Salzburg,
411 Austria. *J Urban Plann Dev* 141:A5014005. doi:10.1061/(ASCE)UP.1943-5444.0000291
- 412 Caldeyro-Stajano M (2004) Simplified hydroponics as an appropriate technology to implement food security
413 in urban agriculture. *Pract Hydroponics Greenh* 76:1-6
- 414 Christensen BT, Elsgaard L (2014) Comparing heavy metal contents in crops receiving mineral fertilisers and
415 animal manure. *Int Fertil Soc: Proc* 761
- 416 Chu LM, Wong MH (1987) Heavy metal contents of vegetable crops treated with refuse compost and sewage
417 sludge. *Plant Soil* 103:191-197. doi:10.1007/BF02370388
- 418 Clemens S, Ma JF (2016) Toxic heavy metal and metalloid accumulation in crop plants and foods. *Annu Rev*
419 *Plant Biol* 67:12.1-12.24. doi:10.1146/annurev-arplant-043015-112301
- 420 Cluis C (2004) Junk-greedy greens: phytoremediation as a new option for soil decontamination. *BioTech J*
421 2:60-67
- 422 Cui YJ, Zhu YG, Zhai RH, Chen DY, Huang YZ, Qiu Y, Liang JZ (2004) Transfer of metals from soil to
423 vegetables in an area near a smelter in Nanning, China. *Environ Intern* 30:785-791.
424 doi:10.1016/j.envint.2004.01.003

425 Davies ZG, Edmondson JL, Heinemeyer A, Leake JR, Gaston KJ (2011) Mapping an urban ecosystem service:
426 quantifying above-ground carbon storage at a city-wide scale. *J App Ecol* 48:1125-1134.
427 doi:10.1111/j.1365-2664.2011.02021.x

428 Douay F, Pelfrène A, Planque J, Fourrier H, Richard A, Roussel H, Girondelot B (2013) Assessment of
429 potential health risk for inhabitants living near a former lead smelter. Part 1: metal concentrations in soils,
430 agricultural crops, and homegrown vegetables. *Environ Monit Assess* 185:3665-3680. doi:10.1007/s10661-
431 012-2818-3

432 Drescher AW (2004) Food for the cities: urban agriculture in developing countries. *Acta Horticult* 643:227-
433 231. doi:10.17660/ActaHortic.2004.643.29

434 EU (2009). Commission Regulation (EU) No 420/2011 of 29 April 2011 amending Regulation (EC) No
435 1881/2006 setting maximum levels for certain contaminants in foodstuffs. [http://eur-](http://eur-lex.europa.eu/legalcontent/EN/TXT/HTML/?uri=CELEX:32011R0420&from=EN)
436 [lex.europa.eu/legalcontent/](http://eur-lex.europa.eu/legalcontent/EN/TXT/HTML/?uri=CELEX:32011R0420&from=EN) EN/TXT/HTML/?uri=CELEX:32011R0420&from=EN. Accessed 5 March
437 2016

438 Fayiga AQ, Ma LQ (2006) Using phosphate rock to immobilize metals in soils and increase arsenic uptake in
439 *Pteris vittata*. *Sci Total Environ* 359:17-25. doi:10.1016/j.scitotenv.2005.06.001

440 Gee GW, Bauder JW (1986) Particle-size analysis. In: Klute A (ed) *Methods of soil analysis, Part 1. Physical*
441 *and mineralogical methods*. Am Soc Agron, Madison, pp 383-411

442 Izquierdo M, De Miguel E, Ortega MF, Mingot J (2015) Bioaccessibility of metals and human health risk
443 assessment in community urban gardens. *Chemosphere* 135:312-318.
444 doi:10.1016/j.chemosphere.2015.04.079

445 Jan FA, Ishaq M, Khan S, Ihsanullah I, Ahmad I, Shakirullah M (2010) A comparative study of human health
446 risks via consumption of food crops grown on wastewater irrigated soil (Peshawar) and relatively clean
447 water irrigated soil (lower Dir). *J Hazard Mater* 179:612-621. doi:10.1016/j.jhazmat.2010.03.047

448 Jean-Soro L, Le Guern C, Bechet B, Lebeau T, Ringeard MF (2015) Origin of trace elements in an urban
449 garden in Nantes, France. *J Soils Sediments* 15:1802-1812. doi:10.1007/s11368-014-0952-y

450 Kabala C, Chodak T, Szerszen L, Karczewska A, Szopka K, Fratzak U (2009) Factors influencing the
451 concentration of heavy metals in soils of allotment gardens in the city of Wroclaw, Poland. *Fresen Environ*
452 *Bull* 118:1118-1124

453 Kumpiene J, Lagerkvist A, Maurice C (2008) Stabilization of As, Cr, Cu, Pb and Zn in soil using amendments-
454 a review. *Waste Manag* 28:215-225. doi:10.1016/j.wasman.2006.12.012

455 Leclercq C, Arcella D, Piccinelli R, Sette S, Le Donne C, Turrini A (2009) The Italian national food
456 consumption survey INRANSCAI 2005–2006: main results in terms of food consumption. *Public Health*
457 *Nutr* 12:2504-2532. doi:10.1017/S1368980009005035

458 Lin BB, Philpott SM, Jha S (2015) The Future of urban agriculture and biodiversity-ecosystem services:
459 Challenges and next steps. *Basic App Ecol* 16:189-201. doi:10.1016/j.baae.2015.01.005

460 Liñero O, Ciudad M, Carrero JA, Nguyen C, de Diego A (2015) Accumulation and translocation of essential
461 and nonessential elements by tomato plants (*Solanum lycopersicum*) cultivated in open-air plots under

462 organic or conventional farming techniques. *J Agric Food Chem* 63:9461-9470.
463 doi:10.1021/acs.jafc.5b03878

464 Luo C, Liu C, Wang Y, Liu X, Li F, Zhang G, Li X (2011) Heavy metal contamination in soils and vegetables
465 near an e-waste processing site, south China. *J Hazard Mater* 186:481-490.
466 doi:10.1016/j.jhazmat.2010.11.024

467 Ma LQ, Komar KM, Tu C, Zhang W, Cai Y, Kenelly ED (2001) A fern that hyper accumulates arsenic. *Nature*
468 409:579-582. doi:10.1038/35054664

469 Massaquoi LD, Ma H, Liu XH, Han PY, Zuo SM, Hua ZX, Liu DW (2015) Heavy metal accumulation in soils,
470 plants, and hair samples: An assessment of heavy metal exposure risks from the consumption of vegetables
471 grown on soils previously irrigated with wastewater. *Environ Sci Pollut Res* 22:18456-18468.
472 doi:10.1007/s11356-015-5131-1

473 Orsini F, Kahane R, Nono-Womdim R, Gianquinto G (2013) Urban agriculture in the developing world: a
474 review. *Agron Sustain Dev* 33:695-720. doi:10.1007/s13593-013-0143-z

475 Orsini F, Gasperi D, Marchetti L, Piovene C, Draghetti S, Ramazzotti S, Bazzocchi GG, Gianquinto G (2014)
476 Exploring the production capacity of rooftop gardens (RTGs) in urban agriculture: the potential impact on
477 food and nutrition security, biodiversity and other ecosystem services in the city of Bologna. *Food Sec*
478 6:781-792. doi:10.1007/s12571-014-0389-6

479 Phelan PE, Kaloush K, Miner M, Golden J, Phelan B, Silva III H, Taylor RA (2015) Urban Heat Island:
480 Mechanisms, Implications, and Possible Remedies. *Ann Rev Environ Resour* 40:285-307.
481 doi:10.1146/annurev-environ-102014-021155

482 Rattan RK, Datta SP, Chhonkar PK, Suribabu K, Singh AK (2005) Longterm impact of irrigation with sewage
483 effluents on heavy metal content in soils, crops and groundwater-a case study. *Agric, Ecosyst Environ*
484 109:310-322. doi:10.1016/j.agee.2005.02.025

485 Rezvani M, Zaefarian F (2011) Bioaccumulation and translocation factors of cadmium and lead in '*Aeluropus*
486 *littoralis*'. *Aust J Agric Eng* 2:114.

487 Rowell DL (2014) *Soil science: Methods and applications*. Taylor and Francis, Routledge, London, UK.

488 Säumel I, Kotsyuk I, Hölscher M, Lenkerei C, Weber F, Kowarik I (2012) How healthy is urban horticulture
489 in high traffic areas? Trace metal concentrations in vegetable crops from plantings within inner city
490 neighbourhoods in Berlin, Germany. *Environ Poll* 165:124-132. doi:10.1016/j.envpol.2012.02.019

491 Singh A, Agrawal M, Marshall FM (2010) The role of organic vs. inorganic fertilizers in reducing
492 phytoavailability of heavy metals in a wastewater-irrigated area. *Ecol Eng* 36:1733-1740.
493 doi:10.1016/j.ecoleng.2010.07.021

494 Szolnoki Z, Farsang A, Puskas I (2013) Cumulative impacts of human activities on urban garden soils: origin
495 and accumulation of metals. *Environ Poll* 177:106-115. doi:10.1016/j.envpol.2013.02.007

496 Tiwari KK, Singh NK, Patel MP, Tiwari MR, Rai UN (2011) Metal contamination of soil and translocation in
497 vegetables growing under industrial wastewater irrigated agricultural field of Vadodara, Gujarat, India.
498 *Ecotoxicol Environ Saf* 74:1670-1677. doi:10.1016/j.ecoenv.2011.04.029

499 Tixier P, de Bon H (2006) Urban horticulture. Cities farming for the future, urban agriculture for green and
500 productive cities. RUAF Foundation, IDRC and IIRR. ETC.-Urban Agriculture, Leusden
501 UM 2014 (=Umweltinstitut München): Pestizide (Pesticides) .URL:
502 <http://www.umweltinstitut.org/themen/landwirtschaft/pestizide/glyphosat/haus-und-kleingarten.html>
503 (accessed 24 February 2016)
504 US-EPA IRIS (2006) United States, Environmental Protection Agency, Integrated Risk Information System.
505 <http://www.epa.gov/iris/substS>
506 Vittori Antisari L, Carbone S, Ferronato C, Simoni A, Vianello G (2012) Leaf washing as an assessment tool
507 to characterize dry atmospheric deposition. *Int J Environ Qual* 9:37-50. doi:10.6092/issn.2281-4485/3737
508 Vittori Antisari L, Ventura F, Simoni A, Piana S, Rossi Pisa P, Vianello G (2013) Assesment of pollutants in
509 wet and dry deposition in a suburban area around a wast-to-energy plants (WEP) in Norther Italy. *J Environ*
510 *Prot* 4:16-25. doi:10.4236/jep.2013.45A003
511 Vittori Antisari L, Orsini F, Marchetti L, Vianello G, Gianquinto G (2015) Heavy metal accumulation in
512 vegetables grown in urban gardens. *Agron Sustain Dev* 35:1139-1147. doi:10.1007/s13593-015-0308-z
513 Voigt A, Leitao TE, Hursthouse AS, Jokinen A, Heller A, Latkowska M, Brito da Luz P, Christ Y, Bechet B,
514 Kulvik M, Langemeyer J (2016) Lessons Learned: Indicators and Good Practice for an Environmentally
515 Friendly Urban Garden. In “Urban Allotment Gardens in Europe” (Eds. Bell, S., Fox-Kamper R, Keshavarz
516 N, Drilling M, Benson M, Voigt A, Caputo S). Taylor and Francis, Routledge, London, UK
517 Warming M, Hansen MG, Holm PE, Magid J, Hansen TH, Trapp S (2015) Does intake of trace elements
518 through urban gardening in Copenhagen pose a risk to human health? *Environ Poll* 202:17-23.
519 doi:10.1016/j.envpol.2015.03.011
520 Yoon J, Cao X, Zhou Q, Ma LQ (2006) Accumulation of Pb, Cu, and Zn in native plants growing on a
521 contaminated Florida site. *Sci Total Environ* 368:456-464. doi:10.1016/j.scitotenv.2006.01.016
522 Zaccone C, Di Caterina R, Rotunno T, Quinto M (2010) Soil-farming system-food-health: effect of
523 conventional and organic fertilizers on heavy metal (Cd, Cr, Cu, Ni, Pb, Zn) content in semolina samples.
524 *Soil Tillage Res* 107:97-105. doi:10.1016/j.still.2010.02.004
525 Zeng F, Ali S, Zhang H, Ouyang Y, Qiu B, Wu F, Zhang G (2011) The influence of pH and organic matter
526 content in paddy soil on heavy metal availability and their uptake by rice plants. *Environ Poll* 159:84-91.
527 doi:10.1016/j.envpol.2010.09.019
528 Zezza A, Tasciotti L (2010) Urban agriculture, poverty, and food security: Empirical evidence from a sample
529 of developing countries. *Food Policy* 35:265-273. doi:10.1016/j.foodpol.2010.04.007
530 Zhang B, Xie G, Zhang C, Zhang J (2012) The economic benefits of rainwater-runoff reduction by urban green
531 spaces: A case study in Beijing, China. *J Environ Manag* 100:65-71. doi:10.1016/j.jenvman.2012.01.015
532 Zhang WH, Cai Y, Tu C, Ma QL (2002) Arsenic speciation and distribution in an arsenic hyperaccumulating
533 plant. *Sci Total Environ* 300:167-177. doi:10.1016/S0048-9697(02)00165-1

535 **Table 1.** Main soil features (pH, CaCO₃, Organic matter and Organic C) and concentrations of macro-
 536 elements (expressed as g kg⁻¹ dry weight) and micro-elements (expressed as mg kg⁻¹ dry weight) in soil and
 537 peat samples added with organic fertilizer or mineral fertilizer. Different letters indicate significant
 538 differences within row at P≤0.05; ns indicates non significant differences at P≤0.05 (n=6).

539

		Peat				Soil			
		Organic		Mineral		Organic		Mineral	
pH (H₂O)	-	6.72	c	6.88	b	8.11	a	7.97	a
CaCO₃	g kg ⁻¹	35.15	b	39.95	b	59.9	a	55.05	a
Organic Matter	%	55.72	a	56.35	a	4.89	b	4.77	b
Organic C	%	22.27	a	22.67	a	1.96	b	1.90	b
Al	g kg ⁻¹	8.27	b	6.89	b	32.78	a	31.09	a
B	g kg ⁻¹	0.02	b	0.02	b	0.03	a	0.03	a
Ba	g kg ⁻¹	0.07	b	0.07	b	0.20	a	0.19	a
Ca	g kg ⁻¹	29.34	a	27.42	a	22.76	b	23.29	b
Fe	g kg ⁻¹	8.28	b	7.27	b	24.47	a	24.32	a
K	g kg ⁻¹	6.04	b	5.02	b	7.55	a	7.01	a
Mg	g kg ⁻¹	7.52	ns	6.70	ns	7.01	ns	7.04	ns
Mn	g kg ⁻¹	0.25	b	0.21	b	0.77	a	0.76	a
Na	g kg ⁻¹	1.87	a	1.80	a	0.81	b	0.72	b
P	g kg ⁻¹	1.63	ns	1.59	ns	1.77	ns	1.73	ns
S	g kg ⁻¹	1.14	a	1.21	a	0.17	b	0.19	b
Si	g kg ⁻¹	0.28	a	0.26	a	0.08	b	0.14	b
Sr	g kg ⁻¹	0.09	ns	0.10	ns	0.11	ns	0.11	ns
Ti	g kg ⁻¹	0.57	ns	0.46	ns	0.63	ns	0.56	ns
As	mg kg ⁻¹	3.63	b	3.56	b	6.81	a	7.23	a
Be	mg kg ⁻¹	0.36	b	0.30	b	1.43	a	1.39	a
Cd	mg kg ⁻¹	<DL		<DL		<DL		<DL	
Co	mg kg ⁻¹	4.71	b	4.15	b	13.13	a	12.78	a
Cr	mg kg ⁻¹	38.70	b	41.44	b	93.50	a	87.98	a
Cu	mg kg ⁻¹	37.98	b	34.56	b	56.46	a	56.13	a
Li	mg kg ⁻¹	12.08	b	10.52	b	45.08	a	44.15	a
Mo	mg kg ⁻¹	4.27	a	4.01	a	<DL		<DL	
Ni	mg kg ⁻¹	17.82	b	15.31	b	46.03	a	45.54	a
Pb	mg kg ⁻¹	18.96	b	15.15	b	48.66	a	40.12	a
Sb	mg kg ⁻¹	1.30	b	1.30	b	1.53	a	1.43	a
Sn	mg kg ⁻¹	1.44	b	1.41	b	4.30	a	3.75	a
V	mg kg ⁻¹	19.44	b	16.32	b	65.61	a	63.46	a
Zn	mg kg ⁻¹	74.13	b	71.01	b	106.3	a	108.8	a

540

541

542 **Table 2.** Contamination indexes of soil-grown vegetables at the experimental site. BioAccumulation Factor,
 543 Translocation Factor, and Health Risk Index for adults (70 kg body weight and 0.22 kg d⁻¹ daily plant
 544 consumption) and children (26 kg body weight and 0.13 kg d⁻¹ daily plant consumption). Different letters
 545 indicate significant differences at P \geq 0.05 (n=6).

Element	Species	BioAccumulation Factor		Translocation Factor		Health Risk Index		Health Risk Index	
						adult		children	
Cr	Fennel	0.100	c	-		0.003	e	0.006	e
	Kale	0.060	e	0.260	a	0.001	c	0.002	c
	Lettuce	0.210	b	-		0.005	b	0.011	b
	Onion	0.060	f	-		0.0001	g	0.000	g
	Radish	0.780	a	-		0.015	a	0.032	a
	Basil	0.080	d	0.260	a	0.002	d	0.004	d
	Tomato	0.030	g	0.440	a	0.0002	f	0.001	f
Cu	Fennel	0.220	d	-		0.105	g	0.219	g
	Kale	0.120	g	0.330	c	0.037	c	0.078	c
	Lettuce	0.270	b	-		0.099	d	0.206	d
	Onion	0.140	f	-		0.064	f	0.133	f
	Radish	0.270	a	-		0.130	a	0.270	a
	Basil	0.260	c	1.170	a	0.114	b	0.236	b
	Tomato	0.220	e	0.520	b	0.076	e	0.159	e
Pb	Fennel	0.020	d	-		0.084	e	0.174	e
	Kale	0.020	e	0.780	a	0.049	d	0.102	d
	Lettuce	0.050	b	-		0.182	b	0.379	b
	Onion	0.020	g	-		0.019	g	0.041	g
	Radish	0.110	a	-		0.457	a	0.951	a
	Basil	0.040	c	0.750	a	0.121	f	0.252	f
	Tomato	0.020	f	0.390	b	0.038	e	0.079	e
Ni	Fennel	0.030	d	-		0.023	a	0.048	a
	Kale	0.030	c	0.660	a	0.019	a	0.040	a
	Lettuce	0.050	b	-		0.038	a	0.080	a
	Onion	0.003	f	-		0.002	a	0.004	a
	Radish	0.080	a	-		0.089	a	0.186	a
	Basil	0.010	e	0.220	b	0.014	a	0.030	a
	Tomato	0.000	g	0.000	c	0.000	a	0.000	a


547 **Figures**

548 **Figure 1.** Location of the experimental garden (X) in respect to main pollution sources, namely the city incinerator (I), Via San Donato (R), A14 motorway (M)
549 and main railway (T).



550

551 **Figure 2.** Accumulation of Cr (a), Cu (b), Pb (c) and Ni (d) (expressed in mg kg⁻¹ dry weight) in edible portions of radish, onion, fennel, lettuce, basil, kale and
 552 tomato according to growing system (soilless on peat in red or traditional on soil in black). Vertical bars represent standard deviation. Symbols (*) indicate
 553 significant differences at P≤0.05 (*), 0.01 (**), or 0.001 (***), whereas ‘ns’ indicates non significant differences at P≤0.05 (n=6).



554

555

556 **Figure 3.** Accumulation of Cr (a), Cu (b), Pb (c) and Ni (d) (expressed in mg kg⁻¹ dry weight) in edible portions of radish, onion, fennel, lettuce, basil, kale and
 557 tomato according to fertilization management (organic in red and mineral in black). Vertical bars represent standard deviation. Symbols (*) indicate significant
 558 differences at P≤0.05 (*), 0.01 (**), or 0.001 (***), whereas 'ns' indicates non significant differences at P≤0.05 (n=6).

