

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Polysaccharides from by-products of the Wonderful and Laffan pomegranate varieties: New insight into extraction and characterization

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Khatib, M., Giuliani, C., Rossi, F., Adessi, A., Al-Tamimi, A., Mazzola, G., et al. (2017). Polysaccharides from by-products of the Wonderful and Laffan pomegranate varieties: New insight into extraction and characterization. FOOD CHEMISTRY, 235, 58-66 [10.1016/j.foodchem.2017.05.041].

Availability:

This version is available at: <https://hdl.handle.net/11585/591676> since: 2017-05-25

Published:

DOI: <http://doi.org/10.1016/j.foodchem.2017.05.041>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Title: Polysaccharides from By-products of the Wonderful and Laffan Pomegranate Varieties: New Insight into extraction and characterization.

Article Type: Research Article (max 7,500 words)

Keywords: mesocarp, prebiotic activity, pectin, size exclusion chromatography, ¹H-NMR

Corresponding Author: Professor Nadia Mulinacci, ASSOCIATE PROFESSOR

Corresponding Author's Institution: UNIVERSITY OF FLORENCE

First Author: Mohamad Khatib

Order of Authors: Mohamad Khatib; Camilla Giuliani; Federico Rossi; Alessandra Adessi; Amal Al-Tamimi ; Giuseppe Mazzola; Diana Di Gioia; Marzia Innocenti; Nadia Mulinacci, ASSOCIATE PROFESSOR

Abstract: The main crude polysaccharides (CPS), extracted from two widely cultivated pomegranate varieties, Laffan and Wonderful, were studied and characterized. We obtained the highest CPS extraction yield (approximatively 10% w/w on dried matter) by 1 h of decoction (ratio 1/40 w/v). The predominant polymers (75-80%) of the CPS samples shown a hydrodynamic volume close to 2000 kDa by size exclusion chromatography and the exocarp and mesocarp profiles were very similar. The proton spectra (¹H-NMR), according to sugar composition and gelling ability, confirmed the main polysaccharide fractions were pectin with different acylation and methylation degree. The CPS from Laffan and Wonderful mesocarp showed prebiotic properties in vitro with Lactobacillus and Bifidobacterium strains. The composition of the decoction (12 % ellagitannins and 10 % of CPS) obtained by a green extraction process of pomegranate by-products, makes it a suitable component of functional food formulations.

This is a preprint version of the article, which has been published in final form in Food chemistry Volume 235, November 2017, Pages 58-66. The final published version is available online at: <https://doi.org/10.1016/j.foodchem.2017.05.041>

1 Polysaccharides from By-products of the Wonderful and Laffan Pomegranate Varieties: New
2 Insight into extraction and characterization.

3

4 Mohamad Khatib, Camilla Giuliani, ²Federico Rossi, ²Alessandra Adessi, ¹Amal Al-Tamimi

5 ³Giuseppe Mazzola, ³Diana Di Gioia, Marzia Innocenti, *Nadia Mulinacci

6

7 *Department of Neuroscience, Psychology, Drug and Child Health, Pharmaceutical and*
8 *Nutraceutical Section, University of Florence, Via Ugo Schiff 6, Sesto Fiorentino, (FI) Italy.*

9 *1 Biology Department of Ecology, Princess Noura University, Al Imam Abdullah Ibn Saud Ibn*
10 *Abdul Aziz Road, Riyadh, Saudi Arabia.*

11 *2 Department of the Sciences of Agrifood Production and the Environment, University of Florence,*
12 *Piazzale delle Cascine, 18 - 50144 Firenze, Italy.*

13 *3 Department of Agricultural Sciences, University of Bologna, Viale Fanin, 44 - 40127*
14 *Bologna, Italy.*

15

16

17 *Corresponding author: Nadia Mulinacci

18 Dipartimento di NEUROFARBA, Università degli Studi di Firenze, Via Ugo Schiff 6, 50019 Sesto
19 Fiorentino, Firenze, Italy.

20 Tel.: 0039-055-4573773

21

22

23 e-mail: nadia.mulinacci@unifi.it

24 **Abstract**

25 The main crude polysaccharides (CPS), extracted from two widely cultivated pomegranate
26 varieties, Laffan and Wonderful, were studied and characterized. We obtained the highest CPS
27 extraction yield (approximately 10% w/w on dried matter) by 1 h of decoction (ratio 1/40 w/v).
28 The predominant polymers (75-80%) of the CPS samples shown a hydrodynamic volume close to
29 2000 kDa by size exclusion chromatography and the exocarp and mesocarp profiles were very
30 similar. The proton spectra (¹H-NMR), according to sugar composition and gelling ability,
31 confirmed the main polysaccharide fractions were pectin with different acylation and methylation
32 degree. The CPS from Laffan and Wonderful mesocarp showed prebiotic properties *in vitro* with
33 *Lactobacillus* and *Bifidobacterium* strains. The composition of the decoction (12 % ellagitannins
34 and 10 % of CPS) obtained by a green extraction process of pomegranate by-products, makes it a
35 suitable component of functional food formulations.

36

37 **Keywords:** mesocarp, prebiotic activity, pectin, size exclusion chromatography, ¹H-NMR

38

1. Introduction

The *Punica granatum* L. (Punicaceae) fruit has been extensively used in the folk medicine of many cultures (Viuda-Martos, Fernández-Lóaez, & Pérez-álvarez, 2010), exhibiting a wide range of potential clinical applications (Viuda-Martos et al., 2010) including antitumor properties (Joseph, Aravind, George, Varghese, & Sreelekha, 2013; Joseph, Aravind, Varghese, Mini, & Sreelekha, 2012; Li, Zhang, & Wang, 2014). Up until now, research indicated that the ellagitannins are the principal bioactive constituents of the different extracts obtained from pomegranate fruit. However, there are few studies regarding the extraction and characterization of the polysaccharide fractions recovered from the different parts of the fruit.

To date, natural polysaccharides have been proven to exert antioxidant, antitumor, immunomodulatory, antimicrobial, antiulcer and hypoglycemic activities (Leung, Liu, Koon, & Fung, 2006; Negi, Jayaprakasha, & Jena, 2003; Schepetkin & Quinn, 2006). A polysaccharide extracted from pomegranate peel has shown significant antioxidant, antiglycation and tyrosinase inhibition properties (Rout & Banerjee, 2007). A more recent paper showed that a galactomannan recovered from the fruit rind of *P. granatum*, exerted *in vitro* immunomodulatory and free radical scavenging activities (Joseph et al., 2012), as well as anticancer activity in mice by reducing the tumor either alone or in combination with doxorubicin (Joseph et al., 2013). One of these studies provided evidence of the non-toxic nature of this plant-derived compound, which was also proposed as an adjuvant or as a single agent for the treatment of cancer (Joseph et al., 2013). Polysaccharides from pomegranate peel have also been reported as able to inhibit the proliferation of U-2 human osteosarcoma cancer cells by inducing apoptosis mainly through a mitochondrial signalling pathway (Li et al., 2014).

At the same time, it is known that polysaccharides are an important part of soluble fermentable dietary fiber. They can exhibit prebiotic activity by stimulating the growth of beneficial bacteria in the colon, thereby contributing to the healthy status of the gut (Di Gioia et al., 2014a; Marotti et al.,

2012). A balanced gut microbial composition confers benefits to the host, due to the modulation of metabolic and immune functions, while microbial imbalances are associated with metabolic disorders and/or diseases (Di Gioia, Aloisio, Mazzola, & Biavati 2014b; Tremaroli & Bäckhed, 2012). Therefore, the maintenance of a correct equilibrium between beneficial microorganisms, mainly belonging to the *Bifidobacterium* and *Lactobacillus* genera, and potentially pathogenic strains, is crucial for host health.

The interest in food processing by-products has increased recently. In particular, waste by-products from pomegranate juice production are being considered for the recovery of bioactive compounds, primarily ellagitannins (Akhtar, Ismail, Fraternale, & Sestili, 2015; Goula & Lazarides, 2015), while polysaccharides have not yet been considered. In literature, it is frequently used to indicate the exocarp (the real peel) and mesocarp together, without making a real distinction between these two parts of the fruit (Viuda-Martos et al., 2010). The main by-products of juice production, the mesocarp (40-50 % of the whole fruit) and exocarp, have not been investigated singularly as potential sources of bioactive polysaccharides, and no data are available on the possible prebiotic properties of these polymers recovered from pomegranate. To the best of our knowledge, none of the studies on pomegranate polysaccharides have taken into account the Wonderful and Laffan varieties, the objects of our investigation.

The aim of this research was to study the polysaccharides from the by-products of Laffan and Wonderful, generated in large amount from these two widely diffused pomegranate varieties. Water extraction processes were applied to efficiently recover the polysaccharides separately from the mesocarp and exocarp. Size exclusion chromatography, chemical hydrolysis and proton nuclear magnetic resonance (¹H-NMR) were used to analyze the characteristics of the principal polysaccharides. For the first time, the prebiotic properties of crude polysaccharides (CPS) from the Wonderful mesocarp were assessed by *in vitro* by testing them on *Bifidobacterium* and *Lactobacillus* strains. The dry decoction is proposed as source of polysaccharides and ellagitannins.

2. Materials and methods

2.1 Materials

The Laffan cultivar (sour-sweet) was harvested from Rif Idlib, Syria in October 2011; the Wonderful cultivar was purchased from Ortofrutta Grosseto (Italy) in October 2013. About 7-10 kg of fresh ripe fruits of both cultivars were used as the source of the exocarp and mesocarp for extraction of polysaccharides.

2.2 Extraction process for recovery of CPS

The exocarp and mesocarp were manually separated from fresh pomegranate fruits, then cut into small pieces and freeze-dried. Both parts were powdered in a grinder and extracted with distilled water as summarized in Table 1. The term dried matter in the text refers to the dry weight of the mesocarp and exocarp. Each solution, recovered after centrifugation (3900 g for 12 min at 5°C) according to the methods listed in Table 1, was supplemented with 2 volumes of ethanol and kept for 3 h at 0°C to induce the precipitation of polysaccharides, which were recovered after a new centrifugation, again at 3900 g for 12 min at 5°C. The further addition of ethanol to the supernatants did not induce the formation of new precipitate. The recovered polysaccharides were freeze-dried, then re-dissolved in a minimum water volume and treated again with 2 volumes of ethanol. The precipitate was then freeze-dried to obtain the crude polysaccharides (CPS) which were successively treated to remove the proteins according to the method reported in Joseph et al., (2012). Briefly, CPS were dissolved in water and extracted with 3 volumes of chloroform. The extraction was repeated several times until the water/chloroform inter-phase became clear. The aqueous phase containing the purified polysaccharides was recovered and freeze-dried to obtain CPSp as summarized in Table 3. CPSp were re-dissolved in water, and 500 µL of this solution were transferred into an ultra-filter device with a cut-off of 10,000 Daltons (Amicon, Millipore, Billerica,

114 MA) and centrifuged at 11,000 g for 15 min. The precipitate was re-suspended in its original
115 volume by adding water; the process was repeated up to 7 times, (as indicated by the supplier), to
116 remove about 99% of possible fouling materials (polar low molecular weight molecules and salts)
117 from the sample. After these cleaning steps, the filter device was placed upside down in a clean
118 microcentrifuge tube for 2 minutes at 1,000 g; 500 μ L of distilled water were then added to dissolve
119 and recover the polysaccharides after filtration.

120

121 2.3 ¹H- NMR analysis

122 The purified polysaccharides from the mesocarp of Wonderful (W-CPSp) and Laffan (L-CPSp)
123 were dialyzed for 48 h at 5°C in a nitrocellulose membrane with a 12-14 kDalton cut-off (Medicell
124 International Ltd, London), and then freeze-dried. The dried samples were dissolved in 1 mL of
125 D₂O and maleic acid was added as internal standard as follows: 6.1 mg W-CPSp and 1.1 mg of
126 maleic acid, 6.3 mg L-CPSp and 1.3 mg of maleic acid, purity grade 98% (Merck, Germany). The
127 ¹H-NMR experiments were carried out using a 400 MHz instrument Advance 400 (Bruker, Bremen,
128 Germany). The quantitative evaluation was done according to reference guidelines (Eurolabs,
129 2014), applying the same protocol previously used for other matrices (Khatib, Pieraccini, Innocenti,
130 Melani, & Mulinacci, 2016).

131

132 2.4 Determination of monosaccharide composition

133 The polysaccharides in Table 2 from Laffan and Wonderful mesocarp and exocarp were
134 hydrolyzed in acidic media (Erbing, Jansson, Widmalm, & Nimmich, 1995). Briefly, 1 mL of a 2 M
135 trifluoroacetic acid water solution was added to 5 mg CPSp, maintained at 120°C for 120 min.
136 Afterwards, samples were cooled on ice, and ultrafiltered at 3500 g for 20 min using 3,000 Daltons
137 cut-off centrifuge filter devices (Amicon Ultra-4, Millipore, Billerica, MA). The supernatant was
138 then dried by a rotavapor at 37 °C, and re-suspended in 1 mL MilliQ-grade water. This

139 evaporation/re-suspension process was repeated three times, with the aim of removing the
140 trifluoroacetic acid that could introduce bias **into** the analysis. The samples were washed twice with
141 MilliQ-grade water, re-dissolved in 1 mL deionized water and then analyzed by ion exchange
142 chromatography using a ICS-2500 ion chromatograph with an ED₅₀ pulsed amperometric detector, a
143 gold working electrode and a Carbopac PA1 250x4mm column, all from Dionex (Sunnyvale, CA,
144 USA). The eluents used were MilliQ-grade water (solution A), 0.185 M sodium hydroxide solution
145 (solution B), and 0.488 M sodium acetate solution (solution C). A gradient elution was used
146 consisting of a first stage (injection time **up** to the 7th min) with the eluent constituted by 84%
147 solution A, 15% solution B, and 1% solution C; a second stage (injection time from the 7th to 13th
148 min) with the eluents constituted by 50% solution B and 50% solution C; and a final stage (injection
149 time from the 13th to the 30th min) with the eluents consisting of 84% solution A, 15% solution B,
150 and 1% solution C. The flow rate was 1 mL min⁻¹. The monosaccharides were detected according to
151 the retention time of pure monosaccharides purchased from Sigma-Aldrich (Milan, Italy) after
152 specific spike injections of the pure monosaccharides; at least three standard injections were
153 repeated in order to obtain a mean retention time of each monosaccharide, and the variance never
154 exceeded 5%.

155

156 2.5 Size Exclusion Chromatography

157 The apparent molecular weight of the polysaccharides of the CPS samples was determined
158 according to a previously reported method (Chen et al., 2014; Colica, Li, Rossi, De Philippis, &
159 Liu, 2015), with some modifications. The samples listed in Table 3 were weighed and dissolved in
160 distilled water, at a concentration of roughly 0.14 mg mL⁻¹. The solution was analyzed using a
161 Varian ProStar HPLC chromatograph (Varian, USA) equipped with a 355 refractive index detector
162 and a Biosep s4000 column (Phenomenex, USA). The samples were analyzed with 30 min runs by
163 HPLC-grade water as eluent at 0.6 mL min⁻¹ flow rate. Blue dextrans (Sigma-Aldrich, USA) at

different molecular weights (approx. 2000 kDa, 1100 kDa, 410 kDa, 150 kDa and 50 kDa) were used as standards for hydrodynamic volume calculation.

2.6 *In vitro* evaluation of the prebiotic activity of CPS

The ability of CPS to induce the growth of beneficial bacteria (prebiotic activity) was assayed using two strains previously isolated from human feces: *Bifidobacterium breve* B632 (Aloisio et al., 2012) and *Lactobacillus plantarum* L12. The latter was isolated from a healthy volunteer (unpublished results) and taxonomic characterization was performed via 16S rDNA amplification and sequencing (Gaggia et al., 2013), this strain is available at the Bologna University, Scardovi Collection of Bifidobacteria. Both strains were stored in lyophilized form. When necessary, they were re-vitalized in de Man Rogosa Sharpe (MRS) medium (Oxoid, Basingstone, UK) supplemented with 0.05% cysteine and incubated in anaerobic conditions at 37°C for 24 h. Anaerobic conditions were created in a capped jar using an anaerobic atmosphere generation system (Anaerocult A, Merck, Darmstadt, Germany).

The MRS medium composition was modified to perform the growth experiment with the pomegranate polysaccharides. The modified medium (m-MRS) did not contain the carbon source (glucose), which was provided by the pomegranate polysaccharides, and had a halved amount of potential growth substrate, such as peptone, yeast extract and meat extract compared to those present in the original medium (peptone, 5 g L⁻¹; yeast extract, 2 g/L, meat extract, 5 g L⁻¹, where the amounts are in m-MRS).

The prebiotic activity was evaluated using CPS at 0.5% (w/v) in m-MRS. A positive growth control was performed using m-MRS with 0.5% glucose and a negative control in m-MRS with no added carbon source. The medium containing CPS as the carbon source was prepared as follows: the m-MRS ingredients were weighed in a flask, dissolved in water and the medium was autoclaved at 120°C for 15 min. A 0.5% (w/v) fiber or glucose at the same concentration were added to the hot

189 medium, stirred, and sterilized again at 102°C for 10 min. This procedure allowed the fiber to
190 dissolve in the medium and, at the same time, to prevent risk of growth of **undesirable**
191 microorganisms. The *B. breve* B632 and *L. plantarum* M12 strains were grown overnight in the
192 respective media, centrifuged, washed in phosphate buffered saline (PBS) and re-suspended in PBS
193 to obtain a solution having an absorbance of 0.7 at 600 nm. This suspension was used to inoculate at
194 2% (v/v) the flasks containing the m-MRS medium plus the fiber, the m-MRS medium plus glucose
195 (positive control) and the m-MRS medium with no additional carbon source (negative control). The
196 flasks were incubated at 37°C in anaerobic conditions for 48 h and a 1 mL culture was sampled
197 from each flask for viable bacterial counts at pre-established times (0, 6, 24, 30 and 48 h of
198 incubation). The sampled amount was mixed with 9 mL of PBS, serially diluted in the same
199 solution and plated on agarized MRS supplemented with cysteine. Following incubation of the
200 plates at 37°C in anaerobic conditions for 24 h, the number of colonies, corresponding to the
201 number of viable cells, was counted. The number of cells expressed as CFU mL⁻¹ were transformed
202 into Log₁₀ value (Log CFU mL⁻¹).

203

204 2.7 Proximate composition and dietary fiber analyses

205 The proximate composition was determined for the decoction from mesocarp of the Wonderful
206 variety. Protein content (PC) was evaluated using the Kjeldhal method: $PC (g/100g) = N \times 6.25$,
207 where N is total nitrogen. The total fat content was determined by Soxhlet extraction, and
208 gravimetrically determined according to ISS protocol (1996/34). Ash content was evaluated by
209 gravimetric assay, according to ISS protocol, (1996/34 method b). Dietary fibers (both soluble and
210 insoluble) were quantified according to AOAC method 991.43 (Determination of soluble, insoluble
211 and total dietary fiber in foods and food products, final approval 1991).

212

213 2.8 Statistical analysis

214 All data in Figure 1, Tables 1 and 3, are presented as mean±SD from triplicate measurements of
215 each measuring point. Statistical significance for evaluating the prebiotic properties of CPS from
216 the mesocarp of Laffan and Wonderful cultivars was calculated within each evaluation time (T6,
217 T24, T30, T48) with a t-test, using the MEANS procedure (SAS).

218

219 3. Results and Discussion

220 3.1 Recovery of polysaccharides: preliminary evaluation on Laffan

221 The decision to study the by-products from Laffan and Wonderful pomegranate varieties, was
222 mainly determined by the high amount produced because the diffusion of the two varieties. The
223 Laffan pomegranate is widely present in Syria but also in Southern Turkey and Israel, while the
224 Wonderful is one of the principal variety cultivated in the Western world. Although a valorisation of
225 the by-products derived from juice production, requires better knowledge of their composition, so
226 far little attention has been addressed to polysaccharides from pomegranates, and the fruits are
227 mainly known for their juice rich in anthocyanins and ellagitannins. Water extraction (sometimes
228 coupled with increased temperature), and subsequent precipitation by adding ethanol, is the most
229 commonly utilized method for recovering polysaccharides from different sources (Huie & Di, 2004;
230 Joseph et al., 2012; Zhu & Liu, 2013). We used a similar procedure on the mesocarp and exocarp
231 separately, to evaluate the polysaccharide content of Wonderful and Laffan. First of all, to select the
232 most efficient extractive procedure, we used the Laffan mesocarp as reference material.

233 To increase the extractive yields, (Table 1), different extraction times, extraction temperatures
234 and dried matter/water ratios, were evaluated. Firstly, 30 and 60 min were set, applying a single or
235 two successive extraction steps, and varying the extractive ratio from 1:15 w/v to 1:40 w/v. The
236 extraction was firstly performed at 25±2 °C as previously proposed for pomegranate (Rout &
237 Banerjee, 2007). A second approach was to pre-treat the dried material with a hydroalcoholic
238 solution to remove the ellagitannins, and then extract the polysaccharides by hot water.

239 To remove part of the impurities co-precipitated after the first ethanol addition, the
240 polysaccharides were re-dissolved in water and precipitated again, adding ethanol to get the CPS
241 listed in Table 1. To verify if this latter step was effective in cleaning the polysaccharides, we
242 evaluated the amount of the impurities by weighing the dried supernatant recovered after the second
243 ethanol addition. The impurities were 5.4% and 7.4% of dried mesocarp for Wonderful and Laffan,
244 respectively, and close to 3% of the dried exocarp for both varieties. These results indicate that the
245 second addition of ethanol was necessary to obtain a cleaner polysaccharide fraction (CPS).
246 As shown in Table 1, the yield in CPS increased from 5% to 8% with a longer extraction time (from
247 30 to 60 min) by applying the same extractive ratio (1:15 w/v). The yield further increased up to
248 10% by applying a single extraction of 60 min and a higher extractive ratio (1/40 w/v). A successive
249 extractive step of 60 min, as well as previous contact of the dried material with water before the
250 decoction, did not increase the recovery of CPS.

251 Overall, the best result in terms of yield and reproducibility, was obtained with a single
252 decoction of 60 min and an extractive ratio of 1/40 p/v (Table 1). Similar recoveries of
253 polysaccharides (10-13%) were reported by Zhu et al., for a pomegranate purchased from a local
254 Chinese market and extracted by hot water (Zhu and Liu, 2013). The same authors have
255 successively proposed an ultrasound-assisted hot water extraction, but obtained similar yields (Zhu,
256 Zhai, Li, Wu, & Li, 2015). In both these studies, and as frequently reported in literature, the authors
257 cited the pomegranate peel but did not specify if the raw material was comprised only of the
258 exocarp or of the mesocarp plus exocarp. Lastly, and in agreement with a previous report (Rout &
259 Banerjee, 2007), we confirmed that extraction with water at a temperature close to 25°C, even when
260 applying long extraction times, gave considerably lower CPS yields (Table 1).

261 Since there are about 12% of ellagitannins in the decoction of Laffan mesocarp (Khatib, 2015),
262 a pretreatment with ethanol 70% v/v was also tested to remove these polar compounds before
263 precipitation of the CPS. Even if the CPS yields are lower than those obtained without applying this

264 treatment, this latter approach can be useful when the objective is to efficiently recover the
265 ellagitannins before precipitation of polysaccharides (Table 1).

266

267 3.2 CPS recovered from the two cultivars

268 After the pre-screening carried out only on the Laffan mesocarp, only the more efficient
269 methods were selected to extract the CPS from the Laffan and Wonderful exocarp and mesocarp
270 (Table 2). Overall, by applying the same extractive method, we obtained similar results from the
271 exocarp and mesocarp of the two varieties. Again, a one-step decoction gave the highest % yields,
272 and the hot water is determinant for maximizing the extractability of CPS because it increases
273 polysaccharide solubility. On the other hand, the extraction carried out at room temperature, was
274 confirmed as the worst. Despite the low yields, this latter method was tested again to verify the
275 effect of temperature on the characteristics of CPS. To this aim, we analyzed the recovered
276 polysaccharides by size exclusion chromatography and compared their profiles with those from
277 CPS obtained by the hot extraction. As shown in Table 2, CPS were mainly located in the mesocarp,
278 with lower values in the exocarp (4.5-4.7%) for both varieties. There was some variability in the
279 method with the CPS amount recovered from exocarp having higher standard deviation values,
280 from 11% to 25%. This finding is attributable to the non-homogeneous thickness of the removed
281 exocarp, still containing residual parts of mesocarp, which is hard to completely remove. Lastly, the
282 decoction of the exocarp, carried out after a previous extraction with ethanol/water (7:3v/v),
283 provided CPS amounts close to 4% and similar to those derived without using the hydroalcoholic
284 solution pre-treatment.

285 It must be emphasized that boiling is a suitable method not only for polysaccharides but also
286 for co-extraction of the ellagitannins in amounts close to 120 mg/g dried decoction (Khatib, 2015).
287 Furthermore, the drying process of the decoction did not require the addition of excipients such as
288 the maltodextrins, commonly used to reduce the final hygroscopicity of the dried herbal extract.

289 This advantage, not frequently observed during the management of herbal products, can be
290 attributable to the presence of CPS in a relatively high amounts.

291 Due to the difficulty of procuring enough fresh Laffan pomegranate during the civil war in
292 Syria, we only determined the nutritional composition of the Wonderful mesocarp decoction. In
293 summary, the total dietary fiber determined by the AOAC.993.41 method was 9.66 % comparable to
294 the CPS content. Moreover, the main fraction was soluble fiber, (6.67 %) fermentable by human
295 microbiota. There was 2.3 % of total proteins and 5.6 % ash, indicating an appreciable amount of
296 minerals.

297

298 3.3 Sugar composition by hydrolysis

299 To verify the purity grade of polysaccharides in terms of the co-presence of oligosaccharides
300 and inorganic salts, the efficiency of the ultrafiltration devices was tested on CPSp from the
301 mesocarp of the two varieties (Table 2). The hydrolysis of CPSp samples before and after the cut-
302 off filtration provided the same results in terms of molar percentage of monosaccharides suggesting
303 that the samples listed in Table 2 did not need further purification by this filter device.

304 The CPSp samples listed in Table 3 were treated with TFA acid to hydrolyze the polysaccharide
305 strands and subsequently determine sugar composition by ionic exchange chromatography,
306 according to a previous method (Erbing et al., 1995). The CPSp samples from both the mesocarp
307 and exocarp showed a very similar composition for both the varieties (Table 3). Hexoses galactose
308 and glucose, deoxysugar rhamnose, and galacturonic acid, were the most abundant monomers,
309 while the main aldopentoses were xylose and arabinose. From our findings it emerges that these two
310 varieties, Laffan from Syria and Wonderful widely diffused throughout the western world, have a
311 very similar compositional profile in terms of polysaccharides. This result is not completely
312 unexpected, and in agreement with a previous work in which it was hypothesized that the
313 Wonderful is derived from the more antique Laffan variety (Goor, 1967).

314 Overall, other reports on pomegranate by-products did not include the varieties selected in this
315 study. In regard to polysaccharides, the literature indicates there is a wide variability in terms of
316 sugar composition depending on the variety, growth site and purity grade of the polysaccharide
317 itself (Normakhtov, Rakhmanberdyeva, & Rakhimov, 1999; Jahfar, Vijayan, & Azadi, 2003) .

318 *3.4 Characterization of the polysaccharide fractions*

319 The CPSs from the mesocarp and exocarp of the two cultivars were analyzed by size exclusion
320 chromatography to determine their apparent molecular weight. Since these polymers may be
321 characterized by a branched structure, often related to the presence of arabinose, galactose and
322 xylose, their size was calculated in terms of hydrodynamic volume, and not in terms of actual
323 molecular weight. The CPS samples were compared to dextrans standards, considering that a 2000
324 kDa fraction has the same hydrodynamic volume as dextran at 2000 kDa molecular weight. The
325 analyzed fractions throughout the text are identified as molecular weight, although with
326 approximation.

327 The data in Figure 1 highlighted that CPS of Laffan and Wonderful mesocarp and exocarp,
328 were of similar molecular weight, since no significant differences were found; all the CPS were
329 characterized by a predominant fraction of about 2000 kDa, accounting for 75.4% of the total. The
330 remaining 24.6% was represented by five minor fractions, the most common being: i) a fraction
331 having a molecular weight between 410 kDa, and 150 kDa (7.4% of total CPS); ii) a fraction having
332 a molecular weight lower than 50 kDa, accounting for 8.9% of total CPS. As expected, more
333 variability was observed for the fractions having small molecular weights (much lower than 50
334 kDa).

335 Few reports are available to date on polysaccharide structure from pomegranate fruit. A first
336 report described a glucofructan extracted from the peel, having 31 kDa molecular weight, that was
337 separated using Sephadex G100 column (Jahfar, et. al., 2003). Sun described a polysaccharide
338 extracted from the rind of a non-specified variety, having a molecular weight of 110 kDa

determined by gel filtration on a Sephadex G200 column and dextrans at different molecular weights as reference standards (Sun, Li, Yan, & Liu, 2010). More recently, a glucomannan was extracted from the rind of a ripe pomegranate fruit and the authors indicated a molecular weight of 110 kDa (Joseph et al., 2012). None of these studies specified which cultivar or variety was investigated.

The present work shows the molecular weight distribution of pomegranate polysaccharides obtained from the Laffan and Wonderful cultivars by using size exclusion chromatography for the first time. We demonstrated that the CPS samples have similar apparent molecular weight distribution with overlapping profiles of the two cultivars and the two parts of the fruit. We also verified that hot extraction (100°C, 1 h) did not modify the CPS composition as demonstrated by the complete overlap of size exclusion chromatography profiles obtained after extraction with cold water and boiling water (Table 2).

Recently, some authors (Moorthy, Maran, Surya, Naganyashree, & Shivamathi, 2015; Pereira et al., 2016) observed the presence of pectin in pomegranate fruit but no spectral data, particularly ¹H-NMR spectra, are reported or discussed. Taking into account these data, we searched for the presence of pectin by analysing of the proton spectra of CPSp from Laffan and Wonderful mesocarp (Figure 2). According to the literature (Bédouet, Courtois, & Courtois, 2003), specific signals indicate the presence of O-methyl and O-acetyl groups typical of pectin and their intensity can be associated with the degree of methylation and acetylation. As shown in Figure 2, the two spectra obtained dissolving comparable amounts of CPSp from the two varieties, clearly revealed signals attributable to O-acetyl groups in the region (of δ 1.98-2.15) and an intense signal ascribable to a singlet of O-methyl groups close to δ 3.7. Both these data and the high percentage of galacturonic acid after the acidic hydrolysis (Table 3), confirm the presence of pectin in Laffan and Wonderful. The singlet at δ 6.31 is due to maleic acid, that added as internal standard permits a preliminary comparison of the degree of methylation and acylation in CPSp from the two varieties. In other

words, the addition of an accurately weighed internal standard can be usefully applied for quantitative purposes. Particularly, the higher intensity observed for the signal at δ 2.15 in the Wonderful spectrum, indicates a higher degree of acylation compared to that of Laffan. The opposite behavior is observed for the signal at δ 1.98 ppm that was more intense in Laffan sample. Analogously, a different degree of methylation is indicated by the signal at δ 3.73, ascribable to a singlet of O-methyl groups (Cui, 2005) at higher intensity in the Laffan sample. Finally, the presence of less intense signals close to δ 1.1 is in agreement with the presence of low amounts of rhamnose units according to hydrolysis results (Table 3). Overall, this rapid measurement, obtained without the need of high magnetic field spectrometer, was able to point out structural differences between W-CPSp and L-CPSp, not highlighted by the size exclusion chromatography technique, showing the same profile for these samples.

Although further studies are needed to elucidate the structure of these polysaccharides, the ^1H -NMR spectra and the sugar composition derived by acidic hydrolysis suggest that the main polysaccharides of pomegranate mesocarp are pectin with different degree of methylation and acetylation. This applied hydrolysis method was recently confirmed as being suitable to guarantee a complete hydrolysis of pectin (Wikiera, Mika, Starzyńska-Janiszewska, & Stodolak, 2015).

We also carried out preliminary tests to evaluate the water-absorption ability of some dried polysaccharide fractions: LM-CPS, WM-CPS and WE-CPS. The adsorbed water ranged from 98.6 to 99.1% of the dried material. Adsorption was rapid and the final samples appeared as clumps with a gel consistency, exhibiting the well-known pectin behaviour.

3.5 *In vitro* evaluation of prebiotic properties

The decoction from mesocarp, was used to recover the CPS for *in vitro* tests of prebiotic properties. We investigated the ability of *B. breve* B632 and *L. plantarum* L12 strains to use crude polysaccharides from pomegranate exocarp and mesocarp as their carbon source and compared this

389 to their growth on glucose, *i.e.* an easily fermentable carbon source. Bifidobacteria and Lactobacilli
390 are able to compete for nutrients with enteric pathogens, to adhere strongly to the intestinal mucosa,
391 thus preventing pathogen adhesion, and to stimulate the development of the mucosal immune
392 system. Moreover, they are known to provide some protection against incoming enteric pathogens
393 in man (Jankowska, Laubitz, Antushevich, Zabielski, & Grzesiuk, 2008; Symonds et al., 2012).

394 Figure 3 shows that both strains grow well on CPS from Laffan and Wonderful mesocarp, being
395 significantly higher ($p < 0.01$) than the negative controls (*i.e.* with no added carbon source) and
396 comparable to that of an easily fermentable carbon source such as glucose added at the same
397 concentration. Growth on the Laffan variety at 24 h was only 0.6 and 0.1 Log CFU/mL lower than
398 that on glucose for *L. plantarum* L12 and *B. breve* B632, respectively. Growth on the Wonderful
399 variety at 24 h was 1.0 and 0.2 Log CFU/ml lower, respectively, than that on glucose for the same
400 strains. After the 24th h of incubation, both strains grown on glucose entered the steady phase,
401 whereas a small decrease in cell survival was observed with CPS as the carbon source. The results
402 shown in Figure 3 clearly indicate that CPS and/or the products of their degradation are not toxic
403 for the assayed strains and, on the contrary, are good growth substrates for them. Growth on the
404 medium with no added carbon source reached only a 1 Log CFU/mL increase at the 24th h
405 compared to the beginning of incubation, thus showing that the m-MRS medium used in the
406 experiments is a valid choice for performing prebiotic activity tests. Furthermore, if we might
407 propose the whole dried decoction for human consumption, due to its easy and rapid preparation,
408 the same is not true for the sub-fractions in Figure 1, whose preparation was longer and more
409 complex. Evaluation of the prebiotic activity of a single fraction from CPS was outside of the scope
410 of our research, but could be object of future investigations.

411 In agreement with our results, a high ability to ferment pectin by human gut microbiota
412 associated with an increase of almost 25 % of *Bifidobacterium* has been demonstrated *in vitro*
413 (Yang, Martinez, Walter, Keshavarzian, & Rose, 2013). Moreover, several studies in the literature
414 (as reviewed by Koropatkin, Cameron, & Martens, 2012) show that the degradation of complex

carbohydrates (glycans and polysaccharides) is a major symbiotic function carried out by microorganisms that inhabit the human distal gut, which increases host nutrition by digesting glycans that the host cannot degrade, providing the host with usable metabolic products such as short-chain fatty acids. Therefore, glycans shape the composition of the gut microbiota. Members of the Firmicutes and Actinobacteria phyla, to which *Lactobacillus* and *Bifidobacteria* spp. belong, possess different glycan acquisition strategies that also involve glycan-degrading enzymes (Mahowald et al., 2009).

4. Conclusions

This work improves the knowledge of the chemical and physical properties of polysaccharides recovered from the typical wastes of the pomegranate fruit, and reveals future perspectives for adding value to these food by-products, produced in large amount but currently discarded. The use of hot water maximized solubility and extractability of the crude polysaccharides from the Laffan and Wonderful varieties. The maximum recovery of polysaccharides was obtained from mesocarp, by a single-step water decoction. At the same time, the boiling process did not modify the molecular size distribution of the polysaccharides as demonstrated by their profiles in size exclusion chromatography, comparable with those obtained by a cold-water extraction. For the first time, the size exclusion chromatography was applied to evaluate the polysaccharides from mesocarp and exocarp of Laffan and Wonderful. A very similar distribution of the apparent molecular weights of the main polysaccharides was highlighted for the two varieties, with chromatographic profiles characterized by a predominant polymer with a hydrodynamic volume close to 2000 kDa, and five other minor fractions. The ¹H-NMR spectra, the sugar composition and the high gelling capacity of some purified polysaccharide fractions of mesocarp, confirmed the presence of pectin as primary component. The use of maleic acid as internal standard was proposed to evaluate the acylation and methylation degree of the main purified polysaccharide fractions. Finally, the crude polysaccharides

440 from Laffan and Wonderful pomegranate mesocarp showed prebiotic properties *in vitro* by serving
441 as an excellent substrate for the growth of potentially probiotic bacteria such as *Lactobacillus* and
442 *Bifidobacterium* strains.

443 We showed that, after a simple decoction of these pomegranate fruit by-products, it was
444 possible to obtain a dry extract rich in polysaccharides with prebiotic activity, associated with a
445 pool of bioactive ellagitannins. This combination of natural compounds can help to valorize these
446 by-products and to enhance the use of pomegranate dry decoction in functional food formulations.

447

448 **Acknowledgments.**

449 This research was partially funded by ECR of Florence within the project, Valorization of by-
450 products from olive oil production and pomegranate fruit processing; code ECRF15 and by
451 PAPARD project (ASI-2014-034-R.O- CUP- F82I14001080005) founded by ASI.

452

453 **References**

- 454 Akhtar, S., Ismail, T., Fraternale, D., & Sestili, P. (2015). Pomegranate peel and peel extracts:
455 Chemistry and food features. *Food Chemistry*, 174, 417–425.
- 456 Aloisio, I., Santini, C., Biavati, B., Dinelli, G., Cencič, A., Chingwaru, W., Mogna, L., & Di Gioia,
457 D. (2012). Characterization of Bifidobacterium spp. strains for the treatment of enteric
458 disorders in newborns. *Applied Microbiology and Biotechnology*, 96(6), 1561–1576.
- 459 Bédouet, L., Courtois, B., & Courtois, J. (2003). Rapid quantification of *O*-acetyl and *O*-methyl
460 residues in pectin extracts. *Carbohydrate Research*, 338(4), 379–383.
- 461 Chen, L., Rossi, F., Deng, S., Liu, Y., Wang, G., Adessi, A., & De Philippis, R. (2014).
462 Macromolecular and chemical features of the excreted extracellular polysaccharides in induced
463 biological soil crusts of different ages. *Soil Biology and Biochemistry*, 78, 1–9.
- 464 Colica, G., Li, H., Rossi, F., De Philippis, R., & Liu, Y. (2015). Differentiation of the characteristics
465 of excreted extracellular polysaccharides reveals the heterogeneous primary succession of
466 induced biological soil crusts. *Journal of Applied Phycology*, 27(5), 1935–1944.
- 467 Cui, S. W. (2005). Structural Analysis of Polysaccharides. In *Food Carbohydrates*. Taylor & Francis
468 (Eds.)
- 469 Di Gioia, D., Strahsburger, E., Lopez de Lacey, A. M., Bregola, V., Marotti, I., Aloisio, I., Biavati,
470 B., & Dinelli, G. (2014a). Flavonoid bioconversion in Bifidobacterium pseudocatenulatum
471 B7003: A potential probiotic strain for functional food development. *Journal of Functional*
472 *Foods*, 7, 671-679.
- 473 Di Gioia, D., Aloisio, I., Mazzola, G., & Biavati, B. (2014b). Bifidobacteria: Their impact on gut
474 microbiota composition and their applications as probiotics in infants. *Applied Microbiology*
475 *and Biotechnology*, 98(2), 563–577.

476

477 Erbing, B., Jansson, P. E., Widmalm, G., & Nimmich, W. (1995). Structure of the Capsular
478 Polysaccharide from the Klebsiella K8 reference strain 1015. *Carbohydrate Research*, 273(2),
479 197–205.

480 Eurolabs Technical Report No. 01/ 2014. *Eurolabs*, (1), 1–20. Retrieved from
481 [http://www.eurolab.org/documents/EUROLAB Technical Report NMR Method Development](http://www.eurolab.org/documents/EUROLAB%20Technical%20Report%20NMR%20Method%20Development%20and%20Validation%20May%202014_final.pdf)
482 [and Validation May 2014_final.pdf](http://www.eurolab.org/documents/EUROLAB%20Technical%20Report%20NMR%20Method%20Development%20and%20Validation%20May%202014_final.pdf)

483 Gaggia, F., Baffoni, L., Di Gioia, D., Accorsi, M., Bosi, S., Marotti, I., Biavati, B., & Dinelli, G.
484 (2013). Inoculation with microorganisms of *Lolium perenne* L.: evaluation of plant growth
485 parameters and endophytic colonization of roots. *New Biotechnology*, 30(6), 695-704.

486 Goor, A. (1967). The history of the pomegranate in the holy land. *Economic Botany*, 21(3), 215–
487 230.

488 Goula, A. M., & Lazarides, H. N. (2015). Integrated processes can turn industrial food waste into
489 valuable food by-products and/or ingredients: The cases of olive mill and pomegranate wastes.
490 *Journal of Food Engineering*, 167, 45–50.

491 Huie, C. W., & Di, X. (2004). Chromatographic and electrophoretic methods for Lingzhi
492 pharmacologically active components. *Journal of Chromatography B*, 812 (1-2), 241–257.

493 Jahfar, M., Vijayan, K. K., & Azadi, P. (2003). Studies on a polysaccharide from the fruit rind of
494 *Punica granatum*. *Research Journal of Chemistry and Environment*, 7(1), 43–50.

495 Jankowska, A., Laubitz, D., Antushevich, H., Zabielski, R. & Grzesiuk, E. (2008). Competition of
496 *Lactobacillus paracasei* with *Salmonella enterica* for Adhesion to Caco-2 Cells. *Journal of*
497 *Biomedicine and Biotechnology*, 2008 (1),1-6.

498 Joseph, M. M., Aravind, S. R., George, S. K., Varghese, S., & Sreelekha, T. T. (2013). A
 499 galactomannan polysaccharide from *Punica granatum* imparts in vitro and in vivo anticancer
 500 activity. *Carbohydrate Polymers*, 98(2), 1466–1475.

501 Joseph, M. M., Aravind, S. R., Varghese, S., Mini, S., & Sreelekha, T. T. (2012). Evaluation of
 502 antioxidant, antitumor and immunomodulatory properties of polysaccharide isolated from fruit
 503 rind of *Punica granatum*. *Molecular Medicine Reports* 5(2), 489–496.

504 Khatib, M. (2015). *Bioactive Compounds Into Edible Syrian Plants: Pomegranate and Capper*.
 505 University of Florence. <http://hdl.handle.net/2158/1045230>

506 Khatib, M., Pieraccini, G., Innocenti, M., Melani, F., & Mulinacci, N. (2016). An insight on the
 507 alkaloid content of Capparis spinosa L. root by HPLC-DAD-MS, MS/MS and ¹H qNMR.
 508 *Journal of Pharmaceutical and Biomedical Analysis*, 123, 53–62.

509 Koropatkin, N. M., Cameron, E. A., & Martens, E. C. (2012). How glycan metabolism shapes the
 510 human gut microbiota. *Nature Reviews Microbiology*, 10(5), 323–335.

511 Leung, M. Y. K., Liu, C., Koon, J. C. M., & Fung, K. P. (2006). Polysaccharide biological response
 512 modifiers. *Immunology Letters*, 105(2), 101–114.

513 Li, J., Zhang, F., & Wang, S. (2014). A polysaccharide from pomegranate peels induces the
 514 apoptosis of human osteosarcoma cells via the mitochondrial apoptotic pathway. *Tumor*
 515 *Biology*, 35 (8), 7475-7482.

516 Mahowald, M. A., Rey, F. E., Seedorf, H., Turnbaugh, P. J., Fulton, R. S., Wollam, A., Shah, N.,
 517 Wang, C., Magrini, V., Wilson, R. K., Cantarel, C. L., Coutinho, P. M., Henrissat, B., Crock, L.
 518 W., Russell, A., Verberkmoes, N. C., Hettich, R. L., & Gordon, J. I. (2009). Characterizing a
 519 model human gut microbiota composed of members of its two dominant bacterial phyla.
 520 *Proceedings of the National Academy of Sciences of the United States of America*, 106 (14),

521 5859–5864.

522 Marotti, I., Bregola, V., Aloisio, I., Di Gioia, D., Bosi, S., Di Silvestro, R., Quinn, R. & Dinelli, G.
523 (2012). Prebiotic effect of soluble fibres from modern and old durum-type wheat varieties on
524 *Lactobacillus* and *Bifidobacterium* strains. *Journal of the Science of Food and Agriculture*,
525 92(10), 2133–2140.

526 Moorthy, I. G., Maran, J. P., Surya, S. M., Naganyashree, S., & Shivamathi, C. S. (2015). Response
527 surface optimization of ultrasound assisted extraction of pectin from pomegranate peel.
528 *International Journal of Biological Macromolecules*, 72, 1323–1328.

529 Negi, P. S., Jayaprakasha, G. K., & Jena, B. S. (2003). Antioxidant and antimutagenic activities of
530 pomegranate peel extracts. *Food Chemistry*, 80(3), 393–397.

531 Normakhtov, R., Rakhmanberdyeva, R. K., & Rakhimov, D. A. (1999). Polysaccharides of the fruit
532 *Punica granatum*. *Chemistry of Natural Compounds*, 35(1), 96–97.

533 Pereira, P. H. F., Oliveira, T. Í. S., Rosa, M. F., Cavalcante, F. L., Moates, G. K., Wellner, N.,
534 Waldron, K.W. & Azeredo, H. M. C. (2016). Pectin extraction from pomegranate peels with
535 citric acid. *International Journal of Biological Macromolecules*, 88, 373–379.

536 Rout, S., & Banerjee, R. (2007). Free radical scavenging, anti-glycation and tyrosinase inhibition
537 properties of a polysaccharide fraction isolated from the rind from *Punica granatum*.
538 *Bioresource Technology*, 98 (16), 3159–3163.

539 Schepetkin, I. A., & Quinn, M. T. (2006). Botanical polysaccharides: Macrophage
540 immunomodulation and therapeutic potential. *International Immunopharmacology*, 6(3), 317–
541 333.

542 Sun, Y., Li, T., Yan, J., & Liu, J. (2010). Technology optimization for polysaccharides (POP)
543 extraction from the fruiting bodies of *Pleurotus ostreatus* by Box-Behnken statistical design.

544 *Carbohydrate Polymers*, 80(1), 242–247.

545 Symonds, E. L., O’Mahony, C., Laphorne, S., O’Mahony, D., Sharry, J. M., O’Mahony, L., &
546 Shanahan, F. (2012). *Bifidobacterium infantis* 35624 protects against salmonella-induced
547 reductions in digestive enzyme activity in mice by attenuation of the host inflammatory
548 response. *Clinical and Translational Gastroenterology*, 3(e15), 1-10

549 Tremaroli, V., & Bäckhed, F. (2012). Functional interactions between the gut microbiota and host
550 metabolism. *Nature*, 489, 242–249.

551 Viuda-Martos, M., Fernández-Lóaez, J., & Pérez-Álvarez, J. A. (2010). Pomegranate and its many
552 functional components as related to human health: a review. *Comprehensive Reviews in Food*
553 *Science and Food Safety*, 9(6), 635–654.

554 Wikiera, A., Mika, M., Starzyńska-Janiszewska, A., & Stodolak, B. (2015). Development of
555 complete hydrolysis of pectins from apple pomace. *Food Chemistry*, 172, 675–680.

556 Yang, J., Martinez, I., Walter, J., Keshavarzian, A., & Rose, D. J. (2013). *In vitro* characterization of
557 the impact of selected dietary fibers on fecal microbiota composition and short chain fatty acid
558 production. *Anaerobe*, 23, 74–81.

559 Zhu, C., & Liu, X. (2013). Optimization of extraction process of crude polysaccharides from
560 Pomegranate peel by response surface methodology. *Carbohydrate Polymers*, 92(2), 1197–
561 1202.

562 Zhu, C. P., Zhai, X. C., Li, L. Q., Wu, X. X., & Li, B. (2015). Response surface optimization of
563 ultrasound-assisted polysaccharides extraction from pomegranate peel. *Food Chemistry*, 177,
564 139–146.

565

566 **Figure Captions**

567 **Figure 1.** Apparent molecular weight distribution by size exclusion chromatography (abscissa) of
568 the CPSP samples from mesocarp and exocarp, data expressed as peak area % of total areas, as
569 mean from triplicate measurements. *W*, *Wonderful*; *L* *Laffan*.

570

571 **Figure 2.** ¹H-NMR spectra in 1mL of D₂O, at room temperature (23±2 °C) for: Laffan mesocarp -
572 CPSP (6.2 mg + 1.3 mg maleic) and Wonderful mesocarp-CPSP (6.05 mg + 1.12 mg maleic acid).
573 *O*-Acetyl, singlet of the acetyl groups; Gal A-OCH₃, singlet of the methoxyl group of galacturonic
574 acid units

575

576 **Figure 3.** Evaluation of prebiotic properties of CPS (5 % in m-MRS medium) from the mesocarp of
577 Laffan and Wonderful cultivars on (a) *L. planctarum* L12 and (b) *Bifidobacterium breve* B632.
578 *C*⁻: growth on m-MRS with no added carbon source; *C*⁺: growth on m-MRS with 0.5% glucose;
579 CFU, colony forming units

580

581 **Table 1.** Applied methods to recover CPS from Laffan mesocarp and corresponding extraction
582 yields (mean values as weight/dried matter); all the determinations were carried out in triplicate
583 except for of **1a** and **1b** methods that were in single.

584 § 24 hrs pretreatment with water before decoction; * 24 hrs pretreatment with 70% ethanol at 25°C.

585

586 **Table 2.** Polysaccharide content in the mesocarp and exocarp of Wonderful and Laffan varieties; the
587 values are a mean of triplicates. * 24 hrs pretreatment with 70% ethanol at 25°C

588

589 **Table 3:** Sugar composition by acidic hydrolysis , *W*, *Wonderful*; *L* , *Laffan*.

590

591

Methods	DM (g)/ solvent (mL)	T (°C)	Time (min)	Yield (%)
1a	1/15	100	30	5
1b	1/15	100	30+30	8
*2a	1/40	100	60	10
*2b	1/40	100	60+ 60	9.8
*2c	1/40	100	60+ 60	9.1
*3a	1/40	25	720	2.0
*3b	1/40	25	1440	3.3
*4a	1/25	100	60	7.8
*\$4b	1/25	100	60	7.2

Table 1. Applied methods to recover CPS from Laffan mesocarp and corresponding extractive yields (*mean values* as w/w DM); *tests carried out in triplicate. \$, *pretreatment with ethanol 70% at 25 °C for 12 h before boiling.*

Extractive methods	Samples	Yield (%)	Yield (%)
		<i>mesocarp</i>	<i>exocarp</i>
2a	Laffan	9.80±0.28	4.47±0.50
	Wonderful	8.0±0.10	4.7±1.15
3b	Laffan	3.7±0.42	1.93±0.23
	Wonderful	3.33±1.15	1.99±0.02
4a	Laffan	7.80±0.28	4.20±0.20
	Wonderful	5.67±0.58	4.13±0.31
4b	Laffan	7.15±0.21	3.93±0.12
	Wonderful	6.70±0.66	4.07±0.31

Table 2. Crude polysaccharides (CPs) content in mesocarp and exocarp of Wonderful and Laffan.

The values are a mean of triplicates and expressed as % on DM.

	Molar %			
Sugars	WM-CPSp	LM-CPSp	WE-CPSp	LE-CPSp
Rhamnose	10.4	7.2	10.8	10.1
Arabinose	4.52	4.04	4.88	4.08
Galactose	5.91	7.31	7.34	7.05
Glucose	14	10.3	11.5	10.9
Xylose	11.2	7.87	9.36	9.3
Fructose	0.41	0.29	0.17	0.2
Galacturonic acid	53.8	63.1	56	58.4

Table 3: Sugar composition of different CPSp samples obtained by acidic hydrolysis




Figure 1.




Figure 2




Figure 3