



Protein S-nitrosylation in photosynthetic organisms: A comprehensive overview with future perspectives / Zaffagnini, Mirko; De Mia, M.; Morisse, S.; Di Giacinto, N.; Marchand, C.H.; Maes, A.; Lemaire, S. D; Trost, PAOLO BERNARDO. In: BIOCHIMICA ET BIOPHYSICA ACTA-PROTEINS AND PROTEOMICS. - ISSN 1570-9639. - ELETTRONICO: 1864:8(2016), pp. 957-966. [10.1016/j.bbapap.2016.02.006]

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Protein S-nitrosylation in photosynthetic organisms: A comprehensive overview with future perspectives

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

*Availability:*

This version is available at: <https://hdl.handle.net/11585/559693> since: 2022-02-11

*Published:*

DOI: <http://doi.org/10.1016/j.bbapap.2016.02.006>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Zaffagnini M., De Mia M., Morisse S., Di Giacinto N., Marchand C.H., Maes A., Lemaire S.D., and Trost P. (2016) Protein S-nitrosylation in photosynthetic organisms: A comprehensive overview with future perspectives. *Biochim Biophys Acta*. 1864(8): 952-66.

The final published version is available online at:  
<https://doi.org/10.1016/j.bbapap.2016.02.006>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

# PROTEIN S-NITROSYLATION IN PHOTOSYNTHETIC ORGANISMS: A COMPREHENSIVE OVERVIEW WITH FUTURE PERSPECTIVES

Zaffagnini<sup>a,1</sup> M., De Mia<sup>b,1</sup> M., Morisse<sup>b</sup> S., Di Giacinto<sup>a</sup> N., Marchand<sup>b</sup> CH., Maes<sup>b</sup> A., Lemaire<sup>b,2</sup> SD., and Trost<sup>a,2</sup> P.

Affiliation:

<sup>a</sup>Laboratory of Plant Redox Biology, Department of Pharmacy and Biotechnology, University of Bologna, 40126 Bologna, Italy.

<sup>b</sup>Sorbonne Universités, UPMC Univ Paris 06, Centre National de la Recherche Scientifique, UMR8226, Laboratoire de Biologie Moléculaire et Cellulaire and des Eucaryotes, Institut de Biologie Physico-Chimique, 75005 Paris, France.

<sup>1</sup>equal contribution

<sup>2</sup>corresponding authors: Prof. Paolo Trost (email: [paolo.trost@unibo.it](mailto:paolo.trost@unibo.it)) and Dr. Stéphane D. Lemaire (email: [stephane.lemaire@ibpc.fr](mailto:stephane.lemaire@ibpc.fr))

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

## ABSTRACT

*Background:* The free radical nitric oxide (NO) and derivative reactive nitrogen species (RNS) play essential roles in cellular redox regulation mainly through protein S-nitrosylation, a redox post-translational modification in which specific cysteines are converted to nitrosothiols.

*Scope of view:* This review aims to discuss the current state of knowledge, as well as future perspectives, regarding protein S-nitrosylation in photosynthetic organisms.

*Major conclusions:* NO, synthesized by plants from different sources (nitrite, arginine), provides directly or indirectly the nitroso moiety of nitrosothiols. Biosynthesis, reactivity and scavenging systems of NO/RNS, determine the NO-based signaling including the rate of protein nitrosylation. Denitrosylation reactions compete with nitrosylation in setting the levels of nitrosylated proteins *in vivo*.

*General significance:* Based on a combination of proteomic, biochemical and genetic approaches, protein nitrosylation is emerging as a pervasive player in cell signaling networks. Specificity of protein nitrosylation and integration among different post-translational modifications are among the major challenges for future experimental studies in the redox biology field.

**KEYWORDS:** cysteine, denitrosylation, nitric oxide, nitrosothiols, redox signaling, S-nitrosylation

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

## 1. INTRODUCTION

Nitric oxide (NO) is a gaseous radical molecule produced in both animals and plants by different biosynthetic systems. Several biological effects are attributed to NO, but in many cases NO itself is not the only responsible of such effects, because NO can react inside the cell with transition metals and other radicals like superoxide and triplet oxygen, giving rise to further nitrogen oxides that are more reactive than NO and are collectively named reactive nitrogen species (RNS). Like reactive oxygen species (ROS), also RNS may be toxic for cells. Toxicity may simply derive from the strong oxidizing character of some RNS such as nitrogen dioxide (N<sub>2</sub>O) or peroxyntirite (ONOO<sup>-</sup>), or it may derive from the capability of other RNS or even metal-NO complexes to indiscriminately transfer an NO moiety to different types of biological molecules that contain a suitable nucleophile. The primary amine of a nucleobase is an example of such suitable nucleophile and DNA can actually be damaged by RNS. In Arabidopsis, NO and derivative RNS have been proposed to restrain root growth via DNA damage-induced cell cycle arrest [1]. Besides DNA, numerous amino acid residues are susceptible to be modified by RNS. Tyrosine, for example, can be converted to 3-nitrotyrosine upon reaction with peroxyntirite. Tyrosine nitration is irreversible in a biological context and is often associated to stress conditions [2-4]. On the other hand, cysteines are the most nucleophilic amino acids and different mechanisms exist by which thiols can be converted into nitrosothiols. Depending on the protein microenvironment, different cysteines show dramatically different reactivity toward RNS or NO, making nitrosylation a specific post-translational modification. Moreover, the nitroso group of nitrosylated cysteines can be removed by common reductants of the cell such as reduced glutathione (GSH) or thioredoxins (TRXs). Thanks to its specificity and reversibility, protein nitrosylation is a post-translational modification with a fundamental role in signaling networks and pervasive biological relevance. Whether NO and RNS trigger signaling cascades through S-nitrosylation of specific cysteines, or they rather damage the cell via an indiscriminate modification of biological molecules, will largely depend on the local concentration of NO that dynamically vary inside the cells as a result of different NO biosynthetic pathways, intricate NO chemistry and a plethora of NO scavenging systems.

Therefore, RNS alone or in combination with ROS play a dual role as toxic and signaling molecules in aerobic organisms. A fundamental concept of redox biology that seems to be even more important in oxygen photosynthetic organisms. In this review, we first present the major mechanisms of NO production/removal and the most representative reactions producing derivative RNS. Subsequently, we discuss the interactions between NO/RNS and protein cysteine residues with special emphasis on the mechanisms of protein nitrosylation/denitrosylation, the methods

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

allowing quantification of protein nitrosothiols and identification of S-nitrosylated targets. Then, we provide some examples of established targets of S-nitrosylation and insights into nitrosylomes in photosynthetic organisms. Finally, future perspectives are presented with special emphasis on the major topics that still remain to be investigated.

## 2. REGULATING REACTIVE NITROGEN SPECIES LEVELS *IN VIVO*

### 2.1 Major Systems of NO production

In animal systems the production of nitric oxide (NO) has been extensively characterized over the last decades. Nowadays it is universally accepted that the main pathway goes through nitric oxide synthase (NOS), a ubiquitous enzyme existing in three isoforms: neuronal NOS (nNOS), endothelial NOS (eNOS), and inducible NOS (iNOS). NOSs are homodimers that catalyze the NADPH-dependent five-electron oxidation of L-arginine to L-citrulline and NO [5, 6]. Beside NOSs, animal cells can also produce NO from nitrite via nitrate reductase (NR), an enzyme that, besides its main reaction in which nitrate is reduced to nitrate by two electrons derived from NADPH, can also catalyze the single-electron reduction of nitrite to NO under anaerobic conditions [7, 8].

In photosynthetic organisms, despite the increasing number of studies on NO since its discovery [9, 10], the sources of NO production are still controversial. Up to now, NO is considered to be mainly produced by two distinct routes, the arginine-dependent and the nitrite-dependent pathways. Paradoxically, the arginine-dependent pathway, involving NOS enzymes, is the most controversial one. Indeed, the direct molecular proof of the existence of NOS in plants is still missing although several studies provided some indirect evidence of the presence of NOS activity [11-14]. For instance, a NOS-like activity was observed in plant extracts [15] and an arginase-deficient mutant with higher levels of arginine was reported to accumulate more NO compared to WT plants [16]. Nevertheless, genetic analyses revealed that the genomes of several photosynthetic organisms do not contain any gene showing significant homology with mammalian NOS, with the notable exception of the alga *Ostreococcus tauri* [17]. In addition, Tun and colleagues [18] showed that an exogenous supply of polyamines (spermine and spermidine) to Arabidopsis seedlings induce immediate release of NO, suggesting that polyamines may be involved in the generation of NO through an Arg-dependent pathway [19]. However, a direct proof of the existence of this pathway is still missing.

Considering the debate around the existence of NOS, the nitrite-dependent pathway is emerging as the main source of NO production in plants [20, 21]. Nitrate reductase (NR), which in plants often prefers NADH as the coenzyme, is the key player of this pathway [22-24]. NR has been first evidenced as a source of NO in Arabidopsis guard cells. Stomata movements are regulated by

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

abscisic acid (ABA) and this hormone is known to interact with NO in many processes and to stimulate its accumulation during stomata regulation [25]. In *Arabidopsis* mutants lacking the two NR isoforms, neither ABA-induced NO production nor ABA-induced stomata closure were observed, providing a first evidence for a role of NR in NO production in plants [22]. At the same time, it has been shown that NR is able to convert, both *in vivo* and *in vitro*, nitrite into NO using saturating NADH as a cofactor [26]. Notably, this enzyme undergoes a regulatory switch based on substrate competition that allows NO production. Nitrate ( $\text{NO}_3^-$ ), which is the high affinity substrate, competes with the low affinity substrate nitrite ( $\text{NO}_2^-$ ). Thus, when  $\text{NO}_2^-$  accumulates due to external signals or stress conditions, nitrate reduction is inhibited and NR preferentially produces NO from nitrite [27]. Nevertheless, nitrite-dependent NO production is not strictly dependent on the activity of NR. In roots, a plasma membrane-bound nitrite/NO reductase (Ni-NOR), distinct from cytosolic NR, has been shown to produce NO in tobacco roots under hypoxic conditions, even though the detailed mechanism is still lacking [28]. Recently, Wei and colleagues observed that *Chlamydomonas* cells deficient for nitrate reductase were able to produce NO from nitrite but the underlying mechanism remains to be elucidated [29].

In addition to these major pathways, several studies described other sources of NO, but their physiological significance is still under debate. For instance, the enzyme xanthine oxidoreductase (XOR) was proposed to constitute a possible additional source of NO [30]. Moreover, other groups suggested non-enzymatic NO sources involving chemical reactions that might occur in specific subcellular compartments such as the apoplast [31] and mitochondria [32] [33].

## 2.2 NO chemistry and production of reactive nitrogen species in a biological context

NO is a free radical with an unpaired electron shared between the nitrogen and the oxygen atom. As a radical, NO is relatively unreactive. It does not react with itself nor it reacts with most biological molecules that have no unpaired electrons, but it tends to combine with other radicals and transition metals [34]. NO is also a small and lipophilic molecule that can easily cross lipid membranes and, thanks to its limited reactivity, is considered relatively long living *in vivo* [35]. With a half life of up to 2 seconds in animal cells, NO may eventually travel over hundreds of microns within and between cells [36, 37]. Under stress conditions, however, NO half life is much shorter because of the prompt reaction with oxygen radicals like superoxide [38] (Figure 1). Although  $\text{H}_2\text{O}_2$  may accumulate in plant cells at much higher concentrations than superoxide, the reaction between NO and  $\text{H}_2\text{O}_2$  is instead prevented by spin forbiddenness [34]. The reactions involving NO inside the cell are extremely important because they give rise to more reactive molecules, particularly nitrogen oxides (e.g.  $\text{NO}_2$ ,  $\text{N}_2\text{O}_3$ ) and peroxyxynitrite ( $\text{ONOO}^-$ ), collectively named reactive nitrogen species (RNS).

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**


These molecules have much higher reactivity than NO and may play a dual role in biological systems, both as signalling molecules and toxic compounds [34, 35, 38-40]

Upon reaction with superoxide, NO forms peroxynitrite (ONOO<sup>-</sup>) (Figure 1; reaction 1) which is not a radical species but nevertheless a strong oxidizing agent and a quite reactive molecule. For instance, tyrosines may react with peroxynitrite and receive a nitrated adduct (NO<sub>2</sub>) in a reaction named tyrosine nitration (Figure 1; reaction 2). Alternatively, peroxynitrite may react with NO generating nitrogen dioxide (NO<sub>2</sub>) and nitrite (NO<sub>2</sub><sup>-</sup>) (Figure 1; reaction 3). Nitrogen dioxide is, like NO, a radical, and can be also produced by the reaction between NO and triplet oxygen (a biradical). However, the reaction: 2NO + O<sub>2</sub> → NO<sub>2</sub> (Figure 1; reaction 4) may be very slow, just because its rate depends on the square of NO concentration (rate = k[NO]<sup>2</sup>[O<sub>2</sub>]). In practical terms, the reaction would proceed very slowly at low NO concentrations and increase exponentially at high NO concentrations. Since both NO and O<sub>2</sub> are lipophilic, this reaction would however proceed much faster [41] within membranes where the substrates may be sequestered at higher concentrations [42].

Whichever the route by which NO<sub>2</sub> is formed, NO<sub>2</sub> may undergo a further radical reaction with NO ending up with dinitrogen trioxide (N<sub>2</sub>O<sub>3</sub>) (Figure 1; reaction 5). Since NO<sub>2</sub> is a good one-electron oxidant, N<sub>2</sub>O<sub>3</sub> has some character of an ionic compound (NO<sup>+</sup>-NO<sub>2</sub><sup>-</sup>). In the presence of a nucleophile, like a thiolate, N<sub>2</sub>O<sub>3</sub> can thus transfer its nitrosonium group (NO<sup>+</sup>) and generate a nitrosothiol (Figure 1; reaction 6).

In spite of the common belief, NO does not react with thiolates which in fact are not radicals. However, nitrosothiols (-SNOs) may readily derive from NO if the thiol is previously oxidized to a thiyl radical. Reactive oxygen species like superoxide or hydroxyl radicals, or RNS like peroxynitrite or NO<sub>2</sub>, may all serve as acceptors of single electrons in thiolate oxidation. Then, thiyl radicals and NO readily combine their unpaired electrons giving rise to nitrosothiols (Figure 1; reaction 7). At difference with the addition of an NO<sup>+</sup> to a thiolate, which is named S-nitrosation (Figure 1; reaction 6), the formation of a nitrosothiol by addition of an NO to a thiyl radical is called S-nitrosylation (Figure 1; reaction 7).

NO may also nitrosylate transition metals including iron centers of proteins such as hemes and, in some cases, iron-sulfur clusters. In these reactions, NO may form irreversible complexes with ferrous iron (Fe<sup>II</sup>) or reversible complexes with ferric iron (Fe<sup>III</sup>). Though unstable, ferric nitrosyl [Fe<sup>III</sup>(NO)] complexes can donate a nitrosonium group to a thiolate (-S<sup>-</sup>) resulting in its S-nitrosation:




*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***



In the special case of hemoglobins (Hb), NO may react with the Fe<sup>II</sup>(O<sub>2</sub>) complex and be rapidly converted into met-hemoglobin (HbFe<sup>III</sup>) and nitrate:



This reaction can be part of a scavenging cycle for NO, once the hemoglobin is reduced back to the HbFe<sup>II</sup> form for further oxygen binding [40].

### 2.3 RNS reactivity and scavenging

The biological activity of NO/RNS is dependent on both spatio-temporal regulation of NO production, NO reactions and NO removal by scavenging systems. Consequently, the systems that control the level of NO and derivative RNS play an important role in regulating the NO-based signaling events.

**Oxygen** – As described in the previous section, NO reacts with oxygen and superoxide (O<sub>2</sub><sup>-</sup>) leading to the formation of nitrogen dioxide (NO<sub>2</sub>) and peroxynitrite (ONOO<sup>-</sup>), respectively. These reactions, besides contributing to modulate NO levels, generate additional RNS that are more reactive than NO itself. While decreasing NO levels, the reaction with oxygen stimulates the formation of protein nitrosothiols via S-nitrosation, and the reaction of NO with superoxide may end up with tyrosine nitration. While reactions of NO with oxygen or superoxide cannot be easily controlled inside the cell, they are predicted to be particularly important in subcellular compartments where the production of oxygen (chloroplasts) and ROS (both chloroplasts and mitochondria) may be massive [43].

**Hemoglobins** – It is well established that NO can form complexes with plant metal-containing proteins and several studies have focused their attention on the interactions between NO and hemoglobin (Hb). In plants, there are three main types of Hb: symbiotic Hb, also named leghemoglobin (Lb), localized in nitrogen-fixing root nodules of leguminous plants; non-symbiotic Hb (nsHb), consisting of two classes based on oxygen affinity (*i.e.* class I has higher affinity for oxygen than class II); and truncated Hb (tHb) [44]. In 1998, Davies and colleagues reported the existence of a complex between ferrous leghemoglobin (LbFe<sup>II</sup>) and NO in intact root nodules or extracts of root nodules [45]. Additional evidence showed that other Lb forms that are produced *in vivo* (*i.e.* oxyLb, LbFeO<sub>2</sub>; and ferrylLb, LbFe<sup>IV</sup>=O), are able to scavenge *in vitro* NO and/or peroxynitrite [46], with concomitant formation of nitrate. The interaction of NO with Hb is not exclusive to Lb. In fact, several

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

studies reported that nsHb from distinct plant species can convert NO into nitrate via an NAD(P)H-dependent reaction [47-49]. The analysis of transgenic plants with altered expression of class-I nsHb sustained the functional role of these proteins in the control of NO bioavailability especially under stress conditions such as hypoxia [48, 50, 51] and pathogen attack [52]. Accordingly, overexpression of class-I nsHb improved plant fitness during hypoxic stress while mutant plants with defective expression of class I nsHb accumulated NO with concomitant decrease of plant growth and enhanced resistance to the hemibiotrophic *P. syringae* and the necrotrophic fungus *Botrytis cinerea*. Altogether, these studies evidenced a strong relation between nsHb proteins and NO suggesting that nsHb can control NO levels with consequent adaptation to stress conditions. Besides Lb and nsHb, also truncated Hb was shown to play a role in NO modulation. In the green alga *Chlamydomonas reinhardtii*, tHb1 was recently shown to bind NO [53] and to modulate NO levels and NR activity [54] while tHb8 was proposed to participate in an NO-dependent signaling pathway required for anaerobic survival [55].

**Glutathione** – An important molecule involved in the control of NO levels *in vivo* is glutathione ( $\gamma$ -L-glutamyl-L-cysteinyl-L-glycine). This molecule is a small tripeptide of 307 Da present in almost all living organisms and widely spread in all cellular compartments [56]. Relatively abundant (1-10 mM), glutathione is considered as the major low molecular weight thiol in cells [57]. It exists in two different redox states: reduced (GSH) and oxidized, the latter bearing a disulfide bridge between two glutathionyl moieties (GSSG). In plant cells, GSH has several antioxidant functions (e.g. regeneration of reduced ascorbate during the ascorbate/glutathione cycle [43]; detoxification of xenobiotics [58] or heavy metals [59], and also participates in NO scavenging by forming nitrosoglutathione (GSNO). The exact mechanism of GSNO formation is still unclear: GSH might be either nitrosated by  $N_2O_3$  or, alternatively, an NO molecule might nitrosylate a glutathionyl radical [60] (see Figure 1). GSNO can be reduced to GSSG and ammonia by the zinc-containing enzyme nitrosoglutathione reductase (GSNOR, for further details see below and paragraph 3.2). Besides modulating NO levels, GSNO is considered as the major reservoir of NO in cells and serves as an important vehicle for NO throughout the cell, enabling NO biological activity to be expanded [61].

**Nitrosoglutathione reductase** – Originally considered as a glutathione-dependent formaldehyde dehydrogenase, GSNO reductase (GSNOR) belongs to the alcohol dehydrogenase protein family III (ADH3 in human) and it was shown to be involved in GSNO catabolism [60, 62-64]. The enzyme catalyzes the NADH-dependent transformation of GSNO into GSSG and ammonia in the presence of GSH [62, 65, 66]. More precisely, GSNO reduction by GSNOR produces an intermediate

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

(glutathione sulfonamide, GSNHOH) that can either react with GSH to produce GSSG and ethanolamine (NH<sub>2</sub>OH) or rearrange and then spontaneously hydrolyze to produce glutathione sulfinate (GSO<sub>2</sub>H) and ammonia [67]. GSNOR is highly conserved between kingdoms and is present in most photosynthetic organisms. In these organisms, GSNOR is generally cytoplasmic and encoded by a single gene, except for few cases known, such as poplar that contains two GSNOR genes [63, 68]. GSNO can act as an NO donor to protein cysteine thiolates to form protein nitrosothiols, and since GSNOR can regulate GSNO cellular homeostasis, it indirectly controls the levels of protein nitrosothiols [62, 69-72]. These functions will be presented with more details in the section dedicated to protein S-nitrosylation and S-nitrosation (see paragraph 3.2).

**Peroxiredoxins** – Peroxiredoxins (PRXs) are small non-heme peroxidases present in diverse compartments and divided into four classes: 2-cys PRX, 1-cys PRX, PRX-Q and type-II PRX [73-75]. Peroxiredoxins catalyze the reduction of H<sub>2</sub>O<sub>2</sub> to water or alkylhydroperoxides (ROOH) to the corresponding alcohol (ROH). PRXs are mainly reduced by thioredoxins but alternative electron donors such as glutathione are possible. Alternatively, PRXs can also scavenge RNS by reducing peroxynitrite into nitrite [76, 77]. PRXs were demonstrated to act as peroxynitrite scavengers under conditions of RNS production in Arabidopsis [78] and bacteria [79].

**Additional RNS scavenging systems** – In addition to the best known RNS-scavenging systems described above, other mechanisms might have a role in the modulation of NO levels in plants. A most likely mechanism involves thioredoxins (TRXs) that are small oxidoreductases known to control the redox state of target proteins through dithiol/disulfide exchange reactions [80, 81]. Besides reducing selected disulfides, dithiolic TRXs can reduce GSNO to GSH and concomitantly release a nitroxyl anion (NO<sup>-</sup>) that immediately gets protonated (HNO). While this reaction is catalyzed by human and plant TRXs *in vitro* [82], there is no clear evidence that TRXs may be involved in GSNO catabolism *in vivo*. Another intriguing mechanism potentially implicated in NO scavenging is protein S-nitrosylation (or S-nitrosation). This thiol-based redox modification involves specific target cysteines on proteins and mainly serves as a signaling mechanism. Being reversible, nitrosylation/nitrosation reactions might contribute to NO/RNS scavenging as long as nitrosothiols formation decreases the intracellular levels of NO and its derivatives. However, this putative function is strictly dependent on the stability of nitrosothiols and may become particularly effective when nitrosylated proteins are subject to denitrosylation reactions that end up with compounds with no nitrosylating activity. This is the case of the GSH-mediated denitrosylation, which forms GSNO which can be reduced (and inactivated) by GSNOR.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

### 3. RNS-PROTEINS INTERACTIONS: METAL LIGATION, TYROSINE NITRATION AND FORMATION OF CYSTEINE NITROSOTHIOLS

In plants, NO and derived RNS are involved in the regulation of several physiological processes [83, 84], and their cellular effects are mediated by three major mechanisms: (1) metal-nitrosylation; (2) tyrosine nitration, and (3) formation of cysteine nitrosothiols, which may occur either by S-nitrosylation or S-nitrosation mechanisms.

NO can directly bind transition metals generating NO-metal complexes [35]. When these metals are located in the prosthetic group of metalloproteins, this reaction can lead to the metal-catalyzed (or metal-assisted) nitrosylation of protein thiols (see 2.2 and 3.1). Among the NO-targeted metalloproteins of animal systems, soluble guanylate cyclase (sGC) is a well-known example [85, 86] together with aconitase [87], hemoglobin and myoglobin [88], NOS and cytochrome P450 [89, 90]. Apart from aconitase, which contains an iron-sulfur cluster, the others are all heme-proteins. Also plants contain heme-proteins that are targeted by NO, including soluble guanylate cyclase [91] besides the already mentioned Hbs [92].

NO and its derivatives also modify specific amino acid residues. Tyrosine is a possible target and the reaction, called nitration, consists in the binding of an NO<sub>2</sub> moiety to the aromatic ring of a tyrosine generally resulting in the modulation of the function of the target protein [93, 94]. The main agent inducing Tyr nitration is peroxynitrite, suggesting that this modification mainly occurs under severe stress conditions (*i.e.* concomitant production of NO and O<sub>2</sub><sup>-</sup>) (Figure 1) [84, 95]. Tyr nitration is implicated in neurodegenerative diseases and cell death pathways and has therefore been extensively studied in mammals [35]. Nevertheless, several studies suggested that this modification may also have important roles in plants [94, 96-101]. At the moment, the number of studies on nitration in plants is limited compared to other post-translational modifications and its role in cell regulation remains unclear because of its irreversibility and limited knowledge on possible underlying mechanisms [35, 84]. Conversely, the formation of cysteine nitrosothiols has been deeply studied in the last years and is considered as the main redox post-translational modification triggered by NO and its derivatives. In animals, it is well-established that it plays a critical role in the control of numerous fundamental processes, regulating the activity, stability and subcellular localization of a large number of target proteins as well as their interactions with DNA/RNA and other protein partners [102-104]. In photosynthetic organisms, formation of protein nitrosothiols has recently emerged as an important mechanism of redox regulation and redox signaling, although much remains to be done [83].

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

### 3.1 Protein S-nitrosothiols: biochemical features and mechanisms of formation

Protein S-nitrosylation consists in the formation of a nitrosothiol (-SNO) after the reversible and covalent linkage of NO to a protein cysteine thiol. The term S-nitrosylation is commonly used as a general term for the formation of S-nitrosothiols, but actually S-nitrosylation should only be applied to the reaction between NO and a thiyl radical, while S-nitrosation describes the reaction between nitrosonium (NO<sup>+</sup>) and a thiolate (Figure 1). However, taking into account the widely accepted use of the term S-nitrosylation as a synonym of "S-nitrosothiol formation", here we will refer to S-nitrosylation whenever the underlying mechanism of S-nitrosothiol formation is not considered, or unknown.

To be suitable for a nitrosylation reaction, cysteine residues must be "reactive", namely in the form of thiyl radicals (-S<sup>•</sup>) or thiolate anions (-S<sup>-</sup>) [105]. The thiyl radical is typically formed under stress conditions (for details see paragraph 2.2) while the thiolate requires a specific protein microenvironment. The thiolate form of a cysteine depends on several factors such as solvent accessibility, the pK<sub>a</sub> of the thiol and the microenvironment surrounding the residue or a combination of these factors [106-109]. The cysteine microenvironment has been also proposed to have an important impact on the sensitivity of a given cysteine to nitrosylation. Notably, the presence of acidic and basic amino acid residues flanking the cysteine is important to favor thiol deprotonation but at the same time they may be involved in the stabilization of the major nitrosylating agent in the cell, GSNO [110]. Based on this property, Stamler and colleagues defined a consensus SNO motif for S-nitrosylation [111]. However, since acidic and basic residues that are in proximity of the reactive Cys in the three-dimensional structure of the protein might not be close in the amino acid sequence, a confident prediction of nitrosylation sites from amino acid sequences was precluded [112].

The pathways leading to the formation of a protein nitrosothiol in a cell are mainly three: direct nitrosylation, metal-mediated nitrosylation and trans-nitrosylation, depending on whether the nitroso group of the SNO derives directly from NO/RNS or is donated by a metal-NO complex or by another nitrosothiol (Figure 2).

**Direct S-nitrosylation** – As mentioned above, direct reactions are represented by two major mechanisms involving the reaction between NO and thiyl radicals (S-nitrosylation *stricto sensu*), and the reaction between dinitrogen trioxide and thiolate anions, where N<sub>2</sub>O<sub>3</sub> acts as a donor of NO<sup>+</sup> (S-nitrosation) (Figure 2, reaction 1). While the former occurs under severe stress conditions that allow the formation of thiyl radicals, the latter typically befalls under physiological conditions, when the redox potential of the cell is not perturbed [113, 114].

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

**Metal-mediated S-nitrosylation** – As already mentioned, metal-mediated nitrosylation involves metal-nitrosyl complexes formed by NO reacting with transition metals of metalloproteins [115] (Figure 2, reaction 2). Once formed, these metal-nitrosyl complexes are responsible for the transfer of NO to a cysteine residue belonging to the same protein (auto-nitrosylation), possibly following an S-nitrosation mechanism as described in part 2.2. To date, the most extensively studied example of an auto-nitrosylated protein is mammalian hemoglobin (Hb) [88] in which NO binding to the heme center allows the protein to auto-catalyze the S-nitrosylation of its Cys93, thereby modulating the enzymatic activity in response to oxygen tension and maximizing oxygen delivery to peripheral tissues.

**Trans-nitrosylation** – Trans-nitrosylation can involve low molecular weight S-nitrosothiols such as GSNO, or nitrosocysteine (CysNO), or a protein-mediated mechanism where an S-nitrosylated protein transfers its NO moiety to another protein [105] (Figure 2, reaction 3). Among the diverse mechanisms of S-nitrosylation, GSNO-mediated trans-nitrosylation is considered the most physiologically relevant. In fact, GSNO is the main NO reservoir and acts as signal transducer, transferring its NO moiety to numerous targets within the cell. Its importance has been firstly clarified in the mouse model, where knocking out GSNOR, an enzyme specifically involved in GSNO breakdown (for further details see paragraph 3.2), results in a high level of nitrosylated proteins, indicating that GSNO is in equilibrium with a pool of SNO-proteins [62, 116]. Similarly, Arabidopsis plants deficient for GSNOR [69, 70] exhibited increased levels of nitrosylated proteins. Conversely, GSNOR overexpression was found to decrease protein nitrosylation levels [71]. Although GSNO-dependent trans-nitrosylation is considered as the main mechanism, a growing interest has recently emerged on protein-mediated trans-nitrosylation. Notably, proteins containing a nitrosylated cysteine may be able to transfer their NO moiety to other proteins. Theoretically, this mechanism may lead to the nitrosylation of proteins that are not themselves target of direct nitrosylation, thereby amplifying the signal. At the moment, there are few examples of trans-nitrosylases in the literature: SNO-hemoglobin, SNO-glyceraldehyde-3-phosphate dehydrogenase, SNO-caspase 3 and SNO-Thioredoxin1 [117-119]. To date, nothing is known in plant systems.

### 3.2 Major mechanisms of denitrosylation

The extent of S-nitrosylation of any given protein depends on the ratio between the rates of nitrosylation and denitrosylation. S-nitrosothiols are generally considered labile entities because of their light- and redox-sensitivity. For example, S-nitrosothiols are often easily reduced by low

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

molecular weight thiols such as reduced glutathione (GSH) [120] (and references therein). However, other S-nitrosothiols may not be easily reverted *in vivo* despite the presence of a high concentrations of reductants in the cell, and seem to be tightly regulated [121]. Proteomic analyses identified a large set of nitrosylated proteins that are relatively stable under normal physiological conditions and are not readily denitrosylated by GSH [122]. Accumulating evidence indicate that specific redox proteins can catalyze the denitrosylation of a subset of nitrosylated proteins thereby contributing to the specificity of S-nitrosylation.

**Non-enzymatic denitrosylation** – The S–NO bond can be broken by the action of intracellular reducing agents such as low molecular weight thiols. Since GSH is present at millimolar concentrations (1-5 mM) in plant cells [123], it is considered to play a major role in the control of SNO homeostasis (Figure 3, reaction 1). However, the reduction of SNOs by GSH is a trans-nitrosylation reaction leading to the formation of GSNO. Consequently, GSH denitrosylates a target protein but the GSNO released could serve as a nitrosylating agent for the same protein or other potential targets. To be effective, the denitrosylating activity of GSH requires the intervention of the enzyme GSNOR, which is the only enzyme able to reduce GSNO without producing NO or RNS (Figure 3, for further details see below). Recently, we provided a clear demonstration of the denitrosylating activity of GSH [124]. In this study, the cytoplasmic GAPDH isoform from *A. thaliana* (AtGAPC1) was effectively denitrosylated by GSH *in vitro*. The nitrosylation state of AtGAPC1 was found under the control of the GSH/GSNO ratio but not affected by GSSG. As a consequence, the nitrosylation state of AtGAPC1 is set by the balance between the GSNO-dependent nitrosylation and GSH-dependent denitrosylation, whichever the redox state of the glutathione pool (GSH/GSSG).

**Enzymatic denitrosylation** – To date, the foremost enzyme that has been clearly demonstrated to play a role in controlling nitrosylation in both animal and plant systems is nitrosogluthione reductase (GSNOR) [68, 121]. Indeed, alteration of GSNOR expression strongly impacts on the level of protein nitrosylation [69-71, 116, 125]. However, this effect is indirect since GSNOR does not display a direct protein-SNO denitrosylase activity but indirectly impact protein nitrosylation by controlling the intracellular GSNO levels [66, 69]. GSNOR catalyzes the reduction of GSNO to GSSG and NH<sub>3</sub> in the presence of GSH (Figure 3, reaction 2). An interesting feature of this enzyme relies in its specificity for NADH unlike all other antioxidant enzymes that use NADPH as the electron donor. GSNOR from different plant sources is totally unable to catalyze GSNO reduction in the presence of NADPH ([66]; authors' personal communication).

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

Another enzymatic pathway involved in SNOs reduction is the TRX system [121, 126]. This system is composed of TRX, thioredoxin reductase (TR) and NADPH and is best known for its role in catalyzing dithiol/disulfide exchange reactions. Different from GSNOR, TRXs can also directly interact with protein SNOs and catalyze protein denitrosylation by means of their active site redox cysteines [120] (Figure 3, reaction 3). Holmgren and colleagues showed that TRX can also reduce GSNO *in vitro*, suggesting that this oxidoreductase might be involved in the control of cell nitrosothiols [127]. The role of TRX as denitrosylase has been clearly demonstrated in human cells, where reduced TRX denitrosylates a subpopulation of mitochondrial caspase-3 constitutively nitrosylated, promoting their activation and consequent apoptosis [128]. The involvement of TRXs as denitrosylating enzymes is not restricted to animal cells and has been recently suggested in the context of the regulation of plant immunity [129]. Two molecular mechanisms have been proposed. In the first [128], the most nucleophilic cysteine of the TRX active site attacks the nitrosylated cysteine of the target protein to form a mixed-disulfide intermediate in which TRX and target are covalently linked and a nitroxyl ( $\text{NO}^-$ ) is released. The mixed-disulfide can then be resolved by the nucleophilic attack of the second TRX cysteine with formation an internal disulfide bond and release of the target protein in the reduced state. In the other possible mechanism [130], the initial nucleophilic attack leads to the nitrosylation of TRX and release of the reduced target. The nitrosylated state of TRX is then resolved by the vicinal cysteine through disulfide bond formation and  $\text{NO}^-$  release (Figure 3). Recently, Spoel and colleagues demonstrated that a plant TRX belonging to the h-type (TRXh5) catalyzes selective NPR1 denitrosylation following the second mechanism in which a nitrosylated TRX intermediate is formed [131]. Both cysteines of the TRXh5 active site were found able to perform the first nucleophilic attack on nitrosylated NPR1. In addition, this study nicely demonstrated that TRXh5 cannot restore the immunity response in the *gsnor1* genetic background, suggesting that the denitrosylation activity of the GSH/GSNOR and the TRXh5 systems are selective and not overlapped.

In summary, denitrosylation reactions are dynamic processes involving reducing agents and/or redox-related enzymes. This process appears to be mainly controlled by two redox systems (*i.e.* GSH and TRX and their associated proteins) that show selectivity towards protein-SNO substrates, allowing specificity of SNOs-dependent signaling.

#### 4. DETECTION OF S-NITROSYLATED PROTEINS AND PROTEOMICS

The quantification of nitrosothiols (-SNOs) and the identification of nitrosylated proteins in biological samples remain a challenge due to the lability of nitrosylated proteins [132]. Indeed, the

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***



stability of SNOs is strongly influenced by several factors including light, metals and reducing agents (e.g. GSH, TRXs, etc., see paragraph 3.2).

#### 4.1 Methods for quantification of protein nitrosothiols

The multiplicity of effects of SNOs has prompted the development of reliable methods to detect these adducts. SNO detection methods can be direct (by which the SNO group is detected as such) or indirect (by which the unstable S–N bond of SNO is broken and either the sulfur or the nitrogen-containing product is captured for detection) [133-135]. Readers interested on direct methods such as mass spectrometry, gold nanoparticles, immuno-histochemical approaches, may find excellent reviews on the subject [133-136]. This review will focus on widely used indirect methods.

The most common indirect method is the Saville method, which is based on the heterolytic cleavage of SNOs by mercury(II) to mercurythiolate and nitrous acid (HNO<sub>2</sub>) [137, 138]. Reaction samples containing protein SNOs are first treated with HgCl<sub>2</sub> in the presence of sulfanilamide (SUF) under acidic conditions. The resulting diazonium salt then reacts with N-(1-naphthyl) ethylenediamine to give an azo dye that can be detected colorimetrically at 540 nm. A correct estimation of SNOs content requires a standard curve generated with low-mass SNOs such as GSNO [139]. While the Saville method is easy to perform and reproducible, the sensitivity of this method is low being in the μM range (0.5-100 μM) [132, 139].

Another mercury-based method consists in the use of the fluorescent probe 2,3-diaminonaphthalene (DAN) or analogous compounds [138-140]. DAN, upon reaction with RNS forms the fluorescent 2,3-naphotriazole [139]. In this assay, SNOs are cleaved by mercury(II) at neutral pH following the addition of DAN under acidic condition and subsequent neutralization, since the DAN triazole fluorescence is highly pH sensitive. The fluorescence is measured with excitation light of 375 nm and emission at 415 nm [139]. Different from the Saville method, the DAN assay exhibits a higher sensitivity, being in the nM range (50 nM-5000 nM).

The combination of these two methods provides a large detection range (*i.e.* from 50 nM to 100 μM SNOs), allowing the determination of the overall concentration of SNO moieties in biological samples [61].

Another representative indirect method is based on chemiluminescence. In this method, NO reacts with ozone to give an excited state of nitrogen dioxide. Upon return to the ground state, nitrogen dioxide emits light, which is then detected by a photomultiplier tube [136, 138].

#### 4.2. Methods of detection of S-nitrosylated proteins and peptides

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

The lability of S-nitrosothiols precludes their direct detection by MALDI-TOF or ESI mass spectrometry since most peptides lose the NO moiety during MS analysis [141, 142]. High throughput proteomic detection of nitrosylated proteins therefore relies on indirect methods mostly based on the replacement of the NO moiety by a more stable tag, such as biotin, followed by an enrichment step. Most studies rely on the Biotin Switch Technique (BST), initially developed in 2001 by Snyder and colleagues [143]. It was the first strategy designed for the detection and identification of S-nitrosylated proteins in a total proteome and it is, by far, the most popular (Figure 4). This method relies on the replacement of the NO moiety of S-nitrosylated cysteine residues by a disulfide-bonded biotin tag in a three step process:

- (1) initial blocking of unmodified cysteine thiols by methyl methanethiosulfonate (MMTS) under denaturing conditions;
- (2) selective conversion of S-nitrosothiols into free thiols by treatment with ascorbate;
- (3) labeling of the nascent thiol groups by thiol/disulfide exchange with the biotinylating reagent N-[6-(biotinamido)hexyl]-3'-(2'-pyridyldithio)-propionamide (Biotin-HPDP).

Steps 2 and 3 are performed simultaneously to allow immediate biotin labeling of nascent thiols. After the biotinylation step, previously S-nitrosylated proteins can be easily detected by immunoblotting with anti-biotin antibodies or purified by avidin-based affinity chromatography and identified by MS after elution with dithiothreitol (DTT) to reduce the disulfide that links the target protein to biotin [144]. Since its invention in 2001, the BST was applied in many biological systems including photosynthetic organisms (see section 5 below). Many variants of the original BST have been proposed. The SNO site identification (SNOSID) approach includes a trypsin digestion step before affinity purification allowing the selective isolation of nitrosylated peptides (Figure 4). Therefore SNOSID allows not only identification of the modified protein but also of the modified cysteine [145]. In the SNO resin-assisted capture (SNO-RAC) strategy, proteins denitrosylated by ascorbate are retained on a Thiopropyl Sepharose 6B thiol-reactive resin and can be eluted by thiol-reducing agents either directly or after on-resin trypsinization for identification of nitrosylated proteins and cysteines respectively [146]. SNORAC appears more sensitive than the BST for high molecular weight proteins and the use of solid phase cysteinyl peptide enrichment, which is well suited for on-resin chemical reactions, may be particularly useful to introduce quantitative labels [146, 147]. Alternate approaches have also been designed based on the direct capture of S-nitrosocysteine residues using organomercury-based chemistry [148], that allows conversion of thiols of formerly S-nitrosylated cysteines to sulfonic acids (+48 Da), using a direct reaction of S-nitrosocysteine residues with gold nanoparticles [149] or using phosphine based direct reaction [134]. Despite promising preliminary results, the methods based on direct capture need further confirmation of their potential

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

for large-scale studies. The BST remains the most reliable approach for large scale proteomic analysis of protein nitrosylation.

Several quantitative BST approaches allowing quantification of nitrosylation levels have also been developed. Quantification is generally performed by combining BST with chemical labeling strategies based on isotope coded probes. Isotope-coded affinity tag (ICAT) allowed monitoring changes in nitrosylation levels in response to cold or salt stress in Arabidopsis [150, 151]. Synthetic isotopic S-nitrosothiol capture reagents (SNOCAP) was used to identify stable nitrosylated proteins that are not reduced in the presence of GSH and are therefore likely reduced through specific denitrosylation pathways [122]. The new isotope-coded cysteine thiol-reactive multiplex reagents cys-TMT [152] and iodo-TMT [153] have been used in combination with BST to investigate nitrosylation in human and murine cells, respectively. Isobaric tag for relative and absolute quantification (iTRAQ) has been used in conjunction with the SNO-RAC method to examine kinetics of intracellular nitrosylation and revealed that denitrosylation is a major determinant of steady-state SNO levels [146]. Besides chemical labeling, quantitative analysis of nitrosylation can be achieved by combining BST with metabolic labeling strategies such as stable isotope labeling with amino acids in cell culture (SILAC) [154, 155].

## 5. ROLES AND TARGETS OF NITROSYLATION IN PLANT PHYSIOLOGY

### 5.1 Targeted studies

Although protein nitrosylation is by far more documented in mammals where it participates in numerous physiological and pathological cellular processes [115, 156], our understanding of the biological importance of nitrosylation in photosynthetic organisms is progressing, as recently extensively reviewed [102, 105, 157]. The major nitrosylated proteins characterized to date in photosynthetic organisms are listed in Table 1 and briefly presented below in their biological context.

Many studies pointed out the importance of nitrosylation in plant immune responses [158-161]. The best studied mechanism implicates the NPR1 (non-expressor of pathogenesis related protein 1) transcription factor and constitutes the most documented example of GSNO-mediated trans-nitrosylation in plants. NPR1 is a key regulator in the signal transduction pathway that leads to systemic acquired resistance in plants [162]. In non-infected cells, NPR1 is located in the cytoplasm, inactive in its oligomeric form stabilized by intermolecular disulfide bonds. Upon redox changes triggered by salicylic acid (SA) produced in response to pathogen infection, NPR1 active monomers are released and translocated into the nucleus where they stimulate the transcriptional activation of defense-related genes through binding to TGA transcription factors, including TGA1 [163, 164]. The

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

reduction of the oligomeric form of NPR1 is catalyzed by TRXs while nitrosylation of Cys156 triggers conformational changes that facilitate oligomerization via disulfide bonding [129]. TRXh5 was recently shown to catalyze the selective denitrosylation of NPR1 ([131]; see 3.2). Moreover, TGA1 was also reported to be nitrosylated by GSNO. Under oxidative conditions, four cysteines contained in TGA1 form intramolecular disulfide bonds locking the protein in an inactive conformation unable to interact with NPR1 [165, 166]. GSNO was found to trigger nitrosylation and glutathionylation of several cysteine residues of TGA1 and to enhance, in the presence of NPR1, the DNA binding activity of TGA1 to the activation-sequence-1 (as-1) element found in the promoter of several defense-related genes [166].

Many studies have reported that a strong interplay between ROS- and NO- (or RNS-) dependent signaling pathways is required to regulate numerous processes, especially responses to biotic and abiotic stresses [49, 167-170]. Modulation of this interplay involves the reciprocal regulation of intracellular ROS and NO (or RNS) levels [169, 171, 172]. During plant immune responses, NO and ROS are considered to act concertedly [171-173]. The regulation of NADPH oxidase constitutes a key example of the control of ROS by NO through nitrosylation [172]. Indeed, in plants, NADPH oxidase is responsible for the generation of ROS during the hypersensitive response [174]. In addition, pathogen infection has been reported to trigger rapid NO synthesis [9, 10]. During the initial response, in the absence of SA, the accumulation of NO facilitates the hypersensitive response [HR]. Then, as the concentration of NO arises, it governs a negative feedback leading to the inactivation of the AtRBOHD NADPH oxidase, the major source of ROS during HR, by nitrosylation of Cys890 [172]. This mechanism was proposed to allow a tight regulation of programmed cell death during the hypersensitive response [102].

Several ROS detoxifying enzymes were also found to be regulated by nitrosylation. For example, Arabidopsis chloroplastic PRXIIIE was found to undergo nitrosylation upon pathogen exposure [175, 176]. Both peroxide and peroxynitrite reductase activities of PRXIIIE were found to be inhibited by nitrosylation of Cys121. This regulation of PRXIIIE was proposed to control the damaging and signaling effects of ONOO<sup>-</sup>. Another good example of ROS/NO crosstalk is cytoplasmic ascorbate peroxidase (APX) that appears to be controlled by NO at multiple levels although contrasting results have been reported in different species. Nitrosylation of tobacco APX was proposed to trigger its ubiquitination and subsequent proteasomal degradation during programmed cell death [177]. By contrast, pea APX1 was activated by nitrosylation of Cys32 and inhibited by tyrosine nitration at Tyr235 [178]. Recently, nitrosylation of Arabidopsis APX1 at Cys32 was shown to increase its activity with hydrogen peroxide and to play an important role during immune responses and abiotic stresses [179]. APX activity was also found to be activated by NO

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

donors in soybean root nodules [180] and auxin control of root growth may act, in part, through denitrosylation of APX1 [181]. While potato dehydroascorbate reductase is inhibited by nitrosylation [182], pea monodehydroascorbate reductase is inhibited by tyrosine nitration [183] and the inhibition of several superoxide dismutases by ONOO<sup>-</sup> was proposed to participate in the control of cell death [184]. Moreover, the nitrosylation of GSNO reductase, that appears to inhibit the enzyme, was suggested to participate in the control of SNO levels [185].

Nitrosylation appears to play an important role in the signaling of many hormones. NO and ROS act in concert together with ABA during stomatal closure [186]. Indeed, NO production induced by ABA, was shown to regulate ABA signaling by nitrosylation of Cys137 of the kinase OST1/SnrK2.6 resulting in inhibition of activity [187]. Nitrosylation of the TIR1 auxin receptor on Cys140 was reported to enhance TIR1 interaction with Aux/IAA proteins, facilitating their degradation and subsequently promoting activation of gene expression [188]. Nitrosylation of the AHP1 at Cys115, a histidine phosphotransfer protein involved in cytokine signaling, represses its phosphorylation and subsequent transfer of the phosphoryl group to downstream ARR response regulators [189].

Other enzymes involved in immune responses have been shown to be regulated by nitrosylation. The carbonic anhydrase SABP3 (salicylic acid binding protein-3), playing an important role in defense responses, was found to be inhibited by nitrosylation of Cys280 [190]. In tobacco, NO is responsible for the inactivation of CDC48, a protein that has an activity associated with the ERAD (ER-associated degradation) pathway, disease resistance, cytokines, cell expansion and differentiation [191-193]. Nitrosylation of the Chaperone like AAA+ ATPase CDC48 occurs on three cysteines (Cys110, Cys526, Cys664) and Cys526, located in the ATP binding site, was found to be responsible for structural changes leading to drastic loss of activity upon nitrosylation [194]. The Arabidopsis Metacaspase-9 (AtMC9) was found to be nitrosylated *in vivo* at catalytic Cys147 and this process was proposed to regulate AtMC9 function in the regulation of PCD [195].

Besides plant immune responses, a number of metabolic enzymes participating in diverse pathways have been also reported to be regulated by nitrosylation in photosynthetic organisms. In most cases nitrosylation was found to inhibit the activity of the modified enzyme. As its mammalian counterpart, Arabidopsis glycolytic GAPDH is inhibited by nitrosylation of its catalytic cysteine [124, 196]. In tobacco, salt stress induced nitrosylation of cytoplasmic GAPDH and activation of the GAPDH-interacting protein NtOSAK (*Nicotiana tabacum* osmotic stress activated protein kinase) [197]. In Arabidopsis, Myb2 DNA binding activity is inhibited by nitrosylation at Cys53 [198, 199] and methionine adenosyltransferase is inhibited by nitrosylation at Cys114 [200]. In Arabidopsis mitochondria, GSNO was found to trigger nitrosylation and glutathionylation of several cysteines of glycine decarboxylase, an enzyme important for glycine catabolism during photorespiration [201]. In

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

addition, ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) also appears to be inhibited by nitrosylation [202, 203].

## 5.2. Proteomic analysis of protein nitrosylation in photosynthetic organisms

In addition to targeted studies, numerous proteomic analyses allowed identification of hundreds of proteins undergoing nitrosylation in a wide range of photosynthetic organisms (reviewed in [114, 204-206]). Different organs and subcellular compartments were analyzed such as peroxisomes [207], mitochondria [201], the nucleus [208] or the apoplast [205]. Several nitrosylomes were also performed on mutants including lines deficient for GSNOR [209] or catalase [71]. While most studies were initially performed adding exogenous NO donors [144, 201, 202], numerous physiological biotic and abiotic stresses have also been investigated including cold stress [151, 203, 205, 210], high light [71], ozone fumigation [211], salinity stress [150, 212, 213], polyamines [214] or in response to diverse pathogens or elicitors [176, 194]. The most recent studies identified 492 proteins and 392 sites in *Chlamydomonas reinhardtii* cells submitted to a 15 minutes GSNO treatment [215] and 926 proteins and 1195 sites in Arabidopsis Col-0 and *gsnor1-3* lines [209]. These studies illustrate the progress made in this type of proteomic analyses. While initial studies focused only on the identification of nitrosylated proteins, nowadays, the nitrosylated cysteines can be identified concomitantly, providing further insights into the exact site of modification. Moreover, the number of identified proteins has increased very significantly lately, mainly because modern mass spectrometers are more sensitive and allow a higher coverage of the proteome. This suggests that low abundant nitrosylated proteins can also be identified. This low level can be either due to a low level of nitrosylation or to the low abundance of the protein. In order to distinguish between these two possibilities and provide further insights into the impact of nitrosylation on the proteome, future studies should include quantitative proteomic approaches to monitor protein abundance and nitrosylation extent (see section 6).

Altogether, these studies suggested the implication of nitrosylation in a myriad of cellular processes and pathways. However, all these proteins identified as nitrosylated by large scale proteomic studies remain putative targets that require further confirmation. Indeed, the nitrosothiols are labile and BST is a very difficult technique with inherent limitations and flaws. A major drawback of BST relies in the identification of false positives due to incomplete blocking or denitrosylation during sample handling that cannot be fully taken into account. The reproducibility of the BST-based proteomic analyses is also unexplored but is expected to be variable due to differences in biological material, growth conditions, experimental design, sample handling, instrument setup and bioinformatic data analysis. Nevertheless despite these considerations, proteomic studies have

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

**When citing, please refer to the published version.**

made an important contribution to the field by revealing that, in photosynthetic organisms, nitrosylation is a widespread post-translational modification that potentially regulates hundreds of proteins involved in a multitude of biological processes. Moreover, one can anticipate that future quantitative proteomic approaches will improve the confidence into data generated by BST-based studies by shedding light on outliers.

## 6. PERSPECTIVES AND FUTURE DIRECTIONS

Considerable progress has been achieved in our understanding of the importance of nitrosylation in photosynthetic organisms. However, although the diversity of the targets is emerging with large scale proteomic analyses, our understanding of the underlying mechanisms and functional roles of protein nitrosylation remains limited. Particular attention should be devoted in the future to explore the determinants of cysteine nitrosylation, the molecular mechanisms controlling nitrosylation levels and the mechanisms of signal integration at the interface between the nitrosylation network and other post-translational modifications and signaling molecules.

### 6.1. Understanding the determinants of cysteine nitrosylation

The ability of a given cysteine to undergo a specific type of redox PTM is governed by multiple factors. Proteomic and biochemical approaches revealed that some cysteines undergo multiple redox modifications, as demonstrated for Cys178 of Chlamydomonas isocitrate lyase [216] and Cys149 of Arabidopsis GAPC1 [217] that both undergo nitrosylation and glutathionylation. In the case of other enzymes, such as the Calvin-Benson cycle enzymes fructose-1,6-bisphosphatase and transketolase, the two modifications occur on distinct cysteine residues [218]. The specificity is primarily linked to the biochemical properties of the cysteine residue that are largely dependent on its microenvironment within the folded protein. This microenvironment can influence the accessibility, the acidity and the nucleophilicity of the residue [107, 108, 219]. The microenvironment of the cysteine is also crucial for the stability of the modification since the additional moieties may trigger local structural changes that will allow accommodating and stabilizing the modification. For example, a cysteine residue can possess the reactivity to undergo nitrosylation but once modified, the presence of a nearby cysteine will favor formation of a disulfide bond that substitutes the initial modification, as recently demonstrated for the yeast Atg4 cysteine protease [220]. Hence, the sites of modification are hardly predictable from the primary structure of proteins as recently underscored [221]. In the future, identification and quantification of numerous sites of modification by proteomics will generate large datasets that should contribute, together with improvements of the 3D-modeling

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

of protein structures, to the development of new algorithms aimed at predicting more accurately redox modification sites.

## 6.2. Molecular mechanisms controlling nitrosylation levels

Besides the properties of the target cysteines, nitrosylation can be influenced by multiple external factors including small molecules and enzymes. Since the level of nitrosylation is dependent on both the forward reactions of nitrosylation and the reverse reactions of denitrosylation, any molecule affecting these reactions will influence the modification state of cysteine residues. Numerous small molecules influence the forward (e.g. NO and its multiple derived species including GSNO) and the reverse (e.g. GSH, cysteine, ascorbate) reactions and alteration of their local concentration or redox state can significantly impact the level of nitrosylation. For example, denitrosylation of Arabidopsis GAPC1 was dependent on the GSH/GSNO ratio [124]. Future studies should be aimed at measuring the *in vivo* dynamics of these small molecules affecting nitrosylation.

Numerous enzymes can also control the level of nitrosylation for a specific set of proteins. Human TRX1 was found to denitrosylate only a subset of nitrosylated proteins [222]. In addition, several enzymes, including TRX and GAPDH, were found to trans-nitrosylate specific target proteins [113, 115]. The specificity of these trans-nitrosylases may allow modification of specific cysteines, even when target cysteines do not appear to be candidates for modification based on their reactivity [125]. These specific sites of modification under the control of diverse enzymes will hardly be predictable and will have to be determined experimentally. Only a few trans-nitrosylating enzymes have been identified recently in mammals. Numerous other activities may remain to be discovered and could constitute an important network of redox signal transduction pathways also in photosynthetic organisms.

## 6.3. Functional role of nitrosylation

Despite the identification of hundreds of putative nitrosylated proteins, the functional role of the modification and its impact on the protein function have only been unraveled for a few of them. Targeted studies will be required to fully understand the functional consequence of nitrosylation for a given protein. However, it should be kept in mind that the nitrosylation of an enzyme involved in a given pathway does not necessarily imply that this pathway is regulated by nitrosylation. Indeed, nitrosylation may have no effect on the activity of the protein, it may only affect a minor pool of the total protein content and nitrosylation of this pool may not be limiting for the pathway flux. Moreover, some nitrosylated proteins may represent moonlighting proteins diverted to new functions unrelated to their known function as shown for GAPDH in mammals [103]. As its mammalian counterpart, plant

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***



GAPDH is nitrosylated and re-localizes to the nucleus under stress conditions but the exact role of nitrosylation remains to be established [196, 223, 224].

#### **6.4 Signal integration: a complex interplay of signaling pathways and molecules**

Despite the complexity of nitrosylation signaling networks and associated regulatory pathways, their contribution to signal transduction and environmental adaptation clearly requires signal integration taking into account multiple other signaling pathways and molecules. In addition to nitrosylation, cysteine residues undergo many other types of post-translational modifications, among which glutathionylation and oxidoreduction of disulfide bonds have been shown to play a major role in redox signaling [225]. These redox modifications are interconnected and influence each other. For example, in *Chlamydomonas reinhardtii*, the targets of glutathionylation, nitrosylation [215] and thioredoxin [226] form a highly complex and interconnected redox network with numerous proteins regulated by several modifications (Figure 5). This complexity is likely required to allow a limited number of simple molecules such as ROS and RNS to play a signaling role. The redox signaling network, which remains largely unexplored, is likely a major component of signal integration and constitutes the molecular signature of the ROS/RNS crosstalk whose importance in cell signaling has been recognized [49, 161, 170]. However, this view (Figure 5) is rather static while the redox network probably involves spatial and temporal regulation of several redox post-translational modifications on hundreds of proteins in a highly dynamic manner. The dynamic nature of the network is most likely playing a central role in the mechanisms of cell signaling.

Although most studies on redox PTMs have been focused on nitrosylation, glutathionylation and oxidoreduction of disulfide bonds, other redox modifications exist and should be considered. These modifications include reversible and non-reversible modifications such as S-sulfhydration, tyrosine nitration, protein carbonylation, sulfenic/sulfonic/sulfinic acids, methionine oxidation and diverse types of S-thiolation like cysteinylolation [2, 108, 225, 227-231]. The importance of these additional redox modifications is emerging and future studies should provide new insights into their functional role and their integration within the redox signaling network. Therefore, understanding the redox network will require quantitative and temporal analysis of multiple redox PTMs. Despite a growing interest during recent years in the analysis of redox PTMs and especially nitrosylation and glutathionylation, these modifications are most exclusively studied independently and qualitatively. Time-resolved quantitative analysis of redox PTMs may also provide insights into the importance of redox PTMs *in vivo* under diverse physiological conditions or in different genetic backgrounds. For example, analysis of mutant lines devoid of key proteins involved directly or indirectly in the regulation of nitrosylation and glutathionylation, such as GSNOR, GR, GRXs TRXs, GAPDH or NO-

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

generating enzymes, will be necessary to understand their role and the importance of the redox PTM they control under basal and stress conditions [72, 158, 232, 233]. Moreover, to date, proteomic analyses were mainly focused on soluble proteins. One can imagine that signal integration can implicate membrane proteins. To unravel these type of proteins, adaptation of current methods targeting redox PTM is necessary and will constitute a challenge for future studies.

In addition, redox modifications are also tightly linked to other signaling networks mediated by non-redox post-translational modifications. For example, in mammals, nitrosylation was reported to interfere with signaling processes mediated by phosphorylation, ubiquitylation, sumoylation, acetylation or palmitoylation [234, 235]. The redox PTM often affects the activity of enzymes controlling these modifications such as kinases and phosphatases in the case of phosphorylation. These observations led to propose that among post-translational modifications that convey cell signals, the breadth of the influence of S-nitrosylation may be comparable with that of phosphorylation and ubiquitylation, where signal crosstalk is established as a central operating principle [234, 236]. Very few studies have addressed this type of crosstalk in plants [227] but it likely plays, as in mammals, an important role in cellular signaling. Besides redox PTMs, the integration of the signal implicates a myriad of other molecules and processes acting at the translational, transcriptional and post transcriptional levels [237]. For example, nitrosylation is clearly linked to signaling pathways controlled by hormones [187, 189] or calcium [238]. A strong effort is therefore required to integrate redox signaling with other signaling pathways and to analyze the cellular responses at multiple levels. The emergence of systems biology approaches will certainly be crucial to rapidly increase our understanding of cell signaling pathways and the mechanisms of adaptation to environmental changes.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

## **ACKNOWLEDGMENTS**

This work was supported by FARB2012 University of Bologna (FARB2012FRAN to MZ), by French National Research Agency (ANR) Grant CYNTHIOL ANR-12-BSV6-0011 (to SDL) and by LABEX DYNAMO ANR-11-LABX-0011 (to SDL).

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

PROTEIN	ORGANISM	TARGET	EFFECT	REFERENCE
NPR1	<i>Arabidopsis thaliana</i>	Cys156	Conformational changes	164, 165
TGA1	<i>Arabidopsis thaliana</i>	Cys260, Cys266	Activation	166, 167
RBOHD	<i>Arabidopsis thaliana</i>	Cys890	Inhibition	172
PRXIIE	<i>Arabidopsis thaliana</i>	Cys121	Inhibition	176
APX1	<i>Nicotiana tabacum</i>	—	Ubiquitination	177
	<i>Pisum sativum</i>	Cys32	Activation	178
	<i>Arabidopsis thaliana</i>	Cys32	Activation	179
DHAR	<i>Solanum tuberosum</i>	Cys20, Cys147	Inhibition	183
GSNOR	<i>Arabidopsis thaliana</i>	—	Inhibition	186
OST1/SnRK2.1	<i>Arabidopsis thaliana</i>	Cys137	Inhibition	188
TIR1	<i>Arabidopsis thaliana</i>	Cys140	Facilitation of protein-protein Interaction	189
AHP1	<i>Arabidopsis thaliana</i>	Cys115	Inhibition	190
SABP3	<i>Arabidopsis thaliana</i>	Cys280	Inhibition	191
CDC48	<i>Nicotiana Tabacum</i>	Cys110, Cys664	Cys526, Inhibition	195
Metacaspase-9 (MC9)	<i>Arabidopsis thaliana</i>	Cys147	Inhibition	196
GAPDH	<i>Arabidopsis thaliana</i>	Cys149	Inhibition	125, 197
	<i>Nicotiana tabacum</i>			198
Myb2	<i>Arabidopsis thaliana</i>	Cys53	Inhibition	199, 200
MAT	<i>Arabidopsis thaliana</i>	Cys1145	Inhibition	201
RubisCO	<i>Arabidopsis thaliana</i>	—	Inhibition	203, 204

**Table 1.** Established nitrosylated proteins in photosynthetic organisms

Protein: abbreviated protein name (see text for further details). Organism: source organism of the protein. Target: Cysteine target residue. Effect: effect of nitrosylation on the protein activity and/or function. Reference: corresponding reference(s) for each protein.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

## FIGURE LEGENDS

**Figure 1. Most relevant reactions in which NO may be involved in a biological context.** All reactions are numbered and explained in the text. Reactions involving only inorganic molecules are indicated by full arrows. Broken arrows represent reactions between a RNS and an amino acid.

**Figure 2. Major mechanisms of protein nitrosothiols formation.** The formation of protein nitrosothiols can occur via three major mechanisms: direct nitrosylation (reaction 1), metal-mediated nitrosylation (reaction 2) or trans-nitrosylation (reaction 3). All reactions are numbered and explained in the text. Abbreviations: GSH, reduced glutathione; GSNO, nitrosoglutathione; Met, metal; Met-NO, metal-nitrosyl complex; PSH, reduced protein; PSNO, nitrosylated protein.

**Figure 3. Major mechanisms of protein denitrosylation.** The reduction of protein nitrosothiols can occur via non-enzymatic mechanisms involving GSH (reaction 1) or via enzymatic mechanisms involving GSNOR (reaction 2, specific reduction of GSNO) or TRX (reaction 3). The reduction of oxidized TRX requires the intervention of NADPH and NTR (reaction 4). Abbreviations: GSH, reduced glutathione; GSNO, nitrosoglutathione; GSNOR, nitrosoglutathione reductase; GSSG, oxidized glutathione; HNO, nitroxyl; NH<sub>3</sub>, ammonia; NTR, NADPH-Thioredoxin reductase; TRX<sub>rd</sub>, reduced thioredoxin; TRX<sub>ox</sub>, oxidized thioredoxin

**Figure 4. Workflow for the analysis of nitrosylated proteins with the BST and SNOSID approaches**

This workflow presents the main steps of the Biotin Switch Technique (BST) [143] and of the SNO Site IDentification approach [145]. In the version of BST presented, the first step consists of blocking with iodoacetamide while MMTS was employed in the original version.

**Figure 5. A complex redox network**

Each ball represents a protein. Data were extracted from large scale studies performed in *Chlamydomonas reinhardtii* on S-nitrosylation [215] S-thiolation/S-glutathionylation [239, 240] and TRX targets [226].

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

## REFERENCES

- [1] S. Bai, M. Li, T. Yao, H. Wang, Y. Zhang, L. Xiao, J. Wang, Z. Zhang, Y. Hu, W. Liu, Y. He, Nitric oxide restrain root growth by DNA damage induced cell cycle arrest in *Arabidopsis thaliana*, *Nitric Oxide*, 26 (2012) 54-60.
- [2] S. Jacques, B. Ghesquiere, F. Van Breusegem, K. Gevaert, Plant proteins under oxidative attack, *Proteomics*, 13 (2013) 932-940.
- [3] S. Saito, A. Yamamoto-Katou, H. Yoshioka, N. Doke, K. Kawakita, Peroxynitrite generation and tyrosine nitration in defense responses in tobacco BY-2 cells, *Plant Cell Physiol*, 47 (2006) 689-697.
- [4] R. Valderrama, F.J. Corpas, A. Carreras, A. Fernandez-Ocana, M. Chaki, F. Luque, M.V. Gomez-Rodriguez, P. Colmenero-Varea, L.A. Del Rio, J.B. Barroso, Nitrosative stress in plants, *FEBS letters*, 581 (2007) 453-461.
- [5] U. Forstermann, W.C. Sessa, Nitric oxide synthases: regulation and function, *Eur Heart J*, 33 (2012) 829-837, 837a-837d.
- [6] A.V. Treuer, D.R. Gonzalez, Nitric oxide synthases, S-nitrosylation and cardiovascular health: from molecular mechanisms to therapeutic opportunities (review), *Molecular medicine reports*, 11 (2015) 1555-1565.
- [7] E.A. Jansson, L. Huang, R. Malkey, M. Govoni, C. Nihlen, A. Olsson, M. Stensdotter, J. Petersson, L. Holm, E. Weitzberg, J.O. Lundberg, A mammalian functional nitrate reductase that regulates nitrite and nitric oxide homeostasis, *Nat Chem Biol*, 4 (2008) 411-417.
- [8] S. Shiva, Mitochondria as metabolizers and targets of nitrite, *Nitric Oxide*, 22 (2010) 64-74.
- [9] M. Delledonne, Y. Xia, R.A. Dixon, C. Lamb, Nitric oxide functions as a signal in plant disease resistance, *Nature*, 394 (1998) 585-588.
- [10] J. Durner, D. Wendehenne, D.F. Klessig, Defense gene induction in tobacco by nitric oxide, cyclic GMP, and cyclic ADP-ribose, *Proc Natl Acad Sci U S A*, 95 (1998) 10328-10333.
- [11] A. Frohlich, J. Durner, The hunt for plant nitric oxide synthase (NOS): is one really needed?, *Plant Sci*, 181 (2011) 401-404.
- [12] J.B. Barroso, F.J. Corpas, A. Carreras, L.M. Sandalio, R. Valderrama, J.M. Palma, J.A. Lupianez, L.A. del Rio, Localization of nitric-oxide synthase in plant peroxisomes, *J Biol Chem*, 274 (1999) 36729-36733.
- [13] F.J. Corpas, J.B. Barroso, A. Carreras, M. Quiros, A.M. Leon, M.C. Romero-Puertas, F.J. Esteban, R. Valderrama, J.M. Palma, L.M. Sandalio, M. Gomez, L.A. del Rio, Cellular and subcellular localization of endogenous nitric oxide in young and senescent pea plants, *Plant Physiol*, 136 (2004) 2722-2733.
- [14] S. Jasid, M. Simontacchi, C.G. Bartoli, S. Puntarulo, Chloroplasts as a nitric oxide cellular source. Effect of reactive nitrogen species on chloroplastic lipids and proteins, *Plant Physiol*, 142 (2006) 1246-1255.
- [15] N.M. Crawford, Mechanisms for nitric oxide synthesis in plants, *J Exp Bot*, 57 (2006) 471-478.
- [16] T. Flores, C.D. Todd, A. Tovar-Mendez, P.K. Dhanoa, N. Correa-Aragunde, M.E. Hoyos, D.M. Brownfield, R.T. Mullen, L. Lamattina, J.C. Polacco, Arginase-negative mutants of *Arabidopsis* exhibit increased nitric oxide signaling in root development, *Plant Physiol*, 147 (2008) 1936-1946.
- [17] N. Foresi, N. Correa-Aragunde, G. Parisi, G. Calo, G. Salerno, L. Lamattina, Characterization of a nitric oxide synthase from the plant kingdom: NO generation from the green alga *Ostreococcus tauri* is light irradiance and growth phase dependent, *Plant Cell*, 22 (2010) 3816-3830.
- [18] N.N. Tun, C. Santa-Catarina, T. Begum, V. Silveira, W. Handro, E.I. Floh, G.F. Scherer, Polyamines induce rapid biosynthesis of nitric oxide (NO) in *Arabidopsis thaliana* seedlings, *Plant Cell Physiol*, 47 (2006) 346-354.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [19] H. Yamasaki, M.F. Cohen, NO signal at the crossroads: polyamine-induced nitric oxide synthesis in plants?, *Trends Plant Sci*, 11 (2006) 522-524.
- [20] L.A. Mur, J. Mandon, S. Persijn, S.M. Cristescu, I.E. Moshkov, G.V. Novikova, M.A. Hall, F.J. Harren, K.H. Hebelstrup, K.J. Gupta, Nitric oxide in plants: an assessment of the current state of knowledge, *AoB PLANTS*, 5 (2013) pls052.
- [21] H. Yamasaki, Y. Sakihama, S. Takahashi, An alternative pathway for nitric oxide production in plants: new features of an old enzyme, *Trends Plant Sci*, 4 (1999) 128-129.
- [22] R. Desikan, R. Griffiths, J. Hancock, S. Neill, A new role for an old enzyme: nitrate reductase-mediated nitric oxide generation is required for abscisic acid-induced stomatal closure in *Arabidopsis thaliana*, *Proc Natl Acad Sci U S A*, 99 (2002) 16314-16318.
- [23] C. Meyer, U.S. Lea, F. Provan, W.M. Kaiser, C. Lillo, Is nitrate reductase a major player in the plant NO (nitric oxide) game?, *Photosynth Res*, 83 (2005) 181-189.
- [24] Y. Morot-Gaudry-Talarmain, P. Rockel, T. Moureaux, I. Quillere, M.T. Leydecker, W.M. Kaiser, J.F. Morot-Gaudry, Nitrite accumulation and nitric oxide emission in relation to cellular signaling in nitrite reductase antisense tobacco, *Planta*, 215 (2002) 708-715.
- [25] J.T. Hancock, S.J. Neill, I.D. Wilson, Nitric oxide and ABA in the control of plant function, *Plant Sci*, 181 (2011) 555-559.
- [26] P. Rockel, F. Strube, A. Rockel, J. Wildt, W.M. Kaiser, Regulation of nitric oxide (NO) production by plant nitrate reductase in vivo and in vitro, *J Exp Bot*, 53 (2002) 103-110.
- [27] K.J. Gupta, A.R. Fernie, W.M. Kaiser, J.T. van Dongen, On the origins of nitric oxide, *Trends Plant Sci*, 16 (2011) 160-168.
- [28] C. Stohr, S. Stremlau, Formation and possible roles of nitric oxide in plant roots, *J Exp Bot*, 57 (2006) 463-470.
- [29] L. Wei, B. Derrien, A. Gautier, L. Houille-Vernes, A. Boulouis, D. Saint-Marcoux, A. Malnoe, F. Rappaport, C. de Vitry, O. Vallon, Y. Choquet, F.A. Wollman, Nitric oxide-triggered remodeling of chloroplast bioenergetics and thylakoid proteins upon nitrogen starvation in *Chlamydomonas reinhardtii*, *Plant Cell*, 26 (2014) 353-372.
- [30] B.L. Wang, X.Y. Tang, L.Y. Cheng, A.Z. Zhang, W.H. Zhang, F.S. Zhang, J.Q. Liu, Y. Cao, D.L. Allan, C.P. Vance, J.B. Shen, Nitric oxide is involved in phosphorus deficiency-induced cluster-root development and citrate exudation in white lupin, *New Phytol*, 187 (2010) 1112-1123.
- [31] P.C. Bethke, M.R. Badger, R.L. Jones, Apoplastic synthesis of nitric oxide by plant tissues, *Plant Cell*, 16 (2004) 332-341.
- [32] E. Planchet, K. Jagadis Gupta, M. Sonoda, W.M. Kaiser, Nitric oxide emission from tobacco leaves and cell suspensions: rate limiting factors and evidence for the involvement of mitochondrial electron transport, *Plant J*, 41 (2005) 732-743.
- [33] S. Rumer, K.J. Gupta, W.M. Kaiser, Plant cells oxidize hydroxylamines to NO, *J Exp Bot*, 60 (2009) 2065-2072.
- [34] T.A. Heinrich, R.S. da Silva, K.M. Miranda, C.H. Switzer, D.A. Wink, J.M. Fukuto, Biological nitric oxide signalling: chemistry and terminology, *Br J Pharmacol*, 169 (2013) 1417-1429.
- [35] D.D. Thomas, J.L. Heinecke, L.A. Ridnour, R.Y. Cheng, A.H. Kesarwala, C.H. Switzer, D.W. McVicar, D.D. Roberts, S. Glynn, J.M. Fukuto, D.A. Wink, K.M. Miranda, Signaling and stress: The redox landscape in NOS2 biology, *Free Radic Biol Med*, 87 (2015) 204-225.
- [36] J.R. Lancaster, Jr., A tutorial on the diffusibility and reactivity of free nitric oxide, *Nitric Oxide*, 1 (1997) 18-30.
- [37] D.D. Thomas, X. Liu, S.P. Kantrow, J.R. Lancaster, Jr., The biological lifetime of nitric oxide: implications for the perivascular dynamics of NO and O<sub>2</sub>, *Proc Natl Acad Sci U S A*, 98 (2001) 355-360.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [38] M.G. Espey, K.M. Miranda, D.D. Thomas, D.A. Wink, Distinction between nitrosating mechanisms within human cells and aqueous solution, *J Biol Chem*, 276 (2001) 30085-30091.
- [39] M.N. Hughes, Chemistry of nitric oxide and related species, *Methods Enzymol*, 436 (2008) 3-19.
- [40] D.D. Thomas, Breathing new life into nitric oxide signaling: A brief overview of the interplay between oxygen and nitric oxide, *Redox biology*, 5 (2015) 225-233.
- [41] X. Liu, M.J. Miller, M.S. Joshi, D.D. Thomas, J.R. Lancaster, Jr., Accelerated reaction of nitric oxide with O<sub>2</sub> within the hydrophobic interior of biological membranes, *Proc Natl Acad Sci U S A*, 95 (1998) 2175-2179.
- [42] I.M. Moller, P.E. Jensen, A. Hansson, Oxidative modifications to cellular components in plants, *Annu Rev Plant Biol*, 58 (2007) 459-481.
- [43] C.H. Foyer, G. Noctor, Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications, *Antioxid Redox Signal*, 11 (2009) 861-905.
- [44] A. Besson-Bard, A. Pugin, D. Wendehenne, New insights into nitric oxide signaling in plants, *Annu Rev Plant Biol*, 59 (2008) 21-39.
- [45] C. Mathieu, S. Moreau, P. Frendo, A. Puppo, M.J. Davies, Direct detection of radicals in intact soybean nodules: presence of nitric oxide-leghemoglobin complexes, *Free Radic Biol Med*, 24 (1998) 1242-1249.
- [46] S. Herold, A. Puppo, Kinetics and mechanistic studies of the reactions of metleghemoglobin, ferrylleghemoglobin, and nitrosylleghemoglobin with reactive nitrogen species, *J Biol Inorg Chem*, 10 (2005) 946-957.
- [47] B.G. Hill, A.N. Higdon, B.P. Dranka, V.M. Darley-Usmar, Regulation of vascular smooth muscle cell bioenergetic function by protein glutathiolation, *Biochim Biophys Acta*, 1797 (2010) 285-295.
- [48] K.H. Hebelstrup, M. van Zanten, J. Mandon, L.A. Voesenek, F.J. Harren, S.M. Cristescu, I.M. Moller, L.A. Mur, Haemoglobin modulates NO emission and hyponasty under hypoxia-related stress in *Arabidopsis thaliana*, *J Exp Bot*, 63 (2012) 5581-5591.
- [49] F. Gross, J. Durner, F. Gaupels, Nitric oxide, antioxidants and prooxidants in plant defence responses, *Front Plant Sci*, 4 (2013) 419.
- [50] C. Dordas, B.B. Hasinoff, A.U. Igamberdiev, N. Manac'h, J. Rivoal, R.D. Hill, Expression of a stress-induced hemoglobin affects NO levels produced by alfalfa root cultures under hypoxic stress, *Plant J*, 35 (2003) 763-770.
- [51] M. Perazzolli, P. Dominici, M.C. Romero-Puertas, E. Zago, J. Zeier, M. Sonoda, C. Lamb, M. Delledonne, *Arabidopsis* nonsymbiotic hemoglobin AHb1 modulates nitric oxide bioactivity, *Plant Cell*, 16 (2004) 2785-2794.
- [52] L.A. Mur, A. Sivakumaran, J. Mandon, S.M. Cristescu, F.J. Harren, K.H. Hebelstrup, Haemoglobin modulates salicylate and jasmonate/ethylene-mediated resistance mechanisms against pathogens, *J Exp Bot*, 63 (2012) 4375-4387.
- [53] E.A. Johnson, S.L. Rice, M.R. Preimesberger, D.B. Nye, L. Gilevicius, B.B. Wenke, J.M. Brown, G.B. Witman, J.T. Lecomte, Characterization of THB1, a *Chlamydomonas reinhardtii* Truncated Hemoglobin: Linkage to Nitrogen Metabolism and Identification of Lysine as the Distal Heme Ligand, *Biochemistry*, (2014).
- [54] E. Sanz-Luque, F. Ocana-Calahorra, A. de Montaigu, A. Chamizo-Ampudia, A. Llamas, A. Galvan, E. Fernandez, THB1, a truncated hemoglobin, modulates nitric oxide levels and nitrate reductase activity, *Plant J*, 81 (2015) 467-479.
- [55] A. Hemschemeier, M. Duner, D. Casero, S.S. Merchant, M. Winkler, T. Happe, Hypoxic survival requires a 2-on-2 hemoglobin in a process involving nitric oxide, *Proc Natl Acad Sci U S A*, 110 (2013) 10854-10859.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***



- [56] C.H. Foyer, G. Noctor, Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria, *Physiol Plant*, 119 (2003) 355-364.
- [57] G. Noctor, A. Mhamdi, S. Chaouch, Y. Han, J. Neukermans, B. Marquez-Garcia, G. Queval, C.H. Foyer, Glutathione in plants: an integrated overview, *Plant Cell Environ*, 35 (2012) 454-484.
- [58] P.G. Board, D. Menon, Glutathione transferases, regulators of cellular metabolism and physiology, *Biochim Biophys Acta*, 1830 (2013) 3267-3288.
- [59] C.S. Cobbett, Phytochelatin biosynthesis and function in heavy-metal detoxification, *Curr Opin Plant Biol*, 3 (2000) 211-216.
- [60] K.A. Broniowska, A.R. Diers, N. Hogg, S-nitrosoglutathione, *Biochim Biophys Acta*, 1830 (2013) 3173-3181.
- [61] F.J. Corpas, J.D. Alche, J.B. Barroso, Current overview of S-nitrosoglutathione (GSNO) in higher plants, *Front Plant Sci*, 4 (2013) 126.
- [62] L. Liu, A. Hausladen, M. Zeng, L. Que, J. Heitman, J.S. Stamler, A metabolic enzyme for S-nitrosothiol conserved from bacteria to humans, *Nature*, 410 (2001) 490-494.
- [63] A. Sakamoto, M. Ueda, H. Morikawa, Arabidopsis glutathione-dependent formaldehyde dehydrogenase is an S-nitrosoglutathione reductase, *FEBS Lett*, 515 (2002) 20-24.
- [64] C.A. Staab, M. Hellgren, J.O. Hoog, Medium- and short-chain dehydrogenase/reductase gene and protein families : Dual functions of alcohol dehydrogenase 3: implications with focus on formaldehyde dehydrogenase and S-nitrosoglutathione reductase activities, *Cell Mol Life Sci*, 65 (2008) 3950-3960.
- [65] J.J. Hedberg, W.J. Griffiths, S.J. Nilsson, J.O. Hoog, Reduction of S-nitrosoglutathione by human alcohol dehydrogenase 3 is an irreversible reaction as analysed by electrospray mass spectrometry, *Eur J Biochem*, 270 (2003) 1249-1256.
- [66] L. Kubienova, D. Kopečný, M. Tylichova, P. Briozzo, J. Skopalova, M. Sebelá, M. Navrátil, R. Tache, L. Luhová, J.B. Barroso, M. Petrivalsky, Structural and functional characterization of a plant S-nitrosoglutathione reductase from *Solanum lycopersicum*, *Biochimie*, 95 (2013) 889-902.
- [67] D.E. Jensen, G.K. Belka, G.C. Du Bois, S-Nitrosoglutathione is a substrate for rat alcohol dehydrogenase class III isoenzyme, *Biochem J*, 331 ( Pt 2) (1998) 659-668.
- [68] S. Xu, D. Guerra, U. Lee, E. Vierling, S-nitrosoglutathione reductases are low-copy number, cysteine-rich proteins in plants that control multiple developmental and defense responses in *Arabidopsis*, *Front Plant Sci*, 4 (2013) 430.
- [69] A. Feechan, E. Kwon, B.W. Yun, Y. Wang, J.A. Pallas, G.J. Loake, A central role for S-nitrosothiols in plant disease resistance, *Proc Natl Acad Sci U S A*, 102 (2005) 8054-8059.
- [70] M. Chaki, R. Valderrama, A.M. Fernandez-Ocana, A. Carreras, M.V. Gomez-Rodriguez, J.R. Pedrajas, J.C. Begara-Morales, B. Sanchez-Calvo, F. Luque, M. Leterrier, F.J. Corpas, J.B. Barroso, Mechanical wounding induces a nitrosative stress by down-regulation of GSNO reductase and an increase in S-nitrosothiols in sunflower (*Helianthus annuus*) seedlings, *J Exp Bot*, 62 (2011) 1803-1813.
- [71] A. Lin, Y. Wang, J. Tang, P. Xue, C. Li, L. Liu, B. Hu, F. Yang, G.J. Loake, C. Chu, Nitric oxide and protein S-nitrosylation are integral to hydrogen peroxide-induced leaf cell death in rice, *Plant Physiol*, 158 (2012) 451-464.
- [72] L. Kubienova, T. Ticha, J. Jahnova, L. Luhová, B. Mieslerova, M. Petrivalsky, Effect of abiotic stress stimuli on S-nitrosoglutathione reductase in plants, *Planta*, 239 (2014) 139-146.
- [73] N. Rouhier, J.P. Jacquot, The plant multigenic family of thiol peroxidases, *Free Radic Biol Med*, 38 (2005) 1413-1421.
- [74] R. Dayer, B.B. Fischer, R.I. Eggen, S.D. Lemaire, The peroxiredoxin and glutathione peroxidase families in *Chlamydomonas reinhardtii*, *Genetics*, 179 (2008) 41-57.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [75] N. Rouhier, C.S. Koh, E. Gelhaye, C. Corbier, F. Favier, C. Didierjean, J.P. Jacquot, Redox based anti-oxidant systems in plants: biochemical and structural analyses, *Biochim Biophys Acta*, 1780 (2008) 1249-1260.
- [76] E. Nogoceke, D.U. Gommel, M. Kiess, H.M. Kalisz, L. Flohe, A unique cascade of oxidoreductases catalyses trypanothione-mediated peroxide metabolism in *Crithidia fasciculata*, *Biol Chem*, 378 (1997) 827-836.
- [77] P.J. Hillas, F.S. del Alba, J. Oyarzabal, A. Wilks, P.R. Ortiz De Montellano, The AhpC and AhpD antioxidant defense system of *Mycobacterium tuberculosis*, *J Biol Chem*, 275 (2000) 18801-18809.
- [78] A. Sakamoto, S. Tsukamoto, H. Yamamoto, M. Ueda-Hashimoto, M. Takahashi, H. Suzuki, H. Morikawa, Functional complementation in yeast reveals a protective role of chloroplast 2-Cys peroxiredoxin against reactive nitrogen species, *Plant J*, 33 (2003) 841-851.
- [79] R. Bryk, P. Griffin, C. Nathan, Peroxynitrite reductase activity of bacterial peroxiredoxins, *Nature*, 407 (2000) 211-215.
- [80] Y. Meyer, C. Belin, V. Delorme-Hinoux, J.P. Reichheld, C. Riondet, Thioredoxin and Glutaredoxin Systems in Plants: Molecular Mechanisms, Crosstalks, and Functional Significance, *Antioxid Redox Signal*, (2012).
- [81] J. Lu, A. Holmgren, The thioredoxin antioxidant system, *Free Radic Biol Med*, 66 (2014) 75-87.
- [82] D. Nikitovic, A. Holmgren, G. Spyrou, Inhibition of AP-1 DNA binding by nitric oxide involving conserved cysteine residues in Jun and Fos, *Biochem Biophys Res Commun*, 242 (1998) 109-112.
- [83] E. Baudouin, J.T. Hancock, Nitric oxide signaling in plants, *Front Plant Sci*, 4 (2013) 553.
- [84] M. Moreau, C. Lindermayr, J. Durner, D.F. Klessig, NO synthesis and signaling in plants--where do we stand?, *Physiol Plant*, 138 (2010) 372-383.
- [85] J.S. Krumenacker, K.A. Hanafy, F. Murad, Regulation of nitric oxide and soluble guanylyl cyclase, *Brain Res Bull*, 62 (2004) 505-515.
- [86] E.R. Derbyshire, M.A. Marletta, Structure and regulation of soluble guanylate cyclase, *Annu Rev Biochem*, 81 (2012) 533-559.
- [87] J.C. Drapier, C. Bouton, Modulation by nitric oxide of metalloprotein regulatory activities, *Bioessays*, 18 (1996) 549-556.
- [88] L. Jia, C. Bonaventura, J. Bonaventura, J.S. Stamler, S-nitrosohaemoglobin: a dynamic activity of blood involved in vascular control, *Nature*, 380 (1996) 221-226.
- [89] O.G. Khatsenko, S.S. Gross, A.B. Rifkind, J.R. Vane, Nitric oxide is a mediator of the decrease in cytochrome P450-dependent metabolism caused by immunostimulants, *Proc Natl Acad Sci U S A*, 90 (1993) 11147-11151.
- [90] J. Stadler, J. Trockfeld, W.A. Schmalix, T. Brill, J.R. Siewert, H. Greim, J. Doehmer, Inhibition of cytochromes P4501A by nitric oxide, *Proc Natl Acad Sci U S A*, 91 (1994) 3559-3563.
- [91] T. Mulaudzi, N. Ludidi, O. Ruzvidzo, M. Morse, N. Hendricks, E. Iwuoha, C. Gehring, Identification of a novel *Arabidopsis thaliana* nitric oxide-binding molecule with guanylate cyclase activity in vitro, *FEBS Lett*, 585 (2011) 2693-2697.
- [92] K.J. Gupta, K.H. Hebelstrup, L.A. Mur, A.U. Igamberdiev, Plant hemoglobins: important players at the crossroads between oxygen and nitric oxide, *FEBS Lett*, 585 (2011) 3843-3849.
- [93] J.M. Souza, G. Peluffo, R. Radi, Protein tyrosine nitration--functional alteration or just a biomarker?, *Free Radic Biol Med*, 45 (2008) 357-366.
- [94] J.C. Begara-Morales, M. Chaki, B. Sanchez-Calvo, C. Mata-Perez, M. Leterrier, J.M. Palma, J.B. Barroso, F.J. Corpas, Protein tyrosine nitration in pea roots during development and senescence, *J Exp Bot*, 64 (2013) 1121-1134.
- [95] L.A. Mur, K.H. Hebelstrup, K.J. Gupta, Striking a balance: does nitrate uptake and metabolism regulate both NO generation and scavenging?, *Front Plant Sci*, 4 (2013) 288.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [96] M. Chaki, R. Valderrama, A.M. Fernandez-Ocana, A. Carreras, M.V. Gomez-Rodriguez, J. Lopez-Jaramillo, J.C. Begara-Morales, B. Sanchez-Calvo, F. Luque, M. Leterrier, F.J. Corpas, J.B. Barroso, High temperature triggers the metabolism of S-nitrosothiols in sunflower mediating a process of nitrosative stress which provokes the inhibition of ferredoxin-NADP reductase by tyrosine nitration, *Plant Cell Environ*, 34 (2011) 1803-1818.
- [97] C. Alvarez, J. Lozano-Juste, L.C. Romero, I. Garcia, C. Gotor, J. Leon, Inhibition of Arabidopsis O-acetylserine(thiol)lyase A1 by tyrosine nitration, *J Biol Chem*, 286 (2011) 578-586.
- [98] M. Chaki, A. Carreras, J. Lopez-Jaramillo, J.C. Begara-Morales, B. Sanchez-Calvo, R. Valderrama, F.J. Corpas, J.B. Barroso, Tyrosine nitration provokes inhibition of sunflower carbonic anhydrase (beta-CA) activity under high temperature stress, *Nitric Oxide*, 29 (2013) 30-33.
- [99] F.J. Corpas, M. Leterrier, J.C. Begara-Morales, R. Valderrama, M. Chaki, J. Lopez-Jaramillo, F. Luque, J.M. Palma, M.N. Padilla, B. Sanchez-Calvo, C. Mata-Perez, J.B. Barroso, Inhibition of peroxisomal hydroxypyruvate reductase (HPR1) by tyrosine nitration, *Biochim Biophys Acta*, 1830 (2013) 4981-4989.
- [100] M. Chaki, P. Alvarez de Morales, C. Ruiz, J.C. Begara-Morales, J.B. Barroso, F.J. Corpas, J.M. Palma, Ripening of pepper (*Capsicum annuum*) fruit is characterized by an enhancement of protein tyrosine nitration, *Ann Bot*, 116 (2015) 637-647.
- [101] M.C. Castillo, J. Lozano-Juste, M. Gonzalez-Guzman, L. Rodriguez, P.L. Rodriguez, J. Leon, Inactivation of PYR/PYL/RCAR ABA receptors by tyrosine nitration may enable rapid inhibition of ABA signaling by nitric oxide in plants, *Sci Signal*, 8 (2015) ra89.
- [102] J. Astier, A. Kulik, E. Koen, A. Besson-Bard, S. Bourque, S. Jeandroz, O. Lamotte, D. Wendehenne, Protein S-nitrosylation: what's going on in plants?, *Free Radic Biol Med*, 53 (2012) 1101-1110.
- [103] M.R. Hara, N. Agrawal, S.F. Kim, M.B. Cascio, M. Fujimuro, Y. Ozeki, M. Takahashi, J.H. Cheah, S.K. Tankou, L.D. Hester, C.D. Ferris, S.D. Hayward, S.H. Snyder, A. Sawa, S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding, *Nat Cell Biol*, 7 (2005) 665-674.
- [104] J. Astier, S. Rasul, E. Koen, H. Manzoor, A. Besson-Bard, O. Lamotte, S. Jeandroz, J. Durner, C. Lindermayr, D. Wendehenne, S-nitrosylation: an emerging post-translational protein modification in plants, *Plant Sci*, 181 (2011) 527-533.
- [105] O. Lamotte, J.B. Bertoldo, A. Besson-Bard, C. Rosnoblet, S. Aime, S. Hichami, H. Terenzi, D. Wendehenne, Protein S-nitrosylation: specificity and identification strategies in plants, *Frontiers in chemistry*, 2 (2014) 114.
- [106] I. Dalle-Donne, R. Rossi, G. Colombo, D. Giustarini, A. Milzani, Protein S-glutathionylation: a regulatory device from bacteria to humans, *Trends Biochem Sci*, 34 (2009) 85-96.
- [107] K.G. Reddie, K.S. Carroll, Expanding the functional diversity of proteins through cysteine oxidation, *Curr Opin Chem Biol*, 12 (2008) 746-754.
- [108] C.C. Winterbourn, M.B. Hampton, Thiol chemistry and specificity in redox signaling, *Free Radic Biol Med*, 45 (2008) 549-561.
- [109] M. Zaffagnini, M. Bedhomme, C.H. Marchand, S. Morisse, P. Trost, S.D. Lemaire, Redox regulation in photosynthetic organisms: focus on glutathionylation, *Antioxid Redox Signal*, 16 (2012) 567-586.
- [110] D.T. Hess, A. Matsumoto, S.O. Kim, H.E. Marshall, J.S. Stamler, Protein S-nitrosylation: purview and parameters, *Nat Rev Mol Cell Biol*, 6 (2005) 150-166.
- [111] J.S. Stamler, E.J. Toone, S.A. Lipton, N.J. Sucher, (S)NO signals: translocation, regulation, and a consensus motif, *Neuron*, 18 (1997) 691-696.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [112] M. Liu, J. Hou, L. Huang, X. Huang, T.H. Heibeck, R. Zhao, L. Pasa-Tolic, R.D. Smith, Y. Li, K. Fu, Z. Zhang, S.H. Hinrichs, S.J. Ding, Site-specific proteomics approach for study protein S-nitrosylation, *Anal Chem*, 82 (2010) 7160-7168.
- [113] B.C. Smith, M.A. Marletta, Mechanisms of S-nitrosothiol formation and selectivity in nitric oxide signaling, *Curr Opin Chem Biol*, 16 (2012) 498-506.
- [114] J. Astier, C. Lindermayr, Nitric oxide-dependent posttranslational modification in plants: an update, *Int J Mol Sci*, 13 (2012) 15193-15208.
- [115] P. Anand, J.S. Stamler, Enzymatic mechanisms regulating protein S-nitrosylation: implications in health and disease, *J Mol Med (Berl)*, 90 (2012) 233-244.
- [116] L. Liu, Y. Yan, M. Zeng, J. Zhang, M.A. Hanes, G. Ahearn, T.J. McMahon, T. Dickfeld, H.E. Marshall, L.G. Que, J.S. Stamler, Essential roles of S-nitrosothiols in vascular homeostasis and endotoxic shock, *Cell*, 116 (2004) 617-628.
- [117] M.D. Kornberg, N. Sen, M.R. Hara, K.R. Juluri, J.V. Nguyen, A.M. Snowman, L. Law, L.D. Hester, S.H. Snyder, GAPDH mediates nitrosylation of nuclear proteins, *Nat Cell Biol*, 12 (2010) 1094-1100.
- [118] T. Nakamura, S. Tu, M.W. Akhtar, C.R. Sunico, S. Okamoto, S.A. Lipton, Aberrant protein s-nitrosylation in neurodegenerative diseases, *Neuron*, 78 (2013) 596-614.
- [119] R. Sengupta, A. Holmgren, Thioredoxin and thioredoxin reductase in relation to reversible s-nitrosylation, *Antioxid Redox Signal*, 18 (2013) 259-269.
- [120] M. Benhar, Nitric oxide and the thioredoxin system: a complex interplay in redox regulation, *Biochim Biophys Acta*, 1850 (2015) 2476-2484.
- [121] M. Benhar, M.T. Forrester, J.S. Stamler, Protein denitrosylation: enzymatic mechanisms and cellular functions, *Nat Rev Mol Cell Biol*, 10 (2009) 721-732.
- [122] J.S. Paige, G. Xu, B. Stancevic, S.R. Jaffrey, Nitrosothiol reactivity profiling identifies S-nitrosylated proteins with unexpected stability, *Chem Biol*, 15 (2008) 1307-1316.
- [123] N. Rouhier, S.D. Lemaire, J.P. Jacquot, The role of glutathione in photosynthetic organisms: emerging functions for glutaredoxins and glutathionylation, *Annu Rev Plant Biol*, 59 (2008) 143-166.
- [124] M. Zaffagnini, S. Morisse, M. Bedhomme, C.H. Marchand, M. Festa, N. Rouhier, S.D. Lemaire, P. Trost, Mechanisms of nitrosylation and denitrosylation of cytoplasmic glyceraldehyde-3-phosphate dehydrogenase from *Arabidopsis thaliana*, *J Biol Chem*, 288 (2013) 22777-22789.
- [125] D. Seth, J.S. Stamler, The SNO-proteome: causation and classifications, *Curr Opin Chem Biol*, 15 (2011) 129-136.
- [126] R. Sengupta, A. Holmgren, The role of thioredoxin in the regulation of cellular processes by S-nitrosylation, *Biochim Biophys Acta*, 1820 (2012) 689-700.
- [127] D. Nikitovic, A. Holmgren, S-nitrosoglutathione is cleaved by the thioredoxin system with liberation of glutathione and redox regulating nitric oxide, *J Biol Chem*, 271 (1996) 19180-19185.
- [128] M. Benhar, M.T. Forrester, D.T. Hess, J.S. Stamler, Regulated protein denitrosylation by cytosolic and mitochondrial thioredoxins, *Science*, 320 (2008) 1050-1054.
- [129] Y. Tada, S.H. Spoel, K. Pajerowska-Mukhtar, Z. Mou, J. Song, C. Wang, J. Zuo, X. Dong, Plant immunity requires conformational changes [corrected] of NPR1 via S-nitrosylation and thioredoxins, *Science*, 321 (2008) 952-956.
- [130] D.A. Stoyanovsky, Y.Y. Tyurina, V.A. Tyurin, D. Anand, D.N. Mandavia, D. Gius, J. Ivanova, B. Pitt, T.R. Billiar, V.E. Kagan, Thioredoxin and lipoic acid catalyze the denitrosation of low molecular weight and protein S-nitrosothiols, *J Am Chem Soc*, 127 (2005) 15815-15823.
- [131] S. Kneeshaw, S. Gelineau, Y. Tada, G.J. Loake, S.H. Spoel, Selective Protein Denitrosylation Activity of Thioredoxin-h5 Modulates Plant Immunity, *Mol Cell*, 56 (2014) 153-162.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [132] E. Bechtold, S.B. King, Chemical methods for the direct detection and labeling of S-nitrosothiols, *Antioxid Redox Signal*, 17 (2012) 981-991.
- [133] M.W. Foster, Methodologies for the characterization, identification and quantification of S-nitrosylated proteins, *Biochim Biophys Acta*, 1820 (2012) 675-683.
- [134] H. Wang, M. Xian, Chemical methods to detect S-nitrosation, *Curr Opin Chem Biol*, 15 (2011) 32-37.
- [135] D. Zhang, I. Macinkovic, N.O. Devarie-Baez, J. Pan, C.M. Park, K.S. Carroll, M.R. Filipovic, M. Xian, Detection of protein S-sulfhydration by a tag-switch technique, *Angew Chem Int Ed Engl*, 53 (2014) 575-581.
- [136] J.B. Mannick, C.M. Schonhoff, Measurement of protein S-nitrosylation during cell signaling, *Methods Enzymol*, 440 (2008) 231-242.
- [137] B. Saville, A scheme for the colorimetric determination of microgram amounts of thiols, *Analyst*, 83 (1958) 670-672.
- [138] N.S. Bryan, M.B. Grisham, Methods to detect nitric oxide and its metabolites in biological samples, *Free Radic Biol Med*, 43 (2007) 645-657.
- [139] D.A. Wink, S. Kim, D. Coffin, J.C. Cook, Y. Vodovotz, D. Chistodoulou, D. Jourdeuil, M.B. Grisham, Detection of S-nitrosothiols by fluorometric and colorimetric methods, *Methods Enzymol*, 301 (1999) 201-211.
- [140] P. Kostka, J.K. Park, Fluorometric detection of S-nitrosothiols, *Methods Enzymol*, 301 (1999) 227-235.
- [141] R. Kaneko, Y. Wada, Decomposition of protein nitrosothiols in matrix-assisted laser desorption/ionization and electrospray ionization mass spectrometry, *J Mass Spectrom*, 38 (2003) 526-530.
- [142] G. Hao, S.S. Gross, Electrospray tandem mass spectrometry analysis of S- and N-nitrosopeptides: facile loss of NO and radical-induced fragmentation, *J Am Soc Mass Spectrom*, 17 (2006) 1725-1730.
- [143] S.R. Jaffrey, H. Erdjument-Bromage, C.D. Ferris, P. Tempst, S.H. Snyder, Protein S-nitrosylation: a physiological signal for neuronal nitric oxide, *Nat Cell Biol*, 3 (2001) 193-197.
- [144] C. Lindermayr, G. Saalbach, J. Durner, Proteomic identification of S-nitrosylated proteins in Arabidopsis, *Plant Physiol*, 137 (2005) 921-930.
- [145] G. Hao, B. Derakhshan, L. Shi, F. Campagne, S.S. Gross, SNOSID, a proteomic method for identification of cysteine S-nitrosylation sites in complex protein mixtures, *Proc Natl Acad Sci U S A*, 103 (2006) 1012-1017.
- [146] M.T. Forrester, J.W. Thompson, M.W. Foster, L. Nogueira, M.A. Moseley, J.S. Stamler, Proteomic analysis of S-nitrosylation and denitrosylation by resin-assisted capture, *Nat Biotechnol*, 27 (2009) 557-559.
- [147] D. Su, A.K. Shukla, B. Chen, J.S. Kim, E. Nakayasu, Y. Qu, U. Aryal, K. Weitz, T.R. Clauss, M.E. Monroe, D.G. Camp, 2nd, D.J. Bigelow, R.D. Smith, R.N. Kulkarni, W.J. Qian, Quantitative site-specific reactivity profiling of S-nitrosylation in mouse skeletal muscle using cysteinyl peptide enrichment coupled with mass spectrometry, *Free Radic Biol Med*, 57 (2013) 68-78.
- [148] P.T. Doulias, J.L. Greene, T.M. Greco, M. Tenopoulou, S.H. Seeholzer, R.L. Dunbrack, H. Ischiropoulos, Structural profiling of endogenous S-nitrosocysteine residues reveals unique features that accommodate diverse mechanisms for protein S-nitrosylation, *Proc Natl Acad Sci U S A*, 107 (2010) 16958-16963.
- [149] A. Faccenda, C.A. Bonham, P.O. Vacratsis, X. Zhang, B. Mutus, Gold nanoparticle enrichment method for identifying S-nitrosylation and S-glutathionylation sites in proteins, *J Am Chem Soc*, 132 (2010) 11392-11394.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [150] A. Fares, M. Rossignol, J.B. Peltier, Proteomics investigation of endogenous S-nitrosylation in Arabidopsis, *Biochem Biophys Res Commun*, 416 (2011) 331-336.
- [151] J. Puyaubert, A. Fares, N. Reze, J.B. Peltier, E. Baudouin, Identification of endogenously S-nitrosylated proteins in Arabidopsis plantlets: effect of cold stress on cysteine nitrosylation level, *Plant Sci*, 215-216 (2014) 150-156.
- [152] C.I. Murray, H. Uhrigshardt, R.N. O'Meally, R.N. Cole, J.E. Van Eyk, Identification and quantification of S-nitrosylation by cysteine reactive tandem mass tag switch assay, *Mol Cell Proteomics*, 11 (2012) M111 013441.
- [153] Z. Qu, F. Meng, R.D. Bomgarden, R.I. Viner, J. Li, J.C. Rogers, J. Cheng, C.M. Greenlief, J. Cui, D.B. Lubahn, G.Y. Sun, Z. Gu, Proteomic quantification and site-mapping of S-nitrosylated proteins using isobaric iodoTMT reagents, *J Proteome Res*, 13 (2014) 3200-3211.
- [154] S.E. Ong, M. Mann, A practical recipe for stable isotope labeling by amino acids in cell culture (SILAC), *Nat Protoc*, 1 (2006) 2650-2660.
- [155] S.E. Ong, B. Blagoev, I. Kratchmarova, D.B. Kristensen, H. Steen, A. Pandey, M. Mann, Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics, *Mol Cell Proteomics*, 1 (2002) 376-386.
- [156] M.W. Foster, D.T. Hess, J.S. Stamler, Protein S-nitrosylation in health and disease: a current perspective, *Trends Mol Med*, 15 (2009) 391-404.
- [157] M. Yu, L. Lamattina, S.H. Spoel, G.J. Loake, Nitric oxide function in plant biology: a redox cue in deconvolution, *New Phytol*, 202 (2014) 1142-1156.
- [158] S.I. Malik, A. Hussain, B.W. Yun, S.H. Spoel, G.J. Loake, GSNOR-mediated de-nitrosylation in the plant defence response, *Plant Sci*, 181 (2011) 540-544.
- [159] M. Yu, B.W. Yun, S.H. Spoel, G.J. Loake, A sleigh ride through the SNO: regulation of plant immune function by protein S-nitrosylation, *Curr Opin Plant Biol*, 15 (2012) 424-430.
- [160] P. Trapet, A. Kulik, O. Lamotte, S. Jeandroz, S. Bourque, V. Nicolas-Frances, C. Rosnoblet, A. Besson-Bard, D. Wendehenne, NO signaling in plant immunity: a tale of messengers, *Phytochemistry*, 112 (2015) 72-79.
- [161] D.E. Frederickson Matika, G.J. Loake, Redox regulation in plant immune function, *Antioxid Redox Signal*, 21 (2014) 1373-1388.
- [162] Z.Q. Fu, X. Dong, Systemic acquired resistance: turning local infection into global defense, *Annu Rev Plant Biol*, 64 (2013) 839-863.
- [163] X. Dong, NPR1, all things considered, *Curr Opin Plant Biol*, 7 (2004) 547-552.
- [164] Z. Mou, W. Fan, X. Dong, Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes, *Cell*, 113 (2003) 935-944.
- [165] C. Despres, C. Chubak, A. Rochon, R. Clark, T. Bethune, D. Desveaux, P.R. Fobert, The Arabidopsis NPR1 disease resistance protein is a novel cofactor that confers redox regulation of DNA binding activity to the basic domain/leucine zipper transcription factor TGA1, *Plant Cell*, 15 (2003) 2181-2191.
- [166] C. Lindermayr, S. Sell, B. Muller, D. Leister, J. Durner, Redox regulation of the NPR1-TGA1 system of Arabidopsis thaliana by nitric oxide, *Plant Cell*, 22 (2010) 2894-2907.
- [167] M. Delledonne, J. Zeier, A. Marocco, C. Lamb, Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease resistance response, *Proc Natl Acad Sci U S A*, 98 (2001) 13454-13459.
- [168] M.D. Keiper, R.I. Grossman, J.A. Hirsch, L. Bolinger, I.L. Ott, L.J. Mannon, C.P. Langlotz, D.L. Kolson, MR identification of white matter abnormalities in multiple sclerosis: a comparison between 1.5 T and 4 T, *AJNR. American journal of neuroradiology*, 19 (1998) 1489-1493.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [169] F. Zaninotto, S. La Camera, A. Polverari, M. Delledonne, Cross talk between reactive nitrogen and oxygen species during the hypersensitive disease resistance response, *Plant Physiol*, 141 (2006) 379-383.
- [170] F. Sevilla, D. Camejo, A. Ortiz-Espin, A. Calderon, J.J. Lazaro, A. Jimenez, The thioredoxin/peroxiredoxin/sulfiredoxin system: current overview on its redox function in plants and regulation by reactive oxygen and nitrogen species, *J Exp Bot*, 66 (2015) 2945-2955.
- [171] D. Wendehenne, Q.M. Gao, A. Kachroo, P. Kachroo, Free radical-mediated systemic immunity in plants, *Curr Opin Plant Biol*, 20 (2014) 127-134.
- [172] B.W. Yun, A. Feechan, M. Yin, N.B. Saidi, T. Le Bihan, M. Yu, J.W. Moore, J.G. Kang, E. Kwon, S.H. Spoel, J.A. Pallas, G.J. Loake, S-nitrosylation of NADPH oxidase regulates cell death in plant immunity, *Nature*, 478 (2011) 264-268.
- [173] C. Scheler, J. Durner, J. Astier, Nitric oxide and reactive oxygen species in plant biotic interactions, *Curr Opin Plant Biol*, 16 (2013) 534-539.
- [174] M.A. Torres, J.L. Dangl, J.D. Jones, Arabidopsis gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense response, *Proc Natl Acad Sci U S A*, 99 (2002) 517-522.
- [175] M.C. Romero-Puertas, M. Laxa, A. Matte, F. Zaninotto, I. Finkemeier, A.M. Jones, M. Perazzolli, E. Vandelle, K.J. Dietz, M. Delledonne, S-nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated tyrosine nitration, *Plant Cell*, 19 (2007) 4120-4130.
- [176] M.C. Romero-Puertas, N. Campostrini, A. Matte, P.G. Righetti, M. Perazzolli, L. Zolla, P. Roepstorff, M. Delledonne, Proteomic analysis of S-nitrosylated proteins in Arabidopsis thaliana undergoing hypersensitive response, *Proteomics*, 8 (2008) 1459-1469.
- [177] M.C. de Pinto, V. Locato, A. Sgobba, C. Romero-Puertas Mdel, C. Gadaleta, M. Delledonne, L. De Gara, S-nitrosylation of ascorbate peroxidase is part of programmed cell death signaling in tobacco Bright Yellow-2 cells, *Plant Physiol*, 163 (2013) 1766-1775.
- [178] J.C. Begara-Morales, B. Sanchez-Calvo, M. Chaki, R. Valderrama, C. Mata-Perez, J. Lopez-Jaramillo, M.N. Padilla, A. Carreras, F.J. Corpas, J.B. Barroso, Dual regulation of cytosolic ascorbate peroxidase (APX) by tyrosine nitration and S-nitrosylation, *J Exp Bot*, 65 (2014) 527-538.
- [179] H. Yang, J. Mu, L. Chen, J. Feng, J. Hu, L. Li, J.M. Zhou, J. Zuo, S-nitrosylation positively regulates ascorbate peroxidase activity during plant stress responses, *Plant Physiol*, 167 (2015) 1604-1615.
- [180] M. Keyster, A. Klein, I. Egbichi, A. Jacobs, N. Ludidi, Nitric oxide increases the enzymatic activity of three ascorbate peroxidase isoforms in soybean root nodules, *Plant Signal Behav*, 6 (2011) 956-961.
- [181] N. Correa-Aragunde, N. Foresi, L. Lamattina, Nitric oxide is a ubiquitous signal for maintaining redox balance in plant cells: regulation of ascorbate peroxidase as a case study, *J Exp Bot*, 66 (2015) 2913-2921.
- [182] H. Kato, D. Takemoto, K. Kawakita, Proteomic analysis of S-nitrosylated proteins in potato plant, *Physiol Plant*, 148 (2013) 371-386.
- [183] J.C. Begara-Morales, B. Sanchez-Calvo, M. Chaki, C. Mata-Perez, R. Valderrama, M.N. Padilla, J. Lopez-Jaramillo, F. Luque, F.J. Corpas, J.B. Barroso, Differential molecular response of monodehydroascorbate reductase and glutathione reductase by nitration and S-nitrosylation, *J Exp Bot*, 66 (2015) 5983-5996.
- [184] C. Holzmeister, F. Gaupels, A. Geerlof, H. Sarioglu, M. Sattler, J. Durner, C. Lindermayr, Differential inhibition of Arabidopsis superoxide dismutases by peroxynitrite-mediated tyrosine nitration, *J Exp Bot*, 66 (2015) 989-999.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

- [185] L. Frungillo, M.J. Skelly, G.J. Loake, S.H. Spoel, I. Salgado, S-nitrosothiols regulate nitric oxide production and storage in plants through the nitrogen assimilation pathway, *Nat Commun*, 5 (2014) 5401.
- [186] J. Bright, R. Desikan, J.T. Hancock, I.S. Weir, S.J. Neill, ABA-induced NO generation and stomatal closure in *Arabidopsis* are dependent on H<sub>2</sub>O<sub>2</sub> synthesis, *Plant J*, 45 (2006) 113-122.
- [187] P. Wang, Y. Du, Y.J. Hou, Y. Zhao, C.C. Hsu, F. Yuan, X. Zhu, W.A. Tao, C.P. Song, J.K. Zhu, Nitric oxide negatively regulates abscisic acid signaling in guard cells by S-nitrosylation of OST1, *Proc Natl Acad Sci U S A*, 112 (2015) 613-618.
- [188] M.C. Terrile, R. Paris, L.I. Calderon-Villalobos, M.J. Iglesias, L. Lamattina, M. Estelle, C.A. Casalongue, Nitric oxide influences auxin signaling through S-nitrosylation of the *Arabidopsis* TRANSPORT INHIBITOR RESPONSE 1 auxin receptor, *Plant J*, 70 (2012) 492-500.
- [189] J. Feng, C. Wang, Q. Chen, H. Chen, B. Ren, X. Li, J. Zuo, S-nitrosylation of phosphotransfer proteins represses cytokinin signaling, *Nat Commun*, 4 (2013) 1529.
- [190] Y.Q. Wang, A. Feechan, B.W. Yun, R. Shafiei, A. Hofmann, P. Taylor, P. Xue, F.Q. Yang, Z.S. Xie, J.A. Pallas, C.C. Chu, G.J. Loake, S-nitrosylation of AtSABP3 antagonizes the expression of plant immunity, *J Biol Chem*, 284 (2009) 2131-2137.
- [191] J. Muller, P. Piffanelli, A. Devoto, M. Miklis, C. Elliott, B. Ortmann, P. Schulze-Lefert, R. Panstruga, Conserved ERAD-like quality control of a plant polytopic membrane protein, *Plant Cell*, 17 (2005) 149-163.
- [192] S. Park, D.M. Rancour, S.Y. Bednarek, In planta analysis of the cell cycle-dependent localization of AtCDC48A and its critical roles in cell division, expansion, and differentiation, *Plant Physiol*, 148 (2008) 246-258.
- [193] J.B. O'Quin, L. Bourassa, D. Zhang, J.M. Shockey, S.K. Gidda, S. Fosnot, K.D. Chapman, R.T. Mullen, J.M. Dyer, Temperature-sensitive post-translational regulation of plant omega-3 fatty-acid desaturases is mediated by the endoplasmic reticulum-associated degradation pathway, *J Biol Chem*, 285 (2010) 21781-21796.
- [194] J. Astier, A. Besson-Bard, O. Lamotte, J. Bertoldo, S. Bourque, H. Terenzi, D. Wendehenne, Nitric oxide inhibits the ATPase activity of the chaperone-like AAA+ ATPase CDC48, a target for S-nitrosylation in cryptogeiin signalling in tobacco cells, *Biochem J*, 447 (2012) 249-260.
- [195] B. Belenghi, M.C. Romero-Puertas, D. Vercammen, A. Brackener, D. Inze, M. Delledonne, F. Van Breusegem, Metacaspase activity of *Arabidopsis thaliana* is regulated by S-nitrosylation of a critical cysteine residue, *J Biol Chem*, 282 (2007) 1352-1358.
- [196] S. Holtgreffe, J. Gohlke, J. Starmann, S. Druce, S. Klocke, B. Altmann, J. Wojtera, C. Lindermayr, R. Scheibe, Regulation of plant cytosolic glyceraldehyde 3-phosphate dehydrogenase isoforms by thiol modifications, *Physiol Plant*, 133 (2008) 211-228.
- [197] I. Wawer, M. Bucholc, J. Astier, A. Anielska-Mazur, J. Dahan, A. Kulik, A. Wyslouch-Cieszynska, M. Zareba-Koziol, E. Krzywinska, M. Dadlez, G. Dobrowolska, D. Wendehenne, Regulation of *Nicotiana tabacum* osmotic stress-activated protein kinase and its cellular partner GAPDH by nitric oxide in response to salinity, *Biochem J*, 429 (2010) 73-83.
- [198] V. Serpa, J. Vernal, L. Lamattina, E. Grotewold, R. Cassia, H. Terenzi, Inhibition of AtMYB2 DNA-binding by nitric oxide involves cysteine S-nitrosylation, *Biochem Biophys Res Commun*, 361 (2007) 1048-1053.
- [199] C.P. Tavares, J. Vernal, R.A. Delena, L. Lamattina, R. Cassia, H. Terenzi, S-nitrosylation influences the structure and DNA binding activity of AtMYB30 transcription factor from *Arabidopsis thaliana*, *Biochim Biophys Acta*, 1844 (2014) 810-817.
- [200] C. Lindermayr, G. Saalbach, G. Bahnweg, J. Durner, Differential inhibition of *Arabidopsis* methionine adenosyltransferases by protein S-nitrosylation, *J Biol Chem*, 281 (2006) 4285-4291.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***



- [201] M.C. Palmieri, C. Lindermayr, H. Bauwe, C. Steinhauser, J. Durner, Regulation of plant glycine decarboxylase by s-nitrosylation and glutathionylation, *Plant Physiol*, 152 (2010) 1514-1528.
- [202] J.K. Abat, A.K. Mattoo, R. Deswal, S-nitrosylated proteins of a medicinal CAM plant *Kalanchoe pinnata*- ribulose-1,5-bisphosphate carboxylase/oxygenase activity targeted for inhibition, *FEBS J*, 275 (2008) 2862-2872.
- [203] J.K. Abat, R. Deswal, Differential modulation of S-nitrosoproteome of *Brassica juncea* by low temperature: change in S-nitrosylation of Rubisco is responsible for the inactivation of its carboxylase activity, *Proteomics*, 9 (2009) 4368-4380.
- [204] J. Puyaubert, E. Baudouin, New clues for a cold case: nitric oxide response to low temperature, *Plant Cell Environ*, 37 (2014) 2623-2630.
- [205] A. Sehwat, R. Deswal, S-nitrosylation analysis in *Brassica juncea* apoplast highlights the importance of nitric oxide in cold-stress signaling, *J Proteome Res*, 13 (2014) 2599-2619.
- [206] I. Kovacs, C. Lindermayr, Nitric oxide-based protein modification: formation and site-specificity of protein S-nitrosylation, *Front Plant Sci*, 4 (2013) 137.
- [207] A.P. Ortega-Galisteo, M. Rodriguez-Serrano, D.M. Pazmino, D.K. Gupta, L.M. Sandalio, M.C. Romero-Puertas, S-Nitrosylated proteins in pea (*Pisum sativum* L.) leaf peroxisomes: changes under abiotic stress, *J Exp Bot*, 63 (2012) 2089-2103.
- [208] M. Chaki, A. Shekariesfahlan, A. Ageeva, A. Mengel, C. von Toerne, J. Durner, C. Lindermayr, Identification of nuclear target proteins for S-nitrosylation in pathogen-treated *Arabidopsis thaliana* cell cultures, *Plant Sci*, 238 (2015) 115-126.
- [209] J. Hu, X. Huang, L. Chen, X. Sun, C. Lu, L. Zhang, Y. Wang, J. Zuo, Site-specific nitrosoproteomic identification of endogenously S-nitrosylated proteins in *Arabidopsis*, *Plant Physiol*, 167 (2015) 1731-1746.
- [210] A. Sehwat, J.K. Abat, R. Deswal, RuBisCO depletion improved proteome coverage of cold responsive S-nitrosylated targets in *Brassica juncea*, *Front Plant Sci*, 4 (2013) 342.
- [211] E. Vanzo, A. Ghirardo, J. Merl-Pham, C. Lindermayr, W. Heller, S.M. Hauck, J. Durner, J.P. Schnitzler, S-nitroso-proteome in poplar leaves in response to acute ozone stress, *PLoS One*, 9 (2014) e106886.
- [212] G. Tanou, P. Filippou, M. Belghazi, D. Job, G. Diamantidis, V. Fotopoulos, A. Molassiotis, Oxidative and nitrosative-based signaling and associated post-translational modifications orchestrate the acclimation of citrus plants to salinity stress, *Plant J*, 72 (2012) 585-599.
- [213] D. Camejo, C. Romero-Puertas Mdel, M. Rodriguez-Serrano, L.M. Sandalio, J.J. Lazaro, A. Jimenez, F. Sevilla, Salinity-induced changes in S-nitrosylation of pea mitochondrial proteins, *J Proteomics*, 79 (2013) 87-99.
- [214] G. Tanou, V. Ziogas, M. Belghazi, A. Christou, P. Filippou, D. Job, V. Fotopoulos, A. Molassiotis, Polyamines reprogram oxidative and nitrosative status and the proteome of citrus plants exposed to salinity stress, *Plant Cell Environ*, 37 (2014) 864-885.
- [215] S. Morisse, M. Zaffagnini, X.H. Gao, S.D. Lemaire, C.H. Marchand, Insight into Protein S-nitrosylation in *Chlamydomonas reinhardtii*, *Antioxid Redox Signal*, 21 (2014) 1271-1284.
- [216] M. Bedhomme, M. Zaffagnini, C.H. Marchand, X.H. Gao, M. Moslonka-Lefebvre, L. Michelet, P. Decottignies, S.D. Lemaire, Regulation by glutathionylation of isocitrate lyase from *Chlamydomonas reinhardtii*, *J Biol Chem*, 284 (2009) 36282-36291.
- [217] M. Bedhomme, M. Adamo, C.H. Marchand, J. Couturier, N. Rouhier, S.D. Lemaire, M. Zaffagnini, P. Trost, Glutathionylation of cytosolic glyceraldehyde-3-phosphate dehydrogenase from the model plant *Arabidopsis thaliana* is reversed by both glutaredoxins and thioredoxins in vitro, *Biochem J*, 445 (2012) 337-347.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

- [218] L. Michelet, M. Zaffagnini, S. Morisse, F. Sparla, M.E. Pérez-Pérez, F. Francia, A. Danon, C.H. Marchand, S. Fermani, P. Trost, S.D. Lemaire, Redox regulation of the Calvin-Benson cycle: something old, something new, *Front Plant Sci*, 4 (2013) 470.
- [219] M. Zaffagnini, M. Bedhomme, S.D. Lemaire, P. Trost, The emerging roles of protein glutathionylation in chloroplasts, *Plant Sci*, 185-186 (2012) 86-96.
- [220] M.E. Perez-Perez, M. Zaffagnini, C.H. Marchand, J.L. Crespo, S.D. Lemaire, The yeast autophagy protease Atg4 is regulated by thioredoxin, *Autophagy*, 10 (2014) 1953-1964.
- [221] M. Chaki, I. Kovacs, M. Spannagl, C. Lindermayr, Computational prediction of candidate proteins for S-nitrosylation in *Arabidopsis thaliana*, *PLoS One*, 9 (2014) e110232.
- [222] M. Benhar, J.W. Thompson, M.A. Moseley, J.S. Stamler, Identification of S-nitrosylated targets of thioredoxin using a quantitative proteomic approach, *Biochemistry*, 49 (2010) 6963-6969.
- [223] M. Zaffagnini, S. Morisse, M. Bedhomme, C.H. Marchand, M. Festa, N. Rouhier, S.D. Lemaire, P. Trost, Mechanism of nitrosylation and denitrosylation of cytosolic glyceraldehyde-3-phosphate dehydrogenase from *Arabidopsis*, *J Biol Chem*, 288 (2013) 22777-22789.
- [224] M. Vescovi, M. Zaffagnini, M. Festa, P. Trost, F. Lo Schiavo, A. Costa, Nuclear accumulation of cytosolic glyceraldehyde-3-phosphate dehydrogenase in cadmium-stressed *Arabidopsis* roots, *Plant Physiol*, 162 (2013) 333-346.
- [225] C.E. Paulsen, K.S. Carroll, Cysteine-mediated redox signaling: chemistry, biology, and tools for discovery, *Chemical reviews*, 113 (2013) 4633-4679.
- [226] S.D. Lemaire, B. Guillon, P. Le Maréchal, E. Keryer, M. Miginiac-Maslow, P. Decottignies, New thioredoxin targets in the unicellular photosynthetic eukaryote *Chlamydomonas reinhardtii*, *Proc Natl Acad Sci U S A*, 101 (2004) 7475-7480.
- [227] I. Lounifi, E. Arc, A. Molassiotis, D. Job, L. Rajjou, G. Tanou, Interplay between protein carbonylation and nitrosylation in plants, *Proteomics*, 13 (2013) 568-578.
- [228] B.D. Paul, S.H. Snyder, H(2)S signalling through protein sulfhydration and beyond, *Nat Rev Mol Cell Biol*, 13 (2012) 499-507.
- [229] T. Finkel, Signal transduction by reactive oxygen species, *J Cell Biol*, 194 (2011) 7-15.
- [230] S.B. Wall, J.Y. Oh, A.R. Diers, A. Landar, Oxidative modification of proteins: an emerging mechanism of cell signaling, *Front Physiol*, 3 (2012) 369.
- [231] M. Lo Conte, K.S. Carroll, The redox biochemistry of protein sulfenylation and sulfinylation, *J Biol Chem*, 288 (2013) 26480-26488.
- [232] S.S. Gill, M. Hasanuzzaman, K. Nahar, A. Macovei, N. Tuteja, Importance of nitric oxide in cadmium stress tolerance in crop plants, *Plant Physiol Biochem*, 63 (2013) 254-261.
- [233] A. Mhamdi, J. Hager, S. Chaouch, G. Queval, Y. Han, L. Tacconnat, P. Saindrenan, H. Gouia, E. Issakidis-Bourguet, J.P. Renou, G. Noctor, *Arabidopsis* GLUTATHIONE REDUCTASE1 plays a crucial role in leaf responses to intracellular hydrogen peroxide and in ensuring appropriate gene expression through both salicylic acid and jasmonic acid signaling pathways, *Plant Physiol*, 153 (2010) 1144-1160.
- [234] D.T. Hess, J.S. Stamler, Regulation by S-nitrosylation of protein post-translational modification, *J Biol Chem*, 287 (2012) 4411-4418.
- [235] J.M. Held, B.W. Gibson, Regulatory control or oxidative damage? Proteomic approaches to interrogate the role of cysteine oxidation status in biological processes, *Mol Cell Proteomics*, 11 (2012) R111 013037.
- [236] C. Lopez-Otin, T. Hunter, The regulatory crosstalk between kinases and proteases in cancer, *Nature reviews. Cancer*, 10 (2010) 278-292.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

- [237] G. Mastrobuoni, S. Irgang, M. Pietzke, H.E. Assmus, M. Wenzel, W.X. Schulze, S. Kempa, Proteome dynamics and early salt stress response of the photosynthetic organism *Chlamydomonas reinhardtii*, *BMC Genomics*, 13 (2012) 215.
- [238] C.J. Wientjes, P. Grossman, A.W. Gaillard, Influence of drive and timing mechanisms on breathing pattern and ventilation during mental task performance, *Biological psychology*, 49 (1998) 53-70.
- [239] M. Zaffagnini, M. Bedhomme, H. Groni, C.H. Marchand, C. Puppo, B. Gontero, C. Cassier-Chauvat, P. Decottignies, S.D. Lemaire, Glutathionylation in the photosynthetic model organism *Chlamydomonas reinhardtii*: a proteomic survey, *Mol Cell Proteomics*, 11 (2012) M111 014142.
- [240] L. Michelet, M. Zaffagnini, H. Vanacker, P. Le Marechal, C. Marchand, M. Schroda, S.D. Lemaire, P. Decottignies, In vivo targets of S-thiolation in *Chlamydomonas reinhardtii*, *J Biol Chem*, 283 (2008) 21571-21578.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**