

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

The Dynamic-Demand Joint Replenishment Problem with Approximated Transportation Costs

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

The Dynamic-Demand Joint Replenishment Problem with Approximated Transportation Costs / Baller, Annelieke C.; Dabia, Said; Dullaert, Wout E.H.; Vigo, Daniele. - In: EUROPEAN JOURNAL OF OPERATIONAL RESEARCH. - ISSN 0377-2217. - STAMPA. - 276:3(2019), pp. S0377221719301055.1013-S0377221719301055.1033. [10.1016/j.ejor.2019.01.070]

Availability:

This version is available at: https://hdl.handle.net/11585/720171 since: 2020-02-01

Published:

DOI: http://doi.org/10.1016/j.ejor.2019.01.070

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version. This is the final peer-reviewed accepted manuscript of:

Annelieke C. Baller, Said Dabia, Wout E.H. Dullaert, Daniele Vigo, **The Dynamic-Demand Joint Replenishment Problem with Approximated Transportation Costs**, European Journal of Operational Research, Volume 276, Issue 3, 2019, Pages 1013-1033, ISSN 0377-2217

The final published version is available online at: https://doi.org/10.1016/j.ejor.2019.01.070

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

The Dynamic-Demand Joint Replenishment Problem with Approximated Transportation Costs

Annelieke C. Baller, Said Dabia, Wout E.H. Dullaert

Department of Information, Logistics and Innovation, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands, a.c.baller@vu.nl, s.dabia@vu.nl, w.e.h.dullaert@vu.nl

Daniele Vigo

Department of Electrical, Electronic and Information Engineering "Guglielmo Marconi", Alma Mater Università di Bologna, 04136 Bologna, Italy, daniele.vigo@unibo.it

Department of Information, Logistics and Innovation, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands

In a vendor-managed inventory setting, a supplier determines the timing and size of replenishments for its customers. In the Dynamic-Demand Joint Replenishment Problem (DJRP), one assumes that the supplier pays a fixed fee for replenishing a customer which often occurs if the supplier outsources transportation. Hence, there is no incentive for the supplier to schedule replenishments for nearby customers in the same period. This results in higher transportation costs for the carrier, decreased vehicle utilization and increased future fees for the supplier. To lower costs for both parties, this paper extends the traditional DJRP to the DJRP with Approximated Transportation Costs (DJRP-AT) by taking transportation considerations into account. Since routing problems are difficult to solve and it is not necessary to know the sequence of the deliveries to the customers as these are outsourced, the transportation costs for a given set of customers are approximated using classical schemes. A solution approach for the DJRP-AT based on Branch-and-Cut-and-Price is validated using test instances from the literature. Results show improvements of 4% on average and up to 14.4% for individual instances compared with the DJRP. Moreover, when the DJRP-AT is compared with the DJRP on instances derived from a real-life case, similar savings are obtained. Comparing the DJRP-AT to an equivalent problem with actual routing costs, the solution values of the DJRP-AT are on average only 0.77% higher showing the value of the approximation.

Key words: logistics; joint replenishment; transportation costs; branch-and-cut-and-price

1. Introduction

During the last decades, Vendor-Managed Inventory (VMI) systems have received a lot of attention in the literature (Andersson et al. 2010). In such a system, a supplier manages the inventory of its customers and arranges the transportation of the replenishments. The supplier bears both the inventory holding and transportation costs and therefore strives to minimize these costs by optimizing inventory and shipping decisions. If the supplier would decide on the replenishments and the routes to deliver the replenishments, the supplier faces a problem known as the Inventory Routing Problem (IRP) (Coelho et al. 2014). However, the transport of the replenishments is often outsourced to a Logistics Service Provider (LSP). As a consequence, the supplier pays a fixed transportation fee for a delivery that is specified in a long-term contract. The supplier therefore faces an optimization problem known as the Joint Replenishment Problem (JRP) (Khouja and Goyal 2008). When customer demand varies over time, this problem is known as the Dynamic-Demand Joint Replenishment Problem (DJRP). The DJRP decides which products to order or customers to serve in which periods of the planning horizon such that demand is satisfied at minimal inventory holding and servicing costs. More specifically, the cost of servicing a group of customers in a given period consists of two components, the first of which is a common set-up cost per period if at least one customer is served in that given period. The second component is a cost for each replenished customer. Because of the common set-up cost per period it can be beneficial for the supplier to have some customers replenished together with other customers even before stock runs low. In that case the inventory holding cost is higher because more inventory is kept, but it allows the supplier to save the fixed fee for a period.

The DJRP encompasses a number of key problem features occurring in real-life applications, but also suffers from a number of drawbacks. First, because the actual transportation costs are not directly considered, the DJRP cannot identify closely situated customers that would be wellsuited for joint replenishment. Hence, customers served in one period may be randomly located in a region, resulting in high actual transportation costs for the LSP. Also, the number of replenishments that the LSP is able to perform on behalf of the supplier can be lower due to large distances between deliveries. Therefore, consideration of proximity of customers could result in better utilization of transportation resources, decrease actual transportation costs for the LSP and eventually decrease transportation fees for the supplier. In short, based on the DJRP, the supplier generates requests that are expensive or hard to fulfill. Second, the DJRP ignores duration constraints. Vehicle capacity constraints have been considered in the JRP literature (see e.g. Anily and Tzur (2005)) and in many routing problems, including most studies on the IRP. However, tour duration constraints have proven to be more binding in several practical applications, such as online ordered package delivery, blood product distribution (Hemmelmayr et al. 2009) and replenishment of ATMs. Tour duration constraints are rarely found in the IRP literature. Finally, the DJRP does not take limited customer storage capacity into account, but in practice, storage capacity is often restricted. To address these shortcomings of the DJRP we propose an extension of the DJRP, the DJRP with Approximated Transportation Costs (DJRP-AT), that explicitly considers transportation costs. Furthermore, the DJRP-AT contains tour duration constraints and limits customer storage capacity. Because determining the optimal delivery tour is computationally expensive, we will approximate the transportation costs in the DJRP-AT by approximating the shortest traveling salesman tour. The only work that we are aware of that includes approximated transportation costs in a VMI setting is Larsen and Turkensteen (2014). The authors consider a VMI setting with stochastic demand and order-up-to-levels at the customers which they solve with a Markov Chain simulation model.

Our research is motivated by ATM replenishment in the Netherlands. A single supplier (vendor) decides on the timing of ATM cash replenishment and on the delivery quantity. The actual ATM replenishment orders per day are outsourced to an LSP, in this application often referred to as Cash-in-Transit company (CIT), that schedules and performs the daily delivery routes. Currently, the supplier pays a fixed fee to the LSP for each ATM replenishment. Therefore, the current replenishment policy ignores the impact that ordering decisions have on distance traveled and vehicle utilization. The supplier is reconsidering the replenishment cost structure to better align decisions. To provide insight for future negotiations between the supplier and the LSP, we examine the benefit of adopting a DJRP and a DJRP-AT perspective. How a new ordering policy is to be incorporated in the contract between the supplier and the LSP, is beyond the scope of this paper. With this work we contribute to the recent stream of publications on ATM replenishment. For example, Van Anholt et al. (2016) develop a heuristic for a pickup and delivery IRP for an advanced type of ATMs. Larrain et al. (2017) focus on a local search based heuristic for an IRP in which stock-outs are allowed and cash is replenished by swapping cassettes.

To solve the DJRP-AT, this paper proposes a compact formulation in which transportation costs and inventory holding costs are minimized. Note that inventory holding costs in the application relate to the value of money or lost interest. The compact formulation is split by applying Dantzig-Wolfe decomposition (Desrosiers and Lübbecke 2005). The resulting Master Problem and Pricing Problem are solved in a Branch-and-Cut-and-Price framework (Nemhauser and Park 1991, Lübbecke and Desrosiers 2005). The Master Problem selects customer subsets to be delivered and determines the corresponding delivery quantities. The Pricing Problem generates these customer subsets using a labeling algorithm with tailored dominance criteria to speed up the process. The solution method is tested on benchmark instances from the literature and on instances derived from a real-life case in ATM replenishment.

The contributions of this paper are threefold. First, the DJRP is extended to incorporate transportation costs, limitations on storage capacity at the customers and restricted tour duration. The results show that the proposed model leads to lower total costs compared with the DJRP. Second, we introduce novel dominance conditions for the labeling algorithm that is used to solve the Pricing Problem. Finally, existing valid inequalities originating from the inventory routing literature are tested, their impact on the integrality gap is demonstrated and it is shown that their effectiveness is different than for other models in which they have been applied.

The remainder of this paper is organized as follows: Section 2 discusses literature on the JRP and DJRP, together with their relation to the IRP. The DJRP-AT is described and modeled in Section 3. Section 4 proposes a decomposition of the model and specifies the Master and Pricing Problems. The algorithm to solve the Pricing Problem, including novel sufficient dominance conditions, the valid inequalities and the branching strategy are presented in Section 5. Section 6 presents the results of the experiments on benchmark instances from the literature, introduces the real-life case and reports on the results of instances derived from the real-life case. Finally, the conclusions and directions for further research are discussed in Section 7.

2. Literature review

The traditional JRP is the problem of minimizing holding and ordering costs, while ensuring that no customer runs out of stock in any period of the planning horizon. The ordering costs consist of a common set-up cost per period and a fixed fee per replenishment. An overview of the literature on the JRP from 1989 to 2005 distinguishes three types of models (Khouja and Goyal 2008): first, the traditional JRP, considers deterministic and static demand. This means that demand is known beforehand and remains the same for every period of the planning horizon. For this problem, analytical expressions have been derived for the minimal total costs and heuristics have been designed to determine the corresponding cyclic replenishment policy. Second, the extension to stationary stochastic demand in which the objective is to minimize the expected total cost. Solution methods mainly consist of using a periodic review policy or a can-order policy. Finally, the JRP with deterministic and dynamic demand (DJRP) in which the demand is known but can vary across periods is discussed. The solution for this type of problem is not necessarily a cyclic replenishment policy as for the traditional JRP. For the DJRP different formulations and heuristic solution methods have been proposed and studied (Webb et al. 1997, Boctor et al. 2004, Narayanan and Robinson 2006, Robinson et al. 2007) and Robinson et al. (2009) have provided an overview of available solution methods. Webb et al. (1997) studied fixed replenishment cycle models for the problem and compared these to optimal solutions that do no constrain the replenishment cycle. Boctor et al. (2004) proposed several linear programming formulations, tested several heuristic solution methods and proposed an improvement procedure that can be used in combination with a heuristic method.

To increase practical relevance of the DJRP, several extensions have been proposed such as capacitated aggregate order size (Anily and Tzur 2005, Federgruen et al. 2007, Narayanan and Robinson 2010), supplier selection (Ventura et al. 2013), supplier selection with discounts (Kang et al. 2016), inventory decisions at the supplier (Solyalı and Süral 2012, Cunha and Melo 2016) as well as inventory decisions and capacitated production at the supplier (Senoussi et al. 2016). A commonly occurring practical constraint is an inventory capacity limit at the customers. However, to our knowledge, this constraint has only been included in one paper on the DJRP (Senoussi et al. 2016); two papers on the traditional JRP also include this constraint (Hoque 2006, Hariga et al. 2013).

The IRP combines an inventory problem and a routing problem: it minimizes inventory holding and routings costs by optimizing replenishments for a set of customers and explicitly determining the delivery routes. The IRP is therefore related to the JRP, yet the IRP is structurally different from the JRP because the routing problem is explicitly solved. Various solution methods for the IRP have been proposed in the literature such as exact methods, matheuristics and metaheuristics (see Coelho et al. (2014) for an overview). Some of the exact solution methods for the IRP rely on the vehicle capacity constraint, for example in the Pricing Problem algorithms and valid inequalities. In our application a tour duration constraint is more appropriate. In some of the heuristic solution methods for the IRP, inventory and routing optimization are considered separately. In a first phase, decisions are made on the inventory policies, often incorporating a fixed replenishment cost per delivery, thus solving a variant of the JRP. In a second phase, routing is optimized given the replenishment decisions of the first phase. Iterative solution schemes have for example been proposed by Cordeau et al. (2015) and Absi et al. (2015).

Some attention has also been paid to the fact that charging a fixed fee for servicing a customer in the DJRP is not always representative for the actual costs involved. A fixed fee per customer replenishment assumes that the costs for replenishing customers are independent, but in practice, this is not always true. Olsen (2008) used the example of using a refrigerated truck for canned food delivery, which increases the marginal replenishment costs of the canned food. Olsen (2008) and Wang et al. (2012) proposed to model the marginal costs with additional fixed fees depending on the combination of items delivered. Senoussi et al. (2016) recognized that actual transportation cost are relevant, however, they assumed that the depot is located far away from a cluster of customers and that the transportation costs between the clustered customers are negligible, therefore the authors assumed that the costs of a tour are fixed. Rahmouni and Hennet (2015) took actual routing costs into account by combining the deterministic and static JRP with the Traveling Salesman Problem (TSP). For each possible subset of customers the actual tour length was computed beforehand by solving a TSP, then a linear programming model was used to select the optimal subsets and to determine the delivery quantities. However, this solution method can only be applied to instances of very limited size since for all combinations of customers the traveling salesman tour has to be computed.

3. Problem description

In the DJRP-AT, a single supplier supplies N customers. The customers face a certain demand per period and have a limited storage capacity, and therefore require replenishments to prevent them from running out of stock. The supplier arranges the customer replenishments in a VMI setting with the objective of minimizing transportation and customer inventory holding costs. The transportation costs in a period are represented by the approximated tour length visiting the replenished customers and a fixed set-up fee for a period if at least one customer is replenished in that period. In the DJRP-AT, constraints are incorporated on the composition of the set of customers served in one period, e.g., the number of customers served or the tour duration. Note that the inventory holding costs at the supplier are not considered since in our practical application there is an infinite supply (similar to Larrain et al. (2017)), but these costs could easily be added.

For the calculation of the transportation costs, consider that these costs must be estimated for a large number of customer subsets, which requires careful balance of approximation accuracy and calculation effort. Also, two sets of customers with the same cardinality, but with customers at different locations should result in different transportation costs. Therefore, based on the short literature review in Appendix A, we adopt the tour length approximation model of Chien (1992):

$$D \approx 0.98\sqrt{RM'},\tag{1}$$

in which R is the area of the smallest rectangle covering both the customers and the depot, and M'is the number of points in the tour (depot and customers). Note that this function underestimates the actual TSP tour length. The transportation costs also include a fixed cost B that is independent of the distance traveled, but is for example a setup cost related to vehicle use. Define binary vector Y to indicate which customers are served and binary variable \hat{Y} to indicate whether any customer is served $(\sum_i Y_i \ge 1)$. The following transportation cost function will be used to approximate the transportation costs for the customers in Y and the depot

$$f(\mathbf{Y}) = B\hat{Y} + 0.98\sqrt{R(\mathbf{Y})M'(\mathbf{Y})}.$$
(2)

To formulate the DJRP-AT, consider the following notation in Boctor et al. (2004). A single depot and a set of $\mathcal{N} = \{1, 2, ..., N\}$ customers are positioned in Euclidean space and there is a finite time horizon $\mathcal{T} = \{1, 2, ..., T\}$. Let Y_{it} denote the binary decision variable that takes value 1 if and only if customer $i \in \mathcal{N}$ is visited in period $t \in \mathcal{T}$. Let Y_t denote the vector $\{Y_{1t}, Y_{2t}, ..., Y_{Nt}\}$. Define X_{it} as the quantity delivered to customer $i \in \mathcal{N}$ in period $t \in \mathcal{T}$ and let I_{it} be the quantity in stock at customer $i \in \mathcal{N}$ at the end of period $t \in \mathcal{T}$. I_{it} should be non-negative, because stock-outs are not allowed. The inventory level is measured at the end of the period assuming the following order of events: delivery of new stock, consumption, inventory calculation. This assumption coincides with JRP literature (Boctor et al. 2004) and with most literature on the IRP (Archetti et al. 2014). I_{i0} denotes the initial inventory level and d_{it} is the dynamic and deterministic demand in period $t \in \mathcal{T}$ at customer $i \in \mathcal{N}$. For the items in stock at a customer $i \in \mathcal{N}$ an inventory holding rate of h_{it} is charged per period $t \in \mathcal{T}$.

We introduce the following additional notation for the DJRP-AT. Each customer $i \in \mathcal{N}$ has a storage capacity u_i . Define for each $i \in \mathcal{N}$ and $t \in \mathcal{T}$ big-M value $M_{it} = \min\left\{u_i, \sum_{s=t}^T d_{it}\right\}$. Furthermore, a single vehicle with unlimited load capacity performs at most one route in each time period, beginning and ending at the depot. Transportation costs in a period are represented by function $f(\cdot)$ as defined in equation (2). Finally, let function $g(\cdot)$ assess the composition of the tour in a given period. We consider two different functions for $g(\cdot)$. First, let $g(\cdot)$ be the approximated tour duration and define $k_D \in \mathbb{R}$ as the maximum tour duration. Second, we let $g(\cdot)$ be the number of customers in a tour and impose that at most $k_M \in \mathbb{N}$ customers can be served in a single tour. These constraints will be referred to as 'subset composition constraints' for the remainder of the paper. DJRP-AT models will only contain one of these two types of constraints to assess the tour composition.

The goal of the DJRP-AT is to minimize inventory holding and transportation costs by selecting, for each period, which customers to replenish, while avoiding stock-out at any customer, without violating the customer's storage capacity restrictions and the additional restrictions on the tour composition. This problem can be formulated as follows:

$$z = \min \sum_{t \in \mathcal{T}} \sum_{i \in \mathcal{N}} h_{it} I_{it} + \sum_{t \in \mathcal{T}} f(\mathbf{Y}_t)$$
(3a)

s.t.
$$I_{it} = I_{i,t-1} - d_{it} + X_{it} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
 (3b)

$$X_{it} \le u_i - I_{i,t-1} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
(3c)

$$X_{it} \le M_{it} Y_{it} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
(3d)

- $g(\mathbf{Y}_t) \le k \qquad \forall t \in \mathcal{T}$ (3e)
- $I_{it} \ge 0 \qquad \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$ (3f)

$$t \ge 0 \qquad \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
 (3g)

$$Y_{it} \in \{0, 1\} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
(3h)

in which $k = k_D$ or $k = k_M$, depending on the applied subset composition constraints. The objective function (3a) minimizes the costs for inventory holding and transportation $f(\cdot)$ defined in (2). The inventory balance for each customer in each period is maintained by constraints (3b). Constraints (3c) ensure that the customer's capacity is not exceeded when a delivery is made and constraints (3d) force the amount delivered to zero if a customer is not visited. Furthermore, constraints (3e) represent the additional constraints on the composition of the subset of customers replenished in a period. Constraints (3f), (3g) and (3h) impose binary and non-negativity constraints on the decision variables.

 X_i

4. Column generation

When considering the complexity of the DJRP-AT with cost function (2), it is important to note that the traditional JRP is not a special case of the DJRP-AT, due to the different cost structure. Hence, although the traditional JRP is NP-complete (Arkin et al. 1989), this conclusion cannot be directly made for the DJRP-AT. Furthermore, analysis of the literature on the complexity of related problems shows that the so-called Pinwheel Scheduling Problem is a special case of the DJRP-AT with cost function (2). The Pinwheel Scheduling Problem is likely to be an NP-complete problem (Jacobs and Longo 2014), but this has not yet been proven despite several attempts. A mapping between the Pinwheel Scheduling Problem and the DJRP-AT, including details on the complexity of the Pinwheel Scheduling Problem, are presented in Appendix B. Because the existence of a polynomial-time algorithm is unlikely and column generation has proven to be efficient for similar problem structures, this solution method will be used to solve the DJRP-AT.

Application of the Dantzig-Wolfe decomposition to problem (3a)-(3h) results in a Master Problem that selects for every period a subset of customers to replenish out of a collection of subsets to minimize the inventory holding and transportation costs. Moreover, the Master Problem optimizes the delivery quantities corresponding to constraints (3b)-(3d) in the compact formulation. The Pricing Problem generates subsets of customers, taking constraints (3e) into account, and is solved for each period separately.

To formulate the Master Problem, let S_t be the collection of subsets of customers that are generated by the Pricing Problem for period $t \in \mathcal{T}$. The binary decision variable Z_{st} equals 1 if subset $s \in S_t$ is selected for period $t \in \mathcal{T}$. For a specific subset $s \in S_t$ the transportation costs c_s for servicing its customers is given by the Pricing Problem. Furthermore, let a_{is} indicate whether customer $i \in \mathcal{N}$ is present in subset $s \in S_t$. The decomposition gives the following Master Problem:

$$z = \min \sum_{t \in \mathcal{T}} \sum_{i \in \mathcal{N}} h_{it} I_{it} + \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} c_s Z_{st}$$
(4a)

s.t.
$$I_{it} = I_{i,t-1} - d_{it} + X_{it}$$
 $\forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$ (4b)

$$X_{it} \le u_i - I_{i,t-1} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$

$$(4c)$$

$$X_{it} \le u_i \sum_{s \in \mathcal{S}_t} a_{is} Z_{st} \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$

$$(4d)$$

$$\sum_{st} Z_{st} \le 1 \qquad \forall t \in \mathcal{T}$$
(4e)

$$I_{it} \ge 0 \qquad \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
(4f)

$$X_{it} \ge 0 \qquad \qquad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
(4g)

$$Z_{st} \in \{0, 1\} \qquad \qquad \forall s \in \mathcal{S}_t, \ \forall t \in \mathcal{T}$$
(4h)

The objective function (4a) aims to minimize total costs. Constraints (4b)-(4d) are equivalent to constraints (3b)-(3d) of the compact formulation. Constraint (4e) ensures that at most one subset of customers is selected for each period. Finally, non-negativity and binary requirements on the decision variables are imposed by constraints (4f)-(4h).

We use column generation to solve the linear programming relaxation of (4a)-(4h) since the total number of variables Z_{st} is exponentially large. Starting with a small subset of all possible columns gives the Restricted Master Problem (RMP) and additional columns with negative reduced cost are generated by repeatedly solving the Pricing Problem. To formulate the Pricing Problem, let us associate the following dual variables with the Master Problem with respect to decision variables Z_{st} . Let π_{it}^1 be a non-positive dual variable associated with constraints (4d) and let π_t^2 be the nonpositive dual variable of constraints (4e). Let us also reuse decision variables Y_{it} from the compact formulation: these variables indicate whether customer $i \in \mathcal{N}$ is replenished in period $t \in \mathcal{T}$ and remind that $\mathbf{Y}_t = \{Y_{1t}, Y_{2t}, \ldots, Y_{Nt}\}$. For a given time period $t \in \mathcal{T}$ the Pricing Problem can be formulated as follows:

min
$$\bar{c}_t(\mathbf{Y}_t) = f(\mathbf{Y}_t) + \sum_{i \in \mathcal{N}} u_i Y_{it} \pi_{it}^1 - \pi_t^2$$
 (5a)

s.t.
$$g(\boldsymbol{Y_t}) \le k$$
 (5b)

$$\boldsymbol{Y_t} \in \{0,1\}^N \tag{5c}$$

The objective (5a) is to minimize the reduced cost $\bar{c}_t(\mathbf{Y}_t)$ while the subset composition constraints are satisfied (5b). The reduced cost consists of the transportation costs of the subset $f(\mathbf{Y}_t)$ and dual terms corresponding to the current solution of the RMP. The subset composition constraints (5b) can, in general, concern any function of the combination of customers in the subset. However, these constraints cannot contain the delivery quantities, since these quantities are determined in the Master Problem. Hence, in our model, a load capacity constraint cannot be in the Pricing Problem, but, for example, a tour duration constraint is possible.

5. Branch-and-Cut-and-Price

The Master Problem and Pricing Problem of Section 4 are solved in a Branch-and-Cut-and-Price framework. In Section 5.1, a tailored labeling algorithm to solve the Pricing Problem per period is described and novel sufficient conditions are presented that provide a dominance criterion to discard labels. Valid inequalities are presented in Section 5.2 and Section 5.3 provides a description of the branching strategy.

5.1. Labeling Algorithm for the Pricing Problem

To solve the Pricing Problem, we propose a tailored labeling algorithm that identifies subsets of customers that will improve the current solution of the RMP. Note that during the process of generating subsets of customers, only the customer combination is relevant, there is no sequential relationship between the customers as opposed to routing problems (Feillet et al. 2004).

Define label $L = \langle s(L), \bar{c}_t(L), g(L) \rangle$ in which s(L) is the subset of customers, $\bar{c}_t(L)$ is the corresponding reduced cost and g(L) represents the value of the function $g(\cdot)$ in the subset composition constraint for subset s(L). Hence, each label corresponds to a subset of customers that is a candidate to be added to the RMP.

The labeling algorithm starts for each customer $i \in \mathcal{N}$ separately and the labels are extended by adding the other customers one by one. The order of the customers in s(L) is not important, since the subset of customers is considered for replenishment, but the order in which they are served is not determined. Hence, each possible subset has to be considered at most once. Therefore, when starting with a label containing customer i and extending with customer j, the inverse order of these customers, starting with j and adding i, does not have to be considered. The labeling algorithm terminates when all possible subsets of customers are considered.

Denote $L \oplus P$ as the resulting label from the extension of label L with the customers in set $P \subseteq \mathcal{N} \setminus s(L)$. The operation to extend a label L with the next customer j is to set $s(L \oplus \{j\}) = s(L) \cup \{j\}$ and to compute $\bar{c}_t(L \oplus \{j\})$ and $g(L \oplus \{j\})$:

$$\bar{c}_{t}(L \oplus \{j\}) = \bar{c}_{t}(L) - f(\mathbf{Y}_{t}(s(L))) + f(\mathbf{Y}_{t}(s(L) \cup \{j\})) + u_{j}\pi_{jt}^{1}$$

$$= f(\mathbf{Y}_{t}(s(L) \cup \{j\})) + \sum_{i \in s(L) \cup \{j\}} u_{i}\pi_{it}^{1} - \pi_{t}^{2}$$
(6)

in which $Y_t(s(L))$ is the vector in which the variables corresponding to s(L) equal 1. If we consider the model with subset composition constraints that set a maximum on the tour duration, we have

$$g(L \oplus \{j\}) = 0.98\sqrt{R(s(L) \cup \{j\})(|s(L)|+1)}$$
(7)

and if we consider the model with subset composition constraints that pose a maximum on the number of customers in the subset, we have

$$g(L \oplus \{j\}) = g(L) + 1 \tag{8}$$

The extended label is feasible if

$$s(L) \cap \{j\} = \emptyset \land g(L \oplus \{j\}) \le k \tag{9}$$

When the number of customers N increases, the maximum number of labels becomes large $(2^N - 1)$. A dominance test will therefore be used to reduce the number of labels.

Denote the set of feasible extensions of label L by E(L) which consists of all combinations of the customers that have not already been considered and for which the extension will satisfy the subset composition constraints. The following definition for dominance holds

DEFINITION 1. Label *L* dominates label *L'* if D.1 $E(L') \subseteq E(L)$ D.2 $\bar{c}_t(L \oplus P) \leq \bar{c}_t(L' \oplus P), \forall P \in E(L')$

The first condition, D.1, states that a feasible extension of L' must also be a feasible extension of L. The second condition, D.2, requires that all feasible extensions of L do not result in worse solutions than the same extensions of L'. These conditions are difficult to check in practice, since all feasible extensions would have to be computed. Therefore, Proposition 1 introduces sufficient conditions for dominance of L over L'.

PROPOSITION 1. Label L dominates label L' if the following conditions hold P.1 $s(L) \subseteq s(L')$ P.2 $g(L) \leq g(L')$ P.3 $\bar{c}_t(L) + \Delta(L, L') \leq \bar{c}_t(L')$

Conditions P.1 and P.2 combined imply condition D.1, such conditions are also used for shortest path problems (Feillet et al. 2004), and condition P.3 implies condition D.2. Before a formal proof for Proposition 1 is presented, an intuitive reasoning for condition P.3 is given and an expression for $\Delta(L, L')$ is derived.

The cost function, and therefore the reduced costs, are dependent on the number of customers and the area in which these customers are located. Consider a comparison of the two labels Land L' for which it holds that $s(L) \subseteq s(L')$, $\bar{c}_t(L) < \bar{c}_t(L')$ and $g(L) \leq g(L')$. One would like to conclude that L dominates L'. However, if a set of customers P is added to both labels, the area that is used in the cost function can increase more for L than for L', i.e., the additional cost of the extension with P is not identical for both labels. This could result in $\bar{c}_t(L \oplus P) > \bar{c}_t(L' \oplus P)$, therefore, it cannot be concluded that L dominates L' since condition D.2 is violated. Hence, a sufficient dominance condition should be stricter than $\bar{c}_t(L) \leq \bar{c}_t(L')$, therefore sufficient condition P.3 is introduced. This will be illustrated in the following example. Consider an instance with four customers, indicated by white nodes and customer index in Figure 1. The depot is indicated by the black node with label D. The current terms for the reduced cost corresponding to each customer are indicated between brackets (suppose $\pi_t = 1000$).

Figure 1 Example of dominance in labeling algorithm.



Now, consider subset $s_1 = \{1,2\}$, the corresponding cost is $f(s_1) = 1000 + 0.98\sqrt{100 \times 200 \times 3} \approx 1240$ and the reduced cost of this subset is $\bar{c}_t(s_1) \approx 1240 - 200 - 100 - 1000 = -60$. Similarly, subset $s_2 = \{1,2,3\}$ has $f(s_2) \approx 1537$ and $\bar{c}_t(s_2) \approx 37$. In this case s_1 and s_2 are comparable, since $s_1 \subset s_2$. Note that $\bar{c}_t(s_1) < \bar{c}_t(s_2)$, hence, one would like to conclude that the label with subset s_1 dominates the label with s_2 . However, suppose customer 4 is added to both subsets. This gives $s_3 = \{1,2,4\}$ with $f(s_3) \approx 1620$ and $\bar{c}_t(s_3) \approx 70$ and $s_4 = \{1,2,3,4\}$ with $f(s_4) \approx 1693$ and $\bar{c}_t(s_4) \approx -57$. Note that the cost increase from subset s_1 to s_3 is larger than from subset s_2 to s_4 . Because $\bar{c}_t(s_3) > \bar{c}_t(s_4)$, we cannot conclude from $\bar{c}_t(s_1) < \bar{c}_t(s_2)$ that the label with s_1 dominates the label with s_2 .

A more strict condition is required and therefore $\Delta(L, L')$ is introduced, representing the maximum difference in costs between two labels. The value of $\Delta(L, L')$ must be sufficient to guarantee $\bar{c}_t(L \oplus P) \leq \bar{c}_t(L' \oplus P)$ ($\forall P \in E(L')$) to conclude that L dominates L'. First, express $\bar{c}_t(L \oplus P)$ and $\bar{c}_t(L' \oplus P)$ in terms of $\bar{c}_t(L)$ and $\bar{c}_t(L')$, respectively. Combined with $\bar{c}_t(L) \leq \bar{c}_t(L')$, an upper bound can be derived for the difference between $\bar{c}_t(L \oplus P)$ and $\bar{c}_t(L' \oplus P)$. The derivation for the following sufficient value of $\Delta(L, L')$ is given in Appendix C:

$$\Delta(L,L') = \phi \sqrt{R(s(L'))|s(L') \cup P| - R(s(L))|s(L) \cup P|}$$
(10)

with $R(\cdot)$ as the area of the smallest rectangle and $P = \mathcal{N} \setminus s(L')$. A formal proof for Proposition 1 with $\Delta(L, L')$ as defined in (10) can now be presented.

Proof of Proposition 1 Assume two labels L and L' with corresponding subsets of customers s = s(L) and s' = s(L') satisfy the conditions in Proposition 1. Given $s \subseteq s'$ and therefore $s \cup P \subseteq s' \cup P$, it holds that if $g(L) \leq g(L')$, then $g(L \oplus P) \leq g(L' \oplus P)$ if $g(\cdot)$ is a monotone function. Hence, if $g(L' \oplus P) \leq k$, then $g(L \oplus P) \leq k$ and condition D.1 is satisfied. To show that condition D.2 holds, note that we have already shown in Appendix C that

$$\bar{c}_t(L \oplus P) - \bar{c}_t(L' \oplus P) \le \bar{c}_t(L) - \bar{c}_t(L') + \Delta(L,L')$$
(11)

Hence, if $\bar{c}_t(L) + \Delta(L, L') \leq \bar{c}_t(L')$, then condition D.2 holds which concludes the proof of Proposition 1. \Box

An overview of the labeling algorithm is provided in Algorithm 1. A label is not extended any further in the labeling algorithm if the subset composition constraint is violated, since no feasible subsets of customers can be found by adding more customers because of the subset composition constraints. Moreover, after extending a label, it is tested whether adding another customer violates a subset composition constraint, in which case an extension to this customer from the current label is not considered in a later stage of the labeling algorithm.

Algorithm 1 Labeling algorithm

Initialize list of improving labels \mathcal{I} and of labels to propagate \mathcal{P}
for $i = 0$ to N do
Create label l containing i and add l to \mathcal{P}
if Reduced cost of <i>l</i> is negative then
Add l to \mathcal{I}
$\mathbf{while} \mathcal{P} \neq \emptyset \mathbf{do}$
Consider a waiting label $p \in \mathcal{P}$
for All customers j with higher index than the last added customer to p do
Extend p with j to q
if q is feasible and p does not dominate q then
$\mathrm{Add}\; q \; \mathrm{to}\; \mathcal{P}$
if Reduced cost of $u < 0$ then
$\mathrm{Add}\; q \; \mathrm{to}\; \mathcal{I}$
Remove p from \mathcal{P}

To accelerate the solution process, a heuristic variant of the labeling algorithm is applied. If the heuristic fails to find any improving subsets, the exact labeling algorithm is applied in which all combinations of customers are considered. In the heuristic pricing instrument, the process is identical to the exact pricing algorithm, but the number of customers to which a label can be extended is limited. Only the extensions to the b customers closest to the last added customer of a subset are evaluated. The initial value of b is small (2) and this value is doubled up to a certain limit (8) as long as no improving subsets of customers are found.

Preliminary experiments showed that adding all columns with negative reduced cost could be time consuming for the algorithm and moreover, many of these columns are not in the final solution. Therefore, at most 10,000 columns are added per call to the Pricing Problem for both the heuristic and the exact pricing instrument.

5.2. Valid inequalities

The formulation of the problem can be strengthened with valid inequalities. Two valid inequalities that were introduced by Archetti et al. (2007) for the IRP are also applicable to the DJRP-AT.

The first inequality states that if a customer $i \in \mathcal{N}$ is not replenished in periods $t - r, t - r + 1, \dots, t$, then the inventory in period t - r - 1 should be sufficient to cover demand of all periods up to t:

(ILB)
$$I_{i,t-r-1} \ge \left(\sum_{j=0}^{r} d_{i,t-j}\right) \left(1 - \sum_{j=0}^{r} \sum_{s \in \mathcal{S}_t} a_{is} Z_{s,t-j}\right) \quad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}, \ \forall r = 0, 1, \dots, t-1$$
(12)

The second inequality gives a lower bound for the number of required visits to a specific customer i up to a period t taking the customer's inventory capacity into account:

(NrVis)
$$\sum_{j=1}^{t} \sum_{s \in \mathcal{S}_t} a_{is} Z_{sj} \ge \left\lceil \frac{\sum_{j=1}^{t} d_{ij} - I_{i0}}{u_i} \right\rceil \quad \forall i \in \mathcal{N}, \ \forall t \in \mathcal{T}$$
 (13)

Preliminary experiments showed that dynamic management of the valid inequalities, i.e., adding them whenever violated, slowed down the execution. Therefore, for all experiments, all valid inequalities are added to the model in the root node of the Branch-and-Bound tree.

5.3. Branching

In the compact formulation (3a)-(3h), the variables indicating whether a customer is served in a certain time period, the assignment variables, are binary decision variables. The delivery quantity and inventory level decision variables are non-negative and continuous. To find binary results for the assignment variables a Branch-and-Bound tree is used, which is explored via a best bound strategy. First, the algorithm branches on the total number of replenishments of a customer $i \in \mathcal{N}$ over all periods $\sum_{t \in \mathcal{T}} Y_{it}$. In the Master Problem variables this can be expressed as $\sum_{t \in \mathcal{T}} \sum_{s \in S_t} a_{is} Z_{st}$. If all customers have an integer number of replenishments, the second branching method branches on whether any customer is replenished in a period t or no customer is replenished $(\sum_{i \in \mathcal{N}} Y_{it})$. Expressed in the Master Problem variables, the corresponding constraints are $\sum_{s \in S_t} z_{st} \leq 0$ and $\sum_{s \in S_t} z_{st} \ge 1$. If no new branches can be identified for the first two branching methods, the algorithm branches on whether a customer $i \in \mathcal{N}$ is visited in a specific period $t \in \mathcal{T}$ or not $(Y_{it} =$ $\sum_{s \in S_t} a_{is} Z_{st}$). This branching method leads to binary solutions for the assignment variables in the compact formulation, hence, if no new branches are identified an integer solution is found. If there is more than one branch candidate, the following strategy to select a branch is followed for each type of branching. For each branch candidate the child nodes are quickly evaluated by solving the LP relaxation given the current set of columns. The branch that maximizes the lower bound is chosen. We consider at most 25 branch candidates in the first 20 nodes of the branch and bound tree and at most 15 candidates in the other nodes.

6. Computational results

The proposed model for the DJRP-AT is analyzed for two types of subset composition constraints: a tour duration constraint and a maximum number of customers served per period (i.e., in a subset). The effectiveness of the valid inequalities presented in Section 5.2 is evaluated for both types of subset composition constraints in Section 6.1. The model with a maximum tour duration most resembles practical cases; this model is tested with different values for the maximum tour duration. The model with a maximum number of customers per subset can be compared to the DJRP with fixed fees (Boctor et al. 2004) and to a variant of the IRP in which the actual routing costs are used. The first comparison provides insight in the potential improvements that can be achieved with the DJRP-AT compared to the DJRP. Section 6.2 explains how the DJRP-AT and the DJRP are compared and provides the results. The second comparison shows how well the DJRP-AT performs compared to an equivalent IRP in which actual routing costs are used, on which Section 6.3 reports the results. Note that results of the DJRP-AT are not comparable to the results of 'standard' IRP as often used in the literature (Coelho et al. 2014) because the constraints are different.

The instances for the IRP, created by Archetti et al. (2007), are used for the computational experiments. Although the IRP differs from the DJRP-AT regarding the cost structure, decision variables and constraints, the IRP instances are used as a base, since they contain most of the data required for the experiments on the DJRP-AT. The time horizon is equal to either 3 or 6 periods and instances with 5, 10, 15 and 20 customers are considered; there are five instances for each combination of number of customers and periods. For each customer, a location is given by two coordinates, both randomly chosen within the interval [0, 500]; demand is randomly selected between 10 and 100, and the customer holding rate is in the interval [0.1, 0.5]. The customer's inventory capacity is the demand of the customers multiplied by either 2 or 3, which is randomly selected. The initial inventory is the customer's capacity minus the demand of the first period. The instances are available online (Coelho n.d.). For the tests on the model including maximal tour duration the maximum (k_D) is set to 600, 800, 1000 and 1200 for all instances. If a maximum is set on the number of customers per subset (k_M) , this maximum depends on the number of customers in the instance. For 5 customers, maxima of 3 and 4 are considered, for 10 customers 5, 6, 7 and 8 are considered as maxima, for 15 customers 7, 8, 9, 10 and 11 are considered, and for instances with 20 customers 10, 11, 12 and 13 are the maxima. The fixed major cost is set to B = 1000.

The RMP is initialized with dummy columns with very high costs to guarantee a feasible solution for the initial linear program. To improve computation times, two heuristics are designed to attempt quickly identifying columns that provide a feasible integer solution. The first heuristic assigns customers to periods in a greedy way, ensuring customers do not run out of stock and respecting the subset composition constraints. If the subset composition constraint is a maximum on the tour duration, a second heuristic is applied if the first one did not yield a feasible integer solution. In the second heuristic, before applying the steps of the first heuristic, the customers are sorted by decreasing distance from the depot. Since the sorting process indirectly takes the subset composition constraint into account, this provides a greater chance of finding a feasible solution. Limited computational experiments showed that the first heuristic provides a better upper bound if a solution is found, therefore this heuristic is applied first.

The algorithm to solve the DJRP-AT as described in the previous sections is implemented using Java and Gurobi 6.5. All tests are performed on a desktop computer running Windows 10, equipped with an eight core Intel(R) Core(TM) i7-6700K, CPU 4.00GHz processor with 24GB of RAM. A single core is used to generate the results and the maximum running time is two hours per instance.

6.1. Effectiveness of valid inequalities

Two valid inequalities are considered to strengthen the formulation for the DJRP-AT. These valid inequalities are tested for both types of subset composition constraints and the results are presented in Table 1. The results are aggregated per number of customers in the instance (N) and the length of the time horizon (T). The average solution time over all tested instances, the number of instances solved, and the average integrality gap for the solved instances are reported for the model without valid inequalities, the model with only the ILB inequalities (equation 12), the model with only NrVis inequalities (equation 13), and the model with both inequalities, respectively. Next to the number of instances solved, the total number of tested instances that have not been proved infeasible is indicated between brackets. The difference between the two numbers gives the number of instances that have not been solved in two hours. The integrality gap is the percentage difference between the optimal binary solution (LB_{best}) and the solution of the relaxation of the model in the root node of the Branch-and-Bound tree (LB_{root}) , which is calculated by $(LB_{best} LB_{root})/LB_{best}$. To test the effectiveness of the valid inequalities, for instances with 15 customers and 3 periods, duration 600 is not considered; for 15 customers and 6 periods, both 600 and 800 are not considered, since most instances would be infeasible. For instances with 15 customers, a maximum number of customers per period (k_M) of 8, 9, 10 and 11 are considered.

The observed integrality gaps, as reported in Table 1, are quite high and decrease if more valid inequalities are added. The high integrality gaps can be partially explained by the formulation of the Master Problem. The linear relaxation of the model allows for satisfying customer demand in a certain period using only a fractional value for the decision variable corresponding to a subset containing this customer. Therefore, the costs of the subset are only fractionally accounted for. Moreover, the linear relaxation allows infeasible combinations of customers to be served in one period in a fractional solution. Therefore, the costs of the fractional solution can be lower, even tough the solution is certainly not feasible. Both of these effects were observed in our experiments. For example, consider instance abs3n5 from Archetti et al. (2007) which contains 5 customers,

			None			Only ILB Only NrVis					ILB and NrVis					
		Time	#	Gap	Time	#	Gap	Time	#	Gap	Time	#	Gap			
N	T	(s)	Solved	(%)	(s)	Solved	(%)	(s)	Solved	(%)	(s)	Solved	(%)			
Cor	nstra	aint on	duration													
5	3	0	20 (20)	28	0	20 (20)	22	0	20 (20)	7	0	20 (20)	7			
10	3	2	18 (18)	42	1	18 (18)	34	1	18 (18)	16	1	18 (18)	16			
15	3	40	13 (13)	44	18	13 (13)	37	17	13 (13)	20	6	13 (13)	20			
5	6	1	20 (20)	25	1	20 (20)	19	1	20 (20)	6	0	20 (20)	3			
10	6	83	16 (16)	31	39	16 (16)	25	71	16 (16)	12	19	16 (16)	9			
15	6	3801	6 (10)	31	3021	7 (10)	26	3316	7 (10)	13	3456	6 (10)	11			
Sun	n		93 (97)			94 (97)			94 (97)			93 (97)				
Сог	ıstra	aint on	number o	f cust	omers											
5	3	0	10 (10)	46	0	10 (10)	42	0	10 (10)	20	0	10 (10)	20			
10	3	2	20 (20)	42	1	20 (20)	34	1	20 (20)	14	1	20 (20)	13			
15	3	30	20 (20)	42	12	20 (20)	34	15	20 (20)	15	7	20 (20)	14			
5	6	3	10 (10)	32	1	10 (10)	26	2	10 (10)	13	0	10 (10)	10			
10	6	86	20 (20)	31	29	20 (20)	24	39	20 (20)	11	20	20 (20)	8			
15	6	4269	12 (20)	30	3212	14 (20)	24	3264	15 (20)	11	2507	17 (20)	11			
Sun	n		92 (100)			94 (100)			95 (100)		97 (100)					

 Table 1
 Effectiveness of valid inequalities for DJRP-AT for two types of constraints.

3 time periods and set $k_D = 600$ as the maximum on the duration. The optimal solution selects customer subset {3} in period 1, subset {1,2} in period 2 and subset {3,4,5} in period 3; the objective value is 4310. In the solution of the relaxed model, subsets {1,2} and {3,4,5} are selected with value 0.5 in both period 1 and period 2. The objective value of this fractional solution is 3397, resulting in an integrality gap of 21%. Note that in the fractional solution all customers can be replenished in both period 1 and 2, while a subset consisting of all customers {1,2,3,4,5} is not a feasible subset given the tour duration constraint. This demonstrates that the fractional model provides the opportunity to select suboptimal subsets of customers, causing high integrality gaps. Also note that the instances with the highest integrality gaps do not necessarily have the highest computation times for both types of subset composition constraints.

For the model with a tour duration constraint, the computational results show that valid inequalities decrease computation times and integrality gaps, but that adding both types of valid inequalities does not always improve computation times, compared with adding one type of inequality. The integrality gaps are best if both types of inequalities are used, but only adding the NrVis inequalities yields almost the same average integrality gap. For this subset composition constraint, the best performance in terms of computation times are found if only the ILB inequalities are added to the model.

For the model with the constraint on the number of customers replenished per period, the results show that both types of valid inequalities improve the efficiency of the model. The average computation time decreases strongly for all instance sizes. For the largest instances (15 customers, 6 time periods) the number of solved instances increases from 12 to 17 out of 20 by adding the valid inequalities and the average integrality gap is only a third of the average gap without inequalities. Furthermore, the integrality gap of the model including only the NrVis inequalities is almost identical to the model with both types of inequalities; however, computation times show that the model including both types of inequalities performs better.

Archetti et al. (2007) concluded that the NrVis inequalities (equation 13) are ineffective for solving their IRP model. It is therefore important to note that, for the DJRP-AT, these valid inequalities lower the integrality gaps substantially and reduce the computation time. Hence, for the DJRP-AT, the effectiveness of these inequalities has been demonstrated for both types of subset composition constraints. The results per instance with all valid inequalities in the model are available in Appendix D for both types of subset composition constraints.

6.2. Comparison DJRP-AT and DJRP

The results of the DJRP-AT with a constraint on the number of customers per tour k_M can be compared with the existing DJRP with fixed fees (Boctor et al. 2004). In this model a common cost is paid for serving at least one customer in a period (B = 1000) and an individual cost m_i is incurred for replenishing each customer. Note that the individual replenishment cost would in practice be given by the contract between the supplier and the LSP, and cannot be changed during the execution of the contract. To assess the impact of the individual replenishment costs, we test several values for m_i , $i \in \mathcal{N}$. For the experiments the individual replenishment cost m_i is either set to the same value for all customers (25 and 100) or set according to one of the following schemes. First, we set m_i to a value proportional to the distance to the depot, which we denote by 'prop'. Second, we define a zone around the depot in which at least one-third of the customers is located ('zones'). For the customers within the zone $m_i = 25$, and for the other customers $m_i = 100$. Third, we divide the total area in four quadrants. If a customer is in the same quadrant as the depot $m_i = 25$, and $m_i = 100$ otherwise.

To make a fair comparison between the DJRP and the DJRP-AT, the DJRP is extended with constraints on the customer's inventory capacity and a constraint on the number of customers served per period. Hence, the difference between this model and the DJRP-AT is the cost structure, i.e., fixed fees for individual replenishments versus transportation costs. Both models result in subsets of customers to be served in each period of the planning horizon and the corresponding delivery quantities. Importantly, it is not possible to directly compare the costs of both models. Therefore, the optimal traveling salesman tours for the resulting customer subsets are computed. The tour costs reflect the actual incurred routing costs. The total costs, for both models consisting of the inventory holding cost, the tour costs and the fixed costs per period, will be compared. First, the results of one instance are studied in more detail. Next, aggregated results over all tested instances are analyzed.

6.2.1. Illustrative result for one instance In this section, one of the tested instances is studied in more detail and the results demonstrate the effect of the DJRP-AT compared with the DJRP. Consider an example containing 10 customers (instance abs3n10), 3 time periods, a maximum of $k_M = 7$ customers per period and an individual replenishment cost of $m_i = 25$. In Figure 2, the routes corresponding with the solution of the DJRP and the DJRP-AT are drawn. The depot is indicated by D and the customers are numbered according to the order of the customers in the instance. The black line represents the performed route in period two of the three periods and the grey line represents the route in period three; no customers are replenished in period one in both solutions. The DJRP solution shows that some customers are replenished multiple times in the planning horizon. Moreover, customers that are located in relatively close proximity, are not necessarily replenished in the same period, e.g., customers 7 and 8. The total costs of the DJRP solution are 4399 consisting of inventory holding cost of 123 and routing costs of 4276, including the fixed fee per period. The DJRP-AT solution is clearly more efficient from a routing point of view. The two routes cover distinct areas of the region in which the customers are located and all customers are replenished only once in the planning horizon. The holding cost, 159, is higher than in the solution of the DJRP. The routing costs, including the fixed fee per period, are 3914 which is lower than the DJRP routing costs. The total costs of DJRP-AT solution are 4073, which is 7.4% lower than the total costs of the DJRP solution.

Figure 2 DJRP and DJRP-AT solutions for instance abs3n10 with T = 3, $k_M = 7$ and $m_i = 25$ (black route in period 2, gray route in period 3).



(a) DJRP solution



(b) DJRP-AT solution

6.2.2. Aggregated results Table 2 compares aggregated results of the DJRP-AT and the DJRP for the instances proposed by Archetti et al. (2007). The first columns indicate the number of customers (N), the length of the time horizon (T), the maximum number of customers (k_M) and the number of instances solved out of the five instances (#). For each combination of N, T and k_M five different individual replenishment costs are considered for the DJRP (m = 25, 100, prop, zones, blocks). The average percentage improvement in total costs, the maximum percentage improvement in total costs and the maximum percentage deterioration in any of the instances of the DJRP-AT, compared with the DJRP, are reported for each combination of parameter values. The percentage improvement in total cost of an instance is computed as (cost DJRP - cost DJRP-AT)/cost DJRP.

					A imp	verage	e cost ent (%)			Maximum cost improvement (%)						Maximum cost deterioration (%)					
						m					m										
N	T	k_M	#*	25	100	prop	zones	blocks	25	100	prop	zones	blocks	25	100	prop	zones	blocks			
5	3	3	5	3.8	2.1	2.1	1.9	1.9	8.7	7.8	7.8	7.8	7.8	0.4	0.4	0.4	0.7	0.7			
5	3	4	5	4.2	2.5	2.5	2.3	2.3	8.6	7.2	7.2	7.2	7.2	-	-	-	1.0	1.0			
10	3	5	5	2.3	3.4	1.5	1.5	1.8	9.5	9.5	9.5	9.5	9.5	-	-	2.2	2.2	0.6			
10	3	6	5	3.6	4.8	2.8	3.6	2.9	7.1	7.6	7.1	7.1	7.1	0.4	0.4	2.1	0.4	2.1			
10	3	7	5	4.6	4.1	4.1	4.4	4.2	7.9	8.1	8.1	7.8	8.1	0.4	0.4	0.4	0.4	0.4			
10	3	8	5	5.9	5.4	5.4	5.7	5.6	11.1	8.9	8.9	10.6	9.4	-	-	-	-	-			
15	3	7	5	10.7	9.4	8.9	8.7	8.2	12.2	11.1	10.8	11.6	12.2	-	-	-	-	-			
15	3	8	5	3.0	1.2	-0.5	0.0	1.1	6.7	3.4	0.4	2.7	5.3	0.5	-	3.0	2.6	2.6			
15	3	9	5	3.6	0.7	0.6	1.7	2.5	5.6	4.6	4.1	4.7	5.7	-	1.5	1.5	0.2	0.2			
15	3	10	5	6.1	3.4	3.3	4.3	5.1	11.0	9.6	9.1	9.6	11.1	-	1.0	1.0	0.1	0.1			
15	3	11	5	8.2	5.6	5.5	6.2	7.0	14.4	14.3	13.9	14.4	13.9	-	2.7	2.7	3.2	3.2			
20	3	10	5	6.8	4.9	1.9	4.3	2.8	8.7	8.7	8.7	8.7	8.7	-	-	0.8	-	0.4			
20	3	11	4	2.2	2.4	2.4	2.7	2.7	6.6	7.4	7.4	8.2	8.2	0.5	-	-	-	-			
20	3	12	5	2.6	1.5	1.5	2.2	2.2	8.5	5.2	5.2	8.1	8.1	0.9	-	-	-	-			
20	3	13	5	4.4	3.4	3.4	3.6	3.6	10.1	8.8	8.8	8.8	8.8	0.9	-	-	-	-			
5	6	3	5	5.8	2.6	4.4	4.5	4.5	12.3	5.9	6.6	6.6	6.6	-	-	-	-	-			
5	6	4	5	3.2	1.8	1.2	1.1	1.4	6.2	4.2	4.1	4.2	4.2	-	0.2	1.5	1.5	1.5			
10	6	5	5	5.3	4.8	4.8	4.8	5.3	10.4	7.9	7.9	8.2	10.4	-	-	-	-	-			
10	6	6	5	5.7	5.7	5.9	5.7	5.8	9.5	8.4	8.1	8.9	8.0	-	-	-	-	-			
10	6	5	5	3.0	2.5	2.2	2.1	2.9	6.9	5.3	5.1	4.2	5.8	0.6	-	-	-	-			
10	6	8	5	4.3	2.6	3.2	3.0	4.5	7.6	4.9	5.4	5.0	7.7	-	-	-	-	-			
15	6	8	3	8.6	5.4	6.2	7.7	7.7	11.3	6.1	7.4	11.3	11.3	-	-	-	-	-			
15	6	9	4	8.1	7.1	6.6	7.1	7.9	11.7	9.5	9.3	9.3	11.6	-	-	-	-	-			
15	6	10	5	3.7	2.4	2.4	3.2	3.4	6.1	6.1	6.3	5.2	6.2	-	-	-	-	-			
15	6	11	5	5.1	3.8	3.1	4.3	4.2	7.6	5.3	4.8	5.9	5.5	-	-	-	-	-			
		Ov	erall	5.0	3.7	3.4	3.9	4.1	14.4	14.3	13.9	14.4	13.9	0.9	2.7	3.0	3.2	3.2			

Table 2 Average and maximum improvement and maximum deterioration of DJRP-AT compared with DJRP.

*Number of instances solved out of 5 for each parameter combination.

The results in Table 2 show that incorporating the approximated transportation costs in the DJRP reduces the total cost with 3.4% - 5.0% on average for different schemes for m and 4%

overall. Individual savings up to 14.4% are achieved. The average improvement between instances with different numbers of customers or periods is similar.

For 494 out of 605 optimally solved instances the DJRP-AT outperforms the DJRP and matches its costs for 51 instances. For only 60 instances the DJRP resulted in a slightly better solution than the DJRP-AT with a maximum cost difference of 3.2%, but only 0.96% on average over all *m* schemes. The cases in which the DJRP resulted in lower costs than the DJRP-AT can be explained by using the approximated transportation cost in the optimization, which does not always lead to the lowest routing and inventory holding cost. Moreover, the DJRP completely ignores customer location when determining replenishments and will mostly serve customers on the day their inventory is exhausted, provided all constraints are respected (except if one day's major cost can be saved). If not all customers can be served on the day their inventory is exhausted, the customers that have the lowest holding costs will be served a day earlier. In this case, the customer's holding costs exert substantial influence on the combination of customers served together in the DJRP solutions. This can, coincidentally, result in favorable combinations of customers regarding the actual routing cost, which can result in better DJRP solutions compared with the DJRP-AT. However, the results show that this scenario is unlikely, since this occurs in a limited number of the instances.

Table 2 suggests that the DJRP-AT performs better on problem instances with a longer planning horizon. The DJRP was only able to find a better solution than the DJRP-AT in five parameter settings of the instances with a six day planning horizon, and the savings were 0.2% - 1.5%. Therefore, we did additional experiments in which we increased the planning horizon from three to six days of the instances with a three day planning horizon. The results do not show clearly that the DJRP-AT performs relatively better on instances with a longer planning horizon.

The improvement in total costs of incorporating the transportation costs in the DJRP slightly decreases if the individual ordering cost m in the DJRP increases from 25 to 100. This can be explained by the fact that if the individual replenishment cost is lower, then the number of replenishments is higher in the DJRP outcomes, resulting in higher actual total routing costs. This effect can also be observed in the number of instances for which an improvement, deterioration, or equal costs are reported (Table 3). The number of instances showing an improvement decreases as the value of the individual replenishment cost m_i increases, compared with the DJRP. However, the percentage of instances for which deterioration must be reported does not increase as m_i increases; instead, the percentage of instances with equal costs for both models increases.

By using proportional costs, zones and blocks based costs instead of the same individual replenishment costs for all customers, the routing costs are better reflected in the DJRP. Indeed, we can observe that on average the improvement of the DJRP-AT over the DJRP is lower than for the DJRP with $m_i = 25$. However, for $m_i = 100$ this is only the case for the proportional individual replenishment costs. The proportional costs best reflect the actual routing costs, in several works in the IRP literature the direct distance is used (as a starting point) to replace actual routing costs (see for example Cordeau et al. (2015) and Absi et al. (2015)). Still, from our results it shows that the DJRP-AT outperforms using proportional costs for replenishing a customer.

		Ν	lumber	r of insta	ances		Percentage							
			m				m							
	25	100	prop	zones	blocks	total	25	100	prop	zones	blocks	total		
Improvement	108	97	92	99	98	494	89%	80%	76%	82%	81%	82%		
Detoriation	8	9	14	14	15	60	7%	7%	12%	12%	12%	10%		
Equal	5	15	15	8	8	51	4%	12%	12%	7%	7%	8%		
Total	121	121	121	121	121		100%	100%	100%	100%	100%			

Table 3Number and percentage of instances that report improvement, deterioration and equal costs.

In conclusion, these results show that in approximately 82% of all solved instances lower total costs can be achieved by using the DJRP-AT, instead of the DJRP. If the individual replenishment costs increase, the improvement that the DJRP-AT can achieve decreases, however, the number of instances with reported deteriorations does not increase. Also, for individual replenishment costs that better reflect the actual routing cost than the same cost for all customers, the DJRP-AT still outperforms the DJRP.

6.3. Comparison DJRP-AT and IRP

To gain further insight in the quality of the solutions of the DJRP-AT, we compare the results of the DJRP-AT to the results of a problem formulation that includes the actual routing problem. This formulation is a variant of the IRP with a constraint on the number of customers in a route, hence, this problem is different than the IRP often addressed in the literature (Coelho et al. 2014). Therefore, to solve this IRP, the labeling algorithm that solves the Pricing Problem of the DJRP-AT is replaced by an Integer Linear Program (ILP) which solves a resource constrained elementary shortest path problem. The resource is the number of customers in the route and note that there are arcs with negative cost, hence, negative cost cycles need to be prevented by adding subtour elimination constraints. A solution of the ILP is a route with corresponding costs that consists of the arc costs and the fixed costs B. We use Gurobi to solve the ILP and apply the 'Solution Pool' option to generate multiple solutions in one iteration. Since the solution method is not especially designed for solving this IRP, only some of the very small instances, with low values for k_M , can be solved within four hours of running time. The differences in computation time are therefore only indicative. Table 4 gives the aggregated results of the comparison between the DJRP-AT and the IRP, the results per instance can be found in Appendix D. As in Section 6.2, the optimal traveling salesman tours are computed to compare the costs. For each combination of parameters, the number of instances solved, the number of instances with equal results for the DJRP-AT and the IRP, the average difference between the DJRP-AT and the IRP, and the average computation times of both models are indicated. For 10 customers, 3 time periods and $k_M = 7$, none of the instances could be solved within four hours of running time. For 10 customers, 6 time periods and $k_M = 6$ only one instance was solved within four hours. In total 35 instances of the IRP are solved to optimality with computation times that are several orders of magnitude higher than those of the DJRP-AT. Out of the 35 instances, for 16 instances the DJRP-AT gives the same result as the IRP. On average, the costs of the Solutions of the IRP are 0.77% lower than the costs of the DJRP-AT. Considering that solutions of the DJRP-AT are found by only using an approximation for the transportation costs, the results are quite close to the exact solutions.

N	T	k_M	Number of instances solved	Number of instances equal result	Average difference (%)	Average time DJRP-AT (s)	Average time IRP (s)
5	3	3	5	3	-0.28	0.0	1.0
5	3	4	5	1	-1.30	0.2	1.6
10	3	5	5	4	-0.54	0.6	1155.8
10	3	6	4	1	-0.56	0.4	9348.8
5	6	3	5	4	-0.23	0.4	4.6
5	6	4	5	0	-1.47	0.4	13.4
10	6	5	5	3	-0.61	11.0	4944.0
10	6	6	1	0	-1.19	34.4	13738.0

 Table 4
 Average difference of DJRP-AT compared with IRP.

6.4. Case study ATM replenishment in Amsterdam

As described in the introduction, our research is motivated by a real-life case in ATM replenishment in the Netherlands, in which a supplier (vendor) decides on which ATMs to replenish per day and an LSP (CIT) designs the routes to perform the replenishments. Currently, the supplier only pays a 'minor' transportation cost to the LSP for each ATM replenishment and no 'major' cost. In this section we use company data to illustrate the benefit of alternative replenishment cost structures based on the DJRP and the proposed DJRP-AT.

Data on ATMs in Amsterdam and the depot location are provided by the supplier. The dataset contains per ATM, the address, storage capacity, dynamic daily demand, and initial inventory level. To use the existing solution framework, the ATM locations are mapped on the Euclidean plane and the demand data is expressed in thousands of Euros. Because the future cost structure parameters are not available and current numbers are not disclosed because of confidentiality reasons, the cost parameters are determined in consultation with the company to reflect the expected ratio between transportation and inventory holding costs. This includes minor (m_i) and major (B) transportation cost and inventory holding rates. A three-day planning horizon is considered appropriate and therefore we select the ATMs that need replenishment within the next three days, which results in 75 ATMs. Based on their geographical locations and postal codes, the set of ATMs is naturally split into four subsets of sizes 16, 19, and two of 20 customers. We let the holding rate vary from 0.1 to 0.3 to represent realistic cost ratios. Based on service times and travel times as observed by the company, we let the maximum number of customers served k_M range from 10 to 13. The same individual replenishment cost m_i is used for all customers, and tests are performed for values 25, 50 and 100. We only use the same values for m_i for all customers to stay close to the real-life case.

We estimate the costs of the current situation at the company by solving the DJRP with the major cost set to zero (B = 0, denoted by DJRP₀). To make a fair comparison with the DJRP with major cost (denoted by DJRP_B) and the DJRP-AT, we subsequently add the major cost B = 1000 for each period in which a replenishment takes place. Note that in practice, the current minor cost should be higher than the future minor cost to cover fixed costs. After computing the solutions of the three models, the optimal traveling salesman tours are computed to compare the costs.

Table 5 reports the percentage cost improvements of the $DJRP_B$ over the $DJRP_0$, and of the DJRP-AT over the DJRP_B. We only report the results for m = 25 and m = 100 since the results for m = 50 are similar to those for m = 100. Comparing DJRP₀ and DJRP_B shows that a substantial cost improvement of 28.6% on average can be obtained, caused by having a route in every period in the $DJRP_0$ while in the $DJRP_B$ often only two routes are used. For the comparison between $DJRP_B$ and DJRP-AT, the results show that for 136 out of 144 cases, the DJRP-AT results in lower total costs than the $DJRP_B$, with a decrease in total costs up to 12.1%. For the remaining eight instances, the DJRP-AT results in slightly higher costs than the DJRP_B, with differences up to 1.7%. The DJRP-AT solutions are, on average, 6.4% better than the DJRP_B solutions. Table 5 shows that the improvement of the DJRP-AT over the $DJRP_B$ decreases slightly for higher holding rates. For the DJRP-AT, it can be beneficial to serve customers earlier in the planning horizon than waiting until the customers run out of stock. However, if the holding rate is higher, serving customers earlier becomes more costly which results in higher costs for the DJRP-AT. Interestingly, varying the individual replenishment cost per ATM does not have a significant impact on the results. Furthermore, Table 5 shows that varying the maximum number of customers has a different impact per region. For Region 1 the results are similar for all values of k_M , while for Region 2, increasing the maximum number of customers to $k_M = 13$ leads to higher costs for the DJRP-AT, compared with the DJRP_B. For Region 2, the lower cost solution for $k_M = 13$ than for $k_M = 12$ of the $DJRP_B$ can be explained by the fact that many customers in the region are located in close

proximity and a few customers are located further away from these clustered customers. In the DJRP-AT solutions, the customers not in the cluster are served on the same day, for all values of k_M . In the DJRP_B solution, for $k_M = 12$, the customers outside the cluster are not all served on one day, which results in high transportation cost, but if $k_M = 13$, these customers are served on the same day, which lowers the total costs of the DJRP_B substantially, as opposed to $k_M = 12$.

Table 5Percentage improvement $DJRP_B$ over $DJRP_0$ (left), and percentage improvement DJRP-AT over $DJRP_B$ (right) in Amsterdam case.

	Region 1 m		Regi	on 2	Regi	on 3	Regi	on 4	Regi	on 1	Regi	on 2	Reg	ion 3	Regi	on 4
h = 0.1			m - m - m		\overline{m}		m		\overline{m}		\overline{m}		\overline{m}		m	
k_M	25	100	25	100	25	100	25	100	25	100	25	100	25	100	25	100
10	30.1	30.1	30.2	30.2	31.1	31.1	28.2	28.2	11.4	11.4	4.0	4.0	1.1	1.1	8.3	8.3
11	30.1	30.1	30.3	30.3	29.3	29.3	27.8	27.8	11.4	11.4	4.4	4.4	4.3	4.3	8.1	8.1
12	30.1	30.1	30.1	30.1	29.2	29.2	24.6	24.6	11.4	11.4	3.1	3.1	4.3	4.3	12.1	12.1
13	30.1	30.1	32.4	32.4	29.3	29.3	25.0	25.0	11.4	11.4	-0.7	-0.7	4.6	4.6	11.5	11.5
h = 0.2	r	m m		n	m		m		m		m		m		m	
k_M	25	100	25	100	25	100	25	100	25	100	25	100	25	100	25	100
10	29.6	29.6	29.1	29.4	30.3	30.3	27.7	27.7	10.8	10.8	3.6	3.6	0.9	0.9	7.7	7.7
11	29.6	29.6	29.2	29.4	28.6	28.6	27.3	27.3	10.8	10.8	2.7	2.7	3.9	3.9	7.3	7.3
12	29.6	29.6	29.0	29.3	28.5	28.5	24.1	24.1	10.8	10.8	2.6	2.6	4.0	4.0	11.3	11.3
13	29.6	29.6	31.4	31.6	28.6	28.6	24.6	24.6	10.8	10.8	-1.2	-1.2	4.0	4.0	10.6	10.6
h = 0.3	r	n	r	n	m		m		m		m		m		n	ı
k_M	25	100	25	100	25	100	25	100	25	100	25	100	25	100	25	100
10	30.1	29.1	26.9	28.6	29.5	29.6	26.9	27.2	9.8	10.3	3.8	3.3	0.7	0.7	7.2	7.2
11	29.8	29.1	27.3	28.7	29.5	27.9	26.5	26.8	10.3	10.3	2.4	2.4	1.4	3.6	6.8	6.8
12	30.0	29.1	27.9	28.5	27.8	27.9	23.4	23.7	10.0	10.3	1.0	2.0	3.7	3.6	11.0	11.0
13	30.0	29.1	27.8	30.8	27.8	28.0	23.8	24.2	10.0	10.3	0.4	-1.7	3.7	3.5	10.3	10.3

The results of DJRP_0 , DJRP_B and DJRP-AT are visualized for Region 1 in Figure 3, with $k_M = 11, m = 100$, and h = 0.2. The cost improvements for this instance are 29.6% and 10.8%, respectively. The results show that, as expected, the DJRP_B saves cheaper solutions than the DJRP_0 and that the routes given by the DJRP-AT are much more efficient than the routes given by the DJRP_B . Overall, the results of the case study show that using a DJRP cost structure could provide significant savings, and the DJRP-AT results in similar improvements for the real-life case as for the artificial benchmark instances.

7. Conclusion

In practice it is regularly the case that a supplier outsources customer deliveries to a Logistics Service Provider (LSP); the supplier often pays a fixed transportation fee to the LSP for this service. Hence, when deciding on the timing of customer replenishments, the supplier often does not take efficiency of the delivery routes into account. To optimize costs, suppliers can use joint replenishment models, such as the Dynamic-Demand Joint Replenishment Problem (DJRP) (Boctor et al.

Figure 3 Inventory based replenishment, DJRP_B and DJRP-AT solutions for Region 1 ($k_M = 11$, m = 100, h = 0.2, black route in period 1, gray route in period 2, dotted route in period 3).



2004) in case of dynamic demand. The DJRP minimizes inventory holding and replenishment costs while ensuring that customers do exhaust their stock. The replenishment costs consist of fixed fees that are independent of the actual routing costs. As a result, we argue that the DJRP is incapable of proposing efficient solutions from a transportation point of view. Although the fee per delivery is fixed, the costs of the inefficient routes are indirectly paid by the supplier via the negotiated delivery fees in the following contract. Inspired by the practical relevance of the DJRP, this paper proposes the Dynamic-Demand Joint Replenishment Problem with Approximated Transportation Costs (DJRP-AT) in which transportation costs are included by approximating the optimal tour length for given subsets of customers. Using the DJRP-AT will result in lower total costs and will increase resource utilization.

We propose a mathematical model for the DJRP-AT which is enriched with two different types of constraints. First, the tour duration is restricted to resemble the limitations encountered in practice. Second, the number of customers served per period is bounded, resulting in the opportunity to investigate the improvement of the proposed DJRP-AT compared with the existing DJRP. The Dantzig-Wolfe decomposition is applied to the proposed compact formulation and the resulting Master and Pricing Problems are solved in a Branch-and-Cut-and-Price framework. The Master Problem defines which subset of customers to replenish per period of the planning horizon and determines the delivery quantities of the customers served. The Pricing Problem generates, via a specially designed labeling algorithm, subsets of customers that can be served in one period and the transportation costs are then approximated for these customer subsets. To increase efficiency, labels need to be discarded during the labeling algorithm. However, existing sufficient dominance rules for shortest path problems are not adequate for discarding labels in the Pricing Problem discarding possible. Costs of the DJRP and DJRP-AT are compared by computing the optimal traveling salesman tours for the subsets of customers selected in the solutions of both models.

To assess the value of the DJRP-AT formulation and solution framework, existing problem instances from the Inventory Routing Problem (IRP) are adjusted for our experiments. The effectiveness of two types of valid inequalities that were proposed for the IRP (Archetti et al. 2007) is tested for both types of extra constraints. The results show that both inequalities are effective for the DJRP-AT, which differs from the results obtained from the IRP. Computational experiments show average improvements of total transportation and inventory holding costs of 3.4% to 5%, respectively. Depending on the individual fixed fee charged in the DJRP, maximum improvements around 14.4% are obtained. The DJRP outperformed the DJRP-AT for only a few instances, due to the approximation of the tour length. The DJRP-AT solutions are also compared to the equivalent IRP solutions for which a different Pricing Problem is implemented. The results show that for the solved instances, the costs of the IRP solutions are on average only 0.77% lower than the costs of the DJRP-AT solutions. The computation times for the IRP are orders of magnitude higher than for the DJRP-AT. Analysis of a real-life case in ATM replenishment shows that significant cost reductions can be achieved for both the LSP and the supplier when using the DJRP-AT.

Computational results with the DJRP-AT show that calculating transportation costs, instead of using fixed fees, in joint replenishment is worthwhile and that approximation of transportation costs works well. Future research could focus on developing novel formulations for this problem, possibly inspired by formulations discussed in Narayanan and Robinson (2006), that may improve the integrality gaps. Moreover, new valid inequalities can be proposed to strengthen the linear relaxation of DJRP-AT models.

Acknowledgments

This work was supported by the Netherlands Organisation for Scientific Research (NWO) [project number 407-13-050].

References

- Absi N, Archetti C, Dauzère-Pérès S, Feillet D (2015) A two-phase iterative heuristic approach for the production routing problem. *Transportation Science* 49(4):784–795.
- Andersson H, Hoff A, Christiansen M, Hasle G, Løkketangen A (2010) Industrial aspects and literature survey: Combined inventory management and routing. *Computers & Operations Research* 37(9):1515– 1536.
- Anily S, Tzur M (2005) Shipping multiple items by capacitated vehicles: An optimal dynamic programming approach. *Transportation Science* 39(2):233–248.

- Archetti C, Bertazzi L, Laporte G, Speranza MG (2007) A Branch-and-Cut algorithm for a vendor-managed inventory-routing problem. *Transportation Science* 41(3):382–391.
- Archetti C, Bianchessi N, Irnich S, Speranza MG (2014) Formulations for an inventory routing problem. International Transactions in Operational Research 21(3):353–374.
- Arkin E, Joneja D, Roundy R (1989) Computational complexity of uncapacitated multi-echelon production planning problems. Operations Research Letters 8(2):61–66.
- Beardwood J, Halton JH, Hammersley JM (1959) The shortest path through many points. *Mathematical Proceedings of the Cambridge Philosophical Society* 55(4):299–327.
- Boctor FF, Laporte G, Renaud J (2004) Models and algorithms for the dynamic-demand joint replenishment problem. *International Journal of Production Research* 42(13):2667–2678.
- Çavdar B, Sokol J (2015) A distribution-free tsp tour length estimation model for random graphs. European Journal of Operational Research 243(2):588–598.
- Chien TW (1992) Operational estimators for the length of a traveling salesman tour. Computers & Operations Research 19(6):469–478.
- Coelho LC (n.d.) Problem instances inventory routing. URL http://www.leandro-coelho.com/instances/ inventory-routing/, accessed: 24 May 2016.
- Coelho LC, Cordeau JF, Laporte G (2014) Thirty years of inventory routing. *Transportation Science* 48(1):1–19.
- Cordeau JF, Laganà D, Musmanno R, Vocaturo F (2015) A decomposition-based heuristic for the multipleproduct inventory-routing problem. *Computers & Operations Research* 55:153–166.
- Cunha JO, Melo RA (2016) On reformulations for the one-warehouse multi-retailer problem. Annals of Operations Research 238(1):99–122.
- Daganzo CF (1984a) The distance traveled to visit n points with a maximum of c stops per vehicle: An analytic model and an application. *Transportation science* 18(4):331–350.
- Daganzo CF (1984b) The length of tours in zones of different shapes. Transportation Research Part B: Methodological 18(2):135–145.
- Dasci A, Verter V (2001) A continuous model for production–distribution system design. *European Journal* of Operational Research 129(2):287–298.
- Desrosiers J, Lübbecke ME (2005) A primer in column generation. Desaulniers G, Desrosiers J, Solomon MM, eds., Column Generation, 1–32 (Springer).
- Eilon S, Watson-Gandy CDT, Christofides N (1971) Distribution management: Mathematical Modelling & Practical Analysis (Griffin London).
- Federgruen A, Meissner J, Tzur M (2007) Progressive interval heuristics for multi-item capacitated lot-sizing problems. Operations Research 55(3):490–502.

- Feillet D, Dejax P, Gendreau M, Gueguen C (2004) An exact algorithm for the elementary shortest path problem with resource constraints: Application to some vehicle routing problems. *Networks* 44(3):216– 229.
- Figliozzi M (2008) Planning approximations to the average length of vehicle routing problems with varying customer demands and routing constraints. Transportation Research Record: Journal of the Transportation Research Board 2089:1–8.
- Franceschetti A, Jabali O, Laporte G (2017) Continuous approximation models in freight distribution management. *TOP* 25(3):413–433.
- Haimovich M, Rinnooy Kan AHG (1985) Bounds and heuristics for capacitated routing problems. Mathematics of Operations Research 10(4):527–542.
- Hariga M, Gumus M, Daghfous A, Goyal SK (2013) A vendor managed inventory model under contractual storage agreement. Computers & Operations Research 40(8):2138–2144.
- Hemmelmayr V, Doerner KF, Hartl RF, Savelsbergh MWP (2009) Delivery strategies for blood products supplies. OR spectrum 31(4):707–725.
- Hindle A, Worthington D (2004) Models to estimate average route lengths in different geographical environments. Journal of the Operational Research Society 55(6):662–666.
- Hoque MA (2006) An optimal solution technique for the joint replenishment problem with storage and transport capacities and budget constraints. *European Journal of Operational Research* 175(2):1033– 1042.
- Jabali O, Gendreau M, Laporte G (2012) A continuous approximation model for the fleet composition problem. *Transportation Research Part B: Methodological* 46(10):1591–1606.
- Jacobs T, Longo S (2014) A new perspective on the windows scheduling problem. *CoRR* arXiv preprint arXiv:1410.7237.
- Kang HY, Lee AHI, Wu CW, Lee CH (2016) An efficient method for dynamic-demand joint replenishment problem with multiple suppliers and multiple vehicles. *International Journal of Production Research* 1–20.
- Khouja M, Goyal S (2008) A review of the joint replenishment problem literature: 1989–2005. European Journal of Operational Research 186(1):1–16.
- Kwon O, Golden B, Wasil E (1995) Estimating the length of the optimal tsp tour: An empirical study using regression and neural networks. *Computers & Operations Research* 22(10):1039–1046.
- Langevin A, Mbaraga P, Campbell JF (1996) Continuous approximation models in freight distribution: An overview. Transportation Research Part B: Methodological 30(3):163–188.
- Langevin A, Soumis F (1989) Design of multiple-vehicle delivery tours satisfying time constraints. Transportation Research Part B: Methodological 23(2):123–138.

- Larrain H, Coelho LC, Cataldo A (2017) A variable MIP neighborhood descent algorithm for managing inventory and distribution of cash in automated teller machines. *Computers & Operations Research* 85:22–31.
- Larsen C, Turkensteen M (2014) A vendor managed inventory model using continuous approximations for route length estimates and markov chain modeling for cost estimates. *International Journal of Produc*tion Economics 157:120–132.
- Lübbecke ME, Desrosiers J (2005) Selected topics in column generation. Operations Research 53(6):1007–1023.
- Narayanan A, Robinson EP (2006) More on 'Models and algorithms for the dynamic-demand joint replenishment problem'. *International Journal of Production Research* 44(2):383–397.
- Narayanan A, Robinson P (2010) Efficient and effective heuristics for the coordinated capacitated lot-size problem. European Journal of Operational Research 203(3):583–592.
- Nemhauser GL, Park S (1991) A polyhedral approach to edge coloring. *Operations Research Letters* 10(6):315–322.
- Olsen AL (2008) Inventory replenishment with interdependent ordering costs: An evolutionary algorithm solution. *International Journal of Production Economics* 113(1):359–369.
- Rahmouni M, Hennet JC (2015) Determining the optimal routes in the multi-product multi-site joint delivery problem. 2015 International Conference on Industrial Engineering and Systems Management (IESM), 1361–1366 (IEEE).
- Robinson EP, Narayanan A, Gao LL (2007) Effective heuristics for the dynamic demand joint replenishment problem. Journal of the Operational Research Society 58(6):808–815.
- Robinson P, Narayanan A, Sahin F (2009) Coordinated deterministic dynamic demand lot-sizing problem: A review of models and algorithms. *Omega* 37(1):3–15.
- Senoussi A, Mouss NK, Penz B, Brahimi N, Dauzère-Pérès S (2016) Modeling and solving a one-supplier multi-vehicle production-inventory-distribution problem with clustered retailers. The International Journal of Advanced Manufacturing Technology 85(5):971–989.
- Shen ZJM, Qi L (2007) Incorporating inventory and routing costs in strategic location models. *European Journal of Operational Research* 179(2):372–389.
- Solyalı O, Süral H (2012) The one-warehouse multi-retailer problem: Reformulation, classification, and computational results. Annals of Operations Research 196(1):517–541.
- Turkensteen M, Klose A (2012) Demand dispersion and logistics costs in one-to-many distribution systems. European Journal of Operational Research 223(2):499–507.
- Van Anholt RG, Coelho LC, Laporte G, Vis IFA (2016) An inventory-routing problem with pickups and deliveries arising in the replenishment of automated teller machines. *Transportation Science* 50(3):1077– 1091.

- Ventura JA, Valdebenito VA, Golany B (2013) A dynamic inventory model with supplier selection in a serial supply chain structure. *European Journal of Operational Research* 230(2):258–271.
- Wang L, He J, Wu D, Zeng YR (2012) A novel differential evolution algorithm for joint replenishment problem under interdependence and its application. *International Journal of Production Economics* 135(1):190–198.
- Webb IR, Buzby BR, Campbell GM (1997) Cyclical schedules for the joint replenishment problem with dynamic demands. *Naval Research Logistics (NRL)* 44(6):577–589.