



Correction to: The model structure of the copper-dependent ammonia Monooxygenase

Francesco Musiani¹ · Valquiria Broll¹ · Elisa Evangelisti¹ · Stefano Ciurli¹

Published online: 27 February 2021
© Society for Biological Inorganic Chemistry (SBIC) 2021

Correction to: JBIC Journal of Biological Inorganic Chemistry (2020) 25:995–1007
<https://doi.org/10.1007/s00775-020-01820-0>

In the original article published, the authors missed to acknowledging an article published by Liew et al. (Mutagenesis of the hydrocarbon monooxygenase indicates a metal centre in subunit-C, and not subunit-B, is essential for copper-containing membrane monooxygenase activity, *Microbiology* **2014**, 160, 1267–1277).

“Thus far, however, the crystal structures have not fully established the location and composition of the pMMO active site [48], but all evidence points to either the CuB or the CuC site for this role. It is the opinion of the authors of the present study that the latter, with its labile water-bound

position, should more logically constitute the enzyme active metal site. This conclusion is strongly supported by site-directed mutagenesis studies on the copper-dependent hydrocarbon monooxygenase (HMO) from *Mycobacterium* NBB4 which provided the first evidence that the C site was essential for activity, whereas mutations in the B site impaired, but did not eliminate, activity (E. F. Liew, D. Tong, N. V. Coleman, A. J. Holmes, *Microbiology* 2014, 160, 1267–1277)”.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00775-020-01820-0>.

✉ Francesco Musiani
francesco.musiani@unibo.it

✉ Stefano Ciurli
stefano.ciurli@unibo.it

¹ Laboratory of Bioinorganic Chemistry, Department of Pharmacy and Biotechnology, University of Bologna, Viale G. Fanin 40, 40127 Bologna, Italy