



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

A Combined Approach Employing Chlorotoxin-Nanovectors and Low Dose Radiation to Reach Infiltrating Tumor Niches in Glioblastoma

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

A Combined Approach Employing Chlorotoxin-Nanovectors and Low Dose Radiation to Reach Infiltrating Tumor Niches in Glioblastoma / Tamborini, Matteo; Locatelli, Erica; Rasile, Marco; Monaco, Ilaria; Rodighiero, Simona; Corradini, Irene; Comes Franchini, Mauro; Passoni, Lorena; Matteoli, Michela. - In: ACS NANO. - ISSN 1936-0851. - STAMPA. - 10:2(2016), pp. 2509-2520. [10.1021/acsnano.5b07375]

Availability:

This version is available at: <https://hdl.handle.net/11585/560803> since: 2016-09-01

Published:

DOI: <http://doi.org/10.1021/acsnano.5b07375>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

A Combined Approach Employing Chlorotoxin-Nanovectors and Low Dose Radiation To Reach Infiltrating Tumor Niches in Glioblastoma

Matteo Tamborini, Erica Locatelli, Marco Rasile, Ilaria Monaco, Simona Rodighiero, Irene Corradini, Mauro Comes Franchini, Lorena Passoni, and Michela Matteoli

***ACS Nano* 2016 10 (2), 2509-2520 .**

The final published version is available online at :
<http://dx.doi.org/10.1021/acsnano.5b07375>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

A COMBINED APPROACH EMPLOYING CHLOROTOXIN-NANOVECTORS AND LOW DOSE RADIATION TO REACH INFILTRATING TUMOR NICHES IN GLIOBLASTOMA

Journal:	<i>ACS Nano</i>
Manuscript ID	nn-2015-07375s.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Tamborini, Matteo; University of Milan, Department of Medical Biotechnology and Translational Medicine Locatelli, Erica; University of Bologna, Department of Industrial Chemistry "Toso Montanari" Rasile, Marco; University of Milan, Department of Medical Biotechnology and Translational Medicine; Humanitas Clinical and Research Center, Monaco, Ilaria; University of Bologna, Department of Industrial Chemistry "Toso Montanari" Rodighiero, Simona; Fondazione Filarete, Corradini, Irene; University of Milan, Department of Medical Biotechnology and Translational Medicine Comes Franchini, Mauro; University of Bologna, Department of Industrial Chemistry "Toso Montanari" Passoni, Lorena; University of Milan, Department of Medical Biotechnology and Translational Medicine; Fondazione Filarete, Matteoli, Michela; Humanitas Clinical and Research Center,

SCHOLARONE™
Manuscripts

1
2
3
4 **A COMBINED APPROACH EMPLOYING CHLOROTOXIN-NANOVECTORS AND LOW DOSE**
5 **RADIATION TO REACH INFILTRATING TUMOR NICHES IN GLIOBLASTOMA**
6
7

8
9 Matteo Tamborini [§], Erica Locatelli [¶], Marco Rasile ^{§#}, Ilaria Monaco [¶], Simona Rodighiero
10 [¥], Irene Corradini [§], Mauro Comes Franchini [¶], Lorena Passoni ^{§¥}, Michela Matteoli ^{* #}
11
12
13
14
15

16 **Affiliations:**
17

18 [§] Department of Medical Biotechnology and Translational Medicine, Via Vanvitelli 32,
19 20129 Milano, Italy
20

21 [¶] Department of Industrial Chemistry “Toso Montanari”, University of Bologna, Viale
22 Risorgimento 4, 40136 Bologna, Italy.
23
24

25 [¥] Fondazione Filarete, Viale Ortles 22/4, 20139 Milano, Italy
26

27 [#] Humanitas Research Hospital, Laboratory of Pharmacology and Brain Pathology, Via
28 Manzoni 56, Rozzano, 20089 Milano, Italy
29

30 ^{*} CNR Institute of Neuroscience, Via Vanvitelli 32, 20129 Milano, Italy
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Glioblastoma multiforme (GBM) is the most aggressive form of glioma, with life expectancy of around two years after diagnosis, due to recidivism and to the blood brain barrier (BBB) limiting the amount of drugs which reach the residual malignant cells, thus contributing to the failure of chemotherapies. To bypass the obstacles imposed by the BBB we investigated the use of nanotechnologies combined with radiotherapy, as a potential therapeutic strategy for GBM. We used poly(lactic-co-glycolic acid) (PLGA) nanoparticles (PNP) conjugated to chlorotoxin (CTX), a peptide reported to bind selectively to glioma cells. Silver nanoparticles were entrapped inside the functionalized nanoparticles (Ag-PNP-CTX), to allow detection and quantification of the cellular uptake by confocal microscopy, both *in vitro* and *in vivo*. *In vitro* experiments performed with different human glioblastoma cell lines showed higher cytoplasmic uptake of Ag-PNP-CTX, with respect to non-functionalized nanoparticles. *In vivo* experiments showed that Ag-NP-CTX efficiently targets the tumor, but are scarcely effective in crossing the blood brain barrier in the healthy brain, where dispersed metastatic cells are present. We show here that a single whole brain x-rays irradiation, performed 20 hrs before nanoparticle injection, enhances the expression of the CTX targets, MMP-2 and CIC-3 and, through BBB permeabilization, potently increases the amount of internalized Ag-PNP-CTX even in dispersed cells, and generated an efficient anti-tumor synergistic effect able to inhibit *in vivo* tumor growth. Notably, the application of Ag-PNP-CTX to irradiated tumor cells decreases the extracellular activity of MMP-2. By targeting dispersed GBM cells and reducing MMP-2 activity, the combined use of CTX-nanovectors with radiotherapy may represent a promising therapeutic approach towards GBM.

KEYWORDS: glioblastoma, chlorotoxin-targeted nanovectors, drug delivery, ionizing radiation, MMP-2

1
2
3
4 Glioblastoma (GBM) is the most common primary brain tumor, highly proliferative and invasive,
5 characterized by remarkable biological heterogeneity and poor response to present treatments.¹
6 Even when treated aggressively, GBM has the tendency to recur and this is the main cause of the
7 poor prognosis. The area of recurrence develops at the borders of the surgical cavity, over 90% of
8 the cases within 2 cm of the original tumor, or at distant sites within the brain at later stage.² The
9 median overall survival of GBM patients with the best currently available therapy is approximately
10 15 months, and the 5-year survival rate is only 9.8% at present.³
11
12
13
14
15

16
17 A major hurdle in the clinical management of GBM is the presence of the blood-brain barrier (BBB)
18 which prevents pharmaceutical molecules entering the brain tissue.^{4,5} Large and advanced brain
19 tumor exhibit disrupted BBB integrity. However peripheral areas of brain tumors contain regions
20 with intact BBB leading to the creation of favorable niches of cancer cell spreading and treatment
21 resistance.⁶ Thus, reaching dispersed and invading GBM cells remains a grand challenge for the
22 successful treatment of GBM.
23
24
25
26

27
28 Given the intrinsic difficulty for most conventional drugs to reach the brain, the use of
29 nanotechnological devices and nanoparticles has been suggested as an innovative therapeutic
30 strategy.⁷ In particular, drug-loaded nanocarriers stably incorporating cytotoxic molecules have
31 been proposed as promising tools. Indeed, nanovectors protect the drugs from metabolic
32 degradation and fast clearance, prolonging circulation lifetime and thus ensuring a higher tumor
33 delivery, reduce unwanted side effects and escape the efflux mechanisms determining tumor cell
34 drug resistance.^{8,9}
35
36
37
38

39
40 Due to their tunable characteristics, controlled- and sustained-release properties, biodegradability
41 and biocompatibility, polymeric nanoparticles (PNPs) represent a powerful nanocarrier for the
42 delivery of drug and/or diagnostic agents. Poly(lactic)-co-glycolic-co-poly ethylene glycol (PLGA-*b*-
43 PEG) copolymer¹⁰ has gained large attention in the past decade, since its approval by Food and
44 Drug Administration (FDA) for biomedical usage due to its ability to create micelles-like
45 nanocarriers displaying an internal hydrophobic PLGA portion, suitable for entrapping lipophilic
46 molecules, and an external hydrophilic PEG forming a stabilizing shell.¹¹ However, PLGA
47 nanoparticles, which are typically 100-200 nm in diameter, even when locally administered by
48 convection-enhanced delivery (CED) have limited mobility and diffusion by convection through the
49 brain interstitial spaces,^{12,13} which are 38-64 nm in normal brain¹⁴ and 70-100 nm in regions with
50 tumor.¹⁵
51
52
53
54
55
56
57
58
59
60

1
2
3 The presence of targeting ligands on surface of drug-loaded nanocarriers enhances tumor-specific
4 intracellular uptake and drug accumulation, thus improving the nanovectors therapeutic efficacy.¹⁶
5 It has been reported that Chlorotoxin (CTX) can function as targeting moiety highly specific for
6 GBM.¹⁷
7
8
9

10 CTX is a 36-amino acid peptide originally isolated from *Leiurus quinquestriatus* venom that
11 specifically and selectively binds to glioma cells.^{17,18} Identified CTX target molecules are the matrix
12 metalloproteinase-2 (MMP-2)¹⁹ and the chloride channel-3 (CIC-3)²⁰ both absent in normal brain
13 tissue but abundantly expressed in glioma cells. CIC-3 and MMP-2 form a protein complex which is
14 targeted by the CTX-peptide, and thought to inhibit glioma cell invasion by induction of the
15 endocytosis of the MMP-2/CIC-3 protein complex.^{19,21}
16
17
18
19

20
21 Although CTX-coupled nanoparticles are considered a promising system for tumor treatment,
22 isolated, infiltrating GBM tumor cells are hardly reached by functionalized nanoparticles, mostly
23 due to the integrity of the BBB at the peripheral tumor sites. Given that ionizing radiations damage
24 BBB integrity^{22,23} and induce MMP-2 expression,²⁴⁻²⁶ we hypothesized that they could enhance
25 CTX-nanovectors, by increasing the CTX target expression and facilitating the nanovector brain
26 penetration through BBB permeabilization. Also, we reasoned that, by promoting MMP-2 inhibition,
27 application of CTX-nanoparticles could in turn limit tumor invasiveness, which is described to
28 increase upon Radiation Therapy (RT).^{27,28} As RT represents one of the mainstays of GBM
29 treatment, here we evaluated whether radiotherapy could benefit by the concomitant administration
30 of CTX-targeted nanovectors and vice versa.
31
32
33
34
35
36
37

38 By intra-tumoral and intra-cellular quantification of CTX-targeted nanovectors we demonstrated
39 that radiations, even at low doses, improve GBM tumor accumulation of CTX-nanovectors and are
40 crucial to reach infiltrating and spreading tumor cells. The increased amount of CTX-nanovectors
41 accumulated after radiation treatment within the tumor was sufficient to determine a detectable and
42 significant reduction of tumor growth and MMP-2 enzymatic activity. Thus, the association of RT
43 with the use of CTX-functionalized drug-loaded nanovectors appears to represent a promising
44 strategy for GBM treatment.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS AND DISCUSSION

Synthesis and Characterization of Ag-PNP-CTX nanovectors

With the intent to quantify chlorotoxin (CTX)-targeted nanovectors reaching the tumor microenvironment, and to track their localization into intracellular compartments, we synthesized poly (lactic)-co-glycolic)-co-poly ethylene glycol (PLGA-*b*-PEG) nanovectors conjugated to CTX and encapsulating silver nanoparticles (Ag-nps), visible with optical microscope imaging in reflection mode (Ag-PNP-CTX).

Ag-PNP-CTX were prepared following a procedure already reported by us (Figure 1 A).²⁹ Firstly, Ag-nps were obtained through reduction of AgNO₃ by glucose and NaOH in presence of polyvinylpyrrolidone (PVP) and later they were coated with the organic ligand 1 in order to obtain lipophilic nanoparticles (Ag-nps-1).³⁰ Native Ag-nps present a hydrodynamic diameter of 35.4 ± 5.3 nm, which increased to 40 ± 7.6 nm after the coating with the organic ligand. A complete characterization of Ag-nps-1 was previously reported.³¹ Ag-nps-1 were then entrapped into PLGA-PEG polymeric nanoparticles (PNP). The nanoprecipitation method was exploited for this purpose and led to the entrapped of Ag-nps-1 in the inner core of the water-soluble PNP. Ag-PNP were then chemically conjugated onto the outer shell with the specific targeting agent CTX, thus giving Ag-PNP-CTX. For this reaction the presence of carboxylic acid groups, derived from PEG chains, onto the surface of Ag-PNP was exploited for the formation of an amide bond with free-amino ending group of CTX.

Ag-PNP-CTX were characterized by Transmission Electron Microscopy (TEM, Figure 1 B). Dynamic light scattering (DLS) revealed particles with diameter equal to 114 ± 2 nm, a low polydispersity index (PDI) (0.21 ± 0.01) and a negative ζ-potential (-31.6 mV). Silver concentration was determined to be 3.81 mM by means of atomic absorption spectroscopy (AAS). The overall Ag-PNP-CTX concentration was established by gravimetric analysis and found to be 8.87 mg/mL.

Optical imaging and quantification, and in vivo biodistribution of encapsulated Ag-nps

A key issue to achieve therapeutic efficacy resides in the amount of nanovectors that localize at the tumor site, and are internalized into the tumor cells. This is particularly relevant for GBM which is a high infiltrating and metastatic tumor, hardly accessible due to the presence of the BBB. Therefore, the use of accurate and quantitative methods to track systemically injected nanovectors is of crucial importance to evaluate and foresee their potential therapeutic efficacy.

1
2
3
4 Taking advantage of the high contrast provided by Ag-nps, we explored Ag-nps as contrast agents
5 for optical microscopy detection and imaging of PNPs without the need to add fluorescent dyes,
6 thus avoiding chemical modification of the outer shell of the nanocarriers, which can modify PNP
7 behavior inside the body. First, the intracellular quantification of Ag-nps contained into PNPs
8 nanovectors was performed in human GBM-derived cell lines by confocal microscopy. Pictures of
9 the nanoparticles alone by reflectance mode imaging are shown in Figure 2 C-E. As hypothesized,
10 intracellular localization and quantification of Ag-nps turned out to be a feasible procedure (Figure
11 2 A, B, F-I). Results showed that conjugation with CTX significantly increases the intracellular
12 accumulation of Ag-nps in GBM cells as shown by comparison between Ag-PNP-CTX and Ag-PNP
13 nanocarriers (Figure 2 L). A competition assay, performed by pretreating GBM cells with increasing
14 concentrations of native CTX peptide, indicated a significantly lower, dose-dependent, uptake of
15 targeted Ag-PNP-CTX. The CTX peptide did not modify the uptake of non-targeted Ag-PNP
16 nanovector (Figure 2 M). Furthermore, in agreement with the described capacity of PLGA
17 nanovectors to undergo endosomal escape,³²⁻³⁴ the intracellular trafficking analysis showed no
18 co-localization of Ag-nps with endosomal and lysosomal fluorescent markers (Figure 1N) implying
19 that CTX-nanovector cargo would be entirely released into the cytoplasm and not sequestered
20 and/or degraded inside cellular organelles.
21
22
23
24
25
26
27
28
29
30

31
32 Ag-nps proved to be a suitable tool also upon biodistribution evaluation in peripheral organs. After
33 Ag-PNP-CTX intra peritoneal (i.p.) injection, animals were sacrificed at selected time points and
34 different tissues including liver, spleen, lung, heart and kidney were collected and examined for the
35 presence of Ag-nps by Inductively Coupled Plasma Mass Spectrometry (ICP-MS). As expected the
36 majority of Ag-PNP-CTX accumulated in the RES organs (liver, ~80%, and spleen, ~ 4% of the
37 injected Ag-nps). Only 2 % of injected Ag-nps accumulated in the lung and a noteworthy
38 nanoparticles washout was observed at 48 hours from nanovector administration. Barely no Ag-
39 PNP-CTX nanovectors accumulate in the heart and in the kidney (Supplementary Figure 1).
40
41
42
43
44

45 Thus Ag-nps appear to represent a valuable imaging tool for single-cell analysis, endowed with
46 superior quality and precision with respect to standard fluorescence due to the high stability and
47 the absence of photobleaching of Ag-nps. Indeed, the light reflection property of Ag-nps allows the
48 acquisition of Ag-nps signal in reflection mode, without affecting the freedom of choice of
49 fluorescent labeled when the localization of diverse proteins have to be studied in multicolor
50 experiments. Moreover a fluorescent moiety, linked or adsorbed onto the external surface, may
51 undergo leakage phenomena from the nanosystem once in the body environment, which can lead
52 in turn to unspecific fluorescence detection in not really targeted cells; on the contrary, Ag-nps
53 being strongly entrapped inside the polymeric matrix, are stable and not subjected to this
54
55
56
57
58
59
60

1
2
3 inconvenience. As a general rule, the use of Ag-nps-loaded nanovectors may represent a valuable
4 tool to measure the delivery potential and/or to define the proper administration route and
5 scheduling of drug-loaded nanovectors of interest.
6
7
8
9

10 **Radiations Boost CTX-targeted nanovector accumulation in GBM tumors**

11
12
13 To verify the possibility to exploit Ag-PNP-CTX in vivo, the amount of Ag-nps accumulating into the
14 tumor and the healthy brain parenchyma was evaluated in an orthotopic mouse model of human
15 GBM (Figure 3) receiving Ag-PNP-CTX by systemic administration (i.p.). U87MG cells were
16 inoculated into the right hemisphere via stereotactic techniques. The GBM-tumor area was defined
17 by immunohistochemical staining for human nestin, a staminal marker overexpressed in glioma
18 stem cells³⁵ (Figure 3 A). Importantly, the U87MG orthotopic tumor was able to originate infiltrative
19 tumor niches. Indeed, in addition to the main tumor mass, groups of nestin-positive infiltrating cells
20 were detectable within the healthy parenchyma at the periphery of the tumor (Figure 3 B and C).
21 Outstandingly, we were able to detect single tumor cells spread into healthy tissue (Figure 3 C),
22 thus increasing the translational relevance of our studies.
23
24
25
26
27
28
29

30 Ag-nps detection and quantification inside the tumor was performed by confocal microscopy
31 imaging on brain cryosections. Ag-nps quantification showed a significant higher level of
32 accumulation in the GBM-tumor compared to healthy brain parenchyma where nearly no Ag-nps
33 were detected (Figure 4, 0 Gy). However, despite CTX favors Ag-PNP-CTX nanovector
34 internalization into GBM cells, CTX is not an agent for BBB crossing, thus nanovector brain
35 biodistribution could be limited by the presence of this barrier. Considering that RT is a mainstay of
36 GBM therapy regimens^{2,3} and that radiations affect BBB permeability,^{22,23} the consequence of a
37 radiation treatment on Ag-PNP-CTX nanovector tumor accumulation was assessed. To adhere
38 with current clinical protocol of RT used to treat GBM patients (60 Gy delivered in fractions of 2 Gy
39 ²) we chose 2 Gy as a standard experimental RT dosage (Figure 4 A and Supplementary Figure
40 2). Interestingly, a further, substantially increased amount (4 to 6 folds) of Ag-nps was detectable in
41 GBM-tumors subjected to radiations compared to the absence of the radiations. A higher quantity
42 of Ag-nps was observed 48 hours from Ag-PNP-CTX injection compared to 24 hours suggestive of
43 an enhanced permeability and retention (EPR) effect, possibly preventing nanovectors wash out
44 and facilitating their accumulation (Figure 4). Noteworthy, radiation treatment did not determine an
45 increased accumulation of Ag-nps in healthy brain parenchyma (Figure 4B).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Radiations alter the blood-brain barrier allowing the targeting of GBM cells infiltrating healthy brain parenchyma

We next aimed to assess whether the lack of Ag-nps in the irradiated brain healthy parenchyma was due to the presence of an intact BBB and to investigate the contribution of BBB integrity in the enhanced accumulation of Ag-PNP-CTX nanovectors in GBM tumors observed upon radiation treatment. To this aim we performed immunostaining for Claudin-5, one of the highly expressed brain endothelial tight junction proteins contributing in maintenance of BBB integrity.³⁶ Claudin-5 levels and pattern of expression were determined by immunohistochemistry on cryosections from normal and U87MG-implanted brain hemispheres treated and non-treated with radiations. Results indicated that a single dose of 2 Gy radiation caused alterations in claudin-5 distribution pattern, which was particularly evident in the healthy brain where the formation of gaps in brain vessel wall, caused by the opening of the tight junctions was clearly detectable (Figure 5 A and B, see also 3D reconstructions). Differently from healthy brain, the tumorigenic GBM microenvironment promotes neo-angiogenesis and disruption of BBB integrity (Figure 5 C and data not shown).³⁷ The permeabilization of the BBB in the tumor area allows in fact penetration and retention of Ag-nps within the non-irradiated GBM-tumor, although at a lesser extent compared to the irradiated one (Figure 4).

While the integrity of the BBB would guarantee reduced nanovectors off-target brain toxicity, it makes infiltrating and metastatic GBM cells inaccessible to nanovectors. Indeed, in the proximity of the peripheral zone of the tumor we observed a normal claudin-5 expression, suggestive of an intact BBB (Figure 5 D). We therefore focused on the tumor periphery in order to investigate the penetration of Ag-nps in dispersed cancer cells located far from the tumor mass, identified by their immunoreactivity to antibodies directed against human nestin. As expected, no Ag-nps accumulation in nestin-positive cells (Figure 5 E, left panel and Supplementary Figure 3). After 2 Gy radiation treatment, however, in line with BBB permeabilization, Ag-nps could be detected in the isolated GBM cells branching into the healthy parenchyma (Fig 5 E, right panel and Supplementary Figure 3).

Despite low doses of radiations were enough to determine *in vivo* tight junction opening (Figure 5B), the presence of a compromised BBB didn't allow significant accumulation of Ag-nps in irradiated healthy brain parenchyma (Figure 4). This is consistent with the presence, in the proximity and surrounding the brain vessels, of pericytes and astrocytes that contribute hindering the passive passage of nanovectors.³⁸ Furthermore, the interstitial fluid pressure³⁹ and the width of the extracellular space impact on nanovector access and penetration within the healthy parenchyma. It was previously reported that the extracellular spaces in normal brain are between

1
2
3 38 and 64 nm.¹⁴ Therefore, 100nm-sized CTX-nanovectors, likewise the ones used in this study,
4 would hardly diffuse and accumulate in healthy brain tissue. It is conceivable that ECM
5 components would physically restrict the motion of nanovector in convection through the
6 extracellular space.
7
8

9 10 11 **Radiation Augments Specific Cellular Uptake and Intracellular Localization by inducing** 12 **MMP-2 and CIC-3 overexpression** 13 14

15
16 The presence of ligands on nanocarrier surface increases the active intracellular uptake
17 determining higher payload delivery inside the cells.¹⁶ Notably, while irradiated cells, both U87MG
18 and T98G lines, incubated in the presence of Ag-PNP-CTX displayed a significant increase in
19 intracellular Ag-nps, no variations were observed in either cell lines when incubated in the
20 presence of non-targeted Ag-PNP nanovectors (Figure 6A and 6B). Given CTX is the ligand for
21 MMP-2 and CIC-3, which are highly expressed in human gliomas,^{17,21} we quantified the
22 expression of these molecules in irradiated tumor, compared with the not irradiated ones. RT is
23 known to induce MMP-2 expression²⁴⁻²⁶ but no data are available yet on CIC-3 expression after
24 radiation treatment.
25
26
27
28
29
30

31
32 Our data indicated that up-regulation of MMP-2 and CIC-3 levels of expression is induced by
33 radiations, as assessed by western-blot analysis (Figure 6B). The effect was particularly evident for
34 CIC-3. Similarly, simultaneous microscopy detection of MMP-2 or CIC-3 and Ag-nps in non-
35 irradiated and irradiated U87MG cells confirmed that radiations increase MMP-2 and CIC-3 levels
36 of expression, which correlate with a higher amount of internalized Ag-nps (Figure 6C and D,
37 Supplementary Figure 4). Notably, the increased expression of MMP-2 and CIC-3 was observed
38 also *in vivo*, in irradiated U87MG xenografts (Figure 7). Despite the results obtained *in vitro*
39 indicated a noteworthy increase of CIC-3 expression compared to MMP-2 after radiation treatment,
40 the *in vivo* CIC-3 and MMP-2 levels were comparable, thus suggesting an equivalent contribution
41 for nanovectors cellular uptake and internalization.
42
43
44
45
46
47

48
49 The lack of Ag-nps accumulation in infiltrating GBM cells in the absence of radiation treatment
50 (Figure 5E) is compatible with the possibility that, besides favoring permeabilization of the BBB,
51 ionizing radiations, by enhancing the expression of cellular proteases, including MMP-2, play a
52 crucial role in degradation and remodeling of brain extra cellular matrix (ECM) thereby facilitating
53 extravasation, convective movements and diffusion of CTX-nanovectors. This is relevant in the
54 clinical setting, where RT is applied not only to the tumor area but extended to a 2 cm surrounding
55 margins where invading GBM stem-like cells escape to surgery and survive to RT.^{40,41} Thus the
56
57
58
59
60

1
2
3 concomitant use of CTX-nanovectors during RT should facilitate and enhance drug-delivery to the
4 invading and metastatic GBM stem-like cells that are the responsible for the majority of the
5 recurrences.
6
7
8
9

10 **CTX-targeted nanovectors function as inhibitor of MMP-2 enzymatic activity**

11
12
13 MMP-2 belongs to a family of endopeptidases, which can cleave almost every component of the
14 ECM proteins. ECM degradation is a key event in the progression, invasion, and metastasis of
15 malignant lesions.⁴² Notably, increase of metastatic tumor activity has been described as a
16 consequence of RT, which enhances MMP-2 expression.²⁵ We therefore asked whether the
17 binding of CTX-nanovectors to MMP-2 could interfere with its catalytic activity. MMP-2 activity was
18 evaluated by zymography in irradiated U87MG cells incubated with non-targeted Ag-PNP and
19 targeted Ag-PNP-CTX nanovectors. Remarkably, MMP-2 catalytic activity was inhibited of about
20 50% exclusively in cells incubated with targeted Ag-PNP-CTX nanovectors while no changes were
21 observed in irradiated cells incubated with non-targeted Ag-PNP nanovectors (Figure 8). Thus,
22 CTX-PNP nanovectors could attain synergistic dual therapeutic effects, tumor-specific intracellular
23 drug-delivery and inhibition of MMP-2 activity. Interestingly, it has been demonstrated that
24 inhibition of MMPs, including MMP-2, provides a potential strategy for improving RT efficacy.^{27,28}
25
26
27
28
29
30
31
32
33
34

35 **Radiations enhance Ag-PNP-CTX cytotoxic activity against U87MG GBM cells**

36
37
38 Recently the possibility to use Ag-nps as anticancer agents for GBM, either alone^{29,43} or in
39 combination with other drugs,³¹ after entrapment in biocompatible PNPs has been explored.
40 Hence, the effect of a combined administration of 2 Gy radiation treatment and Ag-PNP-CTX on
41 U87MG cell growth and viability was investigated.
42
43
44

45
46 Irradiated and non-irradiated U87MG cells were incubated with different concentrations of Ag-PNP-
47 CTX nanovectors for 72 hours and, at the end of the treatment period, cell viability was assessed
48 by MTT test. Parallel U87MG cells were treated with either nanovectors without Ag-nps (PNP-
49 CTX), for polymer and CTX toxicity assessment, or non-targeted nanovectors (Ag-PNP), as control
50 for specific uptake. As expected, no cellular toxicity was observed by incubating U87MG cells with
51 empty PNP-CTX nanovectors, while the presence of CTX on the surface of Ag-PNP nanovectors
52 triggered a dose-dependent cell growth inhibition induced by the encapsulated Ag-nps. More
53 interestingly, a synergic reduction of U87MG cell viability was observed when combined
54 treatments, radiations (2 Gy) and Ag-PNP-CTX nanovectors, were applied (Figure 9A). Pro-
55 apoptotic synergic effect was confirmed *in vivo* in U87MG-transplanted mice. The detection and
56
57
58
59
60

1
2
3 quantification of apoptotic bodies by *in situ* terminal deoxynucleotidyltransferase-UTP nick end
4 labeling (TUNEL) assay indicated a significant apoptosis increase in irradiated U87MG xenografts
5 exposed to radiations and Ag-PNP-CTX nanovectors vs. U87MG xenografts treated with CTX
6 nanovectors or exposed to radiations as single agents (Figure 9B).
7
8

9
10 Corroboration of the TUNEL results was provided by bioluminescence imaging (BLI) analysis
11 performed in luciferase-transfected U87MG (U87MGluc2) xenografts to monitor tumor growth over
12 the treatment period (Figure 9C and D). Eleven days post U87MGluc2 cells implantation (T=0),
13 when the tumor mass was detectable by BLI, mice were divided into 4 groups: untreated mice;
14 mice treated with radiations (2Gy, single dose); mice treated with serial injections of Ag-PNP-CTX
15 single agent (2 doses of 1mg dry matter / mouse, followed by 2 doses of 0,5 mg dry matter /
16 mouse; interval time: 48-72 hours); mice treated with radiations and Ag-PNP-CTX concomitantly.
17
18

19
20 As illustrated in Figure 9C, showing representative images of single animals from each
21 experimental group, and 9D, showing the quantitative analysis of the tumor areas, combined
22 radiations and CTX-nanovectors treatment inhibited tumor growth and progression. No inhibitory
23 effect was observed in mice treated with either radiations or Ag-PNP-CTX alone.
24
25
26
27
28
29

30 The observed increased level of apoptosis together with the *in vivo* inhibition of tumor growth
31 would suggest a positive anti-tumor effect of Ag-nps. Though, more meaningfully, the positive
32 results obtained with Ag-PNP-CTX provide the proof-of-concept of the therapeutic validity of a
33 combination strategy based on radiation therapy and CTX-targeted polymeric nanovectors, which
34 is expected to strengthen the anti-tumor efficacy of any cytotoxic molecules entrapped into PNP-
35 CTX nanovectors.
36
37
38
39
40
41

42 CONCLUSIONS

43
44
45 In this report, we described a nanotechnological strategy for the treatment of GBM that addresses
46 two relevant obstacles to effective therapy: 1) the infiltrative nature of GBM; 2) the increase in
47 MMP-2 expression following radiotherapy. Overall our results showed that (i) although large and
48 advanced GBM tumors disrupt the BBB sufficiently to allow extravasation of engineered
49 nanocarriers, low dose RT (2 Gy, single dose), by inducing overexpression of proteases such as
50 MMP-2, which also represent CTX molecular targets together with the chloride channel ClC-3,
51 render tumor ECM more porous and permeable to nanovectors, determining their significant
52 enhanced accumulation; ii) differently from the main tumor mass, peripheral GBM tumor niches,
53 where more tumorigenic and invading cells reside, are accessible to engineered nanocarriers
54
55
56
57
58
59
60

1
2
3 exclusively upon radiation treatment; *iii*) low dose RT is not enough to determine complete BBB
4 disruption preventing nanocarrier accumulation in irradiated healthy brain parenchyma; *iv*) CTX
5 surface functionalization, besides causing an increased cellular uptake of CTX-targeted
6 nanovectors, inhibits MMP-2 enzymatic activity.
7
8
9

10 Furthermore, silver nanoparticles proved to represent a valuable imaging tool for single-cell
11 analysis both *in vitro* and *in vivo*, with superior quality and precision respect to standard
12 fluorescence. Indeed, Ag-nps were of fundamental importance to demonstrate nanovector
13 localization *in vivo* in peripheral infiltrating GBM tumor niches.
14
15
16
17

18 Having demonstrated that an increased nanovectors accumulation and retention can be achieved
19 at significant lower dose of RT (2 Gy), we provide a strong and relevant rationale for the
20 development of therapeutic strategies able to target tumorigenic and invading GBM cells based on
21 the concomitant administration of RT and CTX-targeted nanovectors. A synergistic therapeutic
22 effect could start from the first fractionated RT dose. Considering that radiations induce the
23 expression of molecules such as MMPs that favor tumor invasion and metastases, the concomitant
24 administration of RT and CTX-targeted nanovectors could represent a significant step forward to
25 GBM management. Indeed, the synergistic effect of CTX-targeted would result from a significant
26 increased accumulation of nanovector therapeutic cargo inside GBM cells and from the inhibition of
27 approximately 50 % of MMP-2 catalytic activity. Thus we conclude that CTX-targeted nanovectors
28 as adjuvants to RT may increase the efficacy of anti-GBM therapies.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Synthesis and characterization of Ag-PNP-CTX nanovectors. 100 mg of PLGA-b-PEG-COOH (7 kDa-3 kDa) and AgNPs-1, both prepared as already reported³⁰, were admixed into 10 mL of dimethylformamide (DMF). This organic phase was mixed with 100 mL of ultrapure water under vigorous stirring, maintaining water/organic ratio 10/1 with a constant removal of the solution. The mixture was kept for 30 min. under vigorous stirring then washed and concentrated by using centrifugal filter devices (Amicon Ultra, Ultracell membrane with 100.000 NMWL, Millipore, USA) to a final volume of 5 mL and finally filtered by using a syringe filters phenex-PES of polyether sulfone (26 mm, 0.20 μ m, Phenomenex, Italy). To a suspension of the so obtained Ag-PNP (5 mL) in PBS (10 mL, 0.01 M), under magnetic stirring, a solution of N-hydroxysulfosuccinimide 2.3 mM (8.7 mL) and a solution of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide 0.28 M (3.6 mL) was added. The reaction was carried out at room temperature for 15 minutes then 50 μ g of Chlorotoxin, dissolved into 1 mL of water, were added and left to react for 24 hours. After that Ag-PNP-CTX were purified and concentrated to a final volume of 5 mL as previously described. Dynamic light scattering (DLS) analysis and ζ -potential values were obtained with a Zetasizer Nano-S (Malvern) instrument, working with a 532 nm laser beam at 25 °C, using standard cuvettes or DTS1060C Clear Disposable zeta cells, and the results expressed as average of three measurements. Atomic absorption analysis was conducted in a SpectraAA 100 Varian instrument, with opportune AgNO₃ standards for calibration line. Final concentration of the suspensions was determined by gravimetric analysis by drying 100 μ L of solution at 130 °C for 24 hours then accurately weighting the residual dry matter amount.

Cell Culture. Human-derived GBM cell line were grown in Minimum Essential Medium Eagle, Alpha Modification (α MEM) (U87MG) or RPMI (A172, T98G) (Lonza, Basel, Switzerland) supplemented with 10% fetal calf serum (FCS, U.S. Origin, Gibco/Life Technologies), 100 U/ml penicillin and 0,1 mg/ml streptomycin. Cells were maintained at 37°C in a humidified incubator with 5% CO₂.

Competitive inhibition assay. U87MG and A172 cell lines were pre-treated for 2 hours with decreasing concentrations of the native CTX peptide (MCMPCFTTDHQMARCDDCCGGKGRGKCYGPQCLCR-NH₂) (H6086, Bachem, Bubendorf, Switzerland) followed by incubation with targeted (Ag-PNP-CTX) or non-targeted (Ag-PNP) nanovectors at Ag concentration of 100 μ M for 16-18 hours (overnight). The intracellular accumulation of Ag nanoparticles (Ag-nps) was determined by confocal laser scanning microscope and 3D imaging analysis performed by IMARIS software (see below).

1
2
3 **Mouse orthotopic xenografts.** Mouse experiments were carried out according to the
4 protocol approved by Italian Minister of Health (protocol number 171/2013-B). CD-1 male nude
5 mice, 6-8 weeks (Charles Rivers Laboratories, Calco, Italy) were housed in specific pathogen free
6 animal house. Sterilized Water and rodent chow were given ad libitum. Mice were injected
7 intracranially into the right striatum with U87MG cells at 5×10^4 cells/mouse or luciferase-transfected
8 U87MGluc2 (gently provided by Laura Cerchia, CNR Naples, Italy) 10^5 cells/mouse. For intra-
9 tumor nanovector accumulation analysis and quantification, U87MG-tumors were allowed to grow
10 for 15 days, and then mice were subject to whole-brain single dose radiation (2 Gy). Ag-PNP-CTX
11 were administered intraperitoneally at a dose of 1mg of dry matter 18 hours after radiations.
12 Animals were euthanized 24 or 48 hours after nanovector injection. For immunofluorescence
13 experiments intracardiac perfusion was carried out with 4% PFA. The harvested brains and organs
14 were frozen and conserve at -80°C .
15
16
17
18
19
20

21 To study the antitumor effect of Ag-PNP-CTX, U87MGluc2-tumors were allowed to growth
22 for 11 days, and then mice were randomized and divided in 4 groups: not treated animals
23 (controls), animal treated only with Ag-PNP-CTX nanovectors, animal treated only with radiations,
24 and animal treated with radiations and Ag-PNP-CTX nanovectors. Mice were subject to whole-
25 brain single dose radiation (2 Gy, T0). Ag-PNP-CTX administration started 18 hours after radiation
26 treatment. Mice received 4 consecutive doses of Ag-PNP-CTX nanovectors (first 2 doses: 1 mg of
27 dry matter; third and fourth doses: 0,5 mg of dry matter) at intervals of 48-72 hours.
28
29
30
31
32

33 ***In vivo* imaging of intracranial tumors.** An IVIS II Imaging System (Caliper LifeSciences-
34 Perkin Elmer-Waltham, Massachusetts, USA) was used to take photograph and luminescent
35 images setting an open filter and binning 8. Exposure time was fixed at 1 minute and 5-10 images
36 were acquired until maximum photon flux achievement. Ten minutes before acquisition, mice were
37 administered intraperitoneal injection of D-luciferin (Perkin Elmer-Waltham), 150 mg/Kg, as a
38 substrate for the luciferase enzyme. Images were captured and quantified with Living Image 4.3.1
39 software (Perkin Elmer-Waltham), based on equivalent regions of interest over the head. Analysis
40 was carried out using the Living imaging software (Perkin Elmer - Waltham, Massachusetts, USA)
41 setting binning 4 and "automatic ROI" (thershold 50%) for all images.
42
43
44
45
46
47

48 **Radiation treatment.** Cells and mice were irradiated with a single dose of 2 Gy using an x-
49 ray biological irradiator operating at 12 mA/190 kv (RADGIL, Gilardoni, Lecco, Italy). Cells were
50 irradiated at a dose rate of 0,65 Gy/min. Mice were irradiated at a dose rate of 0,4 Gy/min in an in-
51 house designed lead irradiation jig that allowed only the head to be exposed and shielded the
52 body.
53
54
55
56
57
58
59
60

1
2
3 ***In vitro* nanovector intracellular uptake.** U87MG , A172, T98G were seeded on glass
4 cover slips (16 mm diameter) at a density of 15.000 cells/coverslip in the presence of 5% FCS.
5 After 24h of culture, the cells were incubated with targeted (Ag-PNP-CTX) or non-targeted (Ag-
6 PNP) nanovectors at Ag concentration of 100 μ M for 16-18 hours (overnight). One hour before cell
7 fixation, Syto Blue 45® fluorescent dye (Life Technologies Corporation, Carlsbad, Ca, USA) was
8 added to the medium (1:1000). After three washes in Dulbecco's Phosphate Buffered Saline (PBS)
9 (Sigma-Aldrich, St. Louis, MO, USA), cells were fixed in 4% paraformaldehyde for 15 min at room
10 temperature. Glasses were mounted with a PBS/Glicerol (Sigma-Aldrich, St. Louis, MO, USA),
11 mixture (1:1) added with DAPI (Sigma-Aldrich, St. Louis, MO, USA), fluorescent stain 1:40.000.
12
13
14
15
16
17

18 **Immunostaining.** Before immunostaining fixed 10-micron-thick frozen brain section were
19 heat antigen retrieved in Sodium Citrate (pH=6). Unspecific binding was blocked with 10% horse
20 serum (Sigma) diluted in PBS/Triton 0,1% (1 hour). Primary antibodies against human-nestin clone
21 2C1.3A11 (1:250 1h RT) (Abcam), human-MMP-2 (1:100 1h RT) (Sigma/Prestige Antibodies),
22 human-CICN3 (1:100 1h RT) (Cell Signaling Technology, Beverly, MA, USA), human-LAMP1 (BD
23 Bioscience) (1:100 1h RT) and mouse-claudin-5 (1:600 24h 4°C) (Millipore) were employed at same
24 experimental conditions for both cell lines and brain cryosections. After primary antibody, samples
25 were incubated with the appropriate secondary antibodies conjugated with one of the indicated
26 fluorophores, Alexa fluor: 488,555, 561 (Invitrogen) (dilution 1:200; 1 hour RT). Early endosome
27 were marked using CellLight Early Endosome-RFP, BacMan 2.0 (Invitrogen) according to the
28 procedure instruction.
29
30
31
32
33
34
35
36

37 **TUNEL assay.** Detection of apoptosis was carried out by *in situ* TUNEL-assay (Click-iT®
38 TUNEL Alexa Fluor® 633 Invitrogen-Life Technologies, Grand Island, NY, USA). TUNEL was
39 performed according to the protocol provided by the manufacturer. Briefly, after slide incubation
40 (10 minutes at 37°C) in TdT reaction buffer, samples were incubated with TdT reaction mixture for
41 60 minutes at 37°C. After washes with PBS1x sections were washed with 3% BSA and 0.1%
42 Triton® X-100 in PBS for 5 minutes. The Click-iT® Plus TUNEL reaction cocktail was added to
43 each slide for 30 minutes at 37°C.
44
45
46
47
48

49 **Confocal microscopy and images analysis.** Confocal microscopy was performed using a
50 Leica TCS SP5 confocal microscope equipped with a resonant scanner (scan speed 8000 Hz) and
51 a HCX PL APO 63x/1.4 OIL objective, a HC PL FLUOTAR 10.0x0.30 DRY and a HC PL FLUOTAR
52 20.0x0.50 DRY. Ag-nanoparticles (nps) were acquired in reflection mode upon 458 nm excitation
53 for *in vitro* cell experiments and upon 488nm excitation for *in vivo* experiments on brain
54 cryosections. Syto blue was excited by the 458 nm laser line, nestin and phalloidin by the 488 nm
55
56
57
58
59
60

1
2
3 line of an Argon laser, claudin, MMP-2 and CIC-3 by the 561 nm laser line of a diode laser. Voxel
4 size was established using Nyquist criteria.

5
6 Intracellular Ag-nps quantification was carried out by 3D imaging analysis by means of IMARIS
7 software (Bitplane, IMARIS) using the “Cells” function and the following set-up: cell
8 detection/smooth filter =0,160; Background Subtraction, sphere diameter=20; Detect Vesicles,
9 Estimated Diameter= 240 nm; Vesicle Quality=18).

10
11 Ag uptake analysis *in vivo* was performed using IMARIS software using the “Spots” function
12 into a fixed area (6500 μm^2) for all fields, murine or human xenograft sites.

13
14 MMP-2 and CLC-3 levels expression analysis was performed using Image J software. In
15 brief, the central, the upper and the lower section of a 6 μm z-stack was analyzed for each image.
16 MMP-2 and CIC-3 Threshold mask was kept constant and using the tool “Analyze Particles” area
17 of MMP-2 and CIC-3 signal was calculated. MMP-2 and CIC-3 area was normalized on number of
18 nuclei present in the slice analyzed.
19
20
21
22
23

24
25 **Western Blotting** . Approximately 2×10^5 U87MG cells were lysed in ice-cold lysis buffer
26 (100 mM Hepes, 2mM EDTA, 1% SDS) supplemented with protease inhibitors (ROCHE) and then
27 stored at -80°C until used. Protein concentrations were determined using a BCA protein assay
28 (Pierce). Equal amounts of protein (20 μg) were loaded on 10% polyacrylamide gels and resolved
29 on SDS-PAGE. Membranes were blocked (1 hour RT) with 5% milk in PBS/tween-20 0,1%
30 (Sigma) and then probed (1 hour RT) with anti-human MMP-2 polyclonal rabbit antibody (1:1000)
31 (Sigma), anti-human CIC-3 monoclonal rabbit antibody (1:1000) (Cell Signaling Technology,
32 Beverly, MA,USA) and anti-human β -actin monoclonal mouse antibody (1:5000) (Sigma). Anti-
33 rabbit and anti-mouse HRP-conjugated secondary antibodies (Pierce, Rockford, IL, USA) were
34 employed at a dilution of 1:5000 for 1 RT. Bands were visualized by Bio-Rad Clarity western ECL
35 substrate (Biorad, Hercules, USA), and detected by ChemiDoc MP system (Biorad, Hercules,
36 USA). All densitometric analyses were carried out using Image Lab software version 5.1 (Biorad
37 Hercules, USA).
38
39
40
41
42
43
44
45

46
47 **Gelatin Zymography**. Supernatant from in vitro grown U87MG cells were eluted in
48 Zymogram Sample Buffer (Biorad, Hercules, USA) according to producer instructions. Equal
49 amount of each samples were loaded in Ready Gel® Zymogram Precast Gels (Biorad, Hercules,
50 USA) and electrophoresis was carried out at fixed voltage of 90 V. The gels were washed twice
51 in distilled water for 10' and twice in Zymogram Renaturation Buffer (Biorad, Hercules, USA) for
52 30' and then incubated for 6 hours at 37°C in Zymogram Development Buffer (Biorad, Hercules,
53 USA). After incubation, gels were stained with Bio-Safe Coomassie Stain (Biorad, Hercules, USA)
54 for 2 hours. Gelatinolytic activity was detected as transparent bands on a blue background. Images
55
56
57
58
59
60

1
2
3 were acquired with chemidoc system (Biorad, Hercules, USA). Densitometry analysis was
4 performed with ImageJ Software.
5
6

7
8 **MTT Assay.** U87MG cells (5,000 cells/well) were plated in 96-well plates with 100 μ l α MEM
9 medium. After 24 h cells were treated with increasing concentrations of Ag-nps encapsulated in the
10 PNP nanovectors, whereas no treatment was done as positive proliferation control. At 72 h, 25 μ l
11 of 5 mg/ml 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Sigma-Aldrich, St
12 Louis, MO) was added directly to the cells followed by an additional 4 hours of incubation, then 100
13 μ L of DMSO was added. The optical density of individual wells was measured at a wavelength of
14 550 nm with the Sunrise apparatus (Tecan Group, Ltd, Mannedorf, Switzerland). All assays were
15 performed in triplicate.
16
17
18
19

20
21 **Statistics.** Data are presented as mean \pm SE. If not indicated mean values were compared
22 by the Student's *t* test (two-tailed, non parametric). Statistics and fitting of experimental data were
23 performed with Graph Pad Prism6.
24
25
26
27
28

29 **ACKNOWLEDGEMENT**

30 We thank Dr. Lorenzo Bello (Humanitas and University of Milano), Dr. Elisabetta Menna (IN-CNR)
31 and Dr. Davide Pozzi (Fondazione Humanitas per la Ricerca) for suggestions. We acknowledge
32 Prof. Giulio Maira and Dr. Maurizio Fornari (Humanitas Research Hospital) for critically reading the
33 manuscript and for continuous constructive discussions. This work was supported by Fondazione
34 "Giancarla Vollaro", Via Filodrammatici 10, Milano, by CNR InterOmics Flagship Project to M.M.
35 (ID B92I15000140001) and by CEN Foundation, Piazza Leonardo da Vinci 32, Milano (Start-up
36 Packages and PhD Program" project, co-funded by Regione Lombardia through the program
37 "Fondo per lo Sviluppo e la Coesione 2007-2013", ex FAS").
38
39
40
41
42
43
44
45

46 **SUPPORTING INFORMATION AVAILABLE:**

47 Ag-NP-CTX biodistribution in peripheral organs, bright field images of cryosections of tumor tissues
48 from U87MG orthotopic xenografts inoculated with Ag-PNPCTX nanovectors, bright field magnified
49 images of cells reported in Figure 6 D.
50
51
52
53
54
55
56
57
58
59
60

REFERENCES AND NOTES

- 1 Bastien, J. I.; McNeill, K. A.; Fine, H. A. Molecular characterizations of glioblastoma, targeted therapy, and clinical results to date. *Cancer*. **2015**, *121*, 502-516.
- 2 Villa, S.; Balana, C.; Comas, S. Radiation and concomitant chemotherapy for patients with glioblastoma multiforme. *Chin J Cancer*. **2014**, *33*, 25-31.
- 3 Stupp, R.; Mayer, M.; Kann, R.; Weder, W.; Zouhair, A.; Betticher, D. C.; Roth, A. D.; Stahel, R. A.; Majno, S. B.; Peters, S.; *et al.* Neoadjuvant chemotherapy and radiotherapy followed by surgery in selected patients with stage IIIB non-small-cell lung cancer: a multicentre phase II trial. *Lancet Oncol*. **2009**, *10*, 785-793.
- 4 Kreuter, J. Drug delivery to the central nervous system by polymeric nanoparticles: what do we know? *Adv Drug Deliv Rev*. **2014**, *71*, 2-14.
- 5 Agarwal, S.; Manchanda, P.; Vogelbaum, M. A.; Ohlfest, J. R.; Elmquist, W. F. Function of the blood-brain barrier and restriction of drug delivery to invasive glioma cells: findings in an orthotopic rat xenograft model of glioma. *Drug Metab Dispos*. **2013**, *41*, 33-39.
- 6 Woodworth, G. F.; Dunn, G. P.; Nance, E. A.; Hanes, J.; Brem, H. Emerging insights into barriers to effective brain tumor therapeutics. *Front Oncol*. **2014**, *4*, 126, doi:10.3389/fonc.2014.00126 .
- 7 Wohlfart, S.; Gelperina, S.; Kreuter, J. Transport of drugs across the blood-brain barrier by nanoparticles. *J Control Release*. **2012**, *161*, 264-273.
- 8 Acharya, S.; Sahoo, S. K. PLGA nanoparticles containing various anticancer agents and tumour delivery by EPR effect. *Adv Drug Deliv Rev*. **2011**, *63*, 170-183.
- 9 Gao, Z.; Zhang, L.; Sun, Y. Nanotechnology applied to overcome tumor drug resistance. *J Control Release*. **2012**, *162*, 45-55.
- 10 Locatelli, E.; Gil, L.; Israel, L. L.; Passoni, L.; Naddaka, M.; Pucci, A.; Reese, T.; Gomez-Vallejo, V.; Milani, P.; Matteoli, M.; *et al.* Biocompatible nanocomposite for PET/MRI hybrid imaging. *Int J Nanomedicine*. **2012**, *7*, 6021-6033.
- 11 Danhier, F.; Ansorena, E.; Silva, J. M.; Coco, R.; Le Breton, A.; Preat, V. PLGA-based nanoparticles: an overview of biomedical applications. *J Control Release*. **2012**, *161*, 505-522.
- 12 Kunwar, S.; Chang, S.; Westphal, M.; Vogelbaum, M.; Sampson, J.; Barnett, G.; Shaffrey, M.; Ram, Z.; Piepmeyer, J.; Prados, M.; *et al.* Phase III randomized trial of CED of IL13-PE38QQR vs Gliadel wafers for recurrent glioblastoma. *Neuro Oncol*. **2010**, *12*, 871-881.
- 13 Sawyer, A. J.; Saucier-Sawyer, J. K.; Booth, C. J.; Liu, J.; Patel, T.; Piepmeyer, J. M.; Saltzman, W. M. Convection-enhanced delivery of camptothecin-loaded polymer nanoparticles for treatment of intracranial tumors. *Drug Deliv Transl Res*. **2011**, *1*, 34-42.
- 14 Thorne, R. G.; Nicholson, C. In vivo diffusion analysis with quantum dots and dextrans predicts the width of brain extracellular space. *Proc Natl Acad Sci U.S.A.* **2006**, *103*, 5567-5572.
- 15 Hobbs, S. K.; Monsky, W. L.; Yuan, F.; Roberts, W. G.; Griffith, L.; Torchilin, V. P.; Jain, R. K. Regulation of transport pathways in tumor vessels: role of tumor type and microenvironment. *Proc Natl Acad Sci U.S.A.* **1998**, *95*, 4607-4612.
- 16 Gao, H.; Yang, Z.; Zhang, S.; Cao, S.; Shen, S.; Pang, Z.; Jiang, X. Ligand modified nanoparticles increases cell uptake, alters endocytosis and elevates glioma distribution and internalization. *Sci Rep*. **2013**, *3*, 2534, doi:10.1038/srep02534.
- 17 Soroceanu, L.; Gillespie, Y.; Khazaeli, M. B.; Sontheimer, H. Use of chlorotoxin for targeting of primary brain tumors. *Cancer Res*. **1998**, *58*, 4871-4879.
- 18 DeBin, J. A.; Maggio, J. E.; Strichartz, G. R. Purification and characterization of chlorotoxin, a chloride channel ligand from the venom of the scorpion. *Am J Physiol*. **1993**, *264*, C361-369.
- 19 Deshane, J.; Garner, C. C.; Sontheimer, H. Chlorotoxin inhibits glioma cell invasion via matrix metalloproteinase-2. *J Biol Chem*. **2003**, *278*, 4135-4144.
- 20 Lui, V. C.; Lung, S. S.; Pu, J. K.; Hung, K. N.; Leung, G. K. Invasion of human glioma cells is regulated by multiple chloride channels including ClC-3. *Anticancer Res*. **2010**, *30*, 4515-4524.
- 21 McFerrin, M. B.; Sontheimer, H. A role for ion channels in glioma cell invasion. *Neuron Glia Biol*. **2006**, *2*, 39-49.


- 1
2
3 22 Yuan, H.; Gaber, M. W.; McColgan, T.; Naimark, M. D.; Kiani, M. F.; Merchant, T. E. Radiation-induced permeability and leukocyte adhesion in the rat blood-brain barrier: modulation with anti-ICAM-1 antibodies. *Brain Res.* **2003**, *969*, 59-69.
- 4
5
6 23 d'Avella, D.; Ciccirello, R.; Albiero, F.; Mesiti, M.; Gagliardi, M. E.; Russi, E.; d'Aquino, A.; Tomasello, F.; d'Aquino, S. Quantitative study of blood-brain barrier permeability changes after experimental whole-brain radiation. *Neurosurgery.* **1992**, *30*, 30-34.
- 7
8
9 24 Kwak, S. Y.; Kim, B. Y.; Ahn, H. J.; Yoo, J. O.; Kim, J.; Bae, I. H.; Han, Y. H. *et al.* Ionizing radiation-inducible miR-30e promotes glioma cell invasion through EGFR stabilization by directly targeting CBL-B. *FEBS J.* **2015** doi:10.1111/febs.13238.
- 10
11
12 25 Park, C. M.; Park, M. J.; Kwak, H. J.; Lee, H. C.; Kim, M. S.; Lee, S. H.; Park, I. C.; Rhee, C. H.; Hong, S. I. Ionizing radiation enhances matrix metalloproteinase-2 secretion and invasion of glioma cells through Src/epidermal growth factor receptor-mediated p38/Akt and phosphatidylinositol 3-kinase/Akt signaling pathways. *Cancer Res.* **2006**, *66*, 8511-8519.
- 13
14
15 26 Lee, W. H.; Warrington, J. P.; Sonntag, W. E.; Lee, Y. W. Irradiation alters MMP-2/TIMP-2 system and collagen type IV degradation in brain. *Int J Radiat Oncol Biol Phys.* **2012**, *82*, 1559-1566.
- 16
17
18 27 Badiga, A. V.; Chetty, C.; Kesanakurti, D.; Are, D.; Gujrati, M.; Klopfenstein, J. D.; Dinh, D. H.; Rao, J. S. MMP-2 siRNA inhibits radiation-enhanced invasiveness in glioma cells. *PLoS One.* **2011**, *6*, e20614, doi:10.1371/journal.pone.002061410-PONE-RA-19742.
- 19
20
21 28 Ulasov, I.; Thaci, B.; Sarvaiya, P.; Yi, R.; Guo, D.; Auffinger, B.; Pytel, P.; Zhang, L.; Kim, C. K.; Borovjagin, A.; *et al.* Inhibition of MMP14 potentiates the therapeutic effect of temozolomide and radiation in gliomas. *Cancer Med.* **2013**, *2*, 457-467.
- 22
23
24 29 Locatelli, E.; Broggi, F.; Ponti, J.; Marmorato, P.; Franchini, F.; Lena, S.; Franchini, M. C. Lipophilic silver nanoparticles and their polymeric entrapment into targeted-PEG-based micelles for the treatment of glioblastoma. *Adv Healthc Mater.* **2012**, *1*, 342-347.
- 25
26
27 30 Gentili, D., Ori, G. & Franchini, MC. Double phase transfer of gold nanorods for surface functionalization and entrapment into PEG-based nanocarriers. *Chem Commun (Camb)* **2009**, *39*, 5874-5876.
- 28
29
30 31 Locatelli, E.; Naddaka, M.; Uboldi, C.; Loudos, G.; Fragogeorgi, E.; Molinari, V.; Pucci, A.; Tsotakos, T.; Psimadas, D.; Ponti, J.; *et al.* Targeted delivery of silver nanoparticles and alisertib: in vitro and in vivo synergistic effect against glioblastoma. *Nanomedicine (Lond).* **2014**, *9*, 839-849.
- 31
32
33 32 Saluja, S. S.; Hanlon, D. J.; Sharp, F. A.; Hong, E.; Khalil, D.; Robinson, E.; Tigelaar, R.; Fahmy, T. M.; Edelson, R. L. Targeting human dendritic cells via DEC-205 using PLGA nanoparticles leads to enhanced cross-presentation of a melanoma-associated antigen. *Int J Nanomedicine.* **2014**, *9*, 5231-5246.
- 34
35
36 33 Shen, H.; Ackerman, A. L.; Cody, V.; Giodini, A.; Hinson, E. R.; Cresswell, P.; Edelson, R. L.; Saltzman, W. M.; Hanlon, D. J. Enhanced and prolonged cross-presentation following endosomal escape of exogenous antigens encapsulated in biodegradable nanoparticles. *Immunology.* **2006**, *117*, 78-88.
- 37
38
39 34 Panyam, J.; Zhou, W. Z.; Prabha, S.; Sahoo, S. K.; Labhasetwar, V. Rapid endo-lysosomal escape of poly(DL-lactide-co-glycolide) nanoparticles: implications for drug and gene delivery. *FASEB J.* **2002**, *16*, 1217-1226.
- 40
41
42 35 Miconi, G.; Palumbo, P.; Raysi Dehcordi, S.; La Torre, C.; Lombardi, F.; Evtoski, Z.; Cimini, A. M.; Galzio, R.; Cifone, M. G.; Cinque, B. Immunophenotypic characterization of human glioblastoma stem cells: correlation with clinical outcome. *J Cell Biochem.* **2015**, *116*, 864-876.
- 43
44
45 36 Jia, W.; Lu, R.; Martin, T. A.; Jiang, W. G. The role of claudin-5 in blood-brain barrier (BBB) and brain metastases (review). *Mol Med Rep.* **2014**, *9*, 779-785.
- 46
47
48 37 Hardee, M. E.; Zagzag, D. Mechanisms of glioma-associated neovascularization. *Am J Pathol.* **2012**, *181*, 1126-1141.
- 49
50
51 38 Engelhardt, S.; Patkar, S.; Ogunshola, O. O. Cell-specific blood-brain barrier regulation in health and disease: a focus on hypoxia. *Br J Pharmacol.* **2014**, *171*, 1210-1230.
- 52
53
54 39 Ernsting, M. J.; Murakami, M.; Roy, A.; Li, S. D. Factors controlling the pharmacokinetics, biodistribution and intratumoral penetration of nanoparticles. *J Control Release.* **2013**, *172*, 782-794.
- 55
56
57 40 Jordan, C. T.; Guzman, M. L.; Noble, M. Cancer stem cells. *N Engl J Med.* **2006**, *355*, 1253-1261.
- 58
59
60 41 Mannino, M.; Chalmers, A. J. Radioresistance of glioma stem cells: intrinsic characteristic or property of the 'microenvironment-stem cell unit'? *Mol Oncol.* **2011**, *5*, 374-386.


1
2
3 43 Bjorklund, M.; Koivunen, E. Gelatinase-mediated migration and invasion of cancer cells. *Biochim Biophys*
4 *Acta.* **2005**, *1755*, 37-69.

5 43 Urbanska, K. ; Pajak, B.; Orzechowski, A; .Sokolowska, J.; Grodzik, M.; Sawosz, E.; Szmidt, M.; Sysa, P. The
6 effect of silver nanoparticles (AgNPs) on proliferation and apoptosis of in ovo cultured glioblastoma
7 multiforme (GBM) cells. *Nanoscale Res Lett.* **2015**, *10*, 98.
8
9
10
11


12 ToC Graphic

13
14
15
16





25 **Figure 1.** (A) Schematic procedure for the synthesis of Ag-PNP-CTX. Representative TEM image
26 (B) DLS analysis (C) and ζ -potential analysis (D) of Ag-PNP-CTX.



1
2
3
4
5 **Figure 2.** Human-derived GBM cell lines U87MG, A172 and T98G were incubated o/n in the
6 presence of targeted (Ag-PNP-CTX) or non-targeted (Ag-PNP) nanovectors at Ag concentration of
7 100 μ M. The intracellular accumulation of Ag nanoparticles (Ag-nps) was determined by confocal
8 laser scanning microscopy (A,B,F,G) and 3D imaging analysis performed by IMARIS software (H
9 and I). (A) and (B) bright field images. (C-E) Pictures of Ag-nps: (C) bright field, (D) reflection mode
10 (as for figures F and G), (E) merge. (F) and (G) images represent the maximum projections of 20
11 optical sections of the cell labelled with Syto Blue 45® (green). Metal Ag-nps have been acquired in
12 reflection mode upon 458 nm excitation and are shown in red and DAPI signal for nuclei detection in
13 blue. (H) and (I) computation 3D images of the peri-nuclear region of cells in panel F and G
14 respectively. (L) Intracellular Ag-nps quantification expressed as mean number of Ag-nps ES
15 normalized on the cell surface (μm^2). About 60 cells / experimental group from 2 or 3 independent
16 experiments were analyzed. Significance *versus* cells incubated with non-targeted nanovector (Ag-
17 PNP): * $p < 0,02$; ** $p < 0,005$. (M) Competitive inhibition of cellular uptake of Ag-PNP-CTX and Ag-PNP
18 nanovectors in U87MG and A172 cells incubated with native CTX peptide at the indicated
19 concentrations (ng/ml). Results are expressed as Ag-nps quantity ES normalized to the Ag-nps
20 detected in cells incubated without native CTX peptide (first columns). Significance *versus* cells
21 incubated in the absence of CTX peptide: U87MG * $p = 0,0334$ ** $p = 0,0057$ *** $p = 0,0004$; A 172 **
22 $p = 0,0052$. (N) Representative confocal microscopy images showing U87-MG cells incubated in the
23 presence of Ag-PNP-CTX nanovectors. Intracellular endosomal and lysosomal compartments were
24 identified using the specific-associated markers Rab5a and LAMP-1 respectively.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58







Figure 3. U87MG (5×10^4) cells were injected intracerebrally into athymic 6-8 weeks old male mice. After 2 weeks, animals were euthanized with intracardiac perfusion of PBS, followed by formaldehyde. GBM xenografts were detected by immune staining for human nestin (green). In blue is the DAPI signal for nuclei detection. (A) representative coronal section 15 days after tumor implantation; (B) and (C) magnified views of the indicated insets.



31 **Figure 4** In vivo accumulation of Ag-PNP-CTX nanovectors at tumor site assessed in U87MG
 32 orthotopic xenografts model. Fifteen days after implantation of tumor cells mice were subjected to
 33 total brain radiation treatment (2Gy). A parallel group of non-irradiated mice (0 Gy) was considered
 34 as control. Ag-PNP-CTX were administered intraperitoneally 18 hours after radiations at a dose of
 35 1mg/mouse. Animals were sacrificed 24 or 48 hours after nanovector injection. Ag-nps quantification
 36 was performed in brain cryosections by confocal microscopy and IMARIS 3D imaging analysis as in
 37 Supplementary Figure 1. (A) Cryosections of U87MG tumor mass from irradiated (2 Gy) and non-
 38 irradiated (0 Gy) mice injected with Ag-PNP-CTX (0Gy). Ag-nps have been acquired in reflection
 39 mode upon 488 nm excitation and are shown in red. In green is the human nestin immunostaining
 40 and the nuclei are in blue (DAPI). (B) Ag-nps quantification is expressed as mean number of Ag-nps
 41 ES in a fixed volume of $6500 \mu\text{m}^3$. About 60 microscopy fields randomly taken from 4 mice in each
 42 experimental group were acquired and analyzed. Significance *versus* healthy brain tissue:
 43 *** $p < 0,0001$
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58



45 **Figure 5.** In vivo BBB integrity and Ag-PNP-CTX nanovector diffusion in peripheral tumor areas. (A-
46 D) Confocal images (maximum projections) of normal and tumor-associated blood vessels. Brain
47 cryosections from U87MG-transplanted mice previously irradiated (2Gy) or non-irradiated (0 Gy) were
48 immunostained for claudin-5 (red) and human nestin (green). Nuclei were revealed by DAPI staining
49 (blue). Lower panels are representative 3D rendering of selected regions of claudin-5 staining carried
50 out by IMARIS software. (E) Ag-nps and nestin co-localization in single cells present at the U87MG
51 tumor periphery from non-irradiated (0Gy, left panel) and irradiated (2Gy, middle and right panel)
52 mice injected with Ag-PNP-CTX. The isolated cell, labeled with antibodies recognizing human nestin
53 (in green), is a tumor cell which infiltrates the healthy mice tissue, which is negative for the human
54 nestin antibody. Ag-nps have been acquired in reflection mode upon 488 nm excitation and are
55 shown in red. In green is the human nestin immunostaining. In the bright field Ag-nps are visible as
56 black dots (middle panel).
57
58



35 **Figure 6 Radiation induced CTX-nanovector intracellular uptake, MMP-2 and CIC-3**
 36 **expression.** (A) Intracellular Ag-nps quantification in irradiated (2 Gy) and non-irradiated (0 Gy)
 37 U87MG or TG98 cells incubated overnight in the presence of Ag-PNP or Ag-PNP-CTX. Results are
 38 expressed as Ag-nps quantity ES normalized to the Ag-nps detected in non-irradiated cells
 39 incubated with Ag-PNP(0 Gy Ag-PNP, first column). For analysis details see also Supplementary
 40 figure 1. Significance *versus* non-irradiated cells: * $p=0,0117$. (B) Western-blot analysis of MMP-2
 41 and CIC-3 expression in irradiated and non-irradiated U87MG cells. One representative experiment
 42 out of 4 performed is reported. Levels of β -actin are reported as control for protein loading. (C)
 43 Densitometry analysis of western-blot bands (panel B) performed by Image-Lab software. Results
 44 normalized on corresponding β -actin. (D) Ag-nps (red) and MMP-2 or CIC-3 (green) signals in
 45 irradiated (2Gy) and non-irradiated (0 Gy) U87MG cells shown as maximum projections of 6 optical
 46 sections. In blue is the nuclear staining (DAPI). Cell borders were defined by phalloidin staining
 47 (gray).
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58





Figure 7. In vivo MMP-2 and CIC-3 expression in U87MG xenograft before and after radiations. (A-C) Representative confocal images of brain cryosections from U87MG-transplanted mice previously irradiated (2Gy) or non-irradiated (0 Gy) and immunostained for human MMP-2 or CIC-3 (red) and nestin (green). Nuclei were revealed by DAPI staining (blue). (B-D) In vivo MMP-2 and CIC-3 level of expression quantification in U87MG xenografts 48 and 72 hours after x-rays treatment. Results are expressed as increased mean MMP-2 and/or CIC-3 fluorescent area respect to non-irradiated samples (0 Gy) ES. About 15 microscopy fields for each experimental condition were randomly acquired and analyzed by ImageJ software. For each single microscopy field analyzed MMP-2 and/or CIC-3 fluorescent area was normalized on nuclei number. MMP-2 significance versus non-irradiated tumor cells * $p=0,0119$; *** $p=0,0005$.



24 **Figure 8.** Inhibition of MMP-2 catalytic activity in U87MG cells induced by Ag-PNP-CTX nanovectors.
 25 U87MG cells previously irradiated (2Gy) were incubated in the presence of Ag-PNP or Ag-PNP-CTX
 26 nanovectors (100 μ M for 16 hours). Inhibition of MMP-2 catalytic activity was evaluated by
 27 zymography. (A) Representative zymography of one out of 2 independent experiments performed.
 28 (B) Densitometry analysis of zymography bands. Results are expressed as mean values from 2
 29 independent experiments normalized to non-irradiated cells (0 Gy) ES. Statistical significance was
 30 evaluated by One-way ANOVA. Significance versus irradiated cells: * $p < 0,05$
 31




Figure 9. Effect of Ag-nps entrapped into PNP-CTX nanovectors on irradiated (2 Gy) and non-irradiated U87MG cells and tumors viability. (A) Upon treatment with the indicated nanoconstructs cell survival fraction was measured by MTT assay. Cells were exposed to increasing concentrations of Ag-nps for 72 hours. The obtained relative values were normalized to the values from the corresponding untreated cells and are shown as percentage survival. Results are expressed as mean percentage of 3 independent experiments performed in triplicates \pm SD. (B) In situ detection of apoptosis by DAPI/TUNEL staining and quantification of apoptotic bodies (1-5 μ m size) in U87MG cryosections from non-irradiated and irradiated (2 Gy) mice injected with Ag-PNP-CTX nanovectors. Animals were sacrificed 48 hours after nanovector injection. Four mice/experimental group were analyzed. Results are expressed as mean values normalized to CTR values \pm ES. (C) Luciferase imaging of representative mice: untreated (CTRL, 4 mice); injected with Ag-PNP-CTX nanovectors as single agent (2 doses of 1 mg/mouse followed by 2 doses of 0,5 mg/mouse, administration interval: 48-72 hours, 5 mice); irradiated (2 Gy single dose, 6 mice); treated with radiations and concomitant Ag-PNP-CTX nanovectors (6 mice). Treatments started 11 days after U87MG_{gluc2} cell implantation (T0=D11). (D) Tumor growth during 11 days of observation measured by area detected setting automatic ROI (50%). Results are expressed as mean values normalized to T0 values \pm SE. Statistical significance was evaluated by Paired T test. Significance: * p <0,05