



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

The role of SPICA-like missions and the Origins Space Telescope in the quest for heavily obscured AGN and synergies with Athena

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Barchiesi L., Pozzi F., Vignali C., Carrera F.J., Vito F., Calura F., et al. (2021). The role of SPICA-like missions and the Origins Space Telescope in the quest for heavily obscured AGN and synergies with Athena. PUBLICATIONS OF THE ASTRONOMICAL SOCIETY OF AUSTRALIA, 38, 1-19 [10.1017/pasa.2021.26].

Availability:

This version is available at: <https://hdl.handle.net/11585/863638> since: 2022-02-22

Published:

DOI: <http://doi.org/10.1017/pasa.2021.26>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Barchiesi, L., Pozzi, F., Vignali, C., Carrera, F., Vito, F., Calura, F., ...Tombesi, F. (2021). The role of SPICA-like missions and the Origins Space Telescope in the quest for heavily obscured AGN and synergies with Athena. *Publications of the Astronomical Society of Australia*, 38, E033.

The final published version is available online at:
<https://doi.org/10.1017/pasa.2021.26>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

The role of SPICA-like missions and the Origins Space Telescope in the quest for heavily obscured AGN and synergies with Athena

L. Barchiesi^{*1,2}, F. Pozzi^{1,2}, C. Vignali^{1,2}, F. J. Carrera³, F. Vito⁴, F. Calura², L. Bisigello², G. Lanzuisi^{1,2}, C. Gruppioni², E. Lusso^{5,6}, I. Delvecchio^{7,8}, M. Negrello⁹, A. Cooray¹⁰, A. Feltre^{2,11}, J. A. Fernández-Ontiveros¹², S. Gallerani⁴, H. Kaneda¹³, S. Oyabu¹⁴, M. Pereira-Santaella¹⁵, E. Piconcelli¹⁶, C. Ricci^{17,18,19}, G. Rodighiero²⁰, L. Spinoglio¹², F. Tombesi^{21,22,23}

(Affiliations can be found after the references)

Abstract

In the BH-galaxy co-evolution framework, most of the star-formation (SF) and the black hole (BH) accretion is expected to take place in highly obscured conditions. The large amount of gas and dust absorbs most of the UV-to-soft-X radiation and re-emits it at longer wavelengths, mostly in the IR. Thus, obscured Active Galactic Nuclei (AGN) are very difficult to identify in optical or X-ray bands, but shine bright in the IR. Moreover, X-ray background (XRB) synthesis models predict that a large fraction of the yet-unresolved XRB is due to the most obscured (Compton thick, CT: $N_{\text{H}} \geq 10^{24} \text{ cm}^{-2}$) of these AGN. In this work, we investigate the synergies between putative IR missions (using *SPICA*, proposed for ESA/M5 but withdrawn in October 2020, and Origins Space Telescope, *OST*, as ‘templates’) and the X-ray mission *Athena*, which should fly in early 2030s, in detecting and characterizing AGN, with a particular focus on the most obscured ones. Using an XRB synthesis model, we estimated the number of AGN and the number of those which will be detected in the X-rays by *Athena*. For each AGN we associated an optical-to-FIR spectral energy distribution (SED) from observed AGN with both X-ray data and SED decomposition, and used these SEDs to check if the AGN will be detected by *SPICA*-like or *OST* at IR wavelengths. We expect that, with the deepest *Athena* and *SPICA*-like (or *OST*) surveys, we will be able to photometrically detect in the IR more than 90 % of all the AGN (down to $L_{2-10\text{keV}} \sim 10^{42} \text{ erg/s}$ and up to $z \sim 10$) predicted by XRB synthesis modeling, and we will detect at least half of them in the X-rays. The spectroscopic capabilities of the *OST* can provide $\approx 51\,000$ and ≈ 3400 AGN spectra with $R = 300$ at $25 - 588 \mu\text{m}$ in the wide and deep surveys, respectively, the last one up to $z \approx 4$. *Athena* will be extremely powerful in detecting and discerning moderate- and high-luminosity AGN, allowing us to properly select AGN even when the mid-IR torus emission is “hidden” by the host-galaxy contribution. We will constrain the intrinsic luminosity and the amount of obscuration for $\sim 20\%$ of all the AGN (and $\sim 50\%$ of those with $L_{2-10\text{keV}} > 3.2 \times 10^{43} \text{ erg/s}$) using the X-ray spectra provided by *Athena WFI*. We find that the most obscured and elusive CT-AGN will be exquisitely sampled by *SPICA*-like mission or *OST* and that *Athena* will allow a fine characterization of the most-luminous ones. This will provide a significant step forward in the process of placing stronger constraints on the yet-unresolved XRB and investigating the BH accretion rate evolution up to very high redshift ($z \geq 4$).

Keywords: galaxies: active – galaxies: evolution – galaxies: star formation – infrared: galaxies – X-rays: diffuse background – techniques: photometric

PREFACE

The articles of this special issue focus on some of the major scientific questions that a future IR observatory will be able to address. We adopt the *SPace Infrared tele-*

scope for Cosmology and Astrophysics (SPICA) design as a baseline to demonstrate how to achieve the major scientific goals in the fields of galaxy evolution, Galactic star formation and protoplanetary disks formation and evolution. The studies developed for the *SPICA* mission

*luigi.barchiesi2@unibo.it

serve as a reference for future work in the field, even though the mission proposal has been cancelled by ESA from its M5 competition.

The mission concept of *SPICA* employs a 2.5 m telescope, actively cooled to below ~ 8 K, and a suite of mid- to far-IR spectrometers and photometric cameras, equipped with state of the art detectors (Roelfsema et al. 2018). In particular the *SPICA* Far-infrared Instrument (SAFARI) is a grating spectrograph with low ($R \sim 200 - 300$) and medium ($R \sim 3000 - 11\,000$) resolution observing modes instantaneously covering the $35 - 210 \mu\text{m}$ wavelength range. The *SPICA* Mid-infrared Instrument (SMI) has three operating modes: a large field of view ($10' \times 12'$) low-resolution $17 - 36 \mu\text{m}$ imaging spectroscopic ($R \sim 50 - 120$) mode and photometric camera at $34 \mu\text{m}$ (SMI-LR), a medium resolution ($R \sim 1300 - 2300$) grating spectrometer covering wavelengths of $18 - 36 \mu\text{m}$ (SMI-MR) and a high-resolution echelle module ($R \sim 29\,000$) for the $10 - 18 \mu\text{m}$ domain (SMI-HR). Finally, B-BOP, a large field of view ($2'.6 \times 2'.6$), three channel, ($70 \mu\text{m}$, $200 \mu\text{m}$ and $350 \mu\text{m}$) polarimetric camera complements the science payload.

1 INTRODUCTION

One of the main open issue in astrophysics is the role of Supermassive Black Holes (SMBH) in shaping the galaxies and the influence that the star formation (SF) has on the properties of the Active Galactic Nuclei (AGN). The scaling relations between the mass of the SMBH and the stellar mass (or luminosity) of the spheroid in which it is hosted (Kormendy & Richstone 1995) or the spheroid velocity dispersion (Ferrarese & Merritt 2000; Gebhardt et al. 2000) suggest a tight link between star-formation activity in the spheroidal components of galaxies and SMBH growth. The BH-galaxy co-evolution paradigm states that an intense phase of SF is triggered by a wet merger, at least for the most luminous and massive systems (Silk & Rees 1998; Di Matteo et al. 2005; Lamastra et al. 2013). A fraction of the gas reservoir of the galaxy is funneled towards the SMBH and turns on the AGN activity. Hydrodynamical simulations (e.g. Di Matteo et al. 2005; Hopkins et al. 2006; Barai et al. 2018), semi-analytic models (e.g. Menci et al. 2008) and observations (e.g. Tombesi et al. 2015; Feruglio et al. 2015; Smith et al. 2019) suggest a scenario where AGN provide feedback that regulates the SF in the host galaxies, thus originating the observed correlations. Such link between SMBH growth and galactic star formation history is supported by the striking similarity of the evolution of the SMBH accretion rate (BHAR) and that of the star formation rate (SFR) densities, especially at high redshifts where the SF mostly occurs in the spheroidal components (Shankar et al. 2009; Fiore et al. 2017; Gruppioni et al. 2020). The evolutions of the SFR and the BHAR follow a parallel evolution, showing both a peak of activity

around $z \approx 1 - 3$, known as the “cosmic noon” (Madau & Dickinson 2014; Heckman & Best 2014; Vito et al. 2018). However, the mechanisms responsible for this similarity are not well understood and the co-evolution between galaxy and SMBH is still a matter of debate. Investigating the BH-galaxy co-evolution requires sizable samples of galaxies and AGN over a broad range of redshift and luminosities, to characterize the accretion and SF histories and disentangle the two contributions within the same source as accurately as possible.

To study the link between star formation and black hole accretion we need, in particular, to investigate the epoch in which the bulk of these growths seems to take place (i.e. the cosmic noon). However, the high quantity of gas and dust that fuels both processes absorbs most of the energy emitted by stars and accreting SMBHs and re-emits it at longer wavelengths, mostly in the IR. The search for and the characterization of these objects in the optical and UV is challenging due to the heavy reddening that affects them (Rowan-Robinson et al. 1997; Hughes et al. 1998). A solution to this problem is to study indirectly the primary emission, by measuring the dust-reprocessed radiation in the IR. A study at such wavelengths allows us to observe the physical processes at work in the obscured regions and, therefore, to derive the SFR density (SFRD) and the BHAR density (BHAD) if we are capable of properly separating them. A direct measure of the dust-obscured SF activity is possible with space-based IR observatories, which do not need to be corrected by dust attenuation.

The mid- to far-IR photometric observations in deep fields (Delvecchio et al. 2014; Schreiber et al. 2015) have provided extremely reliable determination of the SFRD and the BHAD up to redshift $\sim 3 - 4$ (Gruppioni et al. 2013; Magnelli et al. 2013; Delvecchio et al. 2014) using *Herschel* PACS data (observing at 100 and $160 \mu\text{m}$; Poglitsch et al. 2010) and have detected high redshift galaxies (up to $z > 6$; e.g., Riechers et al. 2013; Rowan-Robinson et al. 2016) using *SPIRE*, observing at 250 , 350 and $500 \mu\text{m}$; Griffin et al. 2010). However, the deepest cosmological surveys performed by *Herschel* at high- z have detected only the most luminous galaxies ($L_{\text{IR}} > 10^{12} L_{\odot}$ at $z \geq 3$; Gruppioni et al. 2013) and suffer from the uncertainties of not being able to accurately disentangle the AGN from the SF emissions, as both contributes to the mid- and far-IR emission. Even in the case when Spectral Energy Distribution (SED) fitting allows us to separate the two components, the results usually depend on the adopted technique and modeling. Therefore, the initial phase of the BH-galaxy co-evolution paradigm has been, so far, the most elusive and difficult to track. According to the co-evolution scenario, this heavily obscured phase should be characterized by the presence of a strong starburst and an obscured AGN, both giving a significant contribution to the mid- and far-IR emission (e.g. Page et al. 2004;

Vignali et al. 2009; Lapi et al. 2018).

Compton-thick (CT) AGN, i.e. highly-obscured AGN with column density $\log(N_{\text{H}}/\text{cm}^{-2}) \geq 24$, are also required to explain the spectra of the extragalactic X-ray background (XRB): their flat spectrum is needed to reproduce the hump observed at 20 – 30 keV in the XRB (e.g. Comastri et al. 1995; Gilli et al. 2007; Balantyne et al. 2011). However, the fraction of CT-AGN derived from XRB synthesis models ranges from 10 % to 30 – 50 % (Treister et al. 2009; Gilli et al. 2007; Ueda et al. 2014; Ananna et al. 2019) and suffers from large uncertainties due to degeneracies between several model parameters. Although the expected number of CT-AGN is large, their detection and identification beyond the local universe are challenging and, even in the local universe, their observed fraction ($\sim 5 - 10\%$, e.g. Burlon et al. 2011; Ricci et al. 2015) is significantly lower than models predictions, but X-rays detection bias may be responsible of such low numbers (Burlon et al. 2011).

To select unobscured or mildly-obscured AGN, one of the most effective tools is the observation in the X-ray band, thanks to the radiation coming directly from the AGN innermost regions. However, this selection loses effectiveness with the increasing column density (at $N_{\text{H}} \geq 10^{23} \text{ cm}^{-2}$ the soft continuum is severely depressed) and, even in deep fields, only a limited fraction of the most obscured AGN has been revealed in the X-ray (e.g. Tozzi et al. 2006; Lanzuisi et al. 2013; Marchesi et al. 2016; Del Moro et al. 2017). Moreover, the BHAD estimation from the X-ray luminosity is strongly dependent on the bolometric correction (k_{bol}), that allows to estimate the “total” bolometric power (L_{bol}) of the AGN from the AGN emission in a specific band, but suffers from significant uncertainties in the $k_{\text{bol}} - L_{\text{bol}}$ relation (e.g. Hopkins et al. 2007; Lusso et al. 2010, 2012; Duras et al. 2020). Likewise for obscured SF galaxies, the IR is an effective band in detecting these obscured AGN, thanks to the energy absorbed by the obscuring material and re-emitted at sub-mm wavelengths (e.g. Pozzi et al. 2007). The effectiveness of identifying highly obscured AGN by selecting bright mid-IR sources with faint optical or near-IR emission has been shown in the past by *Spitzer* (e.g. Houck et al. 2005; Weedman et al. 2006; Polletta et al. 2008) and *WISE* (e.g. Mateos et al. 2012; Assef et al. 2013).

To be able to photometrically identify highly-obscured and CT-AGN, both mid-IR and far-IR instruments are necessary, as we need to distinguish between the emission from hot dust ($T > 100 \text{ K}$), mostly due to reprocessed AGN emission from the dusty torus and galactic dust, and the emission from warm/cold dust ($T < 100 \text{ K}$) dominated by the reprocessed SF emission.

A future IR cryogenic cooled space telescope may be able to deliver all of this. We investigated two different IR mission, using as ‘templates’ the proposed, now withdrawn, ESA *SPICA* mission and the NASA concept

study *Origins Space Telescope (OST)*. A *SPICA*-like mission (described extensively in section 2.1) with a mid-IR spectrophotometer will be able to sample the band where the peak of the CT-AGN emission is up to $z \approx 6$ and provide also low-resolution spectra, while far-IR photometric cameras will cover the wavelength range where the SF is the main contributor, thus helping in separating the two emission mechanisms. Detected sources may be followed-up with *SPICA SAFARI*-like instruments to provide low resolution spectra in the mid- and far-IR wavelength range up to $z \approx 3.5$. In these spectra high ionization lines, such as [Ne v] at $14.32 \mu\text{m}$ and $24.32 \mu\text{m}$, [Ne v I] at $7.7 \mu\text{m}$ and [O iv] at $25.89 \mu\text{m}$, would be signatures of AGN presence and their fluxes could be used as proxy of the AGN bolometric luminosities (e.g. Spinoglio & Malkan 1992; García-Bernete et al. 2016; Fernández-Ontiveros et al. 2016). Alternatively an highly capable grating mid- and far-IR spectrometer, as the *OSS* in the *OST* (see section 2.2), will be able to cover instantaneously both the hot and warm/cold dust wavelength regime, thus enabling us to distinguish between the torus and the host-galaxy emission and to estimate, in a reliable way, the AGN bolometric power, the host SF and the BH accretion rate, without the need of additional constraints from other optical-IR instruments or surveys.

To confirm the CT nature of sources, X-ray spectroscopy has been the most reliable method so far. Unfortunately, with the current deepest X-ray surveys, we have only scratched the surface of the obscured AGN population at $z \sim 1 - 4$. Moreover, the identification of an AGN as CT depends strongly on the adopted analysis techniques and modeling, with different techniques giving sometimes different classifications of the same sources (e.g. Castelló-Mor et al. 2013). The best results have come from X-ray deep and ultradeep surveys (e.g. COSMOS, Elvis et al. 2009; Civano et al. 2016, CDF-S, Giacconi et al. 2002; Luo et al. 2008; Xue et al. 2011; Xue 2017) that allowed to detect CT-AGN even at high redshift, but with the disadvantage of having small fields, thus low source statistics. *XMM-Newton* and *Chandra* surveys unveiled only up to few tens of candidates of CT-AGN (e.g. Comastri et al. 2011; Brightman & Ueda 2012; Georgantopoulos et al. 2013; Lanzuisi et al. 2017), while an hard X-ray mission like *NuSTAR* has difficulties in detecting objects beyond $z \sim 1$, due to its sensitivity (Lansbury et al. 2017a,b, the notable exceptions of CT-AGN at $z \sim 2$ detected by Del Moro et al. 2014; Zappacosta et al. 2018).

A breakthrough in selection and identification of CT-AGN is expected with the new ESA X-ray observatory *Athena*, due for launch in early 2030s. *Athena*, with its combination of angular resolution, field of view and collecting area at $\sim 1 \text{ keV}$, is the ideal instrument to perform X-ray surveys and will be more than two orders of magnitude faster than *Chandra* or *XMM-Newton*

(Nandra et al. 2013). The excellent spectral capabilities of *Athena WFI* for surveys will yield samples of the most heavily obscured AGN up to redshifts $z = 4$ and up to 100 times larger than is currently possible. Since $L_{\text{X-ray}} \gtrsim 10^{42} \text{ erg s}^{-1}$ are commonly associated with AGN activity, *Athena* will also be invaluable in quickly identifying moderate- and high-luminous AGN, especially in cases where we can not distinguish between the torus mid-IR emission and the host-galaxy contribution from SED fitting analysis.

The planned launch date for *Athena* is $\gtrsim 2030$. Having, in the same years, a new IR mission dedicated (also) to survey high- z galaxy and AGN would be an extraordinary opportunity to study obscured accretion and to characterize AGN up to high redshift. A possible example of such mission would have been *SPICA*, but also the NASA concept *OST* for an 4.5 K cooled IR space observatory, which has a proposed launch date of ~ 2035 and similar survey strategy and wavelength coverage as *SPICA*. Having data at X-ray and IR wavelengths will be the key to accurately reconstruct the accretion luminosity across cosmic time up to very high redshift ($z \approx 6$). The search for the elusive CT-AGN may decisively benefit from the synergies between *Athena* and future IR mission: while a *SPICA*-like telescope and the *OST* should be able to effectively detect the majority of them (even those too obscured to be detected in X-rays), the X-ray spectra provided by *Athena* will be invaluable to confirm their CT-nature, even in paucity of photons, for a large fraction of them.

The goal of this paper is to study the synergies between *Athena* and future IR surveys in searching and identifying AGN, with a particular focus on the most obscured ones. We will show how the search for these elusive CT-AGN, very difficult to identify and characterize with the current instruments, can benefit from a combined use of IR and X-ray data. In Section 2 we give a brief description of the *SPICA* and *OST* instruments and planned surveys and a comparison with the *Athena* surveys. The simulation of the intrinsic AGN prediction obtained from the XRB synthesis models are presented in Section 3, while the number of those which will be detected by *Athena* are reported in Section 4. In Section 5 we present the optical-to-far-IR SED used to predict the AGN emission and their detectability with *OST* and *SPICA*-like mission. The prediction of the number of AGN that will be detected by the IR surveys, as well as the synergies with *Athena* that we can exploit to study them, are reported in Section 6. In Section 7, we present our conclusions.

Throughout this paper, we adopt the following cosmological parameters: $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_{\text{M}} = 0.3$ and $\Omega_{\Lambda} = 0.7$ (Spergel et al. 2003).

2 INSTRUMENTS AND SURVEYS

2.1 SPICA

SPICA was a joint European-Japanese project to develop a new generation cryogenic infrared space telescope, with the early design dating back to more than two decades ago (e.g. Nakagawa et al. 1998, 2014; Swinyard et al. 2009). Here we discuss the *SMI* and *B-BOP* instruments as it were designed during the ESA M5 selection process and a *SPICA*-like survey strategy.

2.1.1 SMI

SMI is composed of the four channels, *SMI-CAM*, *LR*, *MR* and *HR*. In this paper, we consider *SMI-CAM* and *SMI-LR*, that work simultaneously. *SMI-CAM* is a mid-IR slit viewer photometric camera that covers the $30 - 37 \mu\text{m}$ range and provides $34 \mu\text{m}$ broad-band images with a field of view of $10' \times 12'$ (with the exclusion of the four slits used by the spectrometer). *SMI-LR* is a wide field-of-view multi-slit prism spectrometer, composed of four long slits of $10'$ in length and $3.7''$ in width. It has a resolution of $R = 50 - 120$ and allows spectroscopic surveys in the $17 - 36 \mu\text{m}$ wavelength range. The *SMI-CAM* provides photometric scientific data, and is invaluable in accurately determining the position of the slits on the sky for pointing reconstruction in creating spectral maps. We refer to Kaneda et al. (2017) for further instrument specifications and survey strategies.

2.1.2 B-BOP

The B-BOP polarimetric instrument allows simultaneous imaging observations in three bands, centered at $70 \mu\text{m}$, $200 \mu\text{m}$ and $350 \mu\text{m}$, over an instantaneous field of view from $1.8' \times 1.8'$ to $2.7' \times 2.7'$ at FWHM resolutions of $6''$, $17''$ and $30''$, respectively. B-BOP is two to three orders of magnitude more sensitive than current or planned far-IR/submillimeter polarimeters, e.g. *SHARP* (Li et al. 2008), *HAWC+* (*SOFIA*) (Dowell et al. 1998), *BLAST-TNG* (Galitzki et al. 2014). It provides wide-field $70 - 350 \mu\text{m}$ polarimetric images in Stokes Q and U of comparable quality (in terms of resolution, signal-to-noise ratio, and both intensity and spatial dynamic ranges) to *Herschel* images in Stokes I. More details about the B-BOP instrument are in Rodriguez et al. (2018) and André et al. (2019).

2.1.3 Surveys with SMI

Hereafter, we consider two reference blind spectroscopic surveys:

- a **1deg² Ultradeep survey** with a total observational time of 600 hr, an *SMI-CAM* 5σ sensitivity of $\approx 3 \mu\text{Jy}$ and an *SMI-LR* 5σ sensitivities (high-background case) of $\approx 50 \mu\text{Jy}$ at $20 \mu\text{m}$, $\approx 110 \mu\text{Jy}$ at $30 \mu\text{m}$ and a line sensitivity of $\approx 9.2 \times 10^{-20} \text{ W/m}^2$ (this survey configuration corre-

sponds to the Kaneda et al. 2017 deep survey);

- a **15 deg² Deep survey** with a total observational time of 600 hr, an SMI-CAM 5σ sensitivity of $\approx 13\ \mu\text{Jy}$ and an 5σ SMI-LR sensitivities (high-background case) of $\approx 160\ \mu\text{Jy}$ at $20\ \mu\text{m}$, $\approx 380\ \mu\text{Jy}$ at $30\ \mu\text{m}$ and a line sensitivity of $\approx 33 \times 10^{-20}\ \text{W/m}^2$ (this survey has the same depth of the Kaneda et al. 2017 shallow survey, although over a larger area).

2.2 The Origins Space Telescope

The *OST* (Battersby et al. 2018; Meixner et al. 2019) is a concept study for a Far-Infrared Surveyor mission, the subject of one of four science and technology definition studies supported by NASA for the 2020 Astronomy and Astrophysics Decadal Survey. The *OST* is being designed with the aim of covering a large area of the sky, thus allowing to search for rare objects at low and high redshifts.

OST is composed of a 5.9 m-diameter telescope with a *Spitzer*-like architecture, actively cooled down to 4.5 K. It is designed with three onboard instruments: the *Origins Survey Spectrometer* (OSS) to cover the 25–588 μm wavelength range instantaneously at $R=300$, *MISC-T* (*Mid-Infrared Spectrometer Camera Transit*) a 2.8 – 20 μm transit spectrometer and the *Far-infrared Imager and Polarimeter* (*FIP*), that delivers imaging or polarimetry at 50 μm and 250 μm . In this paper, we take into consideration primarily the OSS instrument, as it is perfectly suited for detecting and characterizing AGN in blind surveys.

2.2.1 OSS

The *Origins Survey Spectrometer* is a $R=300$ spectrometer covering the full 25 – 588 μm wavelength range instantaneously, using six logarithmically-spaced grating modules. True 3D spectral mapping is performed thanks to the 30 to 100 spatial beams that each module couples. Higher spectral resolution can be achieved by inserting into the light-path two mirrors that divert the light into interferometer optics allowing a $R = 43\ 000$ resolution, keeping the full spectral range but working only on single pointing (i.e. losing the survey capabilities). Even higher $R = 300\ 000$ spectral resolution can be achieved using an insertable etalon, further restricting the FoV and reducing the spectral range to 100 – 200 μm . As in this paper we focus on detecting AGN in surveys, we take into consideration only the base $R=300$ spectroscopy mode.

The six OSS bands cover the 25 – 44, 42 – 74, 71 – 124, 119 – 208, 200 – 350, 336 – 589 μm wavelength ranges, with beam sizes of 1.41, 2.38, 4.0, 6.78, 11.3, 19.0 arcsec and 1 hr 5σ sensitivities ($R=300$) of 22, 28, 40, 104, 104, 338 μJy . With the current OSS specs, an improvement of more than a factor 1000 in sensitivity is expected with respect to already flown far-IR observato-

ries (*SOFIA*, *Herschel-SPIRE*, and *Herschel-PACS*). We refer to Meixner et al. (2019) for further instrument specifications.

2.2.2 Survey with OSS

For *OST* we consider two blind blank-field spectroscopic surveys as described in Meixner et al. (2019):

- a **0.5 deg² deep survey** with an observational time of ≈ 1000 hr, a $R=4$, 5σ sensitivity of 4.5, 5.8, 8.4, 21.8, 21.8, 70.9 μJy (for channel 1 to 6) and a $R=300$ 5σ sensitivity of 39, 50, 72, 189, 189, 614 μJy ;
- a **20 deg² wide survey** with an observational time of ≈ 1000 hr, a $R=4$, 5σ sensitivity of 30, 39, 56, 145, 145, 473 μJy (for channel 1 to 6) and a $R=300$ 5σ sensitivity of 262, 336, 483, 1260, 1260, 4094 μJy .

2.3 Athena

Athena (Advanced Telescope for High ENergy Astrophysics ¹ Barcons et al. 2017) is the X-ray observatory mission selected by ESA, within its Cosmic Vision programme, to address the *Hot and Energetic Universe* scientific theme. It is the second L(large)-class mission within that programme and is due for adoption in late 2022 and launch in early 2030s.

Athena will consist of a single large-aperture grazing-incidence X-ray telescope, utilizing a novel technology (High-performance silicon pore optics) developed in Europe, with 12 m focal length and 5 arcsec Half Energy Width (HEW) on-axis angular resolution, degrading gradually to less than 10 arcsec at 30 arcmin off-axis (Bavdaz et al. 2018). There will be two instruments in the focal plane. One is the Wide Field Imager (WFI)² (Meidinger et al. 2018) providing simultaneous sensitive wide-field imaging and spectroscopy ($\text{FWHM} \leq 170\ \text{eV}$ at 7 keV) and high count-rate capability ($>90\%$ throughput and $<1\%$ pile-up for 1 Crab) over the 0.2 to 15 keV energy range. This is achieved through two sets of Silicon-based detectors using DEPFET Active Pixel Sensor technology: the Large Detector Array is a mosaic of 2×2 arrays spanning a $\sim 40 \times 40$ arcmin² Field of View (FoV) oversampling the PSF by more than a factor of two and the Fast Detector is a single array optimised for high count rate applications.

The other instrument is the X-ray Integral Field Unit (X-IFU)³ (Barret et al. 2018) delivering simultaneous spatially resolved (5 arcsec pixels) high-resolution X-ray spectroscopy ($\text{FWHM} < 2.5\ \text{eV}$ below 7 keV) over a limited field of view (~ 5 arcmin equivalent diameter) over the 0.2 to 12 keV energy range, with high count-rate capability (10 eV spectral resolution at 1 Crab inten-

¹<https://www.the-athena-x-ray-observatory.eu/>

²<http://www.mpe.mpg.de/ATHENA-WFI/>

³<http://x-ifu.irap.omp.eu/>

sities with low pile-up and >50% throughput). This performance is based on a large format array of superconducting molybdenum-gold Transition Edge Sensors coupled to absorbers made of gold and bismuth, cooled at ~ 90 mK inside a nested set of cryostats.

2.3.1 Surveys with Athena

Some of the core science objectives of *Athena*⁴ (e.g. finding distant evolved groups of galaxies at $z > 2$, complete the census for high-redshift and heavily obscured AGN) require performing surveys of the X-ray sky with the WFI. These surveys are expected to take a significant part of the observation time during the four years of the nominal mission lifetime. At the time of writing, the nominal number of pointings and exposure times (the so-called ‘‘Tier 2 post-CORE strategy’’) includes an ‘‘ultradeep’’ layer (4×1400 ks), a ‘‘deep’’ layer (3×980 ks + 7×840 ks) and a ‘‘shallow’’ layer (106×84 ks) (see Table 1). Each pointing covers ~ 0.4 deg², so the total area of each layer is 1.6, 4 and 42.5 deg², respectively. The exact places and geometries (single pointings vs. mosaics) of those layers will of course be determined at a later stage by the international astronomical community through the guaranteed time and open time, but they are expected to include regions of the sky with substantial multi-wavelength coverage. An initial estimate is encapsulated in the *Athena* Mock Observing Plan (MOP).

2.4 Surveys comparison

Athena WFI, *OST* OSS, and *SPICA* SMI have planned survey with different layers of depth and area. In this work we compared the different layers, matching them to have similar areas as much as possible. As we compared the expected number of sources per deg² for all the surveys and investigated the fraction of sources that may be detected, the exact area coverage is not critical. We investigate the capabilities of the observatories considering a **DEEP** survey (comparing the *SPICA* ultradeep survey with the *Athena* ultradeep survey and the *OST* deep survey), and a **WIDE** survey (composed of the *SPICA* deep survey, the *Athena* shallow survey and the *OST* wide survey). Table 1 summarizes the main parameters of all the *SPICA*, *OSS* and *Athena* surveys, while Table 2 reports the naming scheme used in this work.

3 SIMULATION OF THE INTRINSIC AGN COUNTS FROM XRB SYNTHESIS MODEL

We have estimated the ‘intrinsic’ AGN number as a function of the redshift z , 2-10 keV intrinsic X-ray lu-

minosities L_x and intrinsic column density N_H based on the Gilli et al. (2007) X-ray background synthesis model. We have started by dividing the (z, L_x, N_H) space in bins, with 21 z bins $\in [0, 10]$, 18 L_x bins with $L_x \in [10^{42}, 1.5 \times 10^{48}]$ erg/s and 6 N_H bins with $\log(N_H/\text{cm}^{-2}) \in [20, 26]$. We, then, make use of the software POMPA⁵ to get the $\log N(> S)$ source density in each of those bins, with S being the 2-10 keV source flux.

From the $N(> S)$ we also obtained the total intensity of each bin (energy received per unit area, unit time and unit sky area). Instead of the composite power law plus reflection spectral model in Gilli et al. (2007), we have adopted here a torus model from Brightman & Nandra (2011) with photon index $\Gamma = 1.9$, aperture angle 30 deg and inclination 80 deg. We have also included an additional scattering component modelled with a power-law with the same photon index and normalization equal to one percent of that of the primary emission.

In addition, we also included ‘normal’ galaxies using the galaxy $\log N - \log S$ curve of Lehmer et al. (2012) and a power-law spectral shape with $\Gamma = 1$ (the median value for the 332 sources with spectral slope not based on upper limits in Lehmer et al. 2012) and $S \in [5 \times 10^{-20}, 5 \times 10^{-16}]$ erg cm⁻² s⁻¹.

Physical fluxes are converted to counts using the latest (at the time of writing) set of FoV-averaged without filter WFI matrices⁶ using *xspec*⁷ (Arnaud 1996).

4 PREDICTIONS OF ATHENA SOURCE COUNTS

When observed by the WFI on board *Athena*, all the sources above are seen against a ‘background’ which includes several components^{8,9}:

- Particle background: due to cosmic rays impacting the detector and the surrounding structure. It is assumed to contribute 6×10^{-4} cts/keV/s/arcmin² with a flat power-law spectrum characterized by a spectral index of $\Gamma = 0$.
- Diffuse Galactic X-ray background: it is assumed to be uniform and to have the spectral shape and intensity from McCammon et al. (2002), which includes two thermal components, one unabsorbed representing emission from the Local Hot Bubble and one with foreground absorption by our own Galaxy representing emission from the hot Galactic

⁵<http://www.bo.astro.it/~gilli/counts.html>

⁶http://www.mpe.mpg.de/ATHENA-WFI/response_matrices.html

⁷<https://heasarc.gsfc.nasa.gov/xanadu/xspec/>

⁸https://www.mpe.mpg.de/ATHENA-WFI/public/resources/background/WFI-MPE-ANA-0010_i7.1_Preparation-of-Background-Files.pdf

⁹Exact matrix used: `athena_wfi_rib2.3_B4C_20190122_wo_filter_FovAvg.rsp`

⁴The full list of science requirements can be found under <https://www.cosmos.esa.int/web/athena/study-documents>

Table 1 Parameters of the *SPICA*, *Athena* and *OST* surveys. The time per field t_{field} are pure integration times, without over-heads. The sensitivities are computed at 5σ . The *SPICA* R=150 sensitivities are reported at $20\mu\text{m}$. The B-BOP sensitivities refer only to the $70\mu\text{m}$ channel, while those for the *OST* refer to channel 1 and channel 6. In this table, the *Athena* survey strategy is simplified, as the deep survey encompasses also the ultradeep pointings and the shallow survey will comprehend also the deep pointings. The *Athena* sensitivities refer to the $2 - 10\text{keV}$ band. The surveys investigated and the naming scheme used in this work are summarized in Table 2.

Instrument	Survey	t_{field} (ks)	Sensitivity (μJy)	t_{tot} (hr)	Area (deg^2)	
<i>SPICA</i>	SMI	ultradeep	67.86	3 (R=4), 50 (R=150)	605	1
	B-BOP		-	60	70	
	SMI	deep	5.22	13 (R=4), 160 (R=150)	482	15
	B-BOP		-	100	350	
<i>OST</i>	OSS	deep	-	4.5 – 79 (R=4), 39.4 – 614 (R=300)	1000	0.5
		wide	-	30 – 470 (R=4), 260 – 4090 (R=300)	1000	20
<i>Athena</i>	WFI	ultradeep	4×1400	$\sim 3 \times 10^{-17} \text{ erg/s/cm}^2$	~ 1600	1.6
		deep	$3 \times 980 + 7 \times 840$	$\sim 4 \times 10^{-17} \text{ erg/s/cm}^2$	~ 2500	4
		shallow	106×84	$\sim 1.2 \times 10^{-16} \text{ erg/s/cm}^2$	~ 2500	42.5

Table 2 Reference surveys used in this work. The surveys composing the reference surveys were matched in similar area coverage.

Instrument	Survey	Area (deg^2)	ref. survey
SPICA	ultradeep	1.0	DEEP 1 deg^2
OST	deep	0.5	
Athena	ultradeep	1.6	
SPICA	deep	15	WIDE 15 deg^2
OST	wide	20	
Athena	shallow	42.5	

halo, in `xspec` parlance: `apc1 + phabs (apc2)` with $kT_1=0.099\text{ keV}$ and $kT_2=0.225\text{ keV}$ and respective `apc` normalizations of 1.7×10^{-6} and 7.3×10^{-7} , corresponding to 1 arcmin^2 ; the column density for the Galactic absorption is assumed to be the same as for extragalactic sources (see below).

- XRB: due to the integrated emission of the above AGN and galaxy populations. It is assumed to have intensity and power-law spectral shape from [McCammon et al. \(2002\)](#) but with $\Gamma = 1.45$ and normalisation $10^{-6}\text{ photons s}^{-1}\text{cm}^{-2}\text{keV}^{-1}$, again normalized to 1 arcmin^2 .
- Stray light: X-ray photons from the two previous components, coming from outside the FoV but impacting the WFI after a single reflection on the mirror. We have modelled it by averaging the contribution of the two previous components over the full WFI FoV and fitting it with two thermal components and a power-law, with the same temperatures and photon index but with free relative normalizations.

All extragalactic components are assumed to undergo absorption by gas in our own Galaxy with hydrogen

column density of $N_{\text{H}} = 1.8 \times 10^{20}\text{ cm}^{-2}$, which is the average Galactic column density of the estimated survey pointings in the MOP and coincides with the foreground column density in [McCammon et al. \(2002\)](#).

Depending on the (z, L_x, N_{H}) and the exposure time, a fraction f_{det} of the sources in each of the bins, described above, will be detected, contributing to resolve a fraction of the full intensity of the extragalactic XRB, thus lowering the remaining (unresolved) background. For each of the exposure time values of the survey layers, we have determined the resolved fraction iteratively starting from an initial fiducial value of 80% and f_{det} is also determined for each bin in each iteration. To decide whether a source is detected above the background we have used Cash statistic C ([Cash 1979](#)): calling T the total source+background counts, B the background counts and t the exposure time, the mode (most probable value) of the source counts is $\hat{s} = \max(0, T - B)$. For a 5σ detection for a single free parameter (s) we required that the Cash statistic improves by $\Delta C = 25$ between $s = 0$ (i.e. no source) and $s = \hat{s}$. For each bin, 100 sources are simulated at the bin centre and f_{det} is the fraction of those sources that are detected. It turns out that the resolved fraction is around 80% and varies a few percent for exposure times above $\sim 10\text{ ks}$. In order to estimate the source+background and the background counts we have assumed a circular extraction area of radius 5.7 arcsec , which is the WFI FoV-weighted average of the *Athena* mirror HEW.

4.1 *Athena* source characterization

By their own nature, X-ray detectors work in photon-counting mode. One of the consequences of this is that they all are "Integral Field Units", providing spectra for all detected sources. For *Athena*, this ranges from the

superb ~ 2.5 eV resolution of X-IFU to the modest (but good for a Si-based detector) 170 eV resolution at 7 keV of WFI. We have simulated 100 spectra at the "centres" of each of the $1 \leq z \leq 4$, $10^{44} \leq L_x(\text{erg/s}) \leq 5 \times 10^{45}$, $24 \leq \log(N_H/\text{cm}^{-2}) \leq 26$ bins and fitted them. If both the fitted luminosity and the column density, obtained from the spectral fitting, are within 30% of the input values, we say that we have "characterized" the source. Finally, by dividing the number of characterized sources by 100 (the number of spectra we simulated) we have estimated the fraction of characterized sources in each bin.

5 AGN AND HOST GALAXY SED

To compute the AGN and host galaxy expected fluxes at different redshifts in the *SPICA* and *OST* bands, we need a proper set of SEDs to model the AGN as a function of different parameters: the intrinsic rest-frame 2 – 10 keV luminosity L_x , the equivalent hydrogen column density N_H and the redshift z .

We started with the AGN and host galaxy SED compilation from Lanzuisi et al. (2017). They used a sample of 2333 X-ray selected (either *XMM-Newton* or *Chandra*) AGN in the COSMOS field (Scoville et al. 2007) with at least 30 X-ray counts from Lanzuisi et al. (2013, 2015), Marchesi et al. (2016) and then selected only those with a $> 3\sigma$ IR detection in one of the *Herschel* bands. The X-ray spectral analysis provided us with column density N_H and absorption-corrected 2 – 10 keV luminosity for each source (see Lanzuisi et al. 2017 and references therein for more details). The final sample was composed of 692 sources X-ray and FIR detected, all with both X-ray spectral properties and SED decomposition. The SEDs were obtained by Delvecchio et al. (2014, 2015), following the recipes described in Berta et al. (2013), using the `sed3fit` code (with a "smooth" torus model), which allowed us to disentangle the AGN and host-galaxy contribution using three components – stellar emission, AGN torus emission and SF-heated dust emission – with photometric points from the UV to sub-mm. We excluded from our sample the sources with 2-10 keV rest-frame luminosities $\log(L_x/\text{erg/s}) < 42$, as SF galaxies without AGN should not be able to exceed this threshold. Moreover, to overcome the degeneracy between the AGN and the SF contribution during the SED fitting, we selected only the SEDs with AGN significance $S_{AGN} \geq 99\%$, evaluated through the F-test between the best-fit reduced χ^2 with and without the AGN component (see Delvecchio et al. 2014). Thus, the final sample is composed of 422 AGN in the redshift range $0 < z \leq 4$, 2-10 keV rest-frame luminosities in the $42 \leq \log(L_x/\text{erg/s}) \leq 45.5$ range and host-galaxy masses $7.9 \leq \log M/M_\odot \leq 12.2$.

With the aim of obtaining template SEDs representative of AGN with different intrinsic rest-frame 2 – 10

Table 3 Naming scheme used in this work for referring to AGN with different amounts of obscuration and X-ray luminosity. As reported in section 5, we used the Lusso et al. (2012) bolometric correction (eq 1) to compute L_{bol} .

$\log(N_H/\text{cm}^{-2})$		
20 – 22		unobscured
22 – 24		obscured
24 – 26		CT-AGN
$\log(L_{\text{bol}}/\text{erg s}^{-1})$	$\log(L_x/\text{erg s}^{-1})$	
43.0 – 44.6	42.0 – 43.5	low-luminosity
44.6 – 49.7	43.5 – 48.2	high-luminosity

keV luminosity and obscuration, we created four L_x bins and four N_H bins. The binning was chosen to maximize the number of sources in each bin, whilst maintaining a good sampling of the L_x and N_H parameter space. The binning used and the number of SEDs in each bin are shown in Table 4. Two of the most extreme bins had no sources in them; for these bins we chose to use SEDs randomly extracted from the two nearest bins (one with the same L_x and one with the same N_H). The binning choice was also driven by the binning we adopted to obtain the number of expected AGN (see Sec 3) and we used the same binning edges, although we merged some of the less populated bins.

Throughout this paper, we assumed a bolometric correction of $k_{\text{bol}} = L_{\text{bol}}/L_x$, eq 1, from Lusso et al. (2012).

$$\log(L_x/L_\odot) = 0.230x + 0.050x^2 + 0.001x^3 + 1.256$$

with $x = \log(L_{\text{bol}}/L_\odot) - 12$

(1)

In this work, we will refer to the predicted AGN in the simulated redshift range $0 < z < 10$ with $20 \leq \log(N_H/\text{cm}^{-2}) < 26$ and $42 \leq \log(L_x/\text{erg/s}) < 48.2$ ($43 \leq \log(L_{\text{bol}}/\text{erg/s}) < 49.7$), thus covering the whole simulated parameter space, as all the AGN. The naming scheme used for referring to AGN with different properties (namely, N_H and luminosity) is reported in Table 3.

6 PREDICTIONS OF IR SOURCE COUNTS

In Sec 3, we predicted the number of expected AGN (per deg^2) as function of z , L_x and N_H . For each z , L_x and N_H bin we randomly extracted 20 template SEDs, from the corresponding bins in Section 5), and assigned them a weight equal to one-twentieth of the total number of predicted AGN. We measured their total (AGN+host) flux densities in the SMI $34\mu\text{m}$ and B-BOP $70\mu\text{m}$, $200\mu\text{m}$ $350\mu\text{m}$ bands (for *SPICA*) and in the six *OSS* bands (for *OST*). We obtained the number sources detectable at 5σ by summing the weights of the sources with flux larger than the survey sensitivity.

Table 4 Number of binned sources, using the compilation of Lanzuisi et al. (2017) as explained in section 5, in each 2-10 keV rest-frame luminosity L_x and amount of obscuration N_H bin. The used N_H bins are: $20 \leq \log(N_H/\text{cm}^{-2}) \leq 22$; $22 < \log(N_H/\text{cm}^{-2}) \leq 23$; $23 < \log(N_H/\text{cm}^{-2}) \leq 24.18$; $24.18 < \log(N_H/\text{cm}^{-2}) \leq 26$. We refer to the sources in the first two L_x bins as low-luminosity ones and to those in the other two as high-luminosity ones. We consider the sources in the first N_H bin as unobscured AGN, those in the second and third bins as obscured and those in the last N_H bin as CT-AGN. For the two low-luminosity CT bins we had no SED available and we chose to use the SED of the nearest bins.

$\log(N_H/\text{cm}^{-2})$	21.5	22.5	23.5	24.5
$42.0 \leq \log(L_x/\text{erg/s}) < 42.9$	21	14	5	-
$42.9 \leq \log(L_x/\text{erg/s}) < 43.5$	43	54	10	-
$43.5 \leq \log(L_x/\text{erg/s}) < 44.2$	48	76	53	3
$44.2 \leq \log(L_x/\text{erg/s}) < 48.2$	23	25	38	6

In case of detection, we also differentiated if we were primarily detecting either the AGN or the host galaxy, on the basis of which of the two components had the highest flux in the band we were considering. In the case of a *SPICA* SMI $34\mu\text{m}$ detection, we considered whether we could also detect the source with the SMI-LR mode (as the SMI-CAM photometric channel operates at the same time with the SMI-LR spectroscopic channel). We used, as SMI-LR sensitivity, a mean value between the sensitivities at the boundaries of the SMI-LR wavelength range. For *OST* *OSS* we always considered both the detection in photometric mode ($R=4$) and in spectroscopic mode ($R=300$).

We iterated this whole process forty times (different SEDs were extracted at each iteration), each time obtaining an estimate for the number of detected sources, AGN/torus and spectroscopic detection. We computed the median of these numbers and used the 16th-84th percentiles as uncertainties. Extracting 20 SEDs for each bin (and not one SED for each of the expected detection in that bin) allowed us to save computational time while maintaining a result resolution of 5%. We chose to iterate the process forty times to have good statistical significance of the median and of the 16th-84th percentiles.

In conclusion, from the IR simulations, we derive, for each bin, the percentage of sources photometrically detected, spectroscopically detected, and for which it is the AGN emission to be primarily detected. From the X-ray simulations, we obtained the total number of AGN in each bin, as well as the fraction of sources photometrically detected and spectroscopically characterized using *Athena WFI*. In each bin, we computed the fraction of sources detected both in the IR and in the X-rays, as the minimum between the detection fraction in the two wavelength ranges. Although, in some bins, there may

be sources detected in the IR and not in the X-rays, alongside those detected in the X-rays and not in the IR, usually the detection fraction in one of the wavelength ranges is much higher than the other, and we can safely consider the minimum as a reliable estimate of the fraction of sources detectable with both instruments.

We summarize the main results of our prediction in Tables 5 and 6. Figures 1, 2, 3, and 4 illustrate the number of photometric detections per deg^2 for *SPICA-SMI* DEEP, *SPICA-SMI* WIDE, *OSS* DEEP, and *OSS* WIDE, respectively. All the figures also report the number of sources detected by *Athena WFI*. The black lines are the total number of AGN expected, the red areas represent those which can be detected using *SPICA* SMI-CAM at $34\mu\text{m}$ (or band 1 of *OSS*), the blue areas those which can be detected in the X-rays by *Athena*, the purple areas are the AGN that can be detected both by *SPICA* SMI-CAM (or *OSS*) and by *Athena*. The red uniform area represents the IR detected sources in which the main component of emission is due to the host-galaxy, while for those represented with the starry red area, the AGN is the main contributor to the detected emission. Figures 5 and 6 have a similar color code, but with the dashed areas representing the sources for which we will have spectroscopic detection in a DEEP survey with a *SPICA* SMI-LR-like instrument or with the *OSS*. Finally, figure 9 (figure 10) shows a comparison of the number of sources that we will be able to detect in a DEEP (WIDE) survey with *SPICA* and *OST*, with the color code indicating the expected number of photometric bands in which we may detect the source (considering SMI-CAM and B-BOP for *SPICA* and the six *OSS* bands for the *OST*).

6.1 Photometric detections

Considering the DEEP survey we will be able to detect in the IR $\gtrsim 90\%$ of all the AGN, approximately half of them will have photometric detection both in the IR and in the X-ray bands; this synergy will allow us to identify the source as an AGN and will help in placing better constraints to its properties. By a DEEP survey, working jointly, a IR cryogenic observatory (like *SPICA* or *OST*) and *Athena* can detect all the AGN up to $z = 4$, *de-facto* completely covering the “cosmic noon” and bringing important insights about the AGN density evolution at higher redshifts. At the same time, the bulk of the intrinsic X-ray emission is produced at the “knee” of the luminosity function ($L_x \sim 5 \times 10^{44}$ erg/s for $z \sim 1 - 4$, Aird et al. 2010) and these sources will be easily detected and characterized by *Athena* up to $z \sim 3$. Moreover, *Athena* will be of paramount importance to identify as AGN even sources for which the IR torus emission is diluted and hidden behind the more powerful host-galaxy contribution. The combined use of data in both bands will be fundamental in obtaining a more robust estimate

Table 5 Percentage of all the AGN, CT-AGN, CT-AGN at $z \leq 4$, and CT-AGN at $z \leq 2$ photometrically detected with various configurations of instruments and surveys. SMI refers to *SPICA SMI-CAM*. OSS refers to the sources detected with the *OST OSS* in photometric mode (R=4) in at least one of the bands, while the number within the parenthesis to those with detection in all the six *OSS* bands. SMI AGN and OSS AGN refer to the direct detection of AGN emission (thus the cases where the AGN is more luminous than the host-galaxy in the considered band) for respectively the *SPICA SMI-CAM* and the *OST OSS* instruments. WFI refers to the source photometrically detected by *Athena WFI*, while SMI+WFI (OSS+WFI) refers to the sources with both *SMI-CAM (OSS)* detection and *Athena* photometric detection.

	Survey	SMI %	SMI AGN %	OSS %	OSS AGN %	WFI %	SMI+WFI %	OSS+WFI %
All AGN	DEEP	88 ± 3	22^{+8}_{-6}	95^{+1}_{-2} (84 \pm 3)	21^{+6}_{-7} (< 1)	52	51	51 (50)
	WIDE	72 ± 5	16 ± 6	83^{+3}_{-4} (60 \pm 6)	15 ± 5 (< 1)	29	27	28 (25)
CT-AGN	DEEP	87 ± 3		96 ± 1 (84 \pm 4)		20	20	20 (20)
	WIDE	71 ± 5		84 ± 3 (59 \pm 6)		4	4	4 (4)
CT $z \leq 4$	DEEP	94 ± 2		99 ± 1 (89 \pm 4)		22	22	22 (22)
	WIDE	80 ± 5		93^{+2}_{-3} (68 \pm 7)		5	5	5 (5)
CT $z \leq 2$	DEEP	99 ± 1		99 ± 1 (92 \pm 3)		28	28	28 (28)
	WIDE	84^{+5}_{-4}		98 ± 1 (78 \pm 6)		6	6	6 (6)

Table 6 Percentage of all the AGN, CT-AGN, CT-AGN at $z \leq 4$, and CT-AGN at $z \leq 2$ spectroscopically detected with various configurations of instruments and surveys. SMISp refers to *SPICA SMI-LR*. OSSSp refers to the sources detected with the *OST OSS* in spectroscopic mode (R=300) in at least one of the bands, while the number within the parenthesis to those with detection in all the six *OSS* bands. WFISp refers to the source spectroscopically detected by *Athena WFI*, while SMISp+WFI (OSSSp+WFI) refers to the sources with both *SMI-LR (OSS)* spectroscopic detection and *Athena* photometric detection.

	Survey	SMISp %	OSSSp %	WFISp %	SMISp+WFI %	OSSSp+WFI %
All AGN	DEEP	38 ± 6	81 ± 3 (55 \pm 6)	20	28	47 (38)
	WIDE	18 ± 5	53 ± 5 (20 \pm 5)	6	10	22 (11)
CT-AGN	DEEP	37 ± 6	82^{+3}_{-2} (55 \pm 6)	1	16	20 (18)
	WIDE	17 ± 5	54 ± 6 (22 \pm 5)	<1	3	4 (4)
CT $z \leq 4$	DEEP	43^{+8}_{-7}	92^{+6}_{-7} (63 \pm 7)	2	18	22 (21)
	WIDE	18 ± 5	63^{+7}_{-6} (26 \pm 7)	<1	4	5 (4)
CT $z \leq 2$	DEEP	55^{+9}_{-7}	98 ± 1 (75 \pm 7)	2	22	28 (27)
	WIDE	20 ± 6	77^{+5}_{-6} (34 \pm 8)	<1	5	6 (6)

of the AGN bolometric power. In particular, on the one hand, obtaining the AGN bolometric power with only X-ray data requires to assume an AGN X-ray bolometric correction, thus introducing large uncertainties in the derived values, due to our limited knowledge of the $k_{bol} - L_{bol}$ relation (Hopkins et al. 2007; Lusso et al. 2010, 2012; Duras et al. 2020). On the other hand, using only an instrument similar to *SPICA SMI-CAM*, it will not be possible to distinguish between the AGN and the galaxy emission, as *SPICA SMI-CAM* can measure only the total (AGN+host-galaxy) emission.

Considering a WIDE survey, the fractions of sources that will be detected are lower of $\sim 20\%$ with respect to the DEEP survey, but the (at least) fifteen times wider area will allow us to detect at least 10 times more sources. Similarly to the DEEP, the WIDE survey may allow us to perfectly cover the “cosmic noon” with the advantage of a ten-folded statistic (at least). In a WIDE survey, *Athena WFI* may lose a significant fraction of the most obscured AGN or of the low-luminosity ones. However, a *SPICA*-like (or *OST*-like) mission should be able to effectively recover these sources, although we may need an effective way (e.g. *SPICA-SAFARI* or *OST* spectroscopic follow-up, see sec 6.2 and 6.3) to characterize their AGN properties (intrinsic bolometric luminosity and amount of obscuration). For this shallower survey it is still valid our statement that, for most of the detected sources at $\log(N_{\text{H}}/\text{cm}^{-2}) \leq 23$, $\log(L_{\text{x}}/\text{erg/s}) > 43$ and $z \leq 3$ (and higher z for higher L_{x}), *Athena* detections would provide essential evidence of the AGN nature of those sources.

Focusing on *OST*, the six spectrophotometric bands of *OSS* will give us a good coverage at and beyond the “cosmic noon” even in photometric mode. Moreover, thanks to the capabilities of *OSS*, for the majority of the sources detected by *Athena*, we will have six photometric points (and maybe up to six spectra) in the mid- and far-IR.

6.2 Spectroscopic detections

We estimated that, exploiting *Athena WFI* spectra, we will be able to recover the intrinsic AGN luminosity and obscuration (with 30% uncertainties) for $\approx 20\%$ of all the AGN and up to $\approx 50\%$ of the high-luminosity ones. Even low-count X-ray spectra will allow reliable N_{H} characterization, as for obscured and CT-AGN, we expect simple spectra with a flat continuum and a possibly strong iron $K\alpha$ line. Furthermore, an X-ray detection at a level of $\sim 2 \times 10^{-16}$ erg cm $^{-2}$ s $^{-1}$ in the 2 – 10 keV band at $z \geq 1$, or simply any detection at $z \gtrsim 2.3$ in the DEEP survey, imply X-ray luminosities $\gtrsim 10^{42}$ erg/s in that band, revealing almost unequivocally an AGN.

Considering a *SMI-LR*-like instrument (figure 5), we find that with a DEEP survey we will be able to detect spectroscopically more than half of all the AGN within

$z = 2$. For the high-luminosity objects, we expect to have some spectroscopic detections even up to $z \approx 4$. Considering a WIDE survey, the fraction of AGN that we should be able to detect spectroscopically drops considerably. For the sources for which a fast low-resolution spectroscopic characterization will not be possible, we may make use of photometric data (such as those produced by *SMI-CAM*- or *B-BOP*-like instruments) and of SED-fitting technique (see section 6.3) to disentangle the AGN and SF emissions.

Considering the *OSS*, we find that, with a DEEP survey, we will be able to detect spectroscopically between $55 \pm 6\%$ and $81 \pm 3\%$ of all the AGN, depending on the *OSS* band considered. Thus, we will have $R=300$ full IR spectra for more than half of all the predicted AGN at $z \leq 10$. For the high-luminosity objects, we expect to have some spectroscopic detections even up to $z > 4$. In the WIDE survey, the fraction of AGN that we should be able to detect spectroscopically lowers of $\approx 30\%$ with respect to the DEEP survey.

Several important galaxy and AGN mid-IR spectral features fall in the wavelength range of a *SPICA SMI-LR*-like (or *OSS*) instrument and can be of fundamental importance in both characterizing AGN and host-galaxy properties and measuring the redshift. In particular, we expect AGN-related high-ionization lines¹⁰, such as [Ne v]14.3 μm and 24.3 μm , [O iv]25.9 μm (e.g. Tommasini et al. 2008, 2010; Feltre et al. 2016), but also SF-related Polycyclic Aromatic Hydrocarbons (PAHs) features (e.g. Leger et al. 1989) and the 9.7 μm and 18 μm silicate features. The 9.7 μm feature is typically associated with type 1 AGN when observed in emission and with type 2 when it is in absorption (with the few notable exceptions of the type 2 quasars of Sturm et al. 2006; Teplitz et al. 2006, which show the silicate feature in emission); moreover, more obscured AGN usually have deeper silicate features, with depths that correlate with the X-ray derived N_{H} (e.g. Wu et al. 2009; Shi et al. 2006, but see also Goulding et al. 2012).

Considering the AGN spectroscopic properties, the most evident mid-IR AGN lines ([Ne v]14.3 μm and 24.3 μm , [O iv]25.9 μm) are redshifted out of a *SMI-LR* spectral range at modest redshift ($z < 1$), and would require follow-ups with either *SPICA-SAFARI* or *SMI-MR* to be properly detected. In particular, exploiting 5 hours pointed observations with a *SPICA SAFARI*-like instruments, for low-luminosity (high-luminosity) unobscured AGN the [Ne v]24.3 μm and [O iv]25.9 μm lines can be studied up to $z \approx 1$ and $z \approx 2$ ($z \approx 3$ and $z \approx 4$). The same lines can be detected up to $z \approx 1.5$ and $z \approx 2.5$ ($z \approx 4$ and $z \geq 4$) for low-luminosity (high-luminosity) obscured AGN (Spinoglio et al. 2017, 2021). For the majority of the sources, the brightest and most

¹⁰The high-ionization potential assures us that the emission can not be associated with star-formation activity and has to be, instead, linked to AGN activity




Figure 1. Number of AGN expected per deg^2 and per $\Delta z = 1$ for a *SPICA*-like DEEP survey. The black lines are the total number of expected AGN, the red areas represent those which can be detected with the *SPICA* SMI-CAM at 34 μm , the blue areas those which can be detected in the X-rays with the *Athena* WFI, the purple areas are the AGN that will be detected both by *SPICA* SMI-CAM and *Athena*. The red uniform area represents the sources detected by *SPICA* in which the main component is due to the host-galaxy emission, while for those represented with the starry red area the AGN is the main contributor to the detected emission. The columns refer to AGN with different amount of obscuration ($20 \leq \log(N_{\text{H}}/\text{cm}^{-2}) \leq 21$; $22 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 23$; $24.18 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 25$, from left to right), the rows to different AGN luminosity ($42.0 \leq \log(L_{\text{x}}/\text{erg/s}) < 42.3$, $42.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 43.2$, $43.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 44.2$, $44.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 45.2$, from top to bottom). For forty times we extracted 20 SED for each bin, measured the flux in each *SPICA* band and compared them with the 5σ sensitivities to compute the number of detectable sources. The median of these forty values are the numbers used in this figure, while the 84th – 16th percentiles are used as uncertainties (not reported in the figure).




Figure 2. Number of AGN expected per deg^2 and per $\Delta z = 1$ for a *SPICA*-like WIDE survey. The lines and areas are coded as in Figure 1.




Figure 3. Number of AGN expected per deg^2 and per $\Delta z = 1$ for a *OST OSS DEEP* survey. The black lines are the total number of expected AGN, the red areas represent those which can be detected in the band 1 by *OST OSS*, the blue areas those which can be detected in the X-rays with the *Athena WFI*, the purple areas are the AGN that will be detected both by *OSS* and *Athena*. The red uniform area represents the sources detected by *OSS* in which the main component is due to the host-galaxy emission, while for those represented with the starry red area the AGN is the main contributor to the detected emission. The columns refer to AGN with different amount of obscuration ($20 \leq \log(N_{\text{H}}/\text{cm}^{-2}) \leq 21$; $22 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 23$; $24.18 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 25$, from left to right), the rows to different AGN luminosity ($42.0 \leq \log(L_{\text{x}}/\text{erg/s}) < 42.3$, $42.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 43.2$, $43.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 44.2$, $44.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 45.2$, from top to bottom). For forty times we extracted 20 SED for each bin, measured the flux in each *OSS* band and compared them with the 5σ sensitivities to compute the number of detectable sources. The median of these forty values are the numbers used in this figure, while the 84^{th} – 16^{th} percentiles are used as uncertainties (not reported in the figure).




Figure 4. Number of AGN expected per deg^2 and per $\Delta z = 1$ for a *OST OSS WIDE* survey. The lines and areas are coded as in Figure 3.




Figure 5. Number of AGN expected per deg^2 and per $\Delta z = 1$ for a *SPICA*-like DEEP survey. The black lines are the total number of AGN expected, the red areas represent those which can be detected with the *SPICA* SMI-CAM at $34\mu\text{m}$, the blue areas those which can be detected in the X-rays by *Athena* SMI, the purple areas are the AGN that will be detected both by *SPICA* SMI-CAM and *Athena*. The dashed areas represent the sources which could be spectroscopically detected with SMI-LR (5σ detection with a resolution $R = 50 - 120$ in the $17 - 36\mu\text{m}$ wavelength range). The columns refer to AGN with different amount of obscuration ($20 \leq \log(N_{\text{H}}/\text{cm}^{-2}) \leq 21$; $22 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 23$; $24.18 < \log(N_{\text{H}}/\text{cm}^{-2}) \leq 25$, from left to right), the rows to different AGN luminosity ($42.0 \leq \log(L_{\text{x}}/\text{erg/s}) < 42.5$, $42.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 43.2$, $43.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 44.2$, $44.9 \leq \log(L_{\text{x}}/\text{erg/s}) < 45.2$, from top to bottom).




Figure 6. Number of AGN expected per deg^2 and per $\Delta z = 1$ for an *OSS* DEEP survey. The lines and areas are coded as in Figure 5, but with the red areas referring to the band 1 of the *OST* *OSS* instrument.

recognizable features may be due to the galaxy emission, in particular the dust continuum and the PAH emission lines (e.g. Lutz et al. 1998; Hollenbach & Tielens 1999; Yan et al. 2007; Fadda et al. 2010; Spinoglio et al. 2021). However, for most cases, we expect the $9.7\mu\text{m}$ silicate feature, to be easily recognizable. The silicate feature can be helpful in estimating the AGN amount of obscuration (e.g. Vignali et al. 2011; La Caria et al. 2019) and can be entirely within a SMI-LR-like instrument wavelength range up to $z \approx 3$. This is particularly important for the CT-AGN, for which, with the exception of the most luminous ones ($L_x \approx 10^{44}$ erg/s for $z < 3$), it will be difficult to obtain the AGN properties using only X-ray spectroscopy. Thus, the analysis of both X-ray and IR spectroscopic data will play a fundamental role in discerning their characteristics.

One of the advantages of using a *SPICA SMI*-like (or *OSS*-like) instrument is that it can provide, at the same time, photometry and spectroscopy, and it can quickly produce low-resolution spectroscopic data for a large number of sources. X-ray spectral analysis is, so far, the best way to characterize the AGN (e.g. to determine the intrinsic luminosity), but it requires a redshift determination. Fortunately, there is a good prospect that redshifts can be determined solely from X-ray spectra (e.g. Simmonds et al. 2018; Peca et al. 2021). We simulated mid-IR spectra to evaluate the possibility of measuring the redshift from a *SMI-LR*-like spectra. In particular, we used a code that exploits the most prominent PAH features to recover the redshift. This implies that it is most suited for sources with sufficient star-formation to produce strong PAH features and low- or moderate-luminosity AGN, as the equivalent width of the PAH features is greatly reduced when the AGN luminosity is high (Voit 1992; Genzel et al. 1998; O’Dowd et al. 2009). We chose ten sources among the ~ 500 of our sample and use their SEDs as spectral templates. The ten sources were chosen so that in our sample there were un-obscured, obscured and CT-AGN, high- and low-luminosity sources, AGN- and host-dominated sources, with emission and absorption $9.7\mu\text{m}$ features. For each template, we simulated sixteen spectra in the redshift range $0.5 \leq z \leq 6$ (as we can see from Fig 5, at $z > 6$ the fraction of spectroscopically detected sources is very small). The simulated spectra were created accordingly to the latest *SMI* specifications. We used a spectral resolution of $R = 50$ at $17\mu\text{m}$ and $R = 120$ at $36\mu\text{m}$ and assumed a linear regime between these two values. We used the LR continuum sensitivity (see section 2.1.3), rescaled to the Ultradeep survey, as standard deviation of the Gaussian noise that we included in the spectra. Figure 7 shows two of the simulated spectra. Finally, we exploit a modified version of the PAHFIT code (Smith et al. 2007) that uses a power-law continuum, several PAH emission lines and silicate absorption, to fit the spectra and provide the source redshift (Negrello et al.




Figure 7. Simulated SMI-LR spectra for a *SPICA*-like DEEP survey. Left panel is for $\log(L_x/\text{erg/s}) = 43.9$, $\log(N_{\text{H}}/\text{cm}^{-2}) = 23.4$ (i.e. moderate-luminous obscured) AGN at $z = 2$; right panel for $\log(L_x/\text{erg/s}) = 44.3$, $\log(N_{\text{H}}/\text{cm}^{-2}) = 24.1$ (i.e. high-luminous) CT-AGN $z = 2$. We used the Lanzuisi et al. (2017) SEDs (inset in the upper left of the plots, where the red line is the AGN component, the blue one is related to the host-galaxy and the black one is the total AGN+galaxy emission) as spectral templates and added a white noise with amplitude based on the LR continuum sensitivity expected for a DEEP survey. The spectra were sampled with a resolution of $R = 50$ at $17\mu\text{m}$ and $R = 120$ at $36\mu\text{m}$.

in prep). As shown in figure 8, we find that we should be able to effectively recover the redshift in case of moderate and luminous type 2 AGN up to $z \sim 3 - 4$. However, the code currently does not use AGN silicate $9.7\mu\text{m}$ emission feature in finding the z and is prone to misinterpret it as a PAH feature when it is strong; however, the code is still able to recover the redshift in case of faint AGN silicate emission line, as long as the silicate line is not mistakenly interpreted as a PAH feature. In particular, excluding three sources with strong Si $9.7\mu\text{m}$ emission features and one source with very low signal-to-noise ratio, we recovered the redshifts with a median error $|z_{\text{best}} - z_{\text{true}}| / (1 + z_{\text{true}}) = 0.02$. In conclusion, a *SPICA SMI*-like instrument can provide us with low-resolution spectra for more than 6 000 AGN in a DEEP survey and, in the cases of sources with prominent host-galaxy PAH lines and low AGN activity (i.e. unable to destroy these lines), measures of their spectroscopic redshifts. Moreover, it should be possible to obtain an estimate of the AGN obscuration from the depth of the silicate $9.7\mu\text{m}$ absorption feature (Wu et al. 2009; Shi et al. 2006 had shown that there is a correlation between these two quantities), that may provide support to the outcomes of X-ray spectroscopy in terms of obscuration (especially for low-statistics obscured AGN).

Focusing on the *OSS* capabilities, this instrument is able to exploit the above mentioned mid-IR lines, but with the advantages of higher spectral resolution and larger wavelength range coverage than a *SPICA*-like instrument. Lines as the $[\text{Ne V}]14.3\mu\text{m}$ and $24.3\mu\text{m}$, $[\text{O IV}]25.9\mu\text{m}$, which are redshifted out of *SMI-LR* spectral range at modest redshift ($z < 1$), thus requiring




Figure 8. Redshift estimate from simulated *SMI-LR*-like spectra using a modified version of the PAHFIT code (Negrello et al. in prep). Left panel is for $\log(L_x/\text{erg/s}) = 43.9$, $\log(N_H/\text{cm}^{-2}) = 23.4$ moderate-luminous obscured AGN; right panel for $\log(L_x/\text{erg/s}) = 44.3$, $\log(N_H/\text{cm}^{-2}) = 24.1$ high-luminous CT-AGN. We find that we should be able to effectively recover the redshift up to $z \approx 3 - 4$; at $z > 4$, there are too few strong PAH features in the *SMI-LR* wavelength range to allow a proper determination of the redshift.

follow-ups with other instruments, can be detected up to very high-redshift, their detection being a matter of instrument sensitivity and not of limited wavelength range. Fine structure lines of O, C, Ne, S, N, Fe, Ar, and Si are keys to probe neutral and ionized gas phases and can be used to estimate redshift, amount of gas, its ionization, presence of outflows and the contribution of SF and AGN to the emission (Meixner et al. 2019; Mordini et al. 2021). The OSS DEEP survey will allow us to detect the PAH up to $z \approx 4$ for galaxies with $\text{SFR} \sim 10M_\odot/\text{yr}$ and over $z > 6$ for those with $\text{SFR} \sim 100M_\odot/\text{yr}$ (Meixner et al. 2019). Regarding the estimate of the BHARD, the high sensitivity and large spectral range of the OSS survey will allow the detection of the [Ne v] line (whose flux correlates with the AGN intrinsic emission), as well as of the [O IV] and [Ne II] line, whose line flux ratio can be used as a diagnostic of AGN power (e.g., Genzel et al. 1998; Lutz et al. 2003; Armus et al. 2006, 2007; Mordini et al. 2021). Gruppioni et al. (2016) well calibrated the relations between [Ne v] (or [O IV]) and AGN bolometric emission in samples of local galaxies. Bonato et al. (2019) explored the spectroscopic survey capabilities of the OSS and find out that, in a DEEP survey, it can detect emission lines associated to SF up to $z \approx 8.5$ and those of AGN emission, in particular [Ne v] $24.3\mu\text{m}$ and [O IV] $25.9\mu\text{m}$, up to $z \approx 3$ and $z \approx 5$, respectively.

6.3 SED fitting

Working jointly, an IR cryogenic observatory, like *SPICA* or *OST*, and *Athena* can detect a huge fraction of the the AGN up to $z = 10$. The combined used of data from both wavelength bands will be fundamental, as with only IR photometric detections, it will not be possible to distinguish between the AGN and the galaxy emission, as *SPICA SMI-CAM* or *OSS* can measure only the total

(AGN+host-galaxy) emission, this in the lucky case of having already identified the source as an AGN. In this regard, a spectrometer similar to *SPICA SMI-LR* or *OSS* may surely help, but we will not be able to spectroscopically detect all the AGN, as the spectroscopic recovery fraction decreases rapidly for $z > 2$. The additional use of other photometric data (such as those of B-BOP-like instruments, or from all the *OSS* bands), along with SED decomposition technique, could be used to overcome this limitation. Even in photometric mode, having six far-IR detections of *OST* will be invaluable in disentangling the AGN and host-galaxy contribution: indeed, these bands cover both the regime where the torus has its peak emission and the regime where the SF is the main contributor to the SED. But we expect that the real game-changer in obscured AGN studies will come from a multi-wavelength approach.

The use of *Athena* data, optical and near-IR photometric datapoints (in fields where these are already available, such as COSMOS and the Euclid Deep Fields), mid-IR and far-IR photometric instruments (similar to those designed for *SPICA*) can provide a never-reached-before coverage of all the AGN and galaxy emission up to high redshift. In addition, with follow-up pointed spectroscopic observations of these targets (photometrically detected, but without a clear separation between galaxy and AGN components) we will be able to distinguish the AGN and host-galaxy emissions using fine-structure lines and PAH features up to $z \sim 4$ (Spinoglio et al. 2021). Therefore, we expect to be able to improve our knowledge of the AGN $k_{bol} - L_{bol}$ relation, similarly to what was done in Lusso et al. (2012), but with a larger sample and to higher redshift. This should enable us to extend the $k_{bol} - L_{bol}$ dynamical range and, having more sources, to reduce its uncertainties.

In figure 9, we show the comparison of the total number of AGN detected by a *SPICA*-like observatory and by the *OST OSS* for a DEEP survey. The color code indicates the number of bands for which we can have photometric detection, considering *SMI-CAM* and the three B-BOP channel for *SPICA* and the six *OSS* bands for *OST*. For $z < 2$ both mission are able to detect almost all the sources in at least one band. While at $z > 2$ *SPICA* begins to lose a fraction of the sources, *OSS* continues to detect them all up to $z \approx 3.5$.

Moreover, having a sufficient number of photometric points, would allow us to break the SF-AGN emission degeneracy and to estimate the AGN properties using SED-fitting decomposition techniques (jointly with additional instruments at different wavelengths) when spectra are not available. Although X-ray photometry and/or low-resolution spectroscopic data may not be used in SED fitting (but see X-CIGALE, Yang et al. 2020), X-ray information could be crucial to select the templates to fit in the IR.

As a quick test for this statement, we simulated




Figure 9. Distribution of the number of AGN expected per deg^2 for a *SPICA*-like (*top panel*) and for the *OST* *OSS* (*bottom panel*) from a DEEP survey. The color code indicates our predictions for the number of photometric detections. For the *SPICA* panel we took into consideration the *SMI-CAM* and the three *B-BOP* channel, while the dashed area represents the sources which can be spectroscopically detected with a *SMI-LR*-like instrument (5σ detection with a resolution $R=50-120$ in the $17-36\mu\text{m}$ wavelength range). For the *OST* panel, we consider the six *OSS* bands in photometric mode ($R=4$), while the dashed line indicates the sources for which we can have $R=300$ spectra in at least one of the *OSS* bands.




Figure 10. Distribution of the number of AGN expected per deg^2 for a *SPICA*-like (*top panel*) and for the *OST* *OSS* (*bottom panel*) from a WIDE survey. The lines and areas are coded as in Figure 9.




Figure 11. SED-fitting probability distribution functions (PDFs) for AGN 392925 ($\log(L_x/\text{erg/s}) = 45.0$, $\log(N_H/\text{cm}^{-2}) = 24.1$) at $z = 1.55$. *Left panel*: AGN bolometric luminosity; *right panel*: SFR in the last 0.1 Gyr. The blue lines are obtained from the SED-fitting using 33 photometric bands from optical to far-IR (in particular, from IR telescope *Spitzer* and *Herschel*). The red lines are the PDF obtained using the same 33 photometric bands plus the four simulated photometric observations of *SPICA SMI-CAM*, *B-BOP1*, *B-BOP2* and *B-BOP3*. The use of the additional *SPICA* photometric points provides better constraints on the AGN bolometric power and, overall, allows us to properly disentangle the AGN and the SF emission.

SPICA-like observations and used *sed3fit* to recover the stellar mass, SFR and AGN bolometric luminosity. In particular, we selected a random sample of seventy sources among the ≈ 550 of the [Lanzuisi et al. \(2017\)](#) compilation that we used in this work and exploit *sed3fit* with up to 33 [Laigle et al. \(2016\)](#) photometric data (from optical to sub-mm including *Spitzer* and *Herschel*, although not all the photometric points were available for all the sources) to obtain the AGN, host-galaxy and total SED. We used the total SED to estimate the *SPICA SMI* and *B-BOP* fluxes, added these four photometric points to the previous 33 and used all of them in new runs of *sed3fit*. We focused on the width of the 84th – 16th percentile of the probability distribution functions (PDF) from *sed3fit* to check for the improvement provided by the use of *SPICA*-like data. We found that for, respectively, 50 and 45 of the 70 sources investigated we have narrower AGN bolometric luminosity and SFR PDFs. The median SF relative error (i.e. $0.5 \times (SFR_{84^{th}} - SFR_{16^{th}}) / SFR_{best}$) improves from 11% to 1% and that of the AGN L_{bol} from 61% to 29%. The improvements in the stellar mass are smaller, since *sed3fit* estimates this parameter exploiting the optical wavelength band, therefore we get only a minor advantage in using more IR data. The major improvements come from the cases where the AGN-SF degeneracies are stronger; as we can see from Figure 11, the additional mid- and far-IR points provided by a *SPICA*-like observatory help in disentangling the two contributions and allow us to obtain better estimates of both the host-galaxy and AGN parameters (e.g. stellar mass, SFR, AGN L_{bol}).

6.4 CT-AGN

The search for CT-AGN is of fundamental importance in view of the still open questions of the missing sources responsible for the XRB and the galaxy co-evolution paradigm. About the former issue, X-ray background synthesis models predict a putative population of CT-AGN challenging to detect with current facilities (Gilli et al. 2007; Gilli 2013; Harrison et al. 2016). In addition, the black-hole-galaxy co-evolution paradigm states that the scarcely known first phases of BH (and galaxy) growth should be associated with very obscured AGN activity and obscured SF, therefore these elusive AGN are of the foremost importance to search for (e.g. Silk & Rees 1998; Di Matteo et al. 2005; Lamastra et al. 2013).

The adopted model for the XRB predicts that low-luminosity CT-AGN form the bulk of the very obscured AGN population. However, due to their large obscuration, it is difficult to detect them with X-ray instruments, while a *SPICA*-like (or *OST*) observatory should be able to effectively sample them with a detection fraction as high as 90% in the DEEP survey and 70% in the WIDE survey.

As we can see from Figure 12, low-luminosity obscured AGN are difficult to detect even with the planned *Athena* facility. In fact, we estimated that *Athena* will be able to detect (light blue) more than half of all the high-luminosity CT-AGN, while the major part of the CT-AGN will be missed ($\sim 20\%$ detection fraction), being low-luminosity sources (~ 4260 of the 5200 CT-AGN per deg^2 have $\log(L_x/\text{erg s}^{-1}) < 43.5$). The characterization of these sources with *Athena WFI* (dotted area) will be challenging due to their low statistics: we may be able to recover L_x and N_H only for $\sim 6\%$ of the high-luminosity ones and $< 1\%$ for the low-luminosity. Sufficiently powerful IR surveys will be capable of detecting the reprocessed emission due to accretion at IR wavelengths. For example, a *SPICA*-like DEEP survey is capable of detecting (red areas) $\approx 90\%$ of the CT-AGN and provide low-resolution spectra (shaded areas) for one third of them. The even more powerful *OST* will allow us to detect photometrically ($R=4$) almost all the CT-AGN and to have $R=300$ spectra for at least half of them. Spectroscopic follow-up with other IR instruments will be fundamental in identifying CT-AGN with photometric detection but without a clear detection with *SMI-LR*-like or *OSS* instruments. Low-luminosity (high-luminosity) CT-AGN can be spectroscopically identified with a *SAFARI*-like instrument up to $z \sim 1.5$ ($z \sim 2.5$) and $z \sim 2.5$ ($z > 4$) via the $[\text{Ne v}]24.3\mu\text{m}$ and the $[\text{O IV}]25.9\mu\text{m}$ lines with 5 hours observations (Spinoglio et al. 2021). Longer exposures (>10 hours) will allow to identify CT-AGN beyond $z \sim 4$. Despite the fact that the X-ray detection of these CT objects will be difficult, the *Athena* contribution to their study remains essential. In fact, only X-ray spectra (for individual sources

or at least stacked X-ray data in case of no detection) can confirm the AGN nature of these sources (i.e., a flat X-ray continuum plus a strong iron line are clear indicators of heavy obscuration).

For the most luminous ones, the combined use of *Athena* X-ray spectroscopy, mid-, and far-IR spectrophotometry (with *SMI-CAM*- and *B-BOP*-like instruments or with the *OSS*) will provide never achieved before insights of the intrinsic AGN emission and bolometric power and opportunities to sample them at redshift higher than the cosmic noon. As shown in Figure 13, at present, there are tensions between BHAD traced by X-rays, and those obtained by simulations, especially at $z > 2$. At $z \sim 4$, the BHAD expected from simulations (Aird et al. 2015; Vito et al. 2018) are ~ 5 times higher than those computed via X-ray surveys, while actual IR-surveys do not go beyond $z \sim 3$. Different arguments, mostly driven by observations in the IR (Gruppioni et al. 2016; Bisigello et al. 2020), predict a higher number of deeply obscured AGN, as for now undetected in the X-ray, that may alleviate this tension.

Exploiting all the capabilities of new generation IR cryogenic observatories (i.e. fast LR spectrophotometry to detect and characterize low-redshift CT-AGN, spectroscopic follow-ups to identify those with only photometric detection and multi-wavelength SED-fitting exploiting already observed deep fields to separate the AGN and host-galaxy emission) will allow us to discover the deeply obscured CT-AGN population up to very high redshift ($z > 4$). The presence of such an obscured population does not violate the limit imposed by the spectral energy density of the XRB, since these sources will provide an almost negligible contribution in the X-ray band (Comastri et al. 2015). The updated picture, in terms of obscured accretion over cosmic time, that the IR missions discussed in this paper will lay out is fundamental in providing inputs to the simulations, which are still affected by considerable uncertainties and depend strongly on the adopted assumptions (e.g., González et al. 2011; Jaacks et al. 2012; Sijacki et al. 2015; Shankar et al. 2013; Volonteri et al. 2016; Thomas et al. 2019).

7 CONCLUSIONS

In this work we have investigated the capabilities of IR deep and wide surveys to detect and characterize AGN, as well as the advantages resulting by the synergies with the future X-ray observatory *Athena*. We used XRB synthesis models to predict the total number of AGN as a function of intrinsic X-ray luminosity, amount of obscuration and redshift. A sample of more than 500 AGN from the COSMOS field with both X-ray spectra and optical-to-FIR SED-fitting has been used to investigate the AGN mid- and far-IR emission and to predict the fraction of AGN that will be detected by instrument similar to *SPICA SMI*, *B-BOP* and the *OST OSS*. We




Figure 12. Number of CT AGN expected per deg^2 and per $\Delta z = 1$ from a DEEP survey with *Athena* (left panels), with a *SPICA*-like observatory (middle panels), and with the *OST OSS* (right panels). The black lines are the total number of expected CT-AGN, the blue areas represent those which can be detected in the X-ray by *Athena WFI* and the dotted areas those for which we will be able to recover the L_x and N_H (with 30% uncertainties) using *Athena WFI* spectroscopy; the red areas represent those which can be detected in the IR with the *SPICA SMI-CAM* at $34\mu\text{m}$. The light red areas indicate that the host galaxy has the largest contribution to the flux in the SMI-CAM detection, while the dark red areas that the AGN is the dominant component. The dashed areas indicate that we will be able to have also mid-IR spectroscopic 5σ detection with *SPICA SMI-LR* with a resolution $R = 50 - 120$ in the $17 - 36\mu\text{m}$ wavelength range. The light orange areas are the sources for which we expect detection in the OSS band 1 ($25 - 44\mu\text{m}$) in photometric mode ($R=4$), the dark orange area those for which the AGN is the dominant component of the detected emission. The dashed areas refer to AGN for which we will have enough flux to have low-resolution ($R=300$) spectra in OSS band 1. For the *OST*, we showed only the prediction for the band 1 of the OSS, but the instrument has five other bands (up to $588\mu\text{m}$), for which we predict a larger number of AGN detected both in photometry and in spectroscopy. The upper row refers to AGN with luminosity $42.0 \leq \log(L_x/\text{erg/s}) < 42.5$, while the lower to those with $43.9 \leq \log(L_x/\text{erg/s}) < 44.2$.




Figure 13. Redshift evolution of the BHAD adapted from Vito et al. (2018). The blue area is a theoretical BHAD curve obtained from Volonteri et al. (2016); Sijacki et al. (2015); Shankar et al. (2013). The red areas are the X-ray-derived BHADs, obtained from observations, from Aird et al. (2015); Ueda et al. (2014); Vito et al. (2014). The yellow area is the IR-derived BHAD from Delvecchio et al. (2014). *SPICA*-like or *OST* missions will allow to sample the BHAD at higher z than actual IR survey and will be able to detect very obscured CT-AGN, missed by X-ray surveys, that should contribute to the total BHAD.

compared the estimated total number of AGN with the fraction of those that will be detected by *Athena*, by a *SPICA*-like observatory and by the *OST*, with the goal of enlightening the synergies between X-ray and IR surveys in terms of detection and characterization of AGN (in particular, of the most obscured ones) up to high redshift (i.e. $z \sim 10$) and down to low luminosity (i.e. $\log(L_x/\text{erg/s}) \sim 42$). In the IR bands, we also differentiated if we were primary detecting either the AGN or the host-galaxy emission. Moreover, we investigated if we were able to spectroscopically identify and characterize the AGN emission with *SMI-LR*-like instruments or exploiting the full $R=300$ spectral resolution of the OSS. In particular, we put emphasis on the elusive CT-AGN, a large fraction of those being missed by the current X-ray surveys, and that will require deep IR surveys for their detection and *Athena* spectra for their characterization. Our main results are the following:

1. An *OST WIDE* survey will be able to photometrically detect $\sim 60\%$ of all the AGN in all the OSS bands and more than ~ 80 in at least one of the bands, while the DEEP survey will detect at least

- 95% of all the AGN at $z \leq 10$ in at least one of the OSS bands and more than $\sim 80\%$ in all of them.
2. Considering an *Athena* and *SPICA*-like DEEP survey, we estimate that we can have both IR and X-ray detections for half of all the AGN and for more than the 80% of the high-luminosity ($\log(L_x/\text{erg/s}) \geq 43.5$) AGN; for these sources just the *Athena* detection will assure us of the presence of AGN activity, even when the mid-IR torus contribution is diluted into the host-galaxy emission. The *Athena* WFI spectral capabilities should also provide X-ray spectra, hence a reliable way to characterize the AGN, for thousands of AGN. This will allow us to estimate the AGN bolometric power and, having such a large number of sources with data in both the wavelength bands, to improve the AGN $L_{\text{mid-IR}} - L_x$ and $k_{\text{bol}} - L_{\text{bol}}$ relations. Moreover, the use of low-resolution spectra (such as those produced by a SMI-LR-like spectrometer), or alternatively, of mid- and far-IR photometric points (available with SMI-CAM- and B-BOP-like instruments) and SED-fitting decomposition technique, will help in constraining the AGN properties even in case of low-statistic X-ray spectra, like those we expect for very obscured and CT-AGN. However, even for sources with low photon statistics, the detection of a flat continuum and/or a strong iron emission line in the *Athena* WFI spectra will be invaluable clues to identify these sources as very obscured AGN.
 3. The synergies of *Athena* with the *OST* are similar to those with a *SPICA*-like mission, but with the advantages of having up to six photometric point in the far-IR, a higher fraction (with respect to a *SPICA*-like mission) of sources with mid-IR spectra and providing R= 300, 25 – 588 μm spectra for $\sim 50\%$ of all the AGN.
 4. The CT-AGN, for which the X-ray detection and characterization are challenging due to their high amount of obscuration, will be very well sampled by deep IR surveys. A *SPICA*-like DEEP survey should detect 86% of the low-luminosity ($\log(L_x/\text{erg/s}) < 43.5$) CT-AGN at $z \leq 10$, while, *OST* can provide with six mid- and far-IR photometric detections for a similar fraction of the AGN, and at least one band detection for more than $\sim 95\%$ of all the CT AGN at $z < 10$. These kinds of missions would allow us to sample higher redshifts than what we are actually capable of at IR wavelengths, and detecting very obscured CT-AGN missed by X-ray surveys, to provide clues and possibly inputs to BHAD simulations. However, for most of these sources, the major contributor to the flux comes from the host galaxy emission, and the challenge will be to recognise them as AGN. In this context, pointed observations, such as those provided by *SPICA* SAFARI-like or *OSS* follow up, can be decisive in detecting the [Ne v]24.3 μm and the [O IV]25.9 μm lines, which would allow us to unambiguously classify the source as an AGN.
 5. For high-luminosity CT-AGN, we will have X-ray detection with *Athena* up to $z \sim 3$. This will allow us to ascertain the AGN nature of these obscured sources and will provide vital input for the selection of IR SED templates and for their spectroscopic characterization.
 6. The use of mid- and far-IR photometric observations (such as those produced by SMI- and B-BOP-like instruments or by the *OST* OSS in R=4 photometric mode) can provide up to four (or six in the case of the *OSS*) photometric data points, which will be unique to improve the reliability of SED-fitting techniques to estimate the AGN properties up and beyond the cosmic noon.
 7. Mid-IR spectra are very helpful to recognize AGN and to separate their emission from the SF-related one. A *SPICA*-like observatory can provide LR spectra (with the SMI-LR instrument) for the majority of the detected objects at $z \leq 2$, and up to $z \approx 4$ for the most luminous sources, as well as follow-up deep spectroscopy with SAFARI for galaxies up to and beyond $z \sim 4$. On the other hand the *OST* OSS can deliver R= 300, 25 – 588 μm spectra for $\sim 50\%$ and $\sim 80\%$ of all the AGN with, respectively, its WIDE and DEEP surveys, the last one up to and beyond $z \approx 4$. The main spectral features to focus on will be the host-galaxy PAH lines and the AGN 9.7 μm silicate feature. We showed that, for what concerns the PAHs, those can be used to quickly recover the source spectroscopic redshift, without requiring specific higher-resolution follow-ups, while the silicate feature, when it is in absorption, jointly with X-ray spectral information, will allow us to assess and quantify the amount of obscuration. An *Athena* DEEP (WIDE) survey will provide X-ray spectral characterization for $\approx 20\%$ ($\approx 6\%$) of all the AGN and up to $\approx 50\%$ ($\approx 21\%$) of the high-luminosity ones, and, with the help of a multi-wavelengths approach, we will be able to characterize the properties even for X-ray low-statistic sources, such as CT-AGN.
- Mid- and far-IR deep surveys, such as those of a *SPICA*-like mission or of the *OST*, will allow us to fully exploit the potential of *Athena* for the measurement of the full Black-Hole Accretion History, back to an age of the Universe of ~ 1 Gyr. This, along with the Star Formation History, will give us a much better understanding of galaxy evolution as compared to what we know today.

8 ACKNOWLEDGEMENTS

This paper is dedicated to the memory of Bruce Swinyard, who initiated the SPICA project in Europe, but unfortunately died on 22 May 2015 at the age of 52. He was ISO-LWS calibration scientist, *Herschel-SPIRE* instrument scientist, first European PI of *SPICA* and first design lead of SAFARI.

We acknowledge the whole *SPICA* Collaboration Team.

FP, CV, CG, LB, and LS acknowledge financial support by the Agenzia Spaziale Italiana (ASI) under the research contract 2018-31-HH.0. FJC acknowledges financial support from the Spanish Ministry MCIU under project RTI2018-096686-B-C21 (MCIU/AEI/FEDER/UE), cofunded by FEDER funds and from the Agencia Estatal de Investigación, Unidad de Excelencia María de Maeztu, ref. MDM-2017-0765. MPS acknowledges support from the Comunidad de Madrid, Spain, through Atracción de Talento Investigador Grant 2018-T1/TIC-11035 and PID2019-105423GA-I00 (MCIU/AEI/FEDER,UE). AF acknowledges the support from grant PRIN MIUR2017-20173ML3WW_001.

We also thank the *SPICA* Science Study Team appointed by ESA and the *SPICA* Galaxy Evolution Working Group. We should also thank here the Athena Science Study Team, also appointed by ESA, the Athena Science Working Groups Formation and growth of earliest SMBH and Understanding the build-up of SMBH and galaxies, and the Athena Community Office

REFERENCES

- Aird, J., Alexander, D. M., Ballantyne, D. R., et al. 2015, *ApJ*, 815, 66
- Aird, J., Nandra, K., Laird, E. S., et al. 2010, *MNRAS*, 401, 2531
- Ananna, T. T., Treister, E., Urry, C. M., et al. 2019, *ApJ*, 871, 240
- André, P., Hughes, A., Guillet, V., et al. 2019, Probing the cold magnetized Universe with SPICA-POL (B-BOP)
- Armus, L., Bernard-Salas, J., Spoon, H. W. W., et al. 2006, *ApJ*, 640, 204
- Armus, L., Charmandaris, V., Bernard-Salas, J., et al. 2007, *ApJ*, 656, 148
- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17
- Assef, R. J., Stern, D., Kochanek, C. S., et al. 2013, *ApJ*, 772, 26
- Ballantyne, D. R., Draper, A. R., Madsen, K. K., Rigby, J. R., & Treister, E. 2011, *ApJ*, 736, 56
- Barai, P., Gallerani, S., Pallottini, A., et al. 2018, *MNRAS*, 473, 4003
- Barcons, X., Barret, D., Decourchelle, A., et al. 2017, *Astronomische Nachrichten*, 338, 153
- Barret, D., Lam Trong, T., den Herder, J.-W., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10699, *Space Telescopes and Instrumentation 2018: Ultraviolet to Gamma Ray*, 106991G
- Battersby, C., Armus, L., Bergin, E., et al. 2018, *Nature Astronomy*, 2, 596
- Bavdaz, M., Wille, E., Ayre, M., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10699, *Space Telescopes and Instrumentation 2018: Ultraviolet to Gamma Ray*, 106990X
- Berta, S., Lutz, D., Santini, P., et al. 2013, *A&A*, 551, A100
- Bisigello, L., Gruppioni, C., Feltre, A., et al. 2020, arXiv e-prints, arXiv:2011.07074
- Bonato, M., De Zotti, G., Leisawitz, D., et al. 2019, *PASA*, 36, e017
- Brightman, M. & Nandra, K. 2011, *MNRAS*, 413, 1206
- Brightman, M. & Ueda, Y. 2012, *MNRAS*, 423, 702
- Burlon, D., Ajello, M., Greiner, J., et al. 2011, *ApJ*, 728, 58
- Cash, W. 1979, *ApJ*, 228, 939
- Castelló-Mor, N., Carrera, F. J., Alonso-Herrero, A., et al. 2013, *A&A*, 556, A114
- Civano, F., Marchesi, S., Comastri, A., et al. 2016, *ApJ*, 819, 62
- Comastri, A., Gilli, R., Marconi, A., Risaliti, G., & Salvati, M. 2015, *A&A*, 574, L10
- Comastri, A., Ranalli, P., Iwasawa, K., et al. 2011, *A&A*, 526, L9
- Comastri, A., Setti, G., Zamorani, G., & Hasinger, G. 1995, *A&A*, 296, 1
- Del Moro, A., Alexander, D. M., Bauer, F. E., et al. 2017, *Frontiers in Astronomy and Space Sciences*, 4, 67
- Del Moro, A., Mullaney, J. R., Alexander, D. M., et al. 2014, *ApJ*, 786, 16
- Delvecchio, I., Gruppioni, C., Pozzi, F., et al. 2014, *MNRAS*, 439, 2736
- Delvecchio, I., Lutz, D., Berta, S., et al. 2015, *MNRAS*, 449, 373
- Di Matteo, T., Springel, V., & Hernquist, L. 2005, *Nature*, 433, 604
- Dowell, C. D., Hildebrand, R. H., Schleuning, D. A., et al. 1998, *The Astrophysical Journal*, 504, 588
- Duras, F., Bongiorno, A., Ricci, F., et al. 2020, *A&A*, 636, A73
- Elvis, M., Civano, F., Vignali, C., et al. 2009, *ApJS*, 184, 158
- Fadda, D., Yan, L., Lagache, G., et al. 2010, *ApJ*, 719, 425
- Feltre, A., Charlot, S., & Gutkin, J. 2016, *MNRAS*, 456, 3354
- Fernández-Ontiveros, J. A., Spinoglio, L., Pereira-Santaella, M., et al. 2016, *ApJS*, 226, 19

- Ferrarese, L. & Merritt, D. 2000, *ApJ*, 539, L9
- Feruglio, C., Fiore, F., Carniani, S., et al. 2015, *A&A*, 583, A99
- Fiore, F., Feruglio, C., Shankar, F., et al. 2017, *A&A*, 601, A143
- Galitzki, N., Ade, P. A. R., Angilè, F. E., et al. 2014, *Journal of Astronomical Instrumentation*, 03, 1440001
- García-Bernete, I., Ramos Almeida, C., Acosta-Pulido, J. A., et al. 2016, *MNRAS*, 463, 3531
- Gebhardt, K., Bender, R., Bower, G., et al. 2000, *ApJ*, 539, L13
- Genzel, R., Lutz, D., Sturm, E., et al. 1998, *ApJ*, 498, 579
- Georgantopoulos, I., Comastri, A., Vignali, C., et al. 2013, *A&A*, 555, A43
- Giacconi, R., Zirm, A., Wang, J., et al. 2002, *ApJS*, 139, 369
- Gilli, R. 2013, *Mem. Soc. Astron. Italiana*, 84, 647
- Gilli, R., Comastri, A., & Hasinger, G. 2007, *A&A*, 463, 79
- González, V., Labbé, I., Bouwens, R. J., et al. 2011, *ApJ*, 735, L34
- Goulding, A. D., Alexander, D. M., Bauer, F. E., et al. 2012, *ApJ*, 755, 5
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3
- Gruppioni, C., Berta, S., Spinoglio, L., et al. 2016, *MNRAS*, 458, 4297
- Gruppioni, C., Béthermin, M., Loiacono, F., et al. 2020, *A&A*, 643, A8
- Gruppioni, C., Pozzi, F., Rodighiero, G., et al. 2013, *MNRAS*, 432, 23
- Harrison, F. A., Aird, J., Civano, F., et al. 2016, *ApJ*, 831, 185
- Heckman, T. M. & Best, P. N. 2014, *ARA&A*, 52, 589
- Hollenbach, D. J. & Tielens, A. G. G. M. 1999, *Reviews of Modern Physics*, 71, 173
- Hopkins, P. F., Hernquist, L., Cox, T. J., et al. 2006, *ApJS*, 163, 1
- Hopkins, P. F., Richards, G. T., & Hernquist, L. 2007, *ApJ*, 654, 731
- Houck, J. R., Soifer, B. T., Weedman, D., et al. 2005, *ApJ*, 622, L105
- Hughes, D. H., Serjeant, S., Dunlop, J., et al. 1998, *Nature*, 394, 241
- Jaacks, J., Choi, J.-H., Nagamine, K., Thompson, R., & Varghese, S. 2012, *MNRAS*, 420, 1606
- Kaneda, H., Ishihara, D., Oyabu, S., et al. 2017, *PASA*, 34, e059
- Kormendy, J. & Richstone, D. 1995, *ARA&A*, 33, 581
- La Caria, M. M., Vignali, C., Lanzuisi, G., Gruppioni, C., & Pozzi, F. 2019, *MNRAS*, 487, 1662
- Laigle, C., McCracken, H. J., Ilbert, O., et al. 2016, *ApJS*, 224, 24
- Lamastra, A., Menci, N., Fiore, F., et al. 2013, *A&A*, 559, A56
- Lansbury, G. B., Alexander, D. M., Aird, J., et al. 2017a, *ApJ*, 846, 20
- Lansbury, G. B., Stern, D., Aird, J., et al. 2017b, *ApJ*, 836, 99
- Lanzuisi, G., Civano, F., Elvis, M., et al. 2013, *MNRAS*, 431, 978
- Lanzuisi, G., Delvecchio, I., Berta, S., et al. 2017, *A&A*, 602, A123
- Lanzuisi, G., Ranalli, P., Georgantopoulos, I., et al. 2015, *A&A*, 573, A137
- Lapi, A., Pantoni, L., Zanisi, L., et al. 2018, *ApJ*, 857, 22
- Leger, A., D’Hendecourt, L., & Defourneau, D. 1989, *A&A*, 216, 148
- Lehmer, B. D., Xue, Y. Q., Brandt, W. N., et al. 2012, *ApJ*, 752, 46
- Li, H., Dowell, C. D., Kirby, L., Novak, G., & Vaillancourt, J. E. 2008, *Appl. Opt.*, 47, 422
- Luo, B., Bauer, F. E., Brandt, W. N., et al. 2008, *ApJS*, 179, 19
- Lusso, E., Comastri, A., Simmons, B. D., et al. 2012, *Monthly Notices of the Royal Astronomical Society*, 425, 623
- Lusso, E., Comastri, A., Vignali, C., et al. 2010, *A&A*, 512, A34
- Lutz, D., Spoon, H. W. W., Rigopoulou, D., Moorwood, A. F. M., & Genzel, R. 1998, *ApJ*, 505, L103
- Lutz, D., Sturm, E., Genzel, R., et al. 2003, *A&A*, 409, 867
- Madau, P. & Dickinson, M. 2014, *ARA&A*, 52, 415
- Magnelli, B., Popesso, P., Berta, S., et al. 2013, *A&A*, 553, A132
- Marchesi, S., Civano, F., Elvis, M., et al. 2016, *ApJ*, 817, 34
- Mateos, S., Alonso-Herrero, A., Carrera, F. J., et al. 2012, *MNRAS*, 426, 3271
- McCammom, D., Almy, R., Apodaca, E., et al. 2002, *ApJ*, 576, 188
- Meidinger, N., Nandra, K., & Plattner, M. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10699, *Space Telescopes and Instrumentation 2018: Ultraviolet to Gamma Ray*, 106991F
- Meixner, M., Cooray, A., Leisawitz, D., et al. 2019, *arXiv e-prints*, arXiv:1912.06213
- Menci, N., Fiore, F., Puccetti, S., & Cavaliere, A. 2008, *ApJ*, 686, 219
- Mordini, S., Spinoglio, L., & Fernández-Ontiveros, J. A. 2021, *arXiv e-prints*, arXiv:2105.04584
- Nakagawa, T., Hayashi, M., Kawada, M., et al. 1998, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 3356, *HII/L2 mission*:

- future Japanese infrared astronomical mission, ed. P. Y. Bely & J. B. Breckinridge, 462–470
- Nakagawa, T., Shibai, H., Onaka, T., et al. 2014, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9143, The next-generation infrared astronomy mission SPICA under the new framework, 91431I
- Nandra, K., Barret, D., Barcons, X., et al. 2013, arXiv e-prints, arXiv:1306.2307
- O’Dowd, M. J., Schiminovich, D., Johnson, B. D., et al. 2009, *ApJ*, 705, 885
- Page, M. J., Stevens, J. A., Ivison, R. J., & Carrera, F. J. 2004, *ApJ*, 611, L85
- Peca, A., Vignali, C., Gilli, R., et al. 2021, *ApJ*, 906, 90
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, 518, L2
- Polletta, M., Weedman, D., Hönic, S., et al. 2008, *ApJ*, 675, 960
- Pozzi, F., Vignali, C., Comastri, A., et al. 2007, *A&A*, 468, 603
- Ricci, C., Ueda, Y., Koss, M. J., et al. 2015, *ApJ*, 815, L13
- Riechers, D. A., Bradford, C. M., Clements, D. L., et al. 2013, *Nature*, 496, 329
- Rodriguez, L., Poglitsch, A., Aliane, A., et al. 2018, *Journal of Low Temperature Physics*, 193, 449
- Roelfsema, P. R., Shibai, H., Armus, L., et al. 2018, *PASA*, 35, e030
- Rowan-Robinson, M., Mann, R. G., Oliver, S. J., et al. 1997, *MNRAS*, 289, 490
- Rowan-Robinson, M., Oliver, S., Wang, L., et al. 2016, *MNRAS*, 461, 1100
- Schreiber, C., Pannella, M., Elbaz, D., et al. 2015, *Astronomy and Astrophysics*, 575, A74
- Scoville, N., Aussel, H., Brusa, M., et al. 2007, *ApJS*, 172, 1
- Shankar, F., Weinberg, D. H., & Miralda-Escudé, J. 2009, *ApJ*, 690, 20
- Shankar, F., Weinberg, D. H., & Miralda-Escudé, J. 2013, *MNRAS*, 428, 421
- Shi, Y., Rieke, G. H., Hines, D. C., et al. 2006, *ApJ*, 653, 127
- Sijacki, D., Vogelsberger, M., Genel, S., et al. 2015, *MNRAS*, 452, 575
- Silk, J. & Rees, M. J. 1998, *A&A*, 331, L1
- Simmonds, C., Buchner, J., Salvato, M., Hsu, L. T., & Bauer, F. E. 2018, *A&A*, 618, A66
- Smith, J. D. T., Draine, B. T., Dale, D. A., et al. 2007, *ApJ*, 656, 770
- Smith, R. N., Tombesi, F., Veilleux, S., Lohfink, A. M., & Luminari, A. 2019, *ApJ*, 887, 69
- Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175
- Spinoglio, L., Alonso-Herrero, A., Armus, L., et al. 2017, *PASA*, 34, e057
- Spinoglio, L. & Malkan, M. A. 1992, *ApJ*, 399, 504
- Spinoglio, L., Mordini, S., Fernández-Ontiveros, J. A., et al. 2021, *PASA*, 38, e021
- Sturm, E., Hasinger, G., Lehmann, I., et al. 2006, *ApJ*, 642, 81
- Swinyard, B., Nakagawa, T., Merken, P., et al. 2009, *Experimental Astronomy*, 23, 193
- Teplitz, H. I., Armus, L., Soifer, B. T., et al. 2006, *ApJ*, 638, L1
- Thomas, N., Davé, R., Anglés-Alcázar, D., & Jarvis, M. 2019, *Monthly Notices of the Royal Astronomical Society*, 487, 5764–5780
- Tombesi, F., Meléndez, M., Veilleux, S., et al. 2015, *Nature*, 519, 436
- Tommasin, S., Spinoglio, L., Malkan, M. A., & Fazio, G. 2010, *ApJ*, 709, 1257
- Tommasin, S., Spinoglio, L., Malkan, M. A., et al. 2008, *ApJ*, 676, 836
- Tozzi, P., Gilli, R., Mainieri, V., et al. 2006, *A&A*, 451, 457
- Treister, E., Urry, C. M., & Virani, S. 2009, *ApJ*, 696, 110
- Ueda, Y., Akiyama, M., Hasinger, G., Miyaji, T., & Watson, M. G. 2014, *ApJ*, 786, 104
- Vignali, C., Piconcelli, E., Lanzuisi, G., et al. 2011, *MNRAS*, 416, 2068
- Vignali, C., Pozzi, F., Fritz, J., et al. 2009, *MNRAS*, 395, 2189
- Vito, F., Brandt, W. N., Yang, G., et al. 2018, *MNRAS*, 473, 2378
- Vito, F., Gilli, R., Vignali, C., et al. 2014, *MNRAS*, 445, 3557
- Voit, G. M. 1992, *MNRAS*, 258, 841
- Volonteri, M., Dubois, Y., Pichon, C., & Devriendt, J. 2016, *MNRAS*, 460, 2979
- Weedman, D., Polletta, M., Lonsdale, C. J., et al. 2006, *ApJ*, 653, 101
- Wu, Y., Charmandaris, V., Huang, J., Spinoglio, L., & Tommasin, S. 2009, *ApJ*, 701, 658
- Xue, Y. Q. 2017, *New Astronomy Review*, 79, 59
- Xue, Y. Q., Luo, B., Brandt, W. N., et al. 2011, *ApJS*, 195, 10
- Yan, L., Sajina, A., Fadda, D., et al. 2007, *ApJ*, 658, 778
- Yang, G., Boquien, M., Buat, V., et al. 2020, *MNRAS*, 491, 740
- Zappacosta, L., Piconcelli, E., Duras, F., et al. 2018, *A&A*, 618, A28

A AFFILIATIONS

¹Dipartimento di Fisica e Astronomia, Università degli Studi di Bologna, via P. Gobetti 93/2, 40129 Bologna, Italy

²INAF-OAS, Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, via Gobetti 93/3, 40129 Bologna, Italy

³Instituto de Física de Cantabria (CSIC-U. Cantabria), Avenida de los Castros, 39005 Santander, Spain

⁴Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

⁵Dipartimento di Fisica e Astronomia, Università di Firenze, via G. Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy

⁶INAF—Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy

⁷CEA, IRFU, DAp, AIM, Université Paris-Saclay, Université Paris Diderot, Sorbonne Paris Cité, CNRS, F-91191 Gif-sur-Yvette, France

⁸INAF - Osservatorio Astronomico di Brera, via Brera 28, I-20121, Milano, Italy & via Bianchi 46, I-23807, Merate, Italy

⁹School of Physics and Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, UK

¹⁰University of California, Irvine, Irvine, CA

¹¹SISSA, Via Bonomea 265, 34136 Trieste, Italy

¹²Istituto di Astrofisica e Planetologia Spaziali - INAF, Rome, Via Fosso del Cavaliere 100, 00133, Roma, Italia

¹³Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan

¹⁴Institute of Liberal Arts and Sciences, Tokushima University, Minami Jousanjima-Machi 1-1, Tokushima, Tokushima 770-8502, Japan

¹⁵Centro de Astrobiología (CSIC-INTA), Ctra. de Ajalvir, Km 4, 28850, Torrejón de Ardoz, Madrid, Spain

¹⁶Osservatorio Astronomico di Roma (INAF), Via Frascati 33, I-00040 Monte Porzio Catone (Roma), Italy

¹⁷Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

¹⁸Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, People's Republic of China

¹⁹George Mason University, Department of Physics & Astronomy, MS 3F3, 4400 University Drive, Fairfax, VA 22030, USA

²⁰Dipartimento di Fisica e Astronomia, Università di Padova, vicolo Osservatorio 3, 35122 Padova, Italy

²¹Department of Physics, University of Rome “Tor Vergata”, Via della Ricerca Scientifica 1, I-00133 Rome, Italy

²²Department of Astronomy, University of Maryland, College Park, MD 20742, USA

²³Astrophysics Science Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA