

Table S1. Strains used in this study. Natural *N. meningitidis* serogroup B strains used in Figure 1 and their main characteristics are indicated. For each of them a *nhba* deletion mutant was generated. * Unassigned. □ Kan, kanamycin, Ery, erythromycin, Clm, chloramphenicol.

Natural Strains	Description	Clonal Complex	CREN	NHBA Peptide Variant	NHBA Isoform
M11205	Clinical isolate, USA 2003	41/44	-	p0001	Long
M11822	Clinical isolate, USA 2004	41/44	-	p0001	Long
NGH38	Clinical isolate, Norway 1988	ua *	-	p0002	Long
MC58	Laboratory-adapted reference strain, UK 1985	32	+	p0003	Long
M10935	Clinical isolate, USA 2003	35	-	p0058	Long
M14933	Clinical isolate, USA 2006	32	+	p0003	Long
M10713	Clinical isolate, USA 2003	41/44	-	p0010	Short
M03279	Clinical isolate, USA 1997	41/44	-	p0011	Short
N16/97	Clinical isolate, Norway 2007	41/44	-	p0029	Long
M11204	Clinical isolate, USA 2003	41/44	-	p0029	Long
M10282	Clinical isolate, USA 2003	41/44	-	p0029	Long
M07-0240679	Clinical isolate, UK 2007	269	-	p0017	Short
M11719	Clinical isolate, USA 2003	162	-	p0020	Short
M16452	Clinical isolate, USA 2007	41/44	-	p0144	Short
M18070	Clinical isolate, USA 2008	162	-	p0020	Short
8047	Clinical isolate, USA 1978	11	-	p0020	Short
Engineered Strains	Description				
C_nhba	Complemented <i>nhba</i> mutant of MC58; EryR and KanR □ [39]				
Ptac_nhba	Complemented mutant of MC58 expressing <i>nhba</i> (MC58 sequence) under the control of an IPTG-inducible Ptac promoter; EryR and ClmR □ [39]				

Pwt_	mCherry	Complemented mutant of MC58 expressing <i>mCherry</i> reporter gene under the control of <i>nhba</i> intergenic region (MC58 sequence); EryR and ClmR
Ptac_	mCherry	Complemented mutant of MC58 expressing <i>mCherry</i> reporter gene under the control of an IPTG-inducible Ptac promoter; EryR and ClmR

Table S2. Oligonucleotides used in this study.

Plasmid	Description	Application
NMB2132RT-F	GGCTTGTATTTGCCCTTC	qRT-PCR
NMB2132RT-R	GCATCTCCTTGCCCTGT	
mCherry_RT1 fw	GCTCAGTTCAGGTAGTCG	qRT-PCR
mCherry_RT1 REV	AGCCCTCAATTCATGTACGG	
pRTNM16sII.F1	GTGGGGAATTGGACAATG	qRT-PCR
pRTNM16sII.R1	CAACAGCCTTCTCCCTG	
mCherry Fw	TCCAAAGGCGAGGAAGATAAC	Generation of Pwt_mCherry fusion: insert
mCherry Rev	CACGGATCCCTGCAGTTATTGTAC AGTCGTCCATG	PCR to amplify the <i>mCherry</i> gene codon- optimized for <i>N. meningitidis</i>
CmR Fw	CTGCAGGGATCCGTGATATAG	
Pwt_14aa Rev	CTTCCTCGCCTTGGAGGAAAAAT ACAAGCCATTGC	Generation of Pwt_mCherry fusion: vector PCR on pBS_nhba background
Insert Fw	CGAAAGTGGGAATCTAGACATTATC GGCGTGATTCA	Subcloning of Pwt_mCherry fusion into pComClmRP _{ind} plasmid
Insert Rev	GAGCATAAAATTAGTAACCTATG TTTTTATTCAAGCAAGTCTTGTAAATT	
Vector Fw	GAATTACAAGACTGCTGAATAAAA ACATAGGTTACTAAAATTATGCTC	Subcloning of Pwt_mCherry fusion into pComClmRP _{ind} plasmid
Vector Rev	CTGAATCACGCCGATAATGTCTAGA TTCCCACTTCG	
NmmCh_NdeI Fw	GAAACAC <u>CATATGATGGTGTCCAAAG</u> GCGAG	Cloning <i>mCherry</i> gene into pCOM_Ptac plasmid
NmmCh_NsiI Rev	GTGTCA <u>ATGCATTATTGTACAGTT</u> CGTCCATGCC	

Table S3. Plasmid used in this study.

Plasmid	Description	Antibiotic Resistance	Reference or Source
pBluescript (pBS)	Cloning vector	Amp	Stratagene
pBS-UDgna2132	Plasmid for the deletion of the <i>nhba</i> gene by homologous recombination	Amp, Ery	[39]
pBS_ <i>nhba</i>	pBS derivative for the <i>in-locus</i> complementation of the <i>nhba</i> gene (MC58 sequence)	Amp, Clm	[37]
pComClmR-Pind	Plasmid for the allelic replacement at chromosomal location between ORFs <i>NMB1428</i> and <i>NMB1429</i> and inducible expression under control of Ptac promoter and lacI repressor	Amp, Clm	[54]
pCOM- <i>Cnhba</i>	Plasmid for the <i>ex-locus</i> complementation of the <i>nhba</i> null mutant. Derivative of pComClmR-Pind containing a copy of <i>nhba</i> (MC58 sequence) under control of the <i>nhba</i> intergenic region.	Amp, Kan	[39]
pCOM-Ptac_ <i>nhba</i>	Plasmid for the <i>ex-locus</i> complementation of the <i>nhba</i> null mutant. Derivative of pComClmR-Pind containing a copy of <i>nhba</i> (MC58 sequence) under control of Ptac promoter and lacI repressor.	Amp, Clm	[39]
pBS_14aa_ <i>mCherry</i>	Derivative of pBS_ <i>nhba</i> . Construct for the <i>in-locus</i> complementation of the <i>mCherry</i> reporter gene under the control of the <i>nhba</i> intergenic region	Amp, Clm	This study
pCOM_14aa_ <i>mCherry</i>	Construct for the <i>ex-locus</i> complementation of the <i>mCherry</i> reporter gene under the control of the <i>nhba</i> intergenic region. Derivative of pComClmR-Pind	Amp, Clm	This study
pCOM_Ptac_ <i>mCherry</i>	Construct for complementation of the <i>mCherry</i> reporter gene under the control of an IPTG-inducible Ptac promoter. Derivative of pComClmR-Pind	Amp, Clm	This study

References

37. Vacca, I.; Del Tordello, E.; Gasperini, G.; Pezzicoli, A.; Di Fede, M.; Rossi Paccani, S.; Marchi, S.; Mubaiwa, T.D.; Hartley-Tassell, L.E.; Jennings, M.P.; et al. Neisserial Heparin Binding Antigen (NHBA) Contributes to the Adhesion of *Neisseria meningitidis* to Human Epithelial Cells. *PLoS ONE* **2016**, *11*, e0162878.
39. Serruto, D.; Spadafina, T.; Ciucchi, L.; Lewis, L.A.; Ram, S.; Tontini, M.; Santini, L.; Biolchi, A.; Seib, K.L.; Giuliani, M.M.; et al. *Neisseria meningitidis* GNA2132, a heparin-binding protein that induces protective immunity in humans. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 3770–3775.
54. Ieva, R.; Alaimo, C.; Delany, I.; Spohn, G.; Rappuoli, R.; Scarlato, V. CrgA is an inducible LysR-type regulator of *Neisseria meningitidis*, acting both as a repressor and as an activator of gene transcription. *J. Bacteriol.* **2005**, *187*, 3421–3430.