

Search for long-lived heavy neutral leptons in proton-proton collision events with a lepton-jet pair associated with a secondary vertex at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search for long-lived heavy neutral leptons (HNLs) using proton-proton collision data corresponding to an integrated luminosity of 138 fb^{-1} collected at $\sqrt{s} = 13$ TeV with the CMS detector at the CERN LHC is presented. Events are selected with a charged lepton originating from the primary vertex associated with the proton-proton interaction, as well as a second charged lepton and a hadronic jet associated with a secondary vertex that corresponds to the semileptonic decay of a long-lived HNL. No excess of events above the standard model expectation is observed. Exclusion limits at 95% confidence level are evaluated for HNLs that mix with electron and/or muon neutrinos. Limits are presented in the mass range of 1–16.5 GeV, with excluded square mixing parameter values reaching as low as 2×10^{-7} . For masses above 11 GeV, the presented limits exceed all previous results in the semileptonic decay channel, and for some of the considered scenarios are the strongest to date.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering

ARXIV EPRINT: [2407.10717](https://arxiv.org/abs/2407.10717)

Contents

1	Introduction	1
2	The CMS experiment and event reconstruction	3
3	Data and simulated samples	5
4	Event selection	7
5	Identification of the SV from an HNL decay	11
6	Background estimation	14
7	Systematic uncertainties	18
8	Results	22
9	Summary	26
	The CMS collaboration	36

1 Introduction

The discovery of neutrino oscillations [1–3] has provided experimental evidence that neutrinos have nonzero masses [4]. Constraints from cosmological probes, such as measurements of the cosmic microwave background [5–7], as well as direct measurements [8, 9], indicate that the neutrino masses are orders of magnitude smaller than the masses of other fermions in the standard model (SM) of particle physics. A possible way to generate gauge-invariant mass terms for the SM neutrinos and explain their small mass scale is the see-saw mechanism, which requires the introduction of new heavy states with right-handed chirality [10–17]. These new states, referred to as heavy neutral leptons (HNLs), can also provide explanations for other open questions in high-energy physics, such as the nature of dark matter [18, 19] or the matter-antimatter asymmetry in the universe [20–22].

The HNL flavor eigenstates are singlets of the SM gauge groups, and thus do not interact with the SM particles through the electroweak or strong interaction. They do, however, form mixed mass eigenstates with the SM neutrinos. This mixing generates HNL-dominated mass eigenstates with effective electroweak couplings, and hence these mass eigenstates can be produced and decay via the electroweak interaction. For an explanation of both neutrino observations and cosmological problems, an SM extension with HNLs generally needs to provide three HNL generations with couplings to all SM neutrino generations [23]. In the following, we will refer to the HNL mass eigenstates and indicate them with the symbol N . A wide range of HNL masses m_N and mixing parameters $V_{\ell N}$ can yield mass values for the SM neutrinos consistent with observed neutrino oscillations and offer an explanation for

the matter-antimatter asymmetry in the universe. In the case of, e.g., $m_N = 10$ GeV, these requirements are found to be fulfilled for a range $10^{-11} < |V_{\ell N}|^2 < 10^{-5}$ [22]; a significant part of which can be probed in proton-proton (pp) collisions at the CERN LHC.

The experimental signature of HNL models has been studied extensively [17, 24–31], and searches have been conducted at the LHC [32–49], at other colliders [50–54], and in fixed-target experiments [55–65], covering a mass range from a few keV to several TeV [31]. The proper lifetime t of HNLs follows an exponential distribution $P(t) = \tau_N^{-1} \exp(-t/\tau_N)$ with mean proper lifetime τ_N depending on m_N and $V_{\ell N}$ as [25]

$$\tau_N \propto m_N^{-5} |V_{\ell N}|^{-2}. \quad (1.1)$$

For $m_N < 20$ GeV, the value of τ_N can become so large that many HNLs will decay after a time t long enough for the HNL decay products to emerge from a secondary vertex (SV) that is spatially displaced from the primary vertex (PV) and separately reconstructable. Collider searches for HNL production can thus be optimized either for short-lived HNL scenarios with prompt signatures, or for long-lived HNL scenarios with displaced signatures.

This paper reports a search for W boson decays to an HNL and a charged lepton ℓ_1 , with a subsequent HNL decay to a second charged lepton ℓ_2 and two quarks. The search is optimized for displaced HNL decays with $1 < m_N < 20$ GeV, where the large boost of the HNL in the W boson decay causes the two quarks to be merged into a single jet that can also encompass ℓ_2 . The experimental signature thus consists of a prompt ℓ_1 originating from the PV, and a nonprompt ℓ_2 encompassed in a jet originating from an SV. We consider only electrons and muons for ℓ_1 and ℓ_2 , and refer to them as “leptons” in the following. Two possible decay modes are considered for the HNLs: lepton-number-conserving (LNC) decays with ℓ_1 and ℓ_2 of opposite sign (OS), and lepton-number-violating (LNV) decays with same-sign (SS) leptons. The case where both LNC and LNV decays are possible is referred to as the “Majorana” nature of the HNL, whereas the “Dirac” nature refers to HNLs with only LNC decays. As a consequence of having twice the number of possible decays, the decay width (τ_N) of a Majorana HNL is twice (half) that of a Dirac HNL with the same m_N and $|V_{\ell N}|^2$. Examples of leading-order (LO) Feynman diagrams of the studied process with either an LNC or LNV decay are shown in figure 1.

The analyzed pp collision data were recorded by the CMS experiment at $\sqrt{s} = 13$ TeV in 2016–2018, and correspond to an integrated luminosity of 138 fb^{-1} . We select events with one prompt lepton and apply a dedicated reconstruction of the SV with an associated nonprompt lepton and a jet. Events are grouped into the four lepton flavor categories ee , $\mu\mu$, $e\mu$, and μe , where the first (second) symbol refers to the flavor of ℓ_1 (ℓ_2). A machine-learning algorithm that uses information related to the nonprompt lepton, the jet, and the SV is trained to distinguish between HNL signal and SM background events. Background contributions from SM processes can arise when a displaced lepton forms an SV by overlapping with a random track, when a photon converts into a lepton pair in the detector material, or when a heavy-flavor hadron decay results in a displaced lepton associated with a genuine SV. These contributions are estimated from control samples in data. For the interpretation of our results, we consider simplified models with a single HNL that couples exclusively to




Figure 1. Examples of LO Feynman diagrams for production and decay of an HNL (indicated with the symbol N) resulting in a final state with two charged leptons and two quarks. In the left diagram, the HNL is a Dirac particle and thus the two charged leptons must have opposite charge. In the right diagram, the HNL is a Majorana particle and the two charged leptons can have the same charge.

electron and/or muon neutrinos (refs. [66–69] discuss limitations of this approach). The ee ($\mu\mu$) channel is used to constrain $|V_{eN}|^2$ ($|V_{\mu N}|^2$) in HNL scenarios with an exclusive coupling to electron (muon) neutrinos, whereas the $e\mu$ and μe channels are used together to constrain $|V_{eN}V_{\mu N}|^2/(|V_{eN}|^2 + |V_{\mu N}|^2)$ in HNL scenarios with couplings to both electron and muon neutrinos [25]. Exclusion limits are derived on the respective $|V_{\ell N}|^2$ values as functions of m_N . Tabulated results are provided in the HEPData record for this analysis [70].

Recent analyses by the CMS experiment have used the same data set to search for HNLs in various mass ranges and final-state signatures. The search presented in ref. [44] targets the same decay channel and m_N range as the one presented in this paper, but uses a deep-neural-network-based jet tagger to identify the displaced decay products without explicitly reconstructing an SV. The explicit SV reconstruction employed in the analysis presented here results in more stringent limits in the region with $m_N > 10$ GeV and high τ_N values. Searches for HNL production and decay resulting in events with three leptons have been performed in the mass range $10 \text{ GeV} < m_N < 1.5 \text{ TeV}$, where all leptons are required to originate from the PV [46], and in the mass range $1 < m_N < 20 \text{ GeV}$, where two of the leptons are required to originate from an SV [40]. Other results target the range $1 < m_N < 3 \text{ GeV}$ by reconstructing HNLs with $c\tau_N > 0.1 \text{ m}$ decaying in the muon system [45] or by searching for HNLs in B meson decays [47]. A summary of all HNL searches performed by the CMS Collaboration is presented in ref. [49].

2 The CMS experiment and event reconstruction

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in refs. [71, 72].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about $4\,\mu\text{s}$ [73]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [74, 75].

Charged-particle tracks are reconstructed across the entire η range of the silicon tracker, which is $|\eta| < 2.5$ with the tracker used in 2016 and $|\eta| < 3.0$ with the new pixel detector installed at the start of 2017 [76]. Track reconstruction enables the detection of charged particles with transverse momentum (p_T) as low as 0.1 GeV and originating from distances up to 60 cm from the beam line. In simulation with 2017 detector conditions, the tracking efficiency for charged particles with $p_T > 0.9$ GeV of better than 90, 60, and 25% for tracks originating at distances smaller than 2, 10, and 48 cm from the beamline, respectively, is found [77]. These charged-particle tracks are then clustered to form vertices. To reconstruct vertices corresponding to pp interactions, the clustering is performed based on the z coordinate of the closest approach of the tracks to the beam line. The vertex position is estimated with an adaptive vertex fitter [78] using the collection of tracks compatible with originating from the same interaction. From the vertices reconstructed in this way, the PV is taken to be the one corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [79].

The impact parameter of a track is defined as the distance of the track from the PV at its point of closest approach. The three-dimensional (3D) impact parameter is referred to as d_{xyz} , the two-dimensional (2D) value in the plane transverse to the beam line as d_{xy} , and the longitudinal value along the beam line as d_z . For nonisolated particles of $1 < p_T < 10$ GeV and $|\eta| < 1.4$, the track resolutions in 2016 were typically 1.5% in p_T and 25–90 (45–150) μm in d_{xy} (d_z) [80]. From 2017 onward, the upgraded tracker measured particles up to $|\eta| < 3.0$ with typical resolutions of 1.5% in p_T and 20–75 μm in d_{xy} for nonisolated particles of $1 < p_T < 10$ GeV [81].

To reconstruct vertices corresponding to the decay of long-lived particles displaced from the PV, we employ a modified version of the inclusive vertex finder (IVF) algorithm [82, 83] originally developed for the identification of displaced B hadron decays. The IVF clusters tracks that have at least six hits in the tracker and $p_T > 0.8$ GeV, and is seeded with tracks that additionally have a 3D impact parameter significance, defined as d_{xyz} divided by its uncertainty, above 1.2. The 3D (2D) distance of the SV to the PV is referred to as Δ_{3D} (Δ_{2D}). All SVs are retained that have a significance of Δ_{3D} (Δ_{2D}) above 0.5 (2.5), while no explicit requirements are applied on the values of Δ_{3D} or Δ_{2D} directly. From the high number of SVs reconstructed with the IVF, we select the candidate vertex corresponding to the HNL decay by requiring that the SV contains the track associated with ℓ_2 , selected as described in section 4. If more than one SV contains the ℓ_2 track, we select the SV with the highest associated p_T .

The particle-flow (PF) algorithm [84] aims to reconstruct and identify each individual particle in an event, with an optimized combination of information from the various elements of the CMS detector. The energy of photons is obtained from the ECAL measurement. The energy of electrons is determined from a combination of the electron momentum at the PV,

the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The energy of muons is obtained from the curvature of the corresponding track. The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

Reconstructed PF particles are clustered into jets with the anti- k_T algorithm [85, 86] using a distance parameter of 0.4. The jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5–10% of the true momentum over the whole p_T spectrum and detector acceptance. Additional pp interactions within the same or nearby bunch crossings (pileup) can contribute with additional tracks and calorimetric energy depositions, increasing the apparent jet momentum. To mitigate this effect, tracks identified to be originating from pileup vertices are discarded and an offset correction is applied to account for remaining contributions [87]. Jet energy corrections are derived from simulation studies so that the average measured energy of jets becomes identical to that of particle-level jets. In situ measurements of the momentum balance in dijet, photon+jet, Z+jet, and multijet events are used to determine any residual differences between the jet energy scale in data and simulation, and appropriate corrections are made [88]. Additional selection criteria are applied to each jet to remove jets potentially dominated by instrumental effects or reconstruction failures [87]. Finally, jets are required to have $p_T > 20$ GeV and $|\eta| < 2.4$.

Electrons are reconstructed within the geometrical acceptance of the silicon strip tracker, which is $|\eta| < 2.5$. The electron momentum is estimated by combining the energy measurement in the ECAL with the momentum measurement in the tracker. The momentum resolution for electrons with $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.6 to 5%. It is generally better in the barrel region than in the endcaps, and also depends on the bremsstrahlung energy emitted by the electron as it traverses the material in front of the ECAL [89, 90].

Muons are measured in the range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive plate chambers. The efficiency to reconstruct and identify muons is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative p_T resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps [91].

3 Data and simulated samples

Data events have been recorded with single-electron and single-muon triggers. For the single-electron trigger, the presence of at least one isolated electron at the HLT is required with a p_T threshold of 27 (32) GeV in 2016 (2017–2018) [92]. For the single-muon trigger, the presence of at least one isolated muon at the HLT is required with a p_T threshold of 24 GeV throughout all data-taking years [93].

Monte Carlo (MC) event simulation is used to generate HNL signal events for the evaluation of the signal selection efficiency, and SM background events for the validation of the background estimation. For each signal and background process, separate simulated event

samples are generated corresponding to the conditions of the three data-taking years. For the 2016 (2017–2018) samples, the NNPDF 3.0 [94] (3.1 [95]) parton distribution functions (PDFs) are used in the simulation of the hard process. All generators are interfaced with the PYTHIA 8.226 (8.230) simulation [96] for parton showering and hadronization, using the CUETP8M1 [97] (CP5 [98]) tune for the underlying event modeling in case of the event modeling for the 2016 (2017–2018) samples. Simulated minimum bias events are superimposed on the generated events to reproduce the pileup distribution obtained from the measured instantaneous luminosity. For all simulated events, the CMS detector response is modeled with the GEANT4 toolkit [99].

Background samples for W boson production in association with jets (W+jets) are generated at next-to-LO (NLO) with the MADGRAPH5_aMC@NLO 2.4.2 event generator [100], including up to two additional partons in the matrix-element calculation. For Drell-Yan (DY) dilepton production, event samples are generated at LO (NLO) with up to four (two) additional partons in the matrix-element calculation with MADGRAPH5_aMC@NLO. Top quark (pair and single production) and diboson (WW, WZ, ZZ, W γ , Z γ) processes are simulated with the POWHEG v2 [101–108] and MADGRAPH5_aMC@NLO event generators at NLO. Samples generated with MADGRAPH5_aMC@NLO at LO (NLO) use the MLM [109] (FxFx [110]) merging algorithm. For the background from events comprised uniquely of jets produced through the strong interaction (QCD multijet events), event samples are generated at LO with PYTHIA.

Signal events for HNL production are simulated at LO with MADGRAPH5_aMC@NLO, using a model that extends the SM particle content by introducing up to three right-handed neutrinos [25, 111, 112]. Event samples are generated for a single HNL mass eigenstate with pure electron neutrino couplings, pure muon neutrino couplings, and equal electron and muon neutrino couplings, covering the parameter ranges $1 < m_N < 20$ GeV and $10^{-9} < |V_{\ell N}|^2 < 10^{-1}$. The cross section of the HNL event samples is calculated at LO with MADGRAPH5_aMC@NLO, and scaled to next-to-NLO (NNLO) precision using a K -factor of 1.089 derived from the comparison of SM W boson production samples generated at LO accuracy with MADGRAPH5_aMC@NLO, using the same settings as the signal simulation, and the cross section calculated at NNLO accuracy with the FEWZ v3.1 program [113–116].

For a fixed value of m_N , we emulate an event sample with a different $|V_{\ell N}|^2$ value by applying a global weight to match the expected cross section and a per-event weight to correct the proper lifetime distribution. The cross section is proportional to $|V_{\ell N}|^2$ [112] and the global weight is accordingly calculated as the ratio of the new to the simulated $|V_{\ell N}|^2$ value. To reweight from τ_N values τ_0 to τ_1 , the per-event weights for events with proper lifetime t are calculated as

$$W(t) = \frac{P(t; \tau_1)}{P(t; \tau_0)} = \frac{\tau_0}{\tau_1} \exp \left[-t \left(\frac{1}{\tau_1} - \frac{1}{\tau_0} \right) \right]. \quad (3.1)$$

The simulation is performed assuming Majorana HNLs, i.e., both LNC and LNV decays are included. To obtain an event sample for Dirac HNLs, we select only events with LNC decays. The production cross section is the same for Majorana and Dirac HNLs, and thus the global weight is only adjusted for the smaller number of generated events in the sample.

Criterion	Prompt		Nonprompt	
	Electrons	Muons	Electrons	Muons
p_T [GeV]	>30 (34)	>30	>7	>7
$ \eta $	<2.5	<2.4	<2.5	<2.4
I_{rel}	<0.1	<0.1	—	—
$ d_{xy} $ [cm]	<0.02	<0.01	>0.02	>0.02
$ d_z $ [cm]	<0.04	<0.1	<10	<10
ID	full	full	reduced	reduced

Table 1. Selection criteria for electrons and muons. Numbers in parentheses indicate values applied in the 2017–2018 data sets, when different from those for 2016.

The per-event weights are calculated with eq. (3.1) but using that τ_1 is twice as large for a Dirac HNL with respect to a Majorana HNL of same m_N and τ_N .

4 Event selection

To select prompt and nonprompt leptons and to reduce the contribution from particles misidentified as leptons, additional identification (ID) criteria are applied to the reconstructed electrons and muons. We employ two standard sets of electron ID criteria defined in ref. [89] that both have an efficiency of 90%, where the first is based on sequential selection of requirements on seven observables (“sequential ID”), and the second is based on a multivariate ID discriminant trained with an extended set of observables (“multivariate ID”). For muons, we use the set of “medium” ID criteria defined in ref. [91] that has an efficiency of 99.5%. In addition, the relative isolation I_{rel} , defined as the scalar sum of the of surrounding particles divided by the lepton p_T , is used to quantify how isolated a reconstructed lepton is from other particles in the event. The isolation sum requires particles to be within a cone of $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 0.3$, where $\Delta\eta$ and $\Delta\phi$ are the η and azimuthal angle differences between the lepton and the particle, respectively. Charged hadrons are included only if they originate from the PV, while all neutral hadrons and photons within the cone are included and additional corrections for the contribution of neutral particles from pileup interactions are applied [89, 91]. Finally, we employ selection criteria for the impact parameters of the tracks associated with the electrons and muons, as defined in section 2.

The selection criteria for prompt and nonprompt electrons and muons are listed in table 1. Prompt electrons and muons are required to have a p_T above a threshold chosen larger than the HLT single-lepton trigger thresholds, to be isolated, and to originate from the PV. Nonprompt electrons and muons are selected with a much lower p_T threshold and no isolation requirement. Instead of consistency with the PV, a minimal requirement on d_{xy} is imposed, and a loose maximal requirement on d_z is used to reduce background contributions from cosmic ray muons and pileup. In addition, we require that electrons and muons pass a set of ID criteria as introduced before, where prompt leptons have to pass the “full” set and nonprompt leptons only a “reduced” set without criteria that are inefficient for displaced




Figure 2. Selection efficiencies of nonprompt electrons (upper) and muons (lower), evaluated in simulated HNL signal events as functions of the generated lepton p_T (left) and transverse displacement of the generated SV (right). Three HNL signal scenarios are shown, with $m_N = 3$ GeV and $|V_{\ell N}|^2 = 9.9 \times 10^{-5}$ (HNL3, corresponding to $c\tau_N = 23$ mm), $m_N = 5$ GeV and $|V_{\ell N}|^2 = 1.6 \times 10^{-6}$ (HNL5, corresponding to $c\tau_N = 92$ mm), and $m_N = 10$ GeV and $|V_{\ell N}|^2 = 5.7 \times 10^{-7}$ (HNL10, corresponding to $c\tau_N = 7$ mm). The error bars in the plot represent statistical uncertainties.

leptons. For prompt electrons, the multivariate ID is used. For nonprompt electrons, the sequential ID is used except for the requirements on the maximum number of missing hits in the pixel detector and the veto on photon conversions, as defined in ref. [89]. We use the medium ID for both prompt and nonprompt muons, but remove the requirements on the fraction of valid tracker hits associated with the muon track and on the fit quality of the global muon track fit [91] for nonprompt muons.

The selection efficiencies for nonprompt electrons and muons are evaluated in HNL signal samples with different $(m_N, |V_{\ell N}|^2)$ scenarios, representing $c\tau_N$ values in the range 7–92 mm, and shown in figure 2 as functions of the generated p_T and the transverse displacement Δ_{2D}

of the generated SV. The simulated HNL signal events used for this evaluation are selected by requiring a generated nonprompt electron (muon) with $p_T > 7$ GeV and $|\eta| < 2.5$ (2.4), and a generated HNL decay vertex with $\Delta_{2D} < 50$ cm. The obtained efficiencies reflect various effects in the electron and muon reconstruction methods and identification criteria, as well as effects caused by different m_N and $c\tau_N$ values. Overall, the better resolution and higher efficiency in the basic reconstruction of muons compared to electrons [89, 91] results in higher selection efficiencies for nonprompt muons when compared to electrons, which leads to a better sensitivity of the analysis to $|V_{\mu N}|^2$ than $|V_{e N}|^2$. At low values of lepton p_T , the track reconstruction in the inner tracker has a worse efficiency and resolution, decreasing especially the efficiency for nonprompt electrons. At high values of lepton p_T , the decay products of the HNL will be boosted and thus the nonprompt lepton is closer to or fully included in the jet from the HNL decay, which reduces the efficiency to pass selection criteria designed to reject hadrons misidentified as leptons. Due to differences in the track reconstruction and combination with ECAL deposits or the muon system track, respectively, the p_T -dependent shape is different between nonprompt electrons and muons. The selection efficiency of nonprompt electrons decreases quickly with larger displacement, since the dedicated electron track reconstruction is very sensitive to missing hits in the inner tracker [89], whereas the muon reconstruction including the muon system track is less dependent on a well-reconstructed track in the inner tracker. For nonprompt muons, a small inefficiency can be seen in the first bin of the efficiency as a function of Δ_{2D} , caused by the minimal requirement on $|d_{xy}|$ listed in table 1.

Events are selected with exactly one prompt electron or muon (ℓ_1), which is required to match geometrically to the lepton reconstructed at the HLT that satisfies the trigger selection. Further, events are required to have exactly one nonprompt electron or muon (ℓ_2), as well as one jet matched to ℓ_2 by the requirement $\Delta R < 0.7$. To reconstruct the HNL decay vertex, the ℓ_2 track is required to be part of an SV reconstructed by the IVF, as described in section 2. The SV reconstruction is validated in simulation by comparing properties of the reconstructed SV with the generated HNL decay vertex. In figure 3, the SV reconstruction efficiency is shown as a function of Δ_{2D} , using HNL signal events with an SV that contains the nonprompt-lepton track and is matched to the generated HNL decay vertex. The vertex matching is defined by the criterion $|\Delta_{3D} - \Delta_{3D}^{\text{gen}}|/\Delta_{3D}^{\text{gen}} < 0.1$, where Δ_{3D}^{gen} is the 3D displacement of the generated HNL decay vertex.

Further requirements are imposed on the invariant mass of the two selected leptons, $m(\ell_1\ell_2) > 10$ GeV, and on their angular separation, $|\Delta\phi(\ell_1, \ell_2)| > 0.4$. These requirements remove poorly reconstructed events, as well as background events originating from low-mass resonances. Events with at least one additional jet separated from ℓ_1 by $\Delta R > 0.4$, as expected for top quark production, are removed and used as an orthogonal event selection for the validation of the background prediction described in section 6. Events where ℓ_1 and ℓ_2 are OS muons and have $85 < m(\mu^\pm\mu^\mp) < 95$ GeV are removed to suppress DY dimuon background contributions. This DY veto is introduced because of a correlation between the observables used in the background estimation observed in the OS dimuon channels, as described in section 6, but is not required for the other channels where no such correlation is observed. The event selection criteria are summarized in table 2.




Figure 3. The SV reconstruction efficiency in simulated HNL signal events as a function of the SV displacement for vertices with a nonprompt electron (left) or muon (right). Three HNL signal scenarios are shown, with $m_N = 3$ GeV and $|V_{\ell N}|^2 = 9.9 \times 10^{-5}$ (HNL3, corresponding to $c\tau_N = 23$ mm), $m_N = 5$ GeV and $|V_{\ell N}|^2 = 1.6 \times 10^{-6}$ (HNL5, corresponding to $c\tau_N = 92$ mm), and $m_N = 10$ GeV and $|V_{\ell N}|^2 = 5.7 \times 10^{-7}$ (HNL10, corresponding to $c\tau_N = 7$ mm). The error bars in the plot represent statistical uncertainties.

Event selection criteria
$N(\text{prompt } \ell_1) = 1$
$N(\text{nonprompt } \ell_2) = 1$
$N(\text{jets}) = 1$
$\Delta R(\ell_2, \text{jet}) < 0.7$
$\ell_2 \in \text{SV}$
$m(\ell_1 \ell_2) > 10 \text{ GeV}$
$ \Delta\phi(\ell_1, \ell_2) > 0.4$
$m(\mu^\pm \mu^\mp) \notin [85, 95] \text{ GeV}$

Table 2. Summary of the event selection criteria.

Separate event categories are defined based on the flavors of ℓ_1 and ℓ_2 (ee, e μ , $\mu\mu$, and $\mu\mu$), and for the cases of SS and OS charges. The invariant mass of ℓ_1 and the tracks belonging to the selected SV, $m(\ell_1, \text{SV})$, is an important observable for the discrimination between signal and background events. For HNL signal events, $m(\ell_1, \text{SV})$ is kinematically limited by the W boson mass since it includes all charged particles originating from the W boson decay in the hard interaction but does not account for potential neutral hadrons and photons formed in the hadronic decay. The distribution of $m(\ell_1, \text{SV})$ predicted from simulated event samples is shown in figure 4, separately for the different event categories.




Figure 4. The $m(\ell_1, \text{SV})$ distribution of predicted events yields after applying the selection summarized in table 1, for the ee (upper left), $\mu\mu$ (upper right), $e\mu$ (lower left), and μe (lower right) categories. The filled histograms display the predicted background yields. The lines show the predicted yields for three HNL signal scenarios, with $m_N = 3$ GeV and $|V_{\ell N}|^2 = 9.9 \times 10^{-5}$ (HNL3), $m_N = 5$ GeV and $|V_{\ell N}|^2 = 1.6 \times 10^{-5}$ (HNL5), and $m_N = 10$ GeV and $|V_{\ell N}|^2 = 5.7 \times 10^{-7}$ (HNL10). The HNL signal yield is normalized to the total background yield. The last bins include the overflow.

5 Identification of the SV from an HNL decay

A multivariate discriminant based on machine learning is employed to distinguish between events with a reconstructed SV originating from an HNL decay and background events where, e.g., the jet accidentally overlaps with the nonprompt lepton. We employ a particle-flow network (PFN) [117], an application of the deep sets theorem [118], that uses per-particle information of the particles associated with the jet as well as per-event information associated with the nonprompt lepton, the jet, and the SV. The PFN architecture consists of two sections, with three fully connected neural network layers of 128 nodes each per section. First, each individual particle is passed through the per-particle section, which is provided with the per-particle information and produces a 256-dimensional output representing the

latent space. The per-particle outputs are added up in a summation layer, and the sums are provided together with event-level variables as inputs to the second section that processes the full event information. The two sections are combined into a single network, which allows for a simultaneous optimization of both the per-particle and per-event sections. The PFN output is a single score value between zero and one, where high (low) values are assigned to signal-like (background-like) events.

Up to 50 PF particles associated with the selected jet are used for the per-particle section. For each particle, the p_T , η , ϕ , charge, particle type as identified by the PF algorithm, impact parameter properties and uncertainties, number of hits in the pixel and strip detectors of the tracker system [80], and the association with tracks and SVs are provided as input variables to the PFN.

The per-event input variables are associated with the selected jet, nonprompt lepton, and SV. For the jet, the kinematic properties and the number of constituent particles are provided. For the nonprompt lepton, the kinematic properties, I_{rel} , impact parameter properties and uncertainties, and the number of hits in the pixel detector are used. Additionally, the ΔR between lepton and jet, the transverse component of the lepton momentum with respect to the jet axis, and the p_T ratio between the lepton and the jet are provided. For the SV, the kinematic properties including the invariant mass m_{SV} of the tracks associated with the SV, the number of tracks, the normalized χ^2 of the reconstruction as determined with the IVF, and the distance from the PV are used.

The PFN is trained with MC samples for the HNL signal, and the SM background samples described in section 3. Only the QCD multijet sample is excluded from the training, because of its large statistical uncertainties. Only events passing the event selection summarized in table 2 are used for the training. To account for the upgrade of the pixel detector in 2017 [76], PFNs are trained separately for the 2016 data set and the combined 2017–2018 data sets. Separate PFNs are trained for nonprompt electrons and muons, and for m_N up to and above 5 GeV. The reason for the split into a low- and high-mass PFN is the strong dependence of some of the input variables for HNL signal events on the m_N scenario, and also heavy-flavor background events are more similar to low-mass than to high-mass HNL events. For the low-mass (high-mass) PFN, the training is performed using an admixture of MC signal samples with HNL masses in the range of 1–5 (6–20) GeV and with different τ_N values.

The PFN training is performed with the KERAS v2.1.5 deep learning library [119] interfaced to TENSORFLOW v1.6 [120]. Parametric rectified linear functions are chosen for the activation function. The dropout parameter is set to a high value of 50%, i.e., a random half of the neurons are dropped from each training iteration to prevent overtraining [121]. We set aside 10% of the training samples as a testing set, which is not used during the training but only for the final validation after the full training is done. Two thirds of the remaining events in the training samples are used for training during an epoch and the other third for validation after the epoch. The PFN is trained for a maximum of 100 epochs, and stopped earlier if the accuracy on the training data does not increase for four full training epochs. This cycle is repeated three times, where a different third of the events is used as the validation set. This K -fold cross-validation method, where each event is used both for training and validation, further reduces the possibility of overtraining [122]. The final model is obtained by taking the




Figure 5. The PFN score distribution of predicted events yields after applying the selection summarized in table 1, for the combined ee and μe (left) or $\mu\mu$ and $e\mu$ (right) categories, using the low-mass (upper) or high-mass (lower) PFNs. The filled histograms display the predicted background yields, where the QCD multijet background displays huge statistical fluctuations for PFN scores above 0.2. The lines show the predicted yields for four HNL signal scenarios, with $m_N = 3$ GeV and $|V_{\ell N}|^2 = 9.9 \times 10^{-5}$ (HNL3), $m_N = 5$ GeV and $|V_{\ell N}|^2 = 1.6 \times 10^{-6}$ (HNL5), $m_N = 6$ GeV and $|V_{\ell N}|^2 = 2.0 \times 10^{-6}$ (HNL6), and $m_N = 10$ GeV and $|V_{\ell N}|^2 = 5.7 \times 10^{-7}$ (HNL10). The HNL signal yield is normalized to the total background yield.

average of the three models with the best accuracy on the validation data of the respective training cycle. The PFN score is shown for simulated HNL and background events in figure 5.

A validation of differences in the shape of the PFN score between data and simulated events is performed to ensure that the signal efficiency is the same in data and simulation. The meson decays $K_S^0 \rightarrow \pi^+\pi^-$ are a representative long-lived decay process that can be used to emulate HNL decays. Following a procedure similar to that described in ref. [123], a pure sample of $Z \rightarrow \mu^+\mu^-$ events is selected, requiring two OS muons with $m(\mu^\pm\mu^\mp)$ within 10 GeV of the Z boson mass [124] to facilitate the data-to-simulation comparison and




Figure 6. Predicted and observed event yields in the PFN score distribution for $K_S^0 \rightarrow \pi^+\pi^-$ decays in $Z \rightarrow \mu^+\mu^-$ events, using the low-mass PFNs for the electron (left) and muon (right) channels, where the $\pi^\pm$ with higher p_T is treated as the lepton. The prediction is scaled to match the overall data yield. The lower panels show the data-to-prediction ratio.

normalization. These events are then required to contain an SV formed by two tracks with an invariant mass within 15 MeV of the K_S^0 meson mass [124]. The K_S^0 meson is typically found within a jet with additional PF candidates. The identification of the SV and jet is performed with the same configuration employed for HNL events. All PF particles in the jet, including those not associated with the K_S^0 meson decay, are used as inputs to the PFN. Since no displaced lepton is present in the K_S^0 meson decay, the $\pi^\pm$ candidate track with the higher p_T is chosen to emulate the displaced lepton for the PFN. In figure 6, the result of this comparison is shown for the low-mass PFNs. The total K_S^0 candidate yield derived from MC samples is normalized to the yield in data, to facilitate the shape comparison of the PFN score. Agreement between the data and the MC prediction within 20% is found over the whole range of the PFN score, both in the electron and muon channels, with an agreement of better than 10% for the last two bins with the highest PFN scores. A similar level of agreement is found for the high-mass PFNs. This validation is used to assign a systematic uncertainty for data-to-simulation differences in the PFN score, as discussed in section 7.

6 Background estimation

After applying all event selection criteria, the main sources of SM background contributions are top quark, W+jets, DY, and QCD multijet production. Their relative contributions vary across the different channels. To further suppress background events and enrich the event sample used for the background estimation with events that have properties similar to expected HNL signal events, all events are required to have $m(\ell_1, \text{SV}) > 10 \text{ GeV}$ and a PFN score of at least 0.2. The selection after these additional requirements will be referred to as the signal region (SR), and is generally different when using low- or high-mass PFNs. Contributions from DY production are found in the ee and $\mu\mu$ categories, where one lepton is

Region	$N(\text{jets})$	PFN score	$m(\ell_1, \text{SV})$ [GeV]
SR-A	=1	$>x$	$>50, <85$
SR-B	=1	$>0.2, <x$	$>50, <85$
SR-C	=1	$>x$	$>10, <50$ or >85
SR-D	=1	$>0.2, <x$	$>10, <50$ or >85
VR-AB	>1	>0.2	$>50, <85$
VR-CD	>1	>0.2	$>10, <50$ or >85
CR-A	≥ 1	$>x$	$>85, <110$
CR-B	≥ 1	$>0.2, <x$	$>85, <110$
CR-C	≥ 1	$>x$	$>10, <50$ or >110
CR-D	≥ 1	$>0.2, <x$	$>10, <50$ or >110

Table 3. Definition of target and sideband regions used in the ABCD background estimation method for the signal (SR), validation (VR), and control (CR) regions. The threshold value x is chosen between 0.97 and 0.998 separately for each event category, as described in the text.

poorly reconstructed and identified as nonprompt, and combined with a random track to form a compatible SV. Background contributions from top quark (W+jets) production typically have one genuine prompt lepton and a second lepton inside a jet that is reconstructed in genuine (accidental) association with an SV. The QCD multijet events contribute especially to the $\mu\mu$ and $e\mu$ categories, and neither the prompt nor nonprompt lepton is genuine. A final background source arises from displaced photon conversions in the detector material and contributes mainly to the ee and μe categories, which have a photon radiated from a prompt lepton and subsequently converted into an e^+e^- pair with one electron receiving most of the momentum.

Background estimates for displaced signatures based on MC simulation are known to be poorly modeled. Thus, we employ a background estimation method using data control samples and rely on the discriminating powers of the $m(\ell_1, \text{SV})$ distribution described in section 4 and the PFN score described in section 5. A target region A is defined by requiring $50 < m(\ell_1, \text{SV}) < 85$ GeV and that the PFN score is above a threshold value x . An ABCD method [125] is then employed to predict the background contribution in region A from the observed yields in three sideband regions B, C, and D, based on inverting one or both of the $m(\ell_1, \text{SV})$ and PFN score requirements. The region definitions are listed in table 3 and shown schematically in figure 7. The expected number of background events in region A, N_A , is then estimated from the observed number of events in the sideband regions as $N_A = N_B N_C / N_D$.

To optimize the values x of the PFN threshold, we consider the signal-over-background ratio in region A as well as the statistical precision of the ABCD background estimate. For the latter, it is important that N_C does not become zero, which would result in an estimate of $N_A = 0$ with a large statistical uncertainty. The result of this optimization are values of $0.97 \leq x \leq 0.998$, chosen separately for each event category. To further increase the sensitivity to the HNL signal, the SR is binned in m_{SV} and the transverse displacement Δ_{2D} between PV and SV. The ABCD background estimation is done separately for each bin in m_{SV} and Δ_{2D} , and separately using the low- and high-mass PFNs.




Figure 7. Illustration of the target and sideband region definitions for the ABCD method applied to the SR, in terms of $N(\text{jets})$, $m(\ell_1, \text{SV})$, and the PFN score.

The ABCD method relies on the absence of correlation between the PFN score and $m(\ell_1, \text{SV})$. To evaluate in data whether this assumption is valid, we form an orthogonal event selection with $N(\text{jets}) > 1$, referred to as validation region (VR). For the first test, we divide the VR into a target region with $50 < m(\ell_1, \text{SV}) < 85 \text{ GeV}$ and a sideband region outside of this interval, as listed in table 3. In figure 8, the PFN scores in data are compared for VR events in the target and sideband regions and, notably, no difference is observed between the shapes.

As further tests, we calculate Pearson correlation coefficients, which measure the strength and direction of the linear relation, to quantify the correlation between $m(\ell_1, \text{SV})$ and the PFN score, as well as p -values to determine the statistical significance of the correlation [126]. This is done separately in flavor channels and $(m_{\text{SV}}, \Delta_{2\text{D}})$ bins, but inclusively in the ABCD classification of the VR. We find correlation coefficients very close to zero in most cases, and also the p -values indicate a weak or negligible linear dependence of the PFN score on $m(\ell_1, \text{SV})$. Only in the OS $\mu\mu$ category of the VR with the low-mass PFN, a relevant statistical dependence is found. We identified in studies with simulated event samples that DY dimuon production is the only background process that exhibits this correlation.

To correct for the bias in the background estimation from the correlation between $m(\ell_1, \text{SV})$ and the PFN score in DY dimuon events, we apply correction factors to the ABCD estimate. The correction factor c_{DY} that needs to be applied to the DY contribution in a bin is calculated as the ratio between the direct DY MC prediction for N_{A} in that bin with the ABCD estimate $N_{\text{B}}N_{\text{C}}/N_{\text{D}}$ evaluated from the same DY MC prediction. Using the fraction f_{DY} of DY MC events relative to the total MC background prediction in that bin, the correction factor to be applied to the ABCD background prediction evaluated from data is $c_{\text{bkg}} = 1 + (c_{\text{DY}} - 1)f_{\text{DY}}$. Each bin of the OS $\mu\mu$ categories when using the low-mass PFNs is corrected separately. The limited size in the DY MC event samples used for the evaluation of c_{bkg} results in a systematic uncertainty, as discussed in section 7.

We evaluate the closure of the ABCD background estimation in data by defining a control region (CR) that does not include the target region SR-A, splitting it into target




Figure 8. The PFN scores in data in the VR, shown for the electron (left) and muon (right) trainings with the low-mass (upper) and high-mass (lower) samples. The distributions are normalized to unity to facilitate a shape comparison. Statistical uncertainties are indicated with error bars and shaded areas. The lower panels show the ratio of the VR-AB to the VR-CD yields.

and sideband regions, and comparing the ABCD background estimate for CR-A with the observed yields. The CR is formed from all events in the SR and VR outside the interval $50 < m(\ell_1, \text{SV}) < 85 \text{ GeV}$, i.e., excluding the target regions of the SR and VR. The inclusion of the VR increases the statistical power of this closure test. The target region of the CR is defined by the requirement $85 < m(\ell_1, \text{SV}) < 110 \text{ GeV}$, and the corresponding sideband definitions are listed in table 3. The comparison between the ABCD estimate for CR-A with the observed yield is done in bins of m_{SV} and $\Delta_{2\text{D}}$, optimized separately for usage with the low- and high-mass PFNs to provide discrimination between signal and background, as well as between different signal hypotheses. For the low-mass analysis, two m_{SV} bins < 2 and $> 2 \text{ GeV}$ are formed, and each is divided into three $\Delta_{2\text{D}}$ bins < 4 , $4\text{--}10$, and $> 10 \text{ cm}$. For the high-mass

Source	Type	Uncertainty [%]
<i>Signal prediction</i>		
NNLO K -factor	Normalization	4
Integrated luminosity	Normalization	1.6
Pileup modeling	Shape	4.6
e (μ) trigger efficiency	Shape	1 (<1)
Prompt e (μ) selection efficiency	Shape	2–4 (1–3)
Nonprompt e (μ) selection efficiency	Shape	1–20 (<1)
Tracking efficiency	Shape	7.3
Jet energy scale & resolution	Shape	1–2
PFN score	Normalization	10
<i>Background prediction</i>		
CR closure	Shape	20–30
DY scale factor (OS $\mu\mu$, low mass)	Shape	20–50

Table 4. Summary of systematic uncertainty sources in the signal and background predictions. Electron- and muon-related uncertainties are listed together, with the values before and inside the parentheses referring to electrons and muons, respectively.

analysis, the two m_{SV} bins are <6 and >6 GeV, each divided by Δ_{2D} as <1 , $1-5$, and >5 cm. For the SS $\mu\mu$ and $e\mu$ channels with the high-mass PFNs, we split the $m_{SV} > 6$ GeV bin into only two Δ_{2D} bins with <1 and >1 cm, to avoid having a $\Delta_{2D} > 5$ cm bin with zero predicted background. The results of the comparison are shown in figures 9–10. A good closure of the background prediction estimate is found within 20–30% when taking the large statistical uncertainty of the data into account. As discussed in section 7, a systematic uncertainty in the background prediction is assigned based on this closure.

7 Systematic uncertainties

The predicted number of background events, as well as the expected number of signal events in the different SR bins for the different event categories, are affected by several sources of systematic uncertainty. Several theoretical and experimental uncertainties affect the HNL signal yields, with some changing only the overall normalization and others also affecting the shape of the signal prediction in the SR bins. For the background yields, two dedicated uncertainties are evaluated that both affect the shape of the prediction. A summary of the different systematic uncertainty sources is given in table 4.

The K -factor applied to the normalization of HNL signal events has an uncertainty of 4%, evaluated by performing variations of the factorization and renormalization scales and of the PDFs in the NNLO W boson production calculation. Additionally, the p_T spectrum of the simulated W boson in SM W+jets production at either NLO or LO accuracy was compared, and all differences were found to be covered by the K -factor uncertainty.




Figure 9. Predicted and observed event yields in the CR for the low-mass PFNs in the SS (upper) and OS (lower) categories, binned by flavor channel, m_{SV} (as specified below the flavor channel), and Δ_{2D} . The hashed bands represent the total uncertainty in the background prediction. The lower panels show the data-to-prediction ratio and the background prediction uncertainty is split into statistical (“stat.”) and systematic (“syst.”) contributions.




Figure 10. Predicted and observed event yields in the CR for the high-mass PFNs in the SS (upper) and OS (lower) categories, binned by flavor channel, m_{SV} (as specified below the flavor channel), and Δ_{2D} . The hashed bands represent the total uncertainty in the background prediction. The lower panels show the data-to-prediction ratio and the background prediction uncertainty is split into statistical (“stat.”) and systematic (“syst.”) contributions.

The integrated luminosity of the three data-taking years is measured with uncertainties between 1.2 and 2.5% [127–129], with a total uncertainty of 1.6% for the combined data set. The modeling uncertainty of pileup interactions is assessed by varying the total pp inelastic cross section in simulation by $\pm 4.6\%$.

The efficiencies of the trigger and prompt-lepton selections are measured in data and simulation with the “tag-and-probe” method applied to $Z \rightarrow \ell^+ \ell^-$ events [130]. Differences between efficiencies in data and simulation are corrected for with scale factors, and the uncertainties in these scale factors are considered for the HNL signal yield. The uncertainties in the trigger scale factors are found to be about 1% (less than 1%) for the single-electron (single-muon) trigger, and in the selection scale factors to be 2–4 (1–3)% for electrons (muons).

The selection efficiency for nonprompt electrons is studied in asymmetric photon conversions events, following the procedure described in ref. [40]. Events with $Z \rightarrow \ell^+ \ell^- \gamma \rightarrow \ell^+ \ell^- e^\pm (e^\mp)$, where $(e^\mp)$ is a low- p_T electron that is not reconstructed in the detector, are selected. Data-to-simulation differences are evaluated as a function of a displacement variable, with different results found for the different years of data taking, corresponding to differences in the material budget and its modeling in the detector simulation. Systematic uncertainties in the range 1–20% are found, largest for small displacements and depending on the data-taking year.

The nonprompt-muon selection efficiency is studied with a tag-and-probe method in $B^\pm \rightarrow J/\psi K^\pm \rightarrow \mu^+ \mu^- K^\pm$ events, where the muon pair from the J/ψ meson decay provides a good proxy to nonprompt muons in HNL decays [40]. The differences between the efficiencies measured in data and simulated events are found to be small, with a systematic uncertainty of less than 1%.

The track and SV reconstruction efficiency is studied using K_S^0 meson decays to two charged particles, which provide an event signature with two displaced tracks originating from a common vertex [123]. We find that reconstruction inefficiencies originate almost entirely from the track reconstruction inefficiency, while the SV reconstruction is almost 100% efficient. Scale factors are applied as a function of p_T and displacement to correct the simulation for differences with data, and we assign half the difference from unity as a systematic uncertainty.

Uncertainties in the momentum scale and resolution of prompt leptons [89, 91] are found to be negligible. Possible discrepancies between data and simulation in the momentum scale and resolution of nonprompt tracks are assessed by comparing the mean values and standard deviations of Gaussian fits to the reconstructed K_S^0 candidate mass, in bins of SV displacement. All discrepancies are found to be below 1% and are neglected.

The systematic uncertainties associated with the jet energy scale and resolution [88] are evaluated by independently scaling the energy of jets up and down by their uncertainty. These variations result in a variation of the HNL signal yield by 1–2%.

The PFN score validation in K_S^0 meson decays, as described in section 5, is used to evaluate a systematic uncertainty that covers differences between data and simulation in the PFN score for HNL events. For high values of the PFN score, discrepancies between data and simulation of up to 10% are observed, consistent for different PFN trainings and data-taking years. Thus, we assign an overall 10% uncertainty in the HNL yield.

The predicted background yields in the target region are subject to statistical uncertainties in the observed event yields in the sideband regions, resulting in uncertainties of up to 50% in the target region yields. Additional systematic uncertainties are assigned, as discussed in section 6 and described in the following. From the agreement between predicted and observed yields in the CR, an uncertainty of 20–30% in the background prediction is considered, separately for each channel. For the additional scale factor applied to the background prediction in the OS $\mu\mu$ low-mass channel, the limited size of the DY MC sample used for the derivation of this scale factor results in an uncertainty of 20–50%, depending on the SR bin.

8 Results

The yields in the four flavor channels are analyzed with both the low- and high-mass PFNs, and binned according to m_{SV} and Δ_{2D} , as defined in section 6. The observed data and the predicted background yields from the ABCD method in these bins are shown in figure 11 (12) for the low-mass (high-mass) PFNs. The number of observed events in data is in agreement with the SM background expectations within the statistical and systematic uncertainties. No significant data excess is found in any final state or SR bin.

For each $(m_N, |V_{\ell N}|^2)$ HNL signal scenario, we evaluate upper limits on the production cross section using the modified frequentist CL_s criterion [131, 132]. A profile likelihood ratio is used as test statistic [133], employing the asymptotic approximation [134] in the limit setting procedure. The binned likelihood is constructed from the observed data yields and the expected signal and background yields in the (m_{SV}, Δ_{2D}) bins. The expected signal yields are estimated from simulation. The ABCD method for the estimation of the background yields discussed in section 6 is implemented in the likelihood similar to the method described in ref. [44]: both the target and the three sideband regions of the SR are included, with possible signal contamination in the sideband regions explicitly considered, background templates constructed from the profit difference between data and expected signal yields in the sideband regions, and nuisance parameters introduced to constrain the background yield in the target region to the ABCD estimate. Further constrained nuisance parameters are introduced for the systematic uncertainties, which include effects on the shapes of the distributions. The binned profile likelihood fits are performed with the CMS statistical analysis tool COMBINE [135], which is based on the ROOFIT [136] and ROOSTATS [137] frameworks.

We derive exclusion limits on $|V_{eN}|^2$, $|V_{\mu N}|^2$, and $|V_{eN}V_{\mu N}|^2/(|V_{eN}|^2 + |V_{\mu N}|^2)$ as functions of m_N , separately for the cases of Majorana and Dirac HNLs, using a grid of points in the $(m_N, |V_{\ell N}|^2)$ parameter space. A signal scenario is excluded if the predicted cross section is incompatible with the observed data at 95% confidence level (CL). The obtained limits are connected with straight lines between neighboring mass points. We consider values of $1 < m_N < 20$ GeV with a step size of 0.5 or 1 GeV. For each mass point, per-event reweighting is employed to interpolate across the entire range of interest in $|V_{\ell N}|^2$, using all available MC samples, as described in section 3. Generated samples with $V_{eN} = V_{\mu N}$ are used for deriving limits in the $e\mu$ and μe channels, but since the HNL production cross section in these channels depends only on $|V_{eN}V_{\mu N}|^2/(|V_{eN}|^2 + |V_{\mu N}|^2)$ [25], the results are applicable also to scenarios with different $|V_{eN}|^2 : |V_{\mu N}|^2$ ratios. The obtained exclusion limits are presented in figure 13.




Figure 12. Predicted and observed SR event yields for the SS (upper) and OS (lower) categories of the high-mass PFNs, binned by flavor channel, m_{SV} (as specified below the flavor channel), and Δ_{2D} . The hashed band represents the total systematic and statistical uncertainty in the background prediction. Signal predictions are shown for two HNL production hypotheses, with $m_N = 6$ GeV and $|V_{\ell N}|^2 = 2.0 \times 10^{-7}$ (HNL6), and $m_N = 10$ GeV and $|V_{\ell N}|^2 = 5.7 \times 10^{-7}$ (HNL10). The lower panels show the data-to-prediction ratio and the background prediction uncertainty is split into statistical (“stat.”) and systematic (“syst.”) contributions.




Figure 13. Exclusion limits at 95% CL on $|V_{eN}|^2$ (upper row), $|V_{\mu N}|^2$ (middle row), and $|V_{eN}V_{\mu N}|^2/(|V_{eN}|^2 + |V_{\mu N}|^2)$ (lower row) as functions of m_N for a Majorana (left) and Dirac (right) HNL. The solid (dashed) black curve indicates the observed (expected) exclusion, where the parameter combinations inside the curve are excluded.

Type	Coupling	m_N [GeV]	95% CL exclusion	
			Lowest value	Highest value
Majorana	$ V_{eN} ^2$	10	9.1×10^{-7}	5.6×10^{-5}
Majorana	$ V_{\mu N} ^2$	14	2.9×10^{-7}	4.8×10^{-6}
Majorana	$ V_{eN}V_{\mu N} ^2/(V_{eN} ^2 + V_{\mu N} ^2)$	12	1.3×10^{-7}	6.7×10^{-6}
Dirac	$ V_{eN} ^2$	11.5	6.9×10^{-7}	6.8×10^{-5}
Dirac	$ V_{\mu N} ^2$	13.5	2.7×10^{-7}	2.1×10^{-5}
Dirac	$ V_{eN}V_{\mu N} ^2/(V_{eN} ^2 + V_{\mu N} ^2)$	12	1.2×10^{-7}	1.8×10^{-5}

Table 5. Comparison of lowest and highest $|V_{\ell N}|^2$ values excluded at 95% CL for Majorana and Dirac HNLs with different coupling scenarios. For each scenario, the m_N value where the lowest $|V_{\ell N}|^2$ value is excluded is shown.

The exclusion limits exhibit both an upper and a lower curve, corresponding to HNL scenarios with short and long lifetimes. The upper curves are evaluated only for $m_N \geq 8$ GeV; the prompt scenarios with lower m_N and high $|V_{\ell N}|^2$ values are not well described by the available HNL signal samples and are already excluded by previous searches, e.g., by the results of the DELPHI Collaboration [50]. The results of this study extend existing exclusion limits into the parameter space characterized by longer lifetimes and, correspondingly, smaller $|V_{\ell N}|^2$ values. In table 5, the observed limits are summarized for those mass points where the lowest values of $|V_{\ell N}|^2$ are excluded per scenario.

Other experiments and searches have established more stringent limits for some of the mass range probed in this analysis. The CHARM experiment [56] has excluded electron (muon) couplings for $1.0 < m_N < 2.1$ (1.9) GeV, with $|V_{\ell N}|^2$ values as low as 10^{-7} . The previous CMS search for HNLs decaying in the muon system [45] found more stringent limits in the mass range 1–2.8 GeV, with electron (muon) couplings excluded with $|V_{\ell N}|^2$ values as low as 0.9 (0.5) $\times 10^{-6}$. The displaced trilepton search [40] and the displaced-jet-tagger analysis in the same final state [44] probe similar m_N and $|V_{\ell N}|^2$ regions as this work. In the mass range 2–10 GeV, the two searches exhibit sensitivities that are comparable to our results. For masses of 10–17 GeV and $|V_{\ell N}|^2$ values below 10^{-5} , the requirement of an SV in this analysis results in the removal of a significant amount of background from the SRs, resulting in more stringent limits when compared to ref. [44]. For masses above 10 GeV and higher coupling values (i.e., shorter lifetime), however, the displaced-jet-tagger approach shows a better sensitivity. Compared to the recent prompt trilepton search [46], the limits on $|V_{eN}|^2$ and $|V_{\mu N}|^2$ presented here are more stringent in the probed mass range by more than one order of magnitude.

9 Summary

A search for long-lived heavy neutral leptons (HNLs) has been presented using proton-proton collision events with one prompt lepton and a system of a nonprompt lepton and a jet associated with a secondary vertex. The data set corresponds to 138 fb^{-1} and was collected by the CMS experiment at the LHC in 2016–2018. Secondary vertices are reconstructed and

matched with nonprompt leptons to find candidate vertices for the HNL decay, and a dedicated machine-learning method is utilized to distinguish between secondary vertices associated with HNL decays and those from background sources. No excess of events above the standard model background prediction obtained from control samples in data is found. Exclusion limits at 95% confidence level are evaluated for different HNL coupling scenarios as functions of the HNL mass and the mixing parameter with standard model neutrinos. The obtained exclusion limits cover HNL masses from 1 to 16.5 GeV and squared mixing parameters as low as 2×10^{-7} , depending on the scenario. These results exceed previous experimental constraints derived in the single-lepton decay channel in the mass range 11–16.5 GeV. For some of the considered coupling scenarios and mass ranges, the presented limits are the strongest to date.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia);

the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC — National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR — Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Data Availability Statement. Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use, and open access policy](#).

Code Availability Statement. The CMS core software is publicly available on [GitHub](#).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] SUPER-KAMIOKANDE collaboration, *Evidence for oscillation of atmospheric neutrinos*, *Phys. Rev. Lett.* **81** (1998) 1562 [[hep-ex/9807003](#)] [[INSPIRE](#)].
- [2] SNO collaboration, *Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory*, *Phys. Rev. Lett.* **89** (2002) 011301 [[nucl-ex/0204008](#)] [[INSPIRE](#)].
- [3] KAMLAND collaboration, *First results from KamLAND: evidence for reactor anti-neutrino disappearance*, *Phys. Rev. Lett.* **90** (2003) 021802 [[hep-ex/0212021](#)] [[INSPIRE](#)].
- [4] S. Bilenky, *Neutrino oscillations: from a historical perspective to the present status*, *Nucl. Phys. B* **908** (2016) 2 [[arXiv:1602.00170](#)] [[INSPIRE](#)].
- [5] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].

- [6] EBOSS collaboration, *Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory*, *Phys. Rev. D* **103** (2021) 083533 [[arXiv:2007.08991](#)] [[INSPIRE](#)].
- [7] Z. Sakr, *A short review on the latest neutrinos mass and number constraints from cosmological observables*, *Universe* **8** (2022) 284 [[INSPIRE](#)].
- [8] J.A. Formaggio, A.L.C. de Gouvêa and R.G.H. Robertson, *Direct measurements of neutrino mass*, *Phys. Rept.* **914** (2021) 1 [[arXiv:2102.00594](#)] [[INSPIRE](#)].
- [9] KATRIN collaboration, *Direct neutrino-mass measurement with sub-electronvolt sensitivity*, *Nature Phys.* **18** (2022) 160 [[arXiv:2105.08533](#)] [[INSPIRE](#)].
- [10] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?, *Phys. Lett. B* **67** (1977) 421 [[INSPIRE](#)].
- [11] T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, in *Proc. Workshop on the Unified Theories and the Baryon Number in the Universe*, Tsukuba, Japan, 13–14 February 1979 [[DOI:10.1142/9789812702210_0019](#)].
- [12] M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, *Conf. Proc. C* **790927** (1979) 315 [[arXiv:1306.4669](#)] [[INSPIRE](#)].
- [13] S.L. Glashow, *The future of elementary particle physics*, *NATO Sci. Ser. B* **61** (1980) 687 [[INSPIRE](#)].
- [14] R.N. Mohapatra and G. Senjanović, *Neutrino mass and spontaneous parity nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].
- [15] J. Schechter and J.W.F. Valle, *Neutrino masses in $SU(2) \times U(1)$ theories*, *Phys. Rev. D* **22** (1980) 2227 [[INSPIRE](#)].
- [16] R.E. Shrock, *General theory of weak leptonic and semileptonic decays. 1. Leptonic pseudoscalar meson decays, with associated tests for, and bounds on, neutrino masses and lepton mixing*, *Phys. Rev. D* **24** (1981) 1232 [[INSPIRE](#)].
- [17] Y. Cai, T. Han, T. Li and R. Ruiz, *Lepton number violation: seesaw models and their collider tests*, *Front. in Phys.* **6** (2018) 40 [[arXiv:1711.02180](#)] [[INSPIRE](#)].
- [18] S. Dodelson and L.M. Widrow, *Sterile-neutrinos as dark matter*, *Phys. Rev. Lett.* **72** (1994) 17 [[hep-ph/9303287](#)] [[INSPIRE](#)].
- [19] A. Boyarsky et al., *Sterile neutrino dark matter*, *Prog. Part. Nucl. Phys.* **104** (2019) 1 [[arXiv:1807.07938](#)] [[INSPIRE](#)].
- [20] M. Fukugita and T. Yanagida, *Baryogenesis without grand unification*, *Phys. Lett. B* **174** (1986) 45 [[INSPIRE](#)].
- [21] E.J. Chun et al., *Probing leptogenesis*, *Int. J. Mod. Phys. A* **33** (2018) 1842005 [[arXiv:1711.02865](#)] [[INSPIRE](#)].
- [22] M. Drewes, Y. Georis and J. Klarić, *Mapping the viable parameter space for testable leptogenesis*, *Phys. Rev. Lett.* **128** (2022) 051801 [[arXiv:2106.16226](#)] [[INSPIRE](#)].
- [23] T. Asaka, S. Blanchet and M. Shaposhnikov, *The ν MSM, dark matter and neutrino masses*, *Phys. Lett. B* **631** (2005) 151 [[hep-ph/0503065](#)] [[INSPIRE](#)].
- [24] F. del Aguila and J.A. Aguilar-Saavedra, *Distinguishing seesaw models at LHC with multi-lepton signals*, *Nucl. Phys. B* **813** (2009) 22 [[arXiv:0808.2468](#)] [[INSPIRE](#)].

- [25] A. Atre, T. Han, S. Pascoli and B. Zhang, *The search for heavy Majorana neutrinos*, *JHEP* **05** (2009) 030 [[arXiv:0901.3589](#)] [[INSPIRE](#)].
- [26] V. Tello et al., *Left-right symmetry: from LHC to neutrinoless double beta decay*, *Phys. Rev. Lett.* **106** (2011) 151801 [[arXiv:1011.3522](#)] [[INSPIRE](#)].
- [27] F.F. Deppisch, P.S. Bhupal Dev and A. Pilaftsis, *Neutrinos and collider physics*, *New J. Phys.* **17** (2015) 075019 [[arXiv:1502.06541](#)] [[INSPIRE](#)].
- [28] S. Pascoli, R. Ruiz and C. Weiland, *Heavy neutrinos with dynamic jet vetoes: multilepton searches at $\sqrt{s} = 14, 27$, and 100 TeV*, *JHEP* **06** (2019) 049 [[arXiv:1812.08750](#)] [[INSPIRE](#)].
- [29] J. Alimena et al., *Searching for long-lived particles beyond the Standard Model at the Large Hadron Collider*, *J. Phys. G* **47** (2020) 090501 [[arXiv:1903.04497](#)] [[INSPIRE](#)].
- [30] A.M. Abdullahi et al., *The present and future status of heavy neutral leptons*, *J. Phys. G* **50** (2023) 020501 [[arXiv:2203.08039](#)] [[INSPIRE](#)].
- [31] C. Antel et al., *Feebly-interacting particles: FIPs 2022 workshop report*, *Eur. Phys. J. C* **83** (2023) 1122 [[arXiv:2305.01715](#)] [[INSPIRE](#)].
- [32] CMS collaboration, *Search for heavy Majorana neutrinos in $\mu^\pm\mu^\pm + \text{jets}$ and $e^\pm e^\pm + \text{jets}$ events in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **717** (2012) 109 [[arXiv:1207.6079](#)] [[INSPIRE](#)].
- [33] CMS collaboration, *Search for heavy Majorana neutrinos in $\mu^\pm\mu^\pm + \text{jets}$ events in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett. B* **748** (2015) 144 [[arXiv:1501.05566](#)] [[INSPIRE](#)].
- [34] ATLAS collaboration, *Search for heavy Majorana neutrinos with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **07** (2015) 162 [[arXiv:1506.06020](#)] [[INSPIRE](#)].
- [35] CMS collaboration, *Search for heavy Majorana neutrinos in $e^\pm e^\pm + \text{jets}$ and $e^\pm\mu^\pm + \text{jets}$ events in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **04** (2016) 169 [[arXiv:1603.02248](#)] [[INSPIRE](#)].
- [36] CMS collaboration, *Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **120** (2018) 221801 [[arXiv:1802.02965](#)] [[INSPIRE](#)].
- [37] CMS collaboration, *Search for heavy Majorana neutrinos in same-sign dilepton channels in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **01** (2019) 122 [[arXiv:1806.10905](#)] [[INSPIRE](#)].
- [38] ATLAS collaboration, *Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS detector*, *JHEP* **10** (2019) 265 [[arXiv:1905.09787](#)] [[INSPIRE](#)].
- [39] LHCb collaboration, *Search for heavy neutral leptons in $W^+ \rightarrow \mu^+ \mu^\pm \text{jet}$ decays*, *Eur. Phys. J. C* **81** (2021) 248 [[arXiv:2011.05263](#)] [[INSPIRE](#)].
- [40] CMS collaboration, *Search for long-lived heavy neutral leptons with displaced vertices in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2022) 081 [[arXiv:2201.05578](#)] [[INSPIRE](#)].
- [41] ATLAS collaboration, *Search for heavy neutral leptons in decays of W bosons using a dilepton displaced vertex in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Phys. Rev. Lett.* **131** (2023) 061803 [[arXiv:2204.11988](#)] [[INSPIRE](#)].
- [42] CMS collaboration, *Probing heavy Majorana neutrinos and the Weinberg operator through vector boson fusion processes in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **131** (2023) 011803 [[arXiv:2206.08956](#)] [[INSPIRE](#)].
- [43] ATLAS collaboration, *Search for Majorana neutrinos in same-sign WW scattering events from pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **83** (2023) 824 [[arXiv:2305.14931](#)] [[INSPIRE](#)].

- [44] CMS collaboration, *Search for long-lived heavy neutral leptons with lepton flavour conserving or violating decays to a jet and a charged lepton*, *JHEP* **03** (2024) 105 [[arXiv:2312.07484](#)] [[INSPIRE](#)].
- [45] CMS collaboration, *Search for long-lived heavy neutral leptons decaying in the CMS muon detectors in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **110** (2024) 012004 [[arXiv:2402.18658](#)] [[INSPIRE](#)].
- [46] CMS collaboration, *Search for heavy neutral leptons in final states with electrons, muons, and hadronically decaying tau leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2024) 123 [[arXiv:2403.00100](#)] [[INSPIRE](#)].
- [47] CMS collaboration, *Search for long-lived heavy neutrinos in the decays of B mesons produced in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2024) 183 [[arXiv:2403.04584](#)] [[INSPIRE](#)].
- [48] ATLAS collaboration, *Search for heavy Majorana neutrinos in $e^{\pm}e^{\pm}$ and $e^{\pm}\mu^{\pm}$ final states via WW scattering in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **856** (2024) 138865 [[arXiv:2403.15016](#)] [[INSPIRE](#)].
- [49] CMS collaboration, *Review of searches for vector-like quarks, vector-like leptons, and heavy neutral leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV at the CMS experiment*, [arXiv:2405.17605](#) [[INSPIRE](#)].
- [50] DELPHI collaboration, *Search for neutral heavy leptons produced in Z decays*, *Z. Phys. C* **74** (1997) 57 [Erratum *ibid.* **75** (1997) 580] [[INSPIRE](#)].
- [51] BELLE collaboration, *Search for heavy neutrinos at Belle*, *Phys. Rev. D* **87** (2013) 071102 [Erratum *ibid.* **95** (2017) 099903] [[arXiv:1301.1105](#)] [[INSPIRE](#)].
- [52] BABAR collaboration, *Search for heavy neutral leptons using tau lepton decays at BaBar*, *Phys. Rev. D* **107** (2023) 052009 [[arXiv:2207.09575](#)] [[INSPIRE](#)].
- [53] BELLE collaboration, *Search for a heavy neutrino in τ decays at Belle*, *Phys. Rev. Lett.* **131** (2023) 211802 [[arXiv:2212.10095](#)] [[INSPIRE](#)].
- [54] BELLE collaboration, *Search for a heavy neutral lepton that mixes predominantly with the tau neutrino*, *Phys. Rev. D* **109** (2024) L111102 [[arXiv:2402.02580](#)] [[INSPIRE](#)].
- [55] WA66 collaboration, *Search for heavy neutrino decays in the BEBC beam dump experiment*, *Phys. Lett. B* **160** (1985) 207 [[INSPIRE](#)].
- [56] CHARM collaboration, *A search for decays of heavy neutrinos in the mass range 0.5 GeV to 2.8 GeV*, *Phys. Lett. B* **166** (1986) 473 [[INSPIRE](#)].
- [57] NA3 collaboration, *Mass and lifetime limits on new longlived particles in 300 GeV/c π^{-} interactions*, *Z. Phys. C* **31** (1986) 21 [[INSPIRE](#)].
- [58] CHARM II collaboration, *Search for heavy isosinglet neutrinos*, *Phys. Lett. B* **343** (1995) 453 [[INSPIRE](#)].
- [59] NuTeV and E815 collaborations, *Search for neutral heavy leptons in a high-energy neutrino beam*, *Phys. Rev. Lett.* **83** (1999) 4943 [[hep-ex/9908011](#)] [[INSPIRE](#)].
- [60] T2K collaboration, *Search for heavy neutrinos with the T2K near detector ND280*, *Phys. Rev. D* **100** (2019) 052006 [[arXiv:1902.07598](#)] [[INSPIRE](#)].
- [61] NA62 collaboration, *Search for heavy neutral lepton production in K^{+} decays to positrons*, *Phys. Lett. B* **807** (2020) 135599 [[arXiv:2005.09575](#)] [[INSPIRE](#)].
- [62] NA62 collaboration, *Search for K^{+} decays to a muon and invisible particles*, *Phys. Lett. B* **816** (2021) 136259 [[arXiv:2101.12304](#)] [[INSPIRE](#)].

- [63] ARGONEUT collaboration, *New constraints on tau-coupled heavy neutral leptons with masses $m_N = 280\text{--}970\text{ MeV}$* , *Phys. Rev. Lett.* **127** (2021) 121801 [[arXiv:2106.13684](#)] [[INSPIRE](#)].
- [64] MICROBOONE collaboration, *Search for long-lived heavy neutral leptons and Higgs portal scalars decaying in the MicroBooNE detector*, *Phys. Rev. D* **106** (2022) 092006 [[arXiv:2207.03840](#)] [[INSPIRE](#)].
- [65] MICROBOONE collaboration, *Search for heavy neutral leptons in electron-positron and neutral-pion final states with the MicroBooNE detector*, *Phys. Rev. Lett.* **132** (2024) 041801 [[arXiv:2310.07660](#)] [[INSPIRE](#)].
- [66] A. Abada, N. Bernal, M. Losada and X. Marcano, *Inclusive displaced vertex searches for heavy neutral leptons at the LHC*, *JHEP* **01** (2019) 093 [[arXiv:1807.10024](#)] [[INSPIRE](#)].
- [67] J.-L. Tastet, O. Ruchayskiy and I. Timiryasov, *Reinterpreting the ATLAS bounds on heavy neutral leptons in a realistic neutrino oscillation model*, *JHEP* **12** (2021) 182 [[arXiv:2107.12980](#)] [[INSPIRE](#)].
- [68] I. Boiarska, A. Boyarsky, O. Mikulenko and M. Ovchinnikov, *Constraints from the CHARM experiment on heavy neutral leptons with tau mixing*, *Phys. Rev. D* **104** (2021) 095019 [[arXiv:2107.14685](#)] [[INSPIRE](#)].
- [69] A. Abada, P. Escribano, X. Marcano and G. Piazza, *Collider searches for heavy neutral leptons: beyond simplified scenarios*, *Eur. Phys. J. C* **82** (2022) 1030 [[arXiv:2208.13882](#)] [[INSPIRE](#)].
- [70] *HEPData record for this analysis*, <https://doi.org/10.17182/hepdata.150534>, (2024).
- [71] CMS collaboration, *The CMS experiment at the CERN LHC*, 2008 *JINST* **3** S08004 [[INSPIRE](#)].
- [72] CMS collaboration, *Development of the CMS detector for the CERN LHC run 3*, 2024 *JINST* **19** P05064 [[arXiv:2309.05466](#)] [[INSPIRE](#)].
- [73] CMS collaboration, *Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13\text{ TeV}$* , 2020 *JINST* **15** P10017 [[arXiv:2006.10165](#)] [[INSPIRE](#)].
- [74] CMS collaboration, *The CMS trigger system*, 2017 *JINST* **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [75] CMS collaboration, *Performance of the CMS high-level trigger during LHC run 2*, 2024 *JINST* **19** P11021 [[arXiv:2410.17038](#)] [[INSPIRE](#)].
- [76] CMS TRACKER GROUP collaboration, *The CMS phase-1 pixel detector upgrade*, 2021 *JINST* **16** P02027 [[arXiv:2012.14304](#)] [[INSPIRE](#)].
- [77] CMS collaboration, *2017 tracking performance plots*, CMS-DP-2017-015, CERN, Geneva, Switzerland (2017).
- [78] R. Frühwirth, W. Waltenberger and P. Vanlaer, *Adaptive vertex fitting*, *J. Phys. G* **34** (2007) N343 [[INSPIRE](#)].
- [79] D. Contardo et al., *Technical proposal for the phase-II upgrade of the CMS detector*, CERN-LHCC-2015-010, CERN, Geneva, Switzerland (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)] [[INSPIRE](#)].
- [80] CMS collaboration, *Description and performance of track and primary-vertex reconstruction with the CMS tracker*, 2014 *JINST* **9** P10009 [[arXiv:1405.6569](#)] [[INSPIRE](#)].
- [81] CMS collaboration, *Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS phase-1 pixel detector*, CMS-DP-2020-049, CERN, Geneva, Switzerland (2020).

- [82] CMS collaboration, *Measurement of $B\bar{B}$ angular correlations based on secondary vertex reconstruction at $\sqrt{s} = 7$ TeV*, *JHEP* **03** (2011) 136 [[arXiv:1102.3194](#)] [[INSPIRE](#)].
- [83] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, 2018 *JINST* **13** P05011 [[arXiv:1712.07158](#)] [[INSPIRE](#)].
- [84] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, 2017 *JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [85] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [86] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [87] CMS collaboration, *Pileup mitigation at CMS in 13 TeV data*, 2020 *JINST* **15** P09018 [[arXiv:2003.00503](#)] [[INSPIRE](#)].
- [88] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, 2017 *JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [89] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, 2021 *JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [90] CMS collaboration, *ECAL 2016 refined calibration and Run2 summary plots*, CMS-DP-2020-021, CERN, Geneva, Switzerland (2020).
- [91] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [92] CMS collaboration, *Single and double electron trigger efficiencies using the full run 2 dataset*, CMS-DP-2020-016, CERN, Geneva, Switzerland (2020).
- [93] CMS collaboration, *Performance of the CMS muon trigger system in proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2021 *JINST* **16** P07001 [[arXiv:2102.04790](#)] [[INSPIRE](#)].
- [94] NNPDF collaboration, *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [95] NNPDF collaboration, *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[arXiv:1706.00428](#)] [[INSPIRE](#)].
- [96] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [97] CMS collaboration, *Event generator tunes obtained from underlying event and multiparton scattering measurements*, *Eur. Phys. J. C* **76** (2016) 155 [[arXiv:1512.00815](#)] [[INSPIRE](#)].
- [98] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, *Eur. Phys. J. C* **80** (2020) 4 [[arXiv:1903.12179](#)] [[INSPIRE](#)].
- [99] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [100] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [101] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].

- [102] S. Frixione, P. Nason and G. Ridolfi, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126 [[arXiv:0707.3088](#)] [[INSPIRE](#)].
- [103] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [104] S. Alioli, P. Nason, C. Oleari and E. Re, *NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*, *JHEP* **09** (2009) 111 [Erratum *ibid.* **02** (2010) 011] [[arXiv:0907.4076](#)] [[INSPIRE](#)].
- [105] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [106] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547 [[arXiv:1009.2450](#)] [[INSPIRE](#)].
- [107] T. Melia, P. Nason, R. Rötsch and G. Zanderighi, *W^+W^- , WZ and ZZ production in the POWHEG BOX*, *JHEP* **11** (2011) 078 [[arXiv:1107.5051](#)] [[INSPIRE](#)].
- [108] P. Nason and G. Zanderighi, *W^+W^- , WZ and ZZ production in the POWHEG-BOX-V2*, *Eur. Phys. J. C* **74** (2014) 2702 [[arXiv:1311.1365](#)] [[INSPIRE](#)].
- [109] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, *Eur. Phys. J. C* **53** (2008) 473 [[arXiv:0706.2569](#)] [[INSPIRE](#)].
- [110] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061 [[arXiv:1209.6215](#)] [[INSPIRE](#)].
- [111] D. Alva, T. Han and R. Ruiz, *Heavy Majorana neutrinos from $W\gamma$ fusion at hadron colliders*, *JHEP* **02** (2015) 072 [[arXiv:1411.7305](#)] [[INSPIRE](#)].
- [112] C. Degrande, O. Mattelaer, R. Ruiz and J. Turner, *Fully-automated precision predictions for heavy neutrino production mechanisms at hadron colliders*, *Phys. Rev. D* **94** (2016) 053002 [[arXiv:1602.06957](#)] [[INSPIRE](#)].
- [113] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $O(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017 [[hep-ph/0609070](#)] [[INSPIRE](#)].
- [114] R. Gavin, Y. Li, F. Petriello and S. Quackenbush, *FEWZ 2.0: a code for hadronic Z production at next-to-next-to-leading order*, *Comput. Phys. Commun.* **182** (2011) 2388 [[arXiv:1011.3540](#)] [[INSPIRE](#)].
- [115] R. Gavin, Y. Li, F. Petriello and S. Quackenbush, *W physics at the LHC with FEWZ 2.1*, *Comput. Phys. Commun.* **184** (2013) 208 [[arXiv:1201.5896](#)] [[INSPIRE](#)].
- [116] Y. Li and F. Petriello, *Combining QCD and electroweak corrections to dilepton production in FEWZ*, *Phys. Rev. D* **86** (2012) 094034 [[arXiv:1208.5967](#)] [[INSPIRE](#)].
- [117] P.T. Komiske, E.M. Metodiev and J. Thaler, *Energy flow networks: deep sets for particle jets*, *JHEP* **01** (2019) 121 [[arXiv:1810.05165](#)] [[INSPIRE](#)].
- [118] M. Zaheer et al., *Deep sets*, in *Proc. 31st Conference on Neural Information Processing Systems (NIPS 2017)*, Long Beach, CA, U.S.A., 4–9 December 2017, p. 3391 [[arXiv:1703.06114](#)] [[INSPIRE](#)].
- [119] F. Chollet et al., *KERAS*, <https://keras.io>, (2015).
- [120] M. Abadi et al., *TENSORFLOW: large-scale machine learning on heterogeneous systems*, <http://tensorflow.org>, (2015) [[DOI:10.5281/zenodo.4724125](#)].

- [121] N. Srivastava et al., *Dropout: a simple way to prevent neural networks from overfitting*, *J. Machine Learning Res.* **15** (2014) 1929 [INSPIRE].
- [122] T. Hastie, R. Tibshirani and J. Friedman, *Model assessment and selection*, in *The elements of statistical learning*, Springer, New York, NY, U.S.A. (2008), p. 219–259 [DOI:10.1007/978-0-387-84858-7_7].
- [123] CMS collaboration, *Displaced tracking and vertexing calibration using neutral K mesons*, CMS-DP-2024-010, CERN, Geneva, Switzerland (2024).
- [124] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2022** (2022) 083C01 [INSPIRE].
- [125] S. Choi and H. Oh, *Improved extrapolation methods of data-driven background estimations in high energy physics*, *Eur. Phys. J. C* **81** (2021) 643 [arXiv:1906.10831] [INSPIRE].
- [126] B. Rémillard, *Tests of independence*, in *International encyclopedia of statistical science*, Springer, Berlin, Heidelberg, Germany (2011), p. 1598–1601 [DOI:10.1007/978-3-642-04898-2_592].
- [127] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [arXiv:2104.01927] [INSPIRE].
- [128] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV*, CMS-PAS-LUM-17-004, CERN, Geneva, Switzerland (2018).
- [129] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, CMS-PAS-LUM-18-002, CERN, Geneva, Switzerland (2019).
- [130] CMS collaboration, *Measurements of inclusive W and Z cross sections in pp collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **01** (2011) 080 [arXiv:1012.2466] [INSPIRE].
- [131] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [hep-ex/9902006] [INSPIRE].
- [132] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [INSPIRE].
- [133] ATLAS, CMS collaborations and LHC Higgs Combination Group, *Procedure for the LHC Higgs boson search combination in Summer 2011*, CMS-NOTE-2011-005, CERN, Geneva, Switzerland (2011).
- [134] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [arXiv:1007.1727] [INSPIRE].
- [135] CMS collaboration, *The CMS statistical analysis and combination tool: COMBINE*, *Comput. Softw. Big Sci.* **8** (2024) 19 [arXiv:2404.06614] [INSPIRE].
- [136] W. Verkerke and D.P. Kirkby, *The RooFit toolkit for data modeling*, *eConf C* **0303241** (2003) MOLT007 [physics/0306116] [INSPIRE].
- [137] L. Moneta et al., *The RooStats project*, *PoS ACAT2010* (2010) 057 [arXiv:1009.1003] [INSPIRE].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
P.S. Hussain , M. Jeitler ², N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck ²,
R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz ²

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen ,
S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat ³, G.P. Van Onsem ,
S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios ,
J. Jaramillo , A. Khalilzadeh , F.A. Khan , K. Lee , M. Mahdavihorrani , A. Malara ,
S. Paredes , M.A. Shahzad , L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx ,
K. Mota Amarilo , A. Samalan , K. Skovpen , N. Van Den Bossche , J. van der Linden ,
L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere ,
I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre , J. Lidrych ,
P. Mastrapasqua , T.T. Tran , S. Wertz 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Alves Gallo Pereira , E. Coelho , G. Correia Silva , C. Hensel ,
T. Menezes De Oliveira , C. Mora Herrera ⁴, A. Moraes , P. Rebello Teles , M. Soeiro ,
A. Vilela Pereira ⁴

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho ,
J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira ⁶, D. De Jesus Damiao ,
S. Fonseca De Souza , R. Gomes De Souza , M. Macedo , J. Martins ⁷, L. Mundim ,
H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁶, L. Calligaris, T.R. Fernandez Perez Tomei, E.M. Gregores,
B. Lopes Da Costa, I. Maietto Silverio, P.G. Mercadante, S.F. Novaes, B. Orzari,
Sandra S. Padula

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov, G. Antchev, R. Hadjiiska, P. Iaydjiev, M. Misheva, M. Shopova,
G. Sultanov

University of Sofia, Sofia, Bulgaria

A. Dimitrov, L. Litov, B. Pavlov, P. Petkov, A. Petrov, E. Shumka

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri, D. Laroze, S. Thakur

Beihang University, Beijing, China

T. Cheng, T. Javaid, L. Yuan

Department of Physics, Tsinghua University, Beijing, China

Z. Hu, Z. Liang, J. Liu, K. Yi^{8,9}

Institute of High Energy Physics, Beijing, China

G.M. Chen¹⁰, H.S. Chen¹⁰, M. Chen¹⁰, F. Iemmi, C.H. Jiang, A. Kapoor¹¹, H. Liao,
Z.-A. Liu¹², R. Sharma¹³, J.N. Song¹², J. Tao, C. Wang¹⁰, J. Wang, Z. Wang¹⁰, H. Zhang,
J. Zhao

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos, Y. Ban, S. Deng, B. Guo, C. Jiang, A. Levin, C. Li, Q. Li, Y. Mao,
S. Qian, S.J. Qian, X. Qin, X. Sun, D. Wang, H. Yang, L. Zhang, Y. Zhao, C. Zhou

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang

Sun Yat-Sen University, Guangzhou, China

Z. You

University of Science and Technology of China, Hefei, China

K. Jaffel, N. Lu

Nanjing Normal University, Nanjing, China

G. Bauer¹⁴, B. Li, J. Zhang

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China

X. Gao ¹⁵

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

F. Ramirez , C. Rendón, M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic, T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, S. Mishra , A. Starodumov ¹⁶, T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

Y. Assran^{17,18}, B. El-mahdy, S. Elgammal¹⁸

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M.A. Mahmoud , Y. Mohammed 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew ,
K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , L. Martikainen ,
M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour ,
P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi ,
L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro ¹⁹, P. Simkina , M. Titov ,
M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , F. Damas ,
O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert ,
R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen ,
C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza ,
A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ²⁰, J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert ,
C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet ,
G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,
P. Depasse , C. Dozen ²¹, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg,
G. Grenier , B. Ille , E. Jourdhuy, I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries,
A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

D. Chokheli , I. Lomidze , Z. Tsamalaidze ¹⁶

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,
A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann ,
M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung 

M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo ²², S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman ²³, K. Borras ²⁴, A. Campbell , A. Cardini , C. Cheng, F. Colombina , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ²³, A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁵, W. Lohmann ²⁶, F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otari, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham ^{27,24}, S. Schnake ²⁴, P. Schütze , C. Schwanenberger ²³, D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁴, C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ²⁸, R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , J.M. Lawhorn , M. Link, A. Lintuluoto , B. Maier , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Papadopoulos²⁸, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath ^{29,30}, K. Márton, A.J. Rádl ³¹, F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkúti ³², M.M.A. Gadallah ³³, Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Novak 

Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhirgra ³⁴, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , K. Sandeep , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁵, S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri ³⁶, D. Kumar ³⁶, P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh, P. Verma, S. Verma , A. Vijay

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , B. Parida , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar ,
G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati ³⁷, C. Kar , D. Maity ³⁸, P. Mal , T. Mishra ,
V.K. Muraleedharan Nair Bindhu ³⁸, K. Naskar ³⁸, A. Nayak ³⁸, S. Nayak, K. Pal, P. Sadangi,
S.K. Swain , S. Varghese ³⁸, D. Vats ³⁸

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya ³⁹, A. Alpana , S. Dube , B. Gomber ³⁹, P. Hazarika , B. Kansal , A. Laha ,
B. Sahu ³⁹, S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi ⁴⁰, A. Jafari ⁴¹, M. Zeinali ⁴²

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani ⁴³, S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie ⁴⁴,
M. Mohammadi Najafabadi , S. Tizchang ⁴⁵

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia ^{a,b}, A. Colaleo ^{a,b}, D. Creanza ^{a,c}, B. D'Anzi ^{a,b}, N. De Filippis ^{a,c},
M. De Palma ^{a,b}, W. Elmetenawee ^{a,b,46}, L. Fiore ^a, G. Iaselli ^{a,c}, L. Longo ^a, M. Louka ^{a,b},
G. Maggi ^{a,c}, M. Maggi ^a, I. Margjeka ^a, V. Mastrapasqua ^{a,b}, S. My ^{a,b}, S. Nuzzo ^{a,b},
A. Pellecchia ^{a,b}, A. Pompili ^{a,b}, G. Pugliese ^{a,c}, R. Radogna ^{a,b}, D. Ramos ^a, A. Ranieri ^a,
L. Silvestris ^a, F.M. Simone ^{a,c}, Ü. Sözbilir ^a, A. Stamerra ^{a,b}, D. Troiano ^{a,b}, R. Venditti ^{a,b},
P. Verwilligen ^a, A. Zaza ^{a,b}

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi ^a, C. Battilana ^{a,b}, D. Bonacorsi ^{a,b}, P. Capiluppi ^{a,b}, A. Castro ^{†,a,b},
F.R. Cavallo ^a, M. Cuffiani ^{a,b}, G.M. Dallavalle ^a, T. Diotallevi ^{a,b}, F. Fabbri ^a,
A. Fanfani ^{a,b}, D. Fasanella ^a, P. Giacomelli ^a, L. Giommi ^{a,b}, C. Grandi ^a, L. Guiducci ^{a,b},
S. Lo Meo ^{a,47}, M. Lorusso ^{a,b}, L. Lunerti ^a, S. Marcellini ^a, G. Masetti ^a, F.L. Navarria ^{a,b},
G. Paggi ^{a,b}, A. Perrotta ^a, F. Primavera ^{a,b}, A.M. Rossi ^{a,b}, S. Rossi Tisbeni ^{a,b},
T. Rovelli ^{a,b}, G.P. Siroli ^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa ^{a,b,48}, A. Di Mattia ^a, A. Lapertosa ^a, R. Potenza ^{a,b}, A. Tricomi ^{a,b,48}, C. Tuve ^{a,b}

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras , G. Barbagli , G. Bardelli , B. Camaiani , A. Cassese , R. Ceccarelli ,
V. Ciulli , C. Civinini , R. D'Alessandro , E. Focardi , T. Kello , G. Latino ,
P. Lenzi , M. Lizzo , M. Meschini , S. Paoletti , A. Papanastassiou^{a,b}, G. Sguazzoni ,
L. Viliani 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola ⁴⁹, D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

P. Chatagnon , F. Ferro , E. Robutti , S. Tosi ^{a,b}

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia , F. Brivio , F. Cetorelli , F. De Guio , M.E. Dinardo , P. Dini ,
S. Gennai , R. Gerosa , A. Ghezzi , P. Govoni , L. Guzzi , M.T. Lucchini ,
M. Malberti , S. Malvezzi , A. Massironi , D. Menasce , L. Moroni , M. Paganoni ,
S. Palluotto , D. Pedrini , A. Perego , B.S. Pinolini^a, G. Pizzati^{a,b}, S. Ragazzi ,
T. Tabarelli de Fatis 

**INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy;
Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale
(SSM)^d, Napoli, Italy**

S. Buontempo , A. Cagnotta , F. Carnevali^{a,b}, N. Cavallo , F. Fabozzi ,
A.O.M. Iorio , L. Lista ^{a,b,50}, P. Paolucci ^{a,28}

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino , P. Azzi , N. Bacchetta ^{a,51}, M. Biasotto ^{a,52}, D. Bisello , P. Bortignon ,
G. Bortolato^{a,b}, A. Bragagnolo , A.C.M. Bulla , R. Carlin , T. Dorigo , S. Fantinel ,
F. Gasparini , U. Gasparini , E. Lusiani , M. Margoni , A.T. Meneguzzo ,
M. Migliorini , J. Pazzini , P. Ronchese , R. Rossin , F. Simonetto , M. Tosi ,
A. Triossi , S. Ventura , M. Zanetti , P. Zotto , A. Zucchetta 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

C. Aimè , A. Braghieri , S. Calzaferri , D. Fiorina , P. Montagna , V. Re ,
C. Riccardi , P. Salvini , I. Vai , P. Vitulo 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal , M.E. Ascioti^{a,b}, G.M. Bilei , C. Carrivale^{a,b}, D. Ciangottini , L. Fanò ,
M. Magherini , V. Mariani , M. Menichelli , F. Moscatelli ^{a,53}, A. Rossi ,
A. Santocchia , D. Spiga , T. Tedeschi 

**INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c,
Pisa, Italy; Università di Siena^d, Siena, Italy**

C.A. Alexe , P. Asenov , P. Azzurri , G. Bagliesi , R. Bhattacharya , L. Bianchini ,
T. Boccali , E. Bossini , D. Bruschini , R. Castaldi , M.A. Ciocci , M. Cipriani 

V. D'Amante ^{a,d}, R. Dell'Orso ^a, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c}, A.C. Marini ^a,
D. Matos Figueiredo ^a, A. Messineo ^{a,b}, M. Musich ^{a,b}, F. Palla ^a, A. Rizzi ^{a,b},
G. Rolandi ^{a,c}, S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, P. Spagnolo ^a,
R. Tenchini ^a, G. Tonelli ^{a,b}, N. Turini ^{a,d}, F. Vaselli ^{a,c}, A. Venturi ^a, P.G. Verдини ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera ^{a,b}, P. Barria ^a, C. Basile ^{a,b}, M. Campana ^{a,b}, F. Cavallari ^a,
L. Cunqueiro Mendez ^{a,b}, D. Del Re ^{a,b}, E. Di Marco ^{a,b}, M. Diemoz ^a, F. Errico ^{a,b},
E. Longo ^{a,b}, J. Mijuskovic ^{a,b}, G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b},
C. Quaranta ^{a,b}, S. Rahatlou ^{a,b}, C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a,
R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, C. Borca ^{a,b}, N. Cartiglia ^a, M. Costa ^{a,b},
R. Covarelli ^{a,b}, N. Demaria ^a, L. Finco ^a, M. Grippio ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a,
F. Luongo ^{a,b}, C. Mariotti ^a, L. Markovic ^{a,b}, S. Maselli ^a, A. Mecca ^{a,b}, L. Menzio ^{a,b},
P. Meridiani ^a, E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a, M.M. Obertino ^{a,b},
G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c}, F. Siviero ^{a,b},
V. Sola ^{a,b}, A. Solano ^{a,b}, A. Staiano ^a, C. Tarricone ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b},
R. White ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^a, G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra ^a, J. Hong ^a, C. Huh ^a, B. Kim ^a, J. Kim ^a, D. Lee ^a, H. Lee ^a, S.W. Lee ^a, C.S. Moon ^a,
Y.D. Oh ^a, M.S. Ryu ^a, S. Sekmen ^a, B. Tae ^a, Y.C. Yang ^a

Department of Mathematics and Physics — GWNu, Gangneung, Korea

M.S. Kim ^a

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak ^a, P. Gwak ^a, H. Kim ^a, D.H. Moon ^a

Hanyang University, Seoul, Korea

E. Asilar ^a, J. Choi ^a, D. Kim ^a, T.J. Kim ^a, J.A. Merlin ^a, Y. Ryou ^a

Korea University, Seoul, Korea

S. Choi ^a, S. Han ^a, B. Hong ^a, K. Lee ^a, K.S. Lee ^a, S. Lee ^a, J. Yoo ^a

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh ^a, S. Yang ^a

Sejong University, Seoul, Korea

H. S. Kim ^a, Y. Kim ^a, S. Lee ^a

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , S. Ko , H. Kwon , H. Lee ,
J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,
I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , C. Munoz Diaz, D. Osite , G. Pikurs, A. Potrebko , M. Seidel ,
D. Sidiropoulos Kontos

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff ⁵⁴, Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo ,
I. Heredia-De La Cruz ⁵⁵, R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, MontenegroI. Bubanja , N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**K. Pozniak , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo, T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff**Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**M. Dordevic , J. Milosevic , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**J. Alcaraz Maestre , Cristina F. Bedoya , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar **Universidad Autónoma de Madrid, Madrid, Spain**J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras ,
I. Gonzalez Caballero , J.R. González Fernández , P. Leguina , E. Palencia Cortezon ,
J. Prado Pico, C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote ,
C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon ,
J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasosa García , R. Lopez Ruiz ,
C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas ,
E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy ⁵⁶, D.D.C. Wickramaratna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna ⁵⁷, K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney ,
A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta ,
E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d’Enterria , A. Dabrowski ,
A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni ,
W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber,
V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban ²⁶, S. Laurila ,
P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman,
A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat ,
M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer ,
M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , M. Rovere , H. Sakulin ,
S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva ,
P. Sphicas ⁵⁸, A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea ,
D. Walter , J. Wanczyk ⁵⁹, J. Wang, S. Wuchterl , P. Zehetner , P. Zejdl , W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶⁰, L. Caminada ⁶⁰, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶⁰, L. Nohte ⁶⁰, T. Rohe 

ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov ⁵⁹, M. Backhaus , G. Bonomelli, A. Calandri , C. Cazzaniga ,
K. Datta , P. De Bryas Dexmiers D’archiac ⁵⁹, A. De Cosa , G. Dissertori , M. Dittmar,
M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer ,
T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani 

L. Marchese , C. Martin Perez , A. Mascellani ⁵⁹, F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , C. Reissel , B. Ristic , F. Riti , R. Seidita , J. Steggemann ⁵⁹, A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶¹, P. Bärtshi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , F. Meng , U. Molinatti , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶², D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ³⁶, S.S. Yu 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , P.s. Chen, Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu ⁶³, E. Eskut , Y. Guler ⁶⁴, E. Gulpinar Guler ⁶⁴, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁶⁵, A. Polatoz , B. Tali ⁶⁶, U.G. Tok , S. Turkcapar , E. Usan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac ⁶⁷

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya ⁶⁸, O. Kaya ⁶⁹, S. Tekten ⁷⁰

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak ^{63,71}, G.G. Dincer ⁶³, Y. Komurcu , S. Sen ⁷²

Istanbul University, Istanbul, Turkey

O. Aydilek ⁷³, B. Hacisahinoglu , I. Hos ⁷⁴, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci ⁶⁶, B. Isildak ⁷⁵, D. Sunar Cerci , T. Yetkin 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw , S. Seif El Nasr-Storey , V.J. Smith , N. Stylianou ⁷⁶, K. Walkingshaw Pass

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball , K.W. Bell , A. Belyaev ⁷⁷, C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis , K. Harder , S. Harper , J. Linacre , K. Manolopoulos , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh , C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller , V. Cacchio , C.A. Carrillo Montoya , G.S. Chahal ⁷⁸, D. Colling , J.S. Dancu , I. Das , P. Dauncey , G. Davies , J. Davies , M. Della Negra , S. Fayer , G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard , G. Iles , C.R. Knight , J. Langford , J. León Holgado , L. Lyons , A.-M. Magnan , S. Mallios , M. Mieskolainen , J. Nash ⁷⁹, M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards , A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage , T. Virdee ²⁸, M. Vojinovic , N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

K. Coldham , J.E. Cole , A. Khan , P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, U.S.A.

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish ⁸⁰, J. Dittmann , K. Hatakeyama , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

Catholic University of America, Washington, DC, U.S.A.

R. Bartek , A. Dominguez , C. Huerta Escamilla , A.E. Simsek , R. Uniyal , A.M. Vargaz Hernandez 

The University of Alabama, Tuscaloosa, Alabama, U.S.A.

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson , C.U. Perez , P. Rumerio ⁸¹, E. Usai , R. Yi 

Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar , C. Cosby , G. De Castro , Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain , I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

Brown University, Providence, Rhode Island, U.S.A.

G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz , J.M. Hogan ⁸², T. Kwon ,
G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , N. Pervan , T. Russell, S. Sagir ⁸³,
F. Simpson , M. Stamenkovic , N. Venkatasubramanian, X. Yan 

University of California, Davis, Davis, California, U.S.A.

S. Abbott , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez ,
M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral ,
G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , Y. Yao , S. Yoo , F. Zhang 

University of California, Los Angeles, California, U.S.A.

M. Bachtis , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko ,
M.A. Iqbal , T. Lam , E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, U.S.A.

R. Clare , J.W. Gary , M. Gordon, G. Hanson , W. Si 

University of California, San Diego, La Jolla, California, U.S.A.

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte ,
L. Giannini , Y. Gu, J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts ,
M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , M. Quinnan ,
B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang ,
A. Yagil 

University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , J. Incandela ,
J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica ,
R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , S. Wang , D. Zhang

California Institute of Technology, Pasadena, California, U.S.A.

S. Bhattacharya , A. Bornheim , O. Cerri, A. Latorre, J. Mao , H.B. Newman ,
G. Reales Gutiérrez, M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , T. Ferguson ,
T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, C. Liu , T. Mudholkar , S. Murthy ,
P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganeli ,
J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner ,
N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, U.S.A.

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan ,
P. Kotamnives, J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom ,
P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick ,
D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira ,
Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer ,
S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig ,
J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijsma ,
B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid ,
K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn ,
J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro ,
C. Pena ⁸⁴, F. Ravera , A. Reinsvold Hall ⁸⁵, L. Ristori , M. Safdari , E. Sexton-Kennedy ,
N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk ,
N.V. Tran , L. Uppegger , E.W. Vaandering , I. Zoi 

University of Florida, Gainesville, Florida, U.S.A.

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, E. Koenig ,
M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher ,
K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi ,
J. Wang 

Florida State University, Tallahassee, Florida, U.S.A.

T. Adams , A. Al Kadhim , A. Askew , S. Bower , V. Hagopian , R. Hashmi , R.S. Kim ,
S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay ,
J. Zhang

Florida Institute of Technology, Melbourne, Florida, U.S.A.

B. Alsufyani, M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy ⁸⁶, M. Hohlmann ,
E. Yanes

University of Illinois Chicago, Chicago, U.S.A., Chicago, U.S.A.

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov ,
C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos ,
A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti,
T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhusseini , D. Blend, K. Dilsiz ⁸⁷, L. Emediato , G. Karaman , O.K. Köseyan ,
J.-P. Merlo, A. Mestvirishvili ⁸⁸, O. Neogi, H. Ogul ⁸⁹, Y. Onel , A. Penzo , C. Snyder,
E. Tiras ⁹⁰

Johns Hopkins University, Baltimore, Maryland, U.S.A.

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou ,
P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean ,
Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu ,
J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹¹, C. Rogan , C. Royon ,
R. Salvatico , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli ,
D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, U.S.A.

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen ,
R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili ,
C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova ⁹², A. Shevelev , L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.

J. Bendavid , I.A. Cali , P.c. Chou , M. D’Alfonso , J. Eysermans , C. Freer ,
G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa ,
L. Lavezzo , Y.-J. Lee , K. Long , C. Meginn, A. Novak , C. Paus , C. Roland ,
G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, U.S.A.

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,
R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado ,
W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, U.S.A.

H. Bandyopadhyay , L. Hay , H.w. Hsia, I. Iashvili , A. Kalogeropoulos , A. Kharchilava ,
M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, U.S.A.

G. Alverson , E. Barberis , J. Bonilla , J. Dervan, Y. Haddad , Y. Han , A. Krishna ,
J. Li , M. Lu , G. Madigan , R. Mccarthy , D.M. Morse , V. Nguyen , T. Orimoto ,
A. Parker , L. Skinnari , D. Wood 

Northwestern University, Evanston, Illinois, U.S.A.

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt ,
A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, U.S.A.

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko ¹⁶, H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, U.S.A.

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, U.S.A.

H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik 

Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer, J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Virdi , F. Wang , A. Wildridge , W. Xie 

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, U.S.A.

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, U.S.A.

A. Bodek , P. de Barbaro , R. Demina , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar, P. Parygin ⁹², R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, U.S.A.

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, U.S.A.

D. Aebi , M. Ahmad , T. Akhter , O. Bouhali ⁹³, R. Eusebi , J. Gilmore , T. Huang ,
T. Kamon ⁹⁴, H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, U.S.A.

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim,
K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, U.S.A.

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, U.S.A.

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Neu 

Wayne State University, Detroit, Michigan, U.S.A.

S. Bhattacharya , P.E. Karchin 

University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts ,
C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless ,
J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida ,
L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi ,
W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin [†], I. Gorbunov , V. Karjavine ,
V. Korenkov , A. Lanev , A. Malakhov , V. Matveev ⁹⁵, V. Palichik , V. Perelygin ,
M. Savina , V. ShalaeV , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin ,
B.S. Yuldashev⁹⁶, A. Zarubin , I. Zhizhin , G. GavriloV , V. Golovtsov , Y. Ivanov ,
V. Kim ⁹⁵, P. Levchenko ⁹⁷, V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov ,
A. Vorobyev[†], Yu. Andreev , A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu ,
D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova , A. Toropin , T. Aushev ,
V. GavriloV , N. Lychkovskaya , A. Nikitenko ^{98,99}, V. Popov , A. Zhokin , R. Chistov ⁹⁵,
M. Danilov ⁹⁵, S. Polikarpov ⁹⁵, V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov ,
E. Boos , V. Bunichev , M. Dubinin ⁸⁴, L. Dudko , A. Ershov , A. Gribushin ,
V. Klyukhin , O. Kodolova ⁹⁹, S. Obraztsov , M. Perfilov, V. Savrin , G. Vorotnikov ,
V. Blinov⁹⁵, T. Dimova ⁹⁵, A. Kozyrev ⁹⁵, O. Radchenko ⁹⁵, Y. Skovpen ⁹⁵, V. Kachanov ,
D. Konstantinov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin ¹⁰⁰

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

V. Chekhovsky, V. Makarenko 

[†] Deceased

- ¹ Also at Yerevan State University, Yerevan, Armenia
- ² Also at TU Wien, Vienna, Austria
- ³ Also at Ghent University, Ghent, Belgium
- ⁴ Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
- ⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil
- ⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- ⁷ Also at UFMS, Nova Andradina, Brazil
- ⁸ Also at Nanjing Normal University, Nanjing, China
- ⁹ Now at The University of Iowa, Iowa City, Iowa, U.S.A.
- ¹⁰ Also at University of Chinese Academy of Sciences, Beijing, China
- ¹¹ Also at China Center of Advanced Science and Technology, Beijing, China
- ¹² Also at University of Chinese Academy of Sciences, Beijing, China
- ¹³ Also at China Spallation Neutron Source, Guangdong, China
- ¹⁴ Now at Henan Normal University, Xinxiang, China
- ¹⁵ Also at Université Libre de Bruxelles, Bruxelles, Belgium
- ¹⁶ Also at an institute or an international laboratory covered by a cooperation agreement with CERN
- ¹⁷ Also at Suez University, Suez, Egypt
- ¹⁸ Now at British University in Egypt, Cairo, Egypt
- ¹⁹ Also at Purdue University, West Lafayette, Indiana, U.S.A.
- ²⁰ Also at Université de Haute Alsace, Mulhouse, France
- ²¹ Also at Istinye University, Istanbul, Turkey
- ²² Also at The University of the State of Amazonas, Manaus, Brazil
- ²³ Also at University of Hamburg, Hamburg, Germany
- ²⁴ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ²⁵ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ²⁶ Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁷ Also at Forschungszentrum Jülich, Jülich, Germany
- ²⁸ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁹ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- ³⁰ Now at Universitatea Babes-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania
- ³¹ Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ³² Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ³³ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³⁴ Also at Punjab Agricultural University, Ludhiana, India
- ³⁵ Also at University of Visva-Bharati, Santiniketan, India
- ³⁶ Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁷ Also at IIT Bhubaneswar, Bhubaneswar, India
- ³⁸ Also at Institute of Physics, Bhubaneswar, India
- ³⁹ Also at University of Hyderabad, Hyderabad, India
- ⁴⁰ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴¹ Also at Isfahan University of Technology, Isfahan, Iran
- ⁴² Also at Sharif University of Technology, Tehran, Iran
- ⁴³ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁴ Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
- ⁴⁵ Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁶ Also at Helwan University, Cairo, Egypt
- ⁴⁷ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁸ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵⁰ Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy
- ⁵¹ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

- ⁵² Also at *Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy*
- ⁵³ Also at *Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy*
- ⁵⁴ Also at *Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia*
- ⁵⁵ Also at *Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico*
- ⁵⁶ Also at *Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka*
- ⁵⁷ Also at *Saegis Campus, Nugegoda, Sri Lanka*
- ⁵⁸ Also at *National and Kapodistrian University of Athens, Athens, Greece*
- ⁵⁹ Also at *Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland*
- ⁶⁰ Also at *Universität Zürich, Zurich, Switzerland*
- ⁶¹ Also at *Stefan Meyer Institute for Subatomic Physics, Vienna, Austria*
- ⁶² Also at *Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France*
- ⁶³ Also at *Near East University, Research Center of Experimental Health Science, Mersin, Turkey*
- ⁶⁴ Also at *Konya Technical University, Konya, Turkey*
- ⁶⁵ Also at *Izmir Bakircay University, Izmir, Turkey*
- ⁶⁶ Also at *Adiyaman University, Adiyaman, Turkey*
- ⁶⁷ Also at *Bozok Universititesi Rektörlüğü, Yozgat, Turkey*
- ⁶⁸ Also at *Marmara University, Istanbul, Turkey*
- ⁶⁹ Also at *Milli Savunma University, Istanbul, Turkey*
- ⁷⁰ Also at *Kafkas University, Kars, Turkey*
- ⁷¹ Now at *Istanbul Okan University, Istanbul, Turkey*
- ⁷² Also at *Hacettepe University, Ankara, Turkey*
- ⁷³ Also at *Erzincan Binali Yildirim University, Erzincan, Turkey*
- ⁷⁴ Also at *Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey*
- ⁷⁵ Also at *Yildiz Technical University, Istanbul, Turkey*
- ⁷⁶ Also at *Vrije Universiteit Brussel, Brussel, Belgium*
- ⁷⁷ Also at *School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom*
- ⁷⁸ Also at *IPPP Durham University, Durham, United Kingdom*
- ⁷⁹ Also at *Monash University, Faculty of Science, Clayton, Australia*
- ⁸⁰ Also at *Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt*
- ⁸¹ Also at *Università di Torino, Torino, Italy*
- ⁸² Also at *Bethel University, St. Paul, Minnesota, U.S.A.*
- ⁸³ Also at *Karamanoğlu Mehmetbey University, Karaman, Turkey*
- ⁸⁴ Also at *California Institute of Technology, Pasadena, California, U.S.A.*
- ⁸⁵ Also at *United States Naval Academy, Annapolis, Maryland, U.S.A.*
- ⁸⁶ Also at *Ain Shams University, Cairo, Egypt*
- ⁸⁷ Also at *Bingol University, Bingol, Turkey*
- ⁸⁸ Also at *Georgian Technical University, Tbilisi, Georgia*
- ⁸⁹ Also at *Sinop University, Sinop, Turkey*
- ⁹⁰ Also at *Erciyes University, Kayseri, Turkey*
- ⁹¹ Also at *Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania*
- ⁹² Now at *another institute or international laboratory covered by a cooperation agreement with CERN*
- ⁹³ Also at *Texas A&M University at Qatar, Doha, Qatar*
- ⁹⁴ Also at *Kyungpook National University, Daegu, Korea*
- ⁹⁵ Also at *another institute or international laboratory covered by a cooperation agreement with CERN*
- ⁹⁶ Also at *Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan*
- ⁹⁷ Also at *Northeastern University, Boston, Massachusetts, U.S.A.*
- ⁹⁸ Also at *Imperial College, London, United Kingdom*
- ⁹⁹ Now at *Yerevan Physics Institute, Yerevan, Armenia*
- ¹⁰⁰ Also at *Universiteit Antwerpen, Antwerpen, Belgium*