



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Effects of 3-month Qigong exercise on heart rate variability and respiration in anxious college students

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Sun, J., Zhuo, J., Chu, H., Wang, J., Chen, T., Li, B., et al. (2024). Effects of 3-month Qigong exercise on heart rate variability and respiration in anxious college students. SCANDINAVIAN JOURNAL OF MEDICINE & SCIENCE IN SPORTS, 34(1), 1-9 [10.1111/sms.14521].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/944676> since: 2024-03-14

*Published:*

DOI: <http://doi.org/10.1111/sms.14521>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Sun J, Zhuo J, Chu H, Wang J, Chen T, Li B, Lu T, Zheng H, Xu Y, Dong J, Cicchella A. Effects of 3-month Qigong exercise on heart rate variability and respiration in anxious college students. *Scandinavian Journal of Medicine & Science in Sports* 2024;34:e14521.

The final published version is available online at:

<https://dx.doi.org/10.1111/sms.14521>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

# 3 Effects of three-month Qigong exercise on heart rate variability 4 and respiration in anxious college students

5 Jingyu Sun<sup>1</sup>, Jiatong Zhuo<sup>1</sup>, Hang Chu<sup>1</sup>, Jing Wang<sup>1</sup>, Tao Chen<sup>1</sup>, Bin Li<sup>1</sup>, Tianfeng Lu<sup>1</sup>, Hao Zheng<sup>1</sup>,  
6 Yaning Xu<sup>1</sup>, Jingmei Dong<sup>1</sup>, Antonio Cicchella<sup>2\*</sup>

7 <sup>1</sup>Sports and Health Research Center, Department of Physical Education, Tongji University, Shanghai,  
8 People's Republic of China

9 <sup>2</sup> International College of Football, Tongji University, Shanghai, People's Republic of China

10 Email: antonio.cicchella@unibo.it

11 **Abstract:** Objective: This longitudinal study aimed to investigate the effects of Qigong on the anxiety state, heart rate  
12 variability (HRV), and breathing of anxious college students. Methods: A total of 37 individuals (18–25 years old) were  
13 randomly allocated to the control ( $n = 19$ ) and intervention ( $n = 18$ ) groups. Qigong interventions were conducted five times  
14 weekly for 12 weeks, with each session lasting 60 min. Hamilton Anxiety Scale, Fatigue Scale 14, Pittsburgh Sleep Quality  
15 Index, and 36-item Short Form Survey (SF-36), HRV, and respiration data were collected before and after the 3-month  
16 intervention Results: Individuals who participated in the three-month Qigong exercise intervention showed a significant  
17 reduction in anxiety, particularly mental anxiety ( $P < 0.05$ ). Subjects in the intervention group presented a decrease in skin  
18 temperature ( $P < 0.05$ ) and an increase in blood volume pulsation ( $P < 0.05$ ). Meanwhile, HRV exhibited a significant  
19 increase in the standard deviation of interbeat interval before and after comparisons ( $P < 0.05$ ) and between the two groups  
20 ( $P = 0.039$ ) and a reduction in the normalized low-frequency power after the intervention. Moreover, the intervention group  
21 experienced increased abdominal breathing depth and abdominal breathing per minute ( $P < 0.05$ ). Conclusion: These findings  
22 indicate that Qigong is an effective mind–body exercise strategy for relieving anxiety. HRV and breathing were improved  
23 accordingly among college students after the completion of the 3-month Qigong program.

24 **Keywords:** Qigong; HRV; respiration; college students; anxiety

---

## 26 1. Introduction

27 Excessive academic, social, and life pressure among college students and increased sedentary time have  
28 serious detrimental effects on their physical and mental health, which can easily lead to negative emotions<sup>1</sup>.  
29 Contemporary college students face more negative emotions than ever before, with the incidence of anxiety,  
30 sleep disorders, and depression reaching 32.73%, 15.43%, and 62.91%, respectively<sup>2</sup>. On a psychological  
31 level, anxiety shows association with a poor mental health<sup>3</sup>; on a medical level, it is associated with an  
32 increased all-cause mortality<sup>4</sup>. These data indicate the necessity of developing a method that can effectively  
33 relieve anxiety in college students. Although some drug treatments have been developed, they may be  
34 accompanied with side effects, and occasionally, their effectiveness is uncertain<sup>5</sup>. By contrast, exercise is an  
35 effective, inexpensive, and simple means of reducing anxiety<sup>6</sup>.

36 Fitness Qigong, as a traditional sport in China, has ameliorative effects on a variety of chronic diseases,  
37 such as diabetes<sup>7</sup>, hypertension<sup>8</sup>, chronic heart disease<sup>9</sup>, and cancer<sup>10</sup>, particularly in reducing anxiety and  
38 depression levels<sup>11</sup>. However, studies on fitness Qigong primarily focused on its effects on the physical fitness  
39 among the elderly<sup>12</sup>, and a limited number are related to anxious college students. Compared with highly  
40 confrontational, high-intensity, and difficult exercise programs, fitness Qigong is more suitable for college  
41 students with poor physical fitness, no exercise habits, and are prone to anxiety symptoms because of its “three  
42 tunes in one” for regulating the body, breath, and mind. Fitness Qigong includes many types, which all have

---

---

43 different fitness effects<sup>13</sup>. Yijinjing enhances physical fitness and prevents diseases through postural  
44 adjustments and breathing exercises, whereas Liuzijue improves the functions of internal organs and alleviates  
45 negative emotions through vocalization<sup>14</sup>. Therefore, this study combined Yijinjing and Liuzijue to develop  
46 an exercise intervention program for anxious college students.

47 Anxiety and negative emotions can be assessed through certain physiological measures, particularly heart  
48 rate variability (HRV)<sup>15</sup>. HRV is a measure of cardiac vagal tone. It can be quantified through the spectral  
49 analysis of interbeat interval (RR)<sup>16</sup>. A low HRV is related to a number of psychopathological states, including  
50 anxiety<sup>17</sup>. However, the effect of Qigong on the HRV of anxious college students remains poorly understood.  
51 Breathing exercises (as long as the inhalation/exhalation ratio is equal) as short as 5 min promote a balance  
52 between sympathetic and parasympathetic activities<sup>18</sup>, which increases HRV and improves  
53 psychophysiological coherence<sup>19</sup>. Therefore, fitness Qigong may reduce anxiety by modulating HRV,  
54 breathing depth, and breathing frequency.

55 In this study, we initially determined the alleviating effect of combined Qigong practices on the anxiety  
56 of college students. We also analyzed the possible mechanisms of fitness Qigong in improving anxiety by  
57 comparing the changes in HRV and respiratory indexes before and after fitness Qigong practice and identified  
58 multidimensional, subjective, and objective measurement and evaluation to illustrate its improvement effect  
59 on the anxiety status of college students.

## 60 2. Methods

### 61 2.1 Participants

62 A total of 68 undergraduate students (30 males and 38 females; mean age = 21.20 years, range: 18–25  
63 years; mean body mass index (BMI) = 22.17, range: 18–30) participated in this experiment (Table 1). Males  
64 and females were grouped together because differences in the results showed no statistical significance for  
65 any of the variables measured. The students were asked to complete a basic questionnaire prior to exercise  
66 intervention. Those who satisfied the following criteria were qualified to participate in the study: (1) aged  
67 between 18 and 25 years; (2) sedentary lifestyle (<150 min of moderate-to-high-intensity physical activity per  
68 week, and >8 h daily sedentary time); (3) severe anxiety (scores in the Hamilton Anxiety Rating Scale [HAM-  
69 A] >14); (4) willingness to participate in the test and sign an informed consent form. Participants were  
70 excluded if they met the following exclusion criteria: (1) diagnosed with a disease that affects the performance  
71 of movements; (2) history of heart disease, severe arrhythmia, or pacemaker use; (3) habitual smoking or  
72 heavy consumption of alcohol or other stimulant beverages; (4) have taken drugs that may affect emotional  
73 reaction, such as antihypertensives, anticholinergics, anxiolytics, or antidepressants, at least 1 month before  
74 the start of the study. Of the included individuals, 26 did not meet the screening criteria and hence were  
75 eliminated from the study (Fig. 1). The remaining 42 participated in the experiment. The subjects were  
76 randomly divided into two groups: the control group, who did not conduct any regular exercise, and the  
77 intervention group, who participated in regular Qigong exercise. Two and three individuals dropped out after  
78 being assigned to the control ( $n = 19$ ) and intervention ( $n = 18$ ) groups.




Figure 1. Participant flow diagram

## 2.2 Experimental procedures

Participants completed the HAM-A for anxiety, Fatigue Scale 14 (FS-14), Pittsburgh Sleep Quality Index (PSQI), and 36-Item Short Form Survey (SF-36). Prior to the test, they were asked to inhibit from drinking caffeinated or alcoholic beverages for at least 24 h. All the subjects were tested individually, and data were collected in a constant-temperature (24 °C–26 °C) and bright lounge between 14:00 and 17:00 pm. Then, the participants were instructed to sit on a chair and rest for 5 min to determine the indicators of multiple physiological parameters (MPPs), HRV, and respiration (pretest). Finally, signals were collected using a biofeedback instrument and transmitted to a laptop for postanalysis. The intervention group adopted the combined Yijinjing and Liuzijue Qigong styles as the training method under the guidance of experienced coaches. The Qigong program lasted 12 weeks, with a frequency of 5 days a week and 60 min a day and including a 10 min preworkout warm-up, 40 min exercise workout, and 10 min cool down. At the end of the 3-month intervention, the subjects were reassessed in accordance with the preintervention steps (posttest). All exercise and measurement procedures were conducted under professional guidance. The study was approved by the Ethics Committee of Tongji University (2021tjdx024).

## 2.3 Measurements

### 2.3.1 Scale evaluation

Prior to Qigong interventions, the participants were assessed using the HAM-A, FS-14, PSQI, and SF-36. HAM-A measures the severity of anxiety symptoms in patients with neuroses and others who have different conditions, primarily adults with anxiety symptoms<sup>20</sup>. FS-14 consists of 14 entries, each of which focuses on a fatigue-related issue. These 14 questions reflect the severity of fatigue from various perspectives, and they are divided into two categories, including physical fatigue and mental fatigue, through principal component analysis<sup>21</sup>. PSQI refers to a quantity scale that is used to evaluate sleep quality in general<sup>22</sup>. SF-36 is a frequently used, well-researched measurement of health. It comprises 36 questions, including eight domains of health, and commonly used to measure a population's life quality<sup>23</sup>.

### 2.3.2 MPP measurement

MPPs, including skin conductance level, skin temperature (TEMP), blood volume pulsation, blood volume amplitude, and pulse, were evaluated using biofeedback instruments (2000x-pert, Biopac Systems Inc.). The skin conduction sensor had two electrodes. The electrode was tightly tied to the end of a finger to maintain a good contact but not extremely tight to allow blood circulation. For MPP measurement, Bioneural Infiniti v5.1 operating platform, which can formulate single-data reports or trend reports of MPP data, was used for data acquisition, analysis, and management.

### 2.3.3 HRV measurement

As an important indicator of cardiac sympathetic–parasympathetic tone, the HRV primarily reflects changes in the RR<sup>24</sup>. HRV can be measured in the time and frequency domains. In time-domain analysis, we used SDNN and PNN50 metrics, which can reflect the total variation and magnitude of HRV, respectively, as measures.

HRV was measured using biofeedback instruments(2000x-pert, Biopac Systems Inc.). R-wave detection was used to identify R-wave peaks in recorded electrocardiographic signals and measure continuous RR during tracking. HRV-related data were collected before and after the Qigong intervention and labeled for separation.

### 2.3.4 Respiration measurement

The respiratory index, including the amplitude and frequency of breathing in the chest and abdomen, was assessed using a biofeedback instrument (2000x-pert, Biopac Corporation). Circumferential changes were measured using ultrasound signals to maintain the accuracy of readings. Breathing measurements included four indicators: (1) maximum and minimum of the abdominal breathing curve (RESPA-A); (2) the number of breaths per minute (calculated from the abdominal respiratory curve [RESPF-A]); (3) maximum and minimum of the thoracic respiratory curve (RESPA-C); (4) the number of breaths per minute, which was calculated on the basis of the chest respiratory curve (RESPF-C). The participants were required to breathe naturally before and after the exercise and coordinate their movements and breathing during Qigong exercises.

## 2.4 Statistics and analysis

Statistical analyses were performed using the SPSS statistical package (IBM, SPSS Statistics 20). The extracted data included participant characteristics, questionnaire scores, HRV, respiration, and MPPs. The means and standard deviation (SD) of key outcome indicators were extracted at baseline and at time point closest to the end of exercise. Levine’s and Shapiro–Wilk tests were used to check the homogeneity of variance and normality of data, respectively. All data were normally distributed. Data that did not meet the test parameters were converted when necessary via regression. Independent sample t-test was used to compare the baseline characteristics of the two groups, and the same was performed for the results of HAM-A, FS-14, PSQI, and SF-36 between the two groups. Independent and paired sample t-tests were conducted to compare the differences between pretesting and posttesting, respectively, of MPP, HRV, and respiration. The confidence level was set at 95% ( $P < 0.05$ ) in all statistical analyses.

## 3. Results

### 3.1 Demographic characteristics and screening criteria for participants in the two groups

The study initially screened 68 sedentary college students, 26 of whom were prescreened out for failing to meet the criteria and 5 of whom dropped out. Finally, 37 students, all of which were anxious young adults, participated in this research. Similarities in their basic information and subhealth levels were considered during grouping to ensure that no significant differences exist in various baseline parameters between the two groups (males and females and control versus intervention groups). Table 1 shows the basic characteristics of the participants between the two groups, which showed no significant differences in their age, sex, body weight, body height, BMI, and education level.

Table 1. Characteristics of participants

Variable	Control ( $n = 19$ )	Intervention ( $n = 18$ )	Total
N (%)	19 (51.4%)	18 (48.6%)	37 (100%)

Age	20.2 ± 3.5	21.7 ± 5.1	20.9 ± 4.3
Sex (M/F)	11/8	7 /11	18/19
Height (cm)	165.2 ± 10.2	164.9 ± 8.4	165.1 ± 9.3
Weight (kg)	62.46 ± 5.25	59.18 ± 4.53	60.86 ± 4.90
BMI (kg/m <sup>2</sup> )	19.76 ± 2.06	21.23 ± 5.82	20.48 ± 3.89
Education level (bachelor degree and postgraduate, %)	89.34	82.11	85.82

151 Data are presented as mean ± SD. BMI, body mass index.

### 152 3.2 Effect of Qigong intervention on anxiety status among college students

153 After the three-month Qigong exercise intervention, significant improvements (decrease in  
154 questionnaires scores) in the total anxiety ( $P = 0.039$ ) and mental anxiety ( $P = 0.004$ ) were observed. Physical  
155 anxiety did not change, which was possibly because the amount of Qigong exercise did not reach the level of  
156 changing physical anxiety for a three-month beginner. Compared with the control group, the intervention  
157 group showed a significant improvement in physical pain ( $P = 0.013$ ) In addition, the intervention group  
158 attained a lower sleep quality score ( $P = 0.027$ ), which indicated an improvement. “Life quality” and “Bodily  
159 pain” showed high SDs, which can be explained by the anxiety conditions of participants, who could have  
160 different perceptions (Table 2).

161 Table 2. Effect of Qigong on anxiety, fatigue, and sleep quality among anxious college students

	Control ( $n = 19$ )	Intervention ( $n = 18$ )	Change (95%)	$P$ value
Total anxiety (Ham-a)	15.67±5.94	11.63±5.50*	4.04 (0.217 to 7.853)	0.039
Physical anxiety (Ham-a)	6.50±3.09	5.89±3.11	0.61 (-1.465 to 2.675)	0.714
Mental anxiety (Ham-a)	9.17±4.29	5.74±5.05*	3.21 (1.094 to 5.332)	0.004
Total fatigue (FS-14)	8.39±2.83	8.16±2.81	0.13 (-1.654 to 2.116)	0.805
Sleep quality (PSQI)	7.61±2.59	6.06±1.58*	1.66 (0.208 to 3.120)	0.027
Life quality (SF-36)	511.51±98.63	523.91±104.3	-5.21 (-50.737 to 40.313)	0.817
Bodily pain (SF-36)	63.29±30.55	84.13±11.28*	-20.83 (-36.866 to -4.795)	0.013

162 Data are presented as mean ± SD, \*  $P < 0.05$  represents statistically significant difference.

### 163 3.3 Effect of Qigong intervention on MPPs of anxious college students

164 Psychological improvements related to anxiety levels also cause changes in basic physiological  
165 indicators. Thus, in this study, the effects of Qigong on these indicators were investigated through  
166 measurement of several parameters. After the Qigong intervention, the TEMP index in the intervention group  
167 was significantly lower than the preintervention level ( $P < 0.05$ ). Meanwhile, the BVP in the intervention  
168 group showed a significant increase after the intervention ( $P < 0.05$ ) and responded more prominently to  
169 Qigong intervention than that in the control group ( $P = 0.029$ ). Other indices exhibited no significant change  
170 between the groups before and after the intervention (Table 3).

171 Table 3. Effects of Qigong on the MPPs of pre- and posttests in two groups (control and intervention) of  
172 anxious college students

Index	Control ( $n = 19$ )			Intervention ( $n = 18$ )			$P$ value
	Pre	Post	Change (95%CI)	Pre	Post	Change (95%CI)	
SCL	1.24±0.70	1.24±0.96	0.00 (-0.649 to 0.653)	1.43±0.83	4.29±6.31	-2.86 (-6.500 to 0.774)	0.114
TEMP	32.04±3.76	30.09±4.40	1.95 (-2.196 to 6.105)	34.43±2.78	31.52±3.81*	2.91 (1.641 to 4.171)	0.620

BVP	49.45±0.45	49.74±1.20	-0.29 (-1.421 to 0.831)	48.03±2.38	51.40±3.61*	-3.36 (-5.935 to -2.881)	0.029
PVA	67.88±22.97	44.14±21.47	23.74 (-5.598 to 53.091)	54.64±23.02	48.53±19.65	6.12 (-12.195 to 24.427)	0.235
PULS	68.66±5.25	67.40±4.19	1.26 (-2.145 to 4.673)	73.93±7.90	74.87±5.13	-0.94 (-6.834 to 4.945)	0.478

173 Data are presented as mean ± SD. \* represents statistically significant difference in pre- and post-test. *P* value indicates differences  
174 in variables between the control and intervention group. SCL, skin conductance level; TEMP, skin temperature; BVP, blood volume  
175 pulsation; PVA, blood volume amplitude, relative maximum recorded range of BVP peak values; PULS, number of heart beats per  
176 minute.

### 177 3.4 Effects of Qigong on HRV of anxious college students

178 In the time-domain analysis, SDNN in the intervention group presented a significant increase in after the  
179 intervention ( $P < 0.05$ ). Furthermore, the changes in SDNN ( $P = 0.039$ ) between the two groups showed a  
180 significant difference after the intervention. Frequency-domain analysis enables the detailed observation of  
181 sympathetic and vagal modulation, which compensates for the poor sensitivity and specificity of time-domain  
182 methods<sup>25</sup>. Therefore, we further analyzed the frequency domain and selected three indicators: the low-  
183 frequency power spectrum band (LF: 0.04–0.15 Hz), high-frequency power spectrum band (HF: 0.15–0.4 Hz),  
184 and LF/HF ratio. The results showed that the normalized LF in the intervention group was significantly lower  
185 after Qigong intervention than before intervention ( $P < 0.05$ ). No statistically significant difference in  
186 normalized HF and LF/HF was observed within and between groups, and the results are reported in Table 4.

187 Table 4. Effects of Qigong on the HRV of the control and intervention groups in the pre- and posttests.

Index	Control ( <i>n</i> = 19)			Intervention ( <i>n</i> = 18)			<i>P</i> value
	Pre	Post	Change (95%CI)	Pre	Post	Change (95%CI)	
SDNN	87.79±71.81	66.91±25.36	20.88 (-43.343 to 85.102)	78.59±39.04	123.61±59.58*	-45.02 (-79.737 to -10.304)	0.335
PNN50	29.90±19.44	30.93±10.03	-1.03 (-17.848 to 15.786)	33.18±23.48	31.93±19.08	1.25 (-12.061 to 14.553)	0.523
nLF	40.80±10.17	44.94±9.76	-3.13 (-14.159 to 7.900)	43.78±10.17	37.77±6.67*	5.01 (1.314 to 8.715)	0.351
nHF	59.18±10.17	56.52±12.36	2.67 (-8.343 to 13.682)	58.68±11.27	60.57±7.52	-1.89 (-7.825 to 4.041)	0.307
LF/HF	0.73±0.29	0.87±0.54	-0.14 (0.195 to 0.326)	0.77±0.38	0.80±0.34	-0.03 (-0.273 to 0.213)	0.195

188 Data are presented as mean ± SD. \* represents statistically significant difference in pre- and post-test. *P* value indicates  
189 differences in variables between the control and intervention group. SDNN, the standard deviation of R-R intervals; PNN50,  
190 the percentage of difference over 50 ms between adjacent normal-normal R-R intervals. nLF, normalized low-frequency  
191 power; nHF, normalized high-frequency power; LF/HF, low-frequency/high-frequency ratio.

### 192 3.5 Effect of Qigong exercise on respiration regulation in anxious college students

193 The two main modes of physiological breathing include thoracic and abdominal breathing, which are  
194 also known as shallow and deep breathing. We analyzed the differences in the amplitude and frequency  
195 between thoracic and abdominal respiration to gain insights into the effect of Qigong exercises on respiratory  
196 regulation. Compared with those in the pretest, RESP-A ( $P < 0.05$ ) decreased, and RESPF-C ( $P < 0.05$ )  
197 increased after the test in the control group. Compared with those in the pretest, RESP-A ( $P < 0.05$ ) decreased,  
198 and RESPA-A ( $P < 0.05$ ), RESPF-A ( $P < 0.05$ ), and RESPF-C ( $P < 0.05$ ) increased in the intervention group  
199 after Qigong exercise, which indicate the important role of Qigong in regulating the amplitude and frequency  
200 of respiration, particularly in abdominal breathing of anxious college students (Table 5).

201 Table 5. Effects of Qigong on the respiration of the control and intervention groups during pre- and posttest  
202 conducted among anxious college students.

Index	Control ( <i>n</i> = 19)			Intervention ( <i>n</i> = 18)			<i>P</i> value
	Pre	Post	Change(95%CI)	Pre	Post	Change(95%CI)	



RESP-A	9.77±0.32	9.18±0.44*	0.58 (0.114 to 1.055)	9.31±0.48	8.88±0.37*	0.43 (0.0816 to 0.778)	0.335
RESPA-A	0.60±0.25	0.70±0.28	-0.10 (-0.246 to 0.046)	0.72±0.28	0.89±0.19*	-0.18 (-0.351 to 0.007)	0.523
RESPF-A	14.10±3.81	15.62±3.89	-1.53 (-3.676 to 0.625)	13.13±3.35	16.18±4.37*	-3.05 (-5.564 to -0.540)	0.351
RESP-C	9.11±1.06	9.22±0.38	-0.11 (-0.948 to 0.726)	8.93±0.78	9.49±0.63	-0.56 (-1.133 to 0.008)	0.307
RESPA-C	0.65±0.38	0.76±0.40	-0.11 (-0.311 to 0.088)	0.80±0.42	0.63±0.23	0.17 (-0.179 to 0.523)	0.195
RESPF-C	13.88±3.84	17.48±2.72*	-3.60 (-0.311 to 0.088)	13.30±3.15	16.53±3.95*	-3.23 (-5.856 to -0.606)	0.825

203 Data are presented as mean ± SD. \* represents statistically significant difference between pre- and post-test in the control and  
204 intervention group. *P* value indicates difference in variables between the control and intervention group. RESP-A, abdominal  
205 breathing; RESPA-A, depth of abdominal breathing; RESPF-A, frequency of abdominal breathing; RESP-C, chest breathing;  
206 RESPA-C, depth of chest breathing; RESPF-C, frequency of chest breathing.

207

#### 208 4. Discussion

209 This study innovatively selected the combination of Yijinjing and Liuzijue from various types of Qigong  
210 exercise to illustrate the effect of breathing training in Qigong on anxiety relief. We observed that Yijinjing  
211 exercises strengthen the body, prevent disease, and improve cognitive function and negative emotion, as also  
212 observed previously from other studies that employed different Qigong techniques<sup>26</sup>. Liuzijue emphasizes the  
213 use of abdominal breathing during practice, which allows for deep and long breaths<sup>27</sup>. Therefore, the addition  
214 of Liuzijue exercises to Yijinjing can enhance the breathing depth through vocal training while retaining the  
215 easy-to-grasp and rhythmically diverse characteristics of Yijinjing, which can relieve anxiety in college  
216 students. Our results revealed that the combined practice of Yijinjing and Liuzijue can effectively improve the  
217 anxiety state and sleep quality among college students. Overall, these findings suggest that a 3-month Qigong  
218 intervention can improve anxiety status, sleep quality, and physical pain among college students with anxiety  
219 disorders.

220 The occurrence of psychological disorders is closely related to physiological factors, such as TEMP and  
221 skin resistance indicators associated with stress levels<sup>28,29</sup>. In this study, short-term Qigong exercises can  
222 considerably reduce psychological stress, as observed in previous works<sup>28</sup> that examined stress-related  
223 physiological indicators and revealed reductions in heart rate and finger temperature. We also observed a  
224 considerable reduction in subjective stress responses. We also showed that Qigong practice achieved and  
225 maintained remarkable decreases in the heart rate, respiratory rate, and systolic blood pressure and maintained  
226 long-term stress reduction. These results are supported by those of previous research, albeit the literature on  
227 this point is limited<sup>29</sup>. In addition, our study indicated a substantial improvement in physical pain perception  
228 of anxious college students after 3-month Qigong training, which is consistent with that of a previous study<sup>30</sup>;  
229 Qigong exercise has also a positive effect on some people with neck pain<sup>31</sup>. Our study adds knowledge to the  
230 understanding of Qigong as a therapeutic tool although its effects on pain relief must be further explored. In  
231 general, these results can be explained as the improvement in autonomic nervous system functioning induced  
232 by the relaxing effect of Qigong.

233 Our findings are in agreement with those of studies performed in the last decades, which provided  
234 considerable evidence showing that Qigong improves autonomic nerves, which increases vagal regulation<sup>32-</sup>  
235 <sup>34</sup>. Most studies have shown that after Qigong intervention, HRV indicators, such as time-domain indicators  
236 (SDNN and RMSSD), increased, and frequency-domain indicators (LF and LF/HF) decreased<sup>35</sup>, which are  
237 consistent with our findings. Therefore, we hypothesized that Qigong may relieve anxiety through HRV  
238 regulation, and it possibly acts on the autonomic innervation system of the heart and lungs. First, the slow  
239 movement patterns of Yijinjing and Liuzijue can alter the autonomic system and modulate its balance toward  
240 parasympathetic dominance, which can improve HRV. Second, Qigong can improve HRV by changing the  
241 breathing pattern and depth. These results are supported by those of previous studies<sup>19</sup>. Finally, the unique  
242 psychological conditioning of Qigong, such as meditation, leads to a decrease in sympathetic excitability and  
243 an increase in vagal excitability, which results in an improved HRV<sup>35</sup>. The abovementioned hypothesis about  
244 Qigong explains its beneficial use as a means of improving HRV, which will contribute to anxiety relief.

---

The adverse effects of stress and negative emotions can be counteracted through different forms of breathing techniques, such as meditation and relaxation<sup>36</sup>. Liuzijue Qigong typically involves training with breathing techniques<sup>27</sup>. Breathing techniques with equal inhalation/exhalation ratios effectively promote the balance between sympathetic and parasympathetic activities and increases HRV<sup>37</sup>. Abdominal breathing is an effective intervention that can increase inspiratory volume and achieve an equal inhalation/exhalation ratio<sup>37</sup>. All these effects have been confirmed by our study, which showed the beneficial effect of two different forms of Qigong. These results can be ascribed to the activation of the autonomic nervous system. A possible effect on the brain might have also caused the improvement of physiological parameters, as demonstrated by a study on the effect of Qigong on increasing alpha-waves in electroencephalogram<sup>38</sup>. The results of the present research showed that the depth and frequency of abdominal breathing among anxious college students were effectively improved through conscious breathing training during Qigong exercise. Therefore, breathing techniques of Qigong can be used as first-line and supplemental treatments for anxiety, depression, and other emotional disorders. In the follow-up study, more attention should be paid to the mechanisms and differences in the effects of various breathing techniques of Qigong exercise. In addition, the physiological importance of breathing training in Qigong for anxious college students should be considered. Although this study provided promising results, it had some limitations, e.g., its inapplicability to college students with other related disorders, such as psychological disorders, frailty, or metabolic syndrome when the applicability and distinction between different kinds of Qigong interventions are considered. In the future, the number of subjects and duration of training intervention must be increased, and different types of Qigong must be associated with various chronic diseases in the form of exercise prescriptions.

## 5. Conclusions and perspectives

Qigong has been studied as a means of improving psychological and associated physiological factors. Compared with the more recent relaxation methods, this form of exercise is grounded in a millenary traditional culture, which piques the interest of people. Qigong is accepted more by eastern (Chinese) populations, which could have added a placebo effect on the results. This study revealed that Qigong exercise may have a vital mood-enhancing effect on anxious college students with a 3-month training, and these changes are affected by the improvement in HRV and breathing regulation. In terms of clinical relevance, this study unveiled the potential effect of two Qigong styles on improving the quality of life of sedentary and anxious university students and ameliorating their basic functional physiological parameters. Follow-up studies should also explore the mechanisms behind such phenomena and compare the differences between traditional Qigong and other exercise modalities in terms of anxiety relief.

## FUNDING INFORMATION

This research was funded by the Shanghai Education Science Research Project (C2021125, C2-2020030), the Scientific Fitness Guidance Project of the National Sports General Administration of China (2017B026), the Tengfei Project of the Shanghai Municipal Sports Bureau (16T006), and the National Undergraduate Innovation and Entrepreneurship Training Program (X2021417).

## CONFLICT OF INTEREST

The authors declare no conflicts of interest.

## DATA AVAILABILITY STATEMENT

The data are available from the corresponding author upon reasonable request.

## AUTHOR CONTRIBUTIONS

Author JYS and JTZ have given substantial contributions to the conception and the design of the manuscript; author HC, JW, BL, TFL, YNX and HZ performed the experiment; author TC performed the

---

289 acquisition of the data; author AC and JMD contributed to analysis and interpretation of the data. All authors  
290 read and approved the final version of the manuscript.

- 291
- 292 1. Akbari HA, Pourabbas M, Yoosefi M, et al. How physical activity behavior affected well-being, anxiety and sleep quality  
293 during COVID-19 restrictions in Iran. *European review for medical and pharmacological sciences*. Dec 2021;25(24):7847-7857.  
294 doi:10.26355/eurrev\_202112\_27632
- 295 2. Duan H, Gong M, Zhang Q, Huang X, Wan B. Research on sleep status, body mass index, anxiety and depression of college  
296 students during the post-pandemic era in Wuhan, China. *Journal of affective disorders*. Mar 15 2022;301:189-192.  
297 doi:10.1016/j.jad.2022.01.015
- 298 3. Casey SM, Varela A, Marriott JP, Coleman CM, Harlow BL. The influence of diagnosed mental health conditions and  
299 symptoms of depression and/or anxiety on suicide ideation, plan, and attempt among college students: Findings from the Healthy  
300 Minds Study, 2018-2019. *Journal of affective disorders*. Feb 1 2022;298(Pt A):464-471. doi:10.1016/j.jad.2021.11.006
- 301 4. Daviu N, Bruchas MR, Moghaddam B, Sandi C, Beyeler A. Neurobiological links between stress and anxiety. *Neurobiology*  
302 *of stress*. Aug 13 2019;11:100191. doi:10.1016/j.ynstr.2019.100191
- 303 5. Koen N, Stein DJ. Pharmacotherapy of anxiety disorders: a critical review. *Dialogues in clinical neuroscience*. Apr 01  
304 2022;13(4):423-37. doi:10.31887/DCNS.2011.13.4/nkoen
- 305 6. McCraty R, Zayas MA. Cardiac coherence, self-regulation, autonomic stability, and psychosocial well-being. *Frontiers in*  
306 *psychology*. Sep 29 2014;5:1090. doi:10.3389/fpsyg.2014.01090
- 307 7. Huang CL, Tai YK, Yang YH, Wang RH. Efficacy of five-element gymnastics in glucose and lipid control in Taiwanese  
308 patients with type 2 diabetes. *Research in nursing & health*. Aug 2012;35(4):419-29. doi:10.1002/nur.21480
- 309 8. Yu AP, Tam BT, Lai CW, et al. Revealing the Neural Mechanisms Underlying the Beneficial Effects of Tai Chi: A  
310 Neuroimaging Perspective. *The American journal of Chinese medicine*. 2018;46(2):231-259. doi:10.1142/s0192415x18500131
- 311 9. Chan CL, Wang CW, Ho RT, et al. A systematic review of the effectiveness of qigong exercise in cardiac rehabilitation. *The*  
312 *American journal of Chinese medicine*. 2012;40(2):255-67. doi:10.1142/s0192415x12500206
- 313 10. Oberoi D, Piedalue KL, Pirbhai H, Guirguis S, Santa Mina D, Carlson LE. Factors related to dropout in integrative oncology  
314 clinical trials: interim analysis of an ongoing comparative effectiveness trial of mindfulness-based cancer recovery and Tai chi/Qigong  
315 for cancer health (The MATCH study). *BMC research notes*. Jul 17 2020;13(1):342. doi:10.1186/s13104-020-05172-5
- 316 11. Yeung A, Chan JSM, Cheung JC, Zou L. Qigong and Tai-Chi for Mood Regulation. *Focus (American Psychiatric Publishing)*.  
317 Jan 24 2018;16(1):40-47. doi:10.1176/appi.focus.20170042
- 318 12. Song R, Grabowska W, Park M, et al. The impact of Tai Chi and Qigong mind-body exercises on motor and non-motor  
319 function and quality of life in Parkinson's disease: A systematic review and meta-analysis. *Parkinsonism & related disorders*. Aug  
320 2017;41:3-13. doi:10.1016/j.parkreldis.2017.05.019
- 321 13. Li L, Huang H, Song J, et al. Network Meta-Analysis of the Effects of Different Types of Traditional Chinese Exercises on  
322 Pulmonary Function, Endurance Capacity and Quality of Life in Patients With COPD. *Frontiers in medicine*. Feb 2 2022;9:806025.  
323 doi:10.3389/fmed.2022.806025
- 324 14. Gao P, Tang F, Liu W, He K, Mo Y. Effect of liuzijue qigong on patients with stable chronic obstructive pulmonary disease:  
325 A systematic review and meta-analysis. *Medicine*. Oct 15 2021;100(41):e27344. doi:10.1097/md.0000000000027344
- 326 15. Cheng YC, Su MI, Liu CW, Huang YC, Huang WL. Heart rate variability in patients with anxiety disorders: A systematic  
327 review and meta-analysis. *Psychiatry and clinical neurosciences*. Jul 2022;76(7):292-302. doi:10.1111/pcn.13356
- 328 16. Porges SW. The polyvagal perspective. *Biological psychology*. Feb 2007;74(2):116-43. doi:10.1016/j.biopsycho.2006.06.009
- 329 17. Kim HG, Cheon EJ, Bai DS, Lee YH, Koo BH. Stress and Heart Rate Variability: A Meta-Analysis and Review of the  
330 Literature. *Psychiatry investigation*. Feb 28 2018;15(3):235-245. doi:10.30773/pi.2017.08.17

- 
- 331 18. Steffen PR, Austin T, DeBarros A, Brown T. The Impact of Resonance Frequency Breathing on Measures of Heart Rate  
332 Variability, Blood Pressure, and Mood. *Frontiers in public health*. Aug 25 2017;5:222. doi:10.3389/fpubh.2017.00222
- 333 19. Magnon V, Dutheil F, Vallet GT. Benefits from one session of deep and slow breathing on vagal tone and anxiety in young  
334 and older adults. *Scientific reports*. Sep 29 2021;11(1):19267. doi:10.1038/s41598-021-98736-9
- 335 20. Maier W, Buller R, Philipp M, Heuser I. The Hamilton Anxiety Scale: reliability, validity and sensitivity to change in anxiety  
336 and depressive disorders. *Journal of affective disorders*. Jan 1988;14(1):61-8. doi:10.1016/0165-0327(88)90072-9
- 337 21. Chalder T, Berelowitz G, Pawlikowska T, et al. Development of a fatigue scale. Feb 1993;37(2):147-153.
- 338 22. Buysse DJ, Reynolds CF, 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for  
339 psychiatric practice and research. *Psychiatry research*. May 1989;28(2):193-213. doi:10.1016/0165-1781(89)90047-4
- 340 23. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection.  
341 *Medical care*. Jun 1992;30(6):473-83.
- 342 24. Soares-Miranda L, Sattelmair J, Chaves P, et al. Physical activity and heart rate variability in older adults: the Cardiovascular  
343 Health Study. *Circulation*. May 27 2014;129(21):2100-10. doi:10.1161/circulationaha.113.005361
- 344 25. Blase KL, van Waning A. Heart Rate Variability, Cortisol and Attention Focus During Shamatha Quiescence Meditation.  
345 *Applied psychophysiology and biofeedback*. Dec 2019;44(4):331-342. doi:10.1007/s10484-019-09448-w
- 346 26. Sun P, Zhang S, Jiang L, et al. Yijinjing Qigong intervention shows strong evidence on clinical effectiveness and  
347 electroencephalography signal features for early poststroke depression: A randomized, controlled trial. *Frontiers in aging neuroscience*.  
348 Aug 30 2022;14:956316. doi:10.3389/fnagi.2022.956316
- 349 27. Gong T, Lu T, Zhang Y, et al. Effects of Liuzijue Qigong Posture on Aerodynamics of Phonation in Healthy Volunteers.  
350 *Journal of voice : official journal of the Voice Foundation*. Feb 16 2022;doi:10.1016/j.jvoice.2021.12.019
- 351 28. Lee MS, Lee MS, Kim HJ, Moon SR. Qigong reduced blood pressure and catecholamine levels of patients with essential  
352 hypertension. *The International journal of neuroscience*. Dec 2003;113(12):1691-701. doi:10.1080/00207450390245306
- 353 29. Skoglund L, Jansson E. Qigong reduces stress in computer operators. *Complementary therapies in clinical practice*. May  
354 2007;13(2):78-84. doi:10.1016/j.ctcp.2006.09.003
- 355 30. Marks R. Qigong and Musculoskeletal Pain. *Current rheumatology reports*. Nov 16 2019;21(11):59. doi:10.1007/s11926-019-  
356 0861-6
- 357 31. Girard J, Girard A. The effects of qigong on neck pain: A systematic review. *Complementary therapies in clinical practice*. Feb  
358 2019;34:23-29. doi:10.1016/j.ctcp.2018.10.013
- 359 32. Tai HC, Chou YS, Tzeng IS, et al. Effect of Tai Chi Synergy T1 Exercise on Autonomic Function, Metabolism, and Physical  
360 Fitness of Healthy Individuals. *Evidence-based complementary and alternative medicine : eCAM*. Jun 27 2018;2018:6351938.  
361 doi:10.1155/2018/6351938
- 362 33. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and  
363 meta-analysis. *JAMA internal medicine*. Mar 2014;174(3):357-68. doi:10.1001/jamainternmed.2013.13018
- 364 34. Larkey L, Kim W, James D, et al. Mind-Body and Psychosocial Interventions May Similarly Affect Heart Rate Variability  
365 Patterns in Cancer Recovery: Implications for a Mechanism of Symptom Improvement. *Integrative cancer therapies*. Jan-Dec  
366 2020;19:1534735420949677. doi:10.1177/1534735420949677
- 367 35. Nesvold A, Fagerland MW, Davanger S, et al. Increased heart rate variability during nondirective meditation. *European  
368 journal of preventive cardiology*. Aug 2012;19(4):773-80. doi:10.1177/1741826711414625
- 369 36. Jerath R, Crawford MW, Barnes VA, Harden K. Self-regulation of breathing as a primary treatment for anxiety. *Applied  
370 psychophysiology and biofeedback*. Jun 2015;40(2):107-15. doi:10.1007/s10484-015-9279-8

---

371 37. MacKinnon S, Gevirtz R, McCraty R, Brown M. Utilizing heartbeat evoked potentials to identify cardiac regulation of vagal  
372 afferents during emotion and resonant breathing. *Applied psychophysiology and biofeedback*. Dec 2013;38(4):241-55. doi:10.1007/s10484-  
373 013-9226-5

374 38. Faber PL, Lehmann D, Tei S, et al. EEG source imaging during two Qigong meditations. *Cognitive processing*. Aug  
375 2012;13(3):255-65. doi:10.1007/s10339-012-0441-4

376

377