



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Mid-term (30 years) changes of soil properties under chestnut stands due to organic residues management:
An integrated study

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Mid-term (30 years) changes of soil properties under chestnut stands due to organic residues management: An integrated study / De Feudis M.; Falsone G.; Vittori Antisari L.. - In: CATENA. - ISSN 0341-8162. - ELETTRONICO. - 198:March 2021(2021), pp. 105021.1-105021.11. [10.1016/j.catena.2020.105021]

This version is available at: <https://hdl.handle.net/11585/792427> since: 2021-01-28

Published:

DOI: <http://doi.org/10.1016/j.catena.2020.105021>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

Manuscript Number: CATENA11740R1

Title: Mid-term (30 years) changes of soil properties under chestnut stands due to organic residues management: An integrated study

Article Type: Research Paper

Keywords: chestnut management; mountain soil; organic carbon dynamics; litter floor, soil development

Corresponding Author: Dr. Mauro De Feudis,

Corresponding Author's Institution: University of Bologna

First Author: Mauro De Feudis

Order of Authors: Mauro De Feudis; Gloria Falsone; Livia Vittori Antisari

Abstract: Chestnut plantations are worldwide distributed and they are often subjected to intensive management practices such as the removal of the organic residues from the soil surface. The present study aimed to investigate the effect of such practices on soil properties at different depths and on nutrient contents in chestnut leaves. To reach our goal, 6 pits down to 30 cm soil depth were dug in European chestnut (*Castanea sativa* Mill.) stands where the organic residues such as burrs, leaves and pruned materials are yearly removed (REM), and 9 pits in stands where the organic residues are chopped and left on soil surface (CONS). Both practices began about 30 years ago. The nutrient contents in leaves were assessed on chestnut trees close to each pit. Our findings showed a more intense soil development in CONS than in REM. At soil surface, CONS had thicker mineral horizon than in REM probably due to the protection acted by the organic residues against erosion. In subsoil, organic residues conservation promoted the organic C (soil organic C, water-extractable organic C, fulvic-like C and soil organic C stock) accumulation which further promoted horizon development. Since the role of soil organic matter (SOM) as source of nutrients, the subsoil of CONS study sites had a greater stock of Ca, P and S and a higher exchangeable Ca content than REM. Furthermore, the higher SOM and nutrient contents in CONS subsoil compared to REM promoted the microbial respiration. The organic residues conservation effects on the surface soil horizon properties were, instead, negligible. Despite the generally better soil conditions in CONS than in REM, no differences were observed for the leaf nutrient contents likely due to the chestnut trees adaptability also to the scarcely fertile soils. Overall, our findings highlighted the importance of organic residues conservation for the improvement in deeper soil horizons of the chemical and biological fertility in chestnut plantations. Furthermore, this research pointed out to pay more attention on subsoil since it is a good indicator of the changes caused by external factors.

The final version has been published in Catena Volume 198, March 2021, n. 105021, DOI <https://dx.doi.org/10.1016/j.catena.2020.105021>

© 2020 Elsevier. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0>)

Highlights

- Organic residues removal (ORR) effect on chestnut stand soil properties was tested
- ORR limits the organic C content and nutrient stocks (C, N, P and S) in subsoil
- CONS management promoted the development of AB and Bw horizons
- The limited organic C deepening due to ORR penalised the subsoil microbial activity
- Organic residues should be conserved to maintain soil quality in chestnut stands

1 **Mid-term (30 years) changes of soil properties under chestnut stands due to organic residues**
2 **management: An integrated study**

3

4 Mauro De Feudis*^{1,2}, Gloria Falsone^{1,2}, Livia Vittori Antisari^{1,2}

5

6 ¹Department of Agricultural and Food Sciences, Alma Mater Studiorum - University of Bologna,
7 Via Fanin, 40; 40127 Bologna, Italy;

8 ²Centro Sperimentale per lo Studio e l'Analisi del Suolo (CSSAS), Alma Mater Studiorum -
9 University of Bologna, Bologna, Italy.

10

11

12

13 *Corresponding author:

14 Mauro De Feudis

15 Department of Agricultural and Food Sciences

16 Alma Mater Studiorum – University of Bologna

17 Via Fanin, 40; 40127 Bologna Italy

18 e-mail: mauro.defeudis2@unibo.it

19

20

21

22

23

24

25

26

27 **Abstract**

28 Chestnut plantations are worldwide distributed and they are often subjected to intensive
29 management practices such as the removal of the organic residues from the soil surface. The present
30 study aimed to investigate the effect of such practices on soil properties at different depths and on
31 nutrient contents in chestnut leaves. To reach our goal, 6 pits down to 30 cm soil depth were dug in
32 European chestnut (*Castanea sativa* Mill.) stands where the organic residues such as burrs, leaves
33 and pruned materials are yearly removed (REM), and 9 pits in stands where the organic residues are
34 chopped and left on soil surface (CONS). Both practices began about 30 years ago. The nutrient
35 contents in leaves were assessed on chestnut trees close to each pit. Our findings showed a more
36 intense soil development in CONS than in REM. At soil surface, CONS had thicker mineral horizon
37 than in REM probably due to the protection acted by the organic residues against erosion. In
38 subsoil, organic residues conservation promoted the organic C (soil organic C, water-extractable
39 organic C, fulvic-like C and soil organic C stock) accumulation which further promoted horizon
40 development. Since the role of soil organic matter (SOM) as source of nutrients, the subsoil of
41 CONS study sites had a greater stock of Ca, P and S and a higher exchangeable Ca content than
42 REM. Furthermore, the higher SOM and nutrient contents in CONS subsoil compared to REM
43 promoted the microbial respiration. The organic residues conservation effects on the surface soil
44 horizon properties were, instead, negligible. Despite the generally better soil conditions in CONS
45 than in REM, no differences were observed for the leaf nutrient contents likely due to the chestnut
46 trees adaptability also to the scarcely fertile soils. Overall, our findings highlighted the importance
47 of organic residues conservation for the improvement in deeper soil horizons of the chemical and
48 biological fertility in chestnut plantations. Furthermore, this research pointed out to pay more
49 attention on subsoil since it is a good indicator of the changes caused by external factors.

50

51 **Keywords:** chestnut management; mountain soil; organic carbon dynamics; litter floor, soil
52 development

53

54 **1. Introduction**

55 Chestnut plantations for nut production are widely distributed throughout the world (Conedera et
56 al., 2016). Among the European countries, Italy is the largest sweet chestnut producer and importer
57 with about 52,000 Mg (FAOSTAT, 2017). In Italy managed chestnut stands showed a decrease
58 from the second half of the twentieth century to the present time principally due to the rural
59 depopulation (Zlatanov et al., 2013), to the presence of more competitive cultures, such as corn and
60 potatoes, and to the spread of the chestnut blight (*Chryphonectria parasitica* [Murr.] Barr) and ink
61 (*Phytophthora cambivora* [Petri] Buism.) diseases (Pezzi et al., 2011). More recently, a pivotal role
62 in the chestnut stands abandonment could have played by the reduced precipitations due to the
63 climate change (Pieri et al., 2017) which have promoted the spread of the chestnut diseases
64 (Waldboth and Oberhuber, 2009). However, in the last decades old chestnut groves have been
65 recovered, also by replacement of coppices with plantations (Beccaro et al., 2009), with an increase
66 of both socio-economic and environmental interest (Martín et al., 2017).

67 Chestnut stand agroecosystems represent today an important landscape component in the
68 mountainous regions of the European Mediterranean basin. The management of the chestnut stands
69 in a sustainable way is essential to improve or to maintain the soil quality in these areas. Soil
70 organic matter (SOM) is the most frequently used indicator of soil quality, as it affects some soil
71 biological, physical and chemical properties, followed by nutrient content and soil microbial
72 biomass and respiration (Zornoza et al., 2015). Unfortunately, studies concerning the effect of
73 above ground biomass management on SOM are missing for the chestnut ecosystems. Conversely,
74 taking in account other tree species, several are the studies monitoring the influence of forest
75 harvesting on soil organic carbon pool (e.g., Homann et al., 2001; Nave et al., 2010). For example,
76 previous studies (James and Harrison, 2016; Johnson and Curtis, 2001; McDonnell et al., 2013;
77 Zetterberg et al., 2016) highlighted how the intense harvesting practices aimed to remove most of
78 the tree biomass significantly reduce the amounts of C and other essential nutrients such as Ca^{2+} ,

79 Mg^{2+} and K^+ in soil. However, in forest ecosystems another key driver of the soil C is the litter
80 layer (Maes et al., 2019). Indeed, litter removal practices could alter the pool sizes of organic
81 compounds and nutrients, and influence the rates of biogeochemical processes which could
82 potentially compromise soil fertility (Xiong et al., 2008). Several studies have found that the
83 presence of plant litter can significantly increase both the soil resistance to flowing water erosion
84 and the water holding capacity (Giménez and Govers, 2008; Liu et al., 2017). Furthermore, the litter
85 floor could be incorporated into the mineral soil horizons through the soil fauna activities (Yavitt et
86 al., 2015), increasing soil aggregate stability, SOM amount and stabilisation, and decreases soil bulk
87 density (Mandal et al., 2004; Mitchell et al., 2018). Since the positive relationship between soil
88 microbial biomass and SOM (van Leeuwen et al., 2017) the removal or alteration of soil organic
89 matter inputs due to the different forest management might change the size and the activity of the
90 soil microbial biomass.

91 Because of the role of soil on plant health, changes of the soil properties could affect plant nutrient
92 contents. For example, a reduced plant nutritional status was found in soil characterized by poor
93 SOM (Nielsen et al., 2014) and nutrient contents (Demchik and Sharpe, 2000) and microbial
94 biomass (Van Der Heijden et al., 2008).

95 In order to implement those agroforestry practices able to preserve or improve tree health and soil
96 quality, we tested the effect of 30 years chestnut organic residues management practices on soil
97 thickness, amount and quality of SOM, nutrient content and microbial biomass and its activity of
98 top- and subsoil layers, and on nutrient concentrations in chestnut tree leaves.

99 To achieve this goal, we selected chestnut stands where the organic residues such as burrs, leaves
100 and pruned materials are yearly removed, and chestnut stands with more conservationist practices
101 where organic residues are chopped and left on the soil surface. It was hypothesized that the organic
102 residues removal *i*) declines *solum* thickness; *ii*) reduces the contents of soil organic carbon and of
103 SOM pools, and the soil organic carbon stock; *iii*) decreases the stocks of N, Ca, Mg, Na, K, Fe, P

104 and S and the availability of Ca, Mg, Na and K; *iv*) has negative impact on soil microbial biomass
105 and respiration; *v*) reduces the nutrient contents in plants.

106

107 **2. Materials and methods**

108 *2.1. Study sites*

109 The study sites included four farms (Figure 1 and Table 1) with uneven-aged chestnut (*Castanea*
110 *sativa* Mill) stands for fruit production. The chestnut stands were recovered about 30 years ago
111 through a more or less severe pruning of the plants, according to the age of the trees and to the
112 severity of parasitic attacks, necessary to stimulate the growth of a new and vigorous canopy.
113 Furthermore, the recovery actions included the selection of the healthiest trees, to maintain a tree
114 spacing of about 10 × 10 m, whose dry, senescent, sick parts and overlapping branches were yearly
115 removed in order to open the canopy and allowing a better fruits production on sunny branches
116 (Figure 1). Therefore, the farms included uneven-aged chestnut stands whose average size of trees
117 was about 30-35 cm reflecting the 40-years old age of shoots. In all study sites the chestnut
118 production was about 40 kg tree⁻¹.

119 The study sites are located on the northern Apennine chain in Italy (Figure 1), their locations and
120 climatic characteristics are reported in Table 1. In all investigated sites, the chestnut plantations
121 stand at an altitude ranging from 650 to 750 m above sea level, they are exposed to south, south–
122 east and on slopes with an inclination ranging from 15° to 30°. In all study sites the soil developed
123 from sandstone with similar elemental compositions (ISPRA, 1999).

124

125 *2.2. Experimental design*

126 The four farms involved in this study were selected in order to compare the soil properties of
127 chestnut stands in which the organic residues removal (ORR) has been made or not. In two farms,
128 the chestnut stands were subjected to yearly ORR, hereafter called REM, due to the use of small
129 blowing devices during the fruits harvesting which remove from the soil surface both fruits and

130 other organic residues. In two other farms, management was more conservationist which consist to
131 leave on the soil surface the organic residues, hereafter called CONS, which are chopped. In CONS
132 study sites the fruits harvesting was manual or using nets put on trees. Both ORR and the more
133 conservationist practices began at the recovery time of the chestnut stands, therefore about 30 years
134 ago, and concerned the whole surface of the stands.

135 The soils were classified as Typic Udorthents for REM study sites and Typic Dystrudepts for
136 CONS ones (Soil Survey Staff, 2014). The description of representative soil profiles are reported in
137 Table S1 of the supplementary materials (Regione Emilia Romagna, 2000).

138 In spring 2017, fifteen pits down to 30 cm soil depth were dug (6 pits for REM and 9 for CONS).
139 Since the high soil variability that generally occurs in mountainous areas (Hoffmann et al., 2014), a
140 different number of pits were dug for each study area to get a better representation of the soil on the
141 entire fields. Each pit was dug at a distance of at least 1 m away from the stem of the trees.

142

143 *2.3. Sampling and analysis*

144 In each dug pit, each identified genetic horizon down to 30 cm of soil depth has been sampled by
145 collecting at least 2 kg of sample and, once in the laboratory, it was air dried and sieved (<2 mm).
146 For the determination of the bulk density (BD), additional soil samples were taken using steel
147 cylinder cores with an internal diameter of 5.0 cm and 5.0 cm height from the 0-10, 10-20 and 20-
148 30 cm soil depth.

149 In order to evaluate the effect of ORR practice on nutrient contents in chestnut leaves, the
150 uppermost fully expanded leaves and without visible injury symptoms were collected from
151 randomly selected branches. Specifically, from 20 to 30 leaves were picked from each tree. The
152 leaves were collected from 4 healthy trees located close to each mini-pit and mixed to composed
153 one sample per pit. The leaf samples were oven-dried at 60 °C until constant weight and ground.

154 The soil particle size distribution was obtained by the pipette method after dispersion of the sample
155 with a sodium hexametaphosphate solution (Gee and Bauder, 1986). The soil samples collected for

156 the determination of BD were oven-dried at 105 °C for 24 h, and then weighed (Blake and Hartge,
157 1986). The dry weight was thereafter divided by volume of the cores. The obtained BD was
158 adjusted by subtracting the mass and the volume of skeleton and roots in order to determine the BD
159 of the fine earth fraction.

160 The soil pH was determined potentiometrically in a 1:2.5 solution ratio in both deionized water
161 ($\text{pH}_{\text{H}_2\text{O}}$) and 1M KCl (pH_{KCl}) suspension. Total organic C (TOC) and total nitrogen (TN) were
162 determined by a CHN elemental analyser (EA 1110 Thermo Fisher, USA) without to be pre-treated
163 with hydrochloric acid due to the absence of carbonates. The total amount of Ca, Mg, Na, K, Fe,
164 Mn, P and S were determined by inductively coupled plasma optical emission spectrometer (ICP-
165 OES, Ametek, Spectro Arcos, Germany) after aqua regia extraction (Vittori Antisari et al., 2014).

166 The cation exchange capacity (CEC) and the exchangeable cation contents were determined
167 according to the method proposed by Orsini and Rémy (1976) and modified by Ciesielski and
168 Sterckeman (1997) using 0.017 M hexamminecobalt(III)chloride as extracting solution and
169 measuring the amounts of Co and exchangeable cations by ICP-OES.

170 Different soil C pools characterized by increasing persistence in soil were sequentially separated
171 according to Agnelli et al. (2014) with some modifications. Briefly, a 1:10 solid:water suspension
172 was shaken on a horizontal shaker for 16 hours at 25°C, centrifuged and the supernatant was separated
173 from the precipitate. From the supernatant two C pools have been further separated by sieving it at
174 53 μm : the particles $>53 \mu\text{m}$ which represented the particulate organic matter (POM), namely plant
175 and animal residues at different stages of decomposition, and the supernatant $<53 \mu\text{m}$ which was
176 further filtered through 0.45 μm filter paper and it represented the water-extractable organic matter
177 (WEOM), namely the more mobile and labile organic matter pool. To the remaining precipitate into
178 the centrifugation tubes, 0.1M NaOH solution was added and the samples were shaken for 24 hours
179 at 25°C and then again centrifuged. The NaOH extract was passed through a 0.45 μm polycarbonate
180 filter, while the remaining precipitate, containing the humin-like (Hum) compounds was washed
181 using deionized water to remove the excess of Na until the pH of the rinsed solution was ≤ 7 . The

182 0.45 μm filtered NaOH extract was acidified to about pH 1.5 with 6M HCl and allowed to settle
183 overnight to separate humic acid-like (HA) and fulvic acid-like (FA) compounds and centrifuged.
184 To remove the excess of Na from the obtained fractions, the supernatant (FA) was dialyzed through
185 1000 Da cut-off membranes (Spectra/Por® Dialysis membrane) against distilled water, while the
186 residual (HA) was washed with 0.002 M HCl. Both purified fractions were freeze-dried. The POM
187 and Hum fractions were instead dried at 40 °C. The organic C (OC) and N contents of POM, FA,
188 HA and Hum were determined by a CHN elemental analyser (EA 1110 Thermo Fisher, USA). The
189 OC and N contents of WEOM were recorded by a TOC-V CPN total organic carbon analyzer
190 (Shimadzu, Japan).

191 The C and N concentrations in leaf samples were measured by the CHN elemental analyser (EA
192 1110 Thermo Fisher, USA). While, for the determination of Ca, Mg, K, Mn, Na, P and S, the
193 ground leaves were mineralized in 4:1 HNO₃ (65 % v/v): H₂O₂ (30% v/v) solution in a microwave
194 oven (START D Microwave Digestion System; Milestone Inc., Sorisole, Bergamo, Italy). The
195 concentrations of the selected elements in the extracts were measured by ICP-OES.

196 The soil respiration during 28 days at 25 °C of incubation was determined according to Falsone et
197 al. (2015) after conditioning of the samples at 60 % of their water holding capacity (WHC) and pre-
198 incubating them for 3 days at 4 °C. After 1-3-7-10-14-21-28 days the beginning of incubation,
199 the amount of CO₂ emitted from incubated soils was measured by alkali (1 M NaOH solution)
200 absorption of the evolved CO₂ and titration of the residual OH⁻ with a standardized HCl solution.
201 While the soil basal respiration (SBR) of each soil sample was computed as the average of the
202 values measured during the incubation period, the cumulative soil basal respiration (RCUM) was
203 expressed as the total amount of CO₂ evolved during the 28 days of incubation.

204 Soil microbial biomass-C (C_{mic}) and microbial biomass-N (N_{mic}) were measured on soil samples
205 at 60% of WHC using chloroform fumigation extraction method with 0.5 M K₂SO₄ solution (Vance
206 et al., 1987). Both fumigated and non-fumigated extracts were analysed using a TOC-V CPN total
207 organic carbon analyzer (Shimadzu, Japan). C_{mic} was calculated as E_C*2.64, where E_C = difference

208 between organic C extracted from fumigated soils and organic C extracted from non-fumigated
209 soils. N_{mic} was calculated as $E_N * 2.22$, where E_N = difference between total N extracted from
210 fumigated soils and total N extracted from non-fumigated soils (Vance et al., 1987).

211 The metabolic (qCO_2) and mineralization (qM) quotients were calculated according to the following
212 equations (Eq. 1 and 2):

$$213 \quad qCO_2 = 100 * SBR / C_{mic} \quad (1)$$

$$214 \quad qM = 28 \text{ days cumulative respiration (RCUM)} / TOC \quad (2)$$

215

216 *2.4. Data analysis*

217 In order to evaluate the effect on soil of the organic residues conservation or removal, the results of
218 the chemical and biological properties are presented separating epipedon (superficial layers) and
219 endopedon (deeper layers), in particular, epipedon (Epi) included superficial A horizons (i.e., A,
220 A1, A2), while endopedon (Endo) included other deeper transition and mineral horizons (i.e., AC,
221 AB, Bw, BC and C).

222 The stock of OC, TN and nutrients are presented for fixed soil-depth intervals (0–10, 10–20 and
223 20–30 cm), taking into account the BD value of each soil layer at fixed depth and the OC, TN and
224 nutrients content of each genetic horizon averaged on the basis of horizon thickness.

225 Two-way analysis of variance was performed to assess the effect of chestnut stand management on
226 the selected physical, chemical and biological parameters and the variation of the selected variables
227 with soil depth. Instead, for the plant samples, one-way analysis of variance was carried out to
228 assess the effect of chestnut stand management on the leaf nutrient concentrations. Since the
229 graphical analysis of residuals showed the absence of normality and homoscedasticity, the all the
230 data were processed by the Kruskal-Wallis rank sum test to identify statistically significant
231 differences and Holm test was performed as multi-comparison test ($p < 0.05$). The results
232 presented are based on mean values and their standard error. The data were analyzed using R
233 software (R Core Team, 2016).

234

235 **3. Results**

236 *3.1. Morphological properties of chestnut stand soils*

237 Our results showed the differences in the *solum* thickness and horizons sequences between REM
238 and CONS study sites.

239 The mean thickness of superficial A horizons (i.e., A, A1, A2), called Epi horizons, was in fact of
240 5.1 and 9.7 cm in REM and CONS, respectively. The deeper mineral horizons (i.e., AC, AB, Bw,
241 BC and C), called Endo horizons, showed a mean thickness of 12.4 and 20.2 cm in REM and
242 CONS, respectively (Table 2). Both in REM and CONS Epi horizons were thinner than Endo ones
243 ($p < 0.05$), and between both practices, CONS allowed a thicker Epi than REM ($p < 0.05$). Similarly,
244 for Endo, CONS showed a larger thickness compared to REM. However, it needs to take in mind
245 that the horizons sequence down to 30 cm was A/AC/C in REM soils, while was A1/A2 or AB or
246 Bw or BC in CONS ones. The horizons sequence of *solum* thus showed weak development in REM
247 because of the lack of B horizons, and furthermore the thickness of *solum* was lower in REM than
248 in CONS ones ($p < 0.05$).

249

250 *3.2. Physicochemical properties of chestnut stand soils*

251 The occurrence of differences in particle size distribution, BD values, pH, CEC and concentration
252 of exchangeable bases between Epi and Endo in the soils of REM and CONS study sites has been
253 determined.

254 Our findings showed that the soils of the selected chestnut orchards have sandy–loam/loam texture,
255 with lower amount of clay in CONS than in REM (Table 2). Furthermore, REM soils had similar
256 particle distribution along the soil depth; in CONS soils, instead, Endo horizons had higher silt
257 content than Epi ones ($p < 0.05$). The bulk density, on average, ranged from 1.02 to 1.47 g cm⁻³, but
258 significant differences neither between soil depth nor between soil management practices have been
259 found.

260 The $\text{pH}_{\text{H}_2\text{O}}$ values ranged from 4.8 to 5.4, the pH_{KCl} ranged from 3.2 to 4.1 with some differences in
261 CONS where the Endo showed higher values compared to Epi (Table 2).

262 In both managements, the CEC did not change between Epi and Endo, however, the soils under
263 REM management showed higher CEC than those under CONS (Figure 2I). Regarding to the
264 exchangeable bases (Ca, Mg, K, Na), the ExCa always showed higher values in Epi than in Endo
265 (Figure 2II), the ExK had higher concentrations in Epi than in Endo only in REM (Figure 1III), and
266 no differences occurred between Epi and Endo for ExMg and ExNa (Figure 2IV, V). Comparing the
267 two management practices, among the exchangeable bases, only the ExCa showed some
268 differences, in particular higher ExCa values were found in Endo of CONS sites compared to Endo
269 of REM (Figure 2II). As expected, the base saturation (BS) values were higher in Epi compared to
270 Endo but only in REM soils (Figure 2VI). In CONS soils, the BS values were similar between Epi
271 and Endo, and no differences occurred between the two management practices (Figure 2VI).

272

273 *3.3. SOM pools of chestnut stand soils*

274 As expected, in both chestnut managements, most of the studied pools of C and N cycles (i.e., total,
275 labile, humic-like and humin substances) were higher in the Epi than in the Endo (Figure 3, Figure
276 4, Figure S1). Comparing the two chestnut stand managements, some differences occurred for the
277 Endo. Specifically, the concentrations of TOC and of organic C related to WEOM and FA were
278 higher in CONS than in REM (Figure 3I, III, Figure 4II). Similarly, the contents of TN and nitrogen
279 related to WEOM were greater in CONS than in REM (Figure 3II, IV).

280

281 *3.4. Microbial biomass amount and respiration of chestnut stand soils*

282 Regarding to the soil microbial biomass, in both chestnut stand managements C_{mic} , N_{mic} , SBR
283 and RCUM values were higher in Epi than in Endo, while no differences occurred for q_R and q_M
284 (Figure 5I–VI). Through the comparison of the two chestnut stand managements, some differences

285 occurred for RCUM and qM. In particular, the RCUM was higher in Endo of CONS than in REM
286 (Figure 5IV), the qM was higher in CONS than in REM both for Epi and Endo (Figure 5VI).

287

288 *3.5. Soil organic C and nutrients stocks, and nutrient concentrations in chestnut tree leaves*

289 The stock of elements of organic origin (i.e., C, N, P and S) generally decreased along soil depth,
290 and in deeper layers (10–20 and 20–30 cm) in CONS chestnut stands had higher values than in
291 REM (Table 3). Nutrients mainly derived by parent material (Ca, Mg, Na, K and Fe) did not instead
292 show significant differences between the two different managements (Table 3). However, a certain
293 Mg and Fe redistribution occurred in deeper layers (10-20 cm and 20-30 cm) of CONS soils, having
294 higher stock values than in 0-10 cm layer, while in REM soils Na stocks was higher in deeper layers
295 than in 0-10 cm (Table 3).

296 Although some differences in soil elements stock occurred, the concentrations of C, N, Ca, Mg, K,
297 Mn, Na, P and S in chestnut leaves did not differ between the two type of organic residue
298 managements (Table 4).

299

300 **4. Discussion**

301 *4.1. Organic residues conservation promotes soil thickness and development in chestnut stands*

302 In the present study, the ORR practice reduced the thickness of superficial A horizons (5.1 vs 9.7
303 cm in REM and CONS, respectively) which could be attributed to higher soil erosion that probably
304 takes place in the former than in the latter study sites. Indeed, as reported by several works (e.g.,
305 Dick et al., 1997; Křeček et al., 2019; Liu et al., 2017) the presence of the litter layer limits the
306 surface runoff and soil erosion. The highest soil erosion that likely occurs in REM study sites could
307 also limit the thickness of *solum* due to losses of pedogenized material (Mokma et al., 1996).
308 Consequently, the formation of more developed B horizons or transitional AB ones was prevented
309 in REM soils. Our findings are in accordance to Świtoniak (2014) and Jankauskas and Fullen
310 (2002) which, through a pedological investigation of soils subjected to different erosion severity,

311 observed weakly developed soil horizon sequences in eroded sites. In contrast, the reduced erosion
312 in CONS study sites might promoted the development of their soils allowing to define the CONS
313 soils as Inceptisol while those of REM as weakly developed Entisols.

314

315 *4.2. Organic residues conservation promotes soil organic matter content and stock in chestnut* 316 *stands*

317 As reported in several works carried out on Italian mountains soils (e.g., De Feudis et al., 2017;
318 Vittori Antisari et al., 2016), the TOC content was higher in organo–mineral horizons (Epi) than in
319 deeper ones due to input of OC from the litterfall (Boča and Miegroet, 2017). However, the ORR
320 practice reduced the soil OC content, in particular we found lower TOC content in Endo of REM
321 than in CONS, in agreement with early studies (e.g., Achat et al., 2015; Tian et al., 2010). This fact
322 might suggest how the ORR limits the incorporation of the organic matter in the subsoil preventing
323 any vertical redistribution of organic C. Conversely, the conservation of the organic residues allows
324 it. Specifically, our findings indicated that the incorporation of the organic matter in the subsoil of
325 CONS could be assigned to the redistribution along depth of WEOM-C and FA-C fractions which
326 showed higher concentrations in Endo of CONS than in that of REM. Both the water–soluble and
327 the fulvic–like substances represent an important source of organic C for the deeper layers of the
328 soils (Aran et al., 2001; Kaiser and Kalbitz, 2012).

329 Because of the importance of SOM on the development of the subsoil horizons even in poorly
330 developed soils such as Entisols and Inceptisols (Falsone et al., 2012), the deepening of the organic
331 substances in CONS, together with the limited soil erosion, likely promoted the development of AB
332 and Bw horizons.

333 Although the lower clay content in CONS than in REM, the OC stock was larger in 10–30 cm soil
334 depth section of CONS than in that of REM. This fact emphasizes the leading role of organic matter
335 input amount and rate on soil OC stock (Orgill et al., 2017). Since the subsoil OC generally has a
336 higher residence time compared to topsoil (Jenkinson et al., 2008; Trumbore, 2009), the ORR

337 seemed to reduce the role of soil as stable sink of C reducing the OC stocks below 10 cm of soil
338 depth.

339 The modification of TOC content and OC stock only in Endo and in 10–30 cm soil depth interval,
340 respectively, might indicate a high sensibility of subsoil OC to management practices. The high
341 sensibility of subsoil OC to external factors, such as management practices, is in accordance with
342 recent studies (Jia et al., 2019; Leuschner et al., 2014; Mobley et al., 2015; Steinmann et al., 2016)
343 which observed meaningful changes of subsoil OC stock caused by climate warming, reforestation
344 and soil managements such as tillage and fertilization.

345

346 *4.3. Organic residues conservation increases soil nutrient contents in chestnut stands*

347 The similar parent material on which the studied soils developed explains the similar stock of
348 nutrients, with exception of P and S, which exhibited higher values in 10 – 30 soil depth section of
349 CONS than of REM probably due to the higher amount of organic matter. Indeed, concerning P, in
350 forest soils most of the soil P is made up of organic P forms (Adams et al., 2018; De Feudis et al.,
351 2016). As the organic P is the most important P source in forest ecosystems (Turner et al., 2014)
352 and the total P concentration generally decline over time (Chen et al., 2015), litter recycling is
353 essential to avoid an accelerated loss of soil P. The nutrient recycling process seemed to affect also
354 S. In fact, similarly to P, the total amount of S in 10 – 30 cm soil depth interval was higher in
355 CONS than in REM. As for P, the organic S compounds represent the most abundant S forms in
356 soils (e.g., Likens et al., 2002; Vannier and Guillet, 1994). Regarding to the other nutrients, such as
357 exchangeable Mg, K and Na, their amounts did not change between the two managements.
358 Exchangeable Ca, instead, had higher contents in Endo of CONS than in that of REM, following the
359 trend of OC content. This is in agreement with Vittori Antisari et al. (2013) who reported that SOM
360 is an important source of available Ca in chestnut stands. The importance of Ca biocycling was
361 further stressed by the lack of differences of Ca stock. In fact, in mountain acid soils the
362 redistribution of Ca-mineral derived is often prevented by the limited Ca-mineral weathering

363 (Ghobadi and Mousavi, 2014), and available Ca derived mainly from litter recycling (Haynes and
364 Swift, 1986; Johnson et al., 1994). Although the higher amounts of some nutrients in CONS than in
365 REM, the lack of differences occurring for the nutrient contents in plant leaves would suggest the
366 unresponsive of chestnut to soil fertility and, therefore, the ability of chestnut plants to grow also on
367 poor sandy-rich soils (Oosterbaan, 1998). The ability of chestnut trees to healthily grow also in soils
368 characterized by low amount of nutrients is confirmed by Ribeiro et al. (2019) which did not find
369 differences in leaf nutrient contents between fertilized and unfertilized *Castanea sativa* plants.

370

371 *4.4. Organic residues conservation improves soil microbial respiration in chestnut stands*

372 The RCUM in Endo of CONS study sites was higher than in REM likely due to the higher C
373 content associated to WEOM (WEOM-C). In fact, because of the most labile organic compounds
374 represent the main substrate for soil microbial communities (Rees and Parker, 2005) and changes of
375 WEOM-C quantity affect both microbial activities and composition (Drenovsky et al., 2004), the
376 enhance of WEOM-C can promote the heterotrophic soil respiration (Rees and Parker, 2005).
377 Furthermore, since the positive effect of WEOM-C on soil microbial activities, the higher qM in
378 CONS than in REM might be attributed to the priming effect, namely a short-term change in
379 microbial mineralization of SOC in response to labile OC inputs (Kuzyakov, 2010). Unlike RCUM
380 and qM, both the microbial biomass and qCO₂ have shown similar values between the two chestnut
381 orchard managements. The lack of differences about microbial biomass and qR might be attributed
382 to the fact that they are driven not only by the amount of TOC but also by other soil properties such
383 as BD, soil moisture and temperature, and C:N ratio (Ren et al., 2018; Stefanowicz et al., 2016; van
384 Leeuwen et al., 2017; Xu et al., 2007) which, in our case, were similar among the study sites.

385

386 *4.5. Overview of the effects of organic residues management on soil properties in chestnut stands*

387 In the present study, the soil erosion occurring in the mid-term (30 years) in REM study sites due to
388 the periodic ORR might promoted the thinning of the A horizons and prevented the development of

389 B or transitional AB horizons. Moreover, ORR practice reduced the SOM content in subsoil of
390 REM study sites because this practice impeded the inclusion of the organic material in soil mineral
391 horizons. Therefore, because of the importance of SOM on the development of soil horizons, the
392 limited amount of organic residues left on soil surface in REM could have further prevented the
393 development of B or transitional AB horizons.

394 The presence of chopped organic residues on soil surface of CONS promoted a greater soil OC
395 stock than in REM. However, it is noteworthy to highlight that the differences about soil OC stock
396 were observed in subsoil layers indicating how the conservation of the organic residues enhanced
397 SOM stabilization. Because of the role of organic matter on soil aggregation, fertility and to store
398 water (de Paul Obade and Lal, 2016), the decline of OC content and stock in subsoil due to ORR
399 practice could instead cause a reduction of soil quality. For example, the higher OC stock in Endo
400 horizons of CONS than in that of REM resulted in a higher amount of those nutrients, such as P and
401 S, whose cycling is related to SOM. Moreover, our findings showed that the conservation of the
402 organic residues on soil surface improves the soil also from the biological point of view. In fact, the
403 higher SOM content and, therefore, the higher amount of labile OC in CONS than in REM
404 increased the activity of soil microbes which are important player on the soil nutrient cycling.

405 Hence, a chestnut stand management based on the conservation of the litter floor should be
406 supported in order to improve the soil quality and, in particular, to increase the volume of *solum*, the
407 SOM and nutrient contents and the soil microbial respiration. However, though not considered in
408 the present work, CONS management could have some negative effects on the spreading of some
409 chestnut diseases. For example, the increase of SOM content could benefit the proliferation of the
410 ink disease (*Phytophthora cinnamomi*) because of its saprotrophic nature (Fonseca et al., 2004).
411 Furthermore, since in CONS study sites the organic residues are chopped, this practice could
412 increase the production costs. Finally, since the ORR and the chipping of the litter floor are yearly
413 performed, both studied managements could have some negative effects on the proliferation of
414 *Torymus sinensis* Kamiyo. In fact, in order to maintain this biological control agent of the chestnut

415 gall wasp, the plant material bearing galls (branches, suckers) should not be removed from the stand
416 at least for two years (Ferracini et al., 2015). More researches need however on the assessment of
417 the phytosanitary consequences and on the role that the SOM and microbial activity can have on the
418 enhancement of chestnuts disease resistance.

419

420 **5. Conclusion**

421 In the mid-term (30 years), the common practice of ORR in recovered century chestnut stands has
422 negative impacts on soil quality. In particular, compared to chestnut stands where the organic
423 residues are chopped and left on soil surface, the ORR practices prevent the development of AB and
424 Bw soil horizons. The limited soil development in REM was likely due to an accelerated soil
425 erosion leading to loss of pedogenized material, and to a negligible incorporation of organic matter
426 from soil surface into the subsoil limiting mineral weathering. Furthermore, the reduced stock of P,
427 S, and organic C and the reduced amount of exchangeable Ca, cumulative basal respiration and
428 mineralization quotient in subsoil of the chestnut stands subjected to ORR practice might suggest
429 that the conservation of the organic residues is necessary to maintain the soil chemical and
430 biological fertility in mountainous ecosystems. The improved soil quality, together with the
431 unchanged nutrient contents in chestnut leaves, would suggest that the switch from the common
432 management practices in more conservative ones matches with the aims of those agroforestry
433 practices addressed to preserve or improve both tree health and environmental quality. Furthermore,
434 the present study highlighted the higher sensibility of the subsoil to the management practices
435 compared to the topsoil. Therefore, for the studies about the effect of management practices in
436 forests ecosystems located in mountainous areas, we suggest to pay more attention on subsoil
437 properties since it is a good indicator of the changes caused by external factors.

438

439 **Acknowledgments**

440 This work was partially funded by the European Agricultural Fund for Rural Development through
441 the Rural Development Programme - Emilia Romagna, focus area 5E, Action Group: CASTANI-
442 CO, support application number 5015571, project name: Carbon sequestration in the system of
443 chestnut orchards for fruit production. We would like to thank Dr. Carla Scotti of I.TER soc. coop a
444 r.l. for her help during soil pits description and sampling.

445

446 **References**

- 447 Achat, D.L., Fortin, M., Landmann, G., Ringeval, B., Augusto, L., 2015. Forest soil carbon is
448 threatened by intensive biomass harvesting. *Sci. Rep.* 5. <https://doi.org/10.1038/srep15991>
- 449 Adams, J.L., Tipping, E., Thacker, S.A., Quinton, J.N., 2018. An investigation of the distribution of
450 phosphorus between free and mineral associated soil organic matter, using density
451 fractionation. *Plant Soil* 427, 139–148. <https://doi.org/10.1007/s11104-017-3478-4>
- 452 Agnelli, A., Bol, R., Trumbore, S.E., Dixon, L., Cocco, S., Corti, G., 2014. Carbon and nitrogen in
453 soil and vine roots in harrowed and grass-covered vineyards. *Agric. Ecosyst. Environ.* 193, 70–
454 82. <https://doi.org/10.1016/j.agee.2014.04.023>
- 455 Aran, D., Gury, M., Jeanroy, E., 2001. Organo-metallic complexes in an Andosol: A comparative
456 study with a Cambisol and Podzol. *Geoderma* 99, 65–79. [https://doi.org/10.1016/S0016-](https://doi.org/10.1016/S0016-7061(00)00064-1)
457 [7061\(00\)00064-1](https://doi.org/10.1016/S0016-7061(00)00064-1)
- 458 Beccaro, G.L., Mellano, M.G., Barrel, A., Trasino, C., 2009. Restoration of old and abandoned
459 chestnut plantations in Northern Italy. *Acta Hort.* 815, 185–190.
460 <https://doi.org/10.17660/ActaHortic.2009.815.24>
- 461 Blake, G.R., Hartge, K.H., 1986. Bulk density, in: *Methods of Soil Analysis, Part 1*. Madison, pp.
462 363–375.
- 463 Boča, A., Miegroet, H. Van, 2017. Can carbon fluxes explain differences in soil organic carbon
464 storage under aspen and conifer forest overstories? *Forests* 8. <https://doi.org/10.3390/f8040118>
- 465 Chen, C.R., Hou, E.Q., Condrón, L.M., Bacon, G., Esfandbod, M., Olley, J., Turner, B.L., 2015.

466 Soil phosphorus fractionation and nutrient dynamics along the Cooloola coastal dune
467 chronosequence, southern Queensland, Australia. *Geoderma* 257–258, 4–13.
468 <https://doi.org/10.1016/j.geoderma.2015.04.027>

469 Ciesielski, H., Sterckeman, T., 1997. Determination of cation exchange capacity and exchangeable
470 cations in soils by means of cobalt hexamine trichloride. Effects of experimental conditions.
471 *Agronomie* 17, 1–7. <https://doi.org/10.1051/agro:19970101>

472 Conedera, M., Tinner, W., Krebs, P., de Rigo, D., Caudullo, G., 2016. *Castanea sativa* in Europe:
473 distribution, habitat, usage and threats, in: *European Atlas of Forest Tree Species*. p. e0125e0+.

474 De Feudis, M., Cardelli, V., Massaccesi, L., Bol, R., Willbold, S., Cocco, S., Corti, G., Agnelli, A.,
475 2016. Effect of beech (*Fagus sylvatica* L.) rhizosphere on phosphorous availability in soils at
476 different altitudes (Central Italy). *Geoderma* 276, 53–63.
477 <https://doi.org/10.1016/j.geoderma.2016.04.028>

478 De Feudis, M., Cardelli, V., Massaccesi, L., Lagomarsino, A., Fornasier, F., Westphalen, D.J.,
479 Cocco, S., Corti, G., Agnelli, A., 2017. Influence of altitude on biochemical properties of
480 European Beech (*Fagus sylvatica* L.) forest soils. *Forests* 8, 1–14.
481 <https://doi.org/10.3390/f8060213>

482 de Paul Obade, V., Lal, R., 2016. A standardized soil quality index for diverse field conditions. *Sci.*
483 *Total Environ.* 541, 424–434. <https://doi.org/10.1016/j.scitotenv.2015.09.096>

484 Demchik, M.C., Sharpe, W.E., 2000. The effect of soil nutrition, soil acidity and drought on
485 northern red oak (*Quercus rubra* L.) growth and nutrition on Pennsylvania sites with high and
486 low red oak mortality. *For. Ecol. Manage.* 136, 199–207. [https://doi.org/10.1016/S0378-1127\(99\)00307-2](https://doi.org/10.1016/S0378-1127(99)00307-2)

488 Dick, R.P., Pankhurst, C., Doube, B.M., Gupta, V., 1997. Soil enzyme activities as integrative
489 indicators of soil health. *Biol. Indic. soil Heal.* 121–156.

490 Drenovsky, R.E., Vo, D., Graham, K.J., Scow, K.M., 2004. Soil water content and organic carbon
491 availability are major determinants of soil microbial community composition. *Microb. Ecol.*

492 48, 424–430. <https://doi.org/10.1007/s00248-003-1063-2>

493 Falsone, G., Bonifacio, E., Zanini, E., 2012b Structure development in aggregates of poorly
494 developed soils through the analysis of the pore system. *Catena* 95, 169–176.
495 <https://doi.org/10.1016/j.catena.2012.02.014>

496 Falsone, G., Marinari, S., Vittori Antisari, L., Vianello, G., 2015. Soil processes related to organic
497 matter modifications following Douglas-fir mature reforestation. *Biol. Fertil. Soils* 51, 277–
498 287. <https://doi.org/10.1007/s00374-014-0971-2>

499 FAO - Food and Agriculture Organization of the United Nations, 2017. Food and Agriculture
500 Organization of the United Nations - Statistic division. URL
501 <http://www.fao.org/faostat/en/#data/QC>

502 Ferracini, C., Gonella, E., Ferrari, E., Saladini, M.A., Picciau, L., Tota, F., Pontini, M., Alma, A.,
503 2015. Novel insight in the life cycle of *Torymus sinensis*, biocontrol agent of the chestnut gall
504 wasp. *BioControl* 60, 169–177. <https://doi.org/10.1007/s10526-014-9633-4>

505 Fonseca, T.F., Abreu, C.G., Parresol, B.R., 2004. Soil compaction and chestnut ink disease. *For.*
506 *Path.* 34, 273–283. <https://doi.org/10.1111/j.1439-0329.2004.00371.x>

507 Gee, G. W., Bauder, J. W., 1986. *Methods of Soil Analysis: Part 1—Physical and Mineralogical*
508 *Methods*. SSSA Book Series. Soil Science Society of America, American Society of
509 *Agronomy*. <https://doi.org/10.2136/sssabookser5.1.2ed.c15>

510 Ghobadi, M.H., Mousavi, S., 2014. The effect of pH and salty solutions on durability of sandstones
511 of the Aghajari Formation in Khuzestan province, southwest of Iran. *Arab. J. Geosci.* 7, 641–
512 653. <https://doi.org/10.1007/s12517-012-0741-0>

513 Giménez, R., Govers, G., 2008. Effects of freshly incorporated straw residue on rill erosion and
514 hydraulics. *Catena* 72, 214–223. <https://doi.org/10.1016/j.catena.2007.05.004>

515 Haynes, R.J., Swift, R.S., 1986. Effects of soil acidification and subsequent leaching on levels of
516 extractable nutrients in a soil. *Plant Soil* 95, 327–336. <https://doi.org/10.1007/BF02374613>

517 Hoffmann, U., Hoffmann, T., Johnson, E.A., Kuhn, N.J., 2014. Assessment of variability and

518 uncertainty of soil organic carbon in a mountainous boreal forest (Canadian Rocky Mountains,
519 Alberta). *Catena* 113, 107–121. <https://doi.org/10.1016/j.catena.2013.09.009>

520 Homann, P.S., Bormann, B.T., Boyle, J.R., 2001. Detecting treatment differences in soil carbon and
521 nitrogen resulting from forest manipulations. *Soil Sci. Soc. Am. J.* 65, 463–469.
522 <https://doi.org/10.2136/sssaj2001.652463x>

523 ISPRA, 1999. CARG Project - Geological and geothematic cartography. URL
524 [www.isprambiente.gov.it/en/projects/soil-and-territory/carg-project-geologic-and-geothematic-](http://www.isprambiente.gov.it/en/projects/soil-and-territory/carg-project-geologic-and-geothematic-cartography)
525 [cartography](http://www.isprambiente.gov.it/en/projects/soil-and-territory/carg-project-geologic-and-geothematic-cartography)

526 James, J., Harrison, R., 2016. The effect of harvest on forest soil carbon: A meta-analysis. *Forests* 7.
527 <https://doi.org/10.3390/f7120308>

528 Jankauskas, B., Fullen, M.A., 2002. A pedological investigation of soil erosion severity on
529 undulating land in Lithuania. *Can. J. Soil Sci.* 82, 311–321. <https://doi.org/10.4141/S01-058>

530 Jenkinson, D.S., Poulton, P.R., Bryant, C., 2008. The turnover of organic carbon in subsoils. Part 1.
531 Natural and bomb radiocarbon in soil profiles from the Rothamsted long-term field
532 experiments. *Eur. J. Soil Sci.* 59, 391–399. <https://doi.org/10.1111/j.1365-2389.2008.01025.x>

533 Jia, J., Cao, Z., Liu, C., Zhang, Z., Lin, L., Wang, Y., Haghypour, N., Wacker, L., Bao, H., Dittmar,
534 T., Simpson, M.J., Yang, H., Crowther, T.W., Eglinton, T.I., He, J.S., Feng, X., 2019. Climate
535 warming alters subsoil but not topsoil carbon dynamics in alpine grassland. *Glob. Chang. Biol.*
536 25, 4383–4393. <https://doi.org/10.1111/gcb.14823>

537 Johnson, A.H., Friedland, A.J., Miller, E.K., Siccama, T.G., 1994. Acid rain and soils of the
538 Adirondacks. III. Rates of soil acidification in a montane spruce-fir forest at Whiteface
539 Mountain, New York. *Can. J. For. Res.* 24, 663–669. <https://doi.org/10.1139/x94-089>

540 Johnson, D.W., Curtis, P.S., 2001. Effects of forest management on soil C and N storage: Meta
541 analysis. *For. Ecol. Manage.* 140, 227–238. [https://doi.org/10.1016/S0378-1127\(00\)00282-6](https://doi.org/10.1016/S0378-1127(00)00282-6)

542 Kaiser, K., Kalbitz, K., 2012. Cycling downwards - dissolved organic matter in soils. *Soil Biol.*
543 *Biochem.* 52, 29–32. <https://doi.org/10.1016/j.soilbio.2012.04.002>

544 Křeček, J., Novakova, J., Palan, L., Pazourkova, E., 2019. Soil conservation in a forested mountain
545 catchment. *Int. J. Environ. Qual.* 33, 27–36.

546 Kuzyakov, Y., 2010. Priming effects: Interactions between living and dead organic matter. *Soil*
547 *Biol. Biochem.* 42, 1363–1371. <https://doi.org/10.1016/j.soilbio.2010.04.003>

548 Leuschner, C., Wulf, M., Bäuchler, P., Hertel, D., 2014. Forest continuity as a key determinant of
549 soil carbon and nutrient storage in beech forests on sandy soils in Northern Germany.
550 *Ecosystems* 17, 497–511. <https://doi.org/10.1007/s10021-013-9738-0>

551 Likens, G.E., Driscoll, C.T., Buso, D.C., Mitchell, M.J., Lovett, G.M., Bailey, S.W., Siccama, T.G.,
552 Reiners, W.A., Alewell, C., 2002. The biogeochemistry of sulfur at Hubbard Brook.
553 *Biogeochemistry* 60, 235–316. <https://doi.org/10.1023/A:1020972100496>

554 Liu, W., Luo, Q., Lu, H., Wu, J., Duan, W., 2017. The effect of litter layer on controlling surface
555 runoff and erosion in rubber plantations on tropical mountain slopes, SW China. *Catena* 149,
556 167–175. <https://doi.org/10.1016/j.catena.2016.09.013>

557 Maes, S.L., Blondeel, H., Perring, M.P., Depauw, L., Brūmelis, G., Brunet, J., Decocq, G., den
558 Ouden, J., Härdtle, W., Hédl, R., Heinken, T., Heinrichs, S., Jaroszewicz, B., Kirby, K.,
559 Kopecký, M., Máliš, F., Wulf, M., Verheyen, K., 2019. Litter quality, land-use history, and
560 nitrogen deposition effects on topsoil conditions across European temperate deciduous forests.
561 *For. Ecol. Manage.* 433, 405–418. <https://doi.org/10.1016/j.foreco.2018.10.056>

562 Mandal, K.G., Misra, A.K., Hati, K.M., Bandyopadhyay, K.K., Ghosh, P.K., 2004. Rice residue-
563 management options and effects on soil properties and crop productivity 2, 224–231.

564 Martín, M.A., Mattioni, C., Cherubini, M., Villani, F., Martín, L.M., 2017. A comparative study of
565 European chestnut varieties in relation to adaptive markers. *Agrofor. Syst.* 91, 97–109.
566 <https://doi.org/10.1007/s10457-016-9911-5>

567 McDonnell, T.C., Sullivan, T.J., Cosby, B.J., Jackson, W.A., Elliott, K.J., 2013. Effects of climate,
568 land management, and sulfur deposition on soil base cation supply in national forests of the
569 southern Appalachian Mountains. *Water. Air. Soil Pollut.* 224. <https://doi.org/10.1007/s11270->

570 013-1733-8

571 Mitchell, E., Scheer, C., Rowlings, D., Conant, R.T., Cotrufo, M.F., Grace, P., 2018. Amount and
572 incorporation of plant residue inputs modify residue stabilisation dynamics in soil organic
573 matter fractions. *Agric. Ecosyst. Environ.* 256, 82–91.
574 <https://doi.org/10.1016/j.agee.2017.12.006>

575 Mobley, M.L., Lajtha, K., Kramer, M.G., Bacon, A.R., Heine, P.R., Richter, D.D., 2015. Surficial
576 gains and subsoil losses of soil carbon and nitrogen during secondary forest development.
577 *Glob. Chang. Biol.* 21, 986–996. <https://doi.org/10.1111/gcb.12715>

578 Mokma, D.L., Fenton, T.E., Olson, K.R., 1996. Effect of erosion on morphology and classification
579 of soils in the North Central United States. *J. Soil Water Conserv.* 51, 171–175.

580 Nave, L.E., Vance, E.D., Swanston, C.W., Curtis, P.S., 2010. Harvest impacts on soil carbon
581 storage in temperate forests. *For. Ecol. Manage.* 259, 857–866.
582 <https://doi.org/10.1016/j.foreco.2009.12.009>

583 Neilsen, G., Forge, T., Angers, D., Neilsen, D., Hogue, E., 2014. Suitable orchard floor
584 management strategies in organic apple orchards that augment soil organic matter and maintain
585 tree performance. *Plant Soil* 378, 325–335. <https://doi.org/10.1007/s11104-014-2034-8>

586 Oosterbaan, A., 1998. Growth of chestnut (*Castanea sativa*) in the Netherlands. *Forestry* 71, 267–
587 270. <https://doi.org/10.1093/forestry/71.3.267>

588 Orgill, S.E., Condon, J.R., Kirkby, C.A., Orchard, B.A., Conyers, M.K., Greene, R.S.B., Murphy,
589 B.W., 2017. Soil with high organic carbon concentration continues to sequester carbon with
590 increasing carbon inputs. *Geoderma* 285, 151–163.
591 <https://doi.org/10.1016/j.geoderma.2016.09.033>

592 Orsini, L., Rémy, J., 1976. Utilisation du chlorure de cobaltihexamine pour la détermination
593 simultanée de la capacité d'échange et des bases échangeables des sols. *Bulletin de*
594 *l'Association Française d'Etude du Sol.* 4 : 269–279. *Bull. l'Association Française d'Etude du*
595 *Sol.* 4 269–279. 4, 269–279.

596 Pezzi, G., Maresi, G., Conedera, M., Ferrari, C., 2011. Woody species composition of chestnut
597 stands in the Northern Apennines: The result of 200 years of changes in land use. *Landsc.*
598 *Ecol.* 26, 1463–1476. <https://doi.org/10.1007/s10980-011-9661-8>

599 Pieri, L., Rondini D., Ventura, F., 2017. Changes in the rainfall–streamflow regimes related to
600 climate change in a small catchment in Northern Italy. *Theor Appl Climatol* 129, 1075–1087.
601 <https://doi.org/10.1007/s00704-016-1834-6>

602 R Core Team, 2016. R: A language and environment for statistical computing.

603 Rees, R.M., Parker, J.P., 2005. Filtration increases the correlation between water extractable
604 organic carbon and soil microbial activity. *Soil Biol. Biochem.* 37, 2240–2248.
605 <https://doi.org/10.1016/j.soilbio.2005.03.024>

606 Regione Emilia Romagna, 2000. Cartografia dei suoli della regione Emilia Romagna 1:50,000
607 (1994 updates 2000). https://geo.regione.emilia-romagna.it/cartografia_sgss/user/viewer.jsp?service=pedologia&bookmark=1%22

608

609 Ren, C., Zhang, W., Zhong, Z.K., Han, X., Yang, G., Feng, Y., Ren, G., 2018. Differential
610 responses of soil microbial biomass, diversity, and compositions to altitudinal gradients
611 depend on plant and soil characteristics. *Sci. Total Environ.* 610–611, 750–758.
612 <https://doi.org/10.1016/j.scitotenv.2017.08.110>

613 Ribeiro, S.L., Fonseca, T.F., Pires, A.L., 2019. Influence of fertilization on growth of young
614 chestnut trees (*Castanea sativa* Mill.) managed for wood production. *Cerne* 25, 357–364.
615 <https://doi.org/10.1590/01047760201925042660>

616 Soil Survey Staff, 2014. *Keys to Soil Taxonomy*, 12th ed. USDA–Natural Resources Conservation
617 Service, Washington.

618 Stefanowicz, A.M., Stanek, M., Nobis, M., Zubek, S., 2016. Species-specific effects of plant
619 invasions on activity, biomass, and composition of soil microbial communities. *Biol. Fertil.*
620 *Soils* 52, 841–852. <https://doi.org/10.1007/s00374-016-1122-8>

621 Steinmann, T., Welp, G., Holbeck, B., Amelung, W., 2016. Long-term development of organic

622 carbon contents in arable soil of North Rhine–Westphalia, Germany, 1979–2015. *Eur. J. Soil*
623 *Sci.* 67, 616–623. <https://doi.org/10.1111/ejss.12376>

624 Świtoniak, M., 2014. Use of soil profile truncation to estimate influence of accelerated erosion on
625 soil cover transformation in young morainic landscapes, North-Eastern Poland. *Catena* 116,
626 173–184. <https://doi.org/10.1016/j.catena.2013.12.015>

627 Tian, D.L., Peng, Y.Y., Yan, W. De, Fang, X., Kang, W.X., Wang, G.J., Chen, X.Y., 2010. Effects
628 of thinning and litter fall removal on fine root production and soil organic carbon content in
629 Masson Pine plantations. *Pedosphere* 20, 486–493. [https://doi.org/10.1016/S1002-](https://doi.org/10.1016/S1002-0160(10)60038-0)
630 [0160\(10\)60038-0](https://doi.org/10.1016/S1002-0160(10)60038-0)

631 Trumbore, S., 2009. Radiocarbon and Soil Carbon Dynamics. *Annu. Rev. Earth Planet. Sci.* 37, 47–
632 66. <https://doi.org/10.1146/annurev.earth.36.031207.124300>

633 Turner, B.L., Wells, A., Condon, L.M., 2014. Soil organic phosphorus transformations along a
634 coastal dune chronosequence under New Zealand temperate rain forest. *Biogeochemistry* 121,
635 595–611. <https://doi.org/10.1007/s10533-014-0025-8>

636 Van Der Heijden, M.G.A., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority: Soil
637 microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.* 11
638 (3), 296 - 310. <https://doi.org/10.1111/j.1461-0248.2008.01199.x>

639 van Leeuwen, J.P., Djukic, I., Bloem, J., Lehtinen, T., Hemerik, L., de Ruiter, P.C., Lair, G.J., 2017.
640 Effects of land use on soil microbial biomass, activity and community structure at different soil
641 depths in the Danube floodplain. *Eur. J. Soil Biol.* 79, 14–20.
642 <https://doi.org/10.1016/j.ejsobi.2017.02.001>

643 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil
644 microbial biomass C. *Soil Biol. Biochem.* 19, 703–707. [https://doi.org/10.1016/0038-](https://doi.org/10.1016/0038-0717(87)90052-6)
645 [0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)

646 Vannier, C., Guillet, B., 1994. Sulphur forms in the organic fractions of an upland forest soil (Mont
647 Lozère, France). *Soil Biol. Biochem.* 26, 149–151. <https://doi.org/10.1016/0038->

648 0717(94)90207-0

649 Vittori Antisari, L., Bianchini, G., Dinelli, E., Falsone, G., Gardini, A., Simoni, A., Tassinari, R.,
650 Vianello, G., 2014. Critical evaluation of an intercalibration project focused on the definition
651 of new multi-element soil reference materials (AMS-MO1 and AMS-ML1). *EQA – Int. J.*
652 *Environ. Qual.* 15, 41–64. <https://doi.org/10.6092/issn.2281-4485/4553>

653 Vittori Antisari, L., Falsone, G., Carbone, S., Vianello, G., 2013. Short-term effects of forest
654 recovery on soil carbon and nutrient availability in an experimental chestnut stand. *Biol. Fertil.*
655 *Soils* 49, 165–173. <https://doi.org/10.1007/s00374-012-0708-z>

656 Vittori Antisari, L., Laudicina, V.A., Falsone, G., Carbone, S., Badalucco, L., Vianello, G., 2016.
657 Native and planted forest species determine different carbon and nitrogen pools in Arenosol
658 developed on Holocene deposits from a coastal Mediterranean area (Tuscany, Italy). *Environ.*
659 *Earth Sci.* 75. <https://doi.org/10.1007/s12665-016-5581-x>

660 Waldboth, W., Oberhuber, W., 2009. Synergistic effect of drought and chestnut blight
661 (*Cryphonectria parasitica*) on growth decline of European chestnut (*Castanea sativa*). *For.*
662 *Path.* 39, 43–55. <https://doi.org/10.1111/j.1439-0329.2008.00562.x>

663 Xiong, Y., Xia, H., Li, Z., Cai, X., Fu, S., 2008. Impacts of litter and understory removal on soil
664 properties in a subtropical *Acacia mangium* plantation in China. *Plant Soil* 304, 179–188.
665 <https://doi.org/10.1007/s11104-007-9536-6>

666 Xu, X., Han, L., Wang, Y., Inubushi, K., 2007. Influence of vegetation types and soil properties on
667 microbial biomass carbon and metabolic quotients in temperate volcanic and tropical forest
668 soils. *Soil Sci. Plant Nutr.* 53, 430–440. <https://doi.org/10.1111/j.1747-0765.2007.00146.x>

669 Yavitt, J.B., Fahey, T.J., Sherman, R.E., Groffman, P.M., 2015. Lumbricid earthworm effects on
670 incorporation of root and leaf litter into aggregates in a forest soil, New York State.
671 *Biogeochemistry* 125, 261–273. <https://doi.org/10.1007/s10533-015-0126-z>

672 Zetterberg, T., Olsson, B.A., Löfgren, S., Hyvönen, R., Brandtberg, P.O., 2016. Long-term soil
673 calcium depletion after conventional and whole-tree harvest. *For. Ecol. Manage.* 369, 102–

674 115. <https://doi.org/10.1016/j.foreco.2016.03.027>

675 Zlatanov, T., Schleppi, P., Velichkov, I., Hinkov, G., Georgieva, M., Eggertsson, O., Zlatanova, M.,
676 Vacik, H., 2013. Structural diversity of abandoned chestnut (*Castanea sativa* Mill.) dominated
677 forests: Implications for forest management. *For. Ecol. Manag.* 291, 326–335.
678 <https://doi.org/10.1016/j.foreco.2012.11.015>

679 Zornoza, R., Acosta, J.A., Bastida, F., Domínguez, S.G., Toledo, D.M., Faz, A., 2015. Identification
680 of sensitive indicators to assess the interrelationship between soil quality, management
681 practices and human health. *Soil* 1, 173–185. <https://doi.org/10.5194/soil-1-173-2015>

682

Figure 1




Figure 2




Figure 3







Figure 4







Figure 5




Figure captions

Figure 1. The localization of investigated sites (REM1, REM2, CONS1 and CONS2) (on left side) and a recovery chestnut stand where it is visible the 10 x 10 m tree spacing (on right side). REM and CONS study sites are chestnut (*Castanea sativa* Mill.) stands with removal and conservation of organic residues, respectively.

Figure 2. Cation exchange capacity (I), exchangeable calcium (II), magnesium (III), potassium (IV) and sodium (V), and base saturation (VI) in epipedon (Epi) and endopedon (Endo) of soils under chestnut (*Castanea sativa* Mill.) stands with removal (REM; white bars) and conservation (CONS; grey bars) of organic residues. Error bars are the standard errors. According to Kruskal–Wallis test, within each soil layer, different lowercase letters indicate significant differences ($p < 0.05$) between the two managements (REM vs CONS), instead, within each management different uppercase letters indicate significant differences ($p < 0.05$) between epipedon and endopedon (Epi vs Endo).

Figure 3. Amounts of total organic carbon (I), total nitrogen (II), C associated to water–extractable organic matter (III) and N associated to water–extractable organic matter (IV) in epipedon (Epi) and endopedon (Endo) of soils under chestnut (*Castanea sativa* Mill.) stands with removal (REM; white bars) and conservation (CONS; grey bars) of organic residues. Error bars are the standard errors. According to Kruskal–Wallis test, within each soil layer, different lowercase letters indicate significant differences ($p < 0.05$) between the two managements (REM vs CONS), instead, within each management different uppercase letters indicate significant differences ($p < 0.05$) between epipedon and endopedon (Epi vs Endo).

Figure 4. Amounts of C associated to particulate organic matter (I), and to fulvic–like (II), humic–like (III) and humin–like (IV) substances in epipedon (Epi) and endopedon (Endo) of soils under chestnut (*Castanea sativa* Mill.) stands with removal (REM; white bars) and conservation (CONS;

grey bars) of the organic residues. Error bars are the standard errors. According to Kruskal–Wallis test, within each soil layer, different lowercase letters indicate significant differences ($p<0.05$) between the two managements (REM vs CONS, instead, within each management different uppercase letters indicate significant differences ($p<0.05$) between epipedon and endopedon (Epi vs Endo).

Figure 5. Content of microbial biomass C (I) and N (II), basal respiration (III), total amount of CO₂ evolved during basal respiration experiments (IV), metabolic (V) and mineralization (VI) quotients in epipedon (Epi) and endopedon (Endo) of soils under chestnut (*Castanea sativa* Mill.) stands with removal (REM; white bars) and conservation (CONS; grey bars) of the organic residues. Error bars are the standard errors. According to Kruskal–Wallis test, within each soil layer, different lowercase letters indicate significant differences ($p<0.05$) between the two managements (REM vs CONS), instead, within each management different uppercase letters indicate significant differences ($p<0.05$) between epipedon and endopedon (Epi vs Endo).

Table 1. The study sites locations and climatic characteristics

Study site	Coordinates (UTM 32 T)		MAAT	MAP
	m E	m N	° C	mm
REM 1	686305	4904858	11.7	1136
REM 2	686751	4904078	11.7	1136
CONS 1	690032	4898682	11.0	1251
CONS 2	618314	4927009	10.9	883

MAAT = mean annual air temperature; MAP = mean annual precipitations.

Table 2[Click here to download Table: Table 2.docx](#)

Table 2. Thickness, amounts of sand, silt and clay, and pH in water ($\text{pH}_{\text{H}_2\text{O}}$) and in KCl (pH_{KCl}) values measured in epipedon (Epi) and endopedon (Endo) of soils under chestnut (*Castanea sativa* Mill.) stands with removal (REM) and conservation (CONS) of the organic residues. Standard error is reported in parentheses. According to Kruskal–Wallis test: *i*) within each management, the uppercase letters indicate significant differences ($p < 0.05$) between soil layers (Epi vs Endo); *ii*) within each soil layer, the lowercase letters indicate significant differences ($p < 0.05$) between the two managements (REM vs CONS).

Layer	Management	Thickness cm	Sand	Silt g kg^{-1}	Clay	$\text{pH}_{\text{H}_2\text{O}}$	pH_{KCl}
Epi	REM	5.1 Bb (0.7)	564 (67)	382 (51)	152 a (21)	4.9 (0.2)	3.5 (0.2)
Epi	CONS	9.7 Ba (1.8)	522 (20)	370 B (15)	89 b (11)	4.8 (0.1)	3.2 B (0.2)
Endo	REM	12.4 Ab (1.8)	545 (58)	322 (46)	132 a (16)	4.9 (0.1)	3.4 (0.1)
Endo	CONS	20.2 Aa (2.1)	467 (13)	461 A (13)	71 b (3)	5.4 (0.1)	4.1 A (0.1)

Table 3[Click here to download Table: Table 3.docx](#)

Table 3. Stock of organic carbon (OC), N, Ca, Mg, Na, K, Fe, P and S measured at 0 – 10, 10 – 20, 20 – 30 cm soil depth intervals under chestnut (*Castanea sativa* Mill.) stands with organic residues removal (REM) and conservation (CONS). Standard error is reported in parentheses. According to Kruskal–Wallis test: *i*) within each management, different uppercase letters indicate significant differences among the soil depth intervals ($p < 0.05$); *ii*) within each soil depth interval, different lowercase letters indicate significant differences ($p < 0.05$) between the two managements (REM vs CONS).

Depth	Management	OC	N	Ca	Mg	Na	K	Fe	P	S
		kg ha ⁻¹								
0 – 10	REM	37183 A	2702 A	1319	3834	313 B	3716	16056	223	207 A
		(8159)	(598)	(255)	(890)	(17)	(642)	(3308)	(36)	(53)
0 – 10	CONS	30734 A	2338 A	2463	4357 B	346	4845	19435 B	295	186
		(3154)	(285)	(349)	(277)	(22)	(304)	(959)	(18)	(17)
10 – 20	REM	9581 Bb	739 Bb	1224	5002	407 A	4720	20897	159 b	84 Bb
		(2202)	(149)	(189)	(1044)	(32)	(801)	(3883)	(21)	(10)
10 – 20	CONS	17242 Ba	1389 Ba	2446	5537 A	393	5552	25697 A	280 a	138 a
		(2390)	(159)	(458)	(215)	(32)	(351)	(1448)	(27)	(12)
20 – 30	REM	6875 Bb	639 Bb	1441	6009	426 A	5431	24214	150 b	72 Bb
		(1354)	(134)	(207)	(1439)	(46)	(1099)	(5288)	(21)	(10)
20 – 30	CONS	13668 Ba	1159 Ba	2393	6027 A	405	5970	27837 A	269 a	118 a
		(1113)	(71)	(448)	(358)	(35)	(421)	(1562)	(14)	(8)

Table 4[Click here to download Table: Table 4.docx](#)

Table 4. Mean \pm standard error of C, N, Ca, Mg, Na, K, Fe, P and S contents in chestnut (*Castanea sativa* Mill.) leaves collected from chestnut stands with organic residues removal (REM) and conservation (CONS).

	REM	CONS
C (g kg ⁻¹)	411 \pm 11	416 \pm 5
N (g kg ⁻¹)	21.0 \pm 1.0	23.7 \pm 1.7
Ca (g kg ⁻¹)	4.31 \pm 1.12	5.22 \pm 0.78
Mg (g kg ⁻¹)	1.94 \pm 0.09	1.73 \pm 0.08
Na (g kg ⁻¹)	0.10 \pm 0.02	0.19 \pm 0.01
K (g kg ⁻¹)	5.63 \pm 0.19	7.03 \pm 0.55
Fe (g kg ⁻¹)	0.10 \pm 0.00	0.13 \pm 0.02
P (g kg ⁻¹)	1.34 \pm 0.17	1.05 \pm 0.07
S (g kg ⁻¹)	0.99 \pm 0.07	1.48 \pm 0.22

Supplementary material for on-line publication only

[Click here to download Supplementary material for on-line publication only: Supplementary materials_new.docx](#)

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: