

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia / Dean, Silas; Pappalardo, Marta; Boschian, Giovanni; Spada, Giorgio; Forenbaher, Stašo; Juračić, Mladen; Felja, Igor; Radić, Dinko; Miracle, Preston T.. - In: QUATERNARY SCIENCE REVIEWS. - ISSN 0277-3791. - STAMPA. - 244:(2020), pp. 106503.1-106503.18. [10.1016/j.quascirev.2020.106503]

Availability:

This version is available at: https://hdl.handle.net/11585/771653 since: 2022-02-14

Published:

DOI: http://doi.org/10.1016/j.quascirev.2020.106503

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Silas Dean, Marta Pappalardo, Giovanni Boschian, Giorgio Spada, Stašo Forenbaher, Mladen Juračić, Igor Felja, Dinko Radić, Preston T. Miracle, *Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia*, Quaternary Science Reviews, Volume 244, 2020, 106503.

Thefinalpublishedversionisavailableonlineat:https://doi.org/10.1016/j.quascirev.2020.106503

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

1	Human adaptation to changing coastal landscapes in the Eastern
2	Adriatic: Evidence from Vela Spila cave, Croatia
3	Silas Dean <sup>1</sup> , Marta Pappalardo <sup>1</sup> *, Giovanni Boschian <sup>2</sup> , Giorgio Spada <sup>3</sup> , Stašo Forenbaher <sup>4</sup> , Mladen Juračić <sup>5</sup> ,
4	Igor Felja <sup>6</sup> , Dinko Radić <sup>7</sup> Preston T. Miracle <sup>8</sup>
5	
6	<sup>1</sup> Department of Earth Sciences, University of Pisa, Italy
7	<sup>2</sup> Department of Biology, University of Pisa, Italy
8	<sup>3</sup> Dipartimento di Scienze Pure e Applicate, Università di Urbino 'Carlo Bo', Urbino, Italy
9	<sup>4</sup> Institute for Anthropological Research, Zagreb, Croatia
10	<sup>5</sup> Croatian Academy of Sciences and Arts, Croatia
11	<sup>6</sup> University of Zagreb, Faculty of Science, Dept. of Geology, Croatia
12	<sup>7</sup> Centar za kulturu, Vela Luka, Croatia
13	<sup>8</sup> Department of Archaeology, University of Cambridge, UK
14	*Corresponding author
15	
16	KEYWORDS: Sea-level change; Environmental change; Human settlement patterns, Bathymetry;

17 Prehistoric archaeology; Late Pleistocene-Holocene; Central Adriatic coast; Croatia

# 18 Abstract

- 19 In this paper new palaeogeographic and archaeological data from the prehistoric cave Vela Spila on
- 20 the island of Korčula in Croatia are combined with new realizations of two glacial isostatic
- 21 adjustment models in order to present relative sea-level change scenarios confronting the
- 22 inhabitants of the cave at different time slices and to show how they experienced and adapted to

≛

1

23 sea-level and climate change from the Late Pleistocene through the Holocene. Our results show that 24 from the Late Upper Palaeolithic until the Mesolithic, humans in the study area would have 25 experienced tens of metres of sea-level rise, at rates in some cases up to 12 mm per year, and, owing 26 to the relatively flat morphology of the now submerged plains, hundreds of meters of horizontal 27 coastline change in the plains to the north and south of the island. This evidence supports the 28 hypothesis that the rapid loss of these plains likely contributed to the human abandonment of the 29 cave after the Palaeolithic for about five thousand years, followed by significant changes in lifestyle 30 and diet in the Mesolithic. Our results have important implications for the study of how past human 31 groups, especially in vulnerable coastal areas, were affected by sea level, climate, and other 32 environmental changes. Vela Spila represents a case study of how changing environment and rising 33 seas can force significant alterations in human societies, even when there is no risk of inundation to 34 the settlement itself.

## 35 **1** Introduction

36 Since prehistory, humans have adapted their settlement and mobility patterns in response to 37 environmental change (Blute, 2008; Bini et al., 2018; Butzer, 2005), using their decision-making 38 skills (Nunn, 2020), or even engineering skills (Galili et al., 2019) to overcome challenges created 39 by modifications to coastlines and their related ecosystems. The Eastern Adriatic region represents a 40 key area to investigate the relationship between environmental changes and cultural responses. The 41 'Great Adriatic Plain' periodically exposed during the Pleistocene provided suitable conditions for 42 human settlement; its loss to rising sea level during the Late Pleistocene and Early Holocene 43 precipitated a significant restructuring of plant, animal, and human communities (Boschian and 44 Fusco, 2007; Miracle, 2007). The palaeoecological and archaeological evidence of the region is 45 reviewed by Pilaar Birch and Linden (2018).

46 Sea-level change has been a major cause of environmental change; it has modified coastal 47 landscapes by dramatically and cyclically altering coastal plains around shallow sea basins, and by 48 influencing the distribution of islands and their distance from the mainland. The outline and area of 49 many present-day islands were different during the Late Pleistocene owing to the last ice age and 50 the relative sea-level (RSL) changes associated with it; the islands' current isolation is unusual 51 during the last 70 ka (Weigelt et al., 2016). Consequently, the distribution of human and animal 52 communities in coastal areas (Carroll et al., 2012; Garg et al., 2018; Hewitt, 2000; McLaughlin et 53 al., 2018) was profoundly affected, resulting in major behavioural and cultural change. A robust 54 reconstruction of shoreline position at different time slices of the Late Pleistocene and Holocene is 55 essential for understanding human settlement and animal dispersal patterns. 56 Following the seminal work by van Andel (1989), reconstructing ancient coastlines in order to 57 elucidate their influence on hominin dispersal and cultural behaviour became an important research 58 goal. In archaeological literature, the reconstruction of ancient coastal landscapes has focused on 59 comparing present-day bathymetric surfaces with global eustatic curves (Dobson, 2014). However,

60 past RSL can be reconstructed more precisely by combining geophysical and geological techniques

61 (Pirazzoli, 2013; De Boer et al., 2014; Shennan et al., 2015).

62 Sea-level reconstructions must be combined with bathymetric data, which have also been widely 63 used to identify landforms, assess palaeoenvironments and palaeoshorelines, and address 64 archaeological questions in the central Mediterranean. Bathymetric mapping and archaeological 65 research have been combined in Sardinia (Palombo et al., 2017) for this purpose. In the Tyrrhenian 66 Sea west of Italy, seismic profiles have been used to detect terraces that may be associated with 67 ancient sea stands (Casalbore et al., 2017). Extensive single- and multibeam data were used 68 (Trincardi et al., 2014) to identify several drowned landforms in the western Adriatic. Bathymetric 69 data from Malta (Foglini et al., 2016) have shown drowned karstic features and river landforms, and 70 can answer questions about prehistoric fauna migration and human settlement. In the NW Adriatic,

71 prehistoric settlement surfaces and lagoon deposits have contributed to landform and RSL 72 reconstructions (Fontana et al., 2017). In the eastern coast of the Adriatic dozens of submerged 73 coastal prehistoric sites have been located, from Istria to southern Dalmatia, and excavations have 74 been initiated on several of those sites (Bekić, 2017; Bekić et al., 2015; Benjamin et al., 2011; 75 Benjamin and Bonsall, 2009; Čelhar et al., 2017; Koncani Uhač and Čuka, 2015; Radić Rossi, 76 2008). The majority of the sites belong to relatively recent prehistoric times (Bronze Age and Iron 77 Age), while only very few of them belong to Copper Age or Neolithic periods. Further excavation, 78 interdisciplinary investigation (collaboration of archaeologists and geologists), and analyses are 79 required before they can be used as accurate indicators of local sea-level change in prehistoric times 80 (Rossi, 2017)

81 This study integrates palaeoenvironmental and archaeological evidence from Vela Spila, a cave 82 situated on the island of Korčula, close to the central Dalmatian coast in Croatia (Fig 1) with a new 83 synthesis of bathymetric datasets. We Palaeogeographic reconstructions were created at time slices 84 relevant for key cultural events during cave occupation phases from the Late Upper Palaeolithic to 85 the Neolithic. The main goal of this work is to investigate the relationship between the shoreline 86 position and the cave. The results provide an understanding of how inhabitants of the cave experienced and adapted to sea-level and climate changes since the Late Pleistocene. 87 88 This study incorporates both existing and new data and approaches that are discussed in light of 89 current research. In particular:

1) Novel archaeological evidence provided in this study compliments previous work on Vela Spila
(Branscombe et al., 2020; Cristiani et al., 2014; Farbstein et al., 2012; Radić, 2018; Rainsford et al.,
2014). Vela Spila (Fig 2) is one of the richest and most promising archaeological sites in the Eastern
Adriatic region, with a Late Pleistocene to Holocene stratigraphic sequence spanning the Late
Upper Palaeolithic (LUP) to the Bronze Age. The deposits include exceptionally well-preserved
cultural, human, animal, and vegetal remains that document important environmental and biological

96 changes during the Pleistocene-Holocene, and cultural changes during the Palaeolithic-Mesolithic97 and Mesolithic-Neolithic transitions.

98 2) Data used in the current study to reconstruct bathymetry around Vela Spila represent a synthesis 99 of available bathymetric datasets for the area, and thus an improvement on the datasets used by 100 previous authors such as Sikora et al. (2014), which used satellite bathymetric data combined with 101 GIS analysis to reconstruct river drainage during the lowstand of the Last Glacial Maximum 102 (LGM). Bathymetric data, as a source of palaeolandscape data, are applicable to our study area 103 because of the karstic nature of the Eastern Adriatic - rainwater sinks into the endokarst instead of 104 creating high sediment flux onto the seabed (Pikelj and Juračić, 2013). Consequently, LGM 105 landforms and topography are still visible in bathymetric data, aside from limited fluvial sedimentation. 106

107 3) Bathymetric data for the seabed around Korčula are combined with two updated geophysical 108 models of glacial-hydro-isostatic adjustment (GIA) realized specifically for the area under 109 investigation in order to reconstruct palaeogeographic phases; all previous authors used in the best 110 case older, less refined models, and Sikora (2014) used a global model. This implies differences of 111 tens of kilometres in the position of the shoreline at specific times. The use of GIA models is valid 112 for this subregion of the Eastern Adriatic because tidal notches documented by Faivre and Butorac 113 (2018) show relatively slower rates of subsidence here than in areas to the north, at least for the 114 Late Holocene.

115 Vela Spila represents a relevant case study on how changing environment and rising seas can force 116 significant changes in human societies, even when there is no direct risk of inundation for the 117 settlement itself.

5

# **118 2 Regional setting**

## 119 2.1 Adriatic sea

120 The Adriatic Sea (Fig 1) is a NW-SE elongated basin of approximately 800 by 200 km that 121 separates the Italian peninsula from the Balkans (Trincardi et al., 2014). This basin can be divided 122 into three sub-basins: the northern, which is 35 m deep on average, the middle with an average 123 depth of 140 m, and the southern with maximum depths reaching 1200 m (Artegiani et al., 1997). 124 This paper is focused on the island of Korčula, which lies on the eastern coast between the middle 125 and southern basins. The Eastern Adriatic bedrock is about 8,000 m of limestone and/or dolomite 126 (Vlahović et al., 2005, 2002) formed in shallow sedimentation basins between the Mid-Late 127 Triassic, Mesozoic and Palaeogene, followed by relatively thin Eocene flysch outcrops. The 128 tectonic deformation pattern of the Eastern Adriatic is characterised by a complex system of faults, 129 thrust faults and overthrusted tiles formed since the Miocene and developed along the so-called Dinaric Strike with NW-SE orientation (Vlahović et al., 2005, 2002). The typical relief pattern of 130 131 mountain and hill ridges parallel to the coastline (Fig 1) derives from this tectonic compressional 132 motif, followed by extensional processes and eventually repeated flooding during the Quaternary 133 sea-level fluctuations. The westernmost Dinarides were submerged by the postglacial RSL rise 134 (Kelletat, 2005) and now form the Croatian islands and the typically rocky Dalmatian Coast (Felja 135 et al., 2015; Pikelj and Juračić, 2013). Significantly for this study area, the Eastern Adriatic coast 136 and its hinterland are karstic and dominated by typical karst landforms, with few significant rivers 137 discharging sediments into the sea, so that the present-day bathymetry closely reflects LGM 138 landforms (Pikelj and Juračić, 2013). This is confirmed in the immediate study area by research 139 (Giglio et al., 2020) in the waters around the present-day Neretva delta, which shows river 140 sedimentation rates quickly approaching 0 mm/a (mm per year) within 10 km offshore. This aspect of the coast contrasts with the wide and flat morphology of the north-eastern Adriatic, which was an
alluvial plain dominated by sandy river discharge when the LGM RSL was 120 m below present
(Trincardi et al., 2014). Though these submerged morphological features in the western Adriatic
have been identified by multibeam imagery (Trincardi et al., 2014), including drowned karst
landscapes (Taviani et al., 2012), public high-resolution data are less available for the rocky Eastern
Adriatic.

## 147 2.2 Central-southern Dalmatia

148 Korčula is an island in central-southern Dalmatia, separated from the Pelješac peninsula by a 149 narrow channel about 40-45 m deep. Like most Dalmatian islands (Kelletat, 2005), it represents the 150 top of a submerged ridge formed by tectonic overthrusting and uplift of reliefs which, in the case of 151 the southern islands, are rotated counterclockwise with reference to the dinaric strike. The island 152 landscape is rugged, controlled by the nature of the underlying Mesozoic limestones and dolomitic 153 limestones, and also characterised by several deep and elongated *polje* oriented along the structural 154 motif and by deep karstic depressions, whose bottoms are filled by terra rossa/crvenica soils 155 (Bonacci et al., 2012). The coasts are almost always steep and rocky. The channel between Korčula 156 and Hvar -the island immediately to the north – is at most about 70 m deep and represents the 157 prolongation of the Neretva river valley towards the Adriatic basin (Sikora et al., 2014). On the 158 westernmost tip of the island, a narrow, winding and shallow inlet protects the Vela Luka harbour. 159 In the now submerged zones between islands, areas at greater depth closer to the LGM lowstand of 160 -120 m (Peltier and Fairbanks, 2006) or -115 m (Sikora et al., 2014) had less exposure time to 161 erosion and other landscape forming processes than areas situated near highstand levels. As a result, 162 most morphology of these submerged areas consist of broad, flat areas; a fact often compounded by 163 the presence of alluvial sediments in areas near palaeo rivers.

#### 164 **2.3 Vela Spila**

165 Vela Spila (Fig. 2) is a large cave situated 121 m above sea level on the western side of Spilski Rat,

166 a low hill overlooking Vela Luka harbour. The cave's single, semi-ellipsoidal chamber, about 50 m

167 by 30 m wide and 17 m high, opens in Cenomanian radiolaritic limestone whose layers dip

168 southwards (Cristiani et al., 2014). The whole inside surface of about 1000  $m^2$  is at present always

169 well-lit through a wide entrance facing west and two large holes in its ceiling.

170 The outside landscape has been strongly affected by human activities over long time scales,

171 probably since the cave was first inhabited during the Late Glacial. Most hillsides have extensive

172 19<sup>th</sup> century terraces for viticulture or growing olive trees, though many are now abandoned. Good

173 quality data about outside soils and their evolution through the end of the Late Glacial to Middle

174 Holocene are difficult to collect in these conditions: the typical features are thick stone walls

175 sustaining terra rossa backfills, alternating with patches of barren denuded limestone. Cave data are

in this case the best way to acquire information about climate and environment evolution in the areasince they are undisturbed by agricultural activity during the last two millennia.

## 178 2.4 Regional relative sea level

179 Global eustatic sea level has risen roughly 120 m since the LGM (Fairbanks, 1989; Peltier and

180 Fairbanks, 2006). GIA models (Lambeck and Purcell, 2005; Peltier, 2004; Peltier et al., 2015) have

181 been used to estimate timing and magnitude of these changes. Datable biological, geological and

archaeological proxies have been used to calibrate these models and provide more precise

183 chronological indications of past RSL (Rovere et al., 2016).

184 In the Eastern Adriatic area, fossil remains of *Lithophyllum sp.* and an ancient harbour (Faivre et al.,

185 2013; Vacchi et al., 2016) provide sea-level indicators in close proximity to the study area. Salt

186 marsh foraminifera have also been studied (Shaw et al., 2018) to reconstruct RSL in the last

187 millenium, and suggest a subsidence rate of 0.45 +/- 0.6 mm/a in Jadrtovac, ~ 100 km northwest of

188 Korčula. A dataset of speleothem dates from submerged caves exists (Juračić and Surić, 2010) but 189 most of these samples predate the LGM; a specimen from ca. 9.2 cal ka BP is relevant to the current 190 study, though these indicators constitute only a maximum possible elevation for RSL. Vacchi et al. 191 (2016) used the ICE-5G (VM2) GIA model (Peltier, 2004) of the last 15 ka to predict a RSL rise of 192 less than 1 m since 4 ka BP, preceded by a much faster increase. However, they also noted that the 193 sea-level points indicate an RSL approximately 1 m lower than the model during the last 3 ka, and 194 hypothesize a general subsidence trend in the wider area. The observational data in this study area 195 (Faivre et al., 2013; Vacchi et al., 2016), however, were mostly limited to the last 3 ka BP. 196 Unfortunately for central-southern Dalmatia no valuable index points from coastal depressions are 197 available to constrain relative sea level, like the ones obtained by Brunović et al. (2019) for the 198 north-eastern Adriatic. In this area Antonioli et al. (2007) applied a methodology similar to the one 199 used in this work using a GIA model by Lambeck & Purcell (2005) that predicts roughly similar 200 changes for the last 12 ka in Istria (north-eastern Adriatic), but with the stabilizing point occurring 201 at ca. 6 ka BP instead of 4 ka BP.

# 202 **3** Materials and methods

## 203 3.1 Bathymetric data

This study utilizes a combination of publicly available bathymetric data sources. These include digitized nautical charts, and SRTM15\_PLUS data from NOAA ERDDAP (Sandwell et al., 2014a), which is a 15" version of the Shuttle Radar Topography Mission (SRTM) dataset containing bathymetric data from a combination of sources, chiefly from satellite altimetry gravity elevations (Smith and Sandwell, 1997). The bathymetric data have been combined with topographic digital elevation data of 1" resolution (~30 m) from the SRTM. Table 1 lists these datasets and their characteristics.

DATASET	Projection	Horizontal Datum	Vertical Datum	Resolution	Туре	Citation
NASA SRTM Shuttle radar	Geographic	WGS84	EGM96	1" or ~22 m at latitude (emerged areas only)	Raster (with vector coastline)	(NASA, 2000)
SRTM15_PLUS	Geographic	WGS84	Unspecified	15" or ~340 m at latitude	Raster	(Olson et al., 2014, 2016; Sandwell et al., 2014a)
Chart soundings	Geographic	Lat/Lon	Presumed to be mareograph derived lowest astronomical tide	Varies: 10 ^1 or 10^2 m distance between points in harbours or channels. 10^3 m distances in deep seas	Vector	(Croatian Geoportal, 2019; Državni Hidrografski Institut, 1998)
				Non homogeneous		

211 *Table 1. Overview of the bathymetric and topographic datasets.* 

The bathymetric and nautical sounding datasets were combined into a single point cloud using QGIS so that the high resolution SRTM 1" and nautical chart data supersedes the SRTM15\_PLUS 15" data in emerged areas, and a raster interpolation was created using the GRASS GIS algorithm v.surf.rst. Satellite bathymetry, with its high vertical resolution but poor horizontal resolution, was also eliminated within ~500 m of all coastlines to favour the more accurate nautical chart soundings available in those areas.

## 218 **3.2 Uncertainties**

The vertical datum for the Croatian chart soundings used in this DEM are based on HRSDM71, i.e. Mean Lower Low Waters of Spring Tide from tide gauges at Dubrovnik, Split, Bakar, Rovinj and Koper (epoch 1971.5) (Domijan et al., 2005; HHI, 2019). A  $\pm 0.5$  m is a cautious uncertainty for these whole-number elevations. The satellite altimetry data is heavily post processed to remove wave noise (Sandwell et al., 2014b; Smith and Sandwell, 1997), and come from electromagnetic pulse return times between satellites and the ocean surface at time of sampling, but uncertainties 225 surrounding the geoid height, GPS technologies, and the ship soundings used for calibration make 226 calculating an overall vertical uncertainty difficult. See Tozen et al. (2019) for further discussion of 227 error distributions. In any case, the ~0.5 km spatial resolution of the satellite derived bathymetry in 228 the SRTM15 dataset and the heterogenous spatial resolution and unspecified XY precision of 229 nautical chart soundings mean that XY errors far outweigh the vertical Z uncertainties created by 230 harmonizing the datasets used in the DEM in this area where the largest recorded tidal ranges are 231 less than 0.5 m (tide-forecast.com, 2019). The present study excludes datapoints within 1 km of 232 coastlines, and gives precedence to chart soundings when they conflict with the satellite data. We 233 The SRTM15\_PLUS dataset in the study area was also validated with the chart soundings and it 234 was noted that in most deep areas (~100 m) the predicted depths correspond well (less than 10 m 235 disparity) with available chart soundings.

## 236 **3.3 GIA modelling**

237 For this work, new realizations of GIA models calculated specifically for Korčula were used. 238 Previous regional studies based on an GIA models (Lambeck and Purcell, 2005), have used 239 calculations of GIA for Istria (Antonioli et al., 2007) and Vis (Vacchi et al., 2016) but these span 240 only the last 12-14 ka BP, while Vela Spila's archaeological record goes back to ca. 20 ka BP. The 241 GIA results were obtained by solving the sea-level equation with the program SELEN developed by 242 Spada and Stocchi (2007). In its current implementation, SELEN (version 4) accounts for the 243 migration of shorelines and rotational effects on sea level (Spada and Galassi, 2017). The 244 realizations derive from two different GIA models, based on different assumptions about the 245 melting chronologies of Late Pleistocene ice sheets, and the viscosity profile of the mantle. The first 246 model is the ICE-6G\_C (VM5a) of Peltier et al. (2015) while the second has been progressively developed by K. Lambeck and co-workers at the Australian National University (ANU) (for more 247 248 information, see e.g., Lambeck et al. (2003) and subsequent refinements). These two realizations

249 are henceforth referred to as I6G\_S4 and ANU\_S4, where S4 is an abbreviation for SELEN (version 250 4). Since calculating and illustrating an uncertainty for a GIA model is difficult owing to the large 251 number of variables, assumptions, and parameters that comprise the final prediction (Melini and 252 Spada, 2019), having two models allows us to use the difference between them as an envelope of 253 uncertainty. The RSL reconstructions, the bathymetric digital elevation map, and the archaeological 254 stratigraphy were analysed together in order to reconstruct the palaeogeographic and environmental 255 conditions during the relevant archaeological phases. A literature review for sea-level indicators in 256 the region was also undertaken in order to provide verification for the models.

## 257 3.4 Archaeological data

258 The most relevant palaeoenvironmental aspects inferred from the observation of the Vela Spila 259 sediments have been summarised from published work (Branscombe et al., 2020; Čečuk and Radić, 260 2005; Cristiani et al., 2014; Farbstein et al., 2012; Rainsford et al., 2014) and complimented with 261 unpublished observations made in the four cave key trenches (Fig 3) during the systematic 262 excavations carried out since 2010. Chronology used in this study is partly derived from published 263 radiocarbon dates collected (Table 2) from previously unpublished dates and published sources 264 (Čečuk and Radić, 2005; Farbstein et al., 2012; Rainsford et al., 2014; Srdoč et al., 1989), and 265 partly from the typology of the archaeological remains, the chronology of which is assessed 266 confidently in Dalmatia (Kaczanowska and Kozłowski, 2018; Komšo, 2008; Kozlowski and 267 Kaczanowska, 2004; Miracle and Forenbaher, 2005). In this study, dates referring to specific  $C^{14}$ 268 results use cal a or cal ka BP, whereas dates specifying general periods, GIA models, archaeological 269 typology, or other methods use a or ka BP.

# 270 **4** Results

## 271 4.1 Palaeoenvironmental evidence from archaeological stratigraphy

The most relevant palaeoenviromental changes from Vela Spila sediments are outlined here from bottom to top in order to relate them to shoreline changes. The general stratigraphy of the Vela Spila archaeological deposit is presented in Fig 4. The cave bedrock was not found in any of the excavated trenches, despite maximum excavation depths of about 5 m below the present-day cave floor, so the question of the oldest occupation of the cave is still open. Table 2 lists the collected <sup>14</sup>C for this site, with calibration results summarised in Fig. 5 and referenced below in the text as well.

Lab Code	14C Age (a BP)	2-sigma calibrated age (a BP)	Method	Material dated	Reference	Attribution	Context
Z-3987	16200±200	19,530- 19,190	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Palaeolithic	LUP A
VERA- 2338	16140±60	19665- 19255	AMS	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP A
VERA- 2339	15690±70	19130- 18775	AMS	AB	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP A
z-3993	14500±100	17950- 17410	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP C
z-3992	14100±100	17587- 17105	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP D
z-3991	13300±100	16280- 15700	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP E
z-3988	12800±100	15660- 14925	Std	WC	This paper*, Rainsford et. al. 2014	Palaeolithic	LUP G
z-3990	12700±100	15500- 14655	Std	WC	This paper	Palaeolithic	LUP G
z-3989	12700±100	15500- 14655	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP G
VERA- 2345	12290±40	14490- 14050	AMS	AB	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	NYT
VERA- 2346	12269±40	14410- 14025	AMS	WC	Čečuk & Radić 2005: 34	Palaeolithic	NYT
Wk- 27370	12457±57	14995- 14230	AMS	AB	This paper*, Rainsford et. al. 2014	Palaeolithic	LUP I
VERA- 2343	8290±35	9425-9135	AMS	AB	This paper	Mesolithic	Meso A
VERA- 2344	8230±35	9395-9030	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso A
VERA- 2341	8200±30	9270-9030	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso A

z-3995	8200±70	9400-9005	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso B
z-3986	8200±70	9400-9005	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	
OxA- 18171	8110±37	9235-8985	AMS	HB	Forenbaher <i>et al.</i> 2013: 591	Mesolithic	Meso B
Wk- 24217	8004±41	9010-8715	AMS	HB	This paper		Burial 2/1986
z-3994	7410±70	8375-8045	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso D
Wk- 24216	7350±39	8305-8025	AMS	AB	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso D
Z-1967	7300±120	8375-7930	Std	WC	Čečuk & Radić 2005: 62	Early Neolithic?	Meso D?
VERA- 2342	7175±35	8045-7935	AMS	AB	This paper	Mesolithic	Meso D?
VERA- 2340	7200±30	8155-7950	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso D?
Z-1968	7000±120	8030-7595	Std	WC	Čečuk & Radić 2005: 82	Early Neolithic?	Early Neo?
Wk- 24219	6735±35	7665-7515	AMS	AB	This paper	Neolithic	Neo C
Wk- 24218	6678 ± 34	7610-7485	AMS	AB	This paper*, Rainsford <i>et al</i> . 2014	Neolithic	Neo C
z-1742	5430±100	6410-5950	Std	WC	Srdoč <i>et al.</i> 1989: 86	Neolithic ?	Neo?

Table 2. Radiocarbon age dataset from Vela Spila. Std: standard method; AMS: accelerator mass spectrometry; WC: wood charcoal; AB: animal bone; HB: human bone. \*Uncalibrated date published for the first time in this paper.

## 278 4.1.1 Palaeolithic

279 The lowermost unit of the sequence (Fig. 4) exposed so far is represented by a 3 m-thick talus 280 made up of unsorted frost slabs with an open-work fabric, and with no traces of human occupation. 281 Even if geochronometric dates are not available for this unit, it can be attributed to the LGM 282 because of its sedimentological characteristics and of the radiocarbon age of the overlying units 283 (Farbstein et al., 2012). This type of sediment is consistent with a cold climatic spell typically connected with an RSL lowstand. 284 285 The sediments testifying to the first human occupation of the cave directly overlie the LGM talus and include Upper Palaeolithic remains <sup>14</sup>C constrained to approximately 19.7-14.0 cal ka BP 286

287 (Fig. 5, Table 2). These strata are composed of frost slabs that decrease in quantity and become finer

288 upwards to the top of the Late Upper Pleistocene units. Alternations of discontinuous fine and

289 coarse-grained layers and/or gravelly lenses, as well as some blocks, testify to warmer and colder 290 climate fluctuations, and to localised deposition of gravel from inhomogeneous gelifraction of the 291 ceiling. The silty loamy fine fraction of the sediments suggests colluvial inputs of windblown 292 sediments previously deposited outside the cave or, when more clayey and reddish, of material 293 deriving from the dismantling of soils. The top of the sequence is marked by a horizon of large 294 limestone slabs overlain by a 15 cm-thick layer of sub-primary tephra locally deformed by frost 295 action, ascribed to the Neapolitan Yellow Tuff (NYT) ejected from a volcano in the Campi Flegrei near Naples (Radić et al., 2007),  ${}^{14}$ C dated to 12.3 ±0.04 ka BP (14.5-14.1 cal ka BP; Tab. 2; Fig. 5) 296 297 (Čečuk and Radić, 2005; Farbstein et al., 2012), which fits reasonably well the 14.9 ±0.4 ka BP 298  $^{40}$ Ar/ $^{39}$ Ar age of the tephra (Deino et al., 2004) and much better its modelled age 14.3–13.9 cal ka 299 BP date obtained by Bayesian method (Blockley et al., 2008).

300 The anthropic component of these layers –abundant organic matter, charcoal, bone and stone 301 tools deriving from intense occupation of the cave – is dominant over non-anthropic material and 302 gives the sediment a very dark, blackish colour value. Archaeological materials include Late Upper 303 Palaeolithic lithics of Epigravettian tradition (Farbstein et al., 2012; Rainsford et al., 2014), sharing 304 strict analogies with other technocomplexes of Dalmatia and its hinterland, e.g. Pupićina peć 305 (Miracle, 2007), Šandalja II (Janković et al., 2012; Karavanić, 2003; Richards et al., 2015), Vlakno 306 (Cvitkušić et al., 2018; Vukosavljević and Perhoč, 2017), Zemunica (Šošić Klindžić et al., 2015), 307 Kopačina (Vukosavljević et al., 2011; Vukosavljević and Perhoč, 2017), Badanj (Whallon, 1999), 308 Crvena Stijena (Basler, 1975; Kozlowski and Kaczanowska, 2004). The large mammal fauna is 309 dominated by Cervus elaphus (red deer), followed by the dry environment taxon Equus hydruntinus 310 (European ass). Other less frequent taxa are Capreolus capreolus (roe deer), Bos/Bison (wild 311 cattle/bison), Sus scrofa (wild boar) and Lepus sp. (hare). A few specimens of Canis lupus (wolf), 312 Felis lynx (lynx), Felis silvestris (wild cat), Vulpes vulpes (red fox) are also present. Smaller 313 vertebrate remains (e.g., rodents, birds, bats, reptiles, fish) are not abundant in the Pleistocene

layers. The rich and abundant faunal assemblage suggests a landscape characterised by wide-open
areas with wooded patches that was resource-rich, productive and able to sustain a large community
of hunters-gatherers.

317 4.1.2 Abandonment phase

318 A complex and not yet fully understood human occupation gap of approximately 4.6 ka 319 occurred between the latest Late Upper Palaeolithic and the earliest Mesolithic phases (Radić, 320 2018). At the foot of the southern wall of the cave, the Mesolithic layers overlie the NYT tephra 321 directly. Conversely, at the centre of the cave huge limestone blocks, up to 2.5 m large, rest on the 322 NYT and are partially covered by about 1.4 m of limestone rubble mixed with reddish silty loam -323 with no cultural remains. The base of Mesolithic layers is here about 1 m lower than at the foot of 324 the wall and overlies the rubble, apparently without sedimentary gaps. This situation suggests that 325 throughout the Late glacial the topography of the inside of the cave had been pre-determined by the 326 shape of the LGM scree, which had partly filled the cave from the western entrance and along the 327 southern wall. The central area was strongly depressed, so that the accumulation of the Late Upper 328 Palaeolithic layers did not level it completely. Later on, the deposition of ceiling breakdown blocks 329 was triggered by frost in some still undated -posterior to NYT - cold phase of the Late Glacial and 330 was followed by rubble and colluvium of reddish silty loam during the subsequent climate 331 improvement. The fine component of these sediments include mica flakes and other minerals 332 typical of the mineral suite of loess or loess-like sediments of the Adriatic area, and their grain-size 333 supports this (Cremaschi, 1990; Wacha et al., 2011a, 2011b). These sediments can be found only in 334 the central part of the cave, for the following two reasons: i) large breakdown blocks and rubble 335 accumulated only under the ceiling opening, which is located above this area; ii) colluvium flowing 336 into the cave from the western entrance, and probably also from the ceiling opening, was 337 preferentially deposited in the topographic low.

#### 338 *4.1.3 Mesolithic*

339 This topographic setting also influenced the deposition of the Mesolithic layers. Thickness and 340 composition of sediments including the Mesolithic cultural remains are variable throughout the 341 cave; the bottom areas of the cave are rich in ash and charcoal remains, whereas the central part of 342 the cave was still slightly depressed when people started occupying the site again around 9.4 cal ka 343 BP. In the centre these layers are somewhat thick (about 1.3 m) and largely composed of fine 344 sediment deriving from the runoff of reworked red soils and/or windblown sediments, whereas they 345 are much thinner (about 20-60 cm) and stony at the foot of the southern wall. These characteristics 346 suggest that the topography of the cave (Fig. 3) inside was different from the present-day one, with 347 a large depression at its centre, and that different areas of the cave were used for specific purposes 348 by humans.

The age of the Mesolithic occupation is ca. 9.4 to 8.0 cal ka BP (Farbstein et al., 2012; Rainsford et al., 2014). Faunal remains from these layers indicate a reduction of the typical Late Pleistocene open environment mammal taxa, with a shift towards associations preferring good vegetative shrubby or woody cover, represented here mostly by *Vulpes vulpes* (fox), *Capreolus capreolus* (roe deer), *Sus scrofa* (wild boar). Among the infrequent remains of other small-size taxa are those of *Erinaceus roumanicus* (Northern white-breasted hedgehog) (Mauch Lenardić et al., 2018).

The decrease in mammal taxa size and a smaller number of ungulate remains testify to a generalised reduction of high-quality food resources, which was replaced by the exploitation of marine taxa. The Mesolithic units include remains of *Delphinidae* sp. (dolphin) and *Monachus monachus* (monk seal), as well as abundant *Scomber japonicus* (mackerel) and other fish taxa (Rainsford et al., 2014). These marine vertebrates are associated with extremely abundant molluscs like *Patella caerulea* and *Patella rustica* (limpet) and Trochidae, mostly *Phorcus turbinatus*  362 (turbinate monodont) (Branscombe et al. 2020). Several other marine mollusc taxa- mostly adapted 363 to rocky coasts – are also well represented, even if less abundant than the aforementioned ones; 364 continental molluscs are also common, as in most Adriatic area sites (Rizner et al., 2009), and 365 mostly represented by Eobania vermiculata (chocolate band snail). Other Mesolithic sites of 366 Dalmatia and neighbouring areas that share similar techno-typological and economic characteristics are Pupićina peć (Miracle, 2007, 2002, 2001), Abri Šebrn (Miracle et al., 2000), Nugljanska peć 367 368 (Pilaar Birch and Miracle 2015), Vela Spilja Lošinj (Pilaar Birch and Miracle 2017), Vlakno cave 369 (Vukosavljević et al., 2014; Cvitkušić et al., 2018), Zemunica (Šošić Klindžić et al., 2015). The 370 remarkable amount of marine resource remains found in the cave, including large quantities of 371 mackerel bones, pierced Columbella (dove snail) shells, and a relatively few large fish bones 372 (possibly associated with deep-sea fishing), indicates an economic shift towards the sea.

## 373 *4.1.4 Neolithic*

374 The age of the Holocene sediments (including the earliest Neolithic remains of Vela Spila 375 ascribed to the Impressa Ware cultural horizon) is bracketed between 8.0 and 6.0 cal ka BP (Čečuk 376 and Radić, 2001). During the Neolithic there was virtually no deposition of clastic sediments 377 derived by soil erosion from the outside of the cave, which is interpreted as a sign of extended 378 vegetation on the slopes, favoured by mild climatic condition and evenly distributed rainfall. 379 Depositional rates were however high inside the cave, because penning of domesticated animals, 380 mostly sheep and goats, produced large quantities of dung that was intentionally burned. Typical 381 sediments of this phase are the so-called "layer-cake" or *fumier* sequences (Boschian and Miracle, 382 2007), characterised by long successions of cyclically alternating black and white layers including 383 ash, charcoal and other products of herbivore metabolism. From the Neolithic onwards, depositional 384 environments in caves change in this way all over the Mediterranean area (Angelucci et al., 2009), 385 demonstrating the new role of caves within a complex agropastoral system. Similar *fumier* 

sequences were excavated in other caves on both sides of the Adriatic Sea, e.g. in several north
Adriatic Karst sites (Boschian and Montagnari-Kokelj, 2000), at Pupićina peć (Boschian, 2006;
Forenbaher and Miracle, 2006) in Istria, at Zemunica Peć (Šošić Klindžić et al., 2015) in central
Dalmatia, and at Grotta dei Piccioni and Grotta S. Angelo (Boschian and Iaconis, 2008) in central
Italy.

Neolithic deposits contain abundant bones of domestic animals (mostly sheep and goats), few wild animal bones, some terrestrial and marine mollusc shells, many potsherds, chert from the Gargano Peninsula (c. 125 km to the southwest) and obsidian from Lipari (c. 500 km to the southwest). Evidence of soil erosion is again apparent in the sequence corresponding with the Late Neolithic onwards. In this case erosion is interpreted as evidence of human activities (agriculture and pastoralism) which caused deforestation.

A long-lasting and intensive Neolithic occupation of the site is represented by a complete
sequence of cultural facies and characteristic pottery of all major styles occurring in the eastern
Adriatic area during this period (Forenbaher et al., 2013), testifying to the sociocultural and
economic aspects of the Neolithic transition (Zvelebil and Lillie, 2000) and the further evolution of
the Neolithic towards Copper Age and Early Bronze Age.

402 Throughout the period of occupation subsistence depended heavily on herding. Faunal
403 assemblages are dominated by domestic animal remains; primarily sheep and goats, while cattle and
404 pigs are less common. Also present in relatively small quantities are terrestrial and marine molluscs,
405 while fish remains are quite rare.

Despite systematic attempts to recover seeds through flotation of soil samples, direct evidence of agriculture remains scarce, but this should not be interpreted as indicating an absence of plant cultivation. Productive arable land tends to be located some distance away from the heavily karstified areas where caves abound, and a similar absence of cultivars was observed in other caves in the region (Borojević et al., 2008; Forenbaher and Miracle, 2006). After the Neolithic, the archaeological sequence continues through the Copper and Bronze Ages (Radić, 2018), with rich
ceramic assemblages and sediments of mostly *terra rossa* colluvium and anthropic components
represented by variable amounts of organic matter and stable sediments.

## 414 **4.2** Sea-level change and palaeogeography around Vela Spila

415 According to the model predictions (Fig. 6), RSL was 101-107 m below present at the start of 416 Vela Spila's Upper Palaeolithic occupational phase, and then climbed at a rate of about 5 to 6 mm/a. 417 Combining the isostatic model with the bathymetric analysis, it is possible to infer that the 418 landscape around the Island of Korčula changed remarkably over the site's prehistoric phases (Table 419 3). Contemporaneous with the first detected Upper Palaeolithic occupation (starting at ~19.7 cal ka 420 BP, Fig. 7A), the cave was ~8-9 km from the sea and 221-228 m above it, while the area south of 421 western Korčula consisted of a broad plain where the present-day Lastovo island group appeared as 422 a cluster of hills. 7-16% of the land within 20 km of the site (a day's round-trip travel at a 423 comfortable walk) was inundated. Korčula was still also connected to the mainland in the east via 424 the Pelješac Peninsula and Hvar Island to the north. To the SW of the present bay of Vela Luka, a 0-425 30 m deep sea existed, and to the NW Vis and Biševo were also not yet islands. South of Vela Luka 426 bay, a natural depression indicated by the bathymetric data might have been filled by a small, 427 shallow lake or marsh a few meters deep and a couple kilometres wide. To the N and NE of present 428 day Korčula, the palaeo-Neretva river flowed in an alluvial plain between Hvar and Korčula, shown 429 in GIS reconstructions carried out by Sikora et al. (2014).

430

Archaeological phase	Start (ka BP)	End (ka BP)	) RSL at start (m)	RSL at end (M)	RSL change (m)	Average rate of change (mm/a)	Average Horizontal Coastline Change (m/a)	Vertical RSL Change in human lifetime (m)	Horizontal coastline change in human lifetime (m)
Palaeolithic	19.7	14.0	-107 to -100	-74 to -72	29 to 33	5 to 6	5	0.2	160

Abandonment	14.0	9.4	-74 to -72	-30 to -16	44 to 55	10 to 12	14	0.4	492
Mesolithic	9.4	8.0	-30 to -16	-13 to -7	9 to 17	7 to 12	0.5 in Vela Luka Bay	0.2 to 0.4	18 in Vela Luka Bay
Neolithic	8.0	6.0	-13 to -7	-3 to -1	6 to 10	3 to 5	0.1 in Vela Luka Bay	0.1 to 0.2	4 in Vela Luka Bay

Table 3. Palaeogeographic changes by archaeological period at Vela Spila. In the Mesolithic and Neolithic, coastal migration around the island becomes insignificant so the rates of incursion into the long, valley-like Vela Luka Bay are noted instead. N.b. the distinction between mm/a (mm per year) and m/a (meters per year).

431 By the end of the Palaeolithic occupation at 14.0 ka BP (Fig. 7B), ca. 5 ka later, RSL had risen 432 29 to 33 m (Table 3), the coastline had advanced 4-10 km into Vela Luka Bay, and Vis and Biševo 433 were likely already islands. At this point, about 40% of land within 20 km of the site was flooded. 434 When the cave was abandoned, sometime after the NYT eruption between 14.9 and 13.9 cal ka BP 435 (Blockley et al., 2008; Deino et al., 2004), any nearby lake or depression south of Vela Luka Bay 436 had already become part of the Adriatic Sea, which had been advancing eastward up to 4 m/a 437 (meters per year) in this area. 438 According to our models, the most dramatic RSL rise and landscape change around Korčula 439 occurred during the roughly 5 ka (14.0-9.4 cal ka BP) period when the cave was abandoned. RSL 440 rose ~44 to 55 m at an average of about 10-13 mm/a and the coastline advanced eastward by as

much as 10 to 12 m/a or more to the north and south of the island (Table 3). Land connections to

442 areas north and south of Korčula, such as present day Lastovo and Hvar, were probably severed

443 during the first millennium of the abandonment phase.

441

Much of the coastlines of the islands advanced to within 1 km of their present position by ca. 12 ka BP, after which horizontal changes in coastlines became much smaller: above the ~70 m isobath, slopes around the island tend to be much steeper (Fig. 8). A land bridge between the Pelješac Peninsula and Korčula may have persisted until between 12 and 10 ka BP, at which time the geography of the area became quite similar to the present-day. When the cave was finally reoccupied at the start of the Mesolithic at ~8.0 ka BP (Fig. 7C, Fig. 9), the surrounding landscape had been totally transformed. About 80% of nearby lands were flooded. 30-50 m deep seas were 2 km away from the cave to the north and south, and a shallow (10-20 m) bay was only 1-2 km to the west. No land connections with other present-day islands existed, though many of the smaller islets around Korčula may still have been connected to it at times. By ca. 8.0 ka BP, at the beginning of the Neolithic, most of the northern and southern coasts of the island were even closer to the presentday shoreline, though some islets may still have been connected, and the inner part of Vela Luka harbour had not been fully submerged.

# 457 **5 Discussion**

## 458 **5.1 Last Palaeolithic occupants facing rapidly inundating planes**

459 The Palaeolithic people at Vela Spila experienced average RSL change rates of 5–6 mm/a, 460 although the ANU\_S4 realization does predict some phases of higher rates. The rate of RSL change 461 increased to 10-12 mm/a during the abandonment phase (Table 3). Hence the Late Upper 462 Palaeolithic coastline was relatively stable compared to that of the subsequent abandonment phase. 463 The cave was only 8–9 km distant from the sea, and the lack of shells in the archaeological 464 stratigraphy during this phase of inhabitation is not necessarily a consequence of distance from the 465 cave but, more likely, a preference for the terrestrial species which had been the traditional Upper 466 Palaeolithic food resource (Miracle, 2007; Mussi and Palombo, 2001; Stewart et al., 2003) and 467 would still have been widely available at this point, given that they occupy a large part of the diet. 468 The palaeo-Neretva's plain north of Korčula became submerged during the abandonment phase at 469 Vela Spila, and in those millennia the high rates of RSL change may not have left time for the 470 formation of submerged landforms such as palaeocoastlines or palaeodeltas (Sikora et al., 2014). 471 During the Palaeolithic phase, only a relatively small area of potential hunting grounds was 472 inundated in the plains north and south of Korčula. Still, the coastline moved east at a rate of nearly 473 5 m/a, so that in the average human lifespan of 35 years, a person would have witnessed ca. 160 474 meters of horizontal coastal change. The rate of change experienced by the Upper Palaeolithic 475 foragers in the Central Adriatic was 6-7 times greater than the present-day estimates of ongoing 476 RSL change (Church et al., 2013; Neumann et al., 2015). It is most likely that this landscape change 477 stressed littoral habitats and limited potential human exploitation of marine resources; however, 478 sediments and cultural remains show that human frequentation continued to be intensive in the 479 cave. Following on the Palaeolithic occupation at Vela Spila, the abandonment phase (from ca. 14.0 480 to 9.4 cal ka BP) appears to have seen even more intense changes in the surrounding 481 palaeogeography.

### 482 **5.2** Environmental reasons for abandoning the cave: rising sea or climate?

483 The deposition into Vela Spila of in-washed NYT tephra (Fig. 4) coincided or preceded 484 approximately the end of the Late Upper Palaeolithic occupation and the beginning of a long gap in 485 the human record (14.0–9.4 cal ka BP), at the start of which the RSL was 74–72 m below present. 486 Since the eruption took place about 300 km SW of Vela Spila, it is worth speculating what role, if 487 any, this event may have played in inducing Upper Palaeolithic people to abandon the cave. 488 Cryoturbation features involving specifically the NYT sediment in the cave suggest that deposition 489 was associated or immediately followed by a cold spell, though it cannot be ascertained if it was 490 consequential to the eruption.

491 Rapid landscape changes caused by a high RSL rise rate may have had a greater influence on 492 human behaviour and settlement pattern change during this phase; predicted RSL changes were 493 much faster than during the Upper Palaeolithic cave occupation. Humans would have witnessed 494 nearly 0.40 m of RSL rise, corresponding to about 500 m of horizontal coastline displacement 495 during a 35-year average lifespan, i.e. about 14 m of horizontal eastward coastal change per year 496 immediately north and south of Korčula. This suggests a highly unstable or marshy littoral zone, rapid flooding of the traditional foraging grounds of the Upper Palaeolithic groups, and stress on the
ecosystem at all trophic levels. Particularly rapid change (about 20 m rise) took place during the
centuries following the Meltwater Pulse 1A (MWP-1A), which has been dated to 14.7–14.3 ka BP
(Deschamps et al., 2012).

501 Volcanic eruptions such as the NYT on the other hand usually have an impact on climate for only a few years, so that their consequences are not easily detected in  $\delta^{18}$ O isotope or other climate 502 503 proxies (Aufgebauer et al., 2012). The NYT eruption also coincides with the end of a colder climate 504 phase highlighted at Lake Ohrid (Vogel et al., 2010), and with increased winter precipitation 505 (Bordon et al., 2009; Aufgebauer et al., 2012). If the settlement was already in a marginal position 506 due to climate changes and to the landscape changes highlighted by this study, a sequence of 507 particularly unfavourable years due to an eruption may have contributed to pushing the Late Upper 508 Palaeolithic people to a tipping point that eventually resulted in abandonment of the site. 509 The first two thousand years of the Vela Spila occupation gap overlap with, in addition to the 510 NYT and Meltwater Pulse 1A, the Bølling–Allerød warming, and were followed by the Younger 511 Dryas major dry/cold event (Gkinis et al., 2014), which is recorded throughout the Balkans in 512 various lacustrine sequences (Cvetkoska et al., 2014). At many Late Upper Palaeolithic sites in the 513 Eastern Adriatic region the Bølling-Allerød warming is associated an increase in of human 514 settlement (Miracle, 2007), and the same pattern is largely true on the Italian side of the Adriatic 515 (Mussi 2001). The occupational sequence at Vela Spila is in stark contrast to this wider pattern. 516 Younger Dryas effects in the Adriatic Plain are indicated at Vela Spila by the occurrence of 517 partially altered loess-like sediments colluviated into the cave. The precise timing of this 518 colluviation is currently unknown; it probably was partly during the Younger Dryas, but mostly 519 during the early Holocene. It is, however, unclear that the Bølling–Allerød or Younger Dryas 520 climate changes played a direct role in forcing the Late Upper Palaeolithic foragers to leave the site, 521 although the Younger Dryas is correlated with cultural change and adaptation throughout Europe

(Aura et al., 2011; Burdukiewicz, 2011; Mussi and Peresani, 2011; Weber et al., 2011). Instead, the
rapid RSL rise concurrent with the meltwater pulse 1A and Bølling–Allerød may have been the
environmental trigger for the abandonment of Vela Spila, since the cave had already been
abandoned by the time of the Younger Dryas at ca. 12 or 13 ka BP (Gkinis et al., 2014).

527 **5.3 Stabilising coastlines and the Mesolithic** 

528 Benjamin et al. (2017) suggest that the Northern Hemisphere cooling during the Younger Dryas 529 slowed the RSL rise in the Mediterranean as in other areas in the world. The GIA models in this 530 work (Fig. 6) also predict a decrease in RSL rise rates following the Younger Dryas, but only few 531 local relative sea-level indicators are available for verification. By ca. 9.4 ka BP, when the model 532 predicts RSL rates were starting to level off, Mesolithic groups re-occupied the cave and continued 533 dwelling in it for the following ~ 1.5 ka. The corresponding units of the cave sequence show 534 evidence of the Preboreal and Boreal humid climate conditions indicated by colluvia. In this period 535 RSL continued rising quickly from between -30 to -16 below present, up to between -13 m and -7 m 536 - i.e. a rise of 9 m to 17 m in approximately 1.5 ka (Table 3), which gives a rate of 7 to 12 mm/a, 537 though this is a slower rate of change than during the abandonment phase. The sea was only about 538 1-2 km distant from the cave at the onset of the Mesolithic, compared to 8-9 km in the Upper 539 Palaeolithic.

The Mesolithic assemblage is dominated by the remains of shellfish, fish and other marine creatures. The exploitation of marine resources by Mesolithic people is widespread in the Mediterranean and the Atlantic, though the Mediterranean is supposed to be less productive (Colonese et al., 2011). In general, resource depression and human population pressure have been considered the main reason for shifting toward marine resources (Colonese et al., 2011), even if there are major differences between regions. However, it does not appear that the exploitation of 546 fish and shellfish was ever the most important food resource in the Adriatic region; isotopic studies 547 on human remains show that "while the Mesolithic people of the Adriatic coast consumed some 548 marine protein, terrestrial protein constituted a significant part of the diet" (Lightfoot et al., 2011, p. 549 82). The balance between the need for resource diversification and expedience of marine resources 550 due to the approaching coastline is not always clear or consistent. As the Mesolithic progresses, 551 there is a decline in the use of fish and an increase in shellfish and snails at Vela Spila, along with a 552 decrease in terrestrial fauna remains. Fishing seems to shift from open-sea to inshore taxa 553 (Rainsford et al. 2014). This is partially in contrast with data from the northern Adriatic where the 554 coastline approached later: here, fishing shifts from freshwater to marine taxa and Early Mesolithic intensive snail exploitation is substituted by consumption of marine species (Boschian and Pitti, 555 556 1984; Cremonesi et al., 1984).

557 These data suggest that, in the northern Adriatic, the need for resource diversification was the 558 driving force towards marine food well before landscape modification due to RSL rise. In contrast, 559 the same need started more or less contemporaneously at Vela Spila, but was immediately oriented 560 towards marine resources because of the earlier advance of the coastline. The decrease in other 561 resources is testified by the disappearance of red deer and other large ungulates from the Mesolithic 562 layers, presumably because of the early Holocene palaeoenvironmental turnover, and because 563 Korčula had become an island where faunal replacement was impossible as access to more productive hunting grounds decreased. 564

The transition between Mesolithic and Neolithic started in the Eastern Adriatic at ca. 8.0 ka BP (Forenbaher et al., 2013). Radiocarbon determinations and cultural assemblages from Vela Spila are consistent with this date. The Mesolithic-Neolithic transition shortly follows the 8.2 ka event, which was a disruption in thermohaline Atlantic circulation owing to meltwater input from North America (Alley and Ágústsdóttir, 2005). The 8.2 ka BP event is associated with an estimated drop in winter temperatures of more than 4° C, at least in Northeastern Greece (Pross et al., 2009) and the sea571 level impact of the event in the study area is approximately 1 m (Kendall et al., 2008), although this 572 perturbation is too rapid and small to appear in the GIA models. Furthermore, because the coastlines 573 were already near their present-day position abutting the high relief slopes of the island (Fig. 8), the 574 RSL rise itself would not have been significant for the inhabitants unless they had built immobile 575 coastal infrastructure, of which no evidence exists. Likewise, no clear fingerprint of the 8.2 event 576 exists in the archaeological record of the site. By the end of the Neolithic at ca. 6.0 ka BP, RSL had 577 risen to within 3 meters of present level, and sea-level changes from then on were much slower 578 (Fig. 6).

# 579 **6** Conclusions

580 In this work we direct evidence from a detailed, extensively excavated archaeological 581 stratigraphy was used to understand human reactions to changing rates of RSL rise and horizontal 582 coastline shift in a coastal landmass that becomes an island.

In the Late Upper Palaeolithic the abandonment of Vela Spila cave was triggered by environmental events that caused the area to be less attractive for humans. In particular, high rates of sea-level rise likely stressed the least observant inhabitants in Vela Spila more sharply than other environmental factors (e.g. climatic forcing) by submerging hunting grounds and reshaping marine and terrestrial ecosystems in the rapidly disappearing Great Adriatic Plain, where the coastline advanced tens of meters per year.

The coastline positions around Korčula island become stabilized only by ~12 ka BP as the rising seas surpassed the 70 m isobath and encountered the steep morphology of the island's sides (Fig. 8), though rapid RSL rise continued. This state, together with other environmental factors, may have fostered the re-occupation of the cave at ca. 9.4 cal ka BP. In the Mesolithic, it was the new insularity of the island that forced humans to exploit all resources available (e.g. seafood and smaller fauna like foxes, rather than larger animals such as red deer). The disadvantages of this
insularity were initially overcome by skilled seafaring and later, more effectively, by the onset of
agriculture and pastoralism that accompanied, after 8 ka BP, a substantial slowing down of sea-level
rise.

This study also emphasizes the importance of potential future research projects aimed at gathering underwater data, for example by more detailed bathymetric surveys or sediment coring in relevant areas around Vela Luka bay such as the plains to the north and south, the palaeo Neretva, and the strait between Korčula and the Pelješac, which could reveal additional drowned landforms and improve resolution for sea-level change histories and the environmental transformation of Korčula from mainland to island.

## 604 7 Acknowledgements

The authors of this paper would like to recognize the EU project no. 692249 "Smart Integration of Genetics with Sciences of the Past in Croatia: Minding and Mending the Gap" (http://mendthegap.agr.hr/) for providing most of the travel and workshop opportunities for many of the authors to perform research for this paper. Archaeological excavations and analyses were also supported by the British Academy (SG101073), the University of Cambridge, and St John's College, Cambridge.

# 611 8 Works Cited

612

Alley, R.B., Ágústsdóttir, A.M., 2005. The 8k event: cause and consequences of a major

614 Holocene abrupt climate change. Quaternary Science Reviews 24, 1123–1149.

615 https://doi.org/10.1016/j.quascirev.2004.12.004

- 616 Andel, T.H.V., 1989. Late Quaternary sea-level changes and archaeology. Antiquity 63, 733–
- 617 745. https://doi.org/10.1017/S0003598X00076869
- 618 Angelucci, D.E., Boschian, G., Fontanals, M., Pedrotti, A., Vergès, J.M., 2009. Shepherds and
- 619 karst: the use of caves and rock-shelters in the Mediterranean region during the Neolithic. World
- 620 Archaeology 41, 191–214. https://doi.org/10.1080/00438240902843659
- 621 Antonioli, F., Anzidei, M., Lambeck, K., Auriemma, R., Gaddi, D., Furlani, S., Orrù, P.,
- 622 Solinas, E., Gaspari, A., Karinja, S., Kovačić, V., Surace, L., 2007. Sea-level change during the
- 623 Holocene in Sardinia and in the northeastern Adriatic (central Mediterranean Sea) from
- archaeological and geomorphological data. Quaternary Science Reviews 26, 2463–2486.
- 625 https://doi.org/10.1016/j.quascirev.2007.06.022
- 626 Artegiani, A., Paschini, E., Russo, A., Bregant, D., Raicich, F., Pinardi, N., 1997. The Adriatic
- 627 Sea General Circulation. Part II: Baroclinic Circulation Structure. J. Phys. Oceanogr. 27, 1515–
- 628 1532. https://doi.org/10.1175/1520-0485(1997)027<1515:TASGCP>2.0.CO;2
- 629 Aufgebauer, A., Panagiotopoulos, K., Wagner, B., Schaebitz, F., Viehberg, F.A., Vogel, H.,
- 630 Zanchetta, G., Sulpizio, R., Leng, M.J., Damaschke, M., 2012. Climate and environmental change
- 631 in the Balkans over the last 17 ka recorded in sediments from Lake Prespa (Albania/F.Y.R. of
- 632 Macedonia/Greece). Quaternary International, Temporal and spatial corridors of Homo sapiens
- 633 sapiens population dynamics during the Late Pleistocene and Early Holocene 274, 122–135.
- 634 https://doi.org/10.1016/j.quaint.2012.02.015
- Aura, J.E., Jordá, J.F., Montes, L., Utrilla, P., 2011. Human responses to Younger Dryas in the
- 636 Ebro valley and Mediterranean watershed (Eastern Spain). Quaternary International 242, 348–359.
- 637 Basler, Đ., 1975. Crvena stijena: zbornik radova. Zajednica kulturnih ustanova, Nikšić.
- Bekić, L., 2017. Šimuni, novo podvodno brončanodobno nalazište u Zadarskoj županiji.
- 639 Diadora: glasilo Arheološkog muzeja u Zadru 31, 41–50.

- 640 Bekić, L., Pešić, M., Scholz, R., Meštrov, M., 2015. Podvodna arheološka istraživanja na
- 641 prapovijesnom nalazištu Pakoštane–Janice. Diadora: glasilo Arheološkog muzeja u Zadru 29, 7–22.
- 642 Benjamin, J., Bekić, L., Komšo, D., Uhač, I.K., Bonsall, C., 2011. Investigating the submerged
- 643 prehistory of the eastern Adriatic: Progress and prospects. Submerged prehistory 193–206.
- 644 Benjamin, J., Bonsall, C., 2009. An underwater archaeological survey in the northern Adriatic
- 645 Sea. International Journal of Nautical Archaeology 38, 163–172.
- 646 Benjamin, J., Rovere, A., Fontana, A., Furlani, S., Vacchi, M., Inglis, R.H., Galili, E., Antonioli,
- 647 F., Sivan, D., Miko, S., Mourtzas, N., Felja, I., Meredith-Williams, M., Goodman-Tchernov, B.,
- 648 Kolaiti, E., Anzidei, M., Gehrels, R., 2017. Late Quaternary sea-level changes and early human
- 649 societies in the central and eastern Mediterranean Basin: An interdisciplinary review. Quaternary
- 650 International 449, 29–57. https://doi.org/10.1016/j.quaint.2017.06.025
- Bini, M., Pappalardo, M., Rossi, V., Noti, V., Amorosi, A., Sarti, G., 2018. Deciphering the
- effects of human activity on urban areas through morphostratigraphic analysis: The case of Pisa,
- Northwest Italy. Geoarchaeology 33, 43–51. https://doi.org/10.1002/gea.21619
- Birch, S.E.P., Linden, M.V., 2018. A long hard road... Reviewing the evidence for
- 655 environmental change and population history in the eastern Adriatic and western Balkans during the
- 656 Late Pleistocene and Early Holocene. Quaternary International, Impacts of gradual and abrupt
- 657 environmental changes on Late glacial to Middle Holocene cultural changes in Europe 465, 177–
- 658 191. https://doi.org/10.1016/j.quaint.2016.12.035
- 659 Blockley, S.P., Ramsey, C.B., Pyle, D.M., 2008. Improved age modelling and high-precision
- 660 age estimates of late Quaternary tephras, for accurate palaeoclimate reconstruction. Journal of
- 661 Volcanology and Geothermal Research 177, 251–262.
- Blute, M., 2008. CULTURAL ECOLOGY. Encyclopedia of Archaeology.
- 663 https://doi.org/10.1016/B978-012373962-9.00098-4

- Bonacci, O., Ljubenkov, I., Knezić, S., 2012. The water on a small karst island: the island of
- 665 Korčula (Croatia) as an example. Environ Earth Sci 66, 1345–1357. https://doi.org/10.1007/s12665-
- 666 011-1345-9
- Bordon, A., Peyron, O., Lézine, A.-M., Brewer, S., Fouache, E., 2009. Pollen-inferred Late-
- 668 Glacial and Holocene climate in southern Balkans (Lake Maliq). Quaternary International, Rhythms
- and Causalities of the Anthropisation Dynamic in Europe between 8500 and 2500 cal BP:
- 670 Sociocultural and/or Climatic Assumptions 200, 19–30.
- 671 https://doi.org/10.1016/j.quaint.2008.05.014
- 672 Borojević, K., Forenbaher, S., Kaiser, T., Berna, F., 2008. Plant use at Grapčeva cave and in the
- eastern Adriatic Neolithic. Journal of Field Archaeology 33, 279–303.
- Boschian, G., 2006. Geoarchaeology of Pupićina cave, in: Miracle, P., Forenbaher, S. (Eds.),
- 675 Prehistoric Herders of Northern Istria: The Archaeology of Pupićina Cave. Archaeological Museum
- 676 of Istria, Pula, pp. 123–162.
- Boschian, G., Fusco, F., 2007. Figuring Out No-One's Land Why Was the Karst Deserted in the
- 678 Late Glacial?, in: Whallon, R. (Ed.), Late Paleolithic Environments and Cultural Relations Around
- 679 the Adriatic. Archaeopress, Oxford, pp. 15-+.
- Boschian, G., Iaconis, M., 2008. Geoarchaeology of the deposits of Grotta dei Piccioni and
- 681 Grotta Sant'Angelo (Abruzzo, Central Italy), in: Boschian, G. (Ed.), Proceedings of the 2nd
- 682 International Conference on Soils and Archaeology, Atti Società Toscana Scienze Naturali, Mem.
- 683 pp. 181–188.
- Boschian, G., Miracle, P., 2007. Shepherds and caves in the karst of Istria (Croatia). Atti della
  Società Toscana di Scienze Naturali 173–180.
- 686 Boschian, G., Montagnari-Kokelj, E., 2000. Prehistoric shepherds and caves in the Trieste Karst
- 687 (Northeastern Italy). Geoarchaeology 15, 331–371. https://doi.org/10.1002/(SICI)1520-
- 688 6548(200004)15:4<331::AID-GEA3>3.0.CO;2-H

689	Boschian, G., Pitti, C., 1984. I livelli mesolitici della Grotta dell'Edera, in: Il mesolitico sul
690	Carso Triestino, Quaderni della Societá per la Preistoria e Protostoria della Regione Friuli-Venezia
691	Giulia. Italo Svevo, Trieste, pp. 143–210.
692	Branscombe, T.L., Bosch, M.D., Miracle, P.T., 2020. Seasonal Shellfishing across the East
693	Adriatic Mesolithic-Neolithic Transition: Oxygen Isotope Analysis of Phorcus turbinatus from Vela
694	Spila (Croatia). Environmental Archaeology 1–14.
695	Burdukiewicz, J.M., 2011. Late Glacial hunter-gatherer reactions to the Younger Dryas cooling
696	event in the southern and eastern Baltic regions of Europe. Quaternary International 242, 302-312.
697	Butzer, K.W., 2005. Environmental history in the Mediterranean world: cross-disciplinary
698	investigation of cause-and-effect for degradation and soil erosion. Journal of Archaeological
699	Science 32, 1773–1800. https://doi.org/10.1016/j.jas.2005.06.001
700	Carroll, F.A., Hunt, C.O., Schembri, P.J., Bonanno, A., 2012. Holocene climate change,
701	vegetation history and human impact in the Central Mediterranean: evidence from the Maltese
702	Islands. Quaternary Science Reviews 52, 24-40. https://doi.org/10.1016/j.quascirev.2012.07.010
703	Casalbore, D., Falese, F., Martorelli, E., Romagnoli, C., Chiocci, F.L., 2017. Submarine
704	depositional terraces in the Tyrrhenian Sea as a proxy for paleo-sea level reconstruction: Problems
705	and perspective. Quaternary International, Quaternary coastal and marine studies in Central
706	Mediterranean 439, 169–180. https://doi.org/10.1016/j.quaint.2016.02.027
707	Čečuk, B., Radić, D., 2005. Višeslojno pretpovijesno nalazište - Vela Luka, Otok Korčula.
708	Višeslojno pretpovijesno nalazište - Vela Luka, Otok Korčula, Korčula.
709	Čečuk, B., Radić, D., 2001. Vela Špilja - preliminarni rezultati dosadašnjih istraživanja, in:
710	Čečuk, B. (Ed.), Arheološka istraživanja na području otoka Korčule i Lastova. Hrvatsko arheološko
711	društvo, Zagreb, pp. 75–118.

- 712 Čelhar, M., Parica, M., Ilkić, M., Vujević, D., 2017. A Bronze Age underwater site near the islet
- 713 of Ričul in northern Dalmatia (Croatia). SKYLLIS-Zeitschrift für maritime und limnische
- 714 Archäologie und Kulturgeschichte 17, 21.
- 715 Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A.,
- 716 Merrifield, M.A., Milne, G.A., Nerem, R.S., Nunn, P.D., Payne, A.J., Pfeffer, W.T., Stammer, D.,
- 717 Unnikrishnan, A.S., 2013. Sea Level Change, in: The Physical Science Basis: Contribution of
- 718 Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate
- 719 Change. pp. 1137–1216.
- 720 Colonese, A.C., Mannino, M.A., Mayer, D.B.-Y., Fa, D.A., Finlayson, J.C., Lubell, D., Stiner,
- 721 M.C., 2011. Marine mollusc exploitation in Mediterranean prehistory: an overview. Quaternary
- 722 international 239, 86–103.
- Cremaschi, M., 1990. Stratigraphy and palaeoenvironmental significance of the loess deposits
  on Susak Island (Dalmatian archipelago). Quaternary International 5, 97–106.
- 725 Cremonesi, G., Meluzzi, C., Pitti, C., Wilkens, B., 1984. Grotta Azzurra: scavi 1982 (nota
- 726 preliminare), in: Il mesolitico sul Carso Triestino, Quaderni della Societá per la Preistoria e
- 727 Protostoria della Regione Friuli-Venezia Giulia. Italo Svevo, Trieste, pp. 21–37.
- 728 Cristiani, E., Farbstein, R., Miracle, P., 2014. Ornamental traditions in the Eastern Adriatic: The
- 729 Upper Palaeolithic and Mesolithic personal adornments from Vela Spila (Croatia). Journal of
- 730 Anthropological Archaeology 36, 21–31. https://doi.org/10.1016/j.jaa.2014.06.009
- 731 Croatian Geoportal, 2019. Topografska karta u mjerilu 1:25000.
- 732 Cvetkoska, A., Levkov, Z., Reed, J.M., Wagner, B., 2014. Late Glacial to Holocene climate
- change and human impact in the Mediterranean: The last ca. 17ka diatom record of Lake Prespa
- 734 (Macedonia/Albania/Greece). Palaeogeography, Palaeoclimatology, Palaeoecology 406, 22–32.
- 735 https://doi.org/10.1016/j.palaeo.2014.04.010

736	Cvitkušić, B., Radović, S., Vujević, D., 2018. Changes in ornamental traditions and subsistence
737	strategies during the Palaeolithic-Mesolithic transition in Vlakno cave. Quaternary International,
738	Quaternary of Croatia 494, 180-192. https://doi.org/10.1016/j.quaint.2017.08.053
739	De Boer, B., Stocchi, P., Van De Wal, R., 2014. A fully coupled 3-D ice-sheet-sea-level model:
740	algorithm and applications. Geoscientific Model Development 7, 2141–2156.
741	Deino, A.L., Orsi, G., de Vita, S., Piochi, M., 2004. The age of the Neapolitan Yellow Tuff
742	caldera-forming eruption (Campi Flegrei caldera – Italy) assessed by 40Ar/39Ar dating method.
743	Journal of Volcanology and Geothermal Research, The Neapolitan Volcanoes: Vesuvius, Campi
744	Flegrei and Ischia 133, 157–170. https://doi.org/10.1016/S0377-0273(03)00396-2
745	Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L., Henderson,
746	G.M., Okuno, J., Yokoyama, Y., 2012. Ice-sheet collapse and sea-level rise at the Bølling warming
747	14,600 years ago. Nature 483, 559–564. https://doi.org/10.1038/nature10902
748	Dobson, J.E., 2014. Aquaterra Incognita: Lost Land Beneath The Sea. Geographical Review
749	104, 123–138. https://doi.org/10.1111/j.1931-0846.2014.12013.x
750	Domijan, N., Leder, N., Čupić, S., 2005. Visinski datumi Republike Hrvatske (Vertical datums
751	of the Republic of Croatia), in: Zbornik radova trećeg hrvatskog kongresa o katastru s
752	međunarodnim sudjelovanjem. Croatia.
753	Drźavni Hidrografski Institut, 1998. Jadransko More, charts MK17, MK-19, MK-20, MK-21,
754	MK-22, MK-23, MK-24, 535.
755	Fairbanks, R.G., 1989. A 17,000-year glacio-eustatic sea level record: influence of glacial
756	melting rates on the Younger Dryas event and deep-ocean circulation. Nature 342, 637-642.
757	https://doi.org/10.1038/342637a0
758	Faivre, S., Bakran-Petricioli, T., Horvatinčić, N., Sironić, A., 2013. Distinct phases of relative
759	sea level changes in the central Adriatic during the last 1500 years — influence of climatic

- variations? Palaeogeography, Palaeoclimatology, Palaeoecology 369, 163–174.
- 761 https://doi.org/10.1016/j.palaeo.2012.10.016
- Faivre, S., Butorac, V., 2018. Recently submerged tidal notches in the wider Makarska area
- 763 (Central Adriatic, Croatia). Quaternary International 494, 225–235.
- 764 https://doi.org/10.1016/j.quaint.2017.07.020
- Farbstein, R., Radić, D., Brajković, D., Miracle, P.T., 2012. First Epigravettian Ceramic
- Figurines from Europe (Vela Spila, Croatia). PLOS ONE 7, e41437.
- 767 https://doi.org/10.1371/journal.pone.0041437
- 768 Felja, I., Fontana, A., Furlani, S., Bajraktarević, Z., Paradžik, A., Topalović, E., Rossato, S.,
- 769 Ćosović, V., Juračić, M., 2015. Environmental changes in the lower Mirna River valley (Istria,
- 770 Croatia) during Upper Holocene. Geologia Croatica 68, 209–224.
- 771 https://doi.org/10.4154/GC.2015.16
- Foglini, F., Prampolini, M., Micallef, A., Angeletti, L., Vandelli, V., Deidun, A., Soldati, M.,
- 773 Taviani, M., 2016. Late Quaternary coastal landscape morphology and evolution of the Maltese
- 774 Islands (Mediterranean Sea) reconstructed from high-resolution seafloor data. Geological Society,
- 775 London, Special Publications 411, 77–95.
- Fontana, A., Vinci, G., Tasca, G., Mozzi, P., Vacchi, M., Bivi, G., Salvador, S., Rossato, S.,
- 777 Antonioli, F., Asioli, A., Bresolin, M., Di Mario, F., Hajdas, I., 2017. Lagoonal settlements and
- relative sea level during Bronze Age in Northern Adriatic: Geoarchaeological evidence and
- paleogeographic constraints. Quaternary International 439, 17–36.
- 780 https://doi.org/10.1016/j.quaint.2016.12.038
- Forenbaher, S., Kaiser, T., Miracle, P.T., 2013. Dating the East Adriatic Neolithic. European
- 782 Journal of Archaeology 16, 589–609. https://doi.org/10.1179/1461957113Y.0000000038

783	Forenbaher,	S., N	Miracle,	P.T.,	2006.	Pupićina	Cave and	l the s	pread of	of farming	g in	the ea	astern
											_		

- 784 Adriatic, in: Miracle, P., Forenbaher, S. (Eds.), Prehistoric Herders of Northern Istria: The
- 785 Archaeology of Pupićina Cave. Archaeological Museum of Istria, Pula, pp. 483–523.
- Galili, E., Benjamin, J., Eshed, V., Rosen, B., McCarthy, J., Horwitz, L.K., 2019. A submerged
- 787 7000-year-old village and seawall demonstrate earliest known coastal defence against sea-level rise.
- 788 PLoS ONE 14. https://doi.org/10.1371/journal.pone.0222560
- Garg, K.M., Chattopadhyay, B., Wilton, P.R., Malia Prawiradilaga, D., Rheindt, F.E., 2018.
- 790 Pleistocene land bridges act as semipermeable agents of avian gene flow in Wallacea. Molecular
- 791 Phylogenetics and Evolution 125, 196–203. https://doi.org/10.1016/j.ympev.2018.03.032
- Giglio, F., Romano, S., Albertazzi, S., Chiarini, F., Ravaioli, M., Ligi, M., Capotondi, L., 2020.
- 793 Sediment Dynamics of the Neretva Channel (Croatia Coast) Inferred by Chemical and Physical
- 794 Proxies. Applied Sciences 10, 807. https://doi.org/10.3390/app10030807
- Gkinis, V., Simonsen, S.B., Buchardt, S.L., White, J.W.C., Vinther, B.M., 2014. Water isotope
- 796 diffusion rates from the NorthGRIP ice core for the last 16,000 years Glaciological and
- paleoclimatic implications. Earth and Planetary Science Letters 405, 132–141.
- 798 https://doi.org/10.1016/j.epsl.2014.08.022
- Govorčin, D.P., Juračić, M., Horvatinčić, N., Onofri, V., 2001. Holocene sedimentation in the
- 800 Soline Channel (Mljet Lakes, Adriatic Sea). Natura Croatica 10, 247–258.
- 801 Hewitt, G., 2000. The genetic legacy of the Quaternary ice ages. Nature 405, 907–913.
- 802 https://doi.org/10.1038/35016000
- 803 HHI, 2019. Hrvatski Hidrografski Institut [WWW Document]. Hrvatski Hidrografski Institut.
- 804 URL http://www.hhi.hr/en/mareo (accessed 12.28.19).
- Janković, I., Ahern, J.C.M., Karavanić, I., Stockton, T., Smith, F.H., 2012. Epigravettian
- 806 Human Remains and Artifacts from Šandalja II, Istria, Croatia. PaleoAnthropology. Official
- 807 Publication of the Paleoanthropology Society. 2012, 87–122.

808	Juračić, M., Surić, M., 2010. Late Pleistocene-Holocene environmental changes-records from
809	submerged speleothems along the Eastern Adriatic coast (Croatia). Geologia Croatica 63, 155–169.
810	Kaczanowska, M., Kozłowski, J.K., 2018. The Mesolithic lithic industries of the eastern
811	Adriatic zone. Folia Quaternaria 191–215.
812	Karavanić, I., 2003. L'industrie aurignacienne de la grotte de Šandalja II (Istrie, Croatie) dans
813	le contexte de la région de l'Est de l'Adriatique. L'Anthropologie 107, 577-602.
814	https://doi.org/10.1016/j.anthro.2003.10.003
815	Kelletat, D.H., 2005. Dalmatian Coasts. Encyclopedia of coastal science, Encyclopedia of earth
816	sciences series.
817	Kendall, R.A., Mitrovica, J.X., Milne, G.A., Törnqvist, T.E., Li, Y., 2008. The sea-level
818	fingerprint of the 8.2 ka climate event. Geology 36, 423–426. https://doi.org/10.1130/G24550A.1
819	Komšo, D., 2008. Mezolitik u Hrvatskoj. Opusc. Archaeol. Rad. Arheol. zavoda 30, 55-92.
820	Koncani Uhač, I., Čuka, I., 2015. M., 2015–Doprinos poznavanju podmorskog eneolitičkog
821	nalazišta u uvali Zambratija, Histria archaeologica 46. Pula 25–73.
822	Kozlowski, J.K., Kaczanowska, M., 2004. Gravettian/Epigravettian sequences in the Balkans
823	and Anatolia. Mediterranean Archaeology and Archaeometry 4, 5–18.
824	Lambeck, K., Purcell, A., 2005. Sea-level change in the Mediterranean Sea since the LGM:
825	model predictions for tectonically stable areas. Quaternary Science Reviews 24, 1969–1988.
826	https://doi.org/10.1016/j.quascirev.2004.06.025
827	Lambeck, K., Purcell, A., Johnston, P., Nakada, M., Yokoyama, Y., 2003. Water-load definition
828	in the glacio-hydro-isostatic sea-level equation. Quaternary Science Reviews 22, 309-318.
829	https://doi.org/10.1016/S0277-3791(02)00142-7
830	Lightfoot, E., Boneva, B., Miracle, P.T., Šlaus, M., O'Connell, T.C., 2011. Exploring the
831	Mesolithic and Neolithic transition in Croatia through isotopic investigations. Antiquity 85, 73–86.
832	https://doi.org/10.1017/S0003598X00067442

- 833 Mauch Lenardić, J., Oros Sršen, A., Radović, S., 2018. Quaternary fauna of the Eastern
- 834 Adriatic (Croatia) with the special review on the Late Pleistocene sites. Quaternary International
- 835 494, 130–151. https://doi.org/10.1016/j.quaint.2017.11.028
- 836 McLaughlin, T.R., Stoddart, S., Malone, C., 2018. Island risks and the resilience of a
- 837 prehistoric civilization. World Archaeology 50, 570–600.
- 838 https://doi.org/10.1080/00438243.2018.1515656
- 839 Melini, D., Spada, G., 2019. Some remarks on Glacial Isostatic Adjustment modelling
- 840 uncertainties. Geophys J Int 218, 401–413. https://doi.org/10.1093/gji/ggz158
- 841 Miracle, P., 2007. The Late Glacial" Great Adriatic Plain":" Garden of Eden" or "No Man's
- 842 Land" during the Epipalaeolithic? A view from Istria (Croatia), in: Late Paleolithic Environments
- and Cultural Relations around the Adriatic, BAR International Series. Archaeopress, pp. 41–51.
- 844 Miracle, P., 2002. Mesolithic meals from Mesolithic middens, in: Miracle, P., Milner, N. (Eds.),
- 845 Consuming Passions and Patterns of Consumption. McDonald Inst of Archeological, pp. 65–88.
- 846 Miracle, P., 2001. Feast or famine? Epipalaeolithic subsistence in the northern Adriatic basin. 1
- 847 28, 177–197. https://doi.org/10.4312/dp.28.10
- 848 Miracle, P., Galanidou, N., Forenbaher, S., 2000. Pioneers in the hills: Early Mesolithic
- foragers at Šebrn Abri (Istria, Croatia). European Journal of Archaeology 3, 293–329.
- 850 https://doi.org/10.1179/146195700807860837
- 851 Miracle, P.T., Forenbaher, S., 2005. Neolithic and Bronze-Age Herders of Pupićina Cave,
- 852 Croatia. Journal of Field Archaeology 30, 255–281.
- 853 Mussi, M., Palombo, M.R., 2001. Les carnivores du Pliocène supérieur et du Pléistocène de la
- péninsule italienne: quelles implications pour le peuplement humain. Brugal, J. Ph., Fosse, P.(eds),
- 855 Actes du XIVe Congrès UISPP «Hommes & Carnivores au Paléolithique», Liège 597–609.
- 856 Mussi, M., Peresani, M., 2011. Human settlement of Italy during the Younger Dryas.
- 857 Quaternary international 242, 360–370.

- NASA, 2000. Shuttle Radar Topography Mission (SRTM) 1 Arc-Second Global Elevation
  Data.
- 860 Neumann, B., Vafeidis, A.T., Zimmermann, J., Nicholls, R.J., 2015. Future Coastal Population
- 861 Growth and Exposure to Sea-Level Rise and Coastal Flooding A Global Assessment. PLOS ONE
- 862 10, e0118571. https://doi.org/10.1371/journal.pone.0118571
- 863 Nunn, P.D., 2020. In Anticipation of ExtirpationHow Ancient Peoples Rationalized and
- Responded to Postglacial Sea Level Rise. Environmental Humanities 12, 113–131.
- 865 https://doi.org/10.1215/22011919-8142231
- 866 Olson, C.J., Becker, J.J., Sandwell, D.T., 2016. SRTM15\_PLUS: Data Fusion of Shuttle Radar
- 867 Topography Mission (SRTM) Land Topography With Measured and Estimated Seafloor
- 868 Topography (NCEI Accession 0150537).
- 869 Olson, C.J., Becker, J.J., Sandwell, D.T., 2014. A new global bathymetry map at 15 arcsecond
- 870 resolution for resolving seafloor fabric: SRTM15\_PLUS. AGU Fall Meeting Abstracts 2014,
- 871 OS34A-03.
- Palombo, M.R., Antonioli, F., Lo Presti, V., Mannino, M.A., Melis, R.T., Orru, P., Stocchi, P.,
- Talamo, S., Quarta, G., Calcagnile, L., Deiana, G., Altamura, S., 2017. The late Pleistocene to
- 874 Holocene palaeogeographic evolution of the Porto Conte area: Clues for a better understanding of
- human colonization of Sardinia and faunal dynamics during the last 30 ka. Quaternary International
- 876 439, 117–140. https://doi.org/10.1016/j.quaint.2016.06.014
- Peltier, W.R., 2004. Global Glacial Isostasy and the Surface of the Ice-Age Earth: The Ice-5g
- 878 (vm2) Model and Grace. Annual Review of Earth and Planetary Sciences 32, 111–149.
- 879 https://doi.org/10.1146/annurev.earth.32.082503.144359
- 880 Peltier, W.R., Argus, D.F., Drummond, R., 2015. Space geodesy constrains ice age terminal
- deglaciation: The global ICE-6G\_C (VM5a) model. Journal of Geophysical Research: Solid Earth
- 882 120, 450–487. https://doi.org/10.1002/2014JB011176

- 883 Peltier, W.R., Fairbanks, R.G., 2006. Global glacial ice volume and Last Glacial Maximum
- 884 duration from an extended Barbados sea level record. Quaternary Science Reviews, Critical
- 885 Quaternary Stratigraphy 25, 3322–3337. https://doi.org/10.1016/j.quascirev.2006.04.010
- Pikelj, K., Juračić, M., 2013. Eastern Adriatic Coast (EAC): Geomorphology and Coastal
- 887 Vulnerability of a Karstic Coast. Journal of Coastal Research 289, 944–957.
- 888 https://doi.org/10.2112/JCOASTRES-D-12-00136.1
- 889 Pirazzoli, P.A., 2013. Geomorphological Indicators, in: Encyclopedia of Quaternary Science.
- 890 Elsevier, pp. 377–384. https://doi.org/10.1016/B978-0-444-53643-3.00133-3
- 891 Pross, J., Kotthoff, U., Müller, U.C., Peyron, O., Dormoy, I., Schmiedl, G., Kalaitzidis, S.,
- 892 Smith, A.M., 2009. Massive perturbation in terrestrial ecosystems of the Eastern Mediterranean
- region associated with the 8.2 kyr B.P. climatic event. Geology 37, 887–890.
- 894 https://doi.org/10.1130/G25739A.1
- Radić, D. (Ed.), 2018. Island of Korčula: From the earliest times to the end of prehistory.
- 896 Centar za kulturu "Vela Luka," Vela Luka, Croatia.
- 897 Radić, D., Lugović, B., Marjanac, L., 2007. Neapolitan Yellow Tuff (NYT) from the leistocene
- sediments in Vela Spila on the island of Korčula: a valuable chronostratigraphic marker of the
- transition from the Palaeolithic to the Mesolithic. Opvscvla archaeologica 31, 20.
- 900 Radić Rossi, I., 2008. Arheološka baština u podmorju Kaštelanskog zaljeva. Archaeologia
  901 Adriatica 2, 489–506.
- 902 Rainsford, C., O'Connor, T., Miracle, P., 2014. Fishing in the Adriatic at the Mesolithic–
- 903 Neolithic transition: Evidence from Vela Spila, Croatia. Environmental Archaeology 19, 311–320.
- 904 https://doi.org/10.1179/1749631414Y.000000018
- 905 Richards, M.P., Karavanić, I., Pettitt, P., Miracle, P., 2015. Isotope and faunal evidence for high
- 906 levels of freshwater fish consumption by Late Glacial humans at the Late Upper Palaeolithic site of

- 907 Šandalja II, Istria, Croatia. Journal of Archaeological Science 61, 204–212.
- 908 https://doi.org/10.1016/j.jas.2015.06.008
- 909 Rizner, M., Vukosavljević, N., Miracle, P., 2009. The paleoecological and paleodietary
- 910 significance of edible land snails (Helix sp.) across the Pleistocene-Holocene transition on the
- 911 eastern Adriatic coast, in: MESO 2005: The 7th International Conference on the Mesolithic in

912 Europe.

- 913 Rossi, I.R., 2017. Proučavanje promjena morske razine na arheološkim nalazištima u priobalju.
- 914 Vjesnik za Arheologiju i Povijest Dalmatinsku 110.
- 915 Rovere, A., Stocchi, P., Vacchi, M., 2016. Eustatic and Relative Sea Level Changes. Curr Clim
- 916 Change Rep 2, 221–231. https://doi.org/10.1007/s40641-016-0045-7
- Sandwell, D.T., Becker, J.J., Smith, W.H.F., 2014a. SRTM15\_PLUS Estimated Topography, 15
  seconds, Global, v1.
- 919 Sandwell, D.T., Müller, R.D., Smith, W.H.F., Garcia, E., Francis, R., 2014b. New global marine
- gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure. Science 346, 65–67.
- 921 https://doi.org/10.1126/science.1258213
- 922 Shaw, T.A., Plater, A.J., Kirby, J.R., Roy, K., Holgate, S., Tutman, P., Cahill, N., Horton, B.P.,
- 923 2018. Tectonic influences on late Holocene relative sea levels from the central-eastern Adriatic
- 924 coast of Croatia. Quaternary Science Reviews 200, 262–275.
- 925 https://doi.org/10.1016/j.quascirev.2018.09.015
- 926 Shennan, I., Long, A.J., Horton, B.P. (Eds.), 2015. Handbook of Sea-Level Research. John
- 927 Wiley & Sons, Ltd, Chichester, UK.
- 928 Sikora, M., Mihanović, H., Vilibić, I., 2014. Paleo-coastline of the Central Eastern Adriatic Sea,
- 929 and Paleo-Channels of the Cetina and Neretva rivers during the last glacial maximum. Acta
- 930 Adriatica 55, 3–18.

- 931 Smith, W.H.F., Sandwell, D.T., 1997. Global Sea Floor Topography from Satellite Altimetry
- 932 and Ship Depth Soundings. Science 277, 1956–1962. https://doi.org/10.1126/science.277.5334.1956
- 933 Šošić Klindžić, R., Radović, S., Težak Gregl, T., Šlaus, M., Perhoć, Z., Altherr, R., Hulina, M.,
- 934 Gerometta, K., Boschian, G., Vukosavljević, N., Ahern, J.C.M., Janković, I., Richards, M.,
- 935 Karavanić, I., 2015. Late Upper Paleolithic, Early Mesolithic and Early Neolithic from the cave site
- 936 Zemunica near Bisko (Dalmatia, Croatia) 12, 3–46.
- 937 Spada, G., Galassi, G., 2017. Extent and dynamic evolution of the lost land aquaterra since the
- 938 Last Glacial Maximum. Comptes Rendus Geoscience 349, 151–158.
- 939 Spada, G., Stocchi, P., 2007. SELEN: A Fortran 90 program for solving the "sea-level
- 940 equation." Computers & Geosciences 33, 538–562. https://doi.org/10.1016/j.cageo.2006.08.006
- 941 Srdoč, D., Obelić, B., Horvatinčić, N., Bronić, I.K., Sliepčević, A., 1989. Rudjer Bošković
- 942 Institute radiocarbon measurements XI. Radiocarbon 31, 85–98.
- 943 Stewart, J.R., Van Kolfschoten, M., Markova, A., Musil, R., 2003. The mammalian faunas of
- Europe during oxygen isotope stage three. Neanderthals and modern humans in the European
- 945 landscape during the last glaciation 60, 103–130.
- 946 Taviani, M., Angeletti, L., Campiani, E., Ceregato, A., Foglini, F., Maselli, V., Morsilli, M.,
- 947 Parise, M., Trincardi, F., 2012. Drowned karst landscape offshore the Apulian margin (Southern
- Adriatic Sea, Italy). Journal of Cave and Karst Studies 74, 197–212.
- 949 https://doi.org/10.4311/2011JCKS0204
- tide-forecast.com, 2019. Tide Times and Tide Charts Worldwide [WWW Document].
- 951 Meteo365.com. URL https://www.tide-forecast.com (accessed 7.27.19).
- Tozer, B., Sandwell, D.T., Smith, W.H.F., Olson, C., Beale, J.R., Wessel, P., 2019. Global
- 953 Bathymetry and Topography at 15 Arc Sec: SRTM15+. Earth and Space Science 6, 1847–1864.
- 954 https://doi.org/10.1029/2019EA000658

955	Trincardi, F., Campiani, E., Correggiari, A., Foglini, F., Maselli, V., Remia, A., 2014.
956	Bathymetry of the Adriatic Sea: The legacy of the last eustatic cycle and the impact of modern
957	sediment dispersal. Journal of Maps 10, 151-158. https://doi.org/10.1080/17445647.2013.864844
958	Vacchi, M., Marriner, N., Morhange, C., Spada, G., Fontana, A., Rovere, A., 2016. Multiproxy
959	assessment of Holocene relative sea-level changes in the western Mediterranean: Sea-level
960	variability and improvements in the definition of the isostatic signal. Earth-Science Reviews 155,
961	172-197. https://doi.org/10.1016/j.earscirev.2016.02.002
962	Vlahović, I., Tišljar, J., Velić, I., Matičec, D., 2005. Evolution of the Adriatic Carbonate
963	Platform: Palaeogeography, main events and depositional dynamics. Palaeogeography,
964	Palaeoclimatology, Palaeoecology 220, 333-360. https://doi.org/10.1016/j.palaeo.2005.01.011
965	Vlahović, I., Tišljar, J., Velić, I., Matičec, D., 2002. The Karst Dinarides are Composed of
966	Relics of a Single Mesozoic Platform: Facts and Consequences. Geologia Croatica 55, 171–183.
967	Vogel, H., Wagner, B., Zanchetta, G., Sulpizio, R., Rosén, P., 2010. A paleoclimate record with
968	tephrochronological age control for the last glacial-interglacial cycle from Lake Ohrid, Albania and
969	Macedonia. J Paleolimnol 44, 295–310. https://doi.org/10.1007/s10933-009-9404-x
970	Vukosavljević, N., Perhoč, Z., 2017. Lithic raw material procurement of the Late Epigravettian
971	hunter-gatherers from Kopačina Cave (island of Brač, Dalmatia, Croatia). Quaternary International,
972	Prehistoric hunter-gatherers and farmers in the Adriatic and neighboring regions 450, 164–185.
973	https://doi.org/10.1016/j.quaint.2016.09.017
974	Vukosavljević, N., Perhoč, Z., Altherr, R., 2014. Pleistocene-Holocene transition in the Vlakno
975	Cave on the island of Dugi otok (Dalmatia, Croatia)-lithic perspective. Prilozi Instituta za
976	arheologiju u Zagrebu 31, 5–72.
977	Vukosavljević, N., Perhoč, Z., Čečuk, B., Karavanić, I., 2011. Late Glacial knapped stone
978	industry of Kopačina Cave. Vjesnik za arheologiju i povijest dalmatinsku 104, 7–54.

979	Wacha, L., Frechen, M., Crnjaković, M., 2011a. The loess chronology of the Island of Susak,
980	Croatia. Quaternary Science International 60, 153–169.
981	Wacha, L., Pavlaković, S.M., Novothny, Á., Crnjaković, M., Frechen, M., 2011b.
982	Luminescence dating of Upper Pleistocene loess from the Island of Susak in Croatia. Quaternary
983	International 234, 50–61.
984	Weber, MJ., Grimm, S.B., Baales, M., 2011. Between warm and cold: Impact of the Younger
985	Dryas on human behavior in Central Europe. Quaternary International 242, 277-301.
986	Weigelt, P., Steinbauer, M.J., Cabral, J.S., Kreft, H., 2016. Late Quaternary climate change
987	shapes island biodiversity. Nature 532, 99.
988	Whallon, R., 1999. The lithic tool assemblages at Badanj within their regional context, in:
989	Bailey, G.N., Adam, E., Panagopoulou, E., Perles, C., Zachos, K. (Eds.), The Palaeolithic
990	Archaeology of Greece and Adjacent Areas, Monograph of the British School at Athens. Presented
991	at the Proceedings of the ICOPAG Conference, British School at Athens, Athens.
992	Wunsam, S., Schmidt, R., Müller, J., 1999. Holocene lake development of two Dalmatian
993	lagoons (Malo and Veliko Jezero, Isle of Mljet) in respect to changes in Adriatic sea level and
994	climate. Palaeogeography, Palaeoclimatology, Palaeoecology 146, 251–281.
995	Zvelebil, M., Lillie, M.C., 2000. Transition to agriculture in Eastern Europe, in: Price, T.D.
996	(Ed.), Europe's First Farmers. Cambridge University Press, New York.
997	
,,,	

# 999 Figures captions

Fig 1. Map of the Adriatic Sea and study area. A: general map of the Adriatic Sea, divided into
three sub-basins, and surrounding areas. B: Korčula and surrounding Islands. Tectonic information:
Korbar, 2009; Surić et al., 2014. Map and relief data: GEBCO and GADM.

Fig 2. Vela Spila and surroundings. Top: View from the cave towards the SW, showing Vela Luka town and harbour (photo from P.T. Miracle). Middle left: View of Vela Luka bay looking northeast. Hvar and the mainland in background. Vela Spila is in the centre hill of the image (photo from G. Boschian) Middle right: Another view from Vela Spila, but looking west out Vela Luka bay. The innermost harbour is visible (photo from F. Rinaldi). Bottom: Interior photomosaic of Vela Spila. The most recently excavated trenches are on the left (photo from G. Boschian).

Fig 3. Plan of Vela Spila cave, with locations of archaeological excavations. 1: dripline of entrance and of windows in ceiling; 2: excavations pre-2000; 3: excavations 2000-2018; A-A' location of the profile in fig. 4.

Fig 4. General stratigraphy of the Vela Spila archaeological deposit. From left to right: cultural horizons and informal Quaternary stratigraphy, photo and main lithostratigraphic subdivisions of vertical profile (A-A' in fig. 3). NYT: Neapolitan Yellow Tuff; red lines: erosive boundaries. Note the wavy limits of NYT and overlying units, due to frost action in sediment.

1016 Fig 5. Summary of calibrated dates from Vela Spila. Note the absence of archaeological 1017 material for dating during the abandonment phase.

Fig 6. Sea level reconstructions for Vela Luka. Sea-level realizations for the Vela Luka area of Korčula. Dashed line: I6G\_S4 realization of the ICE-6G\_C (VM5a) (Peltier et al., 2015). Dotted line: ANU\_S4 realization of Lambeck/ANU (Lambeck et al., 2003; Lambeck and Purcell, 2005). Sea-level indicators from Faivre et al. (2013), Juračić and Surić (2010), and Mljet lakes (Govorčin et al., 2001; Wunsam et al., 1999) shown in green. Archaeological periods for the site reconstructed with 14C dates from Table 2. Exact period delineations are an abstraction, in reality date uncertainties and gradual cultural shifts mean the transitions were nuanced.

1025 Fig 7. Palaeogeography and sea-level change at Vela Spila. Palaeogeographic reconstruction of 1026 the area around Korčula at different phases of Vela Spila's settlement history. A) Start of the Palaeolithic ca. 19.7 cal ka BP. B) Start of the abandonment phase ca. 14.0 cal ka BP. C) Start of the 1027 1028 Mesolithic ca. 9.4 cal ka BP. Vela Spila location shown by red dot. Light blue shows ICE6G 1029 predicted RSL; dark blue shows Lambeck/ANU predicted RSL. Dotted blue line indicates probable 1030 location of palaeo-Neretva river according to Sikora et al.(2014). For each time-slice, a red line or 1031 lines indicates distance to sea from Vela Spila. The crosshatched circle indicates the percent of land 1032 within a 20 km radius from the cave that is inundated. Map elevations interpolated from datasets 1033 listed in Table 1.

Fig 8. Slope vs. depth in Western Korčula. Profile of 15 km transects AB and CD from Fig 5C, showing the higher slopes that predominate above the 70 m isobath (note, Y axis in left panels is exaggerated). Right panel shows slope in degrees vs. depth below sea level along transect CD. Slopes increase above -70 m. This morphology results in negligible horizontal coastline change after ca. 12 ka BP.

Fig 9. A zoomed in view of the palaeo shorelines at the start of the Mesolithic (~9.4 ka BP) at Vela Spila, overlaid on modern orthophotos (Croatian Geoportal, 2019). ANU and ICE models for sea-level are shown. The situation of the site on a hill at ~121 m ASL above the harbour valley is evident. To the south and southeast, where the town of Vela Luka is now situated the western end of the flat karstic depression known as Blatsko Polje can be seen.

1044



≛







m a.s.l. 113.00



OxCal v4.3.2 Bronk Ramsey (2017); r:5 IntCal13 atmospheric curve (Reimer et al 2013)



Figure 6

RSL (m) vs Year BP











## **Declaration of interests**

 $\boxtimes$  The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

## **Declaration of interests**

 $\boxtimes$  The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: