

Additional File 2

Exosomal miR-6126 as a novel therapeutic target for overcoming resistance of anti-cancer effect in hepatocellular carcinoma

Hyemin Hwang^{1,#}, Jimin Kim^{1,#}, Tae-Hun Kim², Yeonju Han¹, Dayoung Choi¹, Sua Cho¹, Seunghwan Kim¹, Sanghee Park¹, Taehyun Park¹, Filippo Piccinini^{3,4}, Won Jong Rhee⁵, Jae-Chul Pyun², Misu Lee^{1,6,*}

¹Division of Life Sciences, College of Life Science and Bioengineering, Incheon National University, Incheon, 22012, Republic of Korea

²Department of Materials Science and Engineering, Yonsei University, 50 Yonsei-Ro, Seodaemun-Gu, Seoul, 03722, Republic of Korea.

³IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) “Dino Amadori”, Meldola, Italy

⁴Department of Medical and Surgical Sciences (DIMEC), University of Bologna, Bologna, Italy

⁵Department of Bioengineering and Nano-Bioengineering, Incheon National University Incheon 22012, Republic of Korea.

⁶Institute for New Drug Development, College of Life Science and Bioengineering, Incheon National University, Incheon, 22012, Republic of Korea

These authors contributed equally to this study.

Corresponding author: Misu Lee, PhD

Incheon National University, Incheon, Republic of Korea

E-mail: misulee@inu.ac.kr; Tel.: +82 32 835 8091; Fax: + 82 32 835 0754




Figure S1. Whole blot images of Fig 1B. The pERK, ERK, and β -actin images are from the same blot, while CD44 images are from the upper blot of the pERK.




Figure S2. Whole blot images of Fig 1D.




Figure S3. Whole blot images of Fig 4D.




Figure S4. Whole blot images of Fig 5D.




Figure S5. Whole blot images of Fig 6C. The HK2 and CD44 images are from the same blot, while the β-actin images are from an same lower blot of the CD44. The pERK and ERK images are from the same blot.