

Supplementary Table S3. Consistency estimates of each individual statement after three rounds consensus for recommendations on the management of acute type B penetrating aortic ulcer (PAU) and intramural haematoma (IMH).

Statement	Cohen's K	P	ICC (95%CI)	P	Overall Consistency
1	0.72	.001	0.89 (0.78-0.95)	<.001	I
2	0.78	<.001	0.89 (0.79-0.95)	<.001	I
3	0.81	<.001	0.75 (0.71-0.95)	<.001	I
4	0.68	.004	0.82 (0.69-0.91)	.001	II
5	0.45	.020	0.60 (0.35-0.80)	.045	III
6	0.70	.002	0.84 (0.72-0.92)	.001	II
7	0.50	<.001	0.64 (0.59-0.69)	<.001	I
8	0.28	.003	0.26 (0.20-0.31)	<.001	II
9	0.37	<.001	0.55 (0.50-0.60)	<.001	I
10	0.31	<.001	0.39 (0.34-0.45)	<.001	I
11	0.52	<.001	0.59 (0.54-0.64)	<.001	I
12	0.43	<.001	0.57 (0.52-0.62)	<.001	I
13	0.31	<.001	0.40 (0.34-0.44)	<.001	I
14	0.38	<.001	0.51 (0.46-0.56)	<.001	I
15	0.34	<.001	0.54 (0.48-0.60)	<.001	I
16	0.28	<.001	0.49 (0.38-0.58)	<.001	I
17	0.38	<.001	0.40 (0.34-0.44)	<.001	I
18	0.38	<.001	0.58 (0.52-0.62)	<.001	I
19	0.30	<.001	0.38 (0.34-0.44)	<.001	I
20	0.42	<.001	0.56 (0.50-0.60)	<.001	I
21	0.32	<.001	0.40 (0.35-0.45)	<.001	I
22	0.59	<.001	0.75 (0.69-0.79)	<.001	I
23	0.45	<.001	0.68 (0.63-0.73)	<.001	I
24	0.34	.001	0.37 (0.32-0.42)	<.001	II
25	0.44	<.001	0.56 (0.51-0.61)	<.001	I
26	0.41	<.001	0.50 (0.45-0.55)	<.001	I
27	0.43	.001	0.36 (0.31-0.41)	<.001	II
28	0.37	<.001	0.61 (0.56-0.66)	<.001	I
29	0.32	<.001	0.43 (0.37-0.47)	<.001	I
30	0.48	.022	0.58 (0.31-0.79)	.048	III