

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

All-sky search for short gravitational-wave bursts in the first part of the fourth LIGO-Virgo-KAGRA observing run

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Abac, A. g., Abouelfettouh, I., Acernese, F., Ackley, K., Adamcewicz, C., Adhicary, S., et al. (2025). All-sky search for short gravitational-wave bursts in the first part of the fourth LIGO-Virgo-KAGRA observing run. PHYSICAL REVIEW D, 112(10), 1-29 [10.1103/wjdz-jdby].

Availability:

This version is available at: <https://hdl.handle.net/11585/1036980> since: 2026-01-14

Published:

DOI: <http://doi.org/10.1103/wjdz-jdby>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

All-sky search for short gravitational-wave bursts in the first part of the fourth LIGO-Virgo-KAGRA observing run

A. G. Abac¹, I. Abouelfettouh², F. Acernese^{3,4}, K. Ackley⁵, C. Adamcewicz⁶, S. Adhicary⁷, D. Adhikari^{8,9}, N. Adhikari¹⁰, R. X. Adhikari¹¹, V. K. Adkins¹², S. Afroz¹³, A. Agapito¹⁴, D. Agarwal¹⁵, M. Agathos¹⁶, N. Aggarwal¹⁷, S. Aggarwal¹⁸, O. D. Aguiar¹⁹, I.-L. Ahrend²⁰, L. Aiello^{21,22}, A. Ain²³, P. Ajith²⁴, T. Akutsu^{25,26}, S. Albanesi^{27,28}, W. Ali^{29,30}, S. Al-Kersh, S. Al-Kersh^{8,9}, C. Allén³¹, A. Allocca^{32,4}, S. Al-Shammari³³, P. A. Altin³⁴, S. Alvarez-Lopez³⁵, W. Amar³¹, O. Amarasinghe³³, A. Amato^{36,37}, F. Amicucci^{38,39}, C. Amra⁴⁰, A. Ananyeva¹¹, S. B. Anderson¹¹, W. G. Anderson¹¹, M. Andia⁴¹, M. Ando⁴², M. Andrés-Carcasona⁴³, T. Andrić^{44,45,8,9}, J. Anglin⁴⁶, S. Ansoldi^{47,48}, J. M. Antelis⁴⁹, S. Antier⁴¹, M. Aoumi⁵⁰, E. Z. Appavuravther^{51,52}, S. Appert¹¹, S. K. Apple⁵³, K. Arai¹¹, A. Araya⁴², M. C. Araya¹¹, M. Arca Sedda^{44,45}, J. S. Areeda⁵⁴, N. Aritomi², F. Armato^{29,30}, S. Armstrong⁵⁵, N. Arnaud⁵⁶, M. Arogeti⁵⁷, S. M. Aronson¹², G. Ashton⁵⁸, Y. Aso^{25,59}, L. Asprea²⁸, M. Assiduo^{60,61}, S. Assis de Souza Melo⁶², S. M. Aston⁶³, P. Astone³⁸, F. Attadio^{39,38}, F. Aubin⁶⁴, K. AultONeal⁶⁵, G. Avallone⁶⁶, E. A. Avila⁴⁹, S. Babak²⁰, C. Badger⁶⁷, S. Bae⁶⁸, S. Bagnasco²⁸, L. Baiotti⁶⁹, R. Bajpai⁷⁰, T. Baka^{71,37}, A. M. Baker⁶, K. A. Baker⁷², G. Baldi^{73,74}, N. Baldicchi^{75,51}, M. Ball⁷⁶, G. Ballardín⁶², S. W. Ballmer⁷⁷, S. Banagiri⁶, B. Banerjee⁴⁴, D. Bankar⁷⁸, T. M. Baptiste¹², P. Baral¹⁰, M. Baratti^{79,80}, J. C. Barayoga¹¹, B. C. Barish¹¹, D. Barker², N. Barman⁷⁸, P. Barneo^{81,82,83}, F. Barone^{84,4}, B. Barr⁸⁵, L. Barsotti³⁵, M. Barsuglia²⁰, D. Barta⁸⁶, A. M. Bartoletti⁸⁷, M. A. Barton⁸⁵, I. Bartos⁴⁶, A. Basalaev^{8,9}, R. Bassiri⁸⁸, A. Basti^{80,79}, M. Bawaj^{75,51}, P. Baxi⁸⁹, J. C. Bayley⁸⁵, A. C. Baylor¹⁰, P. A. Baynard II⁵⁷, M. Bazzan^{90,91}, V. M. Bedakihale⁹², F. Beirnaert⁹³, M. Bejger⁹⁴, D. Belardinelli²², A. S. Bell⁸⁵, D. S. Bellie⁹⁵, L. Bellizzi^{79,80}, W. Benoit¹⁸, I. Bentara⁵⁶, J. D. Bentley⁹⁶, M. Ben Yaala⁵⁵, S. Bera^{97,98}, F. Bergamin³³, B. K. Berger⁸⁸, S. Bernuzzi²⁷, M. Beroiz¹¹, D. Bersanetti²⁹, T. Bertheas⁹⁹, A. Bertolini^{37,36}, J. Betzwieser⁶³, D. Beveridge⁷², G. Bevilacqua¹⁰⁰, N. Bevis¹⁰¹, R. Bhandare¹⁰², R. Bhatt¹¹, D. Bhattacharjee^{103,104}, S. Bhattacharyya¹⁰⁵, S. Bhaumik⁴⁶, V. Biancalana¹⁰⁰, A. Bianchi^{37,106}, I. A. Bilenko¹⁰⁷, G. Billingsley¹¹, A. Binetti¹⁰⁸, S. Bini^{11,73,74}, C. Binu¹⁰⁹, S. Biot¹¹⁰, O. Birnholtz¹¹¹, S. Biscoveanu⁹⁵, A. Bisht⁹, M. Bitossi^{62,79}, M.-A. Bizouard¹¹², S. Blaber¹¹³, J. K. Blackburn¹¹, L. A. Blagg⁷⁶, C. D. Blair^{72,63}, D. G. Blair⁷², N. Bode^{8,9}, N. Boettner⁹⁶, G. Boileau¹¹², M. Boldrini³⁸, G. N. Bolingbroke¹¹⁴, A. Bolliand^{115,40}, L. D. Bonavena⁴⁶, R. Bondarescu⁸¹, F. Bondu¹¹⁶, E. Bonilla⁸⁸, M. S. Bonilla⁵⁴, A. Bonino¹¹⁷, R. Bonnand^{31,115}, A. Borchers^{8,9}, S. Borhanian⁷, V. Boschi⁷⁹, S. Bose¹¹⁸, V. Bossilkov⁶³, Y. Bothra^{37,106}, A. Boudon⁵⁶, L. Bourg⁵⁷, G. Bouyer¹¹⁹, M. Boyle¹²⁰, A. Bozzi⁶², C. Bradaschia⁷⁹, P. R. Brady¹⁰, A. Branch⁶³, M. Branchesi^{44,45}, I. Braun¹⁰³, T. Briant¹²¹, A. Brillet¹¹², M. Brinkmann^{8,9}, P. Brockill¹⁰, E. Brockmueller^{8,9}, A. F. Brooks¹¹, B. C. Brown⁴⁶, D. D. Brown¹¹⁴, M. L. Brozzetti^{75,51}, S. Brunett¹¹, G. Bruno¹⁵, R. Bruntz¹²², J. Bryant¹¹⁷, Y. Bu¹²³, F. Bucci⁶¹, J. Buchanan¹²², O. Bulashenko^{81,82}, T. Bulik¹²⁴, H. J. Bulten³⁷, A. Buonanno^{125,1}, K. Burtzyk², R. Buscicchio^{126,127}, D. Buskulic³¹, C. Buy⁹⁹, R. L. Byer⁸⁸, G. S. Cabourn Davies¹²⁸, R. Cabrita¹⁵, V. Cáceres-Barbosa⁷, L. Cadonati⁵⁷, G. Cagnoli¹²⁹, C. Cahillane⁷⁷, A. Calafat⁹⁷, T. A. Callister¹³⁰, E. Calloni^{32,4}, S. R. Callos⁷⁶, G. Caneva Santoro⁴³, K. C. Cannon⁴², H. Cao³⁵, L. A. Capistran¹³¹, E. Capocasa²⁰, E. Capote^{2,11}, G. Capurri^{80,79}, G. Carapella^{66,132}, F. Carbognani⁶², M. Carlassara^{8,9}, J. B. Carlin¹²³, T. K. Carlson¹³³, M. F. Carney¹⁰³, M. Carpinelli^{126,62}, G. Carrillo⁷⁶, J. J. Carter^{8,9}, G. Carullo^{117,134}, A. Casallas-Lagos¹³⁵, J. Casanueva Diaz⁶², C. Casentini^{136,22}, S. Y. Castro-Lucas¹³⁷, S. Caudill¹³³, M. Cavaglià¹⁰⁴, R. Cavalieri⁶², A. Ceja⁵⁴, G. Cella⁷⁹, P. Cerdá-Durán^{138,139}, E. Cesarini²², N. Chabbra³⁴, W. Chaibi¹¹², A. Chakraborty¹³, P. Chakraborty^{8,9}, S. Chakraborty¹⁰², S. Chalathadka Subrahmanya⁹⁶, J. C. L. Chan¹⁴⁰, M. Chan¹¹³, K. Chang¹⁴¹, S. Chao^{142,141}, P. Charlton¹⁴³, E. Chassande-Mottin²⁰, C. Chatterjee¹⁴⁴, Debarati Chatterjee⁷⁸, Deep Chatterjee³⁵, M. Chaturvedi¹⁰², S. Chaty²⁰, K. Chatzioannou¹¹, A. Chen¹⁴⁵, A. H.-Y. Chen¹⁴⁶, D. Chen¹⁴⁷, H. Chen¹⁴², H. Y. Chen¹¹⁹, S. Chen¹⁴⁴, Yanbei Chen¹⁴⁸, Yitian Chen¹²⁰, H. P. Cheng¹⁴⁹, P. Chessa^{75,51}, H. T. Cheung⁸⁹, S. Y. Cheung⁶, F. Chiadini^{150,132}, G. Chiarini^{8,9,91}, A. Chiba¹⁵¹, A. Chincarini²⁹, M. L. Chiofalo^{80,79}, A. Chiummo^{4,62}, C. Chou¹⁴⁶, S. Choudhary⁷², N. Christensen^{112,152}, S. S. Y. Chua³⁴, G. Ciani^{73,74}, P. Cieciela⁹⁴, M. Cieřlar¹²⁴, M. Cifaldi²², B. Cirok¹⁵³, F. Clara², J. A. Clark^{11,57}, T. A. Clarke⁶, P. Clearwater¹⁵⁴, S. Clesse¹¹⁰, F. Cleva^{112,115}, E. Coccia^{44,45,43}, E. Codazzo^{155,156}, P.-F. Cohadon¹²¹, S. Colace³⁰, E. Colangeli¹²⁸, M. Colleoni⁹⁷, C. G. Collette¹⁵⁷, J. Collins⁶³, S. Colloms⁸⁵, A. Colombo^{158,127}, C. M. Compton², G. Connolly⁷⁶, L. Conti⁹¹, T. R. Corbitt¹², I. Cordero-Carrión¹⁵⁹, S. Corezzi^{75,51}

- N. J. Cornish ¹⁶⁰ I. Coronado,¹⁶¹ A. Corsi ¹⁶² R. Cottingham,⁶³ M. W. Coughlin ¹⁸ A. Couineaux,³⁸ P. Couvares ^{11,57} D. M. Coward,⁷² R. Coyne ¹⁶³ A. Cozzumbo,⁴⁴ J. D. E. Creighton ¹⁰ T. D. Creighton,¹⁶⁴ P. Cremonese ⁹⁷ S. Crook,⁶³ R. Crouch,² J. Csizmazia,² J. R. Cudell ¹⁶⁵ T. J. Cullen ¹¹ A. Cumming ⁸⁵ E. Cuoco ^{166,167} M. Cusinato ¹³⁸ L. V. Da Conceição,¹⁶⁸ T. Dal Canton ⁴¹ S. Dal Pra ¹⁶⁹ G. Dálya ⁹⁹ B. D'Angelo ²⁹ S. Danilishin ^{36,37} S. D'Antonio ³⁸ K. Danzmann,^{9,8,9} K. E. Darroch,¹²² L. P. Dartez ⁶³ R. Das,¹⁰⁵ A. Dasgupta,⁹² V. Dattilo ⁶² A. Daumas,²⁰ N. Davari,^{170,171} I. Dave,¹⁰² A. Davenport,¹³⁷ M. Davies,⁴¹ T. F. Davies,⁷² D. Davis ¹¹ L. Davis,⁷² M. C. Davis ¹⁸ P. Davis ^{172,173} E. J. Daw ¹⁷⁴ M. Dax ¹ J. De Bolle ⁹³ M. Deenadayalan,⁷⁸ J. Degallaix ¹⁷⁵ M. De Laurentis ^{32,4} F. De Lillo ²³ S. Della Torre ¹²⁷ W. Del Pozzo ^{80,79} A. Demagny,³¹ F. De Marco ^{39,38} G. Demasi,^{176,61} F. De Matteis ^{21,22} N. Demos,³⁵ T. Dent ¹⁷⁷ A. Depasse ¹⁵ N. DePergola,¹⁰¹ R. De Pietri ^{178,179} R. De Rosa ^{32,4} C. De Rossi ⁶² M. Desai ³⁵ R. DeSalvo ¹⁸⁰ A. DeSimone,¹⁸¹ R. De Simone,^{150,132} A. Dhani ¹ R. Diab,⁴⁶ M. C. Díaz ¹⁶⁴ M. Di Cesare ^{32,4} G. Dideron,¹⁸² T. Dietrich ¹ L. Di Fiore,⁴ C. Di Fronzo ⁷² M. Di Giovanni ^{39,38} T. Di Girolamo ^{32,4} D. Diksha,^{37,36} J. Ding ^{20,183} S. Di Pace ^{39,38} I. Di Palma ^{39,38} D. Di Piero,^{184,48} F. Di Renzo ⁵⁶ Divyajyoti ³³ A. Dmitriev ¹¹⁷ J. P. Docherty,⁸⁵ Z. Doctor ⁹⁵ N. Doerken,¹⁶⁸ E. Dohmen,² A. Doke,¹³³ A. Domiciano De Souza,¹⁸⁵ L. D'Onofrio ³⁸ F. Donovan,³⁵ K. L. Dooley ³³ T. Dooney,⁷¹ S. Doravari ⁷⁸ O. Dorosh,¹⁸⁶ W. J. D. Doyle,¹²² M. Drago ^{39,38} J. C. Driggers ² L. Dunn ¹²³ U. Dupletsa,⁴⁴ P.-A. Duverne ²⁰ D. D'Urso ^{170,155} P. Dutta Roy,⁴⁶ H. Duval ¹⁸⁷ S. E. Dwyer,² C. Eassa,² M. Ebersold ^{188,31} T. Eckhardt ⁹⁶ G. Eddolls ⁷⁷ A. Effler ⁶³ J. Eichholz ³⁴ H. Einsle,¹¹² M. Eisenmann,²⁵ M. Emma ⁵⁸ K. Endo,¹⁵¹ R. Enficiaud ¹ L. Errico ^{32,4} R. Espinosa,¹⁶⁴ M. C. Espitia,¹⁸⁹ M. Esposito ^{4,32} R. C. Essick ¹⁹⁰ H. Estellés ¹ T. Etzel,¹¹ M. Evans ³⁵ T. Evstafyeva,¹⁸² B. E. Ewing,⁷ J. M. Ezquiaga ¹⁴⁰ F. Fabrizi ^{60,61} V. Fafone ^{21,22} S. Fairhurst ³³ A. M. Farah ¹³⁰ B. Farr ⁷⁶ W. M. Farr ^{191,192} G. Favaro ⁹⁰ M. Favata ¹⁹³ M. Fays ¹⁶⁵ M. Fazio ⁵⁵ J. Feicht,¹¹ M. M. Fejer,⁸⁸ R. Felicetti ^{184,48} E. Fenyvesi ^{86,194} J. Fernandes,¹⁹⁵ T. Fernandes ^{196,138} D. Fernando,¹⁰⁹ S. Ferraiuolo ^{197,39,38} T. A. Ferreira,¹² F. Fidecaro ^{80,79} P. Figura ⁹⁴ A. Fiori ^{79,80} I. Fiori ⁶² M. Fishbach ¹⁹⁰ R. P. Fisher,¹²² R. Fittipaldi ^{198,132} V. Fiumara ^{199,132} R. Flaminio,³¹ S. M. Fleischer ²⁰⁰ L. S. Fleming,²⁰¹ E. Floden,¹⁸ H. Fong,¹¹³ J. A. Font ^{138,139} F. Fontinele-Nunes,¹⁸ C. Foo,¹ B. Fornal ²⁰² K. Franceschetti,¹⁷⁸ F. Frappez,³¹ S. Frasca,^{39,38} F. Frasconi ⁷⁹ J. P. Freed,⁶⁵ Z. Frei ²⁰³ A. Freise ^{37,106} O. Freitas ^{196,138} R. Frey ⁷⁶ W. Frischhertz,⁶³ P. Fritschel,³⁵ V. V. Frolov,⁶³ G. G. Fronzé ²⁸ M. Fuentes-Garcia ¹¹ S. Fujii,²⁰⁴ T. Fujimori,²⁰⁵ P. Fulda,⁴⁶ M. Fyffe,⁶³ B. Gadre ⁷¹ J. R. Gair ¹ S. Galaudage ¹⁸⁵ V. Galdi,²⁰⁶ R. Gamba,⁷ A. Gamboa ¹ S. Gamoji,¹⁸⁰ D. Ganapathy ²⁰⁷ A. Ganguly ⁷⁸ B. Garaventa ²⁹ J. García-Bellido ²⁰⁸ C. García-Quirós ¹⁸⁸ J. W. Gardner ³⁴ K. A. Gardner,¹¹³ S. Garg,⁴² J. Gargiulo ⁶² X. Garrido ⁴¹ A. Garron ⁹⁷ F. Garufi ^{32,4} P. A. Garver,⁸⁸ C. Gasbarra ^{21,22} B. Gateley,² F. Gautier ²⁰⁹ V. Gayathri ¹⁰ T. Gayer,⁷⁷ G. Gemme ²⁹ A. Gennai ⁷⁹ V. Gennari ⁹⁹ J. George,¹⁰² R. George ¹¹⁹ O. Gerberding ⁹⁶ L. Gergely ¹⁵³ Archisman Ghosh ⁹³ Sayantan Ghosh,¹⁹⁵ Shaon Ghosh ¹⁹³ Shrobana Ghosh,^{8,9} Suprovo Ghosh ²¹⁰ Tathagata Ghosh ⁷⁸ J. A. Giaime ^{12,63} K. D. Giardina,⁶³ D. R. Gibson,²⁰¹ C. Gier ⁵⁵ S. Gkaitatzis ^{80,79} J. Glanzer ¹¹ F. Glotin ⁴¹ J. Godfrey,⁷⁶ R. V. Godley,^{8,9} P. Godwin ¹¹ A. S. Goettel ³³ E. Goetz ¹¹³ J. Golomb,¹¹ S. Gomez Lopez ^{39,38} B. Goncharov ⁴⁴ G. González ¹² P. Goodarzi ²¹¹ S. Goode,⁶ A. W. Goodwin-Jones ¹⁵ M. Gosselin,⁶² R. Gouaty ³¹ D. W. Gould,³⁴ K. Govorkova,³⁵ A. Grado ^{75,51} V. Graham ⁸⁵ A. E. Granados ¹⁸ M. Granata ¹⁷⁵ V. Granata ^{212,132} S. Gras,³⁵ P. Grassia,¹¹ J. Graves,⁵⁷ C. Gray,² R. Gray ⁸⁵ G. Greco,⁵¹ A. C. Green ^{37,106} L. Green,²¹³ S. M. Green,¹²⁸ S. R. Green ²¹⁴ C. Greenberg,¹³³ A. M. Gretarsson,⁶⁵ H. K. Griffin,¹⁸ D. Griffith,¹¹ H. L. Griggs ⁵⁷ G. Grignani,^{75,51} C. Grimaud ³¹ H. Grote ³³ S. Grunewald ¹ D. Guerra ¹³⁸ D. Guetta ²¹⁵ G. M. Guidi ^{60,61} A. R. Guimaraes,¹² H. K. Gulati,⁹² F. Gulminelli ^{172,173} H. Guo ¹⁴⁵ W. Guo ⁷² Y. Guo ^{37,36} Anuradha Gupta ²¹⁶ I. Gupta,⁷ N. C. Gupta,⁹² S. K. Gupta,⁴⁶ V. Gupta ¹⁸ N. Gupte,¹ J. Gurs,⁹⁶ N. Gutierrez,¹⁷⁵ N. Guttman,⁶ F. Guzman ¹³¹ D. Haba,²¹⁷ M. Haberland ¹ S. Haino,²¹⁸ E. D. Hall ³⁵ E. Z. Hamilton ⁹⁷ G. Hammond ⁸⁵ M. Haney,³⁷ J. Hanks,² C. Hanna ⁷ M. D. Hannam,³³ O. A. Hannuksela ²¹⁹ A. G. Hanselman ¹³⁰ H. Hansen,² J. Hanson,⁶³ S. Hanumasagar,⁵⁷ R. Harada,⁴² A. R. Hardison,¹⁸¹ S. Harikumar ¹⁸⁶ K. Haris,^{37,71} I. Harley-Trochimczyk,¹³¹ T. Harmark ¹³⁴ J. Harms ^{44,45} G. M. Harry ²²⁰ I. W. Harry ¹²⁸ J. Hart,¹⁰³ B. Haskell,^{94,221,222} C. J. Haster ²¹³ K. Haughian ⁸⁵ H. Hayakawa,⁵⁰ K. Hayama,²²³ M. C. Heintze,⁶³ J. Heinze ¹¹⁷ J. Heinzl,³⁵ H. Heitmann ¹¹² F. Hellman ²⁰⁷ A. F. Helmling-Cornell ⁷⁶ G. Hemming ⁶² O. Henderson-Sapir ¹¹⁴ M. Hendry ⁸⁵ I. S. Heng,⁸⁵ M. H. Hennig ⁸⁵ C. Henshaw ⁵⁷ M. Heurs ^{8,9} A. L. Hewitt ^{224,225} J. Heynen,¹⁵ J. Heyns,³⁵ S. Higginbotham,³³ S. Hild,^{36,37} S. Hill,⁸⁵ Y. Himemoto ²²⁶ N. Hirata,²⁵ C. Hirose,²²⁷ D. Hofman,¹⁷⁵

- B. E. Hogan,⁶⁵ N. A. Holland,^{37, 106} I. J. Hollows¹⁷⁴ D. E. Holz¹³⁰ L. Honet,¹¹⁰ D. J. Horton-Bailey,²⁰⁷ J. Hough⁸⁵ S. Hourihane¹¹ N. T. Howard,¹⁴⁴ E. J. Howell⁷² C. G. Hoy¹²⁸ C. A. Hrishikesh,²¹ P. Hsi,³⁵ H.-F. Hsieh¹⁴² H.-Y. Hsieh,¹⁴² C. Hsiung,²²⁸ S.-H. Hsu,¹⁴⁶ W.-F. Hsu¹⁰⁸ Q. Hu⁸⁵ H. Y. Huang¹⁴¹ Y. Huang⁷ Y. T. Huang,⁷⁷ A. D. Huddart,²²⁹ B. Hughey,⁶⁵ V. Hui³¹ S. Husa⁹⁷ R. Huxford,⁷ L. Iampieri^{39, 38} G. A. Iandolo³⁶ M. Ianni,^{22, 21} G. Iannone¹³² J. Iascau,⁷⁶ K. Ide,²³⁰ R. Iden,²¹⁷ A. Ierardi,^{44, 45} S. Ikeda,¹⁴⁷ H. Imafuku,⁴² Y. Inoue,¹⁴¹ G. Iorio⁹⁰ P. Iosif^{184, 48} M. H. Iqbal,³⁴ J. Irwin⁸⁵ R. Ishikawa,²³⁰ M. Isi^{191, 192} K. S. Isleif²³¹ Y. Itoh^{205, 232} M. Iwaya,²⁰⁴ B. R. Iyer²⁴ C. Jacquet,⁹⁹ P.-E. Jacquet¹²¹ T. Jacquot,⁴¹ S. J. Jadhav,²³³ S. P. Jadhav¹⁵⁴ M. Jain,¹³³ T. Jain,²²⁴ A. L. James¹¹ K. Jani¹⁴⁴ J. Janquart¹⁵ N. N. Janthalur,²³³ S. Jaraba²³⁴ P. Jaranowski²³⁵ R. Jaume⁹⁷ W. Javed,³³ A. Jennings,² M. Jensen,² W. Jia,³⁵ J. Jiang¹⁴⁹ H.-B. Jin^{236, 237} G. R. Johns,¹²² N. A. Johnson,⁴⁶ M. C. Johnston²¹³ R. Johnston,⁸⁵ N. Johnny,^{8, 9} D. H. Jones³⁴ D. I. Jones,²¹⁰ R. Jones,⁸⁵ H. E. Jose,⁷⁶ P. Joshi,⁷ S. K. Joshi,⁷⁸ G. Joubert,⁵⁶ J. Ju,²³⁸ L. Ju⁷² K. Jung²³⁹ J. Junker³⁴ V. Juste,¹¹⁰ H. B. Kabagoz^{63, 35} T. Kajita²⁴⁰ I. Kaku,²⁰⁵ V. Kalogera⁹⁵ M. Kalomenopoulos²¹³ M. Kamiizumi⁵⁰ N. Kanda^{232, 205} S. Kandhasamy⁷⁸ G. Kang²⁴¹ N. C. Kannachel,⁶ J. B. Kanner,¹¹ S. A. KantiMahanty,¹⁸ S. J. Kapadia⁷⁸ D. P. Kapasi⁵⁴ M. Karthikeyan,¹³³ M. Kasprzack¹¹ H. Kato,¹⁵¹ T. Kato,²⁰⁴ E. Katsavounidis,³⁵ W. Katzman,⁶³ R. Kaushik¹⁰² K. Kawabe,² R. Kawamoto,²⁰⁵ D. Keitel⁹⁷ L. J. Kemperman¹¹⁴ J. Kennington⁷ F. A. Kerkow,¹⁸ R. Kesharwani⁷⁸ J. S. Key²⁴² R. Khadela,^{8, 9} S. Khadka,⁸⁸ S. S. Khadkikar,⁷ F. Y. Khalili¹⁰⁷ F. Khan^{8, 9} T. Khanam,¹⁶² M. Khursheed,¹⁰² N. M. Khusid,^{191, 192} W. Kiendrebeogo^{112, 243} N. Kijbunchoo¹¹⁴ C. Kim,²⁴⁴ J. C. Kim,²⁴⁵ K. Kim²⁴⁶ M. H. Kim²³⁸ S. Kim²⁴⁷ Y.-M. Kim²⁴⁶ C. Kimball⁹⁵ K. Kimes,⁵⁴ M. Kinnear,³³ J. S. Kissel² S. Klimenko,⁴⁶ A. M. Knee¹¹³ E. J. Knox,⁷⁶ N. Knust^{8, 9} K. Kobayashi,²⁰⁴ S. M. Koehlenbeck⁸⁸ G. Koekoek,^{37, 36} K. Kohri^{248, 249} K. Kokeyama^{33, 250} S. Koley^{44, 165} P. Kolitsidou¹¹⁷ A. E. Koloniari²⁵¹ K. Komori⁴² A. K. H. Kong¹⁴² A. Kontos²⁵² L. M. Koponen,¹¹⁷ M. Korobko⁹⁶ X. Kou,¹⁸ A. Koushik²³ N. Kouvatsoz⁶⁷ M. Kovalam,⁷² T. Koyama,¹⁵¹ D. B. Kozak,¹¹ S. L. Kranzhoff,^{36, 37} V. Kringel,^{8, 9} N. V. Krishnendu¹¹⁷ S. Kroker,²⁵³ A. Królak^{254, 186} K. Kruska,^{8, 9} J. Kubisz²⁵⁵ G. Kuehn,^{8, 9} S. Kulkarni²¹⁶ A. Kulur Ramamohan³⁴ Achal Kumar,⁴⁶ Anil Kumar,²³³ Praveen Kumar¹⁷⁷ Prayush Kumar²⁴ Rahul Kumar,² Rakesh Kumar,⁹² J. Kume^{256, 257, 42} K. Kuns³⁵ N. Kuntimaddi,³³ S. Kuroyanagi^{208, 258} S. Kuwahara⁴² K. Kwak²³⁹ K. Kwan,³⁴ S. Kwon⁴² G. Lacaille,⁸⁵ D. Laghi^{188, 99} A. H. Laity,¹⁶³ E. Lalande,²⁵⁹ M. Lalleman²³ P. C. Lalremruati,²⁶⁰ M. Landry,² B. B. Lane,³⁵ R. N. Lang³⁵ J. Lange,¹¹⁹ R. Langgin²¹³ B. Lantz⁸⁸ I. La Rosa⁹⁷ J. Larsen,²⁰⁰ A. Lartaux-Vollard⁴¹ P. D. Lasky⁶ J. Lawrence¹⁶⁴ M. Laxen⁶³ C. Lazarte¹³⁸ A. Lazzarini¹¹ C. Lazzaro,^{156, 155} P. Leaci^{39, 38} L. Leali,¹⁸ Y. K. Lecoeuche¹¹³ H. M. Lee²⁶¹ H. W. Lee²⁶² J. Lee,⁷⁷ K. Lee²³⁸ R.-K. Lee¹⁴² R. Lee,³⁵ Sungho Lee²⁴⁶ Sunjae Lee,²³⁸ Y. Lee,¹⁴¹ I. N. Legred,¹¹ J. Lehmann,^{8, 9} L. Lehner,¹⁸² M. Le Jean^{175, 115} A. Lemaître²⁶³ M. Lenti^{61, 176} M. Leonardi^{73, 74, 264} M. Lequime,⁴⁰ N. Leroy⁴¹ M. Lesovsky,¹¹ N. Letendre,³¹ M. Lethuillier⁵⁶ Y. Levin,⁶ K. Leyde,¹²⁸ A. K. Y. Li,¹¹ K. L. Li²⁶⁵ T. G. F. Li,¹⁰⁸ X. Li¹⁴⁸ Y. Li,⁹⁵ Z. Li,⁸⁵ A. Lihos,¹²² E. T. Lin¹⁴² F. Lin,¹⁴¹ L. C.-C. Lin²⁶⁵ Y.-C. Lin¹⁴² C. Lindsay,²⁰¹ S. D. Linker,¹⁸⁰ A. Liu²¹⁹ G. C. Liu²²⁸ Jian Liu⁷² F. Llamas Villarreal,¹⁶⁴ J. Llobera-Querol⁹⁷ R. K. L. Lo¹⁴⁰ J.-P. Locquet,¹⁰⁸ S. C. G. Loggins,²⁶⁶ M. R. Loizou,¹³³ L. T. London,⁶⁷ A. Longo^{60, 61} D. Lopez¹⁶⁵ M. Lopez Portilla,⁷¹ M. Lorenzini^{21, 22} A. Lorenzo-Medina¹⁷⁷ V. Lorette,⁴¹ M. Lormand,⁶³ G. Losurdo^{267, 79} E. Lotti,¹³³ T. P. Lott IV⁵⁷ J. D. Lough^{8, 9} H. A. Loughlin,³⁵ C. O. Lousto¹⁰⁹ N. Low,¹²³ N. Lu³⁴ L. Lucchesi⁷⁹ H. Lück,^{9, 8, 9} D. Lumaca²² A. P. Lundgren^{268, 269} A. W. Lussier²⁵⁹ R. Macas¹²⁸ M. MacInnis,³⁵ D. M. Macleod³³ I. A. O. MacMillan¹¹ A. Macquet⁴¹ K. Maeda,¹⁵¹ S. Maenaut¹⁰⁸ S. S. Magare,⁷⁸ R. M. Magee¹¹ E. Maggio,¹ R. Maggiore,^{37, 106} M. Magnozzi^{29, 30} M. Mahesh,⁹⁶ M. Maini,¹⁶³ S. Majhi,⁷⁸ E. Majorana,^{39, 38} C. N. Makarem,¹¹ D. Malakar¹⁰⁴ J. A. Malaquias-Reis,¹⁹ U. Mali¹⁹⁰ S. Maliakal,¹¹ A. Malik,¹⁰² L. Mallick^{168, 190} A.-K. Malz⁵⁸ N. Man,¹¹² M. Mancarella⁹⁸ V. Mandic¹⁸ V. Mangano^{170, 155} B. Mannix,⁷⁶ G. L. Mansell⁷⁷ M. Manske¹⁰ M. Mantovani⁶² M. Mapelli^{90, 91, 270} C. Marinelli¹⁰⁰ F. Marion³¹ A. S. Markosyan,⁸⁸ A. Markowitz,¹¹ E. Maros,¹¹ S. Marsat⁹⁹ F. Martelli^{60, 61} I. W. Martin⁸⁵ R. M. Martin¹⁹³ B. B. Martinez,¹³¹ D. A. Martinez,⁵⁴ M. Martinez,^{43, 271} V. Martinez¹²⁹ A. Martini,^{73, 74} J. C. Martins¹⁹ D. V. Martynov,¹¹⁷ E. J. Marx,³⁵ L. Massaro,^{36, 37} A. Masserot,³¹ M. Masso-Reid⁸⁵ S. Mastrogiovanni³⁸ T. Matcovich⁵¹ M. Matushechkina^{8, 9} L. Maurin,²⁰⁹ N. Mavalvala³⁵ N. Maxwell,² G. McCarroll,⁶³ R. McCarthy,² D. E. McClelland³⁴ S. McCormick,⁶³ L. McCuller¹¹ S. McEachin,¹²² C. McElhenny,¹²² G. I. McGhee⁸⁵ J. McGinn,⁸⁵ K. B. M. McGowan,¹⁴⁴ J. McIver¹¹³ A. McLeod⁷² I. McMahon¹⁸⁸ T. McRae,³⁴ R. McTeague,⁸⁵ D. Meacher¹⁰ B. N. Meagher,⁷⁷

- R. Mechum,¹⁰⁹ Q. Meijer,⁷¹ A. Melatos,¹²³ C. S. Menoni¹³⁷ F. Mera,² R. A. Mercer¹⁰ L. Mereni,¹⁷⁵ K. Merfeld,¹⁶² E. L. Merilh,⁶³ J. R. Mérou⁹⁷ J. D. Merritt,⁷⁶ M. Merzougui,¹¹² C. Messick¹⁰ B. Mestichelli,⁴⁴ M. Meyer-Conde²⁷² F. Meylahn^{8,9} A. Mhaske,⁷⁸ A. Miani^{73,74} H. Miao,²⁷³ C. Michel¹⁷⁵ Y. Michimura⁴² H. Middleton¹¹⁷ D. P. Mihaylov¹⁰³ S. J. Miller¹¹ M. Millhouse⁵⁷ E. Milotti^{184,48} V. Milotti⁹⁰ Y. Minenkov,²² E. M. Minihan,⁶⁵ L. M. Mir⁴³ L. Mirasola^{155,156} M. Miravet-Tenés¹³⁸ C.-A. Mirutescu⁴³ A. Mishra,²⁴ C. Mishra¹⁰⁵ T. Mishra⁴⁶ A. L. Mitchell,^{37,106} J. G. Mitchell,⁶⁵ S. Mitra⁷⁸ V. P. Mitrofanov¹⁰⁷ K. Mitsuhashi,²⁵ R. Mittleman,³⁵ O. Miyakawa⁵⁰ S. Miyoki⁵⁰ A. Miyoko,⁶⁵ G. Mo³⁵ L. Mobilia^{60,61} S. R. P. Mohapatra,¹¹ S. R. Mohite⁷ M. Molina-Ruiz²⁰⁷ M. Mondin,¹⁸⁰ J. K. Monsalve,¹⁸⁹ M. Montani,^{60,61} C. J. Moore,²²⁴ D. Moraru,² A. More⁷⁸ S. More⁷⁸ C. Moreno¹³⁵ E. A. Moreno³⁵ G. Moreno,² A. Moreso Serra,⁸¹ S. Morisaki^{42,204} Y. Moriwaki¹⁵¹ G. Morras²⁰⁸ A. Moscatello⁹⁰ M. Mould³⁵ B. Mours⁶⁴ C. M. Mow-Lowry^{37,106} L. Muccillo^{176,61} F. Muciaccia^{39,38} D. Mukherjee¹¹⁷ Samanwaya Mukherjee,²⁴ Soma Mukherjee,¹⁶⁴ Subroto Mukherjee,⁹² Suvodip Mukherjee¹³ N. Mukund³⁵ A. Mullavey,⁶³ H. Mullock,¹¹³ J. Mundi,²²⁰ C. L. Mungoli,⁷² M. Murakoshi,²³⁰ P. G. Murray⁸⁵ D. Nabari^{73,74} S. L. Nadji,^{8,9} A. Nagar,^{28,274} N. Nagarajan⁸⁵ K. Nakagaki,⁵⁰ K. Nakamura²⁵ H. Nakano²⁷⁵ M. Nakano,¹¹ D. Nanadoumgar-Lacroze⁴³ D. Nandi,¹² V. Napolano,⁶² P. Narayan²¹⁶ I. Nardecchia²² T. Narikawa,²⁰⁴ H. Narola,⁷¹ L. Naticchioni³⁸ R. K. Nayak²⁶⁰ L. Negri,⁷¹ A. Nela,⁸⁵ C. Nelle,⁷⁶ A. Nelson¹³¹ T. J. N. Nelson,⁶³ M. Nery,^{8,9} A. Neunzert² S. Ng,⁵⁴ L. Nguyen Quynh²⁷⁶ S. A. Nichols,¹² A. B. Nielsen²⁷⁷ Y. Nishino,^{25,42} A. Nishizawa²⁷⁸ S. Nissanke,^{279,37} W. Niu⁷ F. Nocera,⁶² J. Noller,²⁸⁰ M. Norman,³³ C. North,³³ J. Novak^{115,234,281} R. Nowicki¹⁴⁴ J. F. Nuño Siles²⁰⁸ L. K. Nuttall¹²⁸ K. Obayashi,²³⁰ J. Oberling² J. O'Dell,²²⁹ E. Oelker³⁵ M. Oertel^{234,115,282,281} G. Oganessian,^{44,45} T. O'Hanlon,⁶³ M. Ohashi⁵⁰ F. Ohme^{8,9} R. Oliveri^{115,282,281} R. Omer,¹⁸ B. O'Neal,¹²² M. Onishi,¹⁵¹ K. Oohara²⁸³ B. O'Reilly⁶³ M. Orselli^{51,75} R. O'Shaughnessy¹⁰⁹ S. O'Shea,⁸⁵ S. Oshino⁵⁰ C. Osthelder,¹¹ I. Ota¹² D. J. Ottaway¹¹⁴ A. Ouzriat,⁵⁶ H. Overmier,⁶³ B. J. Owen²⁸⁴ R. Ozaki,²³⁰ A. E. Pace⁷ R. Pagano¹² M. A. Page²⁵ A. Pai¹⁹⁵ L. Paiella,⁴⁴ A. Pal,²⁸⁵ S. Pal²⁶⁰ M. A. Palaia^{79,80} M. Pálfi,²⁰³ P. P. Palma,^{39,21,22} C. Palomba³⁸ P. Palud²⁰ H. Pan,¹⁴² J. Pan,⁷² K. C. Pan¹⁴² P. K. Panda,²³³ Shiksha Pandey,⁷ Swadha Pandey,³⁵ P. T. H. Pang,^{37,71} F. Pannarale^{39,38} K. A. Pannone,⁵⁴ B. C. Pant,¹⁰² F. H. Panther,⁷² M. Panzeri,^{60,61} F. Paoletti⁷⁹ A. Paolone^{38,286} A. Papadopoulos⁸⁵ E. E. Papalexakis,²¹¹ L. Papalini^{79,80} G. Papigkiotis²⁵¹ A. Paquis,⁴¹ A. Parisi^{75,51} B.-J. Park,²⁴⁶ J. Park²⁸⁷ W. Parker⁶³ G. Pascale,^{8,9} D. Pascucci⁹³ A. Pasqualetti⁶² R. Passaquieti^{80,79} L. Passenger,⁶ D. Passuello,⁷⁹ O. Patane² A. V. Patel¹⁴¹ D. Pathak,⁷⁸ A. Patra,³³ B. Patricelli^{80,79} B. G. Patterson,³³ K. Paul¹⁰⁵ S. Paul⁷⁶ E. Payne¹¹ T. Pearce,³³ M. Pedraza,¹¹ A. Pele¹¹ F. E. Peña Arellano²⁸⁸ X. Peng,¹¹⁷ Y. Peng,⁵⁷ S. Penn²⁸⁹ M. D. Penuliar,⁵⁴ A. Perego^{73,74} Z. Pereira,¹³³ C. Périgois^{290,91,90} G. Perna⁹⁰ A. Perreca^{73,74,44} J. Perret²⁰ S. Perriès⁵⁶ J. W. Perry,^{37,106} D. Pesios,²⁵¹ S. Peters,¹⁶⁵ S. Petracca,²⁰⁶ C. Petrillo,⁷⁵ H. P. Pfeiffer¹ H. Pham,⁶³ K. A. Pham¹⁸ K. S. Phukon¹¹⁷ H. Phurailatpam,²¹⁹ M. Piarulli,⁹⁹ L. Piccari^{39,38} O. J. Piccinni³⁴ M. Pichot¹¹² M. Piendibene^{80,79} F. Piergiovanni^{60,61} L. Pierini³⁸ G. Pierra³⁸ V. Pierro^{291,132} M. Pietrzak,⁹⁴ M. Pillas¹⁶⁵ F. Pilo⁷⁹ L. Pinard¹⁷⁵ I. M. Pinto^{291,132,292,32} M. Pinto⁶² B. J. Piotrkowski¹⁰ M. Pirello,² M. D. Pitkin^{224,85} A. Placidi⁵¹ E. Placidi^{39,38} M. L. Planas⁹⁷ W. Plastino^{212,22} C. Plunkett³⁵ R. Poggiani^{80,79} E. Polini,³⁵ J. Pomper,^{79,80} L. Pompili¹ J. Poon,²¹⁹ E. Porcelli,³⁷ E. K. Porter,²⁰ C. Posnansky⁷ R. Poulton⁶² J. Powell¹⁵⁴ G. S. Prabhu,⁷⁸ M. Pracchia¹⁶⁵ B. K. Pradhan⁷⁸ T. Pradier⁶⁴ A. K. Prajapati,⁹² K. Prasai²⁹³ R. Prasanna,²³³ P. Prasia,⁷⁸ G. Pratten¹¹⁷ G. Principe^{184,48} G. A. Prodi^{73,74} P. Prosperi,⁷⁹ P. Proposito,^{21,22} A. C. Providence,⁶⁵ A. Puecher¹ J. Pullin¹² P. Puppo,³⁸ M. Pürner¹⁶³ H. Qi¹⁶ J. Qin³⁴ G. Quémener^{173,115} V. Quetschke,¹⁶⁴ L. H. Quiceno,¹⁸⁹ P. J. Quinonez,⁶⁵ N. Qutob,⁵⁷ F. J. Raab² R. Rading,²³¹ I. Rainho,¹³⁸ S. Raja,¹⁰² C. Rajan,¹⁰² B. Rajbhandari¹⁰⁹ K. E. Ramirez⁶³ F. A. Ramis Vidal⁹⁷ M. Ramos Arevalo¹⁶⁴ A. Ramos-Buades^{97,37} S. Ranjan⁵⁷ K. Ransom,⁶³ P. Rapagnani^{39,38} B. Ratto,⁶⁵ A. Ravichandran,¹³³ A. Ray⁹⁵ V. Raymond³³ M. Razzano^{80,79} J. Read,⁵⁴ T. Regimbau,³¹ S. Reid,⁵⁵ C. Reissel,³⁵ D. H. Reitze¹¹ A. I. Renzini,¹¹ A. Renzini¹²⁶ B. Revenu^{294,41} A. Revilla Peña,⁸¹ R. Reyes,¹⁸⁰ L. Ricca¹⁵ F. Ricci^{39,38} M. Ricci^{38,39} A. Ricciardone^{80,79} J. Rice,⁷⁷ J. W. Richardson²¹¹ M. L. Richardson,¹¹⁴ A. Rijal,⁶⁵ K. Riles⁸⁹ H. K. Riley,³³ S. Rinaldi²⁷⁰ J. Rittmeyer,⁹⁶ C. Robertson,²²⁹ F. Robinet,⁴¹ M. Robinson,² A. Rocchi²² L. Rolland³¹ J. G. Rollins¹¹ A. E. Romano¹⁸⁹ R. Romano^{3,4} A. Romero³¹ I. M. Romero-Shaw,²²⁴ J. H. Romie,⁶³ S. Ronchini⁷ T. J. Rooke¹¹⁴ L. Rosa,^{4,32} T. J. Rosauer,²¹¹ C. A. Rose,⁵⁷ D. Rosińska¹²⁴ M. P. Ross⁵³ M. Rossello-Sastre⁹⁷ S. Rowan⁸⁵ S. K. Roy^{191,192} S. Roy¹⁵ D. Rozza^{126,127} P. Ruggi,⁶² N. Ruhama,²³⁹

- E. Ruiz Morales ^{295, 208} K. Ruiz-Rocha,¹⁴⁴ S. Sachdev ⁵⁷ T. Sadecki,² P. Saffarieh ^{37, 106} S. Safi-Harb ¹⁶⁸
M. R. Sah ¹³ S. Saha ¹⁴² T. Sainrat ⁶⁴ S. Sajith Menon ^{215, 39, 38} K. Sakai,²⁹⁶ Y. Sakai ²⁷²
M. Sakellariadou ⁶⁷ S. Sakon ⁷ O. S. Salafia ^{158, 127, 126} F. Salces-Carcoba ¹¹ L. Salconi,⁶² M. Saleem ¹¹⁹
F. Salemi ^{39, 38} M. Sallé ³⁷ S. U. Salunkhe,⁷⁸ S. Salvador ^{173, 172} A. Salvarese,¹¹⁹ A. Samajdar ^{71, 37}
A. Sanchez,² E. J. Sanchez,¹¹ L. E. Sanchez,¹¹ N. Sanchis-Gual ¹³⁸ J. R. Sanders,¹⁸¹ E. M. Sanger ¹
F. Santoliquido ^{44, 45} F. Sarandrea,²⁸ T. R. Saravanan,⁷⁸ N. Sarin,⁶ P. Sarkar,^{8, 9} A. Sasli ²⁵¹ P. Sassi ^{51, 75}
B. Sassolas ¹⁷⁵ R. Sato,²²⁷ S. Sato,¹⁵¹ Yukino Sato,¹⁵¹ Yu Sato,¹⁵¹ O. Sauter ⁴⁶ R. L. Savage ² T. Sawada ⁵⁰
H. L. Sawant,⁷⁸ S. Sayah,¹⁷⁵ V. Scacco,^{21, 22} D. Schaetzl,¹¹ M. Scheel,¹⁴⁸ A. Schiebelbein,¹⁹⁰ M. G. Schiworski ⁷⁷
P. Schmidt ¹¹⁷ S. Schmidt ⁷¹ R. Schnabel ⁹⁶ M. Schneewind,^{8, 9} R. M. S. Schofield,⁷⁶ K. Schouteden ¹⁰⁸
B. W. Schulte,^{8, 9} B. F. Schutz,^{33, 8, 9} E. Schwartz ²⁹⁷ M. Scialpi ²⁹⁸ J. Scott ⁸⁵ S. M. Scott ³⁴
R. M. Sedas ⁶³ T. C. Seetharamu,⁸⁵ M. Seglar-Arroyo ⁴³ Y. Sekiguchi ²⁹⁹ D. Sellers,⁶³ N. Sembo,²⁰⁵
A. S. Sengupta ³⁰⁰ E. G. Seo ⁸⁵ J. W. Seo ¹⁰⁸ V. Sequino,^{32, 4} M. Serra ³⁸ A. Sevrin,¹⁸⁷ T. Shaffer,²
U. S. Shah ⁵⁷ M. A. Shaikh ²⁶¹ L. Shao ³⁰¹ A. K. Sharma ⁹⁷ Preeti Sharma,¹² Prianka Sharma,¹⁰²
Ritwik Sharma,¹⁸ S. Sharma Chaudhary,¹⁰⁴ P. Shawhan ¹²⁵ N. S. Shcheblanov ^{302, 263} E. Sheridan,¹⁴⁴
Z.-H. Shi,¹⁴² M. Shikauchi,⁴² R. Shimomura,³⁰³ H. Shinkai ³⁰³ S. Shirke,⁷⁸ D. H. Shoemaker ³⁵
D. M. Shoemaker ¹¹⁹ R. W. Short,² S. ShyamSundar,¹⁰² A. Sider,¹⁵⁷ H. Siegel ^{191, 192} D. Sigg ²
L. Silenzi ^{36, 37} L. Silvestri ^{39, 169} M. Simmonds,¹¹⁴ L. P. Singer ³⁰⁴ Amitesh Singh,²¹⁶ Anika Singh,¹¹
D. Singh ²⁰⁷ N. Singh ⁹⁷ S. Singh,^{217, 59} A. M. Sintes ⁹⁷ V. Sipala,^{170, 155} V. Skliris ³³ B. J. J. Slagmolen ³⁴
D. A. Slater,²⁰⁰ T. J. Slaven-Blair,⁷² J. Smetana,¹¹⁷ J. R. Smith ⁵⁴ L. Smith ^{85, 184, 48} R. J. E. Smith ⁶
W. J. Smith ¹⁴⁴ S. Soares de Albuquerque Filho,⁶⁰ M. Soares-Santos,¹⁸⁸ K. Somiya ²¹⁷ I. Song ¹⁴² S. Soni ³⁵
V. Sordini ⁵⁶ F. Sorrentino,²⁹ H. Sotani ³⁰⁵ F. Spada ⁷⁹ V. Spagnuolo ³⁷ A. P. Spencer ⁸⁵ P. Spinicelli ⁶²
A. K. Srivastava,⁹² F. Stachurski ⁸⁵ C. J. Stark,¹²² D. A. Steer ³⁰⁶ N. Steinle ¹⁶⁸ J. Steinlechner,^{36, 37}
S. Steinlechner ^{36, 37} N. Stergioulas ²⁵¹ P. Stevens,⁴¹ M. StPierre,¹⁶³ M. D. Strong,¹² A. Strunk,²
A. L. Stuver,^{101, *} M. Suchenek,⁹⁴ S. Sudhagar ⁹⁴ Y. Sudo,²³⁰ N. Sueltmann,⁹⁶ L. Suleiman ⁵⁴ K. D. Sullivan,¹²
J. Sun ²⁴¹ L. Sun ³⁴ S. Sunil,⁹² J. Suresh ¹¹² B. J. Sutton,⁶⁷ P. J. Sutton ³³ K. Suzuki,²¹⁷ M. Suzuki,²⁰⁴
B. L. Swinkels ³⁷ A. Syx ¹¹⁵ M. J. Szczepańczyk ³⁰⁷ P. Szewczyk ¹²⁴ M. Tacca ³⁷ H. Tagoshi ²⁰⁴
K. Takada,²⁰⁴ H. Takahashi ²⁷² R. Takahashi ²⁵ A. Takamori ⁴² S. Takano ³⁰⁸ H. Takeda ^{309, 310}
K. Takeshita,²¹⁷ I. Takimoto Schmiegelow,^{44, 45} M. Takou-Ayaoh,⁷⁷ C. Talbot,¹³⁰ M. Tamaki,²⁰⁴ N. Tamanini ⁹⁹
D. Tanabe,¹⁴¹ K. Tanaka,⁵⁰ S. J. Tanaka ²³⁰ S. Tanioka ³³ D. B. Tanner,⁴⁶ W. Tanner,^{8, 9} L. Tao ²¹¹
R. D. Tapia,⁷ E. N. Tapia San Martın ³⁷ C. Taranto,^{21, 22} A. Taruya ³¹¹ J. D. Tasson ¹⁵² J. G. Tau ¹⁰⁹
D. Tellez,⁵⁴ R. Tenorio ⁹⁷ H. Themann,¹⁸⁰ A. Theodoropoulos ¹³⁸ M. P. Thirugnanasambandam,⁷⁸
L. M. Thomas ¹¹ M. Thomas,⁶³ P. Thomas,² J. E. Thompson ²¹⁰ S. R. Thondapu,¹⁰² K. A. Thorne,⁶³
E. Thrane ⁶ J. Tissino ^{44, 45} A. Tiwari,⁷⁸ Pawan Tiwari,⁴⁴ Praveer Tiwari,¹⁹⁵ S. Tiwari ¹⁸⁸ V. Tiwari ¹¹⁷
M. R. Todd,⁷⁷ M. Toffano,⁹⁰ A. M. Toivonen ¹⁸ K. Toland ⁸⁵ A. E. Tolley ¹²⁸ T. Tomaru ²⁵ V. Tommasini,¹¹
T. Tomura ⁵⁰ H. Tong ⁶ C. Tong-Yu,¹⁴¹ A. Torres-Forne ^{138, 139} C. I. Torrie,¹¹ I. Tosta e Melo ³¹²
E. Tournefier ³¹ M. Trad Nery,¹¹² K. Tran,¹²² A. Trapananti ^{52, 51} R. Travaglini ¹⁶⁷ F. Travasso ^{52, 51}
G. Traylor,⁶³ M. Trevor,¹²⁵ M. C. Tringali ⁶² A. Tripathee ⁸⁹ G. Troian ^{184, 48} A. Trovato ^{184, 48} L. Trozzo,⁴
R. J. Trudeau,¹¹ T. Tsang ³³ S. Tsuchida ³¹³ L. Tsukada ²¹³ K. Turbang ^{187, 23} M. Turconi ¹¹² C. Turski,⁹³
H. Ubach ^{81, 82} N. Uchikata ²⁰⁴ T. Uchiyama ⁵⁰ R. P. Udall ¹¹ T. Uehara ³¹⁴ K. Ueno ⁴² V. Undheim ²⁷⁷
L. E. Uronen,²¹⁹ T. Ushiba ⁵⁰ M. Vacatello ^{79, 80} H. Vahlbruch ^{8, 9} N. Vaidya ¹¹ G. Vajente ¹¹ A. Vajpeyi,⁶
J. Valencia ⁹⁷ M. Valentini ^{106, 37} S. A. Vallejo-Peña ¹⁸⁹ S. Vallerio,²⁸ V. Valsan ¹⁰ M. van Dael ^{37, 315}
E. Van den Bossche ¹⁸⁷ J. F. J. van den Brand ^{36, 106, 37} C. Van Den Broeck,^{71, 37} M. van der Sluys ^{37, 71}
A. Van de Walle,⁴¹ J. van Dongen ^{37, 106} K. Vandra,¹⁰¹ M. VanDyke,¹¹⁸ H. van Haevermaet ²³
J. V. van Heijningen ^{37, 106} P. Van Hove ⁶⁴ J. Vanier,²⁵⁹ M. VanKeuren,¹⁰³ J. Vanosky,² N. van Remortel ²³
M. Vardaro,^{36, 37} A. F. Vargas ¹²³ V. Varma ¹³³ A. N. Vazquez,⁸⁸ A. Vecchio ¹¹⁷ G. Vedovato,⁹¹ J. Veitch ⁸⁵
P. J. Veitch ¹¹⁴ S. Venikoudis,¹⁵ R. C. Venterea ¹⁸ P. Verdier ⁵⁶ M. Vereecken,¹⁵ D. Verkindt ³¹ B. Verma,¹³³
Y. Verma ¹⁰² S. M. Vermeulen ¹¹ F. Vetrano,⁶⁰ A. Veutro ^{38, 39} A. Vicere ^{60, 61} S. Vidyant,⁷⁷
A. D. Viets ⁸⁷ A. Vijaykumar ¹⁹⁰ A. Vilkhya,¹⁰⁹ N. Villanueva Espinosa,¹³⁸ V. Villa-Ortega ¹⁷⁷
E. T. Vincent ⁵⁷ J.-Y. Vinet,¹¹² S. Viret,⁵⁶ S. Vitale ³⁵ H. Vocca ^{75, 51} D. Voigt ⁹⁶ E. R. G. von Reis,²
J. S. A. von Wrangel,^{8, 9} W. E. Vossius,²³¹ L. Vujeva ¹⁴⁰ S. P. Vyatchanin ¹⁰⁷ J. Wack,¹¹ L. E. Wade,¹⁰³
M. Wade ¹⁰³ K. J. Wagner ¹⁰⁹ L. Wallace,¹¹ E. J. Wang,⁸⁸ H. Wang ²¹⁷ J. Z. Wang,⁸⁹ W. H. Wang,¹⁶⁴
Y. F. Wang ¹ G. Waratkar ¹⁹⁵ J. Warner,² M. Was ³¹ T. Washimi ²⁵ N. Y. Washington,¹¹ D. Watarai,⁴²
B. Weaver,² S. A. Webster,⁸⁵ N. L. Weickhardt ⁹⁶ M. Weinert,^{8, 9} A. J. Weinstein ¹¹ R. Weiss,³⁵ L. Wen ⁷²

K. Wette³⁴, J. T. Whelan¹⁰⁹, B. F. Whiting⁴⁶, C. Whittle¹¹, E. G. Wickens¹²⁸, D. Wilken^{8,9,9},
 A. T. Wilkin²¹¹, B. M. Williams¹¹⁸, D. Williams⁸⁵, M. J. Williams¹²⁸, N. S. Williams¹, J. L. Willis¹¹,
 B. Willke^{9,8,9}, M. Wils¹⁰⁸, L. Wilson¹⁰³, C. W. Winborn¹⁰⁴, J. Winterflood⁷², C. C. Wipf¹¹, G. Woan⁸⁵,
 J. Woehler^{36,37}, N. E. Wolfe³⁵, H. T. Wong¹⁴¹, I. C. F. Wong^{219,108}, K. Wong¹⁹⁰, T. Wouters^{71,37},
 J. L. Wright², M. Wright^{85,71}, B. Wu⁷⁷, C. Wu¹⁴², D. S. Wu^{8,9}, H. Wu¹⁴², K. Wu¹¹⁸, Q. Wu⁵³, Y. Wu⁹⁵,
 Z. Wu⁹⁹, E. Wuchner⁵⁴, D. M. Wysocki¹⁰, V. A. Xu²⁰⁷, Y. Xu⁹⁷, N. Yadav²⁸, H. Yamamoto¹¹,
 K. Yamamoto¹⁵¹, T. S. Yamamoto⁴², T. Yamamoto⁵⁰, R. Yamazaki²³⁰, T. Yan¹¹⁷, K. Z. Yang¹⁸,
 Y. Yang¹⁴⁶, Z. Yarbrough¹², J. Yebana⁹⁷, S.-W. Yeh¹⁴², A. B. Yelikar¹⁴⁴, X. Yin³⁵, J. Yokoyama^{316,42},
 T. Yokozawa⁵⁰, S. Yuan⁷², H. Yuzurihara⁵⁰, M. Zanolin⁶⁵, M. Zeeshan¹⁰⁹, T. Zelenova⁶², J.-P. Zendri⁹¹,
 M. Zeoli¹⁵, M. Zerrad⁴⁰, M. Zevin⁹⁵, L. Zhang¹¹, N. Zhang⁵⁷, R. Zhang¹⁴⁹, T. Zhang¹¹⁷, C. Zhao⁷²,
 Yue Zhao¹⁶¹, Yuhang Zhao²⁰, Z.-C. Zhao³¹⁷, Y. Zheng¹⁰⁴, H. Zhong¹⁸, H. Zhou⁷⁷, H. O. Zhu⁷²,
 Z.-H. Zhu^{317,318}, A. B. Zimmerman¹¹⁹, L. Zimmermann⁵⁶, M. E. Zucker^{35,11} and J. Zweizig¹¹

(The LIGO Scientific Collaboration, the Virgo Collaboration, and the KAGRA Collaboration)

¹Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam, Germany

²LIGO Hanford Observatory, Richland, WA 99352, USA

³Dipartimento di Farmacia, Università di Salerno, I-84084 Fisciano, Salerno, Italy

⁴INFN, Sezione di Napoli, I-80126 Napoli, Italy

⁵University of Warwick, Coventry CV4 7AL, United Kingdom

⁶OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia

⁷The Pennsylvania State University, University Park, PA 16802, USA

⁸Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany

⁹Leibniz Universität Hannover, D-30167 Hannover, Germany

¹⁰University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

¹¹LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA

¹²Louisiana State University, Baton Rouge, LA 70803, USA

¹³Tata Institute of Fundamental Research, Mumbai 400005, India

¹⁴Centre de Physique Théorique, Aix-Marseille Université,
 Campus de Luminy, 163 Av. de Luminy, 13009 Marseille, France

¹⁵Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium

¹⁶Queen Mary University of London, London E1 4NS, United Kingdom

¹⁷University of California, Davis, Davis, CA 95616, USA

¹⁸University of Minnesota, Minneapolis, MN 55455, USA

¹⁹Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil

²⁰Université Paris Cité, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France

²¹Università di Roma Tor Vergata, I-00133 Roma, Italy

²²INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy

²³Universiteit Antwerpen, 2000 Antwerpen, Belgium

²⁴International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India

²⁵Gravitational Wave Science Project, National Astronomical Observatory of Japan,
 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan

²⁶Advanced Technology Center, National Astronomical Observatory of Japan,
 2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan

²⁷Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany

²⁸INFN Sezione di Torino, I-10125 Torino, Italy

²⁹INFN, Sezione di Genova, I-16146 Genova, Italy

³⁰Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy

³¹Univ. Savoie Mont Blanc, CNRS, Laboratoire d'Annecy de Physique des Particules - IN2P3, F-74000 Annecy, France

³²Università di Napoli "Federico II", I-80126 Napoli, Italy

³³Cardiff University, Cardiff CF24 3AA, United Kingdom

³⁴OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia

³⁵LIGO Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

³⁶Maastricht University, 6200 MD Maastricht, Netherlands

³⁷Nikhef, 1098 XG Amsterdam, Netherlands

³⁸INFN, Sezione di Roma, I-00185 Roma, Italy

³⁹Università di Roma "La Sapienza", I-00185 Roma, Italy

⁴⁰Aix Marseille Univ, CNRS, Centrale Med, Institut Fresnel, F-13013 Marseille, France

⁴¹Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

⁴²University of Tokyo, Tokyo, 113-0033, Japan

⁴³Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology,
 Campus UAB, E-08193 Bellaterra (Barcelona), Spain

⁴⁴Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy

- ⁴⁵ INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy
- ⁴⁶ University of Florida, Gainesville, FL 32611, USA
- ⁴⁷ Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università di Udine, I-33100 Udine, Italy
- ⁴⁸ INFN, Sezione di Trieste, I-34127 Trieste, Italy
- ⁴⁹ Tecnológico de Monterrey, Escuela de Ingeniería y Ciencias, 64849 Monterrey, Nuevo León, Mexico
- ⁵⁰ Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo,
238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan
- ⁵¹ INFN, Sezione di Perugia, I-06123 Perugia, Italy
- ⁵² Università di Camerino, I-62032 Camerino, Italy
- ⁵³ University of Washington, Seattle, WA 98195, USA
- ⁵⁴ California State University Fullerton, Fullerton, CA 92831, USA
- ⁵⁵ SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom
- ⁵⁶ Université Claude Bernard Lyon 1, CNRS, IP2I Lyon / IN2P3, UMR 5822, F-69622 Villeurbanne, France
- ⁵⁷ Georgia Institute of Technology, Atlanta, GA 30332, USA
- ⁵⁸ Royal Holloway, University of London, London TW20 0EX, United Kingdom
- ⁵⁹ Astronomical course, The Graduate University for Advanced Studies (SOKENDAI),
2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ⁶⁰ Università degli Studi di Urbino “Carlo Bo”, I-61029 Urbino, Italy
- ⁶¹ INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
- ⁶² European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
- ⁶³ LIGO Livingston Observatory, Livingston, LA 70754, USA
- ⁶⁴ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
- ⁶⁵ Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA
- ⁶⁶ Dipartimento di Fisica “E.R. Caianiello”, Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ⁶⁷ King’s College London, University of London, London WC2R 2LS, United Kingdom
- ⁶⁸ Korea Institute of Science and Technology Information, Daejeon 34141, Republic of Korea
- ⁶⁹ International College, Osaka University, 1-1 Machikaneyama-cho, Toyonaka City, Osaka 560-0043, Japan
- ⁷⁰ Accelerator Laboratory, High Energy Accelerator Research Organization (KEK),
1-1 Oho, Tsukuba City, Ibaraki 305-0801, Japan
- ⁷¹ Institute for Gravitational and Subatomic Physics (GRASP),
Utrecht University, 3584 CC Utrecht, Netherlands
- ⁷² OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
- ⁷³ Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
- ⁷⁴ INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
- ⁷⁵ Università di Perugia, I-06123 Perugia, Italy
- ⁷⁶ University of Oregon, Eugene, OR 97403, USA
- ⁷⁷ Syracuse University, Syracuse, NY 13244, USA
- ⁷⁸ Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India
- ⁷⁹ INFN, Sezione di Pisa, I-56127 Pisa, Italy
- ⁸⁰ Università di Pisa, I-56127 Pisa, Italy
- ⁸¹ Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona (UB),
c. Martí i Franquès, 1, 08028 Barcelona, Spain
- ⁸² Departament de Física Quàntica i Astrofísica (FQA),
Universitat de Barcelona (UB), c. Martí i Franquès, 1, 08028 Barcelona, Spain
- ⁸³ Institut d’Estudis Espacials de Catalunya, c. Gran Capità, 2-4, 08034 Barcelona, Spain
- ⁸⁴ Dipartimento di Medicina, Chirurgia e Odontoiatria “Scuola Medica Salernitana”,
Università di Salerno, I-84081 Baronissi, Salerno, Italy
- ⁸⁵ IGR, University of Glasgow, Glasgow G12 8QQ, United Kingdom
- ⁸⁶ HUN-REN Wigner Research Centre for Physics, H-1121 Budapest, Hungary
- ⁸⁷ Concordia University Wisconsin, Mequon, WI 53097, USA
- ⁸⁸ Stanford University, Stanford, CA 94305, USA
- ⁸⁹ University of Michigan, Ann Arbor, MI 48109, USA
- ⁹⁰ Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
- ⁹¹ INFN, Sezione di Padova, I-35131 Padova, Italy
- ⁹² Institute for Plasma Research, Bhat, Gandhinagar 382428, India
- ⁹³ Universiteit Gent, B-9000 Gent, Belgium
- ⁹⁴ Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland
- ⁹⁵ Northwestern University, Evanston, IL 60208, USA
- ⁹⁶ Universität Hamburg, D-22761 Hamburg, Germany
- ⁹⁷ IAC3–IEEC, Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain
- ⁹⁸ Aix-Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France
- ⁹⁹ Laboratoire des 2 Infinis - Toulouse (L2IT-IN2P3), F-31062 Toulouse Cedex 9, France
- ¹⁰⁰ Università di Siena, Dipartimento di Scienze Fisiche,
della Terra e dell’Ambiente, I-53100 Siena, Italy

- ¹⁰¹ Villanova University, Villanova, PA 19085, USA
- ¹⁰² RRCAT, Indore, Madhya Pradesh 452013, India
- ¹⁰³ Kenyon College, Gambier, OH 43022, USA
- ¹⁰⁴ Missouri University of Science and Technology, Rolla, MO 65409, USA
- ¹⁰⁵ Indian Institute of Technology Madras, Chennai 600036, India
- ¹⁰⁶ Department of Physics and Astronomy, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, Netherlands
- ¹⁰⁷ Lomonosov Moscow State University, Moscow 119991, Russia
- ¹⁰⁸ Katholieke Universiteit Leuven, Oude Markt 13, 3000 Leuven, Belgium
- ¹⁰⁹ Rochester Institute of Technology, Rochester, NY 14623, USA
- ¹¹⁰ Université libre de Bruxelles, 1050 Bruxelles, Belgium
- ¹¹¹ Bar-Ilan University, Ramat Gan, 5290002, Israel
- ¹¹² Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Artemis, F-06304 Nice, France
- ¹¹³ University of British Columbia, Vancouver, BC V6T 1Z4, Canada
- ¹¹⁴ OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
- ¹¹⁵ Centre national de la recherche scientifique, 75016 Paris, France
- ¹¹⁶ Univ Rennes, CNRS, Institut FOTON - UMR 6082, F-35000 Rennes, France
- ¹¹⁷ University of Birmingham, Birmingham B15 2TT, United Kingdom
- ¹¹⁸ Washington State University, Pullman, WA 99164, USA
- ¹¹⁹ University of Texas, Austin, TX 78712, USA
- ¹²⁰ Cornell University, Ithaca, NY 14850, USA
- ¹²¹ Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France
- ¹²² Christopher Newport University, Newport News, VA 23606, USA
- ¹²³ OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia
- ¹²⁴ Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
- ¹²⁵ University of Maryland, College Park, MD 20742, USA
- ¹²⁶ Università degli Studi di Milano-Bicocca, I-20126 Milano, Italy
- ¹²⁷ INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy
- ¹²⁸ University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom
- ¹²⁹ Université de Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France
- ¹³⁰ University of Chicago, Chicago, IL 60637, USA
- ¹³¹ University of Arizona, Tucson, AZ 85721, USA
- ¹³² INFN, Sezione di Napoli, Gruppo Collegato di Salerno, I-80126 Napoli, Italy
- ¹³³ University of Massachusetts Dartmouth, North Dartmouth, MA 02747, USA
- ¹³⁴ Niels Bohr Institute, Copenhagen University, 2100 København, Denmark
- ¹³⁵ Universidad de Guadalajara, 44430 Guadalajara, Jalisco, Mexico
- ¹³⁶ Istituto di Astrofisica e Planetologia Spaziali di Roma, 00133 Roma, Italy
- ¹³⁷ Colorado State University, Fort Collins, CO 80523, USA
- ¹³⁸ Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain
- ¹³⁹ Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain
- ¹⁴⁰ Niels Bohr Institute, University of Copenhagen, 2100 København, Denmark
- ¹⁴¹ National Central University, Taoyuan City 320317, Taiwan
- ¹⁴² National Tsing Hua University, Hsinchu City 30013, Taiwan
- ¹⁴³ OzGrav, Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
- ¹⁴⁴ Vanderbilt University, Nashville, TN 37235, USA
- ¹⁴⁵ University of the Chinese Academy of Sciences / International Centre for Theoretical Physics Asia-Pacific, Beijing 100049, China
- ¹⁴⁶ Department of Electrophysics, National Yang Ming Chiao Tung University, 101 Univ. Street, Hsinchu, Taiwan
- ¹⁴⁷ Kamioka Branch, National Astronomical Observatory of Japan, 238 Higashi-Mozumi, Kamioka-cho, Hida City, Gifu 506-1205, Japan
- ¹⁴⁸ CaRT, California Institute of Technology, Pasadena, CA 91125, USA
- ¹⁴⁹ Northeastern University, Boston, MA 02115, USA
- ¹⁵⁰ Dipartimento di Ingegneria Industriale (DIIN), Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ¹⁵¹ Faculty of Science, University of Toyama, 3190 Gofuku, Toyama City, Toyama 930-8555, Japan
- ¹⁵² Carleton College, Northfield, MN 55057, USA
- ¹⁵³ University of Szeged, Dóm tér 9, Szeged 6720, Hungary
- ¹⁵⁴ OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia
- ¹⁵⁵ INFN Cagliari, Physics Department, Università degli Studi di Cagliari, Cagliari 09042, Italy
- ¹⁵⁶ Università degli Studi di Cagliari, Via Università 40, 09124 Cagliari, Italy
- ¹⁵⁷ Université Libre de Bruxelles, Brussels 1050, Belgium
- ¹⁵⁸ INAF, Osservatorio Astronomico di Brera sede di Merate, I-23807 Merate, Lecco, Italy
- ¹⁵⁹ Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain

- ¹⁶⁰ Montana State University, Bozeman, MT 59717, USA
¹⁶¹ The University of Utah, Salt Lake City, UT 84112, USA
¹⁶² Johns Hopkins University, Baltimore, MD 21218, USA
¹⁶³ University of Rhode Island, Kingston, RI 02881, USA
¹⁶⁴ The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
¹⁶⁵ Université de Liège, B-4000 Liège, Belgium
¹⁶⁶ DIFA- Alma Mater Studiorum Università di Bologna, Via Zamboni, 33 - 40126 Bologna, Italy
¹⁶⁷ Istituto Nazionale Di Fisica Nucleare - Sezione di Bologna,
viale Carlo Bertini Pichat 6/2 - 40127 Bologna, Italy
¹⁶⁸ University of Manitoba, Winnipeg, MB R3T 2N2, Canada
¹⁶⁹ INFN-CNAF - Bologna, Viale Carlo Bertini Pichat, 6/2, 40127 Bologna BO, Italy
¹⁷⁰ Università degli Studi di Sassari, I-07100 Sassari, Italy
¹⁷¹ INFN, Laboratori Nazionali del Sud, I-95125 Catania, Italy
¹⁷² Université de Normandie, ENSICAEN, UNICAEN,
CNRS/IN2P3, LPC Caen, F-14000 Caen, France
¹⁷³ Laboratoire de Physique Corpusculaire Caen, 6 boulevard du maréchal Juin, F-14050 Caen, France
¹⁷⁴ The University of Sheffield, Sheffield S10 2TN, United Kingdom
¹⁷⁵ Université Claude Bernard Lyon 1, CNRS, Laboratoire des Matériaux Avancés (LMA),
IP2I Lyon / IN2P3, UMR 5822, F-69622 Villeurbanne, France
¹⁷⁶ Università di Firenze, Sesto Fiorentino I-50019, Italy
¹⁷⁷ IGFAE, Universidad de Santiago de Compostela, E-15782 Santiago de Compostela, Spain
¹⁷⁸ Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy
¹⁷⁹ INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy
¹⁸⁰ California State University, Los Angeles, Los Angeles, CA 90032, USA
¹⁸¹ Marquette University, Milwaukee, WI 53233, USA
¹⁸² Perimeter Institute, Waterloo, ON N2L 2Y5, Canada
¹⁸³ Corps des Mines, Mines Paris, Université PSL, 60 Bd Saint-Michel, 75272 Paris, France
¹⁸⁴ Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy
¹⁸⁵ Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Lagrange, F-06304 Nice, France
¹⁸⁶ National Center for Nuclear Research, 05-400 Świerk-Otwock, Poland
¹⁸⁷ Vrije Universiteit Brussel, 1050 Brussel, Belgium
¹⁸⁸ University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
¹⁸⁹ Universidad de Antioquia, Medellín, Colombia
¹⁹⁰ Canadian Institute for Theoretical Astrophysics,
University of Toronto, Toronto, ON M5S 3H8, Canada
¹⁹¹ Stony Brook University, Stony Brook, NY 11794, USA
¹⁹² Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA
¹⁹³ Montclair State University, Montclair, NJ 07043, USA
¹⁹⁴ HUN-REN Institute for Nuclear Research, H-4026 Debrecen, Hungary
¹⁹⁵ Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India
¹⁹⁶ Centro de Física das Universidades do Minho e do Porto,
Universidade do Minho, PT-4710-057 Braga, Portugal
¹⁹⁷ Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France
¹⁹⁸ CNR-SPIN, I-84084 Fisciano, Salerno, Italy
¹⁹⁹ Scuola di Ingegneria, Università della Basilicata, I-85100 Potenza, Italy
²⁰⁰ Western Washington University, Bellingham, WA 98225, USA
²⁰¹ SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
²⁰² Barry University, Miami Shores, FL 33168, USA
²⁰³ Eötvös University, Budapest 117, Hungary
²⁰⁴ Institute for Cosmic Ray Research, KAGRA Observatory, The University of Tokyo,
5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan
²⁰⁵ Department of Physics, Graduate School of Science,
Osaka Metropolitan University, 3-3-138 Sugimoto-cho,
Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
²⁰⁶ University of Sannio at Benevento, I-82100 Benevento,
Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy
²⁰⁷ University of California, Berkeley, CA 94720, USA
²⁰⁸ Instituto de Física Teórica UAM-CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain
²⁰⁹ Laboratoire d'Acoustique de l'Université du Mans, UMR CNRS 6613, F-72085 Le Mans, France
²¹⁰ University of Southampton, Southampton SO17 1BJ, United Kingdom
²¹¹ University of California, Riverside, Riverside, CA 92521, USA
²¹² Dipartimento di Ingegneria Industriale, Elettronica e Meccanica,
Università degli Studi Roma Tre, I-00146 Roma, Italy
²¹³ University of Nevada, Las Vegas, Las Vegas, NV 89154, USA

- ²¹⁴ University of Nottingham NG7 2RD, UK
- ²¹⁵ Ariel University, Ramat HaGolan St 65, Ari'el, Israel
- ²¹⁶ The University of Mississippi, University, MS 38677, USA
- ²¹⁷ Graduate School of Science, Institute of Science Tokyo,
2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan
- ²¹⁸ Institute of Physics, Academia Sinica, 128 Sec. 2, Academia Rd., Nankang, Taipei 11529, Taiwan
- ²¹⁹ The Chinese University of Hong Kong, Shatin, NT, Hong Kong
- ²²⁰ American University, Washington, DC 20016, USA
- ²²¹ Dipartimento di Fisica, Università degli studi di Milano, Via Celoria 16, I-20133, Milano, Italy
- ²²² INFN, sezione di Milano, Via Celoria 16, I-20133, Milano, Italy
- ²²³ Department of Applied Physics, Fukuoka University,
8-19-1 Nanakuma, Jonan, Fukuoka City, Fukuoka 814-0180, Japan
- ²²⁴ University of Cambridge, Cambridge CB2 1TN, United Kingdom
- ²²⁵ University of Lancaster, Lancaster LA1 4YW, United Kingdom
- ²²⁶ College of Industrial Technology, Nihon University,
1-2-1 Izumi, Narashino City, Chiba 275-8575, Japan
- ²²⁷ Faculty of Engineering, Niigata University, 8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan
- ²²⁸ Department of Physics, Tamkang University, No. 151,
Yingzhuan Rd., Danshui Dist., New Taipei City 25137, Taiwan
- ²²⁹ Rutherford Appleton Laboratory, Didcot OX11 0DE, United Kingdom
- ²³⁰ Department of Physical Sciences, Aoyama Gakuin University,
5-10-1 Fuchinobe, Sagamihara City, Kanagawa 252-5258, Japan
- ²³¹ Helmut Schmidt University, D-22043 Hamburg, Germany
- ²³² Nambu Yoichiro Institute of Theoretical and Experimental Physics (NITEP),
Osaka Metropolitan University, 3-3-138 Sugimoto-cho,
Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
- ²³³ Directorate of Construction, Services & Estate Management, Mumbai 400094, India
- ²³⁴ Observatoire Astronomique de Strasbourg, 11 Rue de l'Université, 67000 Strasbourg, France
- ²³⁵ Faculty of Physics, University of Białystok, 15-245 Białystok, Poland
- ²³⁶ National Astronomical Observatories, Chinese Academic of Sciences,
20A Datun Road, Chaoyang District, Beijing, China
- ²³⁷ School of Astronomy and Space Science, University of Chinese Academy of Sciences,
20A Datun Road, Chaoyang District, Beijing, China
- ²³⁸ Sungkyunkwan University, Seoul 03063, Republic of Korea
- ²³⁹ Department of Physics, Ulsan National Institute of Science and Technology (UNIST),
50 UNIST-gil, Ulju-gun, Ulsan 44919, Republic of Korea
- ²⁴⁰ Institute for Cosmic Ray Research, The University of Tokyo,
5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8582, Japan
- ²⁴¹ Chung-Ang University, Seoul 06974, Republic of Korea
- ²⁴² University of Washington Bothell, Bothell, WA 98011, USA
- ²⁴³ Laboratoire de Physique et de Chimie de l'Environnement,
Université Joseph KI-ZERBO, 9GH2+3V5, Ouagadougou, Burkina Faso
- ²⁴⁴ Ewha Womans University, Seoul 03760, Republic of Korea
- ²⁴⁵ National Institute for Mathematical Sciences, Daejeon 34047, Republic of Korea
- ²⁴⁶ Korea Astronomy and Space Science Institute, Daejeon 34055, Republic of Korea
- ²⁴⁷ Department of Astronomy and Space Science, Chungnam National University,
9 Daehak-ro, Yuseong-gu, Daejeon 34134, Republic of Korea
- ²⁴⁸ Institute of Particle and Nuclear Studies (IPNS),
High Energy Accelerator Research Organization (KEK),
1-1 Oho, Tsukuba City, Ibaraki 305-0801, Japan
- ²⁴⁹ Division of Science, National Astronomical Observatory of Japan,
2-21-1 Osawa, Mitaka City, Tokyo 181-8588, Japan
- ²⁵⁰ Nagoya University, Nagoya, 464-8601, Japan
- ²⁵¹ Department of Physics, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece
- ²⁵² Bard College, Annandale-On-Hudson, NY 12504, USA
- ²⁵³ Technical University of Braunschweig, D-38106 Braunschweig, Germany
- ²⁵⁴ Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland
- ²⁵⁵ Astronomical Observatory, Jagiellonian University, 31-007 Cracow, Poland
- ²⁵⁶ Department of Physics and Astronomy, University of Padova, Via Marzolo, 8-35151 Padova, Italy
- ²⁵⁷ Sezione di Padova, Istituto Nazionale di Fisica Nucleare (INFN), Via Marzolo, 8-35131 Padova, Italy
- ²⁵⁸ Department of Physics, Nagoya University, ES building,
Furocho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
- ²⁵⁹ Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada
- ²⁶⁰ Indian Institute of Science Education and Research, Kolkata, Mohanpur, West Bengal 741252, India

- ²⁶¹ *Seoul National University, Seoul 08826, Republic of Korea*
- ²⁶² *Department of Computer Simulation, Inje University,
197 Inje-ro, Gimhae, Gyeongsangnam-do 50834, Republic of Korea*
- ²⁶³ *NAVIER, École des Ponts, Univ Gustave Eiffel, CNRS, Marne-la-Vallée, France*
- ²⁶⁴ *Gravitational Wave Science Project, National Astronomical
Observatory of Japan (NAOJ), Mitaka City, Tokyo 181-8588, Japan*
- ²⁶⁵ *Department of Physics, National Cheng Kung University,
No.1, University Road, Tainan City 701, Taiwan*
- ²⁶⁶ *St. Thomas University, Miami Gardens, FL 33054, USA*
- ²⁶⁷ *Scuola Normale Superiore, I-56126 Pisa, Italy*
- ²⁶⁸ *Institució Catalana de Recerca i Estudis Avançats, E-08010 Barcelona, Spain*
- ²⁶⁹ *Institut de Física d'Altes Energies, E-08193 Barcelona, Spain*
- ²⁷⁰ *Institut fuer Theoretische Astrophysik, Zentrum fuer Astronomie Heidelberg,
Universitaet Heidelberg, Albert Ueberle Str. 2, 69120 Heidelberg, Germany*
- ²⁷¹ *Institucio Catalana de Recerca i Estudis Avançats (ICREA),
Passeig de Lluís Companys, 23, 08010 Barcelona, Spain*
- ²⁷² *Research Center for Space Science, Advanced Research Laboratories, Tokyo City University,
3-3-1 Ushikubo-Nishi, Tsuzuki-Ku, Yokohama, Kanagawa 224-8551, Japan*
- ²⁷³ *Tsinghua University, Beijing 100084, China*
- ²⁷⁴ *Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France*
- ²⁷⁵ *Faculty of Law, Ryukoku University, 67 Fukakusa Tsukamoto-cho,
Fushimi-ku, Kyoto City, Kyoto 612-8577, Japan*
- ²⁷⁶ *Phenikaa Institute for Advanced Study (PIAS),
Phenikaa University, Yen Nghia, Ha Dong, Hanoi, Vietnam*
- ²⁷⁷ *University of Stavanger, 4021 Stavanger, Norway*
- ²⁷⁸ *Physics Program, Graduate School of Advanced Science and Engineering,
Hiroshima University, 1-3-1 Kagamiyama, Higashihiroshima City, Hiroshima 739-8526, Japan*
- ²⁷⁹ *GRAPPA, Anton Pannekoek Institute for Astronomy and Institute for High-Energy Physics,
University of Amsterdam, 1098 XH Amsterdam, Netherlands*
- ²⁸⁰ *University College London, London WC1E 6BT, United Kingdom*
- ²⁸¹ *Observatoire de Paris, 75014 Paris, France*
- ²⁸² *Laboratoire Univers et Théories, Observatoire de Paris, 92190 Meudon, France*
- ²⁸³ *Graduate School of Science and Technology, Niigata University,
8050 Ikarashi-2-no-cho, Nishi-ku, Niigata City, Niigata 950-2181, Japan*
- ²⁸⁴ *University of Maryland, Baltimore County, Baltimore, MD 21250, USA*
- ²⁸⁵ *CSIR-Central Glass and Ceramic Research Institute, Kolkata, West Bengal 700032, India*
- ²⁸⁶ *Consiglio Nazionale delle Ricerche - Istituto dei Sistemi Complessi, I-00185 Roma, Italy*
- ²⁸⁷ *Department of Astronomy, Yonsei University,
50 Yonsei-Ro, Seodaemun-Gu, Seoul 03722, Republic of Korea*
- ²⁸⁸ *Department of Physics, University of Guadalajara, Av. Revolucion 1500,
Colonia Olimpica C.P. 44430, Guadalajara, Jalisco, Mexico*
- ²⁸⁹ *Hobart and William Smith Colleges, Geneva, NY 14456, USA*
- ²⁹⁰ *INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy*
- ²⁹¹ *Dipartimento di Ingegneria, Università del Sannio, I-82100 Benevento, Italy*
- ²⁹² *Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi", I-00184 Roma, Italy*
- ²⁹³ *Kennesaw State University, Kennesaw, GA 30144, USA*
- ²⁹⁴ *Subatech, CNRS/IN2P3 - IMT Atlantique - Nantes Université,
4 rue Alfred Kastler BP 20722 44307 Nantes CÉDEX 03, France*
- ²⁹⁵ *Departamento de Física - ETSIDI, Universidad Politécnica de Madrid, 28012 Madrid, Spain*
- ²⁹⁶ *Department of Electronic Control Engineering,
National Institute of Technology, Nagaoka College,
888 Nishikatai, Nagaoka City, Niigata 940-8532, Japan*
- ²⁹⁷ *Trinity College, Hartford, CT 06106, USA*
- ²⁹⁸ *Dipartimento di Fisica e Scienze della Terra, Università Degli Studi di Ferrara, Via Saragat, 1, 44121 Ferrara FE, Italy*
- ²⁹⁹ *Faculty of Science, Toho University, 2-2-1 Miyama, Funabashi City, Chiba 274-8510, Japan*
- ³⁰⁰ *Indian Institute of Technology, Palaj, Gandhinagar, Gujarat 382355, India*
- ³⁰¹ *Kavli Institute for Astronomy and Astrophysics, Peking University,
Yiheyuan Road 5, Haidian District, Beijing 100871, China*
- ³⁰² *Laboratoire MSME, Cité Descartes, 5 Boulevard Descartes,
Champs-sur-Marne, 77454 Marne-la-Vallée Cedex 2, France*
- ³⁰³ *Faculty of Information Science and Technology, Osaka Institute of Technology,
1-79-1 Kitayama, Hirakata City, Osaka 573-0196, Japan*
- ³⁰⁴ *NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*

- ³⁰⁵ *Faculty of Science and Technology, Kochi University,
2-5-1 Akebono-cho, Kochi-shi, Kochi 780-8520, Japan*
- ³⁰⁶ *Laboratoire de Physique de l'École Normale Supérieure, ENS, (CNRS,
Université PSL, Sorbonne Université, Université Paris Cité), F-75005 Paris, France*
- ³⁰⁷ *Faculty of Physics, University of Warsaw, Ludwika Pasteura 5, 02-093 Warszawa, Poland*
- ³⁰⁸ *Laser Interferometry and Gravitational Wave Astronomy,
Max Planck Institute for Gravitational Physics, Callinstrasse 38, 30167 Hannover, Germany*
- ³⁰⁹ *The Hakubi Center for Advanced Research, Kyoto University,
Yoshida-honmachi, Sakyou-ku, Kyoto City, Kyoto 606-8501, Japan*
- ³¹⁰ *Department of Physics, Kyoto University, Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan*
- ³¹¹ *Yukawa Institute for Theoretical Physics (YITP), Kyoto University,
Kita-Shirakawa Oiwake-cho, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan*
- ³¹² *University of Catania, Department of Physics and Astronomy, Via S. Sofia, 64, 95123 Catania CT, Italy*
- ³¹³ *National Institute of Technology, Fukui College, Geshi-cho, Sabae-shi, Fukui 916-8507, Japan*
- ³¹⁴ *Department of Communications Engineering, National Defense Academy of Japan,
1-10-20 Hashirimizu, Yokosuka City, Kanagawa 239-8686, Japan*
- ³¹⁵ *Eindhoven University of Technology, 5600 MB Eindhoven, Netherlands*
- ³¹⁶ *Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), WPI,
The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba 277-8583, Japan*
- ³¹⁷ *Department of Astronomy, Beijing Normal University,
Xiniekouwai Street 19, Haidian District, Beijing 100875, China*
- ³¹⁸ *School of Physics and Technology, Wuhan University,
Bayi Road 299, Wuchang District, Wuhan, Hubei, 430072, China*
- (Dated: August 5, 2025)

We present a search for short-duration gravitational-wave transients in data from the first eight months of Advanced LIGO-Virgo-KAGRA's fourth observing run, denoted O4a. We use four analyses which are sensitive to a wide range of potential signals lasting up to a few seconds in the 16–4096 Hz band. Excluding binary black hole merger candidates that were already identified by low-latency analyses, we find no statistically significant evidence for other gravitational-wave transients. We measure the sensitivity of the search for representative signals, including sine-Gaussians, Gaussian pulses, and white-noise bursts with different frequencies and durations, adopting a false alarm rate of 1 per 100 years as detection threshold. Depending on signal type, we find improvements over previous searches by factors of 2 to 10 in terms of sensitivity to strain amplitude and of 90% confidence upper limit on the rate density of sources. We also evaluate a variety of core-collapse supernova models and find that, for some models, the search could have detected gravitational waves from stellar core-collapse throughout the Milky Way. Finally, we consider neutron star f -modes associated with pulsar glitches and find that, assuming a source similar to the Vela Pulsar, the search could have detected a gravitational-wave signal from a glitch with fractional frequency change as small as ~ 2 to 6×10^{-5} depending on the neutron star mass.

I. INTRODUCTION

The Advanced LIGO [1] and Advanced Virgo [2] detectors have proven the ability of large laser interferometers to detect and characterize gravitational waves. Using data from the first three observing runs of these detectors, the LIGO Scientific Collaboration, Virgo Collaboration and KAGRA Collaboration (collectively *LVK*) identified 90 transient signals most likely to be gravitational-wave (GW) events [3–5]. Other researchers have reported several additional GW event candidates in the publicly available data [6–14]. All of the GW signals detected in those runs have matched the distinctive signature expected from compact binary coalescence (CBC) events. A CBC arises from a co-orbiting pair of compact astrophysical objects, specifically black holes or neutron stars, that spiral inward and finally merge due to emission of

gravitational radiation as predicted by the general theory of relativity.

Over 200 additional CBC candidates have been reported from low-latency analyses during the fourth LVK observing run [15]. Many more CBC events will be detected as the LIGO and Virgo detectors improve and as the GW detector network grows with the maturation of the KAGRA detector [16] and construction of a LIGO observatory in India [17, 18] as well as next-generation facilities currently being designed [19, 20]. In parallel, the LVK continues to search for other types of GW signals. For example, transient GWs—also called *GW bursts*—can arise from the strong dynamics of mass and energy in core-collapse supernovae [21–25], magnetars [26–28], *starquakes* of neutron stars [29, 30], non-linear memory [31], hyperbolic encounters between compact objects [32], and cosmic strings [33–35]. The time-frequency structure, the strength and the rate of occurrence of GW transient events depend on the source's internal microphysics, formation channels and population

* Deceased, September 2024.

models, which are not fully understood. It is also possible that standard searches for CBC events, which use matched filtering with modeled waveforms as templates, may miss some binary mergers with very high masses, eccentric orbits and/or extreme spins, which are difficult to detect efficiently with template banks currently in use. Therefore it is essential to search as broadly as possible for GW signals from potential sources, known or unknown.

Search algorithms which do not rely on having modeled waveforms have been developed and refined to meet this need. Different methods are employed to search for generic short-duration GW burst signals (up to a few seconds long) and longer-duration transient signals over the full frequency range in which the detectors are highly sensitive. Generic searches for short-duration [36] and long-duration [37] GW bursts in LIGO-Virgo data from the third observing run (O3) uncovered no significant candidates, other than already-known CBC events, despite having good sensitivity for a wide range of other potential signals. These searches are complemented by dedicated analyses for GW signals from intermediate mass black hole (IMBH) mergers [38] and compact binary systems with significantly eccentric orbits [39], which obtain better sensitivity for those signal classes by making morphological assumptions specific to each.

In this paper we report on a search for short-duration GW bursts in data from the LIGO Hanford and LIGO Livingston Observatories (LHO and LLO, respectively) taken during the first eight months of the fourth observing run (O4), a period denoted O4a. (A companion paper [40] reports on a search for longer-duration transient GW signals.) We give more details about the detectors, the O4a run period and data quality in section II. In section III we describe the three search *pipelines* used in this paper. All three pipelines are based on the Coherent WaveBurst algorithm [41, 42] but use different configurations, time-frequency transforms, and candidate event classification algorithms to obtain a broad coverage of the signal parameter space.

As we report in section IV, this search finds many of the likely CBC signals that have already been reported publicly from low-latency analysis (and which will be verified and discussed in later LVK papers) but our goal here is to reveal any short-duration transient GW signal that is *not* a CBC event. We find no statistically significant candidate when the likely CBC signals are excluded. In section V we evaluate and compare the sensitivities of the analyses using simulations which sample a wide range of potential signals in the detectors' sensitive frequency range. The O4a run was only about half as long in duration as O3, but due to lower noise in the LIGO detectors and some analysis improvements, this early look at the O4 run covers a larger astrophysical volume-time exposure than O3 by a factor of 2 to ~ 10 , depending on the signal morphology.

In section VI we evaluate the sensitivity of this search to modeled signals from core-collapse supernovae and

sudden excitations of neutron stars, providing astrophysical interpretation of our null search results. This *all-sky search* does not rely on observing a supernova light curve or neutron star pulse timing glitch, so it complements multi-messenger searches that are triggered by electromagnetic or neutrino observations, e.g. [43]. We end with some remarks about the ongoing O4 run and prospects for future detections.

II. DETECTORS AND DATA

A. Detectors Operating During O4a

The LIGO detectors underwent significant upgrades and commissioning between the conclusion of the O3 observing run on March 27, 2020 and the beginning of the O4 run on May 24, 2023 [44, 45]. Those efforts reduced the detector noise at almost all frequencies, as shown in Figure 1. The improvement at low frequencies is due to realizing frequency-dependent squeezing [46] and reducing sources of environmental and technical noise, while the improvement at high frequencies is mainly due to increasing the laser power input into the interferometers to 60–75 W from 37 W in O3 [44]. The reduced detector noise enables more sensitive searches for GW signals, and more stringent limits on source populations and GW emission mechanisms from searches that find no significant candidates.

In addition to the LIGO detectors, the KAGRA GW observatory participated in O4a from the beginning until June 20, 2023, at which point it was taken offline to conduct commissioning activities. KAGRA data are not used in this work due to its much lower sensitivity. The Virgo observatory was being upgraded during all of O4a. Virgo resumed joint observation with the LIGO detectors in April 2024 for the continuation of the O4 run.

B. O4a Data

O4a lasted from May 24, 2023 15:00 UTC to January 16, 2024 16:00 UTC. At least one LIGO facility was observing for 83.1% of this period. To carry out the search for GW signals, the four analyses described in section III are applied to coincident data of high quality (see section II C) from both LIGO detectors. Additionally, the analyses require a minimum of 200 seconds of contiguous observing-mode data and discard the first and last 4 or 10 seconds from each period of observing time in each detector. These restrictions reduce the dataset to 126.4 days of dual-coincident, high quality observing-mode data from the LIGO detectors in O4a, representing 53.3% of the total duration of the O4a run.

The LIGO detectors employ photon calibrators to determine the observed GW strain based on fiducial displacements of the end test masses due to radiation pressure from an auxiliary laser system [47, 48]. The




FIG. 1. A comparison of GW detector noise between observatories and observing runs. Top: amplitude spectral densities (ASDs) of the GW strain noise for the LHO (red) and LLO (blue) detectors in the O4 run compared to the O3 run (gray). Virgo’s and KAGRA’s strain sensitivities are not plotted. Bottom: the ratio between each LIGO detector’s O3 and O4 noise ASD as a function of frequency. A larger value indicates a greater reduction in noise between O3 and O4. This ratio comparison is omitted near power line harmonics and near the thermally-driven “violin” resonance modes of optics suspensions above 250 Hz.

frequency-dependent calibration uncertainty varies during the run and is assessed hour by hour. At the beginning of O4a the amplitude uncertainty is typically $\sim 5\%$ or less for LHO and $\sim 3\%$ or less for LLO for frequencies below 1 kHz, rising to $\sim 10\%$ at 4 kHz. For data collected after June 13, 2023 the LLO calibration uncertainty improves, becoming typically $\sim 2\%$ at all frequencies up to 4 kHz. After July 27, 2023 the LHO calibration uncertainty improves, becoming typically $\sim 2\%$ up to 1 kHz, rising to $\sim 5\%$ at 4 kHz.

C. Data Quality

1. Data Quality Vetoes

Detector data quality varies during observing-mode time and occasionally is poor; these periods can sometimes be identified and vetoed. In O4a, two categories of data quality (DQ) vetoes are applied [45]: category 1 vetoes flag time intervals with such severe issues that data taken during these times are not used by search pipelines, while category 2 vetoes represent times when

the data quality is poor but certain time or frequency bands may still contain usable data.

In O4a, modeled searches for gravitational waves from compact binary coalescences exclude only category 1 vetoed times, while the analyses described in this work exclude the times of both category 1 and 2 vetoes, because the lack of a distinct morphology (like the “chirp” of a CBC) for unmodeled bursts complicates the prospect of distinguishing signals from noise.

Category 1 vetoes in O4a include periods when the observing mode state at an observatory was incorrectly activated (i.e., the detectors were not truly ready to observe) and periods of elevated motion in the detector suspensions that significantly degraded the calibration of LLO. Examples of category 2 vetoes used in this work include periods when large noise transients in the GW data were witnessed by voltage monitors in the electrostatic drive system used to control the position of the test masses; these transients are temporally correlated with instrumental artifacts (*glitches*) in the GW data. Additional criteria for vetoing O4a data may be found in [45]. The total amount of observing-mode time vetoed in O4a was 1.66 days at LHO and 3.36 days at LLO.

2. Other Transient Noise

Despite the efforts to isolate the LIGO detectors from noise, they are still subject to occasional noise transients of both known and unknown origin. The most deleterious class of noise transients for this work are *blip glitches* [49]. Like many of the GW waveforms studied herein, blip glitches are short-duration, large-bandwidth transients. Their cause is uncertain. They are present in both LHO and LLO data, but at uncorrelated times. Most are not witnessed by auxiliary monitoring channels and therefore cannot be vetoed. According to Gravity Spy [50], blip glitches occurred at an average rate of 1.24 per hour at LHO and 0.87 per hour at LLO. This is a decrease from the O3 rate of 1.68 and 3.47 blip glitches per hour, respectively. Investigations to determine the cause of blip glitches have included searching for temporal correlations with vibrational transients along the LIGO beamtubes, temporal correlations with cosmic ray strikes, and correlations with environmental data monitors at each site. These investigations have so far not identified the cause of blip glitches in the LIGO detectors, although a small fraction of these glitches are contemporaneous with computer timing errors [45, 49, 51].

Especially at low frequencies, LIGO detector sensitivity can be degraded by stray light beams scattering off vibrating surfaces and interfering with the main beam, causing *scattering glitches* [52]. Prior to O4a, a campaign was conducted to identify and mitigate several sources of scattered light noise [45]. While a direct comparison between the O3 and O4a searches is difficult to make due to differences in search pipeline configurations, the incidence of large signal outliers below 128 Hz was reduced

in O4a compared to O3. This is believed to be a result of improved isolation from scattered light noise.

III. SEARCH ALGORITHMS, PIPELINES, AND ANALYSES

The search algorithms used in this paper are based on the open-source software Coherent WaveBurst (cWB) [41, 42, 53], which is designed to operate without a specific signal model. cWB begins by computing multi-resolution time-frequency (TF) representations of the strain data from the GW detector network [54, 55]. The multi-resolution approach is essential to cover the wide parameter space of potential signals with durations ranging from milliseconds to a few seconds in a frequency range from tens of Hz up to a few kHz. cWB selects the most energetic TF regions, which are referred to as *triggers* if they pass internal pipeline thresholds, and evaluates how potential GW signals would appear in the network of detectors. For each trigger, a likelihood ratio statistic is maximized over all sky directions [56] enabling cWB to reconstruct the source sky localization, the signal waveforms, and morphological properties of the signals such as the duration, the central frequency, etc. The ranking statistic η is essentially the signal-to-noise ratio (SNR) of the trigger, estimated from the coherent response of the network, with a penalty factor related to the incoherent energy residuals in the detectors. The trigger significance is measured in terms of inverse false alarm rate (IFAR), computed by comparing the ranking statistic of each trigger with a *background* distribution. The latter is built by repeating the cWB algorithm on time-shifted data, shifting one detector with respect to the other detector by an amount that breaks any coherence from astrophysical signals. Previous versions of cWB were used to search and reconstruct GW transients in past LVK observing runs (O1 [57, 58], O2 [59, 60], and O3 [36, 37]).

To maximize the discovery potential of our search, we employ two updated versions of cWB to generate triggers, with shorthand names derived from *second-generation* (cWB-2G) and *cross-power* (cWB-XP) development paths, presented in Section III A. After the generation of the triggers, two different machine-learning procedures—a decision-tree algorithm called XGBoost and a Gaussian mixture model (GMM)—classify the triggers and enhance the separation between possible astrophysical transients and detector noises, as described in Section III B.

The total number of search pipelines applied to the O4a dataset is three, illustrated in Figure 2: two use the triggers generated by the cWB-2G algorithm and apply XGBoost (2G+XGB) and GMM (2G+GMM) classification algorithms, respectively, while the third generates triggers with the cWB-XP algorithm and then applies XGBoost (XP+XGB). All three pipelines operate in the frequency band where the detectors are most sensitive,




FIG. 2. The GW burst search reported in this paper uses two algorithms to identify triggers, i.e. coherent excess of power, described in Section III A, and two machine-learning classification algorithms to suppress background, described in Section III B. These algorithm components are combined to construct three *pipelines*: 2G+XGB, XP+XGB, and 2G+GMM. The 2G+XGB pipeline is used in two *analyses*, one for low-frequency signals and the other for high-frequency signals, while each of the other pipelines is used in a single analysis covering a similar low-frequency range. The complementary aspects of these four analyses are discussed in the text.

up to 2048 Hz, aiming to maximize their combined detection efficiency in this parameter space. Although the pipelines perform similarly on average, as shown in Section V, their detection efficiency varies across different waveform morphologies, signal amplitudes, and polarizations. They also respond differently to non-stationary noise. Our simulation studies reveal that each pipeline detects some signals that the others miss. Additionally, the 2G+XGB pipeline performs a dedicated analysis for high-frequency signals, so a total of four *analyses* make up the search reported in this paper.

The 2G+XGB and XP+XGB pipelines have also been used to search for generic GW transients in low latency while the GW detectors were collecting data. The main difference between the configurations used for low-latency analyses and the offline analyses presented here is in the training of the machine-learning algorithms used to classify candidate events versus transient noise (Section III B 1).

A. Identify triggers showing coherent excess-power

The Coherent WaveBurst algorithm has been successfully employed in previous GW burst searches, e.g. [36]. In the following we provide details about the two upgraded versions of cWB employed to generate triggers from the GW strain data.

1. Coherent WaveBurst 2G

The cWB-2G algorithm computes the TF representation with the Wilson-Daubechies-Meyer wavelet transform [54], using multiple TF resolutions to adapt to different signal morphologies. The cWB-2G algorithm improves on the selection of TF pixels that form trigger candidates to improve the sensitivity to low SNR events. After selection, TF pixels are aggregated if they are separated by time and frequency intervals of less than 0.2 s and 640 Hz; this improves the collection of energy for transient events with complex time-frequency structure that appears as separate clusters of TF pixels. cWB-2G is used in two configurations with different sets of TF resolutions to perform low-frequency (16–2028 Hz) and high-frequency (512–4096) analyses. High-frequency analysis triggers with central frequency below 1600 Hz are removed, leaving a 448 Hz overlap with the low-frequency analysis to ensure good coverage of wide-band GW events at intermediate frequencies. A trials factor of 2 is applied to the IFAR of triggers in this overlap region, accounting for the fact that they could be found by both low- and high-frequency analyses.

2. Coherent WaveBurst Cross Power

The cWB-XP algorithm implements two major changes with respect to the cWB version used in the O3 search [36]. First, the Wilson-Daubechies-Meyer wavelet transform is replaced with the multi-resolution *WaveScan* transform based on Gabor wavelets [55] in order to reduce temporal and spectral leakage in the TF representation. For O4a, cWB-XP analyzes the frequency range between 32 Hz and 2048 Hz. After selection, the initial TF clusters are aggregated if they are separated by time and frequency intervals of less than 0.23 s and 64 Hz, respectively. A coherent SNR > 7 threshold is applied to aggregated clusters, resulting in a trigger rate of $O(0.01 \text{ Hz})$. Second, in addition to the excess-power statistic used previously in cWB, cWB-XP applies a cross-power statistic [55] to identify transient events and suppress the noise transient rate. The cross-power amplitude of each TF data sample (or *WaveScan* pixel) is maximized over all possible time-of-flight delays of a GW signal in the detector network. Both statistics follow a predictable half-normal distribution with unit variance expected for quasi-stationary detector noise and are described in Refs. [55, 61].

B. Classify candidate events versus background

The output rate of both cWB-2G and cWB-XP algorithms is dominated by triggers originating from non-stationary detector noise. To increase the confidence of detections, we have replaced the manual tuning of selection thresholds, used through O3 [36], with procedures

exploiting two machine-learning algorithms.

1. Decision tree learning with XGBoost

XGBoost is a supervised decision tree learning algorithm [62] employed to classify cWB-2G and cWB-XP triggers between a noise class and a signal class. XGBoost receives as input tabular values from the two labelled classes and, during the training process, builds a multi-dimensional model which outputs a number ranging from 0, for triggers supposed to belong to the noise class, to 1 for the signal class.

The training dataset uses unmodified background triggers for the noise class, while the signal class is populated with generic white-noise burst (WNB) injections selected to cover the full parameter space of the search [63]. The tabular values used to train the model are a subset of summary features (10 for 2G+XGB and 16 for XP+XGB) that do not depend directly on the morphological characteristics of anticipated signals, in order to maintain the unmodeled nature of the search. The selected summary statistics describe the degree of correlation of the trigger in the detector network, the energy that is coherent and not-coherent in the network, the energy in each TF resolution, and attributes measuring the likeness to known single-cycle glitches. The detailed list of summary features used by 2G+XGB can be found in Ref. [63]. XP+XGB includes in the training also summary statistics quantifying the cross-power and the network alignment factor [61]. In the low-frequency region, the WNB injections used to train the XGBoost model have duration uniformly distributed in $[0.001, 0.5]$ s, central frequency in $[24, 1696]$ Hz, and bandwidth in $[10, 800]$ Hz. In the high-frequency region (512–4096 Hz), the WNB injections used to train the XGBoost model have duration uniformly distributed in $[0.0001, 0.5]$ s, central frequency in $[1200, 3400]$ Hz, and bandwidth in $[100, 1600]$ Hz.

The 2G+XGB and XP+XGB low-latency analyses update their XGBoost model when the properties of the noise distribution change substantially. The WNB simulations and background triggers used to train the models are acquired continuously during the observing run.

For the offline search presented here, cWB-2G triggers are processed by XGBoost models trained on data from limited periods of time. The full dataset is divided into data segments, referred to as chunks, of about 9 days each. For each chunk, an XGBoost model is trained on background and WNB injection triggers acquired on the two adjacent chunks (both sides, apart from end chunks) plus an additional 10% of the same chunk's triggers. The remaining 90% background triggers of the chunk are used to estimate the statistical significance of candidates. This training procedure adapts well to non-stationary properties of the data, and at the same time prevents model overtuning. Once trained, the XGBoost output W_{XGB} is stretched monotonically to enhance sensitivity for events

with W_{XGB} close to unity and used as the ranking statistic for cWB-2G triggers.

In contrast, the classification stage for cWB-XP uses a single XGBoost model trained on data from the entire run along with the simulated WNBs. Approximately 20% of the background triggers are used for training, while the remaining background triggers are used to estimate the statistical significance of candidates. The ranking statistic is the one described in [63, 64]:

$$\eta_r = \eta_0 \cdot W'_{\text{XGB}}, \quad (1)$$

where W'_{XGB} is computed from the output of the XGBoost model [65], including a monotonic stretch and a correction to mitigate the impact of loud blip-like glitches. It is used as a multiplicative factor to re-weight η_0 , the cWB detection statistic that is based on the coherent energy [65].

2. Gaussian mixture model

An alternative supervised machine-learning approach, GMM, is applied to cWB-2G triggers in the 2G+GMM pipeline. GMM learns to model multi-dimensional distributions as superpositions of Gaussians. The application of GMM to enhance GW burst searches is described in Refs. [66–68]. GMM constructs two distinct models, one for the noise and one for the signal, from a subset of 11 cWB-2G trigger summary statistics which undergo re-parameterization in order to have desired Gaussian behaviour. The ranking statistic η is defined by a log-likelihood ratio $\eta = W_s - W_n$, where W_s and W_n are the maximum log-likelihood statistics derived from the signal and noise models, respectively. The training dataset includes background triggers and WNB injections as described above for the XGBoost algorithm. The 2G+GMM pipeline covers the frequency range $[16, 2048]$ Hz and analyzes data segments formed from two adjacent chunks at a time. For each segment, GMM models first use 5% of the background triggers and 20% of WNB simulations to select the optimal number of Gaussians required to efficiently model the data, then 25% of the background triggers and 80% of the WNB simulations to train the models. The remaining 70% of the background triggers are used to evaluate the statistical significance of candidates.

IV. SEARCH RESULTS

In this section we report the results of the four analyses performed using the search pipelines presented above.

While these search methods are designed to explore a wide range of possible signals, the low-frequency analyses in particular are expected to detect CBC events since most of the signal power of a CBC is at frequencies below 1 kHz. However, being agnostic on the signal morphology, GW burst searches have a higher false alarm rate

than template-based methods so their detection capability for CBC signals is somewhat lower, or comparable in the case of IMBH binaries [38].

The results of the four analyses performed are shown in Figure 3. Each panel of that figure shows the cumulative set of event candidates ordered by IFAR and compared to the straight-line expectation for false alarms. The triangles connected with dashed lines show a substantial excess in each low-frequency analysis, representing detected GW candidate events. XP+XGB accumulates about 500 years of background for each chunk, so there is a sharp decrease at IFAR around 500 years in the upper right panel in Fig. 3. In contrast, 2G+XGB and 2G+GMM accumulate different amounts of background statistics for each chunk, so there is a less-sharp edge to the cumulative number of GW candidate events in the two left panels.

The circles connected with solid lines show the distribution after all CBC candidates, identified by low-latency analyses [41, 69–73] with an IFAR larger than one month and reported in public alerts (including S231123cg which was later confirmed as GW231123 [74]), have been excised [75]. No significant individual event remains after that excision, nor any significant cumulative excess in the IFAR distributions. In the high-frequency analysis, where we do not expect to detect CBC signals, the search results are consistent with the expected distribution of false alarms.

The most significant candidate event from the low-frequency analyses is found on 2023-08-30 at 05:07:03 UTC by the 2G+GMM pipeline with IFAR of 1.75 years. The same trigger is identified by 2G+XGB with IFAR of 0.059 year, while it is not identified by XP+XGB. This candidate has an SNR of 19 concentrated at about 930 Hz. Using the method described in [76], we checked for the presence of linear coupling between the network of environmental monitoring sensors [77] and the GW strain data at each LIGO observatory. No vibrational or magnetic transients witnessed by these sensors contaminated the GW data at either site around the time of this candidate event. Although we cannot reject this candidate on the basis of data quality, its statistical significance is not high enough to be claimed as a GW event.

The second most significant low-frequency event candidate has a low significance ($\text{IFAR} \leq 0.2$ year in all analyses), so it was not investigated further.

The most significant candidate of the 2G+XGB high-frequency analysis occurs on 2023-12-23 at 18:25:36 UTC with IFAR of 0.12 year. This candidate has SNR 14.7, central frequency of 3146 Hz and duration of 0.023 s. We did not investigate this candidate further due to its low significance.

To assess the complementarity of the different algorithms, we consider candidate events found by each low-frequency analysis with $\text{IFAR} > 1$ day. As before, we exclude known CBC candidates, so that the remaining low-significance candidates are most likely due to detector noise fluctuations. There are 316 such candidates, of which 106 are found by 2G+XGB, 129 by XP+XGB,

and 137 by 2G+GMM. Comparing their times, we find that only 10 of these candidates are found by all three pipelines, with an additional 4 found by both 2G+XGB and XP+XGB, 5 by XP+XGB and 2G+GMM, and 27 by 2G+XGB and 2G+GMM. The fact that the majority of the low-significance candidates are found by only one pipeline means that the three pipelines are fairly independent in their response to detector noise fluctuations.

V. SEARCH SENSITIVITY AND RATE LIMITS

As in previous iterations of this search, we assess the sensitivity of our analyses to a variety of ad-hoc signals that cover broad ranges of frequency, duration, and bandwidth. Simulated waveforms are generated with chosen sky directions and injected into data from the network of GW detectors with the proper polarization and time delays to mimic a true GW signal. The amplitude distribution is chosen such that for each waveform and each analysis the detection efficiency goes from around 0 to 100%. The distribution in sky direction is uniform in solid angle over the whole sky. The waveform families used are Gaussian pulses (GA) with varying durations parametrized by the Gaussian standard deviation of time τ_{GA} , sine-Gaussians (SG) with varying central frequency f_0 and quality factor Q proportional to the signal duration, and band-limited white-noise bursts parameterized by a central frequency f_0 , bandwidth Δf , and duration τ_{WNB} . Further description of these waveforms can be found in [78]. The amplitude is characterized by a quantity called the root-sum-squared strain amplitude, $h_{\text{rss}} = \sqrt{\int_{-\infty}^{\infty} (h_+^2 + h_\times^2) dt}$, where h_+ and $h_\times$ are GW polarizations in TT-gauge. Analysis is done using data plus simulated signals of each waveform type with nine different amplitudes, generated in terms of h_{rss} following the rule $h_{\text{rss}} = (\sqrt{3})^N \times 5 \times 10^{-23} \text{ Hz}^{-\frac{1}{2}}$ where N is an integer ranging from 0 to 8.

The polarization distribution is waveform dependent. For the GA waveforms it is fixed to linear polarization mimicking linear motion emission. For SG waveforms, we sample different elliptical polarizations determined by an inclination angle assuming a rotating source, which leads to polarization modes

$$\begin{aligned} h_+(t) &= \frac{1}{2}(1 + \cos^2 \iota)A_+(t), \\ h_\times(t) &= \cos(\iota)A_\times(t) \end{aligned} \quad (2)$$

where ι is the inclination angle (effectively the angle between the rotation axis and the line of sight) and $A_{+, \times}(t)$ is the time-dependent amplitude given by the SG (and analogous cosine-Gaussian) waveform. The polarization distribution for SG follows a uniform distribution of $\cos(\iota)$. For WNB signals, the polarization distribution is kept random, representing isotropic emission.

We use a characteristic amplitude, $h_{\text{rss}}^{50\%}$, as a measure of detection sensitivity. $h_{\text{rss}}^{50\%}$ is the injected h_{rss} for which




FIG. 3. Cumulative number of candidate events versus IFAR found in the low-frequency analyses by 2G+XGB (top left), XP+XGB (top right), 2G+GMM (bottom left), and in the high-frequency analysis by 2G+XGB (bottom right). Triangular points indicate the results for all data, while circular points show candidates remaining after CBC candidate signals found in low latency have been excised. The solid line shows the expected mean value of the background, while the shaded regions indicate 68%, 95%, and 99% Poisson uncertainty regions.

the signal recovery efficiency reaches 50%, which we determine by fitting the efficiency tabulated at the discrete values of h_{rss} used in the simulation and interpolating from the fit. We report the results for all the waveforms at $\text{IFAR} \geq 100$ years in Table I. This IFAR detection threshold corresponds to p-value of 3×10^{-3} which is equivalent to a ~ 3 sigma detection in the O4a dataset. Compared to the corresponding O3 results, we see improvements for all the morphologies. We note that the calibration uncertainties (Sec. II B) are not taken into account here. In response to the better low-frequency sensitivity of the detectors during O4 we have introduced lower-frequency SG injections at 36 and 48 Hz. The largest improvement in $h_{\text{rss}}^{50\%}$ is for the GA waveforms, for which the O4a sensitivity is at least 6 times better than O3. This is attributed to both improvements in the search algorithms and better techniques to isolate the very short-duration and loud glitches, which anyway occur with a lower rate in O4a as compared to O3, as noted in Sec. II C 2. For signals below 2 kHz the results are similar for all three low-frequency analyses, meaning that although the pipelines respond fairly differently to transient noise outliers (section IV), the overall sensitivity of these pipelines to a broad variety of waveforms is similar. The 2G+GMM pipeline performs best for GA

waveforms and $h_{\text{rss}}^{50\%}$ is typically within 15% to 50% of the other two pipelines for SG and WNB waveforms. Beyond the fact that $h_{\text{rss}}^{50\%}$ values are similar, each pipeline recovers some number of injected signals that are not identified by the others.

We can define an astrophysically meaningful quantity to interpret $h_{\text{rss}}^{50\%}$ in terms of the energy emitted by a hypothetical source. Consider a standard candle source at fixed distance of r_0 radiating GWs in a narrow frequency range around a central frequency f_0 . The energy radiated by a rotating system like a circular binary (modeled here as an elliptically polarized SG) will be [79]

$$E_{\text{GW}}^{\text{rot}} = \frac{2}{5} \frac{\pi^2 c^3}{G} r_0^2 f_0^2 h_{\text{rss}}^2 \quad (3)$$

where G is the gravitational constant and c is the speed of light. The narrow-band approximation causes very low biases for our SG waveforms ($\leq 3\%$). We use each $h_{\text{rss}}^{50\%}$ in the above equation to obtain this energy as a function of central frequency for $r_0 = 10$ kpc as shown in Table II. As compared to the previous observing run we improve by a factor of 2.5 to 6 across the different frequencies. These improvements result from both the better sensitivity of the detectors and improvements in the search methods.

We can also compute a limit on the rate per unit volume for a hypothetical population of sources producing these waveforms. The generic source is assumed to emit $1 M_{\odot} c^2$ of GW energy where $M_{\odot}$ is the mass of the Sun. Using equation 3 with $E_{\text{GW}} = 1 M_{\odot} c^2$, the relation between the average h_{rss} at the source at distance r and on Earth is then $hr = h_0 r_0$, where we have dropped the subscript rss for brevity. Assuming an intrinsic event rate of R per unit volume per time, the expected number of detected events for an observing time T is

$$\begin{aligned} N_{\text{det}} &= 4\pi RT \int_0^{\infty} dr r^2 \epsilon(r) \\ &= 4\pi RT (h_0 r_0)^3 \int_0^{\infty} dh h^{-4} \epsilon(h) \end{aligned} \quad (4)$$

where $\epsilon(h)$ is the efficiency as a function of amplitude.

In the absence of a detection, the rate limit at 90% confidence is

$$\mathcal{R}_{90\%} = \frac{2.3}{4\pi T (h_0 r_0)^3 \int_0^{\infty} dh h^{-4} \epsilon(h)}. \quad (5)$$

We use the SG waveforms given in Table II for these computations and fix the threshold at $\text{IFAR} \geq 100$ years. Results from the best of the three low-frequency analyses, on a waveform-by-waveform basis, are shown in Figure 4. While the observing time of O4a was approximately half that of O3, the O4a analysis results in lower rate limits for elliptically polarized SG waveforms across all frequencies by a factor of 2 to ~ 10 . For instance, for the SG with $f_0 = 70$ Hz and $Q = 100$, the O3 rate limit was $1.08 \text{ Gpc}^{-3} \text{ yr}^{-1}$ while with O4a we obtain $0.43 \text{ Gpc}^{-3} \text{ yr}^{-1}$.

VI. SENSITIVITY FOR ASTROPHYSICAL SOURCE POPULATIONS

In this section we evaluate the performance of the search using simulated signals with the morphologies expected from core-collapse supernovae (CCSNe) [21–24] and sudden excitations of neutron stars [29, 30], of the type that might be associated with pulsar timing glitches and underlying starquakes [80]. This enables the interpretation of the null results of the present search in terms of astrophysically-motivated sources, most of which were also discussed in the previous LVK search on O3 data [36]. The procedure is based on the measurement of the detection efficiency: simulated signals are injected via software into the detector data streams and the analysis is repeated. Differently from the ad-hoc injections discussed in the previous section, here signal waveforms and amplitudes are informed by GW emission models and assumed spatial distributions of the sources. This search is still all-sky, meaning that the search does not depend on having additional information on the source from other cosmic messengers, such as from electromagnetic (EM) or neutrino observations. If such information is available to constrain the search, that would enable a




FIG. 4. Rate density upper limits (at 90% confidence) in terms of $\text{Gpc}^{-3} \text{ yr}^{-1}$ for elliptical sine-Gaussian injections, assuming standard candle sources emitting $1 M_{\odot} c^2$ of GW energy, for the best among the three low-frequency analyses in the case of O4a. Upper limits for waveforms with different Q values are plotted versus frequency. Gray markers show the O3 results for the SG waveforms listed for O3 in table I.

deeper exploitation of the GW data. The present all-sky search covers the scenario in which the EM emission is obscured or the EM or neutrino signal is intrinsically too weak to be detected. Throughout this section, quoted sensitivities follow the criterion for significant detection of $\text{IFAR} \geq 100$ years.

A. Core-Collapse Supernovae

GW transients from CCSNe can show complex time-frequency structures related to the inner dynamics of the source, see e.g. [23]. After a *bounce* which halts the initial collapse of the core, the continuing emission typically has a stochastic character with spectral features evolving in time. The GW emission is modeled by numerical simulations of the hydrodynamics and microphysics of the source, which provide deeper comprehension of the rich physical processes occurring within the time scale of ~ 1 second after the collapse of the progenitor star. In general terms, the GW emission is dominated by the proto-neutron star, which is excited and deformed by many concurrent processes, including aspherical hydrodynamical flows developing within the central few hundred kilometers and rotation of the collapsing star. Most of these GW features fall inside the signal parameter space targeted by the low-frequency analyses used in the present search.

We evaluate the search's sensitivity to CCSNe using predicted GW waveforms from seven different three-dimensional CCSN simulations. Along with the five mod-

Morphology	50% efficiency $h_{\text{rss}} (\times 10^{-22} \text{ Hz}^{-1/2})$			
	2G+XGB	2G+GMM	XP+XGB	O3
Gaussian pulses (linear)				
$\tau_{\text{GA}} = 0.1 \text{ ms}$	1.9	1.6	1.6	12.6
$\tau_{\text{GA}} = 2.5 \text{ ms}$	1.7	1.9	2.0	16.7
Sine-Gaussian wavelets (elliptical)				
$f_0 = 36 \text{ Hz}, Q = 3$	3.6	5.7	3.7	-
$f_0 = 36 \text{ Hz}, Q = 9$	4.6	4.5	3.9	-
$f_0 = 48 \text{ Hz}, Q = 9$	2.2	3.1	2.1	-
$f_0 = 48 \text{ Hz}, Q = 100$	3.4	3.6	4.3	-
$f_0 = 70 \text{ Hz}, Q = 3$	1.4	1.9	1.5	(2.9)
$f_0 = 70 \text{ Hz}, Q = 100$	1.8	2.2	1.7	(2.7)
$f_0 = 235 \text{ Hz}, Q = 9$	0.6	0.8	0.6	-
$f_0 = 235 \text{ Hz}, Q = 100$	0.9	1.1	0.9	(2.0)
$f_0 = 554 \text{ Hz}, Q = 9$	0.9	1.3	0.9	(2.0)
$f_0 = 849 \text{ Hz}, Q = 3$	1.4	1.8	1.4	(3.6)
$f_0 = 849 \text{ Hz}, Q = 9$	1.2	1.6	1.1	-
$f_0 = 849 \text{ Hz}, Q = 100$	1.1	1.3	1.1	-
$f_0 = 1304 \text{ Hz}, Q = 9$	1.7	2.3	1.7	(3.6)
$f_0 = 1615 \text{ Hz}, Q = 3$	2.9	3.1	2.9	-
$f_0 = 1615 \text{ Hz}, Q = 9$	2.1	2.9	2.2	-
$f_0 = 1615 \text{ Hz}, Q = 100$	1.8	2.0	1.8	(4.3)
$f_0 = 2000 \text{ Hz}, Q = 3$	2.7	-	-	(6.0)
$f_0 = 2477 \text{ Hz}, Q = 9$	3.0	-	-	(7.2)
$f_0 = 3067 \text{ Hz}, Q = 3$	4.3	-	-	(9.8)
White-noise bursts				
$f_0 = 100 \text{ Hz}, \Delta f = 100 \text{ Hz}, \tau_{\text{WNB}} = 0.1 \text{ s}$	0.6	0.8	0.6	1.0
$f_0 = 250 \text{ Hz}, \Delta f = 100 \text{ Hz}, \tau_{\text{WNB}} = 0.1 \text{ s}$	0.6	0.7	0.6	1.0
$f_0 = 750 \text{ Hz}, \Delta f = 100 \text{ Hz}, \tau_{\text{WNB}} = 0.1 \text{ s}$	0.8	0.9	0.8	1.5

TABLE I. The h_{rss} values (in units of $10^{-22} \text{ Hz}^{-1/2}$) for which 50% detection efficiency is achieved with an IFAR threshold of 100 years for each of the injected signal morphologies. The SG waveforms reported in this table have elliptical polarization. “-” denotes waveforms not evaluated for a given pipeline. The SG results with central frequency of 2000 Hz and higher are from the high-frequency analysis. We show the corresponding results for elliptically polarized SG waveforms for O3 in parentheses. Note that Table I in the O3 paper [36] reported values for circularly polarized SG.

Sine-Gaussian parameters	E_{GW} (in units of $10^{-10} M_{\odot} c^2$)			
	2G+XGB	2G+GMM	XP+XGB	O3
$f_0 = 36 \text{ Hz}, Q = 3$	1.4	3.6	1.5	-
$f_0 = 36 \text{ Hz}, Q = 9$	2.3	2.2	1.7	-
$f_0 = 48 \text{ Hz}, Q = 9$	0.9	1.9	0.9	-
$f_0 = 48 \text{ Hz}, Q = 100$	2.3	2.5	3.6	-
$f_0 = 70 \text{ Hz}, Q = 3$	0.8	1.5	0.9	3.5
$f_0 = 70 \text{ Hz}, Q = 100$	1.3	2.0	1.2	3.0
$f_0 = 235 \text{ Hz}, Q = 9$	1.7	3.0	1.7	-
$f_0 = 235 \text{ Hz}, Q = 100$	3.8	5.7	3.8	18.8
$f_0 = 554 \text{ Hz}, Q = 9$	21	44	21	104
$f_0 = 849 \text{ Hz}, Q = 3$	120	200	120	790
$f_0 = 849 \text{ Hz}, Q = 9$	88	157	74	-
$f_0 = 849 \text{ Hz}, Q = 100$	74	104	74	-
$f_0 = 1304 \text{ Hz}, Q = 9$	420	760	420	1870
$f_0 = 1615 \text{ Hz}, Q = 3$	1900	2100	1900	-
$f_0 = 1615 \text{ Hz}, Q = 9$	980	1860	1070	-
$f_0 = 1615 \text{ Hz}, Q = 100$	720	890	720	4100
$f_0 = 2000 \text{ Hz}, Q = 3$	2500	-	-	12200
$f_0 = 2477 \text{ Hz}, Q = 9$	4700	-	-	27000
$f_0 = 3067 \text{ Hz}, Q = 3$	15000	-	-	77000

TABLE II. GW emitted energy E_{GW} in units of $10^{-10} M_{\odot} c^2$ that corresponds to 50% detection efficiency at IFAR ≥ 100 years, assuming a source distance $r_0 = 10 \text{ kpc}$. We present the results for different SG elliptical waveforms with varying central frequency f_0 and quality factor Q for each pipeline used in O4 and also for the waveforms published for O3. “-” denotes waveforms not evaluated for a given pipeline or not for O3.

els analyzed in the all-sky search on O3 data [36], we add two recent models of CCSNe which are likely to lack EM emission:

- Model *40 NR* (*Pan+21 40 NR* from [81]) has a non-rotating, $40 M_{\odot}$ zero age main sequence (ZAMS) mass progenitor with solar metallicity. Black hole (BH) formation occurs at $\sim 0.78 \text{ s}$ after the initial

bounce. The dominant GW emission is associated with the proto-neutron star excitation, shows a rising frequency with time, and peaks at $\sim 2 \text{ kHz}$ before BH formation. A subdominant emission from standing accretion shock instability (SASI, [82–84]) is also present below 200 Hz.

- Model *s18np* (*Pow+20 s18np* from [85]) has a

non-rotating, $18 M_{\odot}$ ZAMS progenitor with solar metallicity. The GW emission from the proto-neutron star peaks at ~ 0.7 kHz with a subdominant emission below 200 Hz from SASI. Differently from model *s18* below, there is no hint for shock revival by the end of the simulation (at ~ 0.56 s after the bounce).

The likely lack of EM emission makes these two models and model *m20p* (see below) particularly interesting as plausible targets for an all-sky burst search.

The five models that were evaluated with the previous all-sky search (see [36] for more details) enable a direct comparison of sensitivity. They cover three cases with GW energy emission in the lower range of predictions and two cases of more extreme models with higher GW emission. The former three models have solar metallicity non-rotating progenitors:

- Model *s9* (*Rad+19 s9* from [86]) from a $9 M_{\odot}$ ZAMS progenitor. This is at the low end of the progenitor mass spectrum for CCSNe.
- Model *s18* (*Pow+19 s18* from [87]) from a $18 M_{\odot}$ ZAMS progenitor. Differently from model *s18np*, this model includes strong aspherical seed perturbations from convective oxygen burning and undergoes shock revival and SN explosion.
- Model *m20p* (*Oco+18 mesa20_3D_pert* from [88]) from a $20 M_{\odot}$ ZAMS progenitor. This model produces a BH remnant without powering EM emission.

Extreme models with a strongly rotating, low-metallicity progenitor are:

- Model *m39* (*Pow+20 m39* from [85]) from a $39 M_{\odot}$ ZAMS progenitor with $1/50$ solar metallicity.
- Model *35OC* (*Obe+20 35OC-RO* from [89]) from a $35 M_{\odot}$ ZAMS progenitor with $1/50$ solar metallicity.

All signals from these models have been injected into the detectors' data streams with a random orientation of the source. The other extrinsic parameters of the source, in particular distance and sky direction, have been drawn from two population models: i) an isotropic model and ii) a Galactic model. The former allows to measure the average detection efficiency as a function of distance for each all-sky analysis. The sky direction is sampled uniformly in solid angle over all directions, while the distance to the source is distributed between 0.01 kpc and 100 kpc. We present in Figure 5 the distances corresponding to 50% and 10% detection efficiency. Our results improve the distance reach by a factor ~ 2 for the five waveforms that are common to this work and the previously published O3 search [36].

We note that the LVK recently performed a dedicated search for GW bursts associated with SN 2023ixf [90]




FIG. 5. Sensitivity of the all-sky GW search to core-collapse supernova events for different emission models. Boxes represent distance ranges between detection efficiencies 50% and 10% at IFAR > 100 years, evaluated using an isotropic distribution of sources, for each low-frequency analysis. We also plot the results from the previous observing run, O3. The current search reaches the Galactic center for three of the emission models we considered. Models *40 NR* and *m20p* form a BH remnant. The difference for models *s18* and *s18np* reflects the impact of assuming diverse aspherical seed perturbations for progenitor stars with equal masses.

using LIGO data from the engineering run preceding the start of O4a. In that case, the published LVK results [43] make use of the known sky position of the source, achieving a somewhat better detection efficiency at equal distance for the four waveform models in common. Compared to the ad-hoc signals discussed in section V, CCSNe signals are more challenging to detect due to their wider time-frequency structure, as discussed in [24, 91].

In the Galactic model, we generate a population of simulated sources with sky directions and distances drawn from a model of the Milky Way stellar mass distribution, including a bulge, a thick stellar disk and a thin stellar disk, whose parameters are taken from [92] and [93]. In this case we present the overall efficiencies of the three low-frequency analyses for each of the CCSN models in Table III. The efficiency is representative of the probability of detection of a Galactic CCSN for each model: we call this CCSN Galactic coverage. For two of the lower-energy CCSN models (*s9* and *m20p*) the Galactic coverage is practically zero. In fact, *s9* has a progenitor at the lower end of the mass spectrum of CCSNe and *m20p* lacks shock revival, which make them detectable only out to ~ 1 kpc. Within this range, the Galactic distribution provides very rare CCSNe. On the other hand, the Galactic coverage is $\sim 90\%$ or higher for the *m39* and *35OC* models, meaning that a CCSN with those properties occurring at a random location in the Milky Way during O4a would likely have been detected by our search.

Model	2G+XGB	2G+GMM	XP+XGB	best O3
<i>s9</i>	< 0.1%	< 0.1%	< 0.1%	< 0.1%
<i>m20p</i>	< 0.1%	< 0.1%	< 0.1%	< 0.1%
<i>s18np</i>	4.6%	2.0%	4.5%	–
<i>40 NR</i>	20.4%	19.6%	22.8%	–
<i>s18</i>	34.1%	20.4%	32.3%	1.2%
<i>m39</i>	89.5%	74.0%	87.4%	69.4%
<i>35OC</i>	97.2%	93.8%	95.7%	89.8%

TABLE III. The fraction of Galactic CCSN signals that are detectable with an IFAR threshold of 100 years or greater by each low-frequency analysis, for different modeled CCSN waveforms. There is a huge difference between models of CC-SNe with lower and higher GW amplitudes. The last column reports the best results on O3 data [36].

B. Isolated Neutron Star Emitters

Among the known pulsars (rotating neutron stars with beamed radio and/or high-energy EM emission that is detected at Earth as pulses at nearly constant intervals), some exhibit sudden increases in pulse rate known as timing *glitches*. Such a change indicates a reconfiguration of the compact object and its magnetosphere with a concurrent release of energy, some of which may go into transient GWs. In this work we consider the detectability of GW transients connected with neutron star timing glitches from the inferred population of nearby neutron stars which are not observed as pulsars because their EM emission is not angled favorably for telescopes on Earth to detect. The GW emission from a perturbed, rotating neutron star is more isotropic and can be detected even from EM-dark pulsars [94]. Transient excitations of neutron stars associated with *observed* EM signatures are best covered by dedicated GW searches, e.g. for the August 2006 glitch of the Vela Pulsar [95] or magnetar flares (see [96] for O3 results).

To investigate the sensitivity of our search to neutron star excitations, we follow the same lines adopted in the previous LVK analysis on O3 data [36]. In particular, we assume that the dominant GW emission is due to a short-duration *f*-mode excitation of the glitching neutron star [95, 97] and we model the related GW burst as a damped sinusoid signal. This choice is not restrictive, however, as our search does not depend on the morphology of the incoming GW signal at first order. Typical frequencies and damping times of these signals are expected to be in the ranges $2 \text{ kHz} \leq \nu_{\text{GW}} \leq 3 \text{ kHz}$ and from tens of milliseconds to as much as half a second respectively, depending on the mass and the equation of state (EoS) of the neutron star [97]. Searches for GW emission with much longer duration have been performed using different methods, see e.g. the LVK results on O3 data [98].

Our all-sky search has not found any significant event in this high-frequency band. The null result can be interpreted under the optimistic assumptions that all the glitch energy is converted into GW energy, $E_{\text{glitch}} = E_{\text{GW}}$, and

that the source’s angular momentum is optimally oriented, so that the incoming GW burst is circularly polarized. We adopt as a standard source a pulsar at a distance of 287 pc and with spin frequency $\nu_s = 11.2 \text{ Hz}$, which are the parameters of the closest known glitching pulsar [99, 100], the Vela Pulsar. This allows us to interpret the GW search in terms of minimum detectable glitch size, $\Delta\nu_s$, following equation 5 in [97]. The Vela Pulsar belongs to the subpopulation of known pulsars which are both spinning rapidly and undergoing larger glitches [99, 100], and thus are delivering glitch energies at the higher end of the observed range, since $E_{\text{glitch}} \propto \nu_s \Delta\nu_s$.

A population of such standard sources is simulated uniformly distributed in the sky and emitted GW signals are estimated according to two EoS models, APR4 (soft) [101] and H4 (hard) [102], for neutron star masses in the range of $1\text{--}2 M_\odot$ in the non-rotating limit [103]. Detectable glitch sizes (requiring $\text{IFAR} \geq 100$ years) are reported as a function of the neutron star mass and EoS in Figure 6. The minimum detectable pulsar glitch size for the O4a run, expressed as a fractional change referenced to the Vela Pulsar’s frequency, $\Delta\nu_s/\nu_s$, ranges from $\sim 2 \times 10^{-5}$ to $\sim 6 \times 10^{-5}$ depending on the neutron star mass. These values are a factor of ~ 3 to 5 better than what was obtained in O3. The actual Vela Pulsar glitches with fractional magnitude $\Delta\nu_s/\nu_s \approx 1\text{--}3 \times 10^{-6}$ approximately every 3 years [104], so this all-sky search is still an order of magnitude above that value. Nevertheless, many known pulsars show glitches in the range $10^{-5}\text{--}10^{-4} \text{ Hz}$ [105–107], therefore our current sensitivity provides a potential discovery channel for yet unknown glitching pulsars which could be closer to Earth and/or spinning faster than the Vela Pulsar.

VII. CONCLUSIONS

We reported the results of a search for gravitational-wave bursts in LIGO data during the LVK O4a observing period using three pipelines based on variants of the cWB algorithm followed by machine-learning stages. Three low-frequency analyses and one high-frequency analysis targeted GW bursts lasting up to a few seconds with frequency content in the range $16\text{--}4096 \text{ Hz}$. By searching as broadly as possible, we sought to uncover new types of gravitational-wave signals from either known or unknown astrophysical objects. No significant candidate was found beyond the CBC candidates already identified by low-latency searches using modeled templates. The most significant new candidate found has a false alarm rate of one per 1.75 years, statistically consistent with the expected background for this dataset.

We evaluated the sensitivity of the search using ad-hoc simulated signals as well as two types of astrophysically motivated signals. The ad-hoc signals consisted of sine-Gaussians with a wide range of frequencies and Q values plus a few Gaussian pulses and white-noise bursts. Our




FIG. 6. Sensitivity of the high-frequency 2G+XGB analysis to neutron star glitches, considering spin frequency and distance of the Vela Pulsar as a representative standard source for soft (APR4) and hard (H4) EoS and assuming an optimally oriented source at random sky directions. For each EoS and mass bin, the dot shows the mean value of the glitch size that yields 50% detection efficiency at IFAR ≥ 100 years, and the error bar represents the variation within the mass bin. On average, a higher-mass neutron star allows for smaller glitches to be detected. The fractional glitch size $\Delta\nu_s/\nu_s$ for a Vela-like pulsar would need to be stronger than ~ 2 to 6×10^{-5} for 50% of the sources to be detected in O4a.

search was capable of detecting such signals at amplitudes typically a factor of 2 or more weaker than the previous published search, which used data from the LVK O3 run. Thus, even with the shorter duration of O4a, our search achieved the greatest time-volume exposure to date for generic short-duration burst signals. Assuming a population of rotating sources, we placed rate density upper limits an order of magnitude lower than the O3 search for signal frequencies between ~ 100 Hz and 1 kHz. We considered seven stellar core-collapse models and found that our search could have detected GW signals throughout most of the Milky Way from two of the models with high-mass star progenitors, with more limited reach for the other models. Finally, we simulated damped sinusoid GW signals representing neutron star f -mode oscillations excited in connection with pulsar timing glitches. Using an optimistic energy argument, we estimated the glitch size (fractional frequency change) that our GW search could have detected, finding that typical glitches from the Vela Pulsar would not have been detectable but a similarly close unseen pulsar with plausible properties could potentially have been detected if it had experienced a timing glitch during O4a.

This search only used data from the first period of the O4 run, from May 24, 2023 to January 16, 2024. After a commissioning break, the O4 run resumed on April 10, 2024 with the Virgo detector collecting data along with the two LIGO detectors. While Virgo has been a factor of ~ 3 less sensitive than LIGO during the O4 run, it often helps greatly to localize event candidates in

the sky by triangulation with the three-detector network [94, 108]. The KAGRA detector rejoined the O4 run in June 2025 with a noise level ~ 7 times less sensitive than Virgo, as commissioning of KAGRA is still ongoing. The O4 run is now expected to end on November 18, 2025 after accumulating about two years of GW data, excluding commissioning breaks during the run. The full O4 data set will enable more-sensitive searches for GW burst signals, and having data from three (or more) sensitive detectors will improve our ability to characterize any candidate which is found. Even if O4 does not reveal any GW burst signals, subsequent observing runs after further detector upgrades, and with new facilities, will enable deeper, longer-duration searches.

ACKNOWLEDGMENTS

This material is based upon work supported by NSF’s LIGO Laboratory, which is a major facility fully funded by the National Science Foundation. The authors also gratefully acknowledge the support of the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO 600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Netherlands Organization for Scientific Research (NWO) for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación (AEI), the Spanish Ministerio de Ciencia, Innovación y Universidades, the European Union NextGenerationEU/PRTR (PRTR-C17.I1), the ICSC - Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by the European Union NextGenerationEU, the Comunitat Autònoma de les Illes Balears through the Conselleria d’Educació i Universitats, the Conselleria d’Innovació, Universitats, Ciència i Societat Digital de la Generalitat Valenciana and the CERCA Programme Generalitat de Catalunya, Spain, the Polish National Agency for Academic Exchange, the National Science Centre of Poland and the European Union – European Regional Development Fund; the Foundation for Polish Science (FNP), the Polish Ministry of Science and Higher Education, the Swiss National Science Foundation (SNSF), the Russian Science Foundation, the European Commission, the European Social Funds (ESF),

the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the French Lyon Institute of Origins (LIO), the Belgian Fonds de la Recherche Scientifique (FRS-FNRS), Actions de Recherche Concertées (ARC) and Fonds Wetenschappelijk Onderzoek – Vlaanderen (FWO), Belgium, the Paris Île-de-France Region, the National Research, Development and Innovation Office of Hungary (NKFIH), the National Research Foundation of Korea, the Natural Sciences and Engineering Research Council of Canada (NSERC), the Canadian Foundation for Innovation (CFI), the Brazilian Ministry of Science, Technology, and Innovations, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Israel Science Foundation (ISF), the US-Israel Binational Science Fund (BSF), the Leverhulme Trust, the Research Corporation, the National Science and Technology Council (NSTC), Taiwan, the United States Department of Energy, and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, INFN and CNRS for provision of computational resources.

This work was supported by MEXT, the JSPS Leading-edge Research Infrastructure Program, JSPS Grant-in-Aid for Specially Promoted Research 26000005,

JSPS Grant-in-Aid for Scientific Research on Innovative Areas 2402: 24103006, 24103005, and 2905: JP17H06358, JP17H06361 and JP17H06364, JSPS Core-to-Core Program A. Advanced Research Networks, JSPS Grants-in-Aid for Scientific Research (S) 17H06133 and 20H05639, JSPS Grant-in-Aid for Transformative Research Areas (A) 20A203: JP20H05854, the joint research program of the Institute for Cosmic Ray Research, University of Tokyo, the National Research Foundation (NRF), the Computing Infrastructure Project of the Global Science experimental Data hub Center (GSDC) at KISTI, the Korea Astronomy and Space Science Institute (KASI), the Ministry of Science and ICT (MSIT) in Korea, Academia Sinica (AS), the AS Grid Center (ASGC) and the National Science and Technology Council (NSTC) in Taiwan under grants including the Science Vanguard Research Program, the Advanced Technology Center (ATC) of NAOJ, and the Mechanical Engineering Center of KEK.

Additional acknowledgements for support of individual authors may be found in the following document: <https://dcc.ligo.org/LIGO-M2300033/public>. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising. We request that citations to this article use “A. G. Abac *et al.* (LIGO-Virgo-KAGRA Collaboration), ...” or similar phrasing, depending on journal convention.

This paper has been assigned LIGO document ID P2400601-v8.

-
- [1] J. Aasi *et al.* (LIGO Scientific Collaboration), Advanced LIGO, *Class. Quant. Grav.* **32**, 074001 (2015), arXiv:1411.4547 [gr-qc].
 - [2] F. Acernese *et al.* (Virgo Collaboration), Advanced Virgo: a second-generation interferometric gravitational wave detector, *Class. Quant. Grav.* **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
 - [3] B. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, *Phys. Rev. X* **9**, 031040 (2019), arXiv:1811.12907 [astro-ph.HE].
 - [4] R. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), GWTC-2.1: Deep extended catalog of compact binary coalescences observed by LIGO and Virgo during the first half of the third observing run, *Phys. Rev. D* **109**, 022001 (2024), arXiv:2108.01045 [gr-qc].
 - [5] R. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo during the Second Part of the Third Observing Run, *Phys. Rev. X* **13**, 041039 (2023), arXiv:2111.03606 [gr-qc].
 - [6] A. H. Nitz, C. Capano, A. B. Nielsen, S. Reyes, R. White, D. A. Brown, and B. Krishnan, 1-OGC: The First Open Gravitational-wave Catalog of Binary Mergers from Analysis of Public Advanced LIGO Data, *Astrophys. J.* **872**, 195 (2019), arXiv:1811.01921 [gr-qc].
 - [7] A. H. Nitz, T. Dent, G. S. Davies, S. Kumar, C. D. Capano, I. Harry, S. Mozzon, L. Nuttall, A. Lundgren, and M. Tápai, 2-OGC: Open Gravitational-wave Catalog of Binary Mergers from Analysis of Public Advanced LIGO and Virgo Data, *Astrophys. J.* **891**, 123 (2020), arXiv:1910.05331 [astro-ph.HE].
 - [8] A. H. Nitz, C. D. Capano, S. Kumar, Y.-F. Wang, S. Kastha, M. Schäfer, R. Dhurkunde, and M. Cabero, 3-OGC: Catalog of Gravitational Waves from Compact-binary Mergers, *Astrophys. J.* **922**, 76 (2021), arXiv:2105.09151 [astro-ph.HE].
 - [9] A. H. Nitz, S. Kumar, Y.-F. Wang, S. Kastha, S. Wu, M. Schäfer, R. Dhurkunde, and C. D. Capano, 4-OGC: Catalog of Gravitational Waves from Compact Binary Mergers, *Astrophys. J.* **946**, 59 (2023), arXiv:2112.06878 [astro-ph.HE].
 - [10] T. Venumadhav, B. Zackay, J. Roulet, L. Dai, and M. Zaldarriaga, New binary black hole mergers in the second observing run of Advanced LIGO and Advanced Virgo, *Phys. Rev. D* **101**, 083030 (2020), arXiv:1904.07214 [astro-ph.HE].
 - [11] B. Zackay, L. Dai, T. Venumadhav, J. Roulet, and M. Zaldarriaga, Detecting gravitational waves with

- disparate detector responses: Two new binary black hole mergers, *Phys. Rev. D* **104**, 063030 (2021), arXiv:1910.09528 [astro-ph.HE].
- [12] S. Olsen, T. Venumadhav, J. Mushkin, J. Roulet, B. Zackay, and M. Zaldarriaga, New binary black hole mergers in the LIGO-Virgo O3a data, *Phys. Rev. D* **106**, 043009 (2022), arXiv:2201.02252 [astro-ph.HE].
- [13] A. K. Mehta, S. Olsen, D. Wadekar, J. Roulet, T. Venumadhav, J. Mushkin, B. Zackay, and M. Zaldarriaga, New binary black hole mergers in the LIGO-Virgo O3b data, *Phys. Rev. D* **111**, 024049 (2025).
- [14] D. Wadekar, J. Roulet, T. Venumadhav, A. K. Mehta, B. Zackay, J. Mushkin, S. Olsen, and M. Zaldarriaga, New black hole mergers in the LIGO-Virgo O3 data from a gravitational wave search including higher-order harmonics, arXiv e-prints, arXiv:2312.06631 (2023), arXiv:2312.06631 [gr-qc].
- [15] Gravitational-Wave Candidate Event Database (GraceDB), <https://gracedb.ligo.org/superevents/public/04/>.
- [16] T. Akutsu *et al.* (KAGRA Collaboration), Overview of KAGRA: Detector design and construction history, *PTEP* **2021**, 05A101 (2021), arXiv:2005.05574 [physics.ins-det].
- [17] B. Iyer *et al.*, *LIGO-India, Proposal of the Consortium for Indian Initiative in Gravitational-wave Observations (IndIGO)*, Tech. Report M1100296 (LIGO-India, 2011) <https://dcc.ligo.org/LIGO-M1100296/public>.
- [18] M. Saleem, J. Rana, V. Gayathri, A. Vijaykumar, S. Goyal, S. Sachdev, J. Suresh, S. Sudhagar, A. Mukherjee, G. Gaur, B. Sathyaprakash, A. Pai, R. X. Adhikari, P. Ajith, and S. Bose, The science case for LIGO-India, *Classical and Quantum Gravity* **39**, 025004 (2022), arXiv:2105.01716 [gr-qc].
- [19] M. Branchesi, M. Maggiore, D. Alonso, *et al.*, Science with the Einstein Telescope: a comparison of different designs, *J. Cos. Astroparticle Phys.* **2023**, 068 (2023), arXiv:2303.15923 [gr-qc].
- [20] M. Evans *et al.*, *A Horizon Study for Cosmic Explorer: Science, Observatories, and Community*, Document CE-P2100003 (Cosmic Explorer Consortium, 2021) <https://dcc.cosmicexplorer.org/CE-P2100003/public>.
- [21] H.-T. Janka, Explosion Mechanisms of Core-Collapse Supernovae, *Annual Review of Nuclear and Particle Science* **62**, 407 (2012), arXiv:1206.2503 [astro-ph.SR].
- [22] S. E. Gossan, P. Sutton, A. Stuver, M. Zanolin, K. Gill, and C. D. Ott, Observing gravitational waves from core-collapse supernovae in the advanced detector era, *Phys. Rev. D* **93**, 042002 (2016).
- [23] E. Abdikamalov, G. Pagliaroli, and D. Radice, Gravitational waves from core-collapse supernovae, in *Handbook of Gravitational Wave Astronomy*, edited by C. Bambi, S. Katsanevas, and K. D. Kokkotas (Springer Nature Singapore, Singapore, 2022) pp. 909–945.
- [24] M. J. Szczepańczyk, J. M. Antelis, M. Benjamin, M. Cavaglia, D. Gondek-Rosińska, T. Hansen, S. Klimenko, M. D. Morales, C. Moreno, S. Mukherjee, G. Nurbek, J. Powell, N. Singh, S. Sitmukhambetov, P. Szewczyk, O. Valdez, G. Vedovato, J. Westhouse, M. Zanolin, and Y. Zheng, Detecting and reconstructing gravitational waves from the next galactic core-collapse supernova in the advanced detector era, *Phys. Rev. D* **104**, 102002 (2021).
- [25] A. Mezzacappa and M. Zanolin, Gravitational waves from neutrino-driven core collapse supernovae: Predictions, detection, and parameter estimation (2024), arXiv:2401.11635 [astro-ph.HE].
- [26] E. Nazari and M. Roshan, Gravitational radiation by magnetic field: application to millisecond magnetars, *MNRAS* **498**, 110 (2020).
- [27] I. Contopoulos, D. Kazanas, and D. B. Papadopoulos, Gravitational waves from the pulsar magnetosphere, *Monthly Notices of the Royal Astronomical Society* **527**, 11198 (2023).
- [28] C. Kouvaris, Gravitational waves from magnetars, *International Journal of Modern Physics D* **34**, 2550037 (2025), arXiv:2406.03513 [astro-ph.HE].
- [29] A. L. Watts and T. E. Strohmayer, Neutron star oscillations and QPOs during magnetar flares, *Advances in Space Research* **40**, 1446 (2007), arXiv:astro-ph/0612252 [astro-ph].
- [30] E. Giliberti and G. Cambiotti, Starquakes in millisecond pulsars and gravitational waves emission, *MNRAS* **511**, 3365 (2022), arXiv:2102.02540 [astro-ph.HE].
- [31] M. Ebersold and S. Tiwari, Search for nonlinear memory from subsolar mass compact binary mergers, *Phys. Rev. D* **101**, 104041 (2020), arXiv:2005.03306 [gr-qc].
- [32] S. Bini, S. Tiwari, Y. Xu, L. Smith, M. Ebersold, G. Principe, M. Haney, P. Jetzer, and G. A. Prodi, Search for hyperbolic encounters of compact objects in the third LIGO-Virgo-KAGRA observing run, *Phys. Rev. D* **109**, 042009 (2024), arXiv:2311.06630 [gr-qc].
- [33] T. Damour and A. Vilenkin, Gravitational Wave Bursts from Cosmic Strings, *Phys. Rev. Lett.* **85**, 3761 (2000), arXiv:gr-qc/0004075 [gr-qc].
- [34] T. Damour and A. Vilenkin, Gravitational wave bursts from cusps and kinks on cosmic strings, *Phys. Rev. D* **64**, 064008 (2001), arXiv:gr-qc/0104026 [astro-ph].
- [35] T. Damour and A. Vilenkin, Gravitational radiation from cosmic (super)strings: Bursts, stochastic background, and observational windows, *Phys. Rev. D* **71**, 063510 (2005), arXiv:hep-th/0410222 [hep-th].
- [36] R. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), All-sky search for short gravitational-wave bursts in the third Advanced LIGO and Advanced Virgo run, *Phys. Rev. D* **104**, 122004 (2021), arXiv:2107.03701 [gr-qc].
- [37] R. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), All-sky search for long-duration gravitational-wave bursts in the third Advanced LIGO and Advanced Virgo run, *Phys. Rev. D* **104**, 102001 (2021), arXiv:2107.13796 [gr-qc].
- [38] R. Abbott *et al.* (LIGO-VIRGO-KAGRA Collaboration), Search for intermediate mass black hole binaries in the third observing run of Advanced LIGO and Advanced Virgo, *Astron. Astrophys.* **659**, A84 (2022), arXiv:2105.15120 [astro-ph.HE].
- [39] A. G. Abac *et al.* (LIGO-Virgo-KAGRA Collaboration), Search for Eccentric Black Hole Coalescences during the Third Observing Run of LIGO and Virgo, *Astrophys. J.* **973**, 132 (2024).
- [40] A. G. Abac *et al.* (LIGO-Virgo-KAGRA Collaboration), All-sky search for long-duration gravitational-wave transients in the first part of the fourth LIGO-Virgo-KAGRA Observing run, arXiv e-print 2507.12282 (2025), arXiv:2507.12282 [astro-ph.HE].
- [41] S. Klimenko, G. Vedovato, M. Drago, F. Salemi, V. Tiwari, G. Prodi, C. Lazzaro, K. Ackley, S. Tiwari, C. Da Silva, *et al.*, Method for detection and recon-

- struction of gravitational wave transients with networks of advanced detectors, *Phys. Rev. D* **93**, 042004 (2016).
- [42] M. Drago, S. Klimenko, C. Lazzaro, E. Milotti, G. Mitselmakher, V. Necula, B. O'Brian, G. A. Prodi, F. Salemi, M. Szczepańczyk, *et al.*, coherent WaveBurst, a pipeline for unmodeled gravitational-wave data analysis, *SoftwareX* **14**, 100678 (2021).
- [43] A. G. Abac *et al.* (LIGO-Virgo-KAGRA Collaboration), Search for Gravitational Waves Emitted from SN 2023ixf, *Astrophys. J.* **985**, 183 (2025), arXiv:2410.16565 [astro-ph.HE].
- [44] E. Capote, W. Jia, N. Aritomi, M. Nakano, V. Xu, *et al.*, Advanced LIGO detector performance in the fourth observing run, *Phys. Rev. D* **111**, 062002 (2025).
- [45] S. Soni *et al.*, LIGO Detector Characterization in the First Half of the Fourth Observing Run, *Classical and Quantum Gravity* **42**, 085016 (2025), arXiv:2409.02831 [astro-ph.IM].
- [46] D. Ganapathy *et al.* (LIGO O4 Detector Collaboration), Broadband Quantum Enhancement of the LIGO Detectors with Frequency-Dependent Squeezing, *Physical Review X* **13**, 041021 (2023).
- [47] L. Sun *et al.*, Characterization of systematic error in Advanced LIGO calibration, *Classical and Quantum Gravity* **37**, 225008 (2020), arXiv:2005.02531 [astro-ph.IM].
- [48] L. Sun *et al.*, Characterization of systematic error in Advanced LIGO calibration in the second half of O3, arXiv e-prints, arXiv:2107.00129 (2021), arXiv:2107.00129 [astro-ph.IM].
- [49] M. Cabero, A. Lundgren, A. H. Nitz, T. Dent, D. Barker, E. Goetz, J. S. Kissel, L. K. Nuttall, P. Schale, R. Schofield, and D. Davis, Blip glitches in Advanced LIGO data, *Classical and Quantum Gravity* **36**, 155010 (2019), arXiv:1901.05093 [physics.ins-det].
- [50] Y. Wu, M. Zevin, C. P. L. Berry, K. Crowston, C. Østerlund, Z. Doctor, S. Banagiri, C. B. Jackson, V. Kalogera, and A. K. Katsaggelos, Advancing Glitch Classification in Gravity Spy: Multi-view Fusion with Attention-based Machine Learning for Advanced LIGO's Fourth Observing Run, arXiv e-print 2401.12913 (2024), arXiv:2401.12913 [gr-qc].
- [51] D. Davis *et al.* (LIGO O3 Detector Collaboration), LIGO detector characterization in the second and third observing runs, *Classical and Quantum Gravity* **38**, 135014 (2021), arXiv:2101.11673 [astro-ph.IM].
- [52] S. Soni *et al.*, Reducing scattered light in LIGO's third observing run, *Classical and Quantum Gravity* **38**, 025016 (2021), arXiv:2007.14876 [astro-ph.IM].
- [53] Coherent WaveBurst, <https://gwburst.gitlab.io/>.
- [54] V. Necula, S. Klimenko, and G. Mitselmakher, Transient analysis with fast Wilson-Daubechies time-frequency transform, in *Journal of Physics: Conference Series*, Vol. 363 (IOP Publishing, 2012) p. 012032.
- [55] S. Klimenko, Wavescan: multiresolution regression of gravitational-wave data, arXiv e-print 2201.01096 (2022), arXiv:2201.01096 [physics.data-an].
- [56] S. Klimenko, S. Mohanty, M. Rakhmanov, and G. Mitselmakher, Constraint likelihood analysis for a network of gravitational wave detectors, *Phys. Rev. D* **72**, 122002 (2005).
- [57] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for short gravitational-wave bursts in the first Advanced LIGO run, *Phys. Rev. D* **95**, 042003 (2017).
- [58] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for long-duration gravitational wave transients in the first Advanced LIGO observing run, *Classical and Quantum Gravity* **35**, 065009 (2018).
- [59] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for short gravitational-wave bursts in the second Advanced LIGO and Advanced Virgo run, *Phys. Rev. D* **100**, 024017 (2019).
- [60] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for long-duration gravitational-wave transients in the second Advanced LIGO observing run, *Phys. Rev. D* **99**, 104033 (2019).
- [61] T. Mishra, S. Bhaumik, V. Gayathri, M. J. Szczepańczyk, I. Bartos, and S. Klimenko, Gravitational waves detected by a burst search in LIGO/Virgo's third observing run, *Phys. Rev. D* **111**, 023054 (2025).
- [62] T. Chen and C. Guestrin, XGBoost: A scalable tree boosting system, in *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (2016) pp. 785–794.
- [63] M. J. Szczepańczyk, F. Salemi, S. Bini, T. Mishra, G. Vedovato, V. Gayathri, I. Bartos, S. Bhaumik, M. Drago, O. Halim, C. Lazzaro, A. Miani, E. Milotti, G. A. Prodi, S. Tiwari, and S. Klimenko, Search for gravitational-wave bursts in the third Advanced LIGO-Virgo run with coherent WaveBurst enhanced by machine learning, *Phys. Rev. D* **107**, 062002 (2023).
- [64] T. Mishra, B. O'Brien, M. Szczepańczyk, G. Vedovato, S. Bhaumik, V. Gayathri, G. Prodi, F. Salemi, E. Milotti, I. Bartos, and S. Klimenko, Search for binary black hole mergers in the third observing run of Advanced LIGO-Virgo using coherent WaveBurst enhanced with machine learning, *Phys. Rev. D* **105**, 083018 (2022).
- [65] T. Mishra, B. O'Brien, V. Gayathri, M. Szczepańczyk, S. Bhaumik, I. Bartos, and S. Klimenko, Optimization of model independent gravitational wave search for binary black hole mergers using machine learning, *Phys. Rev. D* **104**, 023014 (2021).
- [66] V. Gayathri, D. Lopez, R. Pranjali, I. S. Heng, A. Pai, and C. Messenger, Enhancing the sensitivity of transient gravitational wave searches with Gaussian Mixture Models, *Phys. Rev. D* **102**, 104023 (2020).
- [67] D. Lopez, V. Gayathri, A. Pai, I. S. Heng, C. Messenger, and S. K. Gupta, Utilizing gaussian mixture models in all-sky searches for short-duration gravitational wave bursts, *Phys. Rev. D* **105**, 063024 (2022).
- [68] L. Smith, S. Ghosh, J. Sun, V. Gayathri, I. S. Heng, and A. Pai, Enhancing search pipelines for short gravitational-wave transients with Gaussian mixture modeling, *Phys. Rev. D* **110**, 083032 (2024).
- [69] L. Tsukada *et al.*, Improved ranking statistics of the GstLAL inspiral search for compact binary coalescences, *Phys. Rev. D* **108**, 043004 (2023), arXiv:2305.06286 [astro-ph.IM].
- [70] B. Ewing *et al.*, Performance of the low-latency GstLAL inspiral search towards LIGO, Virgo, and KAGRA's fourth observing run, *Phys. Rev. D* **109**, 042008 (2024), arXiv:2305.05625 [gr-qc].
- [71] T. Dal Canton, A. H. Nitz, B. Gadre, G. S. Cabourn Davies, V. Villa-Ortega, T. Dent, I. Harry, and L. Xiao, Real-time Search for Compact Binary Merg-

- ers in Advanced LIGO and Virgo's Third Observing Run Using PyCBC Live, *Astrophys. J.* **923**, 254 (2021), arXiv:2008.07494 [astro-ph.HE].
- [72] Q. Chu *et al.*, SPIIR online coherent pipeline to search for gravitational waves from compact binary coalescences, *Phys. Rev. D* **105**, 024023 (2022), arXiv:2011.06787 [gr-qc].
- [73] C. All  n  , F. Aubin, I. Bentara, D. Buskulic, G. M. Guidi, V. Juste, M. Lethuillier, F. Marion, L. Mobilia, B. Mours, A. Ouzriat, T. Sainrat, and V. Sordini, The MBTA pipeline for detecting compact binary coalescences in the fourth LIGO-Virgo-KAGRA observing run, *Classical and Quantum Gravity* **42**, 105009 (2025), arXiv:2501.04598 [gr-qc].
- [74] A. G. Abac *et al.* (LIGO-Virgo-KAGRA Collaboration), GW231123: a Binary Black Hole Merger with Total Mass $190\text{--}265 M_\odot$, arXiv e-print 2507.08219 (2025), arXiv:2507.08219 [astro-ph.HE].
- [75] The CBC candidates excised from this search will be published in the next update to the LVK Gravitational-Wave Transient Catalog (GWTC) and online in the Gravitational Wave Open Science Center, <http://www.gwosc.org>.
- [76] A. Helmling-Cornell, P. Nguyen, R. Schofield, and R. Frey, Automated evaluation of environmental coupling for Advanced LIGO gravitational wave detections, *Class. Quant. Grav.* **41**, 145003 (2024), arXiv:2312.00735 [gr-qc].
- [77] P. Nguyen, R. Schofield, A. Effler, C. Austin, V. Adya, M. Ball, S. Banagiri, K. Banowetz, C. Billman, C. Blair, *et al.*, Environmental noise in advanced LIGO detectors, *Classical and Quantum Gravity* **38**, 145001 (2021).
- [78] J. Abadie *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for gravitational-wave bursts in the second joint LIGO-Virgo run, *Phys. Rev. D* **85**, 122007 (2012), arXiv:1202.2788 [gr-qc].
- [79] P. J. Sutton, A Rule of Thumb for the Detectability of Gravitational-Wave Bursts, arXiv e-print 1304.0210 (2013), arXiv:1304.0210 [gr-qc].
- [80] This all-sky search also could, in principle, detect IMBH mergers or highly eccentric binary mergers, but we do not evaluate those signal classes in this paper, instead leaving them for future dedicated analyses.
- [81] K.-C. Pan, M. Liebend  rfer, S. M. Couch, and F.-K. Thielemann, Stellar mass black hole formation and multimessenger signals from three-dimensional rotating core-collapse supernova simulations, *Astrophys. J.* **914**, 140 (2021).
- [82] J. M. Blondin, A. Mezzacappa, and C. DeMarino, Stability of standing accretion shocks, with an eye toward core-collapse supernovae, *Astrophys. J.* **584**, 971 (2003).
- [83] J. M. Blondin and A. Mezzacappa, The Spherical Accretion Shock Instability in the Linear Regime, *Astrophys. J.* **642**, 401 (2006).
- [84] T. Foglizzo, P. Galletti, L. Scheck, and H.-T. Janka, Instability of a Stalled Accretion Shock: Evidence for the Advective-Acoustic Cycle, *Astrophys. J.* **654**, 1006 (2007).
- [85] J. Powell and B. M  ller, Three-dimensional core-collapse supernova simulations of massive and rotating progenitors, *MNRAS* **494**, 4665 (2020).
- [86] D. Radice, V. Morozova, A. Burrows, D. Vartanyan, and H. Nagakura, Characterizing the Gravitational Wave Signal from Core-collapse Supernovae, *Astrophys. J. Lett.* **876**, L9 (2019).
- [87] J. Powell and B. M  ller, Gravitational wave emission from 3D explosion models of core-collapse supernovae with low and normal explosion energies, *MNRAS* **487**, 1178 (2019).
- [88] E. P. O'Connor and S. M. Couch, Exploring Fundamentally Three-dimensional Phenomena in High-fidelity Simulations of Core-collapse Supernovae, *Astrophys. J.* **865**, 81 (2018).
- [89] M. Obergaulinger and M.   . Aloy, Magnetorotational core collapse of possible GRB progenitors - I. Explosion mechanisms, *MNRAS* **492**, 4613 (2020).
- [90] K. Itagaki, Transient Discovery Report for 2023-05-19, Transient Name Server Discovery Report **2023-1158**, 1 (2023).
- [91] N. Raza, J. McIver, G. D  lya, and P. Raffai, Prospects for reconstructing the gravitational-wave signals from core-collapse supernovae with advanced ligo-virgo and the bayeswave algorithm, *Phys. Rev. D* **106**, 063014 (2022).
- [92] P. J. McMillan, The mass distribution and gravitational potential of the Milky Way, *MNRAS* **465**, 76 (2017).
- [93] M. Cautun, A. Ben  tez-Llambay, A. J. Deason, C. S. Frenk, A. Fattahi, F. A. G  mez, R. J. J. Grand, K. A. Oman, J. F. Navarro, and C. M. Simpson, The Milky Way total mass profile as inferred from Gaia DR2, *MNRAS* **494**, 4291 (2020).
- [94] D. Lopez, S. Tiwari, M. Drago, D. Keitel, C. Lazzaro, and G. A. Prodi, Prospects for detecting and localizing short-duration transient gravitational waves from glitching neutron stars without electromagnetic counterparts, *Phys. Rev. D* **106**, 103037 (2022), arXiv:2206.14515 [astro-ph.HE].
- [95] J. Abadie *et al.* (LIGO Scientific Collaboration), Search for gravitational waves associated with the August 2006 timing glitch of the Vela pulsar, *Phys. Rev. D* **83**, 042001 (2011).
- [96] R. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), Search for Gravitational-wave Transients Associated with Magnetar Bursts in Advanced LIGO and Advanced Virgo Data from the Third Observing Run, *Astrophys. J.* **966**, 137 (2024).
- [97] W. C. G. Ho, D. I. Jones, N. Andersson, and C. M. Espinoza, Gravitational waves from transient neutron star f -mode oscillations, *Phys. Rev. D* **101**, 103009 (2020).
- [98] R. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), Narrowband Searches for Continuous and Long-duration Transient Gravitational Waves from Known Pulsars in the LIGO-Virgo Third Observing Run, *Astrophys. J.* **932**, 133 (2022).
- [99] P. Caraveo, A. Luca, and R. Mignani, The Distance to the Vela Pulsar Gauged with Hubble Space Telescope Parallax Observations, *Astrophys. J.* **561**, 930 (2001).
- [100] R. N. Manchester, G. B. Hobbs, A. Teoh, and M. Hobbs, VizieR Online Data Catalog: ATNF Pulsar Catalog (Manchester+, 2005), VizieR Online Data Catalog, VII/245 (2005).
- [101] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, Equation of state of nucleon matter and neutron star structure, *Phys. Rev. C* **58**, 1804 (1998).
- [102] B. D. Lackey, M. Nayyar, and B. J. Owen, Observational constraints on hyperons in neutron stars, *Phys. Rev. D* **73**, 024021 (2006).

- [103] D. D. Doneva, E. Gaertig, K. D. Kokkotas, and C. Krüger, Gravitational wave asteroseismology of fast rotating neutron stars with realistic equations of state, *Phys. Rev. D* **88**, 044052 (2013).
- [104] L. Dunn, A. Melatos, C. M. Espinoza, D. Antonopoulou, and R. Dodson, A new small glitch in Vela discovered with a hidden Markov model, *MNRAS* **522**, 5469 (2023).
- [105] Fuentes, J. R., Espinoza, C. M., Reisenegger, A., Shaw, B., Stappers, B. W., and Lyne, A. G., The glitch activity of neutron stars, *A&A* **608**, A131 (2017).
- [106] M. Yu, R. N. Manchester, G. Hobbs, S. Johnston, V. M. Kaspi, M. Keith, A. G. Lyne, G. J. Qiao, V. Ravi, J. M. Sarkissian, R. Shannon, and R. X. Xu, Detection of 107 glitches in 36 southern pulsars, *MNRAS* **429**, 688 (2012).
- [107] C. M. Espinoza, A. G. Lyne, B. W. Stappers, and M. Kramer, A study of 315 glitches in the rotation of 102 pulsars, *MNRAS* **414**, 1679 (2011).
- [108] B. P. Abbott *et al.* (LIGO-Virgo-KAGRA Collaboration), Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA, *Living Reviews in Relativity* **23**, 3 (2020).