



BMJ Open Practice of oxygenation and respiratory support during fiberoptic bronchoscopy: the OxyFOB study protocol

Federico Longhini ,¹ Claudia Crimi,² Alberto Noto,³ Corrado Pelaia,¹ Zuhail Karakurt,⁴ Szymon Skoczyński ,⁵ José Pedro Boléo-Tomé,⁶ João Carlos Winck,⁷ Antonio M Esquinas,⁸ James Melhorn,⁹ Dan Corneci,¹⁰ Pavol Pobeha,¹¹ Vincenzo Bosco,¹ Eugenio Garofalo,¹ Andrea Bruni,¹ Gianmaria Cammarota,¹² Violeta Todorova,¹³ Mariangela Valentina Puci,¹⁴ Giovanni Sotgiu,¹⁴ Konstantinos Kostikas,¹⁵ Salvatore Maurizio Maggiore,¹⁶ Edoardo De Robertis,¹⁷ Begum Ergan,¹⁸ Giovanni Landoni,¹⁹ Rachele Simonte,¹⁷ Stefano Nava,²⁰ Paolo Navalesi,²¹ Raffaele Scala,²² OxyFOB study group

To cite: Longhini F, Crimi C, Noto A, *et al.* Practice of oxygenation and respiratory support during fiberoptic bronchoscopy: the OxyFOB study protocol. *BMJ Open* 2025;**15**:e104747. doi:10.1136/bmjopen-2025-104747

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2025-104747>).

FL and CC contributed equally.

Received 07 May 2025

Accepted 05 August 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Federico Longhini;
flonghini@unicz.it

ABSTRACT

Introduction Flexible bronchoscopy (FB) is widely used for diagnostic and therapeutic procedures in pulmonary medicine. However, FB can cause respiratory and haemodynamic complications, especially in patients with pre-existing lung and/or cardiovascular comorbidities. Despite the range of oxygenation and ventilatory approaches available to prevent these risks, evidence regarding their real-world application and clinical impact is limited. The OxyFOB study aims to assess the prevalence and outcomes of various oxygenation and ventilatory support strategies used during FB across Europe.

Methods and analysis The OxyFOB study is a large, prospective, international, observational cohort study which aims to involve over 10 000 FB procedures across European centres. Eligible participants include all adults undergoing FB for diagnostic, therapeutic or procedural indications. Data are collected via a standardised electronic case report form and encompass demographic information, procedural details and clinical outcomes. The primary endpoint is the prevalence of oxygenation and ventilatory support strategies: conventional oxygen therapy, high-flow oxygen therapy, continuous positive airway pressure, non-invasive ventilation and invasive mechanical ventilation. Secondary outcomes include periprocedural respiratory and haemodynamic events, patient comfort, dyspnoea and postprocedural complications. Statistical analyses include descriptive statistics, subgroup comparisons and multivariate logistic regression.

Ethics and dissemination The study has received ethical approval from the coordinating centre (protocol n. 22/2022 on the 20 January 2022, by the 'Comitato Etico Sezione Area Centro - Regione Calabria') and all participating sites. Informed consent is given from all patients or their legal representatives. Findings will be disseminated through peer-reviewed publications and presentations at international meetings. Data will be managed and made available on reasonable request to support further research.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Large sample size enhances the prospect of wide validity across multiple European centers and settings, to ensure adequate representation across all subpopulations according to the type of flexible bronchoscopy (FB) procedure (eg, bronchoalveolar lavage, brushing, TransBronchial Needle Aspiration (TBNA), Endobronchial Ultrasound (EBUS)).
- ⇒ Real-world observational design reflects routine clinical practice and variability in FB procedures, especially with regard to preliminary guidance to optimise oxygenation and ventilatory strategies for different patient subpopulations.
- ⇒ Comprehensive data collection includes both clinical outcomes and patient-reported measures (eg, dyspnoea, comfort).
- ⇒ Centralised data quality oversight minimises errors and ensures consistency.
- ⇒ As an observational study, causal relationships cannot be firmly established due to potential confounding and selection bias.

Trial registration number ClinicalTrials.gov ID: [NCT05681962](https://clinicaltrials.gov/ct2/show/study/NCT05681962). Registered January 2023.

INTRODUCTION

First described by Ikeda *et al*,¹ flexible bronchoscopy (FB) is a medical procedure used to visualise the lower airways and perform diagnostic or therapeutic interventions in a large spectrum of respiratory diseases. FB is employed in the diagnostic approach to persistent cough, management of lung infections and evaluation of non-infective infiltrates through procedures like bronchoalveolar lavage (BAL), endobronchial brushing and both bronchial and transbronchial lung biopsies in neoplastic and non-malignant



lung disorders, removing mucus plugs or foreign bodies and addressing airway bleeding.

With the proper precautions, FB is a generally safe procedure.² Studies report a mortality rate ranging between 0.01% and 0.04% and a major complication rate between 0.08% and 0.3%.³⁻⁵ However, most patients undergoing FB have underlying conditions that can compromise gas exchange, such as pneumonia, interstitial lung diseases or neoplasms; the arterial partial pressure of oxygen does in fact often decrease to varying degrees, which may increase the risk of respiratory distress.^{6,7} Desaturation may be secondary to a wide array of conditions, such as: alveolar derecruitment, ventilation-perfusion (V/Q) mismatch and alveolar hypoventilation,⁸⁻¹⁰ the latter mainly due to alterations in respiratory pattern as a side effect of sedative drugs¹¹⁻¹⁴ and their effects on the critical closing pressure of the upper airways, which may collapse.^{15,16} Also worth mentioning, intraprocedural suctioning reduces end-expiratory lung volume, which may therefore reduce lung compliance and cause V/Q mismatch and venous admixture.¹⁷⁻¹⁹ The occurrence of FB-related complications is greater during interventional procedures such as bronchial and transbronchial biopsies; moreover, individual team expertise is a further, independent variable influencing the incidence of FB-related complications in clinical practice.^{8,20}

In non-intubated patients, the fibrescope occupies approximately 10% of the tracheal cross-sectional area and 15% of that of the cricoid ring. This partial obstruction increases airway resistance, thereby elevating the patient's work of breathing during the procedure.¹⁷ In patients with bronchial hyperreactivity, peri-procedural bronchospasm can occur as a complication of FB, potentially leading to airway narrowing and respiratory distress.²¹

These respiratory changes typically resolve after FB; however, in cases of severe parenchymal lung diseases, recovery may take anywhere from 15 min to several hours. This delay may necessitate oxygen therapy or escalation of postprocedural respiratory support and unexpected hospital and/or intensive care unit (ICU) admission.^{8,17}

During FB, complex transient and contrasting haemodynamic modifications may also occur: if on one hand cardiac output may increase by up to 50% due to sympathetic stimulation, typically returning to baseline within 15 min after the procedure,^{17,22} on the other changes in intrathoracic pressure, resulting from increased inspiratory effort, affect venous return and afterload, potentially reducing cardiac output.²³ Those events can precipitate heart failure in patients with underlying cardiovascular conditions, and ECG changes may occur in up to 21% of awake patients over 60 years of age.²⁴ Different strategies of oxygenation and ventilatory support have been applied in spontaneously breathing patients and compared in patients receiving FB to prevent respiratory failure or worsening of gas exchange.²⁵ However, despite a relevant number of available studies, most investigations focus on physiological parameters rather than clinically relevant

outcome variables with heterogeneous populations with respect to severity, type of procedure and supportive means.²⁵

To assess this gap, we therefore designed this multi-centre international prospective observational study to assess the real-life prevalence of oxygenation and ventilatory strategies during FB in different case scenarios, stratifying patients by baseline comorbidities, type of FB procedure and hospital setting, in order to explore whether specific subgroups of patients may benefit from distinct oxygenation or ventilatory strategies during the procedure. In addition, we will evaluate the safety, tolerability and clinical outcomes across different oxygenation and ventilatory support strategies.

METHODS AND ANALYSIS

Study design

This is a prospective international observational cohort study conducted across Europe. The study protocol is designed and reported in this article according to the Strengthening the Reporting of Observational Studies in Epidemiology Statement.²⁶ The study was approved by the Ethics Committee of the coordinating centre (protocol n. 22/2022 on 20 January 2022, by the 'Comitato Etico Sezione Area Centro - Regione Calabria'). All participating centres obtained the local ethical committee approval. This study was prospectively registered on ClinicalTrials.gov (NCT05681962; principal investigator: Federico Longhini) in January 2023. The flow chart of the Oxy-FOB study is depicted in figure 1.

Recruitment started on 15 February 2023. The study is currently ongoing with 6852 patients recruited patients in 63 active centres across Europe as per 7 May 2025. Additional centres are being included to enhance representativeness and further strengthen the study's conclusions. We expect to complete the recruitment by the end of 2026.

Study population

We consider eligible all consecutive adult (ie, ≥ 18 years old) patients requiring a FB procedure for the main diagnostic, therapeutic or procedural indications. Diagnostic indications include: evaluation of pulmonary infections, investigation of lung masses or nodules, assessment of interstitial lung diseases, diagnosis of central airway obstruction, evaluation of haemoptysis to localise bleeding and determine its cause, biopsy of mediastinal or hilar lymphadenopathy and the diagnosis and therapeutic lavage of pulmonary alveolar proteinosis.

Therapeutic indications include: airway clearance and secretion management, foreign body removal, balloon dilation of airway stenosis, endobronchial valve placement for lung volume reduction in emphysema, management of haemoptysis, and therapeutic lavage for alveolar proteinosis.

Procedural indications include: BAL for diagnostic purposes, as well as guidance for transbronchial lung biopsy or needle aspiration.




Figure 1 Study flow from patient enrolment to outcome assessment during and after the FB procedure. FB, flexible bronchoscopy; ICU, intensive care unit.

Included subjects will be screened consecutively, and written informed consent will be obtained from all patients, next of kin or legal representatives, as per national laws. Exclusion criteria include pregnancy, breastfeeding, paediatric patients (ie, <18 years old) and patients for whom informed consent cannot be obtained.

Study outcomes and measures

Primary endpoint

The primary outcome of this study is to assess the prevalence of oxygenation and ventilatory strategies during FB in different case scenarios, conducting subgroup analyses based on baseline comorbidities, type of FB procedure and hospital setting. For this purpose, we defined the following oxygenation and ventilatory strategies as follows:

1. Conventional oxygen therapy (COT): administration of oxygen through nasal prongs, oxygen mask (with or without reservoir) and Venturi mask.
2. High flow oxygen therapy (HFOT): administration of high flows (up to 60L/min) of air/oxygen mixtures, heated at temperatures ranging from 31 to 37°C and fully humidified (up to 44mg H₂O/L), providing an inspired oxygen fraction (FiO₂) ranging from 21% to 100%.
3. Continuous positive airway pressure (CPAP): application of positive end-expiratory pressure (PEEP) throughout the whole respiratory cycle with a ventilator, flow generator or Venturi system, through interfaces including, but not limited to, mask or helmet.

4. Non-invasive ventilation (NIV): application of a PEEP and an inspiratory pressure support (PS) triggered by the patient, delivered by a ventilator through a mask or helmet.
5. Invasive mechanical ventilation (IMV): application of a ventilatory assistance in controlled or assisted modes through an endotracheal or tracheostomy tube.

Secondary endpoints

Secondary endpoints of the study include evaluating the safety and tolerability of the procedure through the recording of major periprocedural respiratory and haemodynamic adverse events, as well as patient comfort and dyspnoea during the procedure and collecting clinical outcomes associated with different oxygenation and ventilatory support strategies.

Major periprocedural respiratory and haemodynamic adverse events include:

- ▶ Peripheral desaturation (defined as peripheral oxygen saturation (SpO₂) <90% for at least 10s) or severe desaturation (defined as SpO₂ <80% for any duration).
- ▶ Need for interruption of FB.
- ▶ Haemodynamic instability, defined as the occurrence of hypotensive (ie, a systolic blood pressure <90 mm Hg) or hypertensive (ie, systolic blood pressure >140 mm Hg) events.
- ▶ New-onset cardiac arrhythmias requiring treatment, myocardial infarction and/or electrocardiographic ST alterations.



- Occurrence of neurological events, defined as ischaemic or haemorrhagic stroke, transient ischaemic attack and/or seizures.^{9 25 27}

Clinical outcomes consist of:

1. Need for additional respiratory support beyond that initially applied.
2. Occurrence of procedural complications (including, but not limited to, airway bleeding, pneumothorax, pneumomediastinum, bronchial perforation, bronchospasm or laryngospasm).
3. Need for unplanned admission to the emergency department (ED), hospital ward or high-dependency unit/ICU.
4. For hospitalised patients, in-hospital and/or ICU length of stay and mortality.

Measures

Baseline and demographic data include patient age, gender, height, type of admission (outpatient, inpatient, ICU or ER), comorbidities, Charlson Comorbidity Index, smoking habit, previous need for long-term oxygen therapy or home NIV or tracheoventilation. In addition, we will register the baseline values of SpO₂, systolic and diastolic blood pressure, baseline arterial blood gas analysis (ABGs) if available, type of FB performed and indication for the procedure.

During the procedure, we will record the lowest SpO₂, the occurrence of desaturation or severe desaturation, the need for interruption of FB, lowest and highest intraprocedural systolic blood pressure, new-onset cardiac arrhythmias or pathological conditions and neurological events.

At the end of the FB, vital parameters and ABGs (if available), presence of dyspnoea and patient's comfort during the procedure will also be recorded. Both dyspnoea and patient's comfort will be assessed using the 11-point Numeric Rating Scale, as previously reported.

Since relying solely on clinical indicators can be insufficient to detect early or evolving hypercapnia, particularly in patients with chronic respiratory conditions, we also collected the arterial partial pressure of CO₂, when available as per clinical practice. This approach is critical for accurately assessing ventilatory adequacy, especially when comparing support strategies such as HFOT, CPAP and NIV during bronchoscopy. Since the study protocol was designed in 2022, only a few (<5%) centres were equipped with end-tidal or transcutaneous CO₂ monitoring at that time; therefore, we chose not to include it as a data point to be collected.

Patients will be asked to define the perceived severity of their condition by providing a number between 0 (no discomfort/no dyspnoea) and 10 (worst possible discomfort/worst possible dyspnoea) on a large printed scale including numbers and descriptors.²⁸⁻³¹

In addition, we will also record the need for respiratory support escalation beyond that initially applied and the occurrence of complications as previously described. These outcomes will be checked and recorded within the first 48 hours from FB or hospital discharge.

All intraprocedural adverse events will also be analysed to assess their impact on post-procedural clinical outcomes, including the need for respiratory support escalation, unexpected ED, ward or ICU admission, hospital/ICU lengths of stay and mortality.

Data collection

Data will be collected on an electronic case report form (eCRF) based on the Research Electronic Data Capture secure web application. The eCRF consists of five sections:

1. Demographic and pre-procedural physiological parameters.
2. Details of FB procedure.
3. Intraprocedural parameters.
4. Postprocedural parameters.
5. Clinical outcomes.

Local investigators will screen and report all the FB procedures.

Data quality framework

A dedicated team of investigators will ensure data quality by routinely assessing the integrity, completeness, consistency and accuracy of the existing data.³² All data will undergo thorough verification, and any instances of missing data, potential errors or ambiguous values will prompt contact with local investigators for resolution, ensuring robust and reliable data quality control.³²

Statistical analysis

Statistical analysis is designed by an independent team of expert statisticians, in collaboration with the steering committee. We plan on collecting data from a convenience sample of 10000 procedures. The sample characteristics will be described using absolute frequencies and percentages (qualitative variables), mean and SD or median and IQR (quantitative variables), as appropriate. Data normality will be assessed using the Shapiro-Wilk test. Prevalence will be determined and reported with the corresponding 95% CIs. Additionally, the incidence and type of major periprocedural respiratory and haemodynamic adverse events will be calculated along with their 95% CIs.

Differences in quantitative variables between two groups will be evaluated using the unpaired Student's t-test or the corresponding non-parametric test, after assessing the relative assumptions. In the case of groups >2, these differences will be assessed using analysis of variance or Kruskal-Wallis test. Differences in qualitative variables will be evaluated using Pearson's χ^2 test or Fisher's exact test. Multivariate logistic regression modelling will be performed to determine factors associated with oxygenation and ventilatory supports; any variable with significant univariate test or clinical relevance will be selected as a candidate for multivariate analysis. The level of statistical significance will be set at a p value <0.05.

DISCUSSION

This large prospective observational study aims to describe the current clinical practices on oxygenation and ventilatory strategies during FB across Europe in various clinical scenarios. The results will provide insights into the use of different oxygenation and ventilatory support strategies for patients with different indications for FB and varying severity of respiratory failure. By analysing the occurrence of respiratory and haemodynamic impairments, as well as clinical outcomes, this study will identify key patterns across subpopulations supported by COT, HFOT, CPAP, NIV or IMV.

As an observational cohort study, this study can estimate the incidence of clinically meaningful outcomes such as respiratory and haemodynamic impairments, adverse events and unexpected hospital or ICU admission³³ and will therefore provide data useful to potentially suggest the use of one technique over another. The findings could modify clinical practices and lay the groundwork for future randomised controlled trials (RCTs) to validate the efficacy of specific interventions.³⁴

To the best of our knowledge, this is the largest ongoing study ever conducted in this population, while the current existing literature predominantly consists of studies with sample sizes of fewer than 100 patients.²⁵

Respiratory and haemodynamic impairments during FBs are common events that can worsen the patients' clinical conditions, potentially leading to unplanned hospital admission during out-patient procedures, or ICU admission and escalation of respiratory support after the procedure.²⁵ A previous systematic review showed that patients with poorer baseline lung function have higher oxygen requirements during FB and an increased risk for post-procedural worsening of respiratory failure.²⁵ The pooled data analysis revealed that HFOT performs better than COT in terms of oxygenation outcomes for patients with lower oxygen requirements, while NIV appeared to be more effective than HFOT in those with more severe respiratory failure. However, insufficient evidence prevents definitive recommendations regarding the superiority of one oxygenation strategy over another. Nevertheless, improving intra-procedural oxygenation remains a critical safety concern for patients undergoing FB and has the potential to improve key clinical outcomes, such as reducing the need for hospital or ICU admission due to post-procedural respiratory failure, though these benefits require further investigation.²⁵ Currently, other RCTs are ongoing to compare different strategies in specific patient populations.^{35 36}

Strengths and limitations

Our study has several strengths and limitations worth discussion.

One of the main strengths is the design that reflects real-world practices and outcomes, fully addressing our primary aim. The large sample size enables pooling of data from multiple centres across Europe, ensuring the inclusion of patients from diverse

geographic, cultural and healthcare settings, thereby enhancing the generalisability of the findings.³⁷ The study was designed to ensure adequate representation across all subpopulations according to the type of FB procedure (eg, BAL, brushing, TBNA, EBUS), providing each group with an adequate number of patients.

Although RCTs remain the standard of reference for establishing causality, observational studies provide valuable insights into real-world practice, can help to identify the effect of interventions in specific subpopulations and can serve as a valid reference when planning future RCTs.^{38 39} Therefore, once this study is completed and published, it will provide the basis for RCTs comparing different oxygenation or ventilatory support techniques in specific patient populations, with the aim of reducing the occurrence of respiratory and haemodynamic adverse events during FB. Another strength of this study is the presence of a data quality team that routinely checks the collected data to reduce the possibility of missing or low-quality data.³²

Finally, this study is highly innovative, providing preliminary guidance on optimising oxygenation and ventilatory strategies for different patient subpopulations. The findings could significantly influence and improve clinical practices, helping clinicians to refine their approaches, reduce costs associated with complications and minimise unexpected hospital or ICU admissions.

Concerning limitations, the observational design may be more susceptible to confounding factors, making it challenging to establish causal relationships. Moreover, not all patients may be enrolled, leading to potential selection bias. However, through careful implementation of rigorous data quality measures and the use of advanced statistical methods to adjust for biases and confounders, many of the conventional limitations of observational studies can be mitigated.⁴⁰ Indeed, a selection bias in participating centres may have occurred, limiting the generalisability of the findings. However, participating centres are representative of different levels of care, settings and geography. Furthermore, this study will not gather information on the direct long-term consequences of periprocedural adverse events on specific patient outcomes, as the aim of the study was to prospectively collect data on short-term adverse events.

A further potential limitation of our study is related to the collection of FB data both from stable and acutely unstable critically ill patients in different settings (from the bronchoscopy unit to the ICU), driven by heterogeneous indications and expected outcomes. However, we believe that this may also be considered as a strength, as it contributes to the ambitious goal of building a comprehensive 'road map' for how different physicians manage FB in real-world

practice, providing insights on safer oxygenation tools and tailored approaches for each clinical scenario.

Ethics and dissemination

The study received approval from the Ethical Committee of the coordinating centre (protocol no. 22/2022, approved on 20 January 2022, by the 'Comitato Etico Sezione Area Centro - Regione Calabria') and, in accordance with local regulations, by all relevant local Ethical Committees. Informed consent is obtained from all participants or their consultee before enrolment. For participants lacking capacity, consent is sought from a personal or nominated consultee. The study adheres to good clinical practice guidelines and current data protection regulations, ensuring the ethical conduct of research and the protection of participant rights. The results of the study will be disseminated at the end of enrolment through a manuscript that will be submitted for publication in an international peer-reviewed scientific journal.

Author affiliations

¹Department of Medical and Surgical Sciences, Magna Graecia University of Catanzaro, Catanzaro, Italy

²Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy

³Department of Human Pathology of the adult and evolutive age "Gaetano Barresi", University of Messina, Messina, Italy

⁴University of Health Sciences Süreyyapaşa Chest Diseases and Thoracic Surgery Teaching and Research Hospital, Istanbul, Turkey

⁵Medical University of Silesia, Ruda Slaska, Poland

⁶Hospital Prof. Doutor Fernando Fonseca, ULS Amadora-Sintra, Amadora, Portugal

⁷Universidade do Porto Faculdade de Medicina, Porto, Portugal

⁸Intensive Care Unit, Hospital Morales Meseguer, Murcia, Spain

⁹Respiratory Medicine Unit and NIHR Oxford Biomedical Research Centre, Nuffield Department of Clinical Medicine, Oxford University, Oxford, UK

¹⁰Central Military Emergency University Hospital, University of Medicine and Pharmacy Carol Davila Bucharest, Bucharest, Romania

¹¹Louis Pasteur University Hospital, Kosice, Slovakia

¹²Department of Translational Medicine, University of Novara Faculty of Medicine and Surgery, Novara, Italy

¹³Anesthesia and ICU, Centre Hospitalier Annecy Genevois, Épagny Metz-Tessy, France

¹⁴Clinical Epidemiology and Medical Statistics Unit, Department of Medicine, Surgery and Pharmacy, Sassari University Hospital, Sassari, Italy

¹⁵Respiratory Medicine Department, University of Ioannina Faculty of Medicine, Ioannina, Greece

¹⁶Department of Anesthesiology and Intensive Care Medicine - Miulli University Hospital, LUM University of Casamassima, Acquaviva delle fonti, Italy

¹⁷Department of Medicine and Surgery, University of Perugia, Perugia, Italy

¹⁸Department of Pulmonary and Critical Care, Dokuz Eylul University Faculty of Medicine, Izmir, Turkey

¹⁹Vita-Salute San Raffaele University, Milano, Italy

²⁰Alma Mater Studiorum University of Bologna, Bologna, Italy

²¹Department of Medicine, University of Padua, Padova, Italy

²²San Donato Hospital, Arezzo, Italy

Acknowledgements We also acknowledge members of the OxyFOB Study group as authors.

Collaborators The OxyFOB Study group: Zuhail Karakurt, Baran Gundogus, Gul Erdal Donmez, Dilek Ernam (University of Health Sciences Süreyyapaşa Chest Diseases and Thoracic Surgery Teaching and Research Hospital, Istanbul, Turkey); Sergio Campinha, Ana Oliveira, Maria José Teixeira (Bronchology Unit, Pulmonology Department, Unidade Local de Saúde de Gaia-Espinho, Portugal);

Francisco Correia Gouveia, Catarina Torres Monteiro (Anesthesiology Department, Unidade Local de Saúde de Gaia-Espinho, Portugal); Nurdan Şimşek Veske, Sedat Altın, Gülşah Günlüoğlu, Efsun Gonca Chousein (University of Health Sciences Yedikule Chest Diseases and Thoracic Surgery Teaching Hospital, Pulmonology Department, Yedikule, Turkey); Hüseyin Arıkan, Sait Karakurt (Marmara University School of Medicine, Pulmonology Department, Marmara, Turkey); Donato Lacedonia, Ruggiero Torracco, Pasquale Tondo (Department of Medical and Surgical Sciences, University of Foggia, Italy); Szymon Bialka, Piotr Palaczynski, Justyna Danel (Department of Anaesthesiology and Intensive Care, Faculty of Medical Sciences in Zabrze, Medical University of Silesia, Katowice, Poland); Piotr Jankowiak, Magdalena Kędra, Magdalena Bednarska (Department of Pneumology, Medical University of Gdansk, Gdansk, Poland); Elisiana Carpagnano, Esterina Boniello, Andrea Portacci (Insitute of Respiratory Disease, Department of Translational Biomedicine and Neuroscience, University "Aldo Moro", Bari, Italy); Simone Scarlata, Panaiotis Finamore, Davide Fontata (Department of Internal Medicine, Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy); Aleksander Kania, Natalia Celejewska-Wojcik, Klaudia Rygiel (2nd Department of Medicine, Department of Pulmonology, Faculty of Medicine, Jagiellonian University Medical College, Krakow, Poland); Ewelina Tobiczak, Daria Springer, Szczepan Cofa (Department of Pulmonology, Allergology and Pulmonary Oncology, Poznan University of Medical Sciences, Poznan, Poland); Jessica Maugeri, Agrippino Bellissima (Anesthesia and Intensive Care Unit, ARNASS Garibaldi, Catania, Italy); Mario Tamburrini, Alberto Papi, Luca Morandi (Pulmonary Division, University of Ferrara, St Anna University Hospital, Ferrara, Italy); Cosimo Franco, Sara Chiesa, Nicolò Verti (Interventional Pulmonology Unit, Pulmonology Department/Respiratory Intensive Care, Piacenza, Italy); Michał Zieliński (Katedra i Klinika Chorób Płuc Gruzlicy, Wydział Nauk Medycznych w Zabrze, Śląski Uniwersytet Medyczny w Katowicach, Poland); Michele Mondoni, Carmelo Intravaia, Paolo Carlucci (Respiratory Unit, ASST Santi Paolo e Carlo, Milan, Italy); Demet Karnak, Aydin Çiledağ, Aslihan Gürün Kaya (Ankara University Medical Faculty, Department of Chest Disease, Cebeci-Ankara, Turkey); Giacomo Ghinassi, Luca Guidelli (Pulmonology and RICU, S Donato H, Cardio-toraco-neurovascular Dept Usl Toscana Sudest, Arezzo Italy); Pierachille Santus, Dejan Radovanovic, Marina Saad (Università degli Studi di Milano, Respiratory Diseases Unit, "L. Sacco" University Hospital, Milan, Italy); Catarina Figueiredo Roquete, Ricardo Petinga Fortes (Hospital Prof. Doutor Fernando Fonseca - ULS Amadora-Sintra, Amadora, Portugal); Konstantinos Porpodis, Ioanna Tsiouppou, Kalliopi Lagoudi (Aristotle University of Thessaloniki, Pulmonary department, 'G. Papanikolaou' hospital, Exohi, Thessaloniki); Girolamo Pelaia, Giuseppina Marrazzo, Nicola Montenegro (Respiratory Medicine Unit, University "Magna Graecia" of Catanzaro, Catanzaro, Italy); Davide Biondini, Marco Damin, Paolo Spagnolo (Department of Cardiac, Thoracic, Vascular Sciences and Public Health, University of Padova, Padova, Italy); João Oliveira Rodrigues, Mário Oliveira Pinto, Dionísio Castro Maia (ULS São José e ULS Loures / Odivelas, Portugal); Fotios Sampsonas, Argyrios Tzouvelekis, Georgios Tsirikos (Respiratory Department, Patras University Hospital, Patras, Greece); Stelios Loukides, Thomas Raptakis, Vasileios Papavasileiou (2nd Respiratory Medicine Department of University General Hospital (U.G.H.) "ATTIKON", Chaidari, Greece); Rosanna Vaschetto, Alessandro Tonino (Dipartimento di medicina Traslationale, Università del Piemonte Orientale, Novara, Italy); Filippo Patrucco (AOJ Maggiore della Carità, Malattie dell'apparato respiratorio, Novara, Italy); Ricardo Estevão Gomes, Jorge Soares, Maria João Santana (Serviço de Pneumologia, Unidade Local de Saúde Almada-Seixal, Hospital Garcia de Orta, Almada, Portugal); Giuseppe Neri, Angela Corea, Antonio Caroleo, Antonio Grande, Deborah Veltri, Giulia Costumato, Federica Di Lorenzo, Giuseppe Bonadio, Zaninni Caroleo, Giuseppe Guerriero, Giuseppe Mazza, Federica Mellace, Giulia Perrelli, Silvia Riillo, Giuseppina Ruocco, Antonio Camastra, Giuseppe Gaetano, Aldo Mesiti, Giusy Guzzi, Helenium Mastrangelo (Department of Medical and Surgical Science, Magna Graecia University, Catanzaro, Italy); Lara Pisani, Maria Laura Vega Pittao (Respiratory and Critical Care Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Alma Mater Studiorum, Department of Medical and Surgical Sciences (DIMEC), University of Bologna, Italy); Grigorios Stratakos, Evangelia Koukaki, Nektarios Anagnostopoulos (1st Respiratory Medicine Department, NK University of Athens, "Sotiria" Hospital, Greece); Roberto Dongilli, Emanuele Stirpe, Johanna Köhl (Division of Respiratory Diseases and Intermediate Respiratory Intensive Care Unit, Central Hospital of Bolzano, Bolzano, Italy); Peter Skyba, Eva Sokolová (Department of Respiratory medicine and Tuberculosis, P.J. Šafárik University, Medical faculty and L. Pasteur university hospital in Košice, Slovakia); Maria Alvarenga Santos, Margarida Aguiar (Hospital Beatriz Ângelo, Loures, Portugal); Nikos Zias, Ioannis Papadakis (Naval Hospital, Athens, Greece); Pierpaolo Terragni, Laura Pistidda, Roberto Castagna (Anesthesia and General Intensive Care Unit, Department of Medicine, Surgery and Pharmacy, University of Sassari, Italy); Mihai Popescu (Carol Davila University of Medicine and Pharmacy, Anaesthesia and Intensive Care Department, Bucharest University Emergency

Hospital, Bucharest, Rumania); Paola Rogliani, Emanno Puxeddu, Chiara Frugoni (Unit of Respiratory Medicine, Department of Experimental Medicine, University of Rome "Tor Vergata", Rome, Italy); Salvatore Notaro, Eugenio Piscitelli (UOSD Intensive Care and ECMO, Colli Hospital Company, Mondaldi Hospital, Naples, Italy); Elena Carrasco González, Áurea Higón Cañigral (Intensive Care Unit and Non Invasive Ventilatory Unit, Hospital General Universitario Morales Meseguer, Murcia, Spain); Stefano Turi, Marta Mucchetti, Alessandro Belletti, Giorgia Volontero, Sabina Sommaruga, Roberta Navarra (Department of Anaesthesia and Intensive Care, IRCCS San Raffaele Institute, Milan, Italy); Andrea Alessia Nardo, Giorgio Morana (Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy); Giulia Palmiotto, Antonio Gabriele Iuliano (Department of Pulmonology, ASST - Valle Olona, Busto Arsizio, Italy); Hélder Novais Bastos (Department of Pulmonology, Unidade Local de Saúde de São João, Faculdade de Medicina da Universidade do Porto, Porto, Portugal); Sabino Scolletta, Cesare Biuzzi (Department of Medical Science, Surgery and Neurosciences, Anesthesia and Intensive Care Unit, University Hospital of Siena, Siena, Italy); Claudio Sorino, Denisa Hida (Department of Pulmonology, Sant'Anna Hospital of Como, Italy); Faculty of Medicine and Surgery, University of Insubria, Varese, Italy); Eugenio Arrighi, Paolo Gambardella, Graziella Perri (Department of Pulmonology, "Giovanni Paolo II" Hospital, ASP Catanzaro, Lamezia Terme, Italy); Andrea Cortegiani, Mariachiara Ippolito (Department of Anesthesia, Analgesia, Intensive Care and Emergency, University Hospital Policlinico Paolo Giaccone & Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Palermo, Italy); Francesco Alessandri, Vlad Cristian Sanda, Veronica Zullino (Department of General and Specialistic surgery, "Sapienza" University of Rome, Italy); Monica Rocco, Silvia Fiorelli (Dipartimento di scienze medico chirurgiche e medicina traslazionale, Sapienza University, Rome, Italy); Lisete Rolo Nunes (Medicina Interna e Competência em Emergência Médica, Serviço de Medicina Intensiva do Hospital Prof. Dr. Fernando da Fonseca, Amadora, Portugal); Cesare Gregoretti, Giovanni Misseri, Matteo Piattoli (Intensive Care Unit, Fondazione G. Giglio, Cefalù, Unicamillus International University, Roma, Cefalù, Italy); Isacco Curto (Department of Neurosurgical Anesthesia and Intensive Care, Santa Chiara Regional Hospital, Trento, Italy); Marco Tescione, Eugenio Vadalà, Annalisa Piccolo, Selma Mammine (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Maciej Gnass, Piotr Gniady (Endoscopy Unit, Pulmonary Hospital, Zakopane, Poland); Paschalis Steiropoulos, Fotios Drakopanagiotakis (Department of Pneumology, Medical School, Democritus University of Thrace, Alexandroupolis, Greece); Lucrezia Gianozzi (Pulmonology Department, Hospital Insular, Las Palmas de Gran Canaria, Spain); Christos Chronis (Department of Respiratory Medicine, University Hospital of Ioannina, Ioannina, Greece); Luca Novelli (Respiratory Medicine Unit, ASST Papa Giovanni XXIII, Bergamo, Italy); Kostas Katsoulis (Pulmonary & Microbiology Department, General Army Hospital, Thessaloniki, Greece); Alessandro Giuseppe Fois (Department of Respiratory Diseases, University Hospital Sassari (AOU), Sassari, Italy); Aleksandra Oraczewska (Department of Pneumology, School of Medicine in Katowice, Medical University of Silesia, Katowice, Poland).

Contributors FL, CC, CP, JCW, BE, SN, PN and RS were involved in the conceptualisation and writing of the protocol and drafted the initial version of this manuscript. CP and RS drafted figures, tables and text of the manuscript. AN serves as data manager of the study and is responsible for the electronic CRF. ZK, SS, JPBT, JCW, AME, JM, DC, PP, VT, KK, SMM, EDR, GL, SN and PN serve as National Coordinators and contact persons for all centres across Europe; they are responsible for overseeing protocol development, study setup and trial management. VB, EG, AB, GC and RS are responsible for the data quality check. MVP and GS provide statistical support and developed the statistical analysis plan. FL is the principal investigator, conceived the study concept and developed the initial protocol; he is also the guarantor of the study. All authors are involved in trial conduct, participant recruitment and data collection; they are also considered scientific experts involved in the field. All authors contributed to manuscript drafting and have read and approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests Dr. FL contributed to the development of a new device for non-invasive ventilation not discussed in the present study (European Patent number 3320941). He also received honoraria from Draeger, Intersurgical, Fisher & Paykel, AOP and Medicaire. Dr CC received speaking honoraria from Fisher & Paykel, Philips, Resmed and Vitulaire. Dr. CP has received lecture fees and advisory board fees from AstraZeneca, GlaxoSmithKline, Sanofi-Regeneron, Chiesi. JPBT received travel grants and economical support to attend training courses from Pentax Medical Iberia. Prof. JCW received from Philips-Respironics (fees for lecturing, participation in advisory board, reimbursement for Congress), ResMed

(fees for lecturing) and Fisher-Paykel (fees for lecturing). Dr. PP received speaking honoraria from Angelini Pharma, Philips, Berlin-Chemie, Astra Zeneca and CSL Behring. Dr GC received travel grants from Draeger. GS received speaking fee from Pfizer, Quiagen, AZ and INSMED. Dr. KK received honoraria for presentations and/or consultancy fees from AstraZeneca, Berlin-Chemie, Boehringer Ingelheim, Chiesi, ELPEN, GSK, Guidotti, Menarini, Pfizer, Sanofi, and Specialty Therapeutics. His department has received funding and/or grants from AstraZeneca, Boehringer Ingelheim, Chiesi, ELPEN, GSK, Menarini. He also worked with AstraZeneca as Global Medical Head Respiratory Biologics and he is a member of the GOLD Assembly. SMM received lecture's fee from GE Healthcare. Dr. EDR received honoraria lectures from Draeger, Fisher & Paykel, Baxter and GE. Dr. BE received honoraria lectures from Fisher & Paykel and Breas. Dr. PN received grants/research equipment from Draeger, Intersurgical SPA, and Gilead and speaking fees from Getinge, Mindray, Intersurgical SPA, Gilead, GSK and Draeger. The remaining authors have no conflict of interest to disclose related to the present work.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Federico Longhini <https://orcid.org/0000-0002-6970-7202>
Szymon Skoczynski <https://orcid.org/0000-0003-1796-7659>

REFERENCES

- Ikeda S, Yanai N, Ishikawa S. Flexible bronchofiberscope. *Keio J Med* 1968;17:1–16.
- Zavala DC. Diagnostic fiberoptic bronchoscopy: Techniques and results of biopsy in 600 patients. *Chest* 1975;68:12–9.
- Smyth CM, Stead RJ. Survey of flexible fiberoptic bronchoscopy in the United Kingdom. *Eur Respir J* 2002;19:458–63.
- Suratt PM, Smiddy JF, Gruber B. Deaths and complications associated with fiberoptic bronchoscopy. *Chest* 1976;69:747–51.
- Credle WF, Smiddy JF, Elliott RC. Complications of fiberoptic bronchoscopy. *Am Rev Respir Dis* 1974;109:67–72.
- Antonelli M, Conti G, Riccioni L, et al. Noninvasive Positive-Pressure Ventilation Via Face Mask During Bronchoscopy With BAL in High-Risk Hypoxemic Patients. *Chest* 1996;110:724–8.
- Goldstein RA, Rohatgi PK, Bergofsky EH, et al. Clinical role of bronchoalveolar lavage in adults with pulmonary disease. *Am Rev Respir Dis* 1990;142:481–6.
- Du Rand IA, Blaikley J, Booton R, et al. Summary of the British Thoracic Society guideline for diagnostic flexible bronchoscopy in adults. *Thorax* 2013;68:786–7.
- Longhini F, Pelaia C, Garofalo E, et al. High-flow nasal cannula oxygen therapy for outpatients undergoing flexible bronchoscopy: a randomised controlled trial. *Thorax* 2022;77:58–64.
- Evans EN, Ganeshalingam K, Ebdon P. Changes in oxygen saturation and transcutaneous carbon dioxide and oxygen levels in patients undergoing fiberoptic bronchoscopy. *Respir Med* 1998;92:739–42.
- Strohleit D, Galetin T, Kosse N, et al. Guidelines on analgesedation, monitoring, and recovery time for flexible bronchoscopy: a systematic review. *BMC Pulm Med* 2021;21:198.
- Lee H, Choe YH, Park S. Analgesedation during flexible fiberoptic bronchoscopy: comparing the clinical effectiveness and safety of remifentanyl versus midazolam/propofol. *BMC Pulm Med* 2019;19:240.
- Costa R, Navalesi P, Cammarota G, et al. Remifentanyl effects on respiratory drive and timing during pressure support ventilation and neurally adjusted ventilatory assist. *Respir Physiol Neurobiol* 2017;244:10–6.
- Vaschetto R, Cammarota G, Colombo D, et al. Effects of Propofol on Patient-Ventilator Synchrony and Interaction During Pressure Support Ventilation and Neurally Adjusted Ventilatory Assist*. *Crit Care Med* 2014;42:74–82.
- Eastwood PR, Platt PR, Shepherd K, et al. Collapsibility of the Upper Airway at Different Concentrations of Propofol Anesthesia. *Anesthesiology* 2005;103:470–7.



- 16 Shteamer JW, Dedhia RC. Sedative choice in drug-induced sleep endoscopy: A neuropharmacology-based review. *Laryngoscope* 2017;127:273–9.
- 17 Lindholm C-E, Oilman B, Snyder JV, et al. Cardiorespiratory Effects of Flexible Fiberoptic Bronchoscopy in Critically Ill Patients. *Chest* 1978;74:362–8.
- 18 Matsushima Y, Jones RL, King EG, et al. Alterations in Pulmonary Mechanics and Gas Exchange during Routine Fiberoptic Bronchoscopy. *Chest* 1984;86:184–8.
- 19 Miller EJ. Hypoxemia during Fiberoptic Bronchoscopy. *Chest* 1979;75:103.
- 20 Facciolongo N, Patelli M, Gasparini S, et al. Incidence of complications in bronchoscopy. Multicentre prospective study of 20,986 bronchoscopies. *Monaldi Arch Chest Dis* 2009;71:8–14.
- 21 Stahl DL, Richard KM, Papadimos TJ. Complications of bronchoscopy: A concise synopsis. *Int J Crit Illn Inj Sci* 2015;5:189.
- 22 Lundgren R, Häggmark S, Reiz S. Hemodynamic Effects of Flexible Fiberoptic Bronchoscopy Performed under Topical Anesthesia. *Chest* 1982;82:295–9.
- 23 Cheyne WS, Gelinas JC, Eves ND. The haemodynamic response to incremental increases in negative intrathoracic pressure in healthy humans. *Exp Physiol* 2018;103:581–9.
- 24 Davies L, Mister R, Spence D, et al. Cardiovascular consequences of fibreoptic bronchoscopy. *Eur Respir J* 1997;10:695–8.
- 25 Pelaia C, Bruni A, Garofalo E, et al. Oxygenation strategies during flexible bronchoscopy: a review of the literature. *Respir Res* 2021;22:253.
- 26 Elm E von, Altman DG, Egger M, et al. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335:806–8.
- 27 Maffucci R, Maccari U, Guidelli L, et al. Pulmonologist-Administered Balanced Propofol Analgosedation during Interventional Procedures: An Italian Real-Life Study on Comfort and Safety. *Int J Clin Pract* 2022;2022:1–8.
- 28 Garofalo E, Bruni A, Pelaia C, et al. Evaluation of a New Interface Combining High-Flow Nasal Cannula and CPAP. *Respir Care* 2019;64:1231–9.
- 29 Longhini F, Liu L, Pan C, et al. Neurally-Adjusted Ventilatory Assist for Noninvasive Ventilation via a Helmet in Subjects With COPD Exacerbation: A Physiologic Study. *Respir Care* 2019;64:582–9.
- 30 Longhini F, Pan C, Xie J, et al. New setting of neurally adjusted ventilatory assist for noninvasive ventilation by facial mask: a physiologic study. *Crit Care* 2017;21:170.
- 31 Olivieri C, Longhini F, Cena T, et al. New versus Conventional Helmet for Delivering Noninvasive Ventilation: A Physiologic, Crossover Randomized Study in Critically Ill Patients. *Anesthesiology* 2016;124:101–8.
- 32 Schmidt CO, Struckmann S, Enzenbach C, et al. Facilitating harmonized data quality assessments. A data quality framework for observational health research data collections with software implementations in R. *BMC Med Res Methodol* 2021;21:63.
- 33 Gilmartin-Thomas JF, Liew D, Hopper I. Observational studies and their utility for practice. *Aust Prescr* 2018;41:82–5.
- 34 Totton N, Julious SA, Coates E, et al. Appropriate design and reporting of superiority, equivalence and non-inferiority clinical trials incorporating a benefit-risk assessment: the BRAINS study including expert workshop. *Health Technol Assess* 2023;27:1–58.
- 35 Danel A, Tobiczek E, Warcholiński A, et al. May noninvasive mechanical ventilation and/ or continuous positive airway pressure increase the bronchoalveolar lavage salvage in patients with pulmonary diseases? Randomized clinical trial - Study protocol. *Adv Med Sci* 2023;68:482–90.
- 36 Oraczewska A, Cofta S, Warcholiński A, et al. The use of non-invasive respiratory assistance to facilitate bronchofiberoscopy performance in patients with hypoxemic (type one) respiratory failure - Study protocol. *Adv Med Sci* 2023;68:474–81.
- 37 Gale RP, Zhang MJ, Lazarus HM. The role of randomized controlled trials, registries, observational databases in evaluating new interventions. *Best Pract Res Clin Haematol* 2023;36:101523.
- 38 Faraoni D, Schaefer ST. Randomized controlled trials vs. observational studies: why not just live together? *BMC Anesthesiol* 2016;16:102.
- 39 Ross JS. Randomized Clinical Trials and Observational Studies Are More Often Alike Than Unlike. *JAMA Intern Med* 2014;174:1557.
- 40 Wunsch H, Linde-Zwirble WT, Angus DC. Methods to adjust for bias and confounding in critical care health services research involving observational data. *J Crit Care* 2006;21:1–7.