

Determination of vitamin B₁₂ in dairy products by ultra performance liquid chromatography-tandem mass spectrometry

Elisa Zironi, Teresa Gazzotti, Andrea Barbarossa, Federica Farabegoli, Andrea Serraino, Giampiero Pagliuca
 Laboratory of Analytical Bio-Agroalimentary Chemistry (CABA-Lab), Department of Veterinary Sciences, Alma Mater Studiorum - University of Bologna, Ozzano dell'Emilia (BO), Italy

Abstract

Vitamin B₁₂ is a water-soluble molecule composed of a tetrapyrrolic complex with a cobalt atom at its centre. It is an essential regulatory element, synthesized only by bacteria; for this reason it is present only in food of animal origin and the daily requirement for humans is about 1 to 2 µg. Since milk and dairy products provide a significant dietary cobalamin intake, an ultra performance liquid chromatography-tandem mass spectrometry method was applied to samples collected at different stages along the process of cheese making in order to evaluate the distribution of this molecule. In particular, samples of milk, rennet, whey, ricotta cheese, curd, mozzarella cheese and caciotta cheese were analysed. Results showed a level of vitamin B₁₂ about 10 times higher in whey and ricotta cheese with respect to the milk they are derived from. These data would confirm the tendency of cobalamin to concentrate in the proteic fractions along the cheese production process.

Introduction

Vitamin B₁₂, or cobalamine, is a water-soluble molecule composed of a tetrapyrrolic complex with a cobalt atom at its centre. It is an essential regulatory element, very important to human physiology since it is involved in the red blood cells formation and nervous system functioning, and its deficiency may lead to anaemia and nerve degenerations. Differently from other vitamins belonging to the B class, vitamin B₁₂ is synthesized only by bacteria; for this reason it is present only in food bacterially fermented or obtained from animals that got this cobalamin from their gastrointestinal microflora or diet (Girard *et al.*, 2009).

Food of animal origin is the main source of

this vitamin for humans and daily requirement is about 1 to 2 µg per day (Chassigne and Lobinski, 1998). In foodstuff, vitamin B₁₂ is present in its coenzyme forms: hydroxycobalamin, 5'-deoxyadenosylcobalamin, methylcobalamin, and cyanocobalamin, which occurs only in small amounts naturally (Kelly *et al.*, 2005). Cyanocobalamin is more stable than the other forms which are light sensitive (Kumar *et al.*, 2010). Cobalamins are naturally protected from degradation by protein-binding; in bovine milk, naturally occurring vitamin B₁₂ is associated with the protein fraction, with high affinity to whey protein (Campos-Giménez *et al.*, 2008).

Since milk and dairy products provide a significant dietary cobalamin intake, an ultra performance liquid chromatography-tandem mass spectrometry method (Zironi *et al.*, 2013) was applied to samples collected at several stages along the process of cheese making in order to evaluate the distribution of this molecule in different dairy products.

Materials and Methods

Sample preparation

Samples of milk, rennet, whey, ricotta cheese, curd, mozzarella cheese and caciotta cheese were collected twice along the production processes, which were repeated four times on different days. Routine control on serum reported a pH of 4.9 and a protein content of 1.14%.

Mozzarella was produced using 50 L of unpasteurized milk and natural whey culture as starter. Briefly, the raw milk was heated to 37-38°C, then the whey and rennet were added and left to ripen at 37-38°C for about 4 h. Afterwards, the curd was extracted from the whey, stretched with hot water (90°C) for about 3 min and then molded into the traditional round shape. Each mozzarella weighed about 250 g (Serraino *et al.*, 2013).

The extraction procedure was conducted on 5 g of the various matrices adding sodium acetate buffer and potassium cyanide; samples were submitted to heat treatment that, in presence of cyanide, converts all the less stable cobalamins in cyanocobalamin (Zironi *et al.*, 2013). For solid matrices it has been necessary to introduce a centrifugation step before solid phase extraction (SPE) clean-up, in order to obtain a better separation between the liquid phase and the solid residue. Samples were then purified by a single SPE step and analysed using reverse phase liquid chromatography coupled with tandem mass spectrometry in positive electrospray ionization (ESI+). Methotrexate was used as internal standard for quantification (Lu *et al.*, 2008).

Correspondence: Giampiero Pagliuca, Laboratory of Analytical Bio-Agroalimentary Chemistry (CABA-Lab), Department of Veterinary Sciences, Alma Mater Studiorum - University of Bologna, via Tolara di Sopra 50, 40064 Ozzano dell'Emilia (BO), Italy.
 Tel. +39.051.2097008 - Fax: +39.051.2097037.
 E-mail: giampiero.pagliuca@unibo.it

Key-words: Vitamin B₁₂, Cobalamine, UPLC-MS/MS, Dairy products.

Received for publication: 8 July 2014.
 Revision received: 5 November 2014.
 Accepted for publication: 8 November 2014.

This work is licensed under a Creative Commons Attribution 3.0 License (by-nc 3.0).

©Copyright E. Zironi *et al.*, 2014
 Licensee PAGEPress, Italy
 Italian Journal of Food Safety 2014; 3:4513
 doi:10.4081/ijfs.2014.4513

Analytical conditions

The separation was performed by an acuity ultra-performance liquid chromatographic system consisting of a binary pump, solvent degasser, autosampler and column heater fitted with a Waters HSS T3 column (1.7 µm, 2.1×50 mm) equipped with a guard column with the same packing (Waters Corporation, Milford, MA, USA). Flow rate was 0.3 mL/min and the column temperature was kept at 45°C. Separation was carried out in programmed conditions with mobile phase consisting of water (A) and acetonitrile containing 0.1% formic acid (B), for a total run time of 5 min. The gradient was T₀ min: 90% A, 10% B; T₂ min: 20% A, 80% B; T₃ min to the end: 90% A, 10% B.

The mass spectrometer was a Quattro Premiere XE, a triple quadrupole instrument equipped with an ESCI™ Multi-Mode Ionization Source (Waters Corporation).

The whole analysis was conducted in ESI+ mode using multiple reaction monitoring (MRM). The monitored transitions were *m/z* 678.36→147.10 and 678.36→359.30 for vitamin B₁₂, and *m/z* 455.22→175.13 and 455.22→308.22 for methotrexate (IS).

Results and Discussion

The complexity of vitamin B₁₂ quantification in food matrices has been investigated by several Authors but very few works on milk, and in particular on dairy products, made use of mass spectrometry. Vitamin B₁₂ concentrations found in milk were similar to those described in the literature (Arkbåge *et al.*, 2003), although it was not possible to find data com-




Figure 1. Vitamin B₁₂ concentrations in different samples (n=8, mean value±standard deviation).

parable to our cheese making process.

Results showed a level of vitamin B₁₂ about 10 times higher in whey and ricotta cheese with respect to the milk they are derived from; at the same time, it is interesting to notice a decrease of cobalamin concentration in curd, caciotta cheese and mozzarella cheese (Figure 1). Vitamin B₁₂ affinity for proteins is well known and several kinds of carrier proteins, called transcobalamines, have been identified; these molecules bind cobalamin to protect it from degradation and some analogues are present in bovine milk as well (Fedosov *et al.*, 1995, 1996; Fox and Kelly, 2003; Le *et al.*, 2011).

Conclusions

These data would confirm the tendency of cobalamin to concentrate in the proteic fraction of whey along the cheese production process. The lower value for mozzarella cheese compared to curd is probably due to the process of production, which implies curd

draining and a dilution effect related to the water addition in the stretching phase.

References

- Arkbåge K, Withthöft C, Fondén R, Jägerstad M, 2003, Retention of vitamin B₁₂ during manufacture of six fermented dairy products using a validated radio protein-binding assay. *Int Dairy J* 13:101-9.
- Campos-Giménez, E, Fontannaz P, Trisconi MJ, Kiliç T, Gimenez C, Andrieux P, 2008. Determination of vitamin B₁₂ in food products by liquid chromatography/UV detection with immunoaffinity extraction: single-laboratory validation. *J AOAC Int* 91:786-93.
- Chassigne H, Lobinski R, 1998. Direct species-selective determination of cobalamins by ionspray mass spectrometry and ionspray tandem mass spectrometry. *Analyst* 123:131-7.
- Fedosov SN, Petersen TE, Nexø E, 1995.

Binding of cobalamin and cobinamide to transcobalamin from bovine milk. *Biochemistry* 34:16082-7.

- Fedosov SN, Petersen TE, Nexø E, 1996. Transcobalamin from milk: isolation and physico-chemical properties. *Biochim Biophys Acta* 1292:113-9.
- Fox PF, Kelly AL, 2003. Developments in the chemistry and technology of milk proteins. *Food Aust* 55:231-4.
- Girard CL, Santschi DE, Stabler SP, Allen RH, 2009. Apparent ruminal synthesis and intestinal disappearance of vitamin B₁₂ and its analogs in dairy cows. *J Dairy Sci* 92:4524-9.
- Kelly RJ, Gruner TM, Sykes AR, 2005. Development of a method for the separation of corrinoids in ovine tissues by HPLC. *Biomed Chromatogr* 19:329-33.
- Kumar SS, Chouhan RS, Thakur MS, 2010. Trends in analysis of vitamin B₁₂. *Anal Biochem* 398:139-49.
- Le A, Barton LD, Sanders JT, Zhang Q, 2011. Exploration of bovine milk proteome in colostrum and mature whey using an ion-exchange approach. *J Proteome Res* 10:692-704.
- Lu B, Ren Y, Huang B, Liao W, Cai Z, Tie X, 2008. Simultaneous determination of four water-soluble vitamins in fortified infant foods by ultra-performance liquid chromatography coupled with triple quadrupole mass spectrometry. *J Chromatogr Sci* 46:225-32.
- Serraino A, Giacometti F, Daminelli P, Losio MN, Finazzi G, Marchetti G, Zambrini AV, Rosmini R, 2013. Survival of *Arcobacter butzleri* during production and storage of artisan water buffalo mozzarella cheese. *Foodborne Pathog Dis* 10:820-4.
- Zironi E, Gazzotti T, Barbarossa A, Devicienti C, Scardilli M, Pagliuca G, 2013. Technical note: development and validation of a method using ultra performance liquid chromatography coupled with tandem mass spectrometry for determination of vitamin B-12 concentrations in milk and dairy products. *J Dairy Res* 96:2832-6.