



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

The makers of the Protoaurignacian and implications for Neandertal extinction

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

The makers of the Protoaurignacian and implications for Neandertal extinction / Benazzi, S.; Slon, V.; Talamo, S.; Negrino, F.; Peresani, M.; Bailey, S.E.; Sawyer, S.; Panetta, D.; Vicino, G.; Starnini, E.; Mannino, M.A.; Salvadori, P.A.; Meyer, M.; Paabo, S.; Hublin, J.-J.. - In: SCIENCE. - ISSN 0036-8075. - ELETTRONICO. - 348:6236(2015), pp. 793-796. [10.1126/science.aaa2773]

This version is available at: <https://hdl.handle.net/11585/516010> since: 2021-11-19

Published:

DOI: <http://doi.org/10.1126/science.aaa2773>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

This is the final peer-reviewed accepted manuscript of:

S. Benazzi, V. Slon, S. Talamo, F. Negrino, M. Peresani, S. E. Bailey, S. Sawyer, D. Panetta, G. Vicino, E. Starnini, M. A. Mannino, P. A. Salvadori, M. Meyer, S. Pääbo, J.-J. Hublin, The makers of the Protoaurignacian and implications for Neandertal extinction. *Science*. **348**, 793–796 (2015)

The final published version is available online at:

<https://www.science.org/doi/10.1126/science.aaa2773>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>) When citing, please refer to the published version.

Title: The Makers of the Protoaurignacian and Implications for Neandertal Extinction

Authors: S. Benazzi^{1,2*}, V. Slon³, S. Talamo², F. Negrino⁴, M. Peresani⁵, S.E. Bailey^{2,6}, S. Sawyer³, D. Panetta⁷, G. Vicino⁸, E. Starnini^{9,10}, M.A. Mannino², P.A. Salvadori⁷, M. Meyer³, S. Pääbo³, J-J. Hublin²

Affiliations:

¹Department of Cultural Heritage, University of Bologna, Via degli Ariani 1, 48121 Ravenna, Italy.

²Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany.

³Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany.

⁴Dipartimento di Antichità, Filosofia, Storia e Geografia, Università di Genova, Via Balbi 2, 16126 Genova, Italy.

⁵Sezione di Scienze Preistoriche e Antropologiche, Dipartimento di Studi Umanistici, Corso Ercole I d’Este 32, Università di Ferrara, 44100 Ferrara, Italy.

⁶Center for the Study of Human Origins, Department of Anthropology, New York University, 25 Waverly Place, New York, NY 10003, USA.

⁷CNR Institute of Clinical Physiology, National Research Council, Via G. Moruzzi 1, 56124 Pisa, Italy.

⁸Museo Archeologico del Finale, Chiostri di Santa Caterina, 17024 Finale Ligure Borgo, Italy.

⁹Scuola di Scienze Umanistiche, Dipartimento di Studi Storici, Università di Torino, via S. Ottavio 20, 10124 Torino, Italy.

¹⁰Museo Preistorico Nazionale dei Balzi Rossi, Via Balzi Rossi 9, 18039 Ventimiglia, Italy.

*Correspondence to: stefano.benazzi@unibo.it

Abstract: The Protoaurignacian culture is pivotal to the debate about the timing of the arrival of modern humans in Western Europe and the demise of Neandertals. However, which group is responsible for this culture remains uncertain. Here we investigate dental remains associated with the Protoaurignacian. The lower deciduous incisor from Riparo Bombrini is modern human based on its morphology. The upper deciduous incisor from Grotta di Fumane contains ancient mitochondrial DNA of a modern human type. These teeth are the oldest human remains in an Aurignacian-related archeological context, confirming that by 41,000 cal BP, modern humans bearing Protoaurignacian culture spread into Southern Europe. Since the last Neandertals date to 41,030-39,260 cal BP, we suggest that the Protoaurignacian triggered the demise of Neandertals in this area.

One Sentence Summary: The Protoaurignacian culture is a modern human industry that overlaps in time with the last Neandertals.

Main Text: The timing and pattern of the biological and cultural shifts that occurred in Western Europe around 45,000-35,000 calendar years ago (cal BP) fuel continuing debates among

paleoanthropologists and prehistorians (1-3). During this period, Neandertals were replaced by anatomically modern humans (AMH) (4), and a variety of “transitional” and early Upper Paleolithic cultures emerged. Among them, the Protoaurignacian is crucial to current interpretations regarding the timing of arrival of AMH and their interaction with Neandertals (5-9).

The Protoaurignacian appeared around 42,000 cal BP (8, 10) in Southwest and South-Central Europe (Fig. S1). In addition to the presence of personal ornaments, such as perforated shells and worked bones, the Protoaurignacian is characterized by a dominance of bladelets with typical retouched standardized implements like Font-Yves points and Dufour bladelets produced from unipolar cores (5). This techno-complex has been tentatively linked to the Ahmorian industry of the Levant (6, 9). Since the Ahmorian has been attributed to modern humans (11), it has been suggested that the Protoaurignacian reflects a westward population movement of AMH from the Near East (1, 7). However, because only three non-diagnostic human remains are associated with this culture, it is still uncertain who the makers of the Protoaurignacian were (9, 12). The fossil remains associated with the Protoaurignacian available for study consist of the undiagnostic skeletal fragments of a fetus retrieved from Le Piage rock-shelter (France) (13), for which the stratigraphic integrity of the Châtelperronian/Aurignacian sequence has been questioned (5), and of two deciduous incisors from two Northern Italian sites: Riparo Bombrini (Western Ligurian Alps, Italy) and Grotta di Fumane (Western Lessini Mountains, Italy). The lower left lateral deciduous incisor (Ldi₂; Fig. 1) found in 1976 in Riparo Bombrini (14, 15) (Figs. S2-S5) and the upper right lateral deciduous incisor (Rdi², Fig. 1) labelled Fumane 2, which was retrieved in

1992 from the Protoaurignacian deposit of Grotta di Fumane (15, 16) (Figs. S6, S7), have to date not been conclusively attributed to modern humans or Neandertals.

The crown diameters of deciduous incisors are undiagnostic for Neandertals and modern humans, as is the case also for other tooth classes (2). However, on the basis of the buccolingual crown diameter, the Bombrini specimen is close to the mean of Upper Paleolithic modern humans, whereas Fumane 2 is closer to the Neandertal mean (Table S1). Other than that, the worn deciduous lower incisors do not provide any morphologically diagnostic information. To establish the identity of the makers of the Protoaurignacian, we analyzed the 3D enamel thickness components of the Bombrini Ldi₂ using a digital approach (17), and we were able to investigate DNA from the Fumane 2 specimen (15).

The relative enamel thickness (RET) index has been recognized as an effective taxonomic discriminator between Neandertals and modern humans. Neandertal deciduous and permanent teeth are characterized by significantly thinner enamel relative to dentine volume (18). To facilitate comparisons with the Bombrini specimen, which is affected by wear stage 4, the Neandertal and recent modern human (RMH) di₂ samples were divided into sub-groups based on their degree of wear (from wear stage 1/2 to wear stage 4) (15, 19) (Table 1, Table S2). The Neandertal di₂ RET indices are lower than those of RMH at similar wear stages, and no overlap in the range of variation is observed between the two groups. The RET index of Bombrini is higher than any values obtained for Neandertals (Table S2), despite the missing portion of the enamel cap, and its computed Z-score is close to the modern human mean in wear stage 4 (Table 1).

To test how much the loss of enamel on the mesial side of the Bombrini tooth affects the computed RET value, two RMH specimens were digitally worn and damaged to simulate the condition observed in Bombrini (15) (Fig. S8). The results confirm that tooth wear, at least up to wear stage 4, decreases the RET index by about 10%, while the mesial loss of enamel affects the index by less than 1.6% (much less than the values considered acceptable for intra- and inter-observer error) (2). Therefore, the RET value for the unworn Bombrini Ldi₂ was certainly much higher (Table S3), further supporting its attribution to modern humans.

DNA was extracted from the Fumane 2 tooth, which yielded few mitochondrial DNA (mtDNA) sequences (Table S4). With respect to 63 ‘diagnostic’ positions at which ten Neandertal mitochondrial genomes differ from 311 present-day humans (20), these sequences were of modern human origin (Table S5). To further explore this, we prepared a second DNA extract and two DNA libraries from this specimen, which yielded a total of 335,628 unique mtDNA fragments (Table S4).

The frequencies of cytosine (C) to thymine (T) substitutions at the ends of these fragments (34–37%, Fig. S9), which reflect the deamination of cytosine residues typical of ancient DNA (21, 22), are consistent with results from other specimens of similar age (23–26). Among the fragments carrying terminal C to T substitutions, we estimated the residual present-day DNA contamination to be 3.8% (15).

Using these fragments, we reconstructed a mitochondrial genome of 157-fold coverage (Fig. S10). This mtDNA sequence was aligned to the mtDNAs of 54 present-day humans, ten ancient modern humans, ten Neandertals, two Denisovans, a hominin from Sima de los Huesos (Spain) and a chimpanzee (*Pan troglodytes*). The Fumane 2 mitochondrial genome falls within the variation of modern humans (Fig. 2) and basally in haplogroup R (Table S6), as also observed for the ~45,000 year old AMH specimen from Ust'-Ishim (in western Siberia (26)), a major group of related mtDNAs in Eurasia (27) into which most pre-agricultural mtDNAs in Europe fall (28).

As expected for an ancient specimen (23, 25), the Fumane 2 mtDNA has accumulated fewer nucleotide substitutions than present-day mtDNA (Fig. 2). Using 10 directly dated ancient modern humans (25, 26) as multiple calibration points, we estimated the age of the Fumane 2 terminal node to be 44,599 (95% highest posterior density: 19,755-72,070) years BP. We thus conclude that the Fumane 2 individual carried a mitochondrial genome of a modern human type. This shows that this individual was a modern human, or had at least some ancestors who were modern humans.

Based on recent chronometric data for the Protoaurignacian deposit of Grotta di Fumane, the specimen Fumane 2 is dated to 41,110-38,500 cal BP (10), as recalibrated with IntCal13 (29). Among modern humans in Western Europe, it is currently predated only by the contested Kent's Cavern maxilla (30, 31) and by the 45,000-43,000 cal BP AMH specimens from the Uluzzian levels of Grotta del Cavallo (2), for which a recent taxonomic reassessment is stimulating intense debate (32, 33).

New radiocarbon dates of the Protoaurignacian layers of Riparo Bombrini were obtained from faunal bones recovered during the G. Vicino excavation (five samples) and the more recent 2002-2005 excavations (three charcoals and eight animal bones) (15) (Tables S7-S10). The new dates confirm the integrity of the cemented deposit explored in 1976 by Vicino (which yielded the Ldi₂ tooth), but suggest that some stratigraphic disturbance affected a restricted area explored during the 2002-2005 excavations (supplementary text; Table S7, Fig. S11). The ¹⁴C dates of the Vicino 1976 excavation (Table S7) were incorporated into a Bayesian model for the distribution of ages (34) (Fig. 3). The Protoaurignacian levels (level III to level II) are dated between 40,710 and 35,640 cal BP (68.2% probability), corresponding to a cold phase that marks the onset of Heinrich Stadial 4 (35).

The Bombrini Ldi₂ and potentially the Fumane 2 Rdi² thus represent the oldest AMH remains in an Aurignacian-related (i.e. Protoaurignacian or Early Aurignacian) archaeological context, confirming that by around 41,000 cal BP (68.2% probability), AMH populations bearing Protoaurignacian culture had spread into Europe along the Mediterranean coast. They are similar in age to, or slightly older than, the modern human remains from: Peștera cu Oase (Romania, ca. 40,000 cal BP), which lacks archaeological context; Kostenki 14, Layer III (Russia, ca. 38,000 cal BP), which is possibly Aurignacian; Kostenki 1, Layer III (ca. 38,000 cal BP) which is associated with diagnostic Aurignacian artifacts; Kostenki 14, Layer IVb and Kostenki 17, Layer II that underlie the Campanian Ignimbrite tephra and are of comparable age to the Protoaurignacian in Italy, are tentatively assigned to AMH and also associated with an

assemblage that includes bladelets; La Quina-Aval and Brassemouy (France), which are Early Aurignacian and more recent than 40,000 cal BP (for a review see (9)).

The Protoaurignacian dispersal overlaps in time with late Neandertal populations, as indicated by the 41,030-39,260 cal BP age of the last Mousterian sites (4), and the ~45,000-40,000 cal BP age of the Châtelperronian culture (3), which is currently attributed to Neandertals (36). The Protoaurignacian dispersal may therefore have been a cause (either directly or indirectly) for the extinction of the Neandertals, at least in Northern Italy.

References and Notes :

1. J. F. Hoffecker, Out of Africa: modern human origins special feature: the spread of modern humans in Europe. *Proc. Natl. Acad. Sci. USA* **106**, 16040–16045 (2009).
2. S. Benazzi *et al.*, Early dispersal of modern humans in Europe and implications for Neanderthal behaviour. *Nature* **479**, 525–528 (2011).
3. J.-J. Hublin *et al.*, Radiocarbon dates from the Grotte du Renne and Saint-Césaire support a Neandertal origin for the Châtelperronian. *Proc. Natl. Acad. Sci. USA* **109**, 18743–18748 (2012).
4. T. Higham *et al.*, The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* **512**, 306-309 (2014).
5. J.-G. Bordes, in *The Chronology of the Aurignacian and of the Transitional Technocomplexes: Dating, Stratigraphies, Cultural Implications*, J. Zilhão, F. D'Errico, Eds. (Instituto Português de Arqueologia, 2003), pp. 223-246.

6. O. Bar-Yosef, in *More than Meets the Eye: Studies on Upper Palaeolithic Diversity in the Near East*, A. N. Goring-Morris, A. Belfer-Cohen, Eds. (Oxbow Books, 2003), pp. 265–273.
7. P. Mellars, A new radiocarbon revolution and the dispersal of modern humans in Eurasia. *Nature* **439**, 931–935 (2006).
8. K. Douka, S. Grimaldi, G. Boschian, A. del Lucchese, T. Higham, A new chronostratigraphic framework for the Upper Palaeolithic of Riparo Mochi (Italy). *J. Hum. Evol.* **62**, 286–99 (2012).
9. J-J. Hublin, The Modern Human Colonization of Western Eurasia: When and Where? *Quarter. Sci. Rev.* (2014) <http://dx.doi.org/10.1016/j.quascirev.2014.08.011>.
10. T. Higham *et al.*, Problems with radiocarbon dating the Middle and Upper Palaeolithic transition in Italy. *Quarter. Sci. Rev.* **28**, 1257–1267 (2009).
11. C. A. Bergman, C. B. Stringer, Fifty years after: Egbert, an early upper Palaeolithic juvenile from Ksar Akil, Lebanon. *Paléorient* **15**, 99–111 (1989).
12. N. J. Conard, P. M. Grootes, F. H. Smith, Unexpectedly recent dates for human remains from Vogelherd. *Nature* **430**, 198–201 (2004).
13. S. Beckouche, F. Poplin, in *Le Piage, site préhistorique du Lot*, F. Champagne, R. Espitalié, Eds. (Société Préhistorique Française, 1981), pp. 159–160.
14. V. Formicola, Early aurignacian deciduous incisor from Riparo Bombrini at Balzi Rossi (Grimaldi, Italy). *Riv. Antropol.* **67**, 287–292 (1989).
15. Information on materials and methods and localities is available at the *Science* Web site.
16. G. Bartolomei *et al.*, La Grotte-Abri de Fumane. Un Site Aurignacien au Sud des Alps. *Preistoria Alpina* **28**, 131–179 (1992).

17. S. Benazzi *et al.*, Guidelines for the digital computation of 2D and 3D enamel thickness. *Am. J. Phys. Anthropol.* **153**, 305–313 (2014).
18. R. Macchiarelli *et al.*, How Neanderthal molar teeth grew. *Nature* **444**, 748–751 (2006).
19. S. Molnar, Human tooth wear, tooth function and cultural variability. *Am. J. Phys. Anthropol.* **34**, 175–190 (1971).
20. R. E. Green *et al.*, A Complete Neandertal Mitochondrial Genome Sequence Determined by High-Throughput Sequencing. *Cell* **134**, 416–426 (2008).
21. A. W. Briggs *et al.*, Patterns of damage in genomic DNA sequences from a Neandertal. *Proc. Natl. Acad. Sci. USA* **104**, 14616–14621 (2007).
22. S. Sawyer, J. Krause, K. Guschanski, V. Savolainen, S. Pääbo, Temporal patterns of nucleotide misincorporations and DNA fragmentation in ancient DNA. *PLoS ONE* **7**, e34131 (2012).
23. J. Krause *et al.*, A complete mtDNA genome of an early modern human from Kostenki, Russia. *Curr. Biol.* **20**, 231-236 (2010).
24. J. Krause *et al.*, The complete mitochondrial DNA genome of an unknown hominin from southern Siberia. *Nature* **464**, 894-897 (2010).
25. Q. Fu *et al.*, A revised timescale for human evolution based on ancient mitochondrial genomes. *Curr. Biol.* **23**, 553-559 (2013).
26. Q. Fu *et al.*, Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* **514**, 445-449 (2014).
27. P. Soares *et al.*, Correcting for purifying selection: An improved human mitochondrial molecular clock. *Am. J. Hum. Genet.* **84**, 740-759 (2009).

28. I. Lazaridis *et al.*, Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* **513**, 409-413 (2014).
29. P. J. Reimer *et al.*, IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* **55**, 1869-1887 (2013).
30. T. Higham *et al.*, The earliest evidence for anatomically modern humans in northwestern Europe. *Nature* **479**, 521–524 (2011).
31. M. White, P. Pettitt, Ancient Digs and Modern Myths: The Age and Context of the Kent’s Cavern 4 Maxilla and the Earliest *Homo sapiens* Specimens in Europe. *Eur. J. Arch.* **15**, 392-420 (2012).
32. W. E. Banks, F. d'Errico, J. Zilhão, Human-climate interaction during the Early Upper Paleolithic: testing the hypothesis of an adaptive shift between the Proto-Aurignacian and the Early Aurignacian. *J. Hum. Evol.* **64**, 39–55 (2013).
33. A. Ronchitelli, S. Benazzi, P. Boscato, K. Douka, A. Moroni, Comments on “Human-climate interaction during the Early Upper Paleolithic: testing the hypothesis of an adaptive shift between the Proto-Aurignacian and the Early Aurignacian” by William E. Banks, Francesco d’Errico, João Zilhão. *J. Hum. Evol.* **73**, 107-111 (2014).
34. C. Bronk Ramsey, S. Lee, Recent and Planned Developments of the Program OxCal. *Radiocarbon* **55**, 720-730 (2013).
35. D. Fleitmann *et al.*, Timing and climatic impact of Greenland interstadials recorded in stalagmites from northern Turkey. *Geophys. Res. Lett.* **36** (2009).
36. S. E. Bailey, J-J. Hublin, Dental remains from the Grotte du Renne at Arcy-sur-Cure (Yonne). *J. Hum. Evol.* **50**, 485–508 (2006).

37. D. Panetta, N. Belcari, A. Del Guerra, A. Bartolomei, P. Salvadori, Analysis of image sharpness reproducibility on a novel engineered micro-CT scanner with variable geometry and embedded recalibration software. *Phys. Med.* **28**, 166–173 (2012).
38. L. A. Feldkamp, L. C. Davis, J. W. Kress, Practical cone-beam algorithm. *J. Opt. Soc. Am. A* **6**, 612–619 (1984).
39. P. S. Ungar, K. J. Fennell, K. Gordon, E. Trinkaus, Neandertal incisor beveling. *J. Hum. Evol.* **32**, 407–421 (1997).
40. P. J. Besl, N. D. McKay, A method for registration of 3D shapes. *PAMI* **14**, 239–256 (1992).
41. J. Dabney *et al.*, Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. USA* **110**, 15758–15763 (2013).
42. M.-T. Gansauge, M. Meyer, Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nature Protocols* **8**, 737–748 (2013).
43. A. W. Briggs *et al.*, Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res.* **38**, e87 (2010).
44. M. Meyer *et al.*, A high-coverage genome sequence from an archaic Denisovan individual. *Science* **338**, 222–226 (2012).
45. M. Meyer, M. Kircher, Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb Protoc.* **2010** (6): pdb.prot5448 (2010).
46. M. Kircher, S. Sawyer, M. Meyer, Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res.* **40**, e3 (2012).

47. J. Dabney, M. Meyer, Length and GC-biases during sequencing library amplification: A comparison of various polymerase-buffer systems with ancient and modern DNA sequencing libraries. *BioTechniques* **52**, 87–94 (2012).
48. T. Maricic, M. Whitten, S. Pääbo, Mutiplexed DNA Sequence Capture for Mitochondrial Genomes Using PCR Products. *PLoS ONE* **5**, e14004 (2010).
49. Q. Fu *et al.*, DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl. Acad. Sci. USA* **110**, 2223–2227 (2013).
50. M. Kircher, U. Stenzel, J. Kelso, Improved base calling for the Illumina Genome Analyzer using machine learning strategies. *Genome Biol.* **10**, R83 (2009).
51. G. Renaud, U. Stenzel, J. Kelso, leeHom: adaptor trimming and merging for Illumina sequencing reads. *Nucleic Acids Res.* **42**, e141 (2014).
52. H. Li, R. Durbin, Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).
53. M. Ingman, H. Kaessmann, S. Pääbo, U. Gyllensten, Mitochondrial genome variation and the origin of modern humans. *Nature*. **408**, 708-713 (2000).
54. A. W. Briggs *et al.*, Targeted retrieval and analysis of five Neandertal mtDNA genomes. *Science* **325**, 318-321 (2009).
55. M. Meyer *et al.*, A mitochondrial genome sequence of a hominin from Sima de los Huesos. *Nature* **505**, 403–406 (2014).
56. K. Prüfer *et al.*, The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* **505**, 43–49 (2014).
57. M.-T. Gansauge, M. Meyer, Selective enrichment of damaged DNA molecules for ancient genome sequencing. *Genome Res.* **24**, 1543-1549 (2014).

58. P. Skoglund *et al.*, Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 2229-2234 (2014).
59. D. Reich *et al.*, Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature* **468**, 1053–1060 (2010).
60. S. Horai, K. Hayasaka, R. Kondo, K. Tsugane, N. Takahata, Recent African origin of modern humans revealed by complete sequences of hominoid mitochondrial DNAs. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 532-536 (1995).
61. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-80 (2013).
62. K. Tamura, G. Stecher, D. Peterson, A. Filipski, S. Kumar, MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* **30**, 2725-2729 (2013).
63. A. Kloss-Brandstaetter *et al.*, HaploGrep: a fast and reliable algorithm for automatic classification of mitochondrial DNA haplogroups. *Hum. Mutat.* **32**, 25-32 (2010).
64. M. van Oven, M. Kayser, Updated comprehensive phylogenetic tree of global human mitochondrial DNA variation. *Hum. Mutat.* **30**, E386-E394 (2009).
65. A. J. Drummond, A. Rambaut, BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* **7**, 214 (2007).
66. D. Posada, K. A. Crandall, MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**, 817-818 (1998).
67. S. Talamo, M. Richards, A comparison of bone pretreatment methods for AMS dating of samples >30, 000 BP. *Radiocarbon* **53**, 443-449 (2011).
68. R. Longin, New method of collagen extraction for radiocarbon dating. *Nature* **230**, 241-242 (1971).

69. T. A. Brown, D. E. Nelson, J. S. Vogel, J. R. Southon, Improved Collagen Extraction by modified Longin method. *Radiocarbon* **30**, 171-177 (1988).
70. F. Brock, C. B. Ramsey, T. Higham, Quality assurance of ultrafiltered bone dating. *Radiocarbon* **49**, 187–192 (2007).
71. S. H. Ambrose, Preparation and Characterization of Bone and Tooth Collagen for Isotopic Analysis. *J. Archaeol. Sci.* **17**, 431–451 (1990).
72. G. J. V. Klinken, Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurements. *J. Archaeol. Sci.* **26**, 687-695 (1999).
73. B. Kromer, S. Lindauer, H-A. Synal, L. Wacker, MAMS – A new AMS facility at the Curt-Engelhorn-Centre for Achaeometry, Mannheim, Germany. *Nucl. Instrum. Meth. B* **294**, 11-13 (2013).
74. K. K. Andersen *et al.*, The Greenland Ice Core Chronology 2005, 15–42 ka. Part 1: constructing the time scale. *Quater. Sci. Rev.* **25**, 3246-3257 (2006).
75. A. Svensson *et al.*, The Greenland Ice Core Chronology 2005, 15–42 ka. Part 2: comparison to other records. *Quater. Sci. Rev.* **25**, 3258-3267 (2006).
76. A. Del Lucchese, F. Negrino, S. Simone, G. Vicino, in *Un siècle de construction du discours scientifique en Préhistoire* (26° Congrès Préhistorique de France, Avignon–Bonnieux 21-25, 2004), pp. 177-184.
77. M. Mussi, J. Cinq-Mars, P. Bolduc, Echoes from the mammoth steppe: the case of the Balzi Rossi. *Anal. Praehist. Leid* **31**, 105-124 (1999).
78. G. Vicino, Incisioni rupestri paleolitiche ai Balzi Rossi. *Riv. Ing. Int.* **26**, 51-56 (1971).
79. G. Vicino, Incisioni rupestri paleolitiche ai Balzi Rossi. *Riv. St. Lig.* **38**, 5-26 (1972).

80. G. Vicino, M. Mussi, Arte parietale ai Balzi Rossi: la Grotticella Blanc-Cardini (Ventimiglia, IM). *Origini* **23**, 21-38 (2011).
81. M.A. de Lumley, L'os iliaque anténéanderthalien de la grotte du Prince (Grimaldi, Ligurie italienne). *Bull. Mus. Anthropol. Préhist. Monaco* **18**, 89-112 (1972).
82. G. Vicino, Gli scavi preistorici nell'area dell'ex Casinò dei Balzi Rossi (Nota preliminare). *Riv. Ing. Int.* **27**, 77-97 (1972).
83. G. Vicino, La spiaggia tirreniana dei Balzi Rossi nei recenti scavi della zona dell'ex-casinò. *Atti della XVI Riunione Scientifica, Istituto Italiano di Preistoria e Protostoria, Firenze*, 75-95 (1974).
84. M. Cremaschi, A. Del Lucchese, F. Negrino, C. Ottomano, B. Wilkens, Ventimiglia (Imperia). Località Balzi Rossi. Nuovi dati sulla successione stratigrafica del ciclo interglaciale-glaciale-postglaciale. Scavi 1990. *Bollettino di Archeologia* **8**, 47-50 (1991).
85. A. Bietti, F. Negrino, in *New Approaches to the Study of Early Upper Palaeolithic "Transitional" Industries in Western Eurasia. Transitions Great and Small*, J. Riel-Salvatore, G. A. Clark, Eds. (B.A.R., Oxford, 2007), pp. 41-60.
86. A. Bietti, F. Negrino, L'Aurignacien et le Gravettien du Riparo Mochi, l'Aurignacien du Riparo Bombrini: comparaisons et nouvelles perspectives. *Arch. Inst. Pal. Hum.* **39**, 133-140 (2008).
87. A. Bietti, A. Del Lucchese, F. Negrino, Nuovi studi e ricerche al Riparo Mochi (Balzi Rossi, Ventimiglia, Imperia). *Paleo-express* **7**, 4-6 (2001).
88. A. Bietti, A. Del Lucchese, F. Negrino, E. Spinapolic, in *Archeologia in Liguria, Nuova Serie, I, 2004-2005*, A. Del Lucchese, L. Gambaro, Eds. (Soprintendenza per i Beni Archeologici della Liguria, Genova, 2008), pp. 237-238.

89. A. Tagliacozzo, F. Zeppieri, I. Fiore, E. Spinapolicce, A. Del Lucchese, Archaeozoological evidence of subsistence strategies during the Gravettian at Riparo Mochi (Balzi Rossi, Ventimiglia, Imperia - Italy). *Quat. Int.* **252**, 142-154 (2012).
90. C. Tozzi, F. Negrino, Nouvelles données sur les cultures moustériennes des grottes de Grimaldi. *Arch. Inst. Pal. Hum.* **39**, 101-107 (2008).
91. V. Formicola, P. B. Pettitt, A. Del Lucchese, A Direct AMS Radiocarbon Date on the Barma Grande 6 Upper Paleolithic Skeleton. *Curr. Anthropol.* **45**, 114-118 (2004).
92. V. Formicola, B. Holt, I resti umani del Paleolitico superiore della Liguria Occidentale: Una sintesi sui risultati degli studi recenti. *Bull. Mus. Anthropol. Préhist. Monaco, suppl.* **1**, 96-102 (2008).
93. B. Holt, V. Formicola, Hunters of the Ice Age: The biology of Upper Paleolithic people. *Am. J. Phys. Anthropol.* **S47**, 70-99 (2008).
94. É. Rivière, De l'antiquité de l'homme dans les Alpes-Maritimes (Paris, 1887).
95. L. Cardini, Recenti scavi dell'Istituto Italiano di Paleontologia Umana alla Barma Grande di Grimaldi. *Arch. Antr. Etn.* **68**, 385-389 (1938).
96. G. Vicino, Lo scavo paleolitico al Riparo Bombrini (Balzi Rossi di Grimaldi, Ventimiglia). *Riv. Ing. Int.* **39**, 1-10 (1984).
97. V. Formicola, Un incisivo umano deciduo dal deposito aurignaziano del Riparo Bombrini ai Balzi Rossi. *Riv. Ing. Int.* **39**, 11-12 (1984).
98. D. Arobba, R. Caramiello, Analisi paleobotanica sui sedimenti del riparo Bombrini (Balzi Rossi, Ventimiglia). *Bull. Mus. Anthropol. Préhist. Monaco* **49**, 41-48 (2009).
99. S. Bertola *et al.*, La diffusione del primo Aurignaziano a sud dell'Arco alpino. *Preistoria Alpina* **47**, 17-30 (2013).

- 100.A. Del Lucchese, V. Formicola, B. Holt, F. Negrino, G. Vicino, Riparo Bombrini, Balzi Rossi (Ventimiglia, Imperia). Notizie preliminari degli scavi 2002-2004. *Ligures* **2**, 287-289 (2004).
- 101.F. Negrino, Riparo Bombrini, Balzi Rossi (Ventimiglia, IM): la campagna 2005. *Ligures* **3**, 194-196 (2005).
- 102.F. Negrino, C. Tozzi, Il Paleolitico in Liguria. *Bull. Mus. Anthropol. Préhist. Monaco suppl.* **1**, 21-28 (2008).
- 103.J. Riel-Salvatore, I. C. Ludeke, F. Negrino, B. M. Holt, A spatial analysis of the Late Mousterian levels of Riparo Bombrini (Balzi Rossi, Italy), *Can. J. Archaeol.* **37**, 70-92 (2013).
- 104.S. Bertola, Approccio micropaleontologico discriminante per riconoscere la provenienza alpina o appenninica delle selci della scaglia rossa (Italia centro-settentrionale). *Bull. Mus. Anthropol. Préhist. Monaco* **52**, 17-27 (2012).
- 105.F. Negrino, E. Starnini, in *Les Matières Premières Lithiques en Préhistoire* (Actes de la Table ronde internationale, Aurillac, France, 2003), pp. 235-243.
- 106.M. Martini, E. Sibilia, S. Croci, M. Cremaschi, Thermoluminescence (TL) dating of burnt flints: problems, perspectives and some example of application. *J. Cult. Herit.* **2**, 179-190 (2001).
- 107.M. Peresani *et al.*, Age of the final Middle Palaeolithic and Uluzzian levels at Fumane Cave, Northern Italy, using ^{14}C , ESR, $^{234}\text{U}/^{230}\text{Th}$ and thermoluminescence methods. *J. Arch. Sci.* **35**, 2986–2996 (2008).
- 108.P. F. Cassoli, A. Tagliacozzo, Considerazioni paleontologiche, paleoeconomiche e archeozoologiche sui macromammiferi e gli uccelli dei livelli del Pleistocene superiore del

- Riparo di Fumane (VR) (Scavi 1988-91). *Boll. Mus. Civ. Stor. Nat. Verona* **18**, 349-445 (1991).
- 109.M. Peresani, A new cultural frontier for the last Neanderthals: the Uluzzian in Northern Italy. *Curr. Anthropol.* **49**, 725–731 (2008).
- 110.C.A. Jéquier, M. Romandini, M. Peresani, Les retouchoirs en matières dures animales: une comparaison entre Moustérien final et Uluzzien. *Comptes Rendus Palevol* **11**, 283-292 (2012).
- 111.M. Peresani, Fifty thousand years of flint knapping and tool shaping across the Mousterian and Uluzzian sequence of Fumane cave. *Quat. Int.* **247**, 125-150 (2012).
- 112.M. Peresani *et al.*, Fire-places, frequentations and the environmental setting of the final Mousterian at Grotta di Fumane: a report from the 2006-2008 research. *Quartär.* **58**, 131-151 (2011).
- 113.M. Peresani, I. Fiore, M. Gala, M. Romandini, A. Tagliacozzo, Late Neandertals and the intentional removal of feathers as evidenced from bird bone taphonomy at Fumane cave 44ky BP, Italy. *Proc. Natl. Acad. Sci. U S A* **108**, 3888-3893 (2011).
- 114.M. Peresani, L. Centi, E. Di Taranto, Blades, bladelets and flakes: a case of variability in tool design at the onset of the Middle – Upper Palaeolithic transition in Italy. *Comptes Rendus Palevol* **12**, 211-221 (+corrigendum) (2013).
- 115.M. Romandini, N. Nannini, A. Tagliacozzo, M. Peresani, The ungulate assemblage from layer A9 at Grotta di Fumane, Italy: a zooarchaeological contribution to the reconstruction of Neanderthal ecology. *Quat. Int.* **337**, 11–27 (2014).

- 116.A. Broglio, L'estinzione dell'Uomo di Neandertal e la comparsa dell'Uomo moderno in Europa. Le evidenze della Grotta di Fumane nei Monti Lessini. *Atti Istituto Veneto SS.LL.AA* **45**, 1–44 (1997).
- 117.A. Broglio, F. Gurioli, in *La spiritualité. Etudes et Recherches Archéologiques 106*, M. Otte, Ed. (Université de Liège, Liège, 2004), pp. 97–102.
- 118.C. Peretto *et al.*, Living-Floors and Structures From the Lower Palaeolithic to the Bronze Age. *Coll. Anthropol.* **28**, 63–88 (2004).
- 119.A. Broglio *et al.*, in *Production lamellaires attribuées à l'Aurignacien*, F. Le Brun-Ricalens, Ed. (Luxembourg, Musée National d'Histoire et d'Art, 2005), pp. 415–436.
- 120.A. Broglio, M. De Stefani, A. Tagliacozzo, F. Gurioli, A. Facciolo, in *Kostenki and the early Upper Paleolithic of Eurasia: General trends, local developments*, S. A. Vasil'ev *et al.*, Eds. (Nestor-Historia Saint Petersburg, 2006), pp. 263–268.
- 121.A. Broglio *et al.*, L'art aurignacien dans la décoration de la Grotte de Fumane. *L'Anthropologie* **113**, 753–761 (2009).
- 122.D.H. Ubelaker, *Human Skeletal Remains. Excavations, Analysis, Interpretation* (Aldine, Chicago, 1978).
- 123.S. J. AlQahtani, M.P. Hector, H.M. Liversidge, The London atlas of human tooth development and eruption. *Am. J. Phys. Anthropol.* **142**, 481–490 (2010).
- 124.P. Graziosi, I Balzi Rossi. Itinerari Liguri 2 (Istituto Internazionale di Studi Liguri, Bordighera, 1976).
- 125.I. Crevecoeur *et al.*, The Spy VI child: a newly discovered Neandertal infant. *J. Hum. Evol.* **59**, 641–656 (2010).

- 126.E. Trinkaus, V. A. Ranov, S. Lauklin, Middle paleolithic human deciduous incisor from Khudji, Tajikistan. *J. Hum. Evol.* **38**, 575-583 (2000).
- 127.T. K. III. Black, Sexual dimorphism in the tooth-crown diameters of the deciduous teeth. *Am. J. Phys. Anthropol.* **48**, 77–82 (1978).

Acknowledgments: The authors are grateful to H. Temming for technical support. Research at Fumane is coordinated by the Ferrara University in the framework of a project supported by the Ministry of Culture - Veneto Archaeological Superintendency, public institutions (Lessinia Mountain Community - Regional Natural Park, Fumane Municipality, Veneto Region - Department for Cultural Heritage), and private associations and companies. We are indebted for technical assistance to Lysann Rädisch and Sven Steinbrenner of the Department of Human Evolution at the Max Planck Institute for Evolutionary Anthropology. We thank Alexander Hübner and Chiara Barbieri for their input on the phylogenetic analyses. The data described in the paper are archived in the Supplemental Material. The sequence data for the Fumane 2 specimen has been submitted to GenBank and is available under the accession number KP718913. The Bombrini tooth is housed at the Museo Preistorico Nazionale dei Balzi Rossi (Ventimiglia, Italy), while the Fumane 2 is temporally housed at the Dipartimento di Studi Umanistici, Università di Ferrara (Ferrara, Italy). This research is supported by the Max Planck Society.

Fig. #:

Fig. 1. Three-dimensional digital models of the Protoaurignacian human remains. The Bombrini tooth is a lower left lateral deciduous incisor (Ldi₂), while Fumane 2 is an upper right lateral deciduous incisor (Rdi²). B, buccal; D, distal; L, lingual; M, mesial; O, occlusal. Scale bar, 1 cm.

Fig. 2. Phylogenetic analysis of the Fumane 2 mtDNA genome, inferred using the Neighbor-Joining method. The Fumane 2 mitochondrial genome falls within the variation of modern humans, and outside the variation of Neandertals, Denisovans and a hominin from Sima de los Huesos. The insert shows the branches closest to Fumane 2. Other ancient modern humans are noted in italics. Branch lengths represent the evolutionary distance between individuals, reflected by the number of inferred substitutions per sequence.

Fig. 3. Bayesian model of dates from the 1976 excavation by Vicino at Riparo Bombrini. Bombrini AMS results compared with Grotta di Fumane boundaries created in ref. (4). Bone samples treated with ultrafiltration in grey; shell samples from the Mousterian level (4) in blue. Radiocarbon dates are calibrated in IntCal13 (29) and Marine13 (29) for shell samples. The model and boundaries were calculated using OxCal 4.2 (34) including the performance of the General t-type Outlier Model (34). The results are linked with the (NGRIP) $\delta^{18}\text{O}$ climate record (15). The grey shaded bar denotes Heinrich event 4 (HE4) (35).

Table #:

Table 1. Three-dimensional enamel thickness. Bombrini (Ldi_2) is standardized to Z-scores (for RET index) of the Neandertal and recent modern humans (RMH) di_2 sample in different wear stages

Taxon	Wear stage*	<i>n</i>	AET (mm)		RET (scale free)		Z-scores for RET index
			Mean	Range	Mean	Range	
Bombrini	4		0.29		9.22		
Neandertals	1/2	3	0.29 (0.01)	0.28-0.30	7.88 (0.33)	7.54-8.20	4.06
Neandertals	3	2	0.26 (0.007)	0.26-0.27	6.95 (0.55)	6.56-7.34	4.13
RMH	2	3	0.35 (0.006)	0.35-0.36	11.41 (0.41)	10.97-11.77	-5.34
RMH	3	11	0.31 (0.04)	0.24-0.35	9.98 (1.17)	8.01-11.85	-0.65
RMH	4	4	0.26 (0.04)	0.22-0.32	8.67 (1.4)	6.98-10.40	0.39

Standard deviation is indicated in parenthesis. AET, average enamel thickness index; RET, relative enamel thickness index. * Based on ref. (19).

Supplementary Materials:

Materials and Methods

Supplementary online text

Figures S1-S11

Tables S1-S10

References (37-127)

Author Contributions: S.B., F.N., M.P., M.A.M. and J.J.H. initiated and organized the project.

S.B. collected the fossil and modern human comparative samples for enamel thickness analysis.

D.P. and P.A.S. carried out the microCT scan of the teeth and reconstructed the digital models.

S.B. and S.E.B. provided the morphological description of the fossil sample. S.B. carried out the digital morphometric analysis of the teeth. V.S., M.M. and S.S. carried out aDNA analysis. S.B., J-J.H., V.S., S.S., M.M. and S.P. analyzed the data. S.T. initiated and performed the radiocarbon dating project at Riparo Bombrini. S.B., V.S., S.T., F.N., M.P., S.E.B., G.V., M.A.M., M.M.,

S.P., J-J.H. discussed the results. S.B., V.S., S.T., F.N., M.P., S.E.B., D.P., E.S., M.A.M., M.M.,

S.P., J-J.H. wrote and edited the manuscript.




Fig. 1




Fig. 2




Fig. 3

Modelled date (BP)