



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Human Multidrug Resistance 1 gene polymorphisms and Idiopathic Pulmonary Fibrosis

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Martinelli, M., Scapoli, L., Pacilli, A.M.G., Carbonara, P., Girardi, A., Mattei, G., et al. (2015). Human Multidrug Resistance 1 gene polymorphisms and Idiopathic Pulmonary Fibrosis. JOURNAL OF RESEARCH IN MEDICAL SCIENCES, 20(1), 93-96.

Availability:

This version is available at: <https://hdl.handle.net/11585/542085> since: 2016-06-14

Published:

DOI: <http://doi.org/>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Human Multidrug Resistance 1 gene polymorphisms and Idiopathic Pulmonary Fibrosis

Marcella Martinelli^{1,2}, Luca Scapoli^{1,2}, Angela Maria Grazia Pacilli¹, Paolo Carbonara¹, Ambra Girardi^{1,2}, Gabriella Mattei^{1,2}, Maria Teresa Rodia^{1,2}, Rossella Solmi^{1,2}

¹Department of Experimental, Diagnostic and Specialty Medicine (DIMES), Unit of Histology, Embryology and Applied Biology and ²Centre of Molecular Genetics, CARISBO Foundation, University of Bologna, Italy

Background: For the first time we tested an association between the human multidrug resistance gene 1 (MDR1) polymorphisms (SNPs) and idiopathic pulmonary fibrosis (IPF). Several MDR1 polymorphisms are associated with pathologies in which they modify the drug susceptibility and pharmacokinetics. **Materials and Methods:** We genotyped three MDR1 polymorphisms of 48 IPF patients and 100 control subjects with Italian origins. **Results:** No evidence of association was detected. **Conclusion:** There are 50 known MDR1 SNPs, and their role is explored in terms of the effectiveness of drug therapy. We consider our small-scale preliminary study as a starting point for further research.

Key words: Idiopathic pulmonary fibrosis, multidrug resistance 1 gene, polymorphism

How to cite this article: Martinelli M, Scapoli L, Pacilli A.M.G, Carbonara P, Girardi A, Mattei G, Rodia M.T, Solmi R. Human Multidrug Resistance 1 gene polymorphisms and Idiopathic Pulmonary Fibrosis. J Res Med Sci 2015;20:??-??.

INTRODUCTION

Anti-inflammatory, anti-fibrosis, immunosuppressive treatments (azathioprine, corticosteroids, cyclophosphamide) or antioxidant strategies and other therapeutic agents targeting fibrogenesis (interferons, antagonists of cytokines, growth factors or their receptors and angiogenic agents) are useless for idiopathic pulmonary fibrosis (IPF).^[1] No studies have considered a possible relationship between the human multidrug resistance 1 gene (MDR1) polymorphisms and the therapy outcome in IPF.

Multidrug resistance 1 gene is expressed in epithelial cells of a certain number of organs, including the lung.^[2] The MDR1 phenotype is acquired after administration of drugs^[3] and when expressed, it extrudes chemotherapeutic substances from cells causing the ineffectiveness of the treatment.^[2]

The MDR1 encodes for a plasma membrane MDR protein, the P-glycoprotein (P-gp)^[2] that functions as an energy-dependent membrane efflux pump to maintain cytoplasmic concentrations steady.^[3] P-gp drags out of the cell a number of substances with diverse chemical structures. It is not clear how P-gp recognizes and transports and hence various range of compounds, but it is not based on simply chemical structure.^[4]

Several drugs have not been evaluated as P-gp substrates, so additional unidentified substances could be transported by the P-gp across cell membrane.

Furthermore, variations in P-gp expression and function are associated with the pathology susceptibility^[5] and have important consequences in therapeutic outcome and disease progression.^[6] By now, 50 single nucleotide and 3 insertion/deletion polymorphisms have been mapped in the MDR1 gene.^[7] To evaluate MDR1 as possible risk factor for IPF, we analyzed the three most commonly studied MDR1 polymorphisms in our case record.

MATERIALS AND METHODS

Between 2008 and 2013, 48 consecutive Italian patients with IPF (31 males and 17 females) were enrolled for this study at the Respiratory Division, Sant'Orsola Hospital, University of Bologna. Diagnosis of IPF was made according to ATS/ERS guidelines of 2011.

A total of 100 healthy volunteers (50 males and 50 females), were enrolled as controls. The study was approved by the Ethical Committee of Sant'Orsola-Malpighi General Hospital and complied with the ethical principles for medical research involving human subjects of the Helsinki declaration. Written informed consent was obtained from all patients and healthy controls

Address for correspondence: Dr. Rossella Solmi, Dipartimento di Medicina Specialistica, Diagnostica e Sperimentale, Via Belmeloro, 8, 40126 Bologna, Italy. E-mail: rossella.solmi@unibo.it

Received: 29-10-2013; **Revised:** 04-10-2014; **Accepted:** 12-12-2014

beforehand. Peripheral blood samples were drawn from each individual and DNA extraction was performed using the GenElute Blood Genomic DNA kit (Sigma, Milan, Italy).

Three single-nucleotide polymorphisms (SNPs) mapping respectively on exon 12 (1236C>T; rs1128503), 21 (2677G>T/A; rs2032582), and 26 (3435C>T; rs1045642) of MDR1 gene were selected. Polymorphism selection was based on data available from published investigations. Among all known gene variation, we selected three SNPs that were previously found associated with other pathologies, thus having higher chances to directly alter gene function or to be linked with susceptibility variation.

Each polymorphism was amplified by polymerase chain reaction using flanking primers (supplementary data: additional file 1) and products incubated with a restriction endonuclease, BsU RI, Gsa I and Dpn II (CABRU, Milan, Italy), respectively. Fragments were separated by 10% polyacrylamide gel electrophoresis, subsequently stained with ethidium bromide (supplementary data: additional file 2). To assess the accuracy of the genotyping outputs, one-third of randomly selected samples were blindly tested by a second operator.

The distribution of genotypes in both probands and control groups was tested for deviations from the Hardy-Weinberg equilibrium using Pearson's Chi-square test. Association analyses were performed with the Unphased software v3.1.5 in a Windows Vista operative system.^[8] This included a tests for both allele and genotype association, as well as odds ratio calculation, which was shown in relation to a single reference allele or genotype.

Two and three SNPs combinations were considered for haplotype analysis. Since the likelihood ratio statistics may

be sensitive to rare haplotypes, a frequency threshold of 0.02 was adopted. An overall association analysis, that tested whether any haplotypes are associated (global *P* value), as well as a specific test for each haplotype, was carried out.

RESULTS

Genotype frequency in cases and controls were distributed accordingly with the Hardy-Weinberg equilibrium law.

Allele frequency was similar in case and control groups at all the tested loci. No evidence of the association between IPF and MDR1 genotypes was detected [Table 1].

Multilocus approach could be more sensitive than single locus analysis. For this reason, two and three locus haplotypes were tested for association with IPF. Both the global and the single haplotype tests for association rejected the hypothesis of the association between MDR1 polymorphisms and IPF (data not shown).

DISCUSSION

C1236T, G2677T/A and C3435T are the most commonly studied among the 50 known MDR1 gene SNPs in disease conditions and different populations. We did not find any differences about the frequencies of these three MDR1 polymorphisms between IPF patients and control subjects in our Italian case study. The relatively low number of patients examined could be the cause of false negative results, especially if MDR1 had minor effects on risk to develop IPF. Moreover, additional untyped polymorphisms could have an effect on IPF risk. This is only the start point to evaluate the role of the MDR1 SNPs in IPF considering the large amount of SNPs to analyze and the differences determined by the

Table 1: Case-control association analysis between MDR1 SNPs and IPF


SNP ID	Allele/genotype	IPF case	Control	IPF-frequency	Co-frequency	Chi-square	P	OR (CI)
rs1128503	C	58	111	0.60	0.56	0.64	0.42	Reference
	T	38	89	0.40	0.44			0.82 (0.50-1.34)
	CC	18	33	0.38	0.33			Reference
	CT	22	45	0.46	0.45			0.90 (0.42-1.93)
	TT	8	22	0.17	0.22			0.67 (0.25-1.80)
rs2032582	T	57	108	0.59	0.54	0.76	0.38	Reference
	A	39	92	0.41	0.46			0.80 (0.49-1.32)
	TT	17	32	0.35	0.32			Reference
	TA	23	44	0.48	0.44			0.98 (0.45-2.1)
	AA	8	24	0.17	0.24			0.63 (0.23-1.69)
rs1045642	C	52	100	0.54	0.50	0.45	0.50	Reference
	T	44	100	0.46	0.50			0.85 (0.52-1.38)
	CC	14	29	0.29	0.29			Reference
	CT	24	42	0.50	0.42			1.18 (0.53-2.67)
	TT	10	29	0.21	0.29			0.71 (0.27-1.87)

IPF = Idiopathic pulmonary fibrosis; OR = Odds ratio; CI = Confidence interval; MDR1 = Multidrug resistance 1 gene; SNPs = Single-nucleotide polymorphisms

Additional file 1: Characteristics of polymorphisms

SNP ID	Primer sequences 5'-3'	Nucleotide change ^a	Location	T _a ^b	Restriction enzyme
rs1128503	For TTGAATGAAGAGTTCTGATGTTT Rev CTCTGCATCAGCTGGACTGT	1236C>T	Exon 12	45°C	BsuRI
rs2032582	For TGTTGCTGGACAAGCACTGA Rev TGTGAAGACAATGGCCTGAA	2677G>T	Exon 21	55°C	Gsal
rs1045642	For GGCTCCGAGCACACCTGGGCA Rev GGCCAGAGAGGCTGCCACATGCT	3435C>T	Exon 26	59°C	DpnII

^aMajor allele/minor; ^bAnnealing temperature; Conditions of amplification for 35 cycles were 30 s at 94°C, 30 s at specific T_a, 30 s at 72°C and final extension for 8 min at 72°C; SNP: Single-nucleotide polymorphism



aMajor all

Additional file 2: Electrophoretic pattern on 10% acrylamide gel of multidrug resistance 1 gene polymerase chain reaction-restriction fragment length polymorphism genotyping assays: a) 1236C>T rs1128503 (exon 12), the PCR product of 237 bp has two restriction sites, b) 2677 G>T/A rs2032582 (exon 21), c) 3435C>T rs1045642 (exon 26). Numbers on the side represent fragments' length expressed in base pairs.

variety of ethnic populations.^[9] In fact, significant differences in the frequencies of MDR1 haplotypes have been found among Caucasian, African-American, and Asian-American population.^[6] We believe that further understanding of the correlation between the MDR1 genetic variations and the physiology and biochemistry of P-gp may be critically

important for the development of new treatment modalities and personalized pharmacotherapy for patients with IPF.

AUTHOR'S CONTRIBUTION

The authors' contributions were as follows: M. M. and S. R. conceived and designed the study; P. A.M.G., C. P. and R. S. were responsible for the sample collection; M. M., G. A., M. G., R. M.T., and S. R. conducted the research; S. L. was responsible for the data analysis and interpretation; S. R. wrote the manuscript; M. M. and S. L. were responsible for the critical revision of the manuscript.

ACKNOWLEDGMENTS

This work is funded by University of Bologna (M. M. and S. R.F.O. 2012).

REFERENCES

1. Farkas L, Gauldie J, Voelkel NF, Kolb M. Pulmonary hypertension and idiopathic pulmonary fibrosis: A tale of angiogenesis, apoptosis, and growth factors. *Am J Respir Cell Mol Biol* 2011;45:1-15.
2. Cao L, Owsianik G, Jaspers M, Janssens A, Cuppens H, Cassiman JJ, et al. Functional analysis of CFTR chloride channel activity in cells with elevated MDR1 expression. *Biochem Biophys Res Commun* 2003;304:248-52.
3. Comerford KM, Wallace TJ, Karhausen J, Louis NA, Montalto MC, Colgan SP. Hypoxia-inducible factor-1-dependent regulation of the multidrug resistance (MDR1) gene. *Cancer Res* 2002;62:3387-94.
4. Seelig A. How does P-glycoprotein recognize its substrates? *Int J Clin Pharmacol Ther* 1998;36:50-4.
5. Mealey KL. Therapeutic implications of the MDR-1 gene. *J Vet Pharmacol Ther* 2004;27:257-64.
6. Woodahl EL, Ho RJ. The role of MDR1 genetic polymorphisms in interindividual variability in P-glycoprotein expression and function. *Curr Drug Metab* 2004;5:11-9.
7. Li YH, Wang YH, Li Y, Yang L. MDR1 gene polymorphisms and clinical relevance. *Yi Chuan Xue Bao* 2006;33:93-104.
8. Dudbridge F. Likelihood-based association analysis for nuclear families and unrelated subjects with missing genotype data. *Hum Hered* 2008;66:87-98.
9. Al-Mohizea AM, Alkharfy KM, Bagulb KM, Alghamdi AM, Al-Jenoobi FI, Al-Muhsen S, et al. Genetic variability and haplotype profile of MDR1 in Saudi Arabian males. *Mol Biol Rep* 2012;39:10293-301.

Source of Support: Nil, **Conflict of Interest:** No conflict of interests.