

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

On parameterizations of plane rational curves and their syzygies

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

On parameterizations of plane rational curves and their syzygies / Alessandra Bernardi; Alessandro Gimigliano; Monica Idà. - In: MATHEMATISCHE NACHRICHTEN. - ISSN 0025-584X. - ELETTRONICO. - 289:5/6(2016), pp. 537-545. [10.1002/mana.201500264]

Availability:

This version is available at: <https://hdl.handle.net/11585/548392> since: 2016-07-07

Published:

DOI: <http://doi.org/10.1002/mana.201500264>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Bernardi, A., Gimigliano, A. and Idà, M. (2016), On parameterizations of plane rational curves and their syzygies. Math. Nachr., 289: 537-545.

The final published version is available online at:
<https://doi.org/10.1002/mana.201500264>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

ON PARAMETERIZATIONS OF PLANE RATIONAL CURVES AND THEIR SYZYGIES.

ALESSANDRA BERNARDI, ALESSANDRO GIMIGLIANO, AND MONICA IDÀ

ABSTRACT. Let C be a plane rational curve of degree d and $p : \tilde{C} \rightarrow C$ its normalization. We are interested in the *splitting type* (a, b) of C , where $\mathcal{O}_{\mathbb{P}^1}(-a-d) \oplus \mathcal{O}_{\mathbb{P}^1}(-b-d)$ gives the syzgies of the ideal $(f_0, f_1, f_2) \subset K[s, t]$, and (f_0, f_1, f_2) is a parameterization of C . We want to describe in which cases $(a, b) = (k, d-k)$ ($2k \leq d$), via a geometric description; namely we show that $(a, b) = (k, d-k)$ if and only if C is the projection of a rational curve on a rational normal surface in \mathbb{P}^{k+1} .

1. INTRODUCTION

We work over an algebraically closed ground field K of characteristic 0. We are interested in projective morphisms $f : \mathbb{P}^1 \rightarrow \mathbb{P}^2$, where $f = (f_0, f_1, f_2)$, $f_i \in K[s, t]_d$, which are generically injective and generically smooth over their images; thus obtaining that $f(\mathbb{P}^1)$ is a rational plane curve C parameterized by (f_0, f_1, f_2) .

In particular we are interested in studying the *splitting type* of C , which is the pair $(k, d-k)$ such that the minimal graded free resolution of the ideal (f_0, f_1, f_2) , as a graded S -module, $S = K[s, t]$, has $S(-k-d) \oplus S(k-2d)$ as its first syzygy module (see (2) in the next section).

It is easy to see (e.g. see [GHI13]) that the splitting type of C which we have just defined is also the pair $(k, d-k)$, with $2k \leq d$, such that $f^*(\Omega_{\mathbb{P}^2}(1)) \cong \mathcal{O}_{\mathbb{P}^1}(-k) \oplus \mathcal{O}_{\mathbb{P}^1}(-d+k)$.

Actually, it is well-known that any vector bundle on \mathbb{P}^1 splits as a direct sum of line bundles (see [Bir13, Gro57]), and the determination of the splitting of the pull back $f^*T_{\mathbb{P}^n}$ (or, which is equivalent, of $f^*\Omega_{\mathbb{P}^n}(1)$), together with the problem of the splitting of the normal bundle, is a quite classical and very investigated problem in Algebraic Geometry for rational curves in \mathbb{P}^n ; e.g. for $n \geq 3$ and smooth rational curves see [EVdV81, EVdV82, Ram90, Asc86, Hul81, GS81, Ber14, Ber12, Iar14, CI15]).

From a more applied point of view, this problem, mainly for plane curves, has been studied in relation to the problem of determining the implicit equations of parameterized plane curves (e.g. see [CSC98]), where the splitting type is studied via the equivalent notion of μ -basis, which is defined for the ideal of the graph of C , in the affine case; here we will also use that approach, just with the difference that we will work in $\mathbb{P}^2 \times \mathbb{P}^1$.

Our interest in the splitting type for plane curves have stemmed also by the fact that it is strictly correlated to the problem of finding the minimal free resolution of fat points in the plane (see [GHI09a, GHI09b, GHI09c, GHI13]).

We are interested in finding the relation between the geometry of the curve C and its splitting type. One possibility is to study the relation between the singularities

of C and its splitting type. For example, if C has a point of multiplicity m , then results of Ascenzi [Asc88] show that

$$(1) \quad \min(m, d - m) \leq a \leq \min\left(d - m, \left\lfloor \frac{d}{2} \right\rfloor\right);$$

see also [GHI13]. If $2m + 1 \geq d$, when m is the maximum multiplicity of the points of C , it follows from these bounds that $a = \min(m, d - m)$ and hence $b = \max(m, d - m)$. So we give the following definition.

Definition 1.1. A rational projective plane curve C is *Ascenzi* if it has a point of multiplicity m , with $2m + 1 \geq d$.

Notice that the Ascenzi behavior is not the general one, in the sense that it is possible that a curve of splitting type $(m, d - m)$ does not have any point of multiplicity m . For example the curve C of degree eight, defined by its syzygy matrix A (see (2) below), where:

$$A := \begin{pmatrix} s^3 & s^2t + st^2 & t^3 \\ s^5 + 3s^2t^3 & t^5 + 3s^3t^2 & s^5 + t^5 + st^4 \end{pmatrix},$$

has only double points as singularities while its splitting type is $(3, d - 3)$.

The non-Ascenzi cases are more difficult to handle and only partial results are known (e.g. see [GHI09a, GHI09b, GHI13]).

In [Asc88], the cases $(a, b) = (1, d - 1)$ (any d) and $(a, b) = (2, d - 2)$, $d \geq 3$, are treated; in the first case C has to be Ascenzi, while in the second either C is Ascenzi or it has only nodes as singularities and it is a projection of a smooth curve $D \subset \mathbb{P}^3$ of degree d which is contained in a quadric surface.

Hence another way to relate the geometry of C with its splitting type is to view C as projection of a curve D lying on some special surfaces.

In this paper we carry on this perspective by determining that the right surfaces to look at are rational normal scrolls. Notice that in general rational curves can be contained in rational normal scrolls only in particular cases, as the following remark shows.

Remark 1.2. Let us consider a generic projection $C \subset \mathbb{P}^m$ of a rational normal curve $C_d \subset \mathbb{P}^d$. The curve C has a good postulation, therefore $h^0(\mathcal{I}_C(2)) = \binom{m+2}{2} - 2d - 1$ ([Hir81, BE87]). On the other hand, a rational normal surface $S \subset \mathbb{P}^m$ verifies $h^0(\mathcal{I}_S(2)) = \binom{m-1}{2}$ (see e.g. [Seg84]). Hence, under a generic projection C can possibly lie on a rational normal surface only if $m \geq \frac{2d+1}{3}$.

In this paper we manage to prove, in Theorem 3.1, that for $(a, b) = (k, d - k)$, $d \geq 2k$, the curve C is the projection of a degree d curve $D \subset \mathbb{P}^{k+1}$ which is contained in a rational normal scroll (and we have the Ascenzi case if the rational scroll is a cone and D passes through its vertex with multiplicity $d - k$, otherwise D is smooth).

Hence we relate the splitting type of the plane rational curve with the existence of a rational curve on a scroll in a space of dimension “smaller than expected”. The main idea in order to construct such a scroll is to define it from the “moving line” (in the language of [CSC98]) defined by a syzygy of minimum degree of C .

The paper is organized as follows. In Section 2 we give a first result which connects the splitting type of a plane rational curve with its being a projection of a curve on a rational normal scroll. Moreover, for curves of splitting type $(3, d - 3)$,

we give an explicit parameterization of a curve on a rational normal scroll in \mathbb{P}^4 which projects to our plane curve C . In Section 3 we prove our main result, i.e. Theorem 3.1. We end the paper with Section 4 where we list a few open problems related to these questions.

2. PRELIMINARY RESULTS

We want to study rational plane curves C given via linear systems $\langle f_0, f_1, f_2 \rangle \subset K[s, t]_d$, i.e. g_d^2 's on \mathbb{P}^1 which give a projective morphism, generically one to one, $f : \mathbb{P}^1 \rightarrow \mathbb{P}^2$, whose image is C . We are interested in studying the splitting type of C , i.e. the pair $(k, d - k)$ such that the minimal graded free resolution of $J = (f_0, f_1, f_2)$, as a graded S -module, $S = K[s, t]$, is:

$$(2) \quad 0 \rightarrow S(-k-d) \oplus S(k-2d) \xrightarrow{A = \begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \beta_0 & \beta_1 & \beta_2 \end{pmatrix}} S(-d)^{\oplus 3} \xrightarrow{\langle f_0 \ f_1 \ f_2 \rangle} J \rightarrow 0,$$

where $\deg \alpha_j = k$, $\deg \beta_j = d - k$, and the f_i 's are the 2×2 minors of A , the Hilbert-Burch matrix of J . For this reason we will speak of “splitting type” not only for parameterized rational curves in \mathbb{P}^2 but more generally for finite maps $\psi : \mathbb{P}^1 \rightarrow \mathbb{P}^2$, maybe not generically one to one. More precisely we will say that ψ has splitting type $(h, k - h)$ if the ideal $(\psi_0, \psi_1, \psi_2) \subset K[s, t]$ has minimal free resolution:

$$0 \rightarrow S(-h-k) \oplus S(h-2k) \rightarrow S(-k)^{\oplus 3} \xrightarrow{\langle \psi_0 \ \psi_1 \ \psi_2 \rangle} J \rightarrow 0.$$

Our aim is to relate the splitting $(k, d - k)$ to a geometric property of C , namely to how C can be obtained as a projection of a rational curve $D \subset \mathbb{P}^{k+1}$ which is contained in a rational normal surface $S \subset \mathbb{P}^{k+1}$ (for such surfaces see e.g. [Seg84, EH87]).

The following result describes how curves on rational normal surfaces project to plane curves whose splitting type is related to the scroll itself.

Proposition 2.1. *Let $D \subset \mathbb{P}^{k+1}$ be a smooth rational curve of degree $d \geq 2k$ and assume that D is contained in a rational normal surface $S \subset \mathbb{P}^{k+1}$ and it is a unisecant for the fibers of S . Consider points $P_1, \dots, P_{k-1} \in \mathbb{P}^{k+1} - S$, such that by projecting D in \mathbb{P}^2 from them, the projection is generically 1:1 onto its image C . Then the splitting type of C is $(k', d - k')$, with $k' \leq k$.*

Proof. The fibers of S define a line bundle $\mathcal{L}_S \subset T_{\mathbb{P}^{k+1}|D}$; by [Asc88, Theorem 1.4], we have that

$$\deg \mathcal{L}_S = \deg(g_S^*(\omega)) + 2d,$$

where ω is the universal 2-vector bundle on the grassmannian $G = \mathbb{G}(1, k+1)$ and the Gauss map $g_S : D \rightarrow \mathbb{G}(1, k+1)$ is defined by sending each $P \in D$ to the point of G corresponding to the line of S passing through P .

We get that the image of g_S in $\mathbb{G}(1, k+1)$ is a rational curve Γ of degree k , in fact S is a rational normal scroll in \mathbb{P}^{k+1} , hence $\deg(S) = k$, and also $\deg(\Gamma) = k$ (e.g. see [Cos06, Section 3]).

Now, since $\deg(\Gamma) = k$ and g_S is an isomorphism between \mathbb{P}^1 and its image and $\omega|_{\Gamma} \simeq \mathcal{O}_{\mathbb{P}^1}(-k)$, we have that $\deg g_S^*(\omega) = -k$ (it is the back image of $\mathcal{O}_G(-1)|_{\Gamma}$). Hence \mathcal{L}_S is a subbundle of $T_{\mathbb{P}^{k+1}|D}$ of degree $(2d - k)$.

We get C by a projection $\nu : \mathbb{P}^{k+1} \rightarrow \mathbb{P}^2$ from points outside S , hence none of the lines belonging to the scroll S contracts via ν to a point in \mathbb{P}^2 . Let $\psi : \mathbb{P}^1 \rightarrow D \subset \mathbb{P}^{k+1}$ be a parameterization of D such that $\nu \circ \psi = f$. The image of the bundle \mathcal{L}_S through the map of vector bundles $\psi^*(T_{\mathbb{P}^{k+1}}) \rightarrow f^*(T_{\mathbb{P}^2})$ is the line bundle $\mathcal{O}_{\mathbb{P}^1}(2d - k)$.

Hence we get that $f^*(T_{\mathbb{P}^2}) \simeq \mathcal{O}_{\mathbb{P}^1}(2d - k + \delta) \oplus \mathcal{O}_{\mathbb{P}^1}(k + d - \delta)$, for some δ , $k - 1 \geq \delta \geq 0$. Now, dualizing and twisting by 1

$$f^*(\Omega_{\mathbb{P}^2}(1)) \cong \mathcal{O}_{\mathbb{P}^1}(-k + \delta) \oplus \mathcal{O}_{\mathbb{P}^1}(-d + k - \delta),$$

i.e. the splitting type of C is $(k', d - k')$, with $k' = k - \delta \leq k$. \square

In [Asc88], Ascenzi showed that when the splitting type is $(2, d - 2)$, C can be viewed as the projection of a curve $D \subset \mathbb{P}^3$ contained in a quadric surface S from a point outside S . The surface S and the curve D are smooth except when C is Ascenzi, i.e. it possesses a point of multiplicity $d - 2$; in that case S is a quadric cone and D has multiplicity $d - 2$ at the vertex of S (its only singularity).

Ascenzi gives a direct construction of the quadric S and of the parameterization of D in \mathbb{P}^3 , via the Hilbert-Burch matrix of (f_0, f_1, f_2) . We want to generalize this point of view to any possible splitting type for a reduced, irreducible plane rational curve.

For curves with splitting type $(3, d - 3)$, we are able to strictly follow Ascenzi's idea, in order to construct S and D as in the following

Proposition 2.2. *Let $C \subset \mathbb{P}^2$ be a rational curve of degree $d \geq 6$, with splitting type $(3, d - 3)$; then there is a rational curve D and a rational normal surface S , with $D \subset S \subset \mathbb{P}^4$, such that C is the projection of D from two points not on S . Moreover:*

- *if C is not Ascenzi, then D and S are smooth;*
- *if C is Ascenzi, then S is a cone, D passes through the vertex of S with multiplicity $(d - 3)$ and that one is its only singular point.*

Proof. If the splitting type of C is $(3, d - 3)$, from (2) we get that we can view C as parameterized by the maximal minors of

$$(3) \quad A = \begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \beta_0 & \beta_1 & \beta_2 \end{pmatrix}$$

where $\deg \alpha_j = 3$ and $\deg \beta_j = d - 3$. Now suppose C is not Ascenzi, i.e. it does not possess a point of multiplicity $d - 3$. This implies that its degree 3 syzygy has length 3, i.e. the α_i 's are linearly independent (see [Asc88]), and, without loss of generality, we can suppose that

$$\alpha_0 = s^3, \quad \alpha_1 = s^2t + st^2, \quad \alpha_2 = t^3$$

In fact, if C has no points of multiplicity $d - 3$, then $\alpha_0, \alpha_1, \alpha_2$ span a plane H in the space \mathbb{P}^3 parameterizing forms of degree 3 in $K[s, t]$. Let H meet in three distinct points the rational normal curve C_3 (whose points correspond to forms which are powers of linear forms); via a linear change of coordinates in \mathbb{P}^1 , which leaves C unchanged, we can suppose that those three points correspond to $s^3, (s + t)^3$ and t^3 . Since what we need is to keep the same g_d^2 defining C , we can change the parametrization (f_0, f_1, f_2) (i.e. the generators we pick for the linear space

$\langle f_0, f_1, f_2 \rangle$) and we can work modulo projectivities on $\langle f_0, f_1, f_2 \rangle$; in particular if we consider (from Equation 2):

$$\begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \beta_0 & \beta_1 & \beta_2 \end{pmatrix} \cdot \begin{pmatrix} f_0 \\ f_1 \\ f_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

we can change to:

$$\begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \beta_0 & \beta_1 & \beta_2 \end{pmatrix} \cdot P \cdot P^{-1} \cdot \begin{pmatrix} f_0 \\ f_1 \\ f_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

where P is a 3×3 invertible matrix, and P^{-1} will give a projectivity in the plane $\langle f_0, f_1, f_2 \rangle$, so we can consider the curve given by parameterization:

$$P^{-1} \cdot \begin{pmatrix} f_0 \\ f_1 \\ f_2 \end{pmatrix}, \quad \text{which has syzygy matrix:} \quad \begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \beta_0 & \beta_1 & \beta_2 \end{pmatrix} \cdot P.$$

Hence, modulo two projectivities, one in H and one in $\langle f_0, f_1, f_2 \rangle$, given by appropriate P and P^{-1} , we can suppose $\alpha_0 = s^3$; $\alpha_1 = s^2t + st^2$; $\alpha_2 = t^3$ (since $s^2t + st^2 = 1/3[(s+t)^3 - s^3 - t^3]$).

Notice that if H were to meet C_3 only in two points, say s^3 and t^3 , this would imply that the tangent line to C_3 at one of them, say s^3 , is contained in H ; in this case we can choose α_1 on the tangent line of the rational normal cubic curve at s^3 , hence we can use: $\alpha_0 = s^3$; $\alpha_1 = s^2t$; $\alpha_2 = t^3$, and all what follows could be done with the same procedure (with even easier computations).

On the other hand, it is not possible for H to meet C_3 at only one point, since in this case H would be the osculating plane of C_3 at this point, say s^3 , and all the forms parameterized by H would have s as a common factor, hence $(\alpha_0, \alpha_1, \alpha_2)$ could not be a primitive syzygy.

Now set

$$\begin{cases} \phi_0 = s^2\beta_1 - \beta_0(st + t^2) \\ \phi_1 = \beta_2(s^2 + st) - \beta_1t^2 \\ \phi = \beta_1s^2t - \beta_0st^2 + \beta_2s^2t - \beta_1st^2 \end{cases}.$$

Let us consider our parameterization of C which can be written as:

$$\begin{cases} f_0 = \alpha_1\beta_2 - \alpha_2\beta_1 = s[s^2\beta_1 - \beta_0(st + t^2)] = s\phi_0 \\ f_1 = \alpha_2\beta_0 - \alpha_0\beta_2 = \beta_0t^3 - \beta_2s^3 \\ f_2 = \alpha_0\beta_1 - \alpha_1\beta_0 = t[\beta_2(s^2 + st) - \beta_1t^2] = t\phi_1 \end{cases}.$$

Since

$$t\phi_0 + s\phi_1 = \beta_1s^2t - \beta_0st^2 + \beta_2s^2t - \beta_1st^2 - (\beta_0t^3 - \beta_2s^3) = \phi - f_1,$$

we get:

$$\begin{cases} f_0 = s\phi_0 \\ f_1 = \phi - (t\phi_0 + s\phi_1) \\ f_2 = t\phi_1 \end{cases}.$$

Now we can consider the curve $D \subset \mathbb{P}^4$ given by the parameterization:

$$\begin{cases} x_0 = s\phi_0 \\ x_1 = -t\phi_0 \\ x_2 = -s\phi_1 \\ x_3 = t\phi_1 \\ x_4 = \phi \end{cases}.$$

We get that C can be viewed as the projection of D on the plane $H = \{x_2 = x_4 = 0\}$ from the points $P_1 = (0, 1, -1, 0, 0)$ and $P_2 = (0, 1, 0, 0, -1)$.

We want to show that D is smooth. It not hard (e.g. by using CoCoA [ABL08]) to see that D is contained in three quadric hypersurfaces $\{x_0x_3 - x_1x_2, x_2x_3 - x_2x_4 + x_3x_4 + x_0x_3, x_0x_1 + x_0x_4 - x_1x_4 - x_1x_3\}$, which generate the ideal of a smooth rational normal surface S such that $P_1, P_2 \notin S$. The structure of S is well

known; $\text{Pic}(S)$ is generated by C_0, f , where f is a fiber and C_0 is a curve with $C_0^2 = -e = -1$, $C_0 \cdot f = 1$ and $f^2 = 0$.

It is not hard to check that the lines of the ruling of S are contained in the planes of the ruling of the quadric cone $\mathcal{Q} = \{x_0x_3 - x_1x_2 = 0\}$, and such planes are unisecant (outside its vertex) to D . Hence D has to be of type $(C_0 + yf)$ on S and so it is smooth.

We also have that the hyperplane section of S is $C_0 + 2f$, and, since D has degree d , we have:

$$D \cdot H = -1 + y + 2 = 1 + y = d$$

which yields $D = C_0 + (d - 1)f$.

Now we sketch the construction in the Ascenzi case (similar to what we just did). If C has a point of multiplicity $d - 3$, we can suppose

$$\alpha_0 = s^2(as + bt); \alpha_1 = t^2(cs + dt), \alpha_2 = 0,$$

for some $a, b, c, d \in K$, with $a \neq 0$, $d \neq 0$ (since α_0, α_1 do not have common factors).

Working as in the previous case, we can construct the curve $D \in \mathbb{P}^4$ given by the parameterization:

$$\begin{cases} x_0 = d\beta_2t^3 \\ x_1 = a\beta_2s^3 \\ x_2 = (a' + b)\beta_2s^2t \\ x_3 = (c + d')\beta_2st^2 \\ x_4 = \phi \end{cases}$$

where $\phi = f_2 - (a'\beta_2s^2t + d'\beta_2st^2)$, $a' = a$ if $b = 0$ and $a' = 0$ if $b \neq 0$, $d' = d$ if $c = 0$ and $d' = 0$ if $c \neq 0$. It is easy to check that the curve D lies on a rational normal scroll S which is a cone, with vertex $V = (1, 0, 0, 0, 0)$ and D has a $(d - 3)$ -ple point at V , given by the $d - 3$ zeroes of β_2 . Hence also in this case C is a projection of D as requested, and we are done. \square

This way to directly construct the curve D on a scroll is not so feasible when $d \geq 4$; we will follow another attack in order to generalize this idea in the next section.

3. THE MAIN THEOREM

In this section we state and prove our main result:

Theorem 3.1. *Let $C \subset \mathbb{P}^2$ be a rational curve of degree d . The splitting type of C is $(k, d - k)$, $2 \leq 2k \leq d$, if and only if there is a degree d rational curve, $\mathcal{D} \subset \mathbb{P}^{k+1}$ which is contained in a rational normal surface (scroll) $S_{h,k-h} \subseteq \mathbb{P}^{k+1}$, with $2h \leq k$, the curve C is the projection of \mathcal{D} from $(k - 1)$ points outside $S_{h,k-h}$ and either*

- i) $h > 0$, $S_{h,k-h}$ and \mathcal{D} are smooth (and C is not Ascenzi), or*
- ii) $S_{0,k}$ is a cone, \mathcal{D} passes through the vertex V of the cone with multiplicity $d - k$ and V is its only singular point (this covers also the case $S_{0,1} = \mathbb{P}^2$, for which $C = \mathcal{D}$). In this case C is Ascenzi.*

Moreover \mathcal{D} is uniscant to the fibers of $S_{h,k-h}$ and h is the degree of a syzygy of minimal degree for a syzygy of minimal degree k of C .

Proof. Notice that for $k = 1, 2$, the theorem is known, proved by Ascenzi in [Asc88], while for $k = 3$ it can be deduced from Proposition 2.2 above.

Let C be parameterized by $(f_0, f_1, f_2) \in K[s, t]_d$, then the ideal $(g_0, g_1, g_2) = (x_2 f_0 - x_0 f_2, -x_1 f_2 + x_2 f_1, x_0 f_1 - x_1 f_0) \subset K[x_0, x_1, x_2, s, t]$ defines a curve D in $\mathbb{P}^2 \times \mathbb{P}^1$, which is the graph of the map $\mathbb{P}^1 \rightarrow \mathbb{P}^2$ (the projection of D on the \mathbb{P}^1 factor is 1:1 and surjective while the projection on the \mathbb{P}^2 factor is C).

Let $I_D \subset K[x_0, x_1, x_2, s, t]$ be the ideal of D , we have that a polynomial $A_0 x_0 + A_1 x_1 + A_2 x_2 \in K[x_0, x_1, x_2, s, t]_{1,n}$ vanishes on D if and only if $A_0 f_0 + A_1 f_1 + A_2 f_2$ is identically zero in $K[s, t]$ (in the language of [CSC98], this is a *moving line of degree n for C*), i.e. the spaces $(I_D)_{1,n}$ correspond to the syzygies of degree n of the ideal $(f_0, f_1, f_2) \subset K[s, t]$.

Hence we have that the splitting type of C is $(k, d - k)$, $2k \leq d$, if and only if k is the first integer for which $(I_D)_{1,k} \neq \{0\}$, and we can distinguish two cases: either $2k < d$ or $2k = d$. From what we know about the syzygies of (f_0, f_1, f_2) , if $2k < d$, there is a unique (up to multiplication by constants) form $p \in (I_D)_{1,k}$, namely $p = \alpha_0 x_0 + \alpha_1 x_1 + \alpha_2 x_2$; all elements of $(I_D)_{1,n}$, for $n = k + 1, \dots, d - k - 1$ are of type λp , with $\lambda \in K[s, t]_{n-k}$, and there is an element $q \in (I_D)_{1,d-k} - (p)_{1,d-k}$ such that $(I_D)_{1,d-k} = (p, q)_{1,d-k}$ (of course we can choose q modulo elements of $(p)_{1,d-k}$). If $2k = d$ there are two independent forms $p, p_1 \in (I_D)_{1,k}$ which generate $\bigoplus_l (I_D)_{1,l}$.

Now let us consider the moving line of degree k for C , given by $p = \alpha_0 x_0 + \alpha_1 x_1 + \alpha_2 x_2 \in K[x_0, x_1, x_2, s, t]$; $p = 0$ is the equation of a rational scroll $S \subset \mathbb{P}^2 \times \mathbb{P}^1$: for every $(s, t) \in \mathbb{P}^1$ we have a line in the corresponding \mathbb{P}^2 -fiber.

Lemma 3.2. *The scroll S and the curve D are smooth.*

Proof. Let us consider the partial derivatives of p :

$$\left(\frac{\partial p}{\partial x_0}, \frac{\partial p}{\partial x_1}, \frac{\partial p}{\partial x_2}; \frac{\partial p}{\partial s}, \frac{\partial p}{\partial t} \right) = (\alpha_0, \alpha_1, \alpha_2; p_s, p_t).$$

Since the α_i 's cannot have common factors (they represent a minimal syzygy), we get that in the above vector the first part in the α 's is never zero at any point of

S . From this it is not hard to check that we can find affine charts where the vector of local derivatives is always non-zero, hence S is smooth.

Then, since D is a curve on S which is unisecant for its fibers, D is smooth. \square

We want to show that the scroll $S_{h,k-h}$ which appears in our statement can be obtained from the scroll $S \subset \mathbb{P}^2 \times \mathbb{P}^1$ that we have just defined as $\{p=0\}$, as the image of S under the map given by a suitable linear system on it.

The idea is to consider the ideal $(\alpha_0, \alpha_1, \alpha_2) \subset K[s, t]$, which in turns gives a finite map $\mathbb{P}^1 \rightarrow \mathbb{P}^2$, maybe not generically one to one. Let us consider the splitting type of $(\alpha_0, \alpha_1, \alpha_2)$, say $(h, k-h)$, $2h \leq k$, where h is the degree of the minimal syzygy $(\gamma_0, \gamma_1, \gamma_2)$; we will get as before a curve $\Gamma \in \mathbb{P}^2 \times \mathbb{P}^1$ and an ideal (p_1, q_1) , where

$$q_1 = \gamma_0 x_0 + \gamma_1 x_1 + \gamma_2 x_2, \quad q_2 = \delta_0 x_0 + \delta_1 x_1 + \delta_2 x_2,$$

and

$$\begin{pmatrix} \gamma_0 & \gamma_1 & \gamma_2 \\ \delta_0 & \delta_1 & \delta_2 \end{pmatrix}$$

is the syzygy matrix of $(\alpha_0, \alpha_1, \alpha_2)$. Hence the δ_i 's give a primitive syzygy of degree $k-h$ for $(\alpha_0, \alpha_1, \alpha_2)$.

Lemma 3.3. *When we consider S as a Hirzebruch surface $S = \mathbb{F}_e$, with $C_0^2 = -e$, we have $e = k - 2h$ and C_0 is the graph of the curve in \mathbb{P}^2 parameterized by $(\gamma_0, \gamma_1, \gamma_2)$. Moreover, we have that the graph of the curve parameterized by $(\delta_0, \delta_1, \delta_2)$ is $C_1 \sim C_0 + ef$ (on S), and the curve D is, as a divisor on S , $D \sim C_0 + (d-h)f$.*

Proof of the Lemma. The unisecant divisors on S are in 1-1 correspondence with syzygies of $(\alpha_0, \alpha_1, \alpha_2)$ (this is quite obvious, see also [CSC98]), and for any such syzygy (A_0, A_1, A_2) , with $A_i \in K[s, t]_n$, $n \geq h$, we have:

$$(A_0, A_1, A_2) = \lambda(\gamma_0, \gamma_1, \gamma_2) + \mu(\delta_0, \delta_1, \delta_2)$$

where $\lambda \in K[s, t]_{n-h}$, $\mu \in K[s, t]_{n-k+h}$. For any such syzygy, the graph of the curve parameterized by (A_0, A_1, A_2) in \mathbb{P}^2 is a divisor on S which is unisecant to its fibers (and is cut out in $\mathbb{P}^2 \times \mathbb{P}^1$ by $x_0 A_1 - x_1 A_0, x_0 A_2 - x_2 A_0, x_1 A_2 - x_2 A_1$).

When $h \neq k/2$ we consider the unique effective divisor C_0 which is associated to the minimal syzygy $(\gamma_0, \gamma_1, \gamma_2)$, we have that for any $n > h$, we get divisors which are linearly equivalent to $C_0 + (n-h)f$ (where f is a fiber); hence C_0 is the only unisecant divisor class which has $H^0(\mathcal{O}_S(C_0)) = 1$ and which does not contain divisors containing fibers, i.e. we have $S \cong \mathbb{F}_e$, as a Hirzebruch surface, with $C_0^2 = -e$. Moreover, we get that all the effective divisors given by syzygies of degree $h+1 \leq n \leq k-h-1$ are not irreducible and are given by C_0 plus $n-h$ fibers (the zeroes of λ); the first value of n for which we have a class of divisors $C_0 + (n-h)f$ whose generic element is an irreducible one, is $n = k-h$, for which we have C_1 given by the graph of $(\delta_0, \delta_1, \delta_2)$. Notice that the curve C_1 is actually linearly equivalent to $C_0 + ef$, since they differ by a principal divisor that, locally, is of the form $\frac{x_i \delta_j - x_j \delta_i}{\lambda(x_u \gamma_v - x_v \gamma_u)}$. Hence, for what we know on rational ruled surfaces (e.g. see [Seg84], or [Har77, Ch. V, Thm 2.17]), $C_1 \sim C_0 + (k-2h)f$, $e = k-2h$, and $C_0 \cdot C_1 = 0$.

If we consider the curve D , this is the graph of (f_0, f_1, f_2) , which is a syzygy of degree d for $(\alpha_0, \alpha_1, \alpha_2)$, hence we have $D \sim C_0 + (d - h)f$.

When $h = k/2$ the choice of the C_0 is not unique, in fact we can choose as $(\gamma_0, \gamma_1, \gamma_2)$ and $(\delta_0, \delta_1, \delta_2)$ any basis of the module of syzygies of degree h , and then we can proceed as in the previous case (here $C_0 \sim C_1$). \square

Once the lemma is true, we can consider the linear system $H \sim C_0 + (k - h)f$ on S ; this is very ample if $h > 0$, and it will give an embedding of S as a surface of degree $H^2 = C_0^2 + 2(k - h) = 2h - k + 2k - 2h = k$ in \mathbb{P}^{k+1} (i.e. as a rational normal scroll); the embedded scroll is of type $S_{h, k-h}$, i.e. it is given by the lines joining two common parameterizations for two rational normal curves, which are the embedding of C_0 and C_1 , of degrees $h, k - h$, in a $\mathbb{P}^h, \mathbb{P}^{k-h}$, respectively.

In the case $h = 0$, the linear system is not very ample, but it still gives a map to \mathbb{P}^{k+1} ; it only contracts C_0 to a point, and we still get a rational normal scroll $S_{0, k}$, which is a cone.

Notice also that the case $h = 0$ corresponds to the case when C is Ascenzi, since $(\gamma_0, \gamma_1, \gamma_2)$ has degree 0, hence, modulo a projective change of coordinates, we can assume $\alpha_2 = 0$, so $(\alpha_0, \alpha_1, 0)$ has a syzygy of degree 0, namely $(0, 0, 1)$; in this case, $C_0 = \{x_0 = x_1 = 0\}$, $(\delta_0, \delta_1, \delta_2) = (-\alpha_1, \alpha_0, 0)$ and the scroll we get is the cone $S_{0, k}$. The curve \mathcal{D} , since $D \cdot C_0 = -e + d = d - k$, is singular in the vertex of the cone, with multiplicity $d - k$. Moreover, in the case $k = 1$, $S_{0, 1}$ is \mathbb{P}^2 itself.

Now we have to check that the curve $\mathcal{D} \subset S_{h, k-h} \subset \mathbb{P}^{k+1}$ that we have constructed really projects to $C \subset \mathbb{P}^2$.

The embedding of S is defined by the linear system $|H| = |C_0 + (k - h)f|$, which corresponds to syzygies of $(\alpha_0, \alpha_1, \alpha_2)$ of degree k ; if $(A_{0,i}, A_{1,i}, A_{2,i})$, $i = 0, \dots, k + 1$ is a base for the module of syzygies of degree k , to any of them it is associated a divisor on S which is the graph of the curve in \mathbb{P}^2 parameterized by $(A_{0,i}, A_{1,i}, A_{2,i})$; such divisor can be locally defined by one of the three functions

$$x_2 A_{1,i} - x_1 A_{2,i}, \quad x_0 A_{2,i} - x_2 A_{0,i}, \quad x_0 A_{1,i} - x_1 A_{0,i};$$

hence the map $\phi : S \rightarrow \mathbb{P}^{k+1}$ given by $|H|$ is defined, locally, by such equations.

Consider the three trivial syzygies: $(0, \alpha_2, -\alpha_1)$, $(\alpha_2, 0, -\alpha_0)$, $(\alpha_1, -\alpha_0, 0)$ as the last three syzygies of our base; they define a plane $\Pi \cong \mathbb{P}^2$ inside our \mathbb{P}^{k+1} . We want to check that the projection of \mathbb{P}^{k+1} on Π maps \mathcal{D} onto our starting curve C .

The three local equations we saw above, for each of these syzygies, can be taken to be (by considering that on S we have $x_0 \alpha_0 + x_1 \alpha_1 + x_2 \alpha_2 = 0$):

$$\begin{aligned} x_0 \alpha_2, \quad x_0 \alpha_1, \quad -x_2 \alpha_2 - x_1 \alpha_1 &= x_0 \alpha_0; & x_1 \alpha_2, \quad -x_2 \alpha_2 - x_0 \alpha_0 &= x_1 \alpha_1, \quad x_1 \alpha_0; \\ -x_1 \alpha_1 - x_0 \alpha_0 &= x_2 \alpha_2, \quad x_2 \alpha_1, \quad x_2 \alpha_0. \end{aligned}$$

Since $(\alpha_0, \alpha_1, \alpha_2)$ cannot have common factors, at any point $(x_0, x_1, x_2; s, t) \in S$, there is at least one $\alpha_i(s, t) \neq 0$; so its image under the projection can be written $(\alpha_i(s, t)x_0, \alpha_i(s, t)x_1, \alpha_i(s, t)x_2)$. Hence the projection map from $\phi(S) = S_{h, k-h}$ to Π can actually be written as: $\phi'(x_0, x_1, x_2; s, t) \mapsto (x_0, x_1, x_2)$, so the image of the curve \mathcal{D} is actually our starting curve C .

Notice that, by Proposition 2.1, C cannot be the projection of a curve D on a rational normal scroll in $\mathbb{P}^{k'+1}$ with $k' < k$. \square

Remark 3.4. We would like to notice that the construction developed in the proof of Theorem 3.1 can be extended in order to see our curve $\mathcal{D} \subset \mathbb{P}^{k+1}$ as the projection

of a rational normal curve.

If we consider, on S , the linear system \mathcal{L} which is given by the divisors in $|C_0 + (d-h)f|$ which have $(d-k+1)$ fixed points $P_1, \dots, P_{d-k}, P_{d-k+1}$, where P_1 and P_{d-k+1} are on the same line and P_1, \dots, P_{d-k} are on $C_0 \cap D$, we can embed S into \mathbb{P}^d as a rational normal scroll $S_{h,d-h-1}$ and the image of D will be a rational normal curve. By projecting $S_{h,d-h-1}$ from $d-k-1$ other points $P_{d-k+2}, \dots, P_{2d-k}$, where P_i and P_{d-k+i} are on the same line, we will get the scroll $S_{h,k-h} \subset \mathbb{P}^{k+1}$ of our previous construction.

4. OPEN PROBLEMS

In this section we give a list of open problems.

4.1. The center of projection. As outlined in Remark 3.4 it would be interesting to understand the relation between the splitting type of the plane curve C and the geometry of the linear space Π , with $\Pi \simeq \mathbb{P}^{d-3}$ which is the center of the projection that sends the rational normal curve $C_d \subset \mathbb{P}^d$ to $C \subset \mathbb{P}^2$. What we know, by Remark 3.4, is that there is a rational normal scroll containing C_d and intersecting Π in $d-k-1$ points.

4.2. Singularities and splitting type. Consider linear systems $\mathcal{L}(d; m_1, \dots, m_s)$ of plane curves of degree d and multiplicities at least m_i at generic points $P_i \in \mathbb{P}^2$, $i = 1, \dots, s$, such that the generic curve C in $\mathcal{L}(d; m_1, \dots, m_s)$ is rational. Find in which cases $(d; m_1, \dots, m_s)$ determines the splitting type of C . Of course a well known example when this happens is the Ascenzi case (e.g. a generic curve in $\mathcal{L}(d; d-1, 1^s)$, $s \leq 2d$, has splitting type $(1, d-1)$). A different (non-Ascenzi) example of such a situation is given by $\mathcal{L}(8; 3^7)$, i.e. curves of degree $d = 8$ with seven points of multiplicities $m_i = 3$. In this case the splitting type of a generic curve C in this linear system has to be $(3, 5)$ (e.g. see [GHI09b]).

Notice that just knowing the number and multiplicities of the singularities of C in general is not enough to determine the splitting type of C . For example the generic projection in the plane of a degree 8 rational normal curve has only nodes as singularities and splitting type $(4, 4)$. On the other hand the example given in the Introduction is of a plane curve degree 8 with only nodes as singularities but splitting type $(3, 5)$, and, by Ascenzi's results, there is a plain curve C which is a generic projection of a rational curve D in \mathbb{P}^3 of type $(1, 7)$ on a smooth quadric, which has splitting type $(2, 6)$ and only nodes as singularities. Clearly, in all cases, the nodes are not supported at generic points.

A deep analysis of the possibilities to describe singularities of certain rational plane curves via its parameterizations and its syzygies can be found in [CKPU13]

4.3. Curves in higher dimensional spaces. Generalize the type of construction that we give in Theorem 3.1 for curves in \mathbb{P}^n with $n \geq 3$, i.e. give an explicit connection between the parameterization of C , its splitting type and the rational scrolls containing C or a curve D in higher space whose projection is C . Notice that the idea of μ -basis is considered in [CSC98] also in the case of curves in higher spaces.

REFERENCES

- [ABL08] J. Abbott, A.M. Bigatti, and G. Lagorio. CoCoA-5: a system for doing Computations in Commutative Algebra. Available at <http://cocoa.dima.unige.it>, 2008.
- [Asc86] M.G. Ascenzi. The restricted tangent bundle of a rational curve on a quadric in \mathbb{P}^3 . *Proc. Amer. Math. Soc.*, 98(4):561–566, 1986.
- [Asc88] M.G. Ascenzi. The restricted tangent bundle of a rational curve in \mathbb{P}^2 . *Comm. Algebra*, 16(11):2193–2208, 1988.
- [BE87] E. Ballico and Ph. Ellia. The maximal rank conjecture for non special curves in \mathbb{P}^n . *Math. Z.*, 196:355–367, 1987.
- [Ber12] Alessandro Bernardi. Apolar ideal and normal bundle of rational curves. <http://arxiv.org/abs/1203.4972>, 03 2012.
- [Ber14] Alessandro Bernardi. Normal bundle of rational curves and Waring decomposition. *J. Algebra*, 400:123–141, 2014.
- [Bir13] G. Birkhoff. A theorem on matrices of analytic functions. *Math. Ann.*, 74(1):122–133, 1913.
- [CI15] D. A. Cox and A. A. Iarrobino. Strata of rational space curves. *Comput. Aided Geom. Des.*, 32(C):50–68, January 2015.
- [CKPU13] D. Cox, A.R. Kustin, C. Polini, and B. Ulrich. *A study of singularities on rational curves via syzygies*. (Memoirs of the American Mathematical Society; no.1045, 2013.
- [Cos06] I. Coskun. Degenerations of surface scrolls and the gromov-witten invariants of grassmannians. *J. Algebraic Geometry*, 15:223–284, 2006.
- [CSC98] D. Cox, T.W. Sederburg, and F. Chen. The moving line ideal basis of planar rational curves. *Computer Aided Geometric Design*, 15:803–827, 1998.
- [EH87] David Eisenbud and Joe Harris. On varieties of minimal degree (a centennial account). In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 3–13. Amer. Math. Soc., Providence, RI, 1987.
- [EVdV81] D. Eisenbud and A. Van de Ven. On the normal bundles of smooth rational space curves. *Math. Ann.*, 256(4):453–463, 1981.
- [EVdV82] D. Eisenbud and A. Van de Ven. On the variety of smooth rational space curves with given degree and normal bundle. *Invent. Math.*, 67(1):89–100, 1982.
- [GHI09a] A Gimigliano, B. Harbourne, and M. Idà. Betti numbers for fat point ideals in the plane: a geometric approach. *Trans. Amer. Math. Soc.*, 361:1103–1127, 2009.
- [GHI09b] A Gimigliano, B. Harbourne, and M. Idà. The role of the cotangent bundle in resolving ideals of fat points in the plane. *Journal of Pure and Applied Algebra*, 213:203–214, 2009.
- [GHI09c] A Gimigliano, B. Harbourne, and M. Idà. Stable postulation and stable ideal generation: Conjectures for fat points in the plane. *Bull. Belg. Math. Soc. Simon Stevin*, 16(5):853–860, 2009.
- [GHI13] A Gimigliano, B. Harbourne, and M. Idà. On plane rational curves and the splitting of the tangent bundle. *Ann. Sc. Norm. Super. Pisa VI. Sci.*, XII(5):1–35, 2013.
- [Gro57] A. Grothendieck. Sur la classification des fibrés holomorphes sur la sphère de riemann. *Amer. J. Math.*, 79:121–138, 1957.
- [GS81] F. Ghione and G. Sacchiero. Normal bundle of rational curves in \mathbb{P}^3 . *Manuscripta Math.*, 33(2):111–128, 1980/81.
- [Har77] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Hir81] A. Hirschowitz. Sur la postulation générique des courbes rationnelles. *Acta. Math.*, 146:209–230, 1981.
- [Hul81] K. Hulek. The normal bundle of a curve on a quadric. *Math. Ann.*, 258:201–206, 1981.
- [Iar14] A. A. Iarrobino. Strata of vector spaces of forms in $R = \mathbb{K}[x, y]$, and of rational curves in \mathbb{P}^k . *Bull. Braz. Math. Soc. (N.S.)*, 45(4):711–725, 2014.
- [Ram90] L. Ramella. La stratification du schéma de Hilbert des courbes rationnelles de \mathbf{P}^n par le fibré tangent restreint. *C. R. Acad. Sci. Paris Sér. I Math.*, 311(3):181–184, 1990.
- [Seg84] C. Segre. *Sulle rigate razionali in uno spazio lineare qualunque*. Loescher, 1884.

(Alessandra Bernardi) DIPARTIMENTO DI MATEMATICA, UNIV. BOLOGNA, ITALY

E-mail address: `alessandra.bernardi5@unibo.it`

(Alessandro Gimigliano) DIPARTIMENTO DI MATEMATICA E CIRM, UNIV. BOLOGNA, ITALY

E-mail address: `Alessandr.Gimigliano@unibo.it`

(Monica Idà) DIPARTIMENTO DI MATEMATICA, UNIV. BOLOGNA, ITALY

E-mail address: `monica.ida@unibo.it`