



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Galeati, G., Zannoni, A., Spinaci, M., Bucci, D., Ostanello, F., Panarese, S., et al. (2016). Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure. *ANIMAL SCIENCE JOURNAL*, 87(4), 511-516 [10.1111/asj.12465].

Availability:

This version is available at: <https://hdl.handle.net/11585/550122> since: 2019-07-17

Published:

DOI: <http://doi.org/10.1111/asj.12465>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the peer reviewed version of the following article:

Galeati G, Zannoni A, Spinaci M, Bucci D, Ostanello F, Panarese S, Tamanini C, Sarli G. Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure. *Anim Sci J.* 2016;87(4):511-6.

which has been published in final form at
<https://doi.org/10.1111/asj.12465>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

2 **Porcine circovirus type 2 detection in *in vitro* produced porcine blastocysts after**
3 **virus sperm exposure**

4 Giovanna GALEATI*, Augusta ZANNONI, Marcella SPINACI, Diego BUCCI, Fabio

5 OSTANELLO, Serena PANARESE, Carlo TAMANINI and Giuseppe SARLI

6 Department of Veterinary Medical Sciences, University of Bologna, Bologna, Italy

7

8 **Running title:** PCV2 and porcine blastocysts

9

10 *Corresponding author. Tel.: +39 051 2097910; fax: +39 051 2097899.

11 Dipartimento di Scienze Mediche Veterinarie (DIMEVET)

12 Via Tolara di Sopra, 50

13 40064 Ozzano dell'Emilia, Bologna, Italy

14 E-mail address: giovanna.galeati@unibo.it

15

16 **ABSTRACT**

17 This study was aimed at assessing the capability of semen experimentally infected with
18 porcine circovirus type 2 (PCV2) to produce porcine blastocysts PCR positive for
19 PCV2. Embryos were obtained from *in vitro* maturation (IVM) and *in vitro* fertilization
20 (IVF) of porcine oocytes or by parthenogenesis. Sperm suspension was exposed to
21 PCV2b and utilized for IVF. PCV2 spiked semen did not reveal any reduction in sperm
22 viability and motility but its ability to produce infected blastocysts was irrelevant as
23 only one out of 15 blastocysts obtained by IVF were PCV2b; however 2 blastocysts
24 were PCV2a positive. Furthermore, the presence of PCV2 was demonstrated also in
25 embryos obtained by parthenogenesis (1 out of 17 was PCV2b and one PCV2a positive).
26 Even if PCV2 firmly attaches to the surface of spermatozoa, experimentally spiked
27 sperm were not effective in infecting oocytes during IVF and in producing PCR positive
28 embryos. The infected blastocysts we obtained derived most probably from infected
29 oocytes recovered at abattoir.

30

31 **Keywords**

32 Blastocysts, IVM-IVF, PCV2, pig, spermatozoa.

33

34 INTRODUCTION

35 Porcine circovirus type 2 (PCV2) related conditions cause economic losses to the pig
36 industry worldwide. Apart from post-weaning multisystemic wasting syndrome
37 (PMWS), PCV2 is associated with a number of conditions collectively known as
38 porcine circovirus diseases (PCVD) (Segales et al. 2004, 2005; Chae 2005). A recent
39 line of studies has focused on PCV2-associated reproductive failure, including irregular
40 return to oestrus, failure of pregnancy progressing to abortion, or reduced litter size with
41 PCV2 being detected in aborted and stillborn foetuses (O'Connor et al. 2001;
42 Ladekjaer-Mikkelsen et al. 2001; Sanchez et al. 2001). Despite the fact that PCV2 is
43 now being regarded as a potentially important emerging pathogen in swine
44 reproduction, its possible interaction with oocytes, spermatozoa and embryos has not
45 been deeply investigated.

46 In general, oocytes may become infected before ovulation by contact with infectious
47 agents present in either granulosa cells or follicular fluid and after fertilization during
48 the passage through the oviduct and the uterus (Bielanski et al. 2004). These Authors
49 showed that only a small proportion of cumulus–oocyte complexes (COCs) recovered
50 from PCV2 seropositive gilts is positive for viral DNA. They suggested that COCs may
51 have the potential for the production of PCV2-contaminated embryos when follicular
52 fluid containing COCs from infected animals is pooled for *in vitro* fertilization (IVF) as
53 already observed for bovine herpesvirus-1 (BHV-1) and bovine viral diarrhoea virus
54 (BVDV) during *in vitro* production of bovine embryos (Bielanski et al. 1993).

55 In contrast, all *in vivo* fertilized, uterine stage, embryos (developmental range: from four
56 cells to hatched blastocysts) from infected sows tested were negative for PCV2 DNA
57 (Bielanski et al. 2004). These authors suggested that this could be due to the fact that,

58 despite the excretion of some pathogenic agents into the reproductive tract, uterine stage
59 embryos may not be associated with infectious agents and/or they may be rendered free
60 from infectious agents by a simple sequential washing procedure (Stringfellow 1998).
61 Recently PCV2 nucleic acid has been detected in semen samples from healthy and
62 experimentally infected boars but information regarding the fraction in which PCV2
63 occurs in semen are conflicting. Pal et al. (2008) showed PCV2 DNA to be present
64 mainly in the cell fraction as opposed to the seminal plasma fraction, whereas Kim et al.
65 (2001) found that PCV2 DNA is mainly present in seminal fluid and nonsperm cell
66 fractions and in only few percent of sperm head.

67 Due to a recent increase in the use of pig as a model in modern biotechnology
68 applications such as *in vitro* fertilization, *in vitro* embryo production and sperm sexing
69 technology, the potential for infectious disease transmission by pig gametes is a topic of
70 considerable interest.

71 To the best of our knowledge, no studies have been performed to date on the possibility
72 of inducing PCV2 infection through infected sperm cell.

73 Basing on above mentioned observations, this study was set up to evaluate the ability of
74 PCV2 experimentally infected semen to produce blastocysts PCR positive for PCV2
75 from IVM (*in vitro maturation*) - IVF (*in vitro fertilization*) oocytes.

76

77 **MATERIALS AND METHODS**

78 Unless otherwise specified, all the reagents were purchased from Sigma–Aldrich
79 (Milan, Italy).

80

81 **Virus**

82 The virus used was a PCV2b strain isolated from an outbreak of PMWS in Italy and
83 genotyped as reported by Hesse et al. (2008). It was propagated in circovirus-free PK15
84 cells and its identity was confirmed by PCR analysis and reactivity with specific PCV2
85 monoclonal antibodies. The viral suspension titre was determined by growing serial
86 dilutions in circovirus-free PK15 cell monolayers, followed by immunofluorescent
87 labelling for viral antigen.

88

89 **Semen processing and exposure of boar spermatozoa to PCV2b**

90 The boar used to collect semen had a proven fertility, was housed since five
91 years in our facility (without contact with other pigs) for experimental purposes
92 and was serologically negative to PCV2. Sperm-rich fraction of ejaculate was
93 collected by gloved hand technique and extended in equal volumes of AndrohepTM
94 (Minitub, Tiefenbach, Germany). The semen resulted negative by PCR to PCV2 DNA
95 (Sarli et al. 2012).

96 Sperm suspension containing 3×10^9 spermatozoa/100 mL of Androhep was incubated
97 for 4 h at 16°C with 10 mL of a PCV2b viral suspension containing $10^{3.9}$ TCID₅₀/mL.

98

99 **Semen evaluation**

100 To evaluate a potential negative effect of viral suspension on sperm cells, viability of
101 spermatozoa was analyzed before and after the above described incubation with viral
102 suspension. Briefly, 25 µL of semen were incubated with 2 µL of a 300 µM Propidium
103 Iodide (PI) stock solution and 2 µL of a 10 µM SYBR-14 stock solution (both obtained
104 from the Live/dead sperm viability kit; Invitrogen, Eugene, OR, USA), for 5 min at
105 37°C in the dark. After incubation, 10 µL of sperm suspensions were analyzed with a

106 Nikon Eclipse epifluorescence microscope using a double-band-pass filter for green and
107 red fluorescence. The spermatozoa with green or red fluorescence on the head were
108 considered live or dead, respectively. At least 200 spermatozoa were scored for each
109 sample.

110

111 ***In vitro* maturation (IVM) of cumulus-oocyte-complexes**

112 Ovaries were collected at a local abattoir and transported to the lab within 2 h in a
113 thermos filled with physiological saline at 30-35°C. Cumulus oocyte complexes (COCs)
114 from follicles 3-6 mm in diameter were aspirated using a 18 gauge needle attached to a
115 10 mL disposable syringe. Under a stereomicroscope, intact COCs were selected and
116 transferred into a petri dish (35 mm, Nunclon, Roskilde, Denmark) prefilled with 2 mL
117 of modified PBS supplemented with 0.4% BSA. Only COCs with complete and dense
118 cumulus oophorus were used. After three washes in NCSU 37 (Petters & Wells 1993)
119 supplemented with 5.0 mg/mL insulin, 0.57 mM cysteine, 10 ng/mL epidermal growth
120 factor, 50 µM β-mercaptoethanol and 10% PCV2-PCR-negative porcine follicular fluid
121 (IVM medium), groups of 50 COCs were transferred to a Nunc 4-well multidish
122 containing 500 µL of the same medium per well and cultured at 39°C in humidified
123 atmosphere of 5% CO₂/7% O₂ in air. For the first 22 h of *in vitro* maturation, the IVM
124 medium was supplemented with 1.0 mM dibutyryl cyclic adenosine monophosphate
125 (db-cAMP), 10 IU/mL eCG (Folligon, Intervet, Boxmeer, The Netherlands) and 10
126 IU/mL hCG (Corulon, Intervet). For the last 20 h COCs were transferred to fresh
127 maturation medium (Funahashi et al. 1997). Part of the oocytes was washed in PBS,
128 treated with pronase, in order to remove zona pellucida, rinsed twice in nuclease free
129 water, frozen in 12 pools of 50 and stored at -80°C until PCR analysis.

130

131 Collection of immature oocytes, cumulus cells and follicular fluid

132 COCs and follicular fluids from 80-90 follicles, 3-6 mm in diameter, were aspirated and
133 placed in a petri dish. COCs were then transferred to another petri dish containing
134 500 μ L of modified PBS supplemented with 0.4% BSA; the remainig follicular fluids
135 were centrifuged and the supernatants were stored at -80°C until PCR analysis.

136 Oocytes were mechanically denuded from cumulus cells, then treated with pronase and
137 frozen at -80°C. Cumulus cells were counted using a Thoma's hemocytometer then
138 centrifuged and the pellets stored at -80°C until PCR analysis.

139

140 *In vitro* fertilization (IVF) of cumulus-oocyte-complexes

141 Matured oocytes were fertilized with infected (infected group; INF) and non-infected
142 (control group; CTR) spermatozoa as follows. Sperm suspensions were washed twice in
143 Brackett&Olifant medium (Bracket & Olifant 1975) supplemented with 12% fetal calf
144 serum (FCS; Gibco, Invitrogen, Monza, Italy) and 0.7 mg/mL caffeine (IVF medium).
145 Sperm concentrations were evaluated and 50 matured oocytes, freed from cumulus cells
146 by gentle repeated pipetting, were transferred to 500 μ l IVF medium containing 1×10^6
147 sperm/mL. After 90 min of co-culture, oocytes were transferred to fresh IVF medium
148 and cultured for an additional 18 h period. Oocytes were then washed twice in NCSU-
149 23 and cultured in 500 μ L of the same medium. On day 5 post fertilization, 250 μ L of
150 medium were replaced with fresh pre-equilibrated NCSU-23 containing 20% (v/v) FCS
151 to reach a final FCS concentration of 10% (v/v). At day 7 post-fertilization, embryos
152 were examined by stereomicroscope and blastocysts were treated with pronase, in order
153 to remove zona pellucida, rinsed twice in nuclease free water, individually frozen in 5

154 μL water and stored at -80°C until PCR analysis. Aliquots of infected spermatozoa used
155 for the IVF trials were washed twice with PBS and stored at -80°C until PCR analysis.

156

157 **Parthenogenetic activation**

158 After IVM, oocytes were denuded as described above, washed three times in IVF
159 medium and then parthenogenetically activated according to the method described by
160 Boquest et al. (2002) slightly modified. Briefly, the oocytes were transferred to IVF
161 medium containing 5 mM ionomycin for 5 min, then washed twice and incubated in
162 NCSU-23 containing 2 mM 6-dimethylaminopurine (6-DMAP) for 3 h at 39°C .
163 Presumptive parthenotes were washed twice in NCSU-23 and cultured in groups of 50
164 in 500 μL of the same medium. On day 5 postactivation, 250 μL of medium were
165 replaced with fresh pre-equilibrated NCSU-23 containing 20% (v/v) FCS to reach a
166 final FCS concentration of 10% (v/v). At day 7 post-activation, percent of blastocysts
167 was determined and treated as described above.

168

169 **Effect of sperm washing on virus presence in experimentally infected spermatozoa**

170 Sperm suspension containing 3×10^9 spermatozoa/100 mL of Androhep, was incubated
171 for 4 h at 16°C with 10 mL of a PCV2b viral suspension containing $10^{3.9}\text{TCID}_{50}/\text{mL}$.
172 Then 3 aliquots of 1 mL of sperm suspension (30×10^6 spermatozoa) were washed with
173 10 mL PBS 5 times. After that spermatozoa were stored at -80°C until PCR analysis.

174

175 **PCV2 detection in blastocysts, oocytes, cumulus cells and follicular fluid**

176 Blastocysts and pool of oocytes were lysed as previously described (Spinaci et al. 2007)
177 and 2 μL were used to perform the first round of PCR for PCV2. Cumulus cells (110000

178 to 880000 cells/pool) were submitted to DNA extraction with ZR Genomic DNA Tissue
179 Miniprep (Zymo Research, Corp. Irvine, CA, USA) according to manufacturer's
180 instruction. Follicular fluid pools were assayed as follows: 200 μ L samples underwent
181 DNA isolation by using a commercially available kit "NucleoSpin Tissue" (Macherey-
182 Nagel, Düren, Germany) in accordance to manufacturer's instruction. Twenty ng of
183 DNA from cumulus cells and follicular fluid were used for first PCR round for PCV2.
184 The detection of PCV2a or PCV2b was conducted by nested-polymerase chain reaction
185 (n-PCR). In the first round of PCR, the outer primers used were specific for PCV2. A
186 forward primer 5'-CAGTTCGTCACC-3' (designed using the Beacon Designer 2.07
187 Software, premier Biosoft International, Palo Alto, CA, USA) and reverse primers 5'-
188 CCGCACCTTCGGATATACTGTC -3' primer (Ouradani et al. 2000) were used.
189 Amplification of DNA was achieved by 35 cycles of 95°C for 1 min, 55°C for 1 min
190 and 72°C for 1 min.

191 In the second PCR run (n-PCR), a common forward primer, 5'-
192 CACGGATATTGTAGTCCTGGT-3' (Ouradani et al. 2000) and different reverse
193 primers sequences, were used to amplify either PCV2a 5'-
194 GGGGGACCAACAAAATCTC-3' or PCV2b (5'-GGGGCTCAAACCCCGCTC-3')
195 (Hesse et al. 2008). Two μ L of first PCR were used to performed the n-PCR.
196 Amplification of DNA was achieved by 35 cycles of 95°C for 1 min, 62°C for 1 min
197 and 72°C for 1 min. The reaction mixtures contained 3 mM MgCl₂; 200mM of each
198 dNTP, 0.2 μ M of each primer; 0.5U of Go Taq Polymerase (Promega Corp, Madison,
199 WI, USA). All reactions were performed in duplicate. PCR products were analyzed on
200 2% agarose gel stained with ethidium bromide under UV light. The expected PCR
201 products of the first and n-PCR were 627 bp or 391 bp respectively. The specificity of

202 the primers used was demonstrated by PCR amplification of PCV2a and PCV2b control
203 viruses ($10^{3.3}$ TCID₅₀/mL and $10^{3.9}$ TCID₅₀/mL for PCV2a and PCV2b respectively).

204 The sensitivity of the PCR was estimated through serial dilutions from 1×10^9 to 10^5
205 DNA copies of reference positive control (PCV2a and PCV2b). The PCR (first round)
206 had sensitivity of 1×10^5 whereas the n-PCR was 10 times more sensitive (1×10^4).

207 Amplification of β -Actin gene was performed for all sample as internal control as
208 previously described (Forni et al. 2003).

209

210 **PCV2 detection in sperm samples**

211 Aliquots of sperms (6×10^6 spermatozoa) were resuspended with denaturing buffer (20
212 mM TrisHCl; EDTA 20mM; DTT 80mM; SDS 4%; NaCl 200mM; proteinase K 20
213 mg/mL) and incubated for 3h at 55°C then submitted to genomic DNA extraction by
214 using ZR Genomic DNA (Zymo Research, Corp. Irvine, CA, USA) according to
215 manufacturer's instruction. The first round of PCR was performed with 50 ng of
216 extracted DNA and PCR product was analyzed on 2% agarose gel stained with ethidium
217 bromide under UV light.

218

219 **Experimental design**

220 Four experiments were performed:

221 Experiment 1: evaluation of PCV2 presence in blastocysts obtained by IVF with *in vitro*
222 infected or control semen;

223 Experiment 2: evaluation of PCV2 presence in blastocysts obtained by parthenogenesis;

224 Experiment 3: evaluation of PCV2 presence in *in vitro* matured oocytes and immature
225 oocytes, cumulus cells and follicular fluid;

226 Experiment 4: evaluation of PCV2 presence in *in vitro* infected spermatozoa after
227 increasing washing steps.

228

229 **Statistical analysis**

230 Data on the number of blastocyst obtained were pooled from two replicates and
231 analyzed using SPSS 11 (SPSS Inc., Chicago, IL, USA) with chi-squared test; the level
232 of significance was set at $P < 0.05$.

233

234 **RESULTS**

235

236 **Sperm viability**

237 Sperm viability was not affected by the treatment with viral suspension: it was $82.97 \pm$
238 3.24% before incubation and $81.44 \pm 2.8\%$ (mean \pm SE) after incubation.

239

240 **Experiment 1: evaluation of PCV2 presence in blastocysts obtained by *in vitro*** 241 **infected semen**

242 Sperm cells exposed for 4 h to PCV2b were PCR positive. Exposure of spermatozoa to
243 PCV2b did not exert any effect on the percentage of embryos that developed to
244 blastocyst stage (Table 1). In CTR group, 4 out of 15 blastocysts tested were PCV2a
245 positive. In INF group, 1 out of 15 blastocysts screened were PCV2b positive and 2
246 PCV2a positive (Table 2, Fig. 1).

247

248 **Experiment 2: evaluation of PCV2 presence in blastocysts obtained by** 249 **parthenogenesis**

250 At day 7 the percent of blastocysts parthenogenetically activated was 23.92%.

251 One out of 17 blastocysts parthenogenetically activated was PCV2b positive and one
252 PCV2a positive.

253

254 **Experiment 3: evaluation of PCV2 presence in *in vitro* matured oocytes and**
255 **immature oocytes, cumulus cells and follicular fluid**

256 None of the twelve pools of fifty *in vitro* matured oocytes was positive to PCV2b or
257 PCV2a. On the basis of these results PCV2b or PCV2a positivity was assayed in six
258 more consistent pools of immature oocytes, cumulus cells and follicular fluid (from 80-
259 90 follicles). The results obtained are presented in Table 3 and Figure 2.

260

261 **Experiment 4: evaluation of PCV2 presence in *in vitro* infected spermatozoa after**
262 **increasing washing steps**

263

264 As shown in Figure 3 a sequential washing procedure was not effective in removing
265 virus from spermatozoa.

266

267 **DISCUSSION**

268 The purpose of the present work was to evaluate the ability of PCV2 experimentally
269 spiked semen to produce blastocysts PCR positive for PCV2. Embryos were obtained
270 from *in vitro* maturation and fertilization of porcine oocytes from ovaries collected at a
271 local abattoir.

272 To our knowledge, the present study is the first one in which spermatozoa have been
273 contaminated with PCV2 and utilized for IVF under experimental conditions.

274 Our results clearly demonstrate that PCV2 firmly attaches to the surface of spermatozoa
275 (as demonstrated in experiment 4) but they don't seem to be effective in infecting the
276 oocytes during IVF and in producing infected embryos.

277 The infected blastocysts we obtained derived most probably from infected oocytes
278 obtained at abattoir; in fact, 6 out of 30 blastocysts tested were PCV2a positive.
279 Moreover parthenogenetic blastocysts PCV2b or PCV2a positive were obtained. Similar
280 contamination has been observed also by Bureau et al. (2005).

281 In order to confirm this hypothesis, we evaluated the presence of PCV2a or PCV2b in
282 different pools of fifty *in vitro* matured oocytes but none of the pools was positive.
283 When more consistent oocyte pools were assayed, few of them resulted positive for
284 PCV2a or PCV2b. Moreover, PCV2 positivity was also recorded in follicular fluids and
285 cumulus cells from pool of follicles.

286 These results confirm that oocyte from ovaries collected at the slaughterhouse can be
287 infected and therefore could generate infected embryos. Why experimentally spiked
288 semen was not effective in producing infected embryos in our study is not clear.

289 A hypothesis could be that the virus bound to sperm cells could be entrapped, during the
290 zona pellucida crossing, in the ZP network of filaments with pores/holes and meshes or
291 to the presence of carbohydrates and glycoproteins that can function as receptors for
292 viruses. The PCV2 capsid is constituted by capsomeres, proteins with a high affinity for
293 glycosilated proteins (Misinzo et al. 2006). Since porcine ZP is composed of
294 glycoproteins (pZPA, pZPB, pZPC) (Harris et al. 1994), an interaction between the viral
295 capsid and these glycoproteins might be responsible for the sticky behaviour of the
296 porcine ZP towards PCV2.

297 It is also important to note that Bureau et al. (2005) reported that hyaluronidase is able
298 to remove PCV2 from contaminated embryos. In fact, those authors demonstrated that
299 when trypsin was used as proteolytic enzyme in washing buffers, more virions remained
300 bound to the zona pellucida than when hyaluronidase was used.

301 It's well known that sperm head surface is involved in acrosome reaction which results
302 in the release of digestive enzymes (including hyaluronidase) required for zona
303 penetration (Yanagimachi et al. 1994; Flesch & Gadella 2000). Therefore we can
304 hypothesize that, during the fertilization process, membrane capacitation modifications
305 first and acrosome reaction subsequently, with hyaluronidase and other enzymes release,
306 could have induced, at least in part, the detach of PCV2 from sperm head plasma
307 membrane. More recently Blomqvist et al. (2011) showed that Single Layer
308 Centrifugation (SLC) on Androcoll-P, followed by swim-up, is effective in removing
309 PCV2 virus from boar semen. Moreover, Bucci et al. (2013) demonstrated that SLC
310 using Androcoll-P induces some capacitation-related changes in boar sperm membrane,
311 possibly as a consequence of the complete removal of seminal plasma and the partial
312 removal of cholesterol from sperm membrane due to Androcoll centrifugation.

313 Basing on the above mentioned observations, we tested the grade of spermatozoa
314 infection after 2 h at 39°C in capacitation medium and acrosome reaction induction by
315 adding A23187 as previously described by Bucci et al. (2012). Spermatozoa subjected
316 to *in vitro* induction of capacitation and acrosome reaction showed a reduction of PCV2
317 amount (unpublished data). These results could explain, at least in part, the inability of
318 sperm cell to produce infected blastocysts.

319 In conclusion, the present study shows that PCV2 attaches to the surface of spermatozoa
320 but it seems not to have the ability to carry the virus into the oocyte at fertilization and
321 to produce infected embryos.

322

323 **Conflict of interest statement**

324 None of the authors of this manuscript have any conflict of interest to declare.

325

326 **Acknowledgements**

327 The work was supported by a Bologna University grant (Ricerca Fondamentale
328 Orientata).

329 We thank P. Cordioli (Istituto Zooprofilattico Sperimentale della Lombardia e
330 dell'Emilia- Romagna) for providing PCV2 isolates.

331

332 **REFERENCES**

333

334 Bielanski A, Laroche R, Algire J, Magar R. 2004. Distribution of PCV-2 DNA in the
335 reproductive tract, oocytes and embryos of PCV-2 antibody-positive pigs. *Veterinary*
336 *Record* **155**, 597–598.

337

338 Bielanski A, Loewen KS, Del Campo RM, Sirard MA, Willadsen S. 1993. Isolation of
339 bovine herpesvirus-1 (BHV-1) and bovine viral diarrhoea virus (BVDV) in association
340 with the *in vitro* production of bovine embryos. *Theriogenology* **40**, 531-538.

341

342

- 343 Blomqvist G, Persson M, Wallgren M, Wallgren P, Morrell JM. 2011. Removal of virus
344 from boar semen spiked with porcine circovirus type 2. *Animal Reproduction Science*
345 **126**, 108–114.
- 346
- 347 Boquest AC, Grupen CG, Harrison SJ, 2002. Production of cloned pigs from cultured
348 fetal fibroblast cells. *Biology of Reproduction* **66**, 1283–1287.
- 349
- 350 Brackett BG, Oliphant G. 1975. Capacitation of rabbit spermatozoa *in vitro*. *Biology of*
351 *Reproduction* **12**, 260–274.
- 352
- 353 Bucci D, Galeati G, Tamanini C, Vallorani C, Rodriguez-Gil JE, Spinaci M. 2012.
354 Effect of sex sorting on CTC staining, actin cytoskeleton and tyrosine phosphorylation
355 in bull and boar spermatozoa. *Theriogenology* **77**, 1206-1216.
- 356
- 357 Bucci D, Spinaci M, Morrell J, Vallorani C, Tamanini C, Guidetti R, Galeati G. 2013.
358 Effects of single layer centrifugation with Androcoll-P on boar sperm. *Animal*
359 *Reproduction Science* **138**, 276-281.
- 360
- 361 Bureau M, Dea S, Sirard MA. 2005. Evaluation of virus decontamination techniques for
362 porcine embryos produced *in vitro*. *Theriogenology* **63**, 2343-2355.
- 363
- 364 Chae C. 2005. A review of porcine circovirus 2-associated syndromes and diseases.
365 *Veterinary Journal* **169**, 326–336.
- 366

- 367 Flesch FM, Gadella BM. 2000. Dynamics of the mammalian sperm plasma membrane
368 in the process of fertilization. *Biochimica et Biophysica Acta* **1469**, 197–235.
- 369
- 370 Forni M, Zannoni A, Tamanini C, Bacci ML. 2003. Opposite regulation of clusterin and
371 LH receptor in the swine corpus luteum during luteolysis. *Reproduction Nutrition and*
372 *Development* **43**,517–25.
- 373
- 374 Funahashi H, Cantley TC, Day BN. 1997. Synchronization of meiosis in porcine oocyte
375 by exposure to dibutyryl cyclic adenosine monophosphate improves developmental
376 competence following *in vitro* fertilization. *Biology of Reproduction* **57**, 49-53.
- 377
- 378 Harris JD, Hibler DW, Fontenot GK, Hsu KT, Yurewicz EC, Sacco AG. 1994. Cloning
379 and characterization of zona pellucida genes and cDNAs from a variety of mammalian
380 species: the ZPA, ZPB and ZPC gene families. *DNA Sequence* **4**, 361-393.
- 381
- 382 Hesse R, Kerrigan M, Rowland RR. 2008. Evidence for recombination between PCV2a
383 and PCV2b in the field. *Virus Research* **132**, 201–207.
- 384
- 385 Kim J, Han DU, Choi C, Chae C. 2001. Differentiation of porcine circovirus (PCV)-1
386 and PCV-2 in boar semen using a multiplex nested polymerase chain reaction. *Journal*
387 *of Virological Methods* **98**, 25–31.
- 388
- 389 Ladekjaer-Mikkelsen AS, Nielsen J, Storgaard T, Bøtner A, Allan G, McNeilly F. 2001.
390 Transplacental infection with PCV-2 associated with reproductive failure in a gilt.
391 *Veterinary Record* **148**, 759–760.

392

393 Misinzo G, Delputte PL, Meerts P, Lefebvre DJ, Nauwynck HJ. 2006. Porcine
394 circovirus 2 uses heparan sulfate and chondroitin sulfate B glycosaminoglycans as
395 receptors for its attachment to host cells. *Journal of Virology* **80**, 3487-3494.

396

397 O'Connor B, Gauvreau H, West K, Bogdan J, Ayroud M, Clark EG, Konoby C, Allan
398 G, Ellis JA. 2001. Multiple porcine circovirus 2-associated abortions and reproductive
399 failure in a multisite swine production unit. *Canadian Veterinary Journal* **42**, 551-553.

400

401 Ouardani M, Wilson L, Jetté R, Montpetit C, Dea S. 2000. Multiplex PCR for detection
402 and typing of porcine circoviruses. *Journal of Clinic Microbiology* **38**, 1707-1714.

403

404 Pal N, Huang YW, Madson DM, Kuster C, Meng XJ, Halbur PG, Opriessing T. 2008.
405 Development and validation of a duplex realtime PCR assay for the simultaneous
406 detection and quantification of porcine circovirus type 2 and an internal control on
407 porcine semen samples. *Journal of Virological Methods* **149**, 217-225.

408

409 Petters RM, Wells KD. 1993. Culture of pig embryos. *Journal Reproduction Fertility* **48**,
410 61-73.

411

412 Sanchez RE, Nauwynck HJ, McNeilly F, Allan GM, Pensaert MB. 2001. Porcine
413 circovirus 2 infection in swine fetuses inoculated at different stages of gestation.
414 *Veterinary Microbiology* **83**, 169-176.

415

- 416 Sarli G, Morandi F, Panarese S, Bacci B, Ferrara D, Bianco C, Fusaro L, Bacci ML,
417 Galeati G, Dottori M, Bonilauri P, Lelli D, Leotti G, Vila T, Joisel F, Allan G, Benazzi
418 C, Ostanello F. 2012. Reproduction in porcine circovirus type 2 (PCV2) seropositive
419 gilts inseminated with PCV2b spiked semen. *Acta Veterinaria Scandinavica* **54**, 51.
420
- 421 Segalés J, Allan GM, Domingo M. 2005. Porcine circovirus diseases. *Animal Health*
422 *Research Reviews* **6**, 119–142.
423
- 424 Segalés J, Rosell C, Domingo M. 2004. Pathological findings associated with naturally
425 acquired porcine circovirus type 2 associated disease. *Veterinary Microbiology* **98**, 137–
426 149.
427
- 428 Spinaci M, Merlo B, Zannoni A, Iacono E, De Ambrogi M, Turba ME, Zambelli D.
429 2007. In vitro production of cat blastocysts of predetermined sex using flow
430 cytometrically sorted semen. *Theriogenology* **67**, 872-877.
431
- 432 Stringfellow DA. 1998. Recommendations for the sanitary handling of in-vivo-derived
433 embryos. In: Stringfellow DA, Seidel SM (eds), *Manual of the International Embryo*
434 *Transfer Society*, pp. 79–84. Int. Embryo Transfer Soc. Savoy, IL.
435
- 436 Yanagimachi R, Knobil E, Neill JD. 1994. Mammalian Fertilization. In: Knobil E, Neill
437 JD (eds), *The Physiology of Reproduction* (2nd edn), pp. 189–317. Raven Press Ltd,
438 New York.
439

440 **Table 1** Blastocysts obtained by IVF with infected (INF) or non-infected spermatozoa
441 (CTR).

442

443

Groups	No. oocytes	No. blastocysts (%)
CTR	250	36 (14.4)
INF	250	33 (13.2)

444

445 **Table 2** Evaluation of PCV2a or PCV2b presence in blastocysts obtained by IVF.

446

Groups	No. blastocysts examined	No. blastocysts positive	
		PCV2a	PCV2b
CTR	15	4	0
INF	15	2	1

447

448

449 **Table 3** PCV2a and PCV2b positivity in the six pools of immature oocytes, cumulus
450 cells and follicular fluid.

451

Pool n°	PCV2a positivity			PCV2b positivity		
	Oocytes	Cumulus cells	Follicular fluids	Oocytes	Cumulus cells	Follicular fluids
1	-	-	-	-	-	+
2	-	-	-	-	-	-
3	-	-	+	-	-	-
4	-	+	-	+	+	-
5	+	-	-	-	+	+
6	-	-	-	-	-	+

452

453

454

455 **Figures captions**

456

457 **Figure 1** Representative image of agarose gels of nested PCR for PCV2a and PCV2b
458 products (**A, B**) and β -actin (**C**) of blastocysts. **A:** CTR blastocysts; **B:** blastocysts
459 obtained after IVF with PCV2b experimentally spiked semen. Lanes 1-5; 10-14:
460 blastocysts; Lane 7: PCV2a positive control; Lane 8: PCV2b positive control; Lane 6:
461 negative control of PCR; MWM: DNA molecular weight marker.

462


463 **Figure 2** Representative image of agarose gels of nested PCR for PCV2a and PCV2b
464 products of different pools (n=6; Lanes 1-6) of: oocytes (**A**); cumulus cells (**B**); follicular
465 fluid (**C**). Lane 7: DNA molecular weight marker. Lane 8: PCV2a and PCV2b positive
466 control.

467

468

469 **Figure 3** Representative image of agarose gel of PCV2 PCR of infected spermatozoa
470 (Lane 4) and washed infected spermatozoa (Lanes 1-3). Lane 5: negative control of
471 PCR; Lane 6: PCV2b positive control; MWM: molecular weight marker.


472



473

474 **Figure 1**


475



476

477 **Figure 2**

478



480 **Figure 3**

481