



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Growth traits, carcass and meat quality of heavy pigs receiving different enrichment tools

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

*Availability:*

This version is available at: <https://hdl.handle.net/11585/601507> since: 2017-06-20

*Published:*

DOI: <http://doi.org/>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)



**Official Journal of  
the Animal Science  
and Production  
Association (ASPA)**

\*

**ISSN 1594-4077**

**eISSN 1828-051X**

\*

**[http://www.  
tandfonline.com/tjas](http://www.tandfonline.com/tjas)**

\*

**Italian Journal of  
Animal Science**

\*

**volume 16**

**supplement 1**

**2017**

# **italian journal of animal science**

**ASPA 22<sup>nd</sup> CONGRESS**

**Perugia, June 13-16, 2017**

**Book of Abstracts**

**Guest Editors: Massimo Trabalza-Marinucci (Coordinator),  
Cesare Castellini, Emiliano Lasagna, Stefano Capomaccio,  
Katia Cappelli, Simone Ceccobelli, Andrea Giontella**

# Italian Journal of Animal Science

Official Journal of the Animal Science  
and Production Association

ISSN 1594-4077

eISSN 1828-051X

## Editor-in-Chief

Rosanna Scipioni, Department of Life Sciences, University of Modena and Reggio Emilia (Italy)

## Deputy Editor

Marcello Mele, University of Pisa (Italy)

## Associate Editors

Umberto Bernabucci, University of Tuscia, Viterbo (Italy)

Oreste Franci, University of Firenze (Italy)

Nicolò Pietro Paolo Macciotta, University of Sassari (Italy)

Antonino Nizza, University of Napoli (Italy)

Giuliana Parisi, University of Firenze (Italy)

Giovanni Savoini, University of Milano (Italy)

## Section Editors

Alessandro Agazzi, University of Milano (Italy)

Susana Alves, Technical University of Lisboa (Portugal)

Paolo Bani, Università Cattolica del S. Cuore, Piacenza (Italy)

Luca M. Battaglini, University of Torino (Italy)

Fulvia Bovera, University of Napoli (Italy)

Rui José Branquinho Bessa, Technical University of Lisboa (Portugal)

Luigi Calamari, Università Cattolica del S. Cuore, Piacenza (Italy)

Cesare Castellini, University of Perugia (Italy)

Alessio Cecchinato, University of Padova (Italy)

Beniamino T. Cenci Goga, University of Perugia (Italy)

Giulio Cozzi, University of Padova (Italy)

Juan Vicente Delgado Bermejo, University of Córdoba (Spain)

Luca Fontanesi, University of Bologna (Italy)

Flaviana Gottardo, University of Padova (Italy)

Michael A. Grashorn, University of Hohenheim (Germany)

Juan J. Lóor, University of Illinois at Urbana-Champaign (USA)

Anna Nudda, University of Sassari (Italy)

Manuela Renna, University of Torino (Italy)

Evangelia Sossidou, Veterinary Research Institute, Thessaloniki (Greece)

Giuseppe Stradaoli, University of Udine (Italy)

Eleni Tsiplakou, Agricultural University of Athens (Greece)

Hai-Jun Zhang, Chinese Academy of Agricultural Sciences, Beijing (China)



**Taylor & Francis**  
Taylor & Francis Group

Feed restriction (R *vs* AL) reduced the daily FI (7%,  $p < .001$ ) and the number of visits (27%,  $p < .001$ ), but increased FI per visit (20%,  $p = .001$ ) and the eating rate (10%,  $p = .032$ ).

The dietary amino acid reduction applied in late finishing (LAA *vs* CAA) increased the daily FI (10.5%,  $p = .037$ ), tended to increase the eating rate (19.6%,  $p = .06$ ) with both feeding regimes, and tended to interact ( $p < .10$ ) with the feeding regime for other eating traits. In fact, under AL conditions, dietary amino acid reduction increased the daily FI and the visit frequency, but reduced the FI per visit, where the opposite occurred under R conditions.

The coefficient of variation of all the behaviour traits (ranging 0.18 to 0.66) was always larger than that found for the daily FI (0.10). Strong relationships between FI per visit and the visit frequency ( $R^2 = 0.93$ ), and between the eating rate and the time spent eating ( $R^2 = 0.80$ ) were found. The behavioural flexibility of the pig in the attempt to achieve their desired feed intake was evidenced.

#### Acknowledgements

The research was funded by the University of Padua under grant number CPDR143385/14 and DOR1655405/16. The authors would like to thank Veronesi SPA (Quinto di Valpantena, Verona, Italy) for financial and technical support, and Aldo Gini Foundation (Via Portello, 15, Padova, Italy) for supporting the research activities of Msc Giuseppe Carcò, during his stage at the University of Zaragoza.

## O102

### Growth traits, carcass and meat quality of heavy pigs receiving different enrichment tools

Eleonora Nannoni, Luca Sardi, Marika Vitali, Giovanna Martelli

Dipartimento di Scienze Mediche Veterinarie, Alma Mater Studiorum University of Bologna, Italy  
Contact: [eleonora.nannoni2@unibo.it](mailto:eleonora.nannoni2@unibo.it)

Two separate and independent trials were carried out to investigate the effect of innovative enrichment devices on Italian heavy pigs' growth parameters, carcass traits, meat and ham quality. In Trial 1, 20 pigs received a hanging metal chain (C1) as environmental enrichment, and 20 received wood (poplar) logs (WL) placed inside a metal rack installed on one side of the pen. In trial 2, 20 pigs received a metal chain (C2), whereas 20 animals received a specifically formulated edible block (EB) placed inside the same metal rack described above. Enrichments were always available to pigs. Animals were kept on slatted floors and liquid-fed twice a day. Trials started when animals were approximately 80 days old

(average Body Weight – BW = 28 kg) and ended at slaughtering (average BW = 158 kg, age: 285 days in Trial 1 and 280 days in Trial 2). Growth parameters, carcass traits and meat quality data were collected, (including hams weight during dry-curing) and submitted to analysis of variance. The pen (5 pigs) was taken as the experimental unit for growth parameters, the individual was used as the experimental unit for carcass and meat traits.

In both trials, no significant differences were observed as concerns growth parameters (final BW, average daily gain, feed consumption, feed conversion ratio). Only minor differences were detected in carcass traits, with WL carcasses having higher lean meat percentage (51.06 *vs* 49.86%,  $p < .05$ ), lower backfat thickness (22.55 *vs* 25.15 mm,  $p < .05$ ) but higher drip losses (1.23 *vs* 1.00%,  $p < .05$ ) if compared to C1. EB carcasses had lower loin thickness if compared to C2 (59.20 *vs* 64.50 mm,  $p < .05$ ). In both trials, these slight differences did not affect the overall yield in lean cuts, the overall meat water holding capacity or any other quality trait of meat (pH, colour, Warner-Bratzler shearing force). Ham weight losses during dry-curing were not affected by the experimental treatment ( $p > .05$ ).

Our results show that providing heavy pigs with either poplar logs or an edible block did not affect their growth parameters, overall carcass traits, meat or ham quality; this observation supports the conclusion that such innovative and destructible/ingestible enrichment tools could be used in substitution of the metal chain. Further useful information will be available once the observation of the behavioural video-recordings will be completed, to get more insights on the type of interactions carried out by pigs on each enrichment device.

#### Acknowledgements

The research was funded by Progetto AGER, grant n° 2011-0280.

## O103

### Seed-based vaccine immunogens administered in feed for the control of verocytotoxic *Escherichia coli* infection in pig livestock

Luciana Rossi<sup>1</sup>, Serena Reggi<sup>2</sup>, Giovanni L. Alborali<sup>3</sup>, Paolo Trevisi<sup>4</sup>, Diana Luise<sup>4</sup>, Angela Lombardi<sup>1</sup>, Eugenio Demartini<sup>1</sup>, Antonella Baldi<sup>1</sup>

<sup>1</sup>Dipartimento di Scienze Veterinarie per la Salute, la Produzione Animale e la Sicurezza Alimentare, University of Milano, Italy

<sup>2</sup>Plant techno s.r.l, Vicomosciano, Italy