

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Relative abundance of heat shock proteins and clusterin transcripts in spermatozoa collected form boar routinely utilised in an artificial insemination centre: preliminary results

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Zannoni, A., Bernardini, C., Zaniboni, A., Ferlizza, E., Ventrella, D., Bacci, M.L., et al. (2017). Relative abundance of heat shock proteins and clusterin transcripts in spermatozoa collected form boar routinely utilised in an artificial insemination centre: preliminary results. VETERINARY RESEARCH COMMUNICATIONS, 41(3), 233-239 [10.1007/s11259-017-9689-6].

Availability:

This version is available at: https://hdl.handle.net/11585/607061 since: 2021-11-29

Published:

DOI: http://doi.org/10.1007/s11259-017-9689-6

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Relative abundance of heat shock proteins and clusterin transcripts in spermatozoa collected from boar routinely utilised in an artificial insemination centre: preliminary results

A Zannoni, C Bernardini, A Zaniboni, E Ferlizza, D Ventrella, M L Bacci, M Forni

Vet Res Commun 2017 Sep;41(3):233-239.

doi: 10.1007/s11259-017-9689-6. Epub 2017 Apr 21.

The final published version is available online at: <u>10.1007/s11259-017-9689-6</u>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

1	
2	Relative abundance of heat shock proteins and clusterin transcripts in spermatozoa collected from boar
3	routinely utilised in an artificial insemination centre: preliminary results
4	
5	A. Zannoni *, C. Bernardini, A. Zaniboni, E. Ferlizza, D. Ventrella, M. L. Bacci, M. Forni
6	
7	Department of Veterinary Medical Sciences – DIMEVET- University of Bologna, Via Tolara di Sopra 50,
8	Ozzano dell'Emilia, 40064 Bologna, Italy
9	
10	
11	
12	
13	
14	Corresponding author A. Zannoni
15	Tel.: +39.051.2097911; fax +39.051.2097899; e-mail: augusta.zannoni@unibo.it (A. Zannoni)
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	

32 Abstract

It is widely accepted that mature sperm contains RNA. The first hypothesis was that sperm RNAs have no functions of their own but are simply residues of spermatogenesis reflecting the events that occurred during their formation in the testes. More recently new discoveries have essentially expanded these views, showing that sperm mRNAs constitute a population of stable full-length transcripts, many of which are selectively retained during spermatogenesis and delivered to oocytes contributing to early embryo development. It is well known that semen quality can be influenced by occasional physical stress, infection, and variation in temperature and the definition of new markers for evaluation of semen could offer knowledge about the fertility potential of a semen sample. The aim of the present study was to evaluate the presence and the relative quantity of transcripts and protein of heat shock protein 70 (HSP70), 90 (HSP90) and clusterin (CLU) in Percoll-selected spermatozoa collected from seven adult boars of proven fertility routinely employed for artificial insemination. Our results showed the presence of HSP70, HSP90 and CLU transcripts with different level of expression: high for HSPs and low for CLU transcripts. The transcript level of both HSPs are similar among selected spermatozoa derived from high quality sperm with the exception of one boar that showed a reduced content of HSP70 and HSP90 mRNA together with a lower semen quality. At protein level, both HSPs were detected with similar amount among all seven boars whilst no band was evidenced for CLU protein.

Keywords: boar; clusterin; HSP70; HSP90; sperm

- -

62

63

64 **1. Introduction**

It is widely accepted that mature sperm contains RNA. As male gametes have been considered 65 transcriptionally inert cells because of the substitution of histones with protamines, the first hypothesis 66 67 was that sperm RNAs have no functions of their own but are simply residues of spermatogenesis reflecting the events that occurred during their formation in the testes (Krawetz 2005). However, other 68 discoveries have essentially expanded these views, showing that sperm mRNAs constitute a population of 69 stable full-length transcripts, many of which are selectively retained during spermatogenesis (Ostermeier 70 71 et al. 2002; Hamatani 2012). Although the functions of sperm RNAs await full clarification, it has been 72 suggested that sperm transcriptional profiles provide a biomarker for male infertility and an increasing 73 number of studies in humans have demonstrated that the sperm mRNA profile may serve as a molecular 74 diagnostic marker for evaluating male fertility (Jodar et al. 2013).

75 Messenger RNAs have been found in the mature mammalian spermatozoa of several species (Dadoune et 76 al. 2005; Gilbert et al. 2007; Yang et al. 2009). In swine, transcriptome analysis of ejaculated spermatozoa has been performed, identifying many mRNAs involved in biological processes, molecular 77 78 functions and cellular components (Yang et al. 2009). More recently, the presence of various transcripts 79 has been investigated in order to describe mRNA involvement in capacitation and early embryo development in pigs (Hwang et al. 2013). It is well known that semen quality can be influenced by 80 81 occasional physical stress, infection, and variation in temperature (Althouse et al. 2000; Althouse and Lu 2005; Kunavongkrit et al. 2005; Knox et al. 2008; Althouse 2008) and the definition of new markers for 82 evaluation of semen could offer knowledge about the fertility potential of a semen sample. Semen quality 83 84 assessment represents a fundamental step for obtaining successful artificial insemination. In addition to 85 the conventional semen parameter evaluation, an increasing number of innovative molecular markers, at 86 proteomic and transcriptomic level, have been used to identify more reliable markers of sperm quality and fertility. In bulls, mRNA abundance of specific sperm proteins has been related to a fertility index, 87 88 allowing the determination of biomarkers able to predict the fertility of subjects (Feugang et al. 2010; 89 Kasimanickam et al. 2012).

In order to contribute to define new potential biomarkers of porcine semen quality, we selected three proteins whose presence is well documented in sperm cells, heat shock protein (HSP) 70, HSP90 and clusterin (CLU). The presence of HSP70 and HSP90 proteins in porcine spermatozoa has been described

93 as playing a role in sperm-oocyte interaction and sperm motility respectively (Huang et al. 2000b; Spinaci et al. 2005; Volpe et al. 2008). No correlation between HSP70 and semen quality was found (Turba et al. 94 95 2007) while HSP90 was investigated only in relation to the quality of cryopreserved semen (Wang et al. 2014; Zhang et al. 2015). Clusterin (CLU), a pleiotropic glycoprotein mainly present in seminal plasma 96 97 has also been found to be associated with spermatozoa in various species (Sylvester et al. 1991; Ibrahim et al. 1999; Ibrahim et al. 2001; Han et al. 2012). In human and bull sperm, clusterin mainly exists on the 98 99 surface of immature, low motile or morphologically abnormal spermatozoa and therefore its presence has been proposed as a marker of poor sperm quality (O'Bryan et al. 1994; Ibrahim et al. 2000; Muciaccia et 100 101 al. 2010).

Until now the presence and the levels of the transcripts of these important proteins (HSP70, HSP90 and
CLU) in pig sperm cells have not been clearly described. Only a single paper reported the presence of
CLU transcript, suggesting a role of the paternal mRNA in early embryo development (Kempisty et al.
2008).

106 Thus, the objectives of this study were to evaluate first the presence of HSP70, HSP90 and CLU 107 transcripts in ejaculated spermatozoa of seven boars of proven fertility, and to verify if the relative 108 abundance of transcripts and their related proteins.

109

110 **2. Materials and methods**

111

112 **2.1** Animal semen collection and quality evaluation parameters

Seven healthy Large White boars, (2-7-year-olds), housed under the same condition (temp. $20 \pm 2C$) on a commercial farm and routinely employed for artificial insemination (AI) were used in this study. The semen parameters were evaluated in order to confirm the quality of the ejaculated routinely. The average farrowing rate was 86% with a range of 84% to 89%.

- 117 Semen was collected every 3 to seven days, but only every fortnight was it used for analysis.
- 118 For this study, three ejaculated/boar (21 ejaculated in total) were collected and evaluated; all the samples
- 119 were collected from December 2013 to January 2014, under favourable environmental conditions.
- 120

121 **2.2** Semen collection and evaluation of semen quality

Sperm samples (n= 21) were collected by an experienced technician using the gloved-hand technique and a dummy. The sperm-rich fraction was collected in a pre-warmed thermos and was diluted 1:1 (v:v) with swine fertilization medium (SFM - home made extender as previously reported (Fantinati et al. 2009)) and immediately transported to the laboratory for evaluation of semen parameters.

126 The concentration of each ejaculate was assessed after spermatozoa immobilization in 1.5 M NaCl 127 solution under a phase-contrast microscope at 400x using a Thoma counting chamber. Three replicates 128 for each ejaculate were analysed.

The spermatozoa viability was evaluated by eosin-nigrosin staining. A working solution was prepared with two parts of 10% (w/v) eosin and one part of 5% (w/v) nigrosin. The sperm suspension and the working solution were mixed 1:1 (v:v) and the viability was evaluated. At least 200 cells were observed (Nikon Eclipse E600, Nikon Corporation, Japan). Three replicates for each ejaculate were analysed.

The sperm motility was subjectively assessed by visual estimation under a phase contrast microscope equipped with a heated plate. Three replicates for each ejaculate were analysed. A total of 10 fields were evaluated.

- 136
- 137

2.3 Percoll density gradient of swine spermatozoa

In order to collect highly motile and morphologically normal spermatozoa, without somatic cell contaminants or debris, the spermatozoa samples, previously diluted with SFM, were separated by centrifugation through a discontinuous Percoll (Sigma-Aldrich St Louis, MO, USA) density gradient (45/90, v/v; (Matàs et al. 2011)). A bilayer of 2 ml each of Percoll solution with different density (45 and 90 in the bottom) was prepared in 15 ml conic centrifuge tubes; then diluted semen samples (5- 6 x 10⁷ sperms/2ml/tube) were added at the top, care being taken to maintain the integrity of the layers.

Samples were then centrifuged for 30 min at 700g. The sperm pellet was collected from the 45/90interfaces, and then washed twice with DPBS (1ml). Sperm sample concentration, viability and motility was then evaluated, as described above. Aliquots of $5x10^6$ spermatozoa for each ejaculate were frozen and stored at -80°C until RNA and protein analysis.

- 148
- 149

2.4 RNA isolation and quantitative real-time PCR (qRT-PCR) for HSP70, HSP90 and CLU

150Total RNA from purified spermatozoa samples (5x106) was extracted using NucleoSpin RNA II151(Macherey Nagel, Düren, Germany) and triplicates for each ejaculate were analysed. The lysis of the cells152was carried out with Tissue Lyser TL (QIAGEN, Hilden, Germany) in the presence of two beads (50Hz

153 for 5 min). Total RNA was extracted according to the manufacturer's instructions, except for double treatment with DNase (instead of single DNase treatment as suggested). RNA was spectrophotometrically 154 155 quantified (DeNovix Inc, Wilmington, DE, USA) and its quality was determined by gel electrophoresis on 1% agarose. Then 1 µg of RNA was reverse-transcribed to cDNA using an iScript cDNA Synthesis 156 Kit (Bio-RAD Laboratories Inc., California, USA), in a final volume of 20 µl. Real time quantitative PCR 157 was performed using an iCycler Thermal Cycler (Bio-RAD). Primers for swine HSP70, HSP90 and CLU 158 were designed using Beacon Designer 2.07 software (Premier Biosoft International, Palo Alto, CA, 159 160 USA). Primers for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were used to normalize mRNA levels across different samples. Primer sequences are shown in Table 1. 161

A master mix of the following reaction components was prepared in the end-concentrations indicated: 0.5 μ l forward primer (0.2 μ M), 0.5 μ l reverse primer (0.2 μ M), 9.5 μ l water and 12.5 μ l IQ SYBR Green BioRad Supermix (Bio-RAD). Two μ l of cDNA were added to 23 μ l of the master mix. All samples were analysed in duplicate. The real-time PCR protocol employed was: initial denaturation for 3 min at 95°C, 40 cycles at 95°C for 15 sec and at 60°C for 30 sec, followed by a melting step with slow heating from 55°C to 95°C at a rate of 0.5C/sec. Primer specificity as well as real-time efficiency was evaluated in previously collected porcine testicular tissue (positive control).

169 Contamination of epithelial cells, germ cells and leucocyte (CDH1, KIT and CD4) in spermatozoa RNA 170 was preliminary verified using specific primers (table 1); the positive control used for the assay was 171 porcine testicular cDNA or cDNA synthetized from RNA purified from blood (Kumar et al., 2010).

172 The gene expression level of HSP70, HSP90 and CLU was evaluated as Δ Ct (threshold Cycle) = (Ct 173 reference gene – Ct gene of interest) which directly correlates with the expression level.

174

175 **2.5 Western blot for HSP70, HSP90 and CLU**

176 The spermatozoa samples $(5x10^6 / \text{sample})$ were resuspended in SDS buffer (Tris-HCl 62.5 mM pH 6.8; 177 SDS 2%, glycerol 20%) and duplicates for each ejaculate were analysed. Proteins from spermatozoa (1x10⁶ for HSP 70 and HSP90, 2x10⁶ for CLU) were separated by NuPage 10% Bis-Tris Gel (Gibco-178 Invitrogen, Paisley, UK) for 50 min at 200 V, and then electrophoretically transferred onto a 179 nitrocellulose membrane. HSP70, HSP90 detection was performed as previously reported (Volpe et al. 180 181 2008), while CLU western blotting was performed as previously reported (Zannoni et al. 2015). Briefly, the nitrocellulose membranes were blocked with 3% milk powder in PBS-T20 (phosphate buffer saline-182 183 0.1% tween-20) for one hour at room temperature. The membranes were then incubated overnight at 4°C

184	with a 1:1000 dilution of monoclonal anti-human HSP70 (C92F3A-5 Stressgen Biotechnologies Corp.
185	Victoria, BC, Canada), monoclonal anti-human HSP90 (AC88 Stressgen) and monoclonal anti-human
186	CLU (05-354 Upstate Millipore Corp. Temecula, CA, USA), in Tris Buffered Saline-T20 (TBS-T20
187	20mM Tris-HCl, pH 7.4, 500 mM NaCl, 0.1% T-20). After several washings with PBS-T20 the
188	membranes were incubated with the secondary biotin-conjugate antibody and then with a 1:1000 dilution
189	of an anti-biotin horseradish peroxidase (HRP)-linked antibody. All western blots were developed using
190	chemiluminescent substrate (Super Signal West Pico Chemiluminescent Substrate, Pierce Biotechnology,
191	Inc., Rockford, IL, USA) according to the manufacturer's instructions. All membranes were stripped and
192	reprobed for β -tubulin in order to normalize the results. The intensity of the resultant bands was acquired
193	using Fluor-S TM Multimager and quantified by Quantity One software (Bio-RAD). The relative protein
194	content was determined by the density of the resultant bands and expressed in arbitrary units (AU)
195	relative to the β -tubulin content, using the Quantity One software (Bio-RAD).
196	
197	2.6 Statistical analysis
198	Data were analysed by one-way analysis of variance (ANOVA) and post hoc Tukey HSD test. The
199	statistical significance was set at $P < 0.05$.
200	
201	3. Results
202	
203	3.1 Sperm parameters: viability, motility and concentration
204	Data on viability, motility and spermatozoa concentration are shown in Fig.1. All the ejaculates collected
205	from six out of the seven boars showed ejaculates of high sperm quality. One boar (B2) showed seminal
206	parameters significantly different from the others and below acceptability for use in artificial
207	insemination (AI) (Fig. 1). After Percoll separation the values of motility (92 \pm 3.7 %) and viability (93
208	± 2.4 %) were similar for all boars.
209	
210	3.2 Quantitative real-time PCR (qRT-PCR) for HSP70, HSP90 and CLU
211	Quantitative PCR experiments performed to verify the presence of contaminants mRNA showed that the
212	selected mRNA markers were not detectable.
213	Quantitative PCR data demonstrated that the HSP70 and HSP90 mRNAs were abundant in all samples
214	whilst CLU mRNA was very low. No statistically significant differences were observed in the gene

expression level of HSP70, HSP90 and CLU among boars, with the exception of one boar (B2) that 215

216 showed a reduced content of HSP70 and HSP90 mRNA (Fig.2).

217 3.3 Western blots for HSP70, HSP90 and CLU

- Western blot analysis for HSPs showed in all samples a specific band (Fig. 3B) as previously reported by 218 219 us (Volpe et al. 2008). No statistically significant differences were observed in the expression levels of 220 the protein among the seven boars (Fig. 3A).
- No signal was observed for CLU protein in sperm cell samples, but only in the positive control of aortic 221 222 porcine endothelial cells (data not shown).
- 223

224 4. Discussion

225 A growing interest is directed towards the presence of mRNA in mammalian ejaculated spermatozoa. Some findings suggested that these transcripts can code for proteins, which are essential in early embryo 226 development or for cellular function. Other studies reported that a semen quality evaluation could be 227 228 derived from RNA profiling which might provide clinical markers for male infertility (Jodar et al. 2013). 229 In swine species, over the past 10 years, transcriptome analysis has identified many mRNAs involved in different biological processes (Yang et al. 2009). More recently, the presence of various transcripts has 230 231 been investigated in order to clarify their involvement in early embryo development (Hwang et al. 2013).

The present study aimed to evaluate the expression of HSP70, HSP90 and CLU in spermatozoa of boar 232 233 routinely employed in AI.

Our study considered seven boars of different ages. One of these boars (B2), at the time of the study, 234 235 presented seminal parameters unsuitable for AI because of transient genitourinary infection but was 236 nonetheless included in the study to obtain Percoll-selected spermatozoa from low-quality semen.

237 Results showed the presence of HSP70, HSP90 and CLU transcripts. In the literature a CLU transcript has been reported (Kempisty et al. 2008) while only the presence of a transcript member of the HSP70 238 family (HSP70.2) has been described in porcine spermatozoa (Yang et al. 2009). The number of 239 transcripts differed, with HSP70 and HSP90 relatively highly abundant with respect to the chosen 240 housekeeping gene while CLU was very low or undetectable in some samples. 241

242 At the protein level, we detected similar amounts of HSP70 and 90 among all seven boars, but the CLU specific band was not detectable in any sample. A previous paper showed how an aggressive 243 manipulation such as cell sorting does not affect the quantity of the expressed protein in boar 244

spermatozoa (Spinaci et al.,2010), therefore we are confident that a much more mild procedure, such as
Percoll selection, will not influence the protein levels.

Even if our data may be considered preliminary, it's interesting to note that the HSPs transcript seem to be more informative than HSP protein in relation to the sperm quality. Indeed, six boars with high sperm quality showed high and similar level of HSPs transcripts while the boar (B2), with low sperm quality, showed a reduced level of HSPs transcripts. The existence of the correlation, if confirmed in future studies, allows us to speculate that, even if selected and able to fertilize, sperm cells deriving from a poorquality semen sample have characteristics that make them unfit to sustain embryonic development.

Previous findings about the absence or low correlation between HSP70 protein level and sperm motility and viability were reported (Huang et al. 2000a; Turba et al. 2007) being that HSP70 seems to be more correlated with morphological abnormalities (Huszar et al. 2000; Turba et al. 2007).

It is important to underline that, in the present work, protein detection was performed after Percoll 256 257 selection of spermatozoa, and, therefore the lack of differences in HSPs content among the different boars may be related to the selection of only highly motile, good-quality spermatozoa, probably all rich in 258 259 HSPs. The opposite may obtain concerning the absence of CLU protein. CLU is reported in the literature as a marker of poor sperm quality and is essentially detectable only in low motile and morphologically 260 261 abnormal spermatozoa in humans and in bulls (O'Bryan et al. 1994; Ibrahim et al. 2000; Martínez-262 Heredia et al. 2008; Muciaccia et al. 2010) moreover the low number of CLU transcripts detected in our 263 Percoll-selected sample agrees with a recent paper that showed a higher content of the transcript in 264 motility-impaired bull semen compared with a good-quality one (Kumar et al. 2015).

Since sperm HSPs and CLU, like other proteins, are probably synthesized prior to chromatin 265 condensation, their presence cannot be related to the residual mRNA present in the mature cells. 266 Although the presence of transcripts should be representative of past cellular events during 267 268 spermatogenesis (Ostermeier et al. 2002; Lambard et al. 2004; Dadoune et al. 2005) the presence of relatively abundant amounts of HSP70 and HSP90 mRNA in Percoll-selected spermatozoa could also be 269 related to the ability to perform active translation of new proteins in developing embryos before maternal 270 271 genome activation. The importance of heat shock proteins in developing embryos is well known (Neuer et al. 2000; Zhang et al. 2011; Driver and Khatib 2013), though the paternal RNA contribution to protein 272 synthesis must be more precisely defined. A further study with the employment of an higher number of 273 274 boars and with the possibility to utilize a CASA system to evaluate semen parameters, will be useful to confirm our preliminary data and verify the correlation between the transcripts and semen quality and to 275

276	evaluate if these transcripts are active and related to embryonic development after fertilization, as
277	hypothesized by some authors (Ostermeier et al. 2004; Kempisty et al. 2008; Hwang et al. 2013).
278	
279	Conclusion
280	This is the first report on differential abundance of HSP70, 90 and clusterin transcripts in selected porcine
281	spermatozoa: the level of expression of HSP was high while CLU expression was very low. The
282	quantitative analysis of mRNA of HSPs that selected spermatozoa derived from high quality of the semen
283	show similar and high level of expression suggesting a potential role of these transcripts as reliable
284	markers of semen quality.
285	
286	
287	Compliance with ethical standards
288	
289	Conflict of interest The authors declare that there is no conflict of interest.
290	
291	Ethical approval All applicable international and institutional guide-lines for the care and use of animals
292	were followed
293	
294	
295	Acknowledgements
296	This study was supported by grant from Bologna University (RFO, Ricerca Fondamentale Orientata). The
297	authors wish to thank Dr Diego Bucci for his technical support.
298	
299	References
300	Althouse GC (2008) Sanitary procedures for the production of extended semen. Reprod Domest Anim 43:374–
301	378. doi: 10.1111/j.1439-0531.2008.01187.x
302	Althouse GC, Kuster CE, Clark SG, Weisiger RM (2000) Field investigations of bacterial contaminants and their
303	effects on extended porcine semen. Theriogenology 53:1167-1176. doi: 10.1016/S0093-691X(00)00261-2
304	Althouse GC, Lu KG (2005) Bacteriospermia in extended porcine semen. Theriogenology 63:573-584. doi:
305	10.1016/j.theriogenology.2004.09.031
306	Dadoune JP, Pawlak A, Alfonsi MF, Siffroi JP (2005) Identification of transcripts by macroarrays, RT-PCR and

307 in situ hybridization in human ejaculate spermatozoa. Mol Hum Reprod 11:133–140. doi:

308 10.1093/molehr/gah137

Driver AM, Khatib H (2013) Physiology and endocrinology symposium: heat shock proteins: potentially
 powerful markers for preimplantation embryonic development and fertility in livestock species. J Anim

311 Sci 91:1154–1161. doi: 10.2527/jas.2012-5928

- 312 Fantinati P, Zannoni A, Bernardini C, et al (2009) Evaluation of swine fertilisation medium (SFM) efficiency in
- 313 preserving spermatozoa quality during long-term storage in comparison to four commercial swine
- 314 extenders. Animal 3:269–274. doi: 10.1017/S1751731108003443
- Feugang JM, Rodriguez-Osorio N, Kaya A, et al (2010) Transcriptome analysis of bull spermatozoa:
- 316 implications for male fertility. Reprod Biomed Online 21:312–324. doi: 10.1016/j.rbmo.2010.06.022
- 317 Gilbert I, Bissonnette N, Boissonneault G, et al (2007) A molecular analysis of the population of mRNA in

318 bovine spermatozoa. Reproduction 133:1073–1086. doi: 10.1530/REP-06-0292

- 319 Hamatani T (2012) Human spermatozoal RNAs. Fertil Steril 97:275–281. doi: 10.1016/j.fertnstert.2011.12.035
- 320 Han Z, Wang Z, Cheng G, et al (2012) Presence, localization, and origin of clusterin in normal human
- 321 spermatozoa. J Assist Reprod Genet 29:751–757. doi: 10.1007/s10815-012-9779-x
- Huang SY, Kuo YH, Lee YP, et al (2000a) Association of heat shock protein 70 with semen quality in boars.
 Anim Reprod Sci 63:231–240.
- 324 Huang SY, Kuo YH, Tsou HL, et al (2000b) The decline of porcine sperm motility by geldanamycin, a specific

inhibitor of heat-shock protein 90 (HSP90). Theriogenology 53:1177–1184. doi: 10.1016/S0093691X(00)00262-4

Huszar G, Stone K, Dix D, Vigue L (2000) Putative creatine kinase M-isoform in human sperm is identified as
the 70-kilodalton heat shock protein HspA2. Biol Reprod 63:925–932.

Hwang JY, Mulligan BP, Kim H-M, et al (2013) Quantitative analysis of sperm mRNA in the pig: relationship
with early embryo development and capacitation. Reprod Fertil Dev 25:807–817. doi: 10.1071/RD12160

with early employ development and capacitation. Reprod Fertil Dev 25.807–817. doi: 10.1071/RD12100

- Ibrahim NM, Foster DN, Crabo BG (2001) Localization of clusterin on freeze-preserved bull spermatozoa before
 and after glass wool-sephadex filtration. J Androl 22:891–902. doi: 10.1002/j.1939-4640.2001.tb02594.x
- 333 Ibrahim NM, Gilbert GR, Loseth KJ, Crabo BG (2000) Correlation between clusterin-positive spermatozoa

determined by flow cytometry in bull semen and fertility. J Androl 21:887–894.

335 Ibrahim NM, Troedesson MHT, Foster DN, et al (1999) Reproductive tract secretions and bull spermatozoa

- 336 contain different clusterin isoforms that cluster cells and inhibit complement induced cytolysis. J Androl
- 337 20:230–240.

- 338 Jodar M, Selvaraju S, Sendler E, et al (2013) The presence, role and clinical use of spermatozoal RNAs. Hum
- 339 Reprod Update 19:604–624. doi: 10.1093/humupd/dmt031
- Kasimanickam V, Kasimanickam R, Arangasamy A, et al (2012) Association between mRNA abundance of
 functional sperm function proteins and fertility of Holstein bulls. Theriogenology 78:2007–2019. doi:
- 342 10.1016/j.theriogenology.2012.07.016
- Kempisty B, Antosik P, Bukowska D, et al (2008) Analysis of selected transcript levels in porcine spermatozoa,
 oocytes, zygotes and two-cell stage embryos. Reprod Fertil Dev 20:513–518.
- 345 Knox R, Levis D, Safranski T, Singleton W (2008) An update on North American boar stud practices.
- 346 Theriogenology 70:1202–1208. doi: 10.1016/j.theriogenology.2008.06.036
- Krawetz SA (2005) Paternal contribution: new insights and future challenges. Nat Rev Genet 6:633–642. doi:
 10.1038/nrg1654
- 349 Kumar S, Deb R, Singh U, et al (2015) Bovine circadian locomotor output cycles kaput (CLOCK) and clusterin
- 350 (CLU) mRNA quantitation in ejaculated crossbred bull spermatozoa. Reprod Domest Anim 50:505–509.
- 351 doi: 10.1111/rda.12522
- Kunavongkrit A, Suriyasomboon A, Lundeheim N, et al (2005) Management and sperm production of boars
 under differing environmental conditions. Theriogenology 63:657–667. doi:
- 354 10.1016/j.theriogenology.2004.09.039
- 355 Lambard S, Galeraud-Denis I, Martin G, et al (2004) Analysis and significance of mRNA in human ejaculated
- 356 sperm from normozoospermic donors: relationship to sperm motility and capacitation. Mol Hum Reprod
- 357 10:535–541. doi: 10.1093/molehr/gah064
- 358 Martínez-Heredia J, de Mateo S, Vidal-Taboada JM, et al (2008) Identification of proteomic differences in
- 359 asthenozoospermic sperm samples. Hum Reprod 23:783–791. doi: 10.1093/humrep/den024
- 360 Matàs C, Vieira L, Garcia-Vàzquez FA, et al (2011) Effects of centrifugation through three different
- 361 discontinuous Percoll gradients on boar sperm function. Anim Reprod Sci 127:62–72. doi:
- 362 10.1016/j.anireprosci.2011.06.009
- 363 Muciaccia B, Sette C, Paronetto MP, et al (2010) Expression of a truncated form of KIT tyrosine kinase in
- human spermatozoa correlates with sperm DNA integrity. Hum Reprod 25:2188–2202. doi:
- 365 10.1093/humrep/deq168
- Neuer A, Spandorfer SD, Giraldo P, et al (2000) The role of heat shock proteins in reproduction. Hum Reprod
 Update 6:149–159.
- 368 O'Bryan MK, Murphy BF, Liu DY, et al (1994) The use of anticlusterin monoclonal antibodies for the

- 369 combined assessment of human sperm morphology and acrosome integrity. Hum Reprod 9:1490–6.
- 370 Ostermeier GC, Dix DJ, Miller D, et al (2002) Spermatozoal RNA profiles of normal fertile men. Lancet

371 360:772–777. doi: 10.1016/S0140-6736(02)09899-9

- Ostermeier GC, Miller D, Huntriss JD, et al (2004) Reproductive biology: delivering spermatozoan RNA to the
 oocyte. Nature 429:154. doi: 10.1038/429154a
- Spinaci M, Volpe S, Bernardini C, et al (2005) Immunolocalization of heat shock protein 70 (Hsp 70) in boar
 spermatozoa and its role during fertilization. Mol Reprod Dev 72:534–541. doi: 10.1002/mrd.20367
- Spinaci M, Vallorani C, Bucci D, Bernardini C, Tamanini C, Seren E, Galeati G (2010) Effect of liquid storage
 on sorted boar spermatozoa. Theriogenology 74:741-748.
- 378 Sylvester SR, Morales C, Oko R, Griswold MD (1991) Localization of sulfated glycoprotein-2 (clusterin) on

379 spermatozoa and in the reproductive tract of the male rat. Biol Reprod 45:195–207.

- 380 Turba ME, Fantinati P, Bernardini C, et al (2007) Relationships between innovative and traditional parameters to
- 381 investigate semen quality in pigs. Anim Reprod Sci 99:72–81. doi: 10.1016/j.anireprosci.2006.04.045
- Volpe S, Galeati G, Bernardini C, et al (2008) Comparative immunolocalization of heat shock proteins (Hsp)-60,
- 383 -70, -90 in boar, stallion, dog and cat spermatozoa. Reprod Domest Anim 43:385–392. doi:
- 384 10.1111/j.1439-0531.2007.00918.x
- Wang P, Wang Y-F, Wang H, et al (2014) HSP90 expression correlation with the freezing resistance of bull
 sperm. Zygote 22:239–245. doi: 10.1017/S096719941300004X
- 387 Yang CC, Lin YS, Hsu CC, et al (2009) Identification and sequencing of remnant messenger RNAs found in
- 388 domestic swine (*Sus scrofa*) fresh ejaculated spermatozoa. Anim Reprod Sci 113:143–155. doi:

389 10.1016/j.anireprosci.2008.08.012

- Zannoni A, Zaniboni A, Bernardini C, et al (2015) Expression fo clusterin isoforms in LPS treated porcine aortic
 endothelial cells. Pak Vet J 33:236–238.
- 392 Zhang B, Peñagaricano F, Driver A, et al (2011) Differential expression of heat shock protein genes and their
- 393 splice variants in bovine preimplantation embryos. J Dairy Sci 94:4174–4182. doi: 10.3168/jds.2010-4137
- 394 Zhang X-G, Hu S, Han C, et al (2015) Association of heat shock protein 90 with motility of post-thawed sperm

395 in bulls. Cryobiology 70:164–169. doi: 10.1016/j.cryobiol.2014.12.010

- 396
- 397

```
Table caption
```

399

400	Table 1 Forward and reverse primer sequences, RT-PCR length (bp) and accession numbers (Acc. No.) in
401	the NCBI database.
402	
403	Figure legends
404	
405	Fig. 1 Percentage of viability and motility and concentration values of ejaculated spermatozoa from different
406	boars (B1-B7) evaluated before density gradient. The data are expressed as mean \pm S.D. Different letters
407	indicate statistically significant differences ($P < 0.05$, ANOVA post hoc Tukey test) among boars.
408	
409	Fig. 2 Relative gene expression of HSP70, HSP90 and clusterin (CLU) in ejaculated spermatozoa, after
410	Percoll selection, of different boars (B1-7). mRNA data are presented as Δ Ct method (reference gene Ct –
411	gene of interest Ct). Error bars represent the range of expression. Different lower case or different capital
412	letters indicate statistically significant differences among boars ($P < 0.05$, ANOVA post hoc Tukey test)
413	within HSP70 (white histograms, lower case) or within HSP90 (grey histograms, capital letters) gene
414	expression. No statistically significant differences were observed in CLU gene expression (black
415	histograms).
416	GI = interest gene (HSP70 or HSP90 or CLU).
417	
418	Fig. 3 A) Expression of HSP70 (white histograms) and HSP90 (grey histograms) in ejaculated spermatozoa,
419	after Percoll selection, of different boars (B1-7). The data are represented as mean \pm SEM of relative protein
420	content (AU = arbitrary units). B) Representative images of western blotting of HSP70 and 90 in ejaculated
421	spermatozoa. MWM molecular weight marker.