

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Preliminary tests on in vitro activity of different pure and commercial compounds against Saprolegnia spp.

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Availability:

This version is available at: https://hdl.handle.net/11585/622818 since: 2019-01-14

Published:

DOI: http://doi.org/

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)



ParaFishControl

www.parafishcontrol.eu

PRELIMINARY TESTS ON *IN VITRO* ACTIVITY OF DIFFERENT PURE AND COMMERCIAL COMPOUNDS AGAINST *SAPROLEGNIA* SPP.

Galuppi R., Tedesco P., Ciardo M.L., Gustinelli A., Menconi V., Caffara M., Fioravanti M.L. Department of Veterinary Medical Sciences, Alma Mater Studiorum Università di Bologna, Italy



Introduction

Oomycetes of the genus *Saprolegnia* are responsible for severe economic losses in freshwater aquaculture. Compounds of proven activity against *Saprolegnia* spp. (i.e. malachite green) are potentially hazardous to human health and the environment, and their use is forbidden by European regulations. Therefore, the demand for new treatments pushes towards the selection of more safe and environmentally friendly products.

In the present work, in vitro activity of two pure compounds (Oxalic acid; Sodium percarbonate) and two commercial products (Actidrox[®], De Marco, Italy; Virkon[®] S, Dupont) was tested on different strains of Saprolegnia. Malachite green was used as reference compound.



protocol I



Methodology

Preliminary trials were performed using two protocols available in the literature by Alderman (1982, Journal of Fish Diseases 8:289-298): one screening method in Agar (**protocol I**) aimed at assessing the *minimum* inhibitory concentration (MIC) and one hour bath in aqueous solution of mycelium growing on polycarbonate membrane (protocol II), to assess the minimum lethal concentration (MLC). Two field strains of Saprolegnia spp. isolated from rainbow trout (Oncorhynchus mykiss) and brown trout (Salmo trutta) and one reference strain of S. parasitica (CBS 223.65 furnished by CSIC-RJB) have been tested in triplicate per each concentration.

Results

Our results show that oxalic acid and Virkon[®]

are effective in inhibiting the growth of the

mycelium, although at concentrations too

high to be applied in the field (MIC 1000

mg/L). Actidrox[®] showed a different activity

between the two methods (MIC 5000 mg/L;

MLC 500 mg/L), possibly due to its

mechanism of action, that requires presence

of water. Tested concentrations of Sodium

percarbonate were effective only in slowing

down the mycelium growth.



protocol II







Average diameters of the colonies with different concentrations of tested compounds



	MIC	MLC
compound	agar	water
malachite green	5 mg/L	5 mg/L
oxalic acid	1000 mg/L	5000 mg/L
		(1000 mg/L for CBS 233.65 strain)
Actidrox®	5000 mg/L	500 mg/L
sodium percarbonate	> 5000 mg/L	> 5000 mg/L
Virkon®	1000 mg/L	1000 mg/L





Average radial growth of hyphae beyond the filter area after immersion treatment with different concentrations of tested compounds





Mycelium growth after 6 days at 3 different concentrations of malachite green

Examples of *Saprolegnia* growth on agar with different concentrations of malachite green (left) and oxalic acid (right) after 24 h of incubation

Conclusions

Further *in vitro* trials will be necessary, considering a wider range of promising compounds. The combination of the two methods (inoculation in agar and contact in aqueous solution) represents a good investigation approach for screening the activity of different molecules and products against *Saprolegnia* spp.



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 634429. This output reflects the views only of the author(s), and the European Union cannot be held responsible for any use which may be made of the information contained therein.