



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Preliminary tests on in vitro activity of different pure and commercial compounds against *Saprolegnia* spp.

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Availability:

This version is available at: <https://hdl.handle.net/11585/622818> since: 2019-01-14

Published:

DOI: <http://doi.org/>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Preliminary tests on *in vitro* activity of different pure and commercial compounds against *Saprolegnia* spp.

Roberta G¹, Tedesco P¹, Ciardo M¹, Gustinelli A¹, Menconi V¹, Caffara M¹, **Fioravanti M¹**

¹*Department of Veterinary Medical Sciences, Alma Mater Studiorum Università di Bologna, Ozzano Emilia, Italy*

Introduction: Oomycetes of the genus *Saprolegnia* are responsible for severe economic losses in freshwater aquaculture. Compounds of proven activity against *Saprolegnia* spp. (i.e. malachite green) are potentially hazardous to human health and the environment, and their use is forbidden by European regulations. Therefore, the demand for new treatments pushes towards the selection of more safe and environmentally friendly products.

In the present work, *in vitro* activity of two pure compounds (Oxalic acid; Sodium percarbonate) and two commercial products (Actidrox®, De Marco, Italy; Virkon® S, Dupont) was tested on different strains of *Saprolegnia*. Malachite green was used as reference compound.

Methodology: Preliminary trials were performed using two protocols available in the literature by Alderman: one screening method in Agar aimed at assessing the *minimum inhibitory concentration* (MIC) and one hour bath in aqueous solution of mycelium growing on polycarbonate membrane, to assess the *minimum lethal concentration* (MLC). Two field strains of *Saprolegnia* spp. isolated from rainbow trout (*Oncorhynchus mykiss*) and brown trout (*Salmo trutta*) and one reference strain of *S. parasitica* (CBS 223.65 furnished by CSIC-RJB) have been tested in triplicate per each concentration.

Results: Our results show that oxalic acid and Virkon® are effective in inhibiting the growth of the mycelium, although at concentrations too high to be applied in the field (MIC 1000 mg/ L). Actidrox® showed a different activity between the two methods (MIC 5000 mg/ L; MLC 500 mg/ L), possibly due to its mechanism of action, that requires presence of water. Tested concentrations of Sodium percarbonate were effective only in slowing down the mycelium growth.

Conclusions: Further *in vitro* trials will be necessary, considering a wider range of promising compounds. The combination of the two methods (inoculation in agar and contact in aqueous solution) represents a good investigation approach for screening the activity of different molecules and products against *Saprolegnia* spp.

Funding of presentation:

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 634429. This publication reflects the views only of the author, and the European Commission cannot be held responsible for any use which may be made of the information contained therein.