Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Combined approaches for the sensory "targetization" of volatile compounds in virgin olive oils by SPME-GC-FID

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:
Availability:
This version is available at: https://hdl.handle.net/11585/648844 since: 2018-11-06
Published:
DOI: http://doi.org/
Terms of use:
Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

Combined approaches for the sensory "targetization" of volatile compounds in virgin olive oils by SPME-GC-FID

Enrico Valli¹, Diego Luis Garcia Gonzalez², Ramon Aparicio-Ruiz², Enrico Casadei¹,
Sara Barbieri¹, Alessandra Bendini¹, Tullia Gallina Toschi¹

¹Alma Mater Studiorum - Università di Bologna, Bologna, Italy,

²Instituto de la Grasa, Sevilla, Spain

It is well known that specific volatile compounds are strictly related with the positive and negative sensory attributes of virgin olive oils, such as fruitiness and defects, thus being key molecules for the product quality and the consumers' acceptability. The objective of this research is to link the volatile compounds with the sensory attributes through a joint approach that could support the organoleptic analysis of virgin olive oils.

With this aim, a wide set of commercial olive oils was previously assessed (EU Reg. 1227/2016) by six sensory panels, resulting in different commercial categories (extra virgin, virgin and lampante) and characterized by various intensities of fruitiness and defects. The headspace volatile profiles data of the UNIBO and CSIC units, obtained by SPME-GC-FID, were statistically elaborated through a number of treatments and a subsequent univariate and multivariate analysis. The approach herein discussed was able to detect the most robust markers of specific sensory defects, thus being possibly applied to all the volatiles data sets (SPME-GC-MS, FGC-e-nose, TDU-GC-MS) of the OLEUM project. The final goal was to find the most relevant analytes to be considered in a method for the determination of volatile compounds in olive oils and also useful for the formulation of sensory reference standards.

This work was developed in the context of the project OLEUM "Advanced solutions for assuring authenticity and quality of olive oil at global scale" funded by the European Commission within the Horizon 2020 Programme (2014–2020, grant agreement no. 635690).