



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Nonuniformly elliptic energy integrals with p, q -growth

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Cupini, G., Marcellini, P., Mascolo, E. (2018). Nonuniformly elliptic energy integrals with p, q -growth. *NONLINEAR ANALYSIS*, 177, 312-324 [10.1016/j.na.2018.03.018].

Availability:

This version is available at: <https://hdl.handle.net/11585/649528.3> since: 2018-11-13

Published:

DOI: <http://doi.org/10.1016/j.na.2018.03.018>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Cupini, Giovanni, Paolo Marcellini, e Elvira Mascolo. «Nonuniformly Elliptic Energy Integrals with p, q -Growth». *Nonlinear Analysis* 177 (dicembre 2018): 312–24.

The final published version is available online at:
<https://doi.org/10.1016/j.na.2018.03.018>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)


When citing, please refer to the published version.



Contents lists available at ScienceDirect

Nonlinear Analysis

www.elsevier.com/locate/na



Nonuniformly elliptic energy integrals with p, q -growth

Giovanni Cupini^a, Paolo Marcellini^{b,*}, Elvira Mascolo^b

^a *Dipartimento di Matematica, Università di Bologna, Piazza di Porta S. Donato 5, 40126 Bologna, Italy*

^b *Dipartimento di Matematica e Informatica "U. Dini", Università di Firenze Viale Morgagni 67/A, 50134 Firenze, Italy*

ARTICLE INFO

Article history:

Received 25 March 2018

Accepted 28 March 2018

Communicated by Enzo Mitidieri

Carlo Sbordone is a leader in Calculus of Variations and PDEs, as well as a master in boosting all aspects of mathematical projects and events. He was among the first one to appreciate Neil Trudinger's work [26] on nonuniformly linear elliptic operators and to find application – for instance – to homogenization [21,22] and not only to this subject [5]. It is an honor for us to dedicate this work to Carlo on the occasion of his seventieth birthday.

MSC:

primary 49N60

secondary 35J60

Keywords:

Nonuniformly elliptic equations

p, q -growth conditions

Local boundedness

ABSTRACT

We study the local boundedness of minimizers of a nonuniformly energy integral of the form $\int_{\Omega} f(x, Dv) dx$ under p, q -growth conditions of the type

$$\lambda(x)|\xi|^p \leq f(x, \xi) \leq \mu(x)(1 + |\xi|^q)$$

for some exponents $q \geq p > 1$ and with nonnegative functions λ, μ satisfying some summability conditions. We use here the original notation introduced in 1971 by Trudinger [26], where $\lambda(x)$ and $\mu(x)$ had the role of the minimum and the maximum eigenvalues of an $n \times n$ symmetric matrix $(a_{ij}(x))$ and

$$f(x, \xi) = \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j$$

was the energy integrand associated to a linear nonuniformly elliptic equation in divergence form. In this paper we consider a class of energy integrals, associated to nonlinear nonuniformly elliptic equations and systems, with integrands $f(x, \xi)$ satisfying the general growth conditions above.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

In the recent mathematical literature a large interest received the energy integral

$$\int_{\Omega} \{|Du(x)|^p + |x|^\alpha |Du(x)|^q\} dx, \quad (1.1)$$

* Corresponding author.

E-mail addresses: giovanni.cupini@unibo.it (G. Cupini), marcellini@math.unifi.it (P. Marcellini), mascolo@math.unifi.it (E. Mascolo).

where $1 < p \leq q$, $\alpha > 0$, Ω is an open set in \mathbb{R}^n , $n \geq 2$, $0 \in \Omega$, $u : \Omega \rightarrow \mathbb{R}^m$, $m \geq 1$, and Du is the gradient of u . See for instance [1,3,4,14,15,16,24,27]. To explain the point of view of this research, we denote as ξ the gradient variable and

$$f(x, \xi) := |\xi|^p + |x|^\alpha |\xi|^q ;$$

then $f : \Omega \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ is a convex function with respect to $\xi \in \mathbb{R}^{m \times n}$ which satisfies the growth conditions

$$|\xi|^p \leq f(x, \xi) \leq c(1 + |\xi|^q) \tag{1.2}$$

and also

$$|x|^\alpha |\xi|^q \leq f(x, \xi) \leq c(1 + |\xi|^q) \tag{1.3}$$

for some positive constant c and for every $\xi \in \mathbb{R}^{m \times n}$. With respect to the condition (1.2) we can say that $f(x, \xi)$ enters in the p, q -growth regularity theory and, by the recent results in [6,8,10], we know that any local minimizer of the energy integral in (1.1) is locally bounded in Ω if $p < n$ and

$$q < p^* =: \frac{np}{n-p}$$

(with $q \leq p^*$ if $m = 1$). However for the model integral (1.1) our point of view here is to take under consideration also condition (1.3). With the symbols described more in details below, we have that, if $\alpha < q$, then $\lambda^{-1} := |x|^{-\alpha} \in L^r_{loc}(\Omega)$ for some r such that $\frac{n}{q} < r < \frac{n}{\alpha}$. This allows us to apply Theorem 1.1 and to obtain that every local minimizer of the energy integral (1.1) is locally bounded.

Let us summarize the discussion above: any local minimizer of the energy integral (1.1) is locally bounded in Ω if either the integrand $f(x, \xi)$ satisfies the p, q -growth in (1.2) and $q < p^*$, or if $f(x, \xi)$ is nonuniformly elliptic as in (1.3) and $\alpha < q$. Of course here it would be interesting to unify the two cases and to give a sole condition. This is one of our aims in what we propose below.

In this paper we consider the general case of nonuniformly elliptic energy integrals of p, q -growth of the type

$$\mathcal{F}(v; \Omega) = \int_{\Omega} f(x, Dv) dx \tag{1.4}$$

with

$$\begin{cases} \lambda(x)|\xi|^p \leq f(x, \xi) \leq \mu(x) (|\xi|^q + 1) \\ \lambda^{-1} \in L^r_{loc}(\Omega), \quad \mu \in L^s_{loc}(\Omega) \end{cases} \tag{1.5}$$

for $1 < p \leq q$, for some exponents $r \in [1, \infty]$, $s \in (1, \infty]$ and for every ξ in $\mathbb{R}^{m \times n}$. The integrand $f(x, \xi)$ also satisfies the condition

$$f(x, \xi) = g(x, |\xi|) , \tag{1.6}$$

where $g : \Omega \times [0, \infty) \rightarrow [0, \infty)$ is a convex Δ_2 function of class C^1 with respect to the last variable, with $g(x, 0) = g_t(x, 0) = 0$.

Theorem 1.1. *Let us consider the energy integral (1.4) under the assumptions (1.5), (1.6), with $r \in [1, \infty]$, $s \in (1, \infty]$ (if $p \leq 2$ then we also require $r > \frac{1}{p-1}$) and*

$$\frac{1}{pr} + \frac{1}{qs} + \frac{1}{p} - \frac{1}{q} < \frac{1}{n} ; \tag{1.7}$$

then every local minimizer of the energy integral (1.4) is locally bounded.

We observe that in the relevant case $r = s = \infty$ then condition (1.7) can be equivalently written in the form $q < p^*$. We also point out the connection of our result with the pioneering and celebrated paper by Trudinger [26], where it is studied the linear case of nonuniformly elliptic second order equations (of course with $p = q = 2$). Indeed, if $p = q$, then (1.7) becomes

$$\frac{1}{r} + \frac{1}{s} < \frac{p}{n},$$

which is exactly the assumption on the summability exponents appearing in [26] in the particular case $p = q = 2$.

We refer to the next Section 2 for more details. There we also fix the notation and we give the complete statement of our main result, see Theorem 2.1, including a precise estimate for the L^∞ norm. In Section 3 we give some preliminary results, while Section 4 is devoted to the proof of Theorem 2.1.

For completeness, the regularity theory under p, q -growth conditions have been considered nowadays by many authors. See for instance [1–4, 11, 14, 15, 17, 18, 20] and recently [12] with geometric constraints under p, q -growth conditions; see also the survey [23] on regularity under non standard growth.

Our research collaboration on regularity in the calculus of variations under p, q -growth conditions was originated in 2009 and focused in particular on the local boundedness of local minimizers [6–10]. A comparison with the previous papers and in particular with the most recent [10] shows that here we consider the vector-valued case (with respect to the scalar one in [10]) and we assume here quite weaker summability conditions on the coefficients.

2. The main results

Let us consider the functional

$$\mathcal{F}(v; \Omega) = \int_{\Omega} f(x, Dv) \, dx, \tag{2.1}$$

with $\Omega \subset \mathbb{R}^n$, $n \geq 2$, open set, and $f : \Omega \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$, $m \geq 1$, a Carathéodory function. We assume that there exist measurable functions $\lambda, \mu : \Omega \rightarrow [0, \infty)$ such that

$$\lambda(x)|\xi|^p \leq f(x, \xi) \leq \mu(x)(|\xi|^q + 1), \quad 1 < p \leq q, \tag{2.2}$$

for a.e. $x \in \Omega$ and every $\xi \in \mathbb{R}^{m \times n}$, where

$$\lambda^{-1} \in L^r_{loc}(\Omega), \quad \mu \in L^s_{loc}(\Omega),$$

for some exponents $r \in [1, \infty]$ and $s \in (1, \infty]$.

With simple computations (see Proposition 3.1), for every open set Ω' compactly contained in Ω we deduce that

$$\begin{aligned} \|\lambda^{-1}\|_{L^r(\Omega')}^{-1} \|Dv\|_{L^{\frac{pr}{r+1}}(\Omega'; \mathbb{R}^{m \times n})}^p &\leq \int_{\Omega'} f(x, Dv) \, dx \leq \|\mu\|_{L^s(\Omega')} \|Dv\|_{L^{\frac{qs}{s-1}}(\Omega'; \mathbb{R}^{m \times n})}^q \\ &\quad + \|\mu\|_{L^1(\Omega')}, \end{aligned} \tag{2.3}$$

and therefore we have

$$W^{1, \frac{pr}{r+1}}(\Omega'; \mathbb{R}^m) \supset W^{1, \mathcal{F}}(\Omega'; \mathbb{R}^m) \supset W^{1, \frac{qs}{s-1}}(\Omega'; \mathbb{R}^m),$$

where $W^{1, \mathcal{F}}(\Omega'; \mathbb{R}^m)$ denotes the set of maps u of finiteness of the integral; i.e.,

$$W^{1, \mathcal{F}}(\Omega'; \mathbb{R}^m) := \{u \in W^{1,1}(\Omega'; \mathbb{R}^m) : \mathcal{F}(u; \Omega') < \infty\}.$$

It is clear that if we fix appropriate conditions at the boundary of a fixed Ω' , then from standard direct methods of the calculus of variations we derive existence of minimizers in $W^{1, \frac{pr}{r+1}}(\Omega'; \mathbb{R}^m)$; at this stage we need the condition $\frac{pr}{r+1} > 1$.

Of course any minimizer belongs also to $W^{1, \mathcal{F}}(\Omega'; \mathbb{R}^m)$.

In the vector-valued case, as suggested by well known counterexamples by De Giorgi [13], Giusti–Miranda [19], Šverák–Yan [25], generally some structure conditions are required for everywhere regularity. Thus, as usual in this theory, we assume that f is a radial function with respect to the gradient variable ξ . Precisely:

there exists $g : \Omega \times [0, \infty) \rightarrow [0, \infty)$ such that, for a.e. $x \in \Omega$,

$$f(x, \xi) = g(x, |\xi|) \text{ is a convex and } C^1 \text{ function with respect to } \xi \in \mathbb{R}^{m \times n}. \tag{2.4}$$

We also assume the so-called Δ_2 -condition holds:

there exists $\gamma \geq 1$ such that

$$f(x, t\xi) \leq t^\gamma f(x, \xi) \tag{2.5}$$

for every $t > 1$, for a.e. x and every ξ . This condition implies that $W^{1, \mathcal{F}}(\Omega'; \mathbb{R}^m)$ is a vector space.

We study the regularity of local minimizers of \mathcal{F} . We recall that a function $u \in W^{1, \frac{pr}{r+1}}_{\text{loc}}(\Omega; \mathbb{R}^m)$ is a local minimizer of \mathcal{F} if

$$\mathcal{F}(u; \Omega') \leq \mathcal{F}(u + \varphi; \Omega')$$

for every open set Ω' compactly contained in Ω and for all $\varphi \in W^{1, \mathcal{F}}(\Omega; \mathbb{R}^m)$ with $\text{supp } \varphi \Subset \Omega'$.

Before stating our regularity result, we recall that, given a real number $\ell \geq 1$, ℓ^* is its Sobolev exponent, i.e.

$$\ell^* := \begin{cases} \frac{n\ell}{n-\ell} & \text{if } \ell < n, \\ \text{any number } > \ell & \text{if } \ell \geq n \end{cases}$$

and ℓ' is the conjugate exponent of ℓ , i.e., $\frac{1}{\ell} + \frac{1}{\ell'} = 1$. Moreover, as usual, $\frac{1}{\infty}$ has to be read as 0.

Theorem 2.1. *Let us assume that (2.2), (2.4) and (2.5) hold under the summability conditions*

$$\lambda^{-1} \in L^r_{\text{loc}}(\Omega), \quad \mu \in L^s_{\text{loc}}(\Omega),$$

for some $r \in [1, \infty]$, $s \in (1, \infty]$ such that

$$\frac{1}{pr} + \frac{1}{qs} + \frac{1}{p} - \frac{1}{q} < \frac{1}{n}. \tag{2.6}$$

If $p \leq 2$ we also require $r > \frac{1}{p-1}$. Then, every local minimizer $u \in W^{1, \frac{pr}{r+1}}_{\text{loc}}(\Omega; \mathbb{R}^m)$ of \mathcal{F} is locally bounded and there exists a positive constant c , depending on p, q, n, m, γ , such that

$$\|u\|_{L^\infty(B_{R_0/2}(x_0); \mathbb{R}^m)} \leq c \left\{ R_0^{-q} \|\lambda^{-1}\|_{L^r(B_{R_0})} \|\mu\|_{L^s(B_{R_0})} \right\}^{\vartheta_1} \| |u| + 1 \|_{L^{qs'}(B_{R_0}(x_0))}^{\vartheta_2}, \tag{2.7}$$

for every $x_0 \in \Omega$ and $B_{R_0}(x_0) \Subset \Omega$, $0 < R_0 \leq 1$, where $\sigma := \frac{pr}{r+1}$ ($\sigma := p$ if $r = \infty$) and

$$\vartheta_1 := \frac{\sigma^*}{p(\sigma^* - qs')}, \quad \vartheta_2 := \frac{q(\sigma^* - ps')}{p(\sigma^* - qs')}.$$

Remark 2.2. If $r = s = \infty$ inequality (2.6) reduces to $q < p^*$, see e.g. [7,8,10]. Moreover, we observe that the assumption on the summability exponent of μ can be written as

$$\left(\frac{\sigma^*}{q}\right)' < s \leq \infty, \tag{2.8}$$

and that, if $p = q$, then it becomes

$$\frac{1}{s} + \frac{1}{r} < \frac{p}{n}.$$

Remark 2.3. Trudinger [26] considered nonuniformly elliptic second order differential equations, with $p = q = 2$, under the following assumptions on λ and μ

$$\lambda^{-1} \in L^r_{\text{loc}}(\Omega), \quad \lambda^{-1}\mu^2 \in L^s_{\text{loc}}(\Omega) \quad \frac{1}{s} + \frac{1}{r} < \frac{2}{n}.$$

His condition on μ is slightly stronger than ours; in fact in our context $\lambda(x) \leq 2\mu(x)$ a.e. in Ω (by (2.2); also in [26] $\lambda(x) \leq \mu(x)$) and thus

$$\lambda^{-1}\mu^2 \in L^s_{\text{loc}}(\Omega) \Rightarrow \mu \in L^s_{\text{loc}}(\Omega).$$

3. Preliminary results

We start this section with a precise statement and proof of (2.3).

Proposition 3.1. Assume that (2.2) holds, where $\lambda^{-1} \in L^r_{\text{loc}}(\Omega)$, $\mu \in L^s_{\text{loc}}(\Omega)$ for some exponents $r \in [1, \infty]$ and $s \in (1, \infty]$ (if $p \leq 2$ then we also require $r \geq \frac{1}{p-1}$). Then for every $v \in W^{1, \frac{pr}{r+1}}_{\text{loc}}(\Omega; \mathbb{R}^m)$ and every open $\Omega' \Subset \Omega$ the inequalities (2.3) hold; i.e.,

$$\|\lambda^{-1}\|^{-1}_{L^r(\Omega')} \|Dv\|^p_{L^{\frac{pr}{r+1}}(\Omega'; \mathbb{R}^{m \times n})} \leq \int_{\Omega'} f(x, Dv) dx \leq \|\mu\|_{L^s(\Omega')} \|Dv\|^q_{L^{\frac{qs}{s-1}}(\Omega'; \mathbb{R}^{m \times n})} + \|\mu\|_{L^1(\Omega')}.$$

Proof. Consider $r \in [1, \infty)$, $v \in W^{1, \frac{pr}{r+1}}_{\text{loc}}(\Omega; \mathbb{R}^m)$ and an open set $\Omega' \Subset \Omega$. By Hölder inequality with exponents $\frac{r+1}{r}$ and $r+1$,

$$\int_{\Omega'} |Dv|^{\frac{pr}{r+1}} dx = \int_{\Omega'} \lambda^{\frac{r}{r+1}} |Dv|^{\frac{pr}{r+1}} \lambda^{-\frac{r}{r+1}} dx \leq \left(\int_{\Omega'} \lambda |Dv|^p dx\right)^{\frac{r}{r+1}} \left(\int_{\Omega'} (\lambda^{-1})^r dx\right)^{\frac{1}{r+1}}.$$

Therefore, using the left inequality in (2.2)

$$\int_{\Omega'} f(x, Dv) dx \geq \int_{\Omega'} \lambda |Dv|^p dx \geq \left(\int_{\Omega'} |Dv|^{\frac{pr}{r+1}} dx\right)^{\frac{r+1}{r}} \left(\int_{\Omega'} (\lambda^{-1})^r dx\right)^{-\frac{1}{r}}$$

and the left inequality in (2.3) follows.

If $r = \infty$, that is $\frac{r}{r+1} = 1$, consider $v \in W^{1,p}_{\text{loc}}(\Omega; \mathbb{R}^m)$. The left inequality in (2.3) follows by

$$\int_{\Omega'} |Dv|^p dx \leq \|\lambda^{-1}\|_{L^\infty(\Omega')} \int_{\Omega'} \lambda |Dv|^p dx$$

and the left inequality in (2.2).

Let us now prove the right inequality in (2.3).

Let us first consider $s \in (1, \infty)$. If $v \notin W^{1, \frac{qs}{s-1}}(\Omega')$ the inequality trivially holds. Let us assume $v \in W^{1, \frac{qs}{s-1}}(\Omega')$. By Hölder inequality with exponents s and $\frac{s}{s-1}$, we obtain

$$\int_{\Omega'} \mu |Dv|^q dx \leq \left(\int_{\Omega'} \mu^s dx \right)^{\frac{1}{s}} \left(\int_{\Omega'} |Dv|^{\frac{qs}{s-1}} dx \right)^{\frac{s-1}{s}}.$$

Using the right inequality in (2.2) we conclude.

Let us now consider $s = \infty$, that is $\frac{s}{s-1} = 1$. If $v \notin W^{1,q}(\Omega')$ the inequality trivially holds. Let us assume $v \in W^{1,q}(\Omega')$. We obtain

$$\int_{\Omega'} \mu |Dv|^q dx \leq \|\mu\|_{L^\infty(\Omega')} \int_{\Omega'} |Dv|^q dx.$$

Using the right inequality in (2.2) we conclude. \square

The next lemma is about a Poincaré – Sobolev type inequality.

Lemma 3.2. Consider a bounded open set $\Omega \subset \mathbb{R}^n$ and let p and r be such that $p > 1$ and $r \in [1, \infty]$ (if $p \leq 2$ then we also require $r \geq \frac{1}{p-1}$). Let $v \in W_0^{1, \frac{pr}{r+1}}(\Omega; \mathbb{R}^m)$ ($v \in W_0^{1,p}(\Omega; \mathbb{R}^m)$ if $r = \infty$) and let $\lambda : \Omega \rightarrow [0, \infty)$ be a measurable function such that $\lambda^{-1} \in L^r(\Omega)$. Then there exists a positive constant c such that

$$\left\{ \int_{\Omega} |v|^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} \leq c \|\lambda^{-1}\|_{L^r(\Omega)} \int_{\Omega} \lambda |Dv|^p dx, \tag{3.1}$$

where $\sigma := \frac{pr}{r+1}$ ($\sigma := p$ if $r = \infty$).

Proof. First, assume that $r < \infty$. By applying the Poincaré inequality and Hölder inequality with exponents $\frac{r+1}{r}, r+1$ we obtain

$$\begin{aligned} \left\{ \int_{\Omega} |v|^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} &\leq c \left\{ \int_{\Omega} |Dv|^{\sigma} dx \right\}^{\frac{p}{\sigma}} \leq c \left\{ \int_{\Omega} \left(\lambda^{\frac{r}{r+1}} |Dv|^{\sigma} \right) \lambda^{-\frac{r}{r+1}} dx \right\}^{\frac{p}{\sigma}} \\ &\leq c \|\lambda^{-1}\|_{L^r(B_R)} \int_{\Omega} \lambda |Dv|^p dx. \end{aligned}$$

Let us now discuss the case $r = \infty$, that implies $\sigma = p$. We have

$$\begin{aligned} \left\{ \int_{\Omega} |v|^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} &\leq c \int_{\Omega} |Dv|^p dx \leq c \int_{B_R} \lambda^{-1} (\lambda |Dv|^p) dx \\ &\leq c \|\lambda^{-1}\|_{L^\infty(\Omega)} \int_{\Omega} \lambda |Dv|^p dx. \end{aligned}$$

This concludes the proof of (3.1). \square

The following lemma deals with well known properties of the convex functions satisfying (2.2), (2.4) and (2.5).

Lemma 3.3. Assume (2.2), (2.4), (2.5). Then

- (i) $f(x, t\xi) \leq \max\{1, t^\gamma\} f(x, \xi)$ for every $t > 0$ and every $\xi \in \mathbb{R}^{m \times n}$,
- (ii) $f(x, \xi + \eta) \leq 2^{\gamma-1} (f(x, \xi) + f(x, \eta))$ for every $\xi, \eta \in \mathbb{R}^{m \times n}$,
- (iii) $g_t(x, t) \leq \gamma g(x, t)$ for every $t \geq 0$.

We also remark that a Euler’s equation holds true.

Proposition 3.4. Assume (2.2), (2.4), (2.5) and let u be a local minimizer of (2.1). Then

$$\int_{\Omega} \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, Du) \varphi_{x_i}^\alpha dx = 0 \tag{3.2}$$

for all $\varphi \in W^{1,\mathcal{F}}(\Omega; \mathbb{R}^m)$, $\text{supp } \varphi \Subset \Omega$.

Proof. Let $\varphi \in W^{1,\mathcal{F}}(\Omega; \mathbb{R}^m)$, $\text{supp } \varphi \Subset \Omega$. By (2.4) also $-\varphi$ is in $W^{1,\mathcal{F}}(\Omega; \mathbb{R}^m)$. By Lemma 3.3 we get $u + t\varphi \in W^{1,\mathcal{F}}(\Omega; \mathbb{R}^m)$ for every $t \in \mathbb{R}$. By the local minimality of u ,

$$\mathcal{F}(u) \leq \mathcal{F}(u + t\varphi) \quad \forall t \in \mathbb{R}.$$

To prove (3.2) it suffices to prove that

$$\left. \frac{d}{dt} \mathcal{F}(u + t\varphi) \right|_{t=0} = \int_{\Omega} \left. \frac{d}{dt} f(x, Du(x) + tD\varphi(x)) \right|_{t=0} dx.$$

To prove this, we need to prove that

$$\left| \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, Du + tD\varphi) \varphi_{x_i}^\alpha \right| \leq H(x) \quad \forall t \in (-1, 1)$$

with $H \in L^1(\text{supp } \varphi)$. By the convexity,

$$f(x, \xi_0) - f(x, 2\xi_0 - \xi) \leq \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, \xi_0) (\xi_i^\alpha - (\xi_0)_i^\alpha) \leq f(x, \xi) - f(x, \xi_0).$$

If $\xi_0 = Du(x) + tD\varphi(x)$, $\xi = Du(x) + (1+t)D\varphi(x)$, we have $2\xi_0 - \xi = Du(x) + (t-1)D\varphi(x)$ and

$$\begin{aligned} f(x, Du + tD\varphi) - f(x, Du + (t-1)D\varphi) &\leq \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, Du + tD\varphi) \varphi_{x_i}^\alpha \\ &\leq f(x, Du + (1+t)D\varphi) - f(x, Du + tD\varphi). \end{aligned}$$

Therefore, since f is non-negative,

$$\left| \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, Du + tD\varphi) \varphi_{x_i}^\alpha \right| \leq f(x, Du + (1+t)D\varphi) + f(x, Du + (t-1)D\varphi).$$

Using again the convexity and Lemma 3.3 we get

$$\begin{aligned} f(x, Du + (1+t)D\varphi) &\leq tf(x, Du + 2D\varphi) + (1-t)f(x, Du + D\varphi) \\ &\leq f(x, Du + 2D\varphi) + f(x, Du + D\varphi) \leq 2^\gamma (f(x, Du + D\varphi) + f(x, D\varphi)) \\ &\leq 2^{2\gamma} (f(x, Du) + f(x, D\varphi)) \end{aligned}$$

and, similarly,

$$\begin{aligned} f(x, Du + (t-1)D\varphi) &\leq tf(x, Du) + (1-t)f(x, Du - D\varphi) \\ &\leq f(x, Du) + f(x, Du - D\varphi) \\ &\leq 2^\gamma (f(x, Du) + f(x, -D\varphi)). \end{aligned}$$

We have so proved that

$$\left| \sum_{i=1}^n \sum_{\alpha=1}^m \frac{\partial f}{\partial \xi_i^\alpha}(x, Du + tD\varphi) \varphi_{x_i}^\alpha \right| \leq 2^{3\gamma} (f(x, Du) + f(x, D\varphi) + f(x, -D\varphi)) =: H(x).$$

Since $u, \varphi, -\varphi \in W^{1,\mathcal{F}}$, we conclude. \square

4. Proof of the main results

We first state a lemma useful for the proof of Theorem 2.1. In the statement, the functions λ, μ , and the exponents p, q, r , are the same of the statement of Theorem 2.1. Moreover, $x_0 \in \Omega$ and $0 < R_0 \leq 1$ are such that $B_{R_0} := B_{R_0}(x_0) \Subset \Omega$. Fixed $R \in (0, R_0]$, the function $\eta \in C_c^\infty(B_R(x_0))$ denotes a cut-off function satisfying

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ in } B_\rho(x_0), \quad |D\eta| \leq \frac{2}{R-\rho}, \tag{4.1}$$

where $0 < \rho < R$.

Lemma 4.1. *Let $u \in W_{loc}^{1, \frac{pr}{r+1}}(\Omega; \mathbb{R}^m)$ be a local minimizer of (2.1). Then for every $\beta \geq 0$*

$$\int_{B_R} \lambda(x)(|u|+1)^{p\beta} |Du|^p \eta^q dx \leq \frac{c_1}{(R-\rho)^q} \int_{B_R} \mu(x)(|u|+1)^{q+p\beta} dx \tag{4.2}$$

for some c_1 depending on n, m, p, q, γ , but independent of β, u, R and ρ .

Proof. We begin using Proposition 3.4, with a suitable test function.

Let us approximate the identity function $\text{id} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with an increasing sequence of C^1 functions $h_k : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, with the following properties:

$$h_k(t) = 0 \quad \forall t \in [0, \frac{1}{k}], \quad h_k(t) = k \quad \forall t \in [k+1, +\infty], \quad 0 \leq h'_k(t) \leq 2 \text{ in } [0, \infty). \tag{4.3}$$

Fix $k \in \mathbb{N}$ and $\beta \geq 0$. Let $\Phi_k^{(\beta)} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be the increasing function defined as follows

$$\Phi_k^{(\beta)}(t) := h_k(t^{p\beta}). \tag{4.4}$$

Define $\varphi_k^{(\beta)} : B_R(x_0) \rightarrow \mathbb{R}^m$,

$$\varphi_k^{(\beta)}(x) := \Phi_k^{(\beta)}(|u(x)|)u(x)[\eta(x)]^q. \tag{4.5}$$

We have that φ_k is in $W^{1,f}(B_R(x_0); \mathbb{R}^m)$, $\text{supp } \varphi \Subset B_R(x_0)$ (see the proof of Lemma 5.1 in [8]).

Let us consider the Euler's equation (3.2) with test function $\varphi_k^{(\beta)}$. From now on, we write φ_k and Φ_k in place of $\varphi_k^{(\beta)}$ and $\Phi_k^{(\beta)}$, respectively. We obtain

$$\begin{aligned} I_1 + I_2 &:= \sum_{j=1}^n \sum_{\alpha=1}^m \int_{B_R} \frac{\partial f}{\partial \xi_j^\alpha}(x, Du) u_{x_j}^\alpha \Phi_k(|u|) \eta^q dx \\ &+ \sum_{j=1}^n \sum_{\alpha, \beta=1}^m \int_{B_R} \frac{\partial f}{\partial \xi_j^\alpha}(x, Du) u^\alpha \frac{u^\beta}{|u|} u_{x_j}^\beta \Phi'_k(|u|) \eta^q dx \\ &\leq q \left| \sum_{j=1}^n \sum_{\alpha=1}^m \int_{B_R} \frac{\partial f}{\partial \xi_j^\alpha}(x, Du) \Phi_k(|u|) u^\alpha \eta^{q-1} \eta_{x_j} dx \right| =: I_3. \end{aligned} \tag{4.6}$$

Now, we separately estimate I_1, I_2, I_3 .

ESTIMATE OF I_1

As far as I_1 is concerned, we use that $f(x, \cdot)$ is convex. Thus,

$$I_1 \geq \int_{B_R} (f(x, Du) - f(x, 0)) \Phi_k(|u|) \eta^q dx.$$

Using (2.2), we get

$$I_1 \geq \int_{B_R} f(x, Du) \Phi_k(|u|) \eta^q dx - \int_{B_R} \mu \Phi_k(|u|) \eta^q dx. \tag{4.7}$$

ESTIMATE OF I_2

We claim that $I_2 \geq 0$. Indeed, by (2.4), $\frac{\partial f}{\partial \xi_j^\alpha}(x, Du) = g_t(x, |Du|) \frac{u_{x_j}^\alpha}{|Du|}$. Therefore

$$\sum_{j=1}^n \sum_{\alpha,\beta=1}^m \frac{\partial f}{\partial \xi_j^\alpha}(x, Du) u^\alpha u^\beta u_{x_j}^\beta = \sum_{j=1}^n g_t(x, |Du|) \frac{\left(\sum_{\alpha=1}^m u^\alpha u_{x_j}^\alpha\right)^2}{|Du|} \geq 0.$$

Thus, by the monotonicity of Φ_k we have

$$I_2 = \int_{B_R} \sum_{j=1}^n g_t(x, |Du|) \frac{\left(\sum_{\alpha=1}^m u^\alpha u_{x_j}^\alpha\right)^2}{|Du| |u|} \Phi'_k(|u|) \eta^q dx \geq 0. \tag{4.8}$$

ESTIMATE OF I_3

By (2.4) and (4.1) we have

$$I_3 \leq \frac{2mq}{R-\rho} \int_{A_{R^-} \cup A_{R^+}} g_t(x, |Du|) |u| \Phi_k(|u|) \eta^{q-1} dx, \tag{4.9}$$

where

$$A_{R^-} := B_R \cap \left\{ \eta \neq 0, |Du| \leq \frac{2mqL|u|}{\eta(R-\rho)} \right\}$$

and

$$A_{R^+} := B_R \cap \left\{ \eta \neq 0, |Du| > \frac{2mqL|u|}{\eta(R-\rho)} \right\}$$

with $L > 0$ to be chosen later.

By (2.2), (2.4) and the assumption $x \in A_{R^-}$ the following inequality follows:

$$g_t(x, |Du|) \frac{2mq|u|}{\eta(R-\rho)} \leq \frac{1}{L} g_t \left(x, \frac{2mqL|u|}{\eta(R-\rho)} \right) \frac{2mqL|u|}{\eta(R-\rho)},$$

and, by Lemma 3.3(iii), (2.4) and (2.2),

$$\frac{1}{L} g_t \left(x, \frac{2mqL|u|}{\eta(R-\rho)} \right) \frac{2mqL|u|}{\eta(R-\rho)} \leq \frac{\gamma}{L} g \left(x, \frac{2mqL|u|}{\eta(R-\rho)} \right) \leq \frac{\gamma}{L} \mu(x) \left\{ \left(\frac{2mqL|u|}{\eta(R-\rho)} \right)^q + 1 \right\}.$$

Therefore

$$\begin{aligned} & \frac{2mq}{R-\rho} \int_{A_{R^-}} g_t(x, |Du|) |u| \Phi_k(|u|) \eta^{q-1} dx \\ & \leq \gamma \left(\frac{2mq}{R-\rho} \right)^q \int_{B_R} \mu(x) L^{q-1} |u|^q \Phi_k(|u|) dx + \frac{\gamma}{L} \int_{B_R} \mu(x) \Phi_k(|u|) \eta^q dx. \end{aligned} \tag{4.10}$$

Let us now deal with A_{R^+} . For a.e. $x \in A_{R^+}$, by Lemma 3.3(iii), (2.4) and (2.2) it follows

$$g_t(x, |Du|) \frac{2mq|u|}{\eta(R-\rho)} \leq \frac{1}{L} g_t(x, |Du|) |Du| \leq \frac{\gamma}{L} f(x, Du),$$

thus

$$\frac{2mq}{R-\rho} \int_{A_R^+} g_t(x, |Du|) |u| \Phi_k(|u|) \eta^{q-1} dx \leq \frac{\gamma}{L} \int_{B_R} f(x, Du) \Phi_k(|u|) \eta^q dx. \tag{4.11}$$

By (4.9), (4.10) and (4.11) we obtain

$$\begin{aligned} I_3 &\leq \gamma \left(\frac{2mq}{R-\rho} \right)^q \int_{B_R} \mu(x) L^{q-1} |u|^q \Phi_k(|u|) dx \\ &\quad + \frac{\gamma}{L} \int_{B_R} f(x, Du) \Phi_k(|u|) \eta^q dx + \frac{\gamma}{L} \int_{B_R} \mu(x) \Phi_k(|u|) \eta^q dx. \end{aligned} \tag{4.12}$$

By (4.7), (4.8), (4.10) and (4.12) we get

$$\begin{aligned} \left(1 - \frac{\gamma}{L}\right) \int_{B_R} f(x, Du) \Phi_k(|u|) \eta^q dx &\leq \gamma \left(\frac{2mq}{R-\rho} \right)^q \int_{B_R} \mu(x) L^{q-1} |u|^q \Phi_k(|u|) dx \\ &\quad + \left(1 + \frac{\gamma}{L}\right) \int_{B_R} \mu(x) \Phi_k(|u|) \eta^q dx. \end{aligned}$$

Choosing $L = 2\gamma$ and using (2.2), we get

$$\begin{aligned} \int_{B_R} f(x, Du) \Phi_k(|u|) \eta^q dx &\leq (2\gamma)^q \left(\frac{2mq}{R-\rho} \right)^q \int_{B_R} \mu(x) |u|^q \Phi_k(|u|) dx \\ &\quad + 3 \int_{B_R} \mu(x) \Phi_k(|u|) dx. \end{aligned} \tag{4.13}$$

Inequalities (2.2) and (4.13) imply

$$\int_{B_R} \lambda(x) |Du|^p \Phi_k(u) \eta^q dx \leq \frac{c_0}{(R-\rho)^q} \int_{B_R} \mu(x) \{|u|^q + 1\} \Phi_k(|u|) dx,$$

where we also used $R_0 \leq 1$. We recall that $\Phi_k = \Phi_k^{(\beta)}$ and we explicitly notice that c_0 is independent of β , ρ and R . Using the monotone convergence theorem we let k go to $+\infty$ and by the definition of Φ we obtain

$$\begin{aligned} \int_{B_R} \lambda(x) |u|^{p\beta} |Du|^p \eta^q dx &\leq \frac{c_0}{(R-\rho)^q} \int_{B_R} \mu(x) \{|u|^{q+p\beta} + |u|^{p\beta}\} dx \\ &\leq \frac{2c_0}{(R-\rho)^q} \int_{B_R} \mu(x) (|u| + 1)^{q+p\beta} dx. \end{aligned}$$

In particular, if $\beta = 0$:

$$\int_{B_R} \lambda(x) |Du|^p \eta^q dx \leq \frac{2c_0}{(R-\rho)^q} \int_{B_R} \mu(x) (|u| + 1)^q dx.$$

Thus, (4.2) follows. \square

Proof of Theorem 2.1. Let $u \in W_{loc}^{1, \frac{pr}{r+1}}(\Omega; \mathbb{R}^m)$ be a local minimizer of (2.1). Consider $x_0 \in \Omega$ and $0 < R_0 \leq 1$, such that $B_{R_0} := B_{R_0}(x_0) \Subset \Omega$. Fix also $0 < \rho < R \leq R_0$ and consider a cut-off function η satisfying (4.1). We split the proof into two steps.

Step 1. First we prove that, if $\delta \geq 1$ and $|u|^\delta \in W^{1, \sigma}(B_R)$, then

$$\begin{aligned} \|(|u| + 1)^\delta\|_{L^{\sigma^*}(B_\rho)} &\leq c \frac{q + \delta}{(R-\rho)^{q/p}} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \\ &\quad \times \| |u| + 1 \|_{L^{qs'}(B_{R_0})}^{\frac{q-p}{p}} \|(|u| + 1)^\delta\|_{L^{qs'}(B_R)}. \end{aligned} \tag{4.14}$$

To prove the above inequality, we notice that for any $\beta := \delta - 1 \geq 0$ we have that

$$\begin{aligned} \int_{\Omega} |D((|u| + 1)^{\beta+1}\eta^q)|^p \lambda(x) dx &\leq \int_{\Omega} (q\eta^{q-1}|D\eta|)^p (|u| + 1)^{(\beta+1)p} \lambda(x) dx \\ &+ (\beta + 1)^p \int_{\Omega} \lambda(x)\eta^{qp}(|u| + 1)^{p\beta}|Du|^p dx. =: J_1 + J_2. \end{aligned} \tag{4.15}$$

To estimate J_1 we observe that

$$J_1 \leq c \frac{q^p}{(R - \rho)^p} \int_{B_R} (|u| + 1)^{(\beta+1)p} \lambda(x) dx \leq c \frac{q^p}{(R - \rho)^p} \int_{B_R} (|u| + 1)^{q+\beta p} \lambda(x) dx. \tag{4.16}$$

Since $\eta^{qp} \leq \eta^q$, we can estimate J_2 using Lemma 4.1. Thus,

$$J_2 \leq c \frac{(\beta + 1)^p}{(R - \rho)^q} \int_{B_R} \mu(x)(|u| + 1)^{q+p\beta} dx. \tag{4.17}$$

By Lemma 3.2 applied to $v = (|u| + 1)^{\beta+1}\eta^q$ the inequality (3.1) holds, that is

$$\left\{ \int_{\Omega} ((|u| + 1)^{\beta+1}\eta^q)^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} \leq c \|\lambda^{-1}\|_{L^r(B_{R_0})} \int_{\Omega} \lambda(x) |D((|u| + 1)^{\beta+1}\eta^q)|^p dx. \tag{4.18}$$

Collecting this inequality, (4.15), (4.16) and (4.17), we get

$$\begin{aligned} \left\{ \int_{\Omega} ((|u| + 1)^{\beta+1}\eta^q)^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} &\leq c \|\lambda^{-1}\|_{L^r(B_{R_0})} \frac{q^p}{(R - \rho)^p} \int_{B_R} \lambda(x)(|u| + 1)^{q+\beta p} dx \\ &+ c \|\lambda^{-1}\|_{L^r(B_{R_0})} \frac{(\beta + 1)^p}{(R - \rho)^q} \int_{B_R} \mu(x)(|u| + 1)^{q+p\beta} dx. \end{aligned}$$

By (2.2) $\lambda \leq 2\mu$ a.e., and $R_0 \leq 1$, so we get

$$\left\{ \int_{\Omega} ((|u| + 1)^{\beta+1}\eta^q)^{\sigma^*} dx \right\}^{\frac{p}{\sigma^*}} \leq c \|\lambda^{-1}\|_{L^r(B_{R_0})} \frac{q^p + (\beta + 1)^p}{(R - \rho)^q} \int_{B_R} \mu(x)(|u| + 1)^{q+\beta p} dx. \tag{4.19}$$

By Hölder inequality and $\mu \in L^s_{loc}(\Omega)$ we obtain

$$\int_{B_R} \mu(x)(|u| + 1)^{q+\beta p} dx \leq \|\mu\|_{L^s(B_{R_0})} \left(\int_{B_R} (|u| + 1)^{(q+\beta p)s'} dx \right)^{\frac{1}{s'}}. \tag{4.20}$$

If $p = q$, (4.19) and (4.20) give

$$\|(|u| + 1)^{\beta+1}\|_{L^{\sigma^*}(B_{\rho})} \leq c \frac{q + \beta}{R - \rho} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \|(|u| + 1)^{\beta+1}\|_{L^{ps'}(B_R)},$$

that is (4.14) holds for $q = p$ and $\delta = \beta + 1$.

If $p < q$, let us apply Hölder inequality in the last integral in (4.20):

$$\begin{aligned} \left(\int_{B_R} (|u| + 1)^{(q+\beta p)s'} dx \right)^{\frac{1}{s'}} &= \left(\int_{B_R} (|u| + 1)^{(q-p)s'} (|u| + 1)^{(\beta+1)ps'} dx \right)^{\frac{1}{s'}} \\ &\leq \left(\int_{B_R} (|u| + 1)^{qs'} dx \right)^{\frac{q-p}{qs'}} \left(\int_{B_R} (|u| + 1)^{(\beta+1)qs'} dx \right)^{\frac{p}{qs'}}. \end{aligned} \tag{4.21}$$

Then, (4.19), (4.20) and (4.21) give

$$\begin{aligned} \|(|u| + 1)^{\beta+1}\|_{L^{\sigma^*}(B_{\rho})} &\leq c \frac{q + \beta + 1}{(R - \rho)^{q/p}} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \\ &\times \| |u| + 1 \|_{L^{qs'}(B_{R_0})}^{\frac{q-p}{p}} \|(|u| + 1)^{\beta+1}\|_{L^{qs'}(B_R)}. \end{aligned}$$

Since $\delta = \beta + 1$, we get that (4.14) holds also for $p < q$.

Step 2. Let us define $G(x) := \max\{1, |u(x)|\}$. Now, we prove the boundedness of G , and then of u , using the Moser’s iteration technique. The inequality (4.14) implies that for any $\delta \geq 1$,

$$\|G^\delta\|_{L^{\sigma^*}(B_\rho)} \leq c \frac{q + \delta}{(R - \rho)^{q/p}} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \|G\|_{L^{qs'}(B_{R_0})}^{\frac{q-p}{p}} \|G^\delta\|_{L^{qs'}(B_R)}. \tag{4.22}$$

Now, we prove the boundedness of G , and then of u , using the Moser’s iteration technique. For all $h \in \mathbb{N}$ define $\delta_h = \left(\frac{\sigma^*}{qs'}\right)^{h-1}$, $R_h = R_0/2 + R_0/2^h$ and $\rho_h = R_{h+1}$. Notice that the choice of δ_h has been done in such a way that $\delta_1 = 1$ and $\delta_h \sigma^* = \delta_{h+1} qs'$. By (4.22), replacing δ , R and ρ with δ_h , R_h and ρ_h , respectively, we have that $G \in L^{\delta_h qs'}(B_{R_h})$ implies $G \in L^{\delta_{h+1} qs'}(B_{R_{h+1}})$. Precisely,

$$\begin{aligned} \|G\|_{L^{\delta_{h+1} qs'}(B_{R_{h+1}})} &= \|G\|_{L^{\delta_h \sigma^*}(B_{R_{h+1}})} = \|G^{\delta_h}\|_{L^{\sigma^*}(B_{R_{h+1}})}^{\frac{1}{\delta_h}} \\ &\leq \left\{ c \frac{q + \delta_h}{(R_h - \rho_h)^{q/p}} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \|G\|_{L^{qs'}(B_{R_0})}^{\frac{q-p}{p}} \right\}^{\frac{1}{\delta_h}} \|G\|_{L^{\delta_h qs'}(B_{R_h})} \end{aligned} \tag{4.23}$$

holds true for every h . For instance, if $h = 1$, we get

$$\|G\|_{L^{\sigma^*}(B_{3R_0/4})} \leq c \frac{q + 1}{R_0^{q/p}} \|\lambda^{-1}\|_{L^r(B_{R_0})}^{\frac{1}{p}} \|\mu\|_{L^s(B_{R_0})}^{\frac{1}{p}} \|G\|_{L^{qs'}(B_{R_0})}^{\frac{q-p}{p}} \|G\|_{L^{qs'}(B_{R_0})}.$$

Notice that the right hand side is finite, because $u \in W^{1,\sigma}$, so $u \in L^{\sigma^*}$ and, by (2.8), $qs' < \sigma^*$. Since

$$\sum_{h=1}^{\infty} \frac{1}{\delta_h} = \sum_{i=0}^{\infty} \left(\frac{qs'}{\sigma^*}\right)^i = \frac{\sigma^*}{\sigma^* - qs'},$$

an iterated use of (4.23) implies the existence of a constant c such that

$$\|G\|_{L^\infty(B_{R_0/2}(x_0))} \leq c \left(\frac{\|\lambda^{-1}\|_{L^r(B_{R_0})} \|\mu\|_{L^s(B_{R_0})}}{R_0^q} \right)^{\frac{\sigma^*}{p(\sigma^* - qs')}} \|G\|_{L^{qs'}(B_{R_0}(x_0))}^{\frac{\sigma^*(q-p)}{p(\sigma^* - qs')}} \|G\|_{L^{qs'}(B_{R_0}(x_0))}.$$

The inequality above implies that u is in $L^\infty(B_{R_0/2}(x_0); \mathbb{R}^m)$ and the estimate (2.7). \square

Acknowledgment

The authors are members of Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

[1] P. Baroni, M. Colombo, G. Mingione, Harnack inequalities for double phase functionals, *Nonlinear Anal.* 121 (2015) 206–222.
 [2] M. Carozza, J. Kristensen, A. Passarelli di Napoli, Regularity of minimizers of autonomous convex variational integrals, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 13 (2014) 1065–1089.
 [3] M. Colombo, G. Mingione, Bounded minimisers of double phase variational integrals, *Arch. Ration. Mech. Anal.* 218 (2015) 219–273.
 [4] M. Colombo, G. Mingione, Regularity for double phase variational problems, *Arch. Ration. Mech. Anal.* 215 (2015) 443–496.
 [5] D. Cruz-Uribe, P. Di Gironimo, C. Sbordone, On the continuity of solutions to degenerate elliptic equations, *J. Differential Equations* 250 (2011) 2671–2686.
 [6] G. Cupini, P. Marcellini, E. Mascolo, Regularity under sharp anisotropic general growth conditions, *Discrete Contin. Dyn. Syst. Ser. B* 11 (2009) 66–86.
 [7] G. Cupini, P. Marcellini, E. Mascolo, Local boundedness of solutions to quasilinear elliptic systems, *Manuscripta Math.* 137 (2012) 287–315.

- [8] G. Cupini, P. Marcellini, E. Mascolo, Local boundedness of solutions to some anisotropic elliptic systems, in: *Recent Trends in Nonlinear Partial Differential Equations. II. Stationary Problems*, in: *Contemp. Math.*, vol. 595, Amer. Math. Soc., Providence, RI, 2013, pp. 169–186.
- [9] G. Cupini, P. Marcellini, E. Mascolo, Existence and regularity for elliptic equations under p, q -growth, *Adv. Differential Equations* 19 (2014) 693–724.
- [10] G. Cupini, P. Marcellini, E. Mascolo, Regularity of minimizers under limit growth conditions, *Nonlinear Anal.* 153 (2017) 294–310.
- [11] A. Dall’Aglio, E. Mascolo, L^∞ estimates for a class of nonlinear elliptic systems with nonstandard growth, *Atti Semin. Mat. Fis. Univ. Modena* 50 (2002) 65–83.
- [12] C. De Filippis, Higher integrability for constrained minimizers of integral functionals with (p, q) -growth in low dimension, *Nonlinear Anal.* 170 (2018) 1–20.
- [13] E. De Giorgi, Un esempio di estremali discontinue per un problema variazionale di tipo ellittico, *Boll. Univ. Mat. Ital.*(4) 1 (1968) 135–137.
- [14] M. Eleuteri, P. Marcellini, E. Mascolo, Lipschitz continuity for energy integrals with variable exponents, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* 27 (2016) 61–87.
- [15] M. Eleuteri, P. Marcellini, E. Mascolo, Lipschitz estimates for systems with ellipticity conditions at infinity, *Ann. Mat. Pura E Appl.* (4) 195 (2016) 1575–1603.
- [16] M. Eleuteri, P. Marcellini, E. Mascolo, Regularity for scalar integrals without structure conditions, *Adv. Calc. Var.* (2018). <http://dx.doi.org/10.1515/acv-2017-0037>. (in press).
- [17] N. Fusco, C. Sbordone, Some remarks on the regularity of minima of anisotropic integrals, *Comm. Partial Differential Equations* 18 (1993) 153–167.
- [18] F. Giannetti, A. Passarelli di Napoli, Higher differentiability of minimizers of variational integrals with variable exponents, *Math. Z.* 280 (2015) 873–892.
- [19] E. Giusti, M. Miranda, Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni, *Boll. Unione Mat. Ital.* 2 (1968) 1–8.
- [20] F. Leonetti, P.V. Petricca, Bounds for vector valued minimizers of some relaxed functionals, *Complex Var. Elliptic Equ.* 58 (2013) 221–230.
- [21] P. Marcellini, C. Sbordone, An approach to the asymptotic behaviour of elliptic–parabolic operators, *J. Math. Pures Appl.* 56 (1977) 157–182.
- [22] P. Marcellini, C. Sbordone, Homogenization of nonuniformly elliptic operators, *Appl. Anal.* 8 (1978) 101–113.
- [23] G. Mingione, Regularity of minima: an invitation to the dark side of the calculus of variations, *Appl. Math.* 51 (2006) 355–426.
- [24] V. Rădulescu, Q. Zhang, Double phase anisotropic variational problems and combined effects of reaction and absorption terms, *J. Math. Pures Appl.* (2018) (in press).
- [25] V. Šverák, X. Yan, A singular minimizer of a smooth strongly convex functional in three dimensions, *Calc. Var. Partial Differential Equations* 10 (2000) 213–221.
- [26] N.S. Trudinger, On the regularity of generalized solutions of linear, nonuniformly elliptic equations, *Arch. Ration. Mech. Anal.* 42 (1971) 50–62.
- [27] V.V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.* 50 (1986) 675–710.