

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Uniqueness and decay in local thermal non-equilibrium double porosity thermoelasticity

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Uniqueness and decay in local thermal non-equilibrium double porosity thermoelasticity / Franca Franchi, Barbara Lazzari, Roberta Nibbi, Brian Straughan. - In: MATHEMATICAL METHODS IN THE APPLIED SCIENCES. - ISSN 1099-1476. - STAMPA. - 41:16(2018), pp. 6763-6771. [10.1002/mma.5190]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/654348> since: 2021-02-26

*Published:*

DOI: <http://doi.org/10.1002/mma.5190>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**F. Franchi; B. Lazzari; R. Nibbi; B. Straughan – *Uniqueness and decay in local thermal non-equilibrium double porosity thermoelasticity* –**

**Mathematical Methods in the Applied Sciences, vol. 41 (2018), pp. 6763—6771.**

The final published version is available online at: <https://doi.org/10.1002/mma.5190>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

**RESEARCH ARTICLE**

# Uniqueness and decay in local thermal non-equilibrium double porosity thermoelasticity

Franca Franchi<sup>1</sup> | Barbara Lazzari\*<sup>1</sup> | Roberta Nibbi<sup>1</sup> | Brian Straughan<sup>2</sup>

<sup>1</sup>Dipartimento di Matematica, Università di Bologna, Italy, Piazza di porta San Donato 5, 40126 Bologna

<sup>2</sup>Department of Mathematical Sciences, Durham University, UK, DH1 3LE

**Correspondence**

\*B. Lazzari, Dipartimento di Matematica, Piazza di porta San Donato 5, 40126 Bologna Italy, Email: barbara.lazzari@unibo.it

**Abstract**

This paper studies a model for thermoelasticity where the body has a double porosity structure. There are the usual pores associated to a porous body, herein called macro pores. In addition, the solid skeleton contains cracks or fissures which give rise to a micro porosity. The fully anisotropic situation is analyzed. We firstly establish uniqueness of a solution to the boundary-initial value problem when the elastic coefficients are sign-indefinite and are required to satisfy only major symmetry. Furthermore, in the quasi-equilibrium case, where the solid acceleration is neglected, we demonstrate that a solution to the boundary-initial value problem with zero boundary conditions will decay to zero in a certain sense, under the assumption that there are no sources and external body force involved.

**KEYWORDS:**

double porosity, thermoelasticity, local-thermal non equilibrium, uniqueness, decay

## 1 | INTRODUCTION

Driven by the many real life applications, elastic materials with a double porosity structure have become the subject of intensive current research.

A double porosity elastic material is a solid which contains pores on a macro scale and pores on a much smaller scale. These are referred to as macro pores and micro pores, respectively. There are many application areas and these are detailed in Straughan<sup>1</sup>, but to give an idea of the vastness we mention oil recovery, Olusola et al.<sup>2</sup>; hydraulic fracturing for gas, Kim and Moridis<sup>3</sup>; bone recovery and bone replacement, Svanadze and Scalia<sup>4</sup>.

In this article we focus on an anisotropic theory for a double porosity elastic material under the effect of local thermal non-equilibrium, cf. Svanadze<sup>5</sup>. Local thermal non-equilibrium is where the solid and the fluid in a porous body may have different temperatures. In studies of fluid motion in a fixed porous body this concept was introduced by Nield<sup>6</sup> and striking differences have been observed by comparison with the single temperature theory, see e.g. Rees et al.<sup>7</sup>, see also Straughan<sup>8</sup>.

The basic double porosity theory we use is based on a single temperature model developed extensively by Svanadze<sup>9,10</sup>, Ciarletta et al.<sup>11</sup>, Svanadze and Scalia<sup>4</sup>, Scarpetta and Svanadze<sup>12</sup>, and within the local thermal non-equilibrium isotropic theory developed and used in Svanadze<sup>5</sup>. Thermal effects in porous elastic material are very important since cracking may occur due to thermally induced stress, see e.g. Siratovich et al.<sup>13</sup>. Thermally induced cracking in a porous elastic body is thus intimately connected to a double porosity theory.

In the first part of the paper we establish uniqueness of a solution without requiring any sign-definiteness of the elastic coefficients. We require only symmetry of elasticities. This is important since, for example, Xinchun and Lakes<sup>14</sup> and Ha et al.<sup>15</sup> have demonstrated that many materials have negative Poisson ratio. In fact, recent studies have abandoned positive-definiteness

of the elastic coefficients in favour of the weaker condition of strong ellipticity, see e.g. Xinchun and Lakes<sup>14</sup>, Chiriță and Ciarletta<sup>16</sup>, Chiriță et al.<sup>17</sup>, Chiriță and Danescu<sup>18</sup>, Chiriță and Ghiba<sup>19</sup>. Thus, for an anisotropic body use of only the major symmetry is important.

In the quasi-equilibrium case where the solid acceleration is absent, see e.g. Scarpetta and Svanadze<sup>12</sup>, the decay in time in a certain sense is established for a solution of the double porosity local thermal non-equilibrium problem, under the assumption that there are no sources and external body force involved.

## 2 | THE GOVERNING EQUATIONS

The porous elastic body consists of a solid elastic skeleton, macro pores, and micro pores in the skeleton. The temperature is allowed to be different in the skeleton, in the fluid in the macro pores and in the fluid in the micro pores. The relevant variables are the elastic displacement,  $u_i$ , the pressure in the macro pores,  $p$ , the pressure in the micro pores,  $q$ , the temperature in the solid skeleton,  $\theta_1$ , the temperature of the fluid in the macro pores,  $\theta_2$ , and the temperature of the fluid in the micro pores,  $\theta_3$ . Each variable is a function of  $\mathbf{x}$  and  $t$  and the elastic body occupies a bounded domain  $\Omega \subset \mathbb{R}^3$ , which has boundary sufficiently smooth to allow application of the divergence theorem.

The basic equations for a double porosity thermoelastic body in local thermal non-equilibrium are derived in the isotropic case by Svanadze<sup>5</sup>. We here generalize this and present the fully anisotropic system of equations, namely

$$\begin{aligned} \rho \ddot{u}_i &= (a_{ijhk} u_{h,k})_{,j} - (\beta_{ij} p)_{,j} - (\gamma_{ij} q)_{,j} \\ &\quad - (a_{ij}^{(1)} \theta_1)_{,j} - (a_{ij}^{(2)} \theta_2)_{,j} - (a_{ij}^{(3)} \theta_3)_{,j} + \rho f_i, \\ \alpha \dot{p} &= (k_{ij} p_{,j})_{,i} - \beta_{ij} \dot{u}_{i,j} - \gamma(p - q) + \rho s_1, \\ \beta \dot{q} &= (m_{ij} q_{,j})_{,i} - \gamma_{ij} \dot{u}_{i,j} + \gamma(p - q) + \rho s_2, \\ a_1 \dot{\theta}_1 &= (\kappa_{ij} \theta_{1,j})_{,i} - a_{ij}^{(1)} \dot{u}_{i,j} - d_1(\theta_1 - \theta_2) - d_2(\theta_1 - \theta_3) + \rho r_1, \\ a_2 \dot{\theta}_2 &= (\omega_{ij} \theta_{2,j})_{,i} - a_{ij}^{(2)} \dot{u}_{i,j} + d_1(\theta_1 - \theta_2) - d_3(\theta_2 - \theta_3) + \rho r_2, \\ a_3 \dot{\theta}_3 &= (\pi_{ij} \theta_{3,j})_{,i} - a_{ij}^{(3)} \dot{u}_{i,j} + d_3(\theta_2 - \theta_3) + d_2(\theta_1 - \theta_3) + \rho r_3, \end{aligned} \quad (1)$$

where a superposed dot denotes  $\partial/\partial t$ , standard indicial notation is employed with  $_{,i}$  denoting  $\partial/\partial x_i$ , and  $f_i, s_1, s_2, r_1, r_2, r_3$  are the body force, sources for the pressure and externally supplied heat sources, respectively. The coefficients may all depend on  $\mathbf{x}$ , with  $\rho > 0$  being the density of the solid elastic skeleton and also  $\alpha, \beta, \gamma, d_1, d_2, d_3, a_1, a_2$  and  $a_3$  are positive. The tensors  $\beta_{ij}, \gamma_{ij}, a_{ij}^{(1)}, a_{ij}^{(2)}, a_{ij}^{(3)}$  are all symmetric. Likewise the coefficients  $k_{ij}, m_{ij}, \kappa_{ij}, \omega_{ij}$  and  $\pi_{ij}$  are all symmetric tensors.

Equations (1) hold on  $\Omega \times (0, T]$  for some  $T < \infty$ . The boundary of  $\Omega$  is denoted by  $\Gamma$  and on  $\Gamma$  the conditions are

$$\begin{aligned} u_i &= h_i(\mathbf{x}, t), \quad p = p^B(\mathbf{x}, t), \quad q = q^B(\mathbf{x}, t), \\ \theta_1 &= \alpha_1(\mathbf{x}, t), \quad \theta_2 = \alpha_2(\mathbf{x}, t), \quad \theta_3 = \alpha_3(\mathbf{x}, t), \end{aligned} \quad (2)$$

holding on  $\Gamma \times [0, T]$ , where  $h_i, p^B, q^B, \alpha_1, \alpha_2, \alpha_3$  are prescribed functions. In addition, the following initial conditions hold

$$\begin{aligned} u_i(\mathbf{x}, 0) &= v_i(\mathbf{x}), \quad \dot{u}_i(\mathbf{x}, 0) = w_i(\mathbf{x}), \\ p(\mathbf{x}, 0) &= P(\mathbf{x}), \quad q(\mathbf{x}, 0) = Q(\mathbf{x}), \\ \theta_1(\mathbf{x}, 0) &= \Theta_1(\mathbf{x}), \quad \theta_2(\mathbf{x}, 0) = \Theta_2(\mathbf{x}), \quad \theta_3(\mathbf{x}, 0) = \Theta_3(\mathbf{x}), \end{aligned} \quad (3)$$

with  $\mathbf{x} \in \Omega$  and for given functions  $v_i, w_i, P, Q$  and  $\Theta_\alpha, \alpha = 1, 2, 3$ .

The boundary-initial value problem given by (1)–(3) is denoted by  $\mathcal{P}$ .

## 3 | UNIQUENESS

In this section we require the elastic coefficients to satisfy the symmetries

$$a_{ijhk} = a_{jihk} = a_{hkij}, \quad (4)$$

although we stress that no definiteness is required, and we require that  $k_{ij}$ ,  $m_{ij}$ ,  $\kappa_{ij}$ ,  $\omega_{ij}$  and  $\pi_{ij}$  are semi-positive-definite.

To establish uniqueness we let  $(u_i^\kappa, p^\kappa, q^\kappa, \theta_1^\kappa, \theta_2^\kappa, \theta_3^\kappa)$ ,  $\kappa = 1, 2$  be two solutions to  $\mathcal{P}$  which satisfy (2) and (3) for the same boundary and initial functions  $h_i, p^B, q^B, \alpha_1, \alpha_2, \alpha_3, v_i, w_i, P, Q, \Theta_1, \Theta_2, \Theta_3$ , and which satisfy (1) for the same body force  $f_i$  and for the same source terms  $s_1, s_2, r_1, r_2, r_3$ . Define the difference solution  $\Lambda = (u_i, p, q, \theta_1, \theta_2, \theta_3)$  by

$$\begin{aligned} u_i &= u_i^1 - u_i^2, & p &= p^1 - p^2, & q &= q^1 - q^2, \\ \theta_1 &= \theta_1^1 - \theta_1^2, & \theta_2 &= \theta_2^1 - \theta_2^2, & \theta_3 &= \theta_3^1 - \theta_3^2. \end{aligned}$$

By subtraction one verifies that  $\Lambda$  satisfies the equations

$$\begin{aligned} \rho \ddot{u}_i &= (a_{ijhk} u_{h,k})_{,j} - (\beta_{ij} p)_{,j} - (\gamma_{ij} q)_{,j} \\ &\quad - (a_{ij}^{(1)} \theta_1)_{,j} - (a_{ij}^{(2)} \theta_2)_{,j} - (a_{ij}^{(3)} \theta_3)_{,j}, \\ \alpha \dot{p} &= (k_{ij} p_{,j})_{,i} - \beta_{ij} \dot{u}_{i,j} - \gamma(p - q), \\ \beta \dot{q} &= (m_{ij} q_{,j})_{,i} - \gamma_{ij} \dot{u}_{i,j} + \gamma(p - q), \\ a_1 \dot{\theta}_1 &= (\kappa_{ij} \theta_{1,j})_{,i} - a_{ij}^{(1)} \dot{u}_{i,j} - d_1(\theta_1 - \theta_2) - d_2(\theta_1 - \theta_3), \\ a_2 \dot{\theta}_2 &= (\omega_{ij} \theta_{2,j})_{,i} - a_{ij}^{(2)} \dot{u}_{i,j} + d_1(\theta_1 - \theta_2) - d_3(\theta_2 - \theta_3), \\ a_3 \dot{\theta}_3 &= (\pi_{ij} \theta_{3,j})_{,i} - a_{ij}^{(3)} \dot{u}_{i,j} + d_3(\theta_2 - \theta_3) + d_2(\theta_1 - \theta_3), \end{aligned} \tag{5}$$

in  $\Omega \times (0, T]$ , together with the homogeneous boundary conditions

$$\begin{aligned} u_i &= 0, & p &= 0, & q &= 0, \\ \theta_1 &= 0, & \theta_2 &= 0, & \theta_3 &= 0, \end{aligned} \tag{6}$$

on  $\Gamma \times [0, T]$ , and the homogeneous initial conditions

$$\begin{aligned} u_i(\mathbf{x}, 0) &= 0, & \dot{u}_i(\mathbf{x}, 0) &= 0, \\ p(\mathbf{x}, 0) &= 0, & q(\mathbf{x}, 0) &= 0, \\ \theta_1(\mathbf{x}, 0) &= 0, & \theta_2(\mathbf{x}, 0) &= 0, & \theta_3(\mathbf{x}, 0) &= 0, \end{aligned} \tag{7}$$

for  $\mathbf{x} \in \Omega$ .

The uniqueness proof proceeds by a logarithmic convexity argument.

We firstly define the variables  $\eta(\mathbf{x}, t)$ ,  $\zeta(\mathbf{x}, t)$ ,  $\phi(\mathbf{x}, t)$ ,  $\psi(\mathbf{x}, t)$ ,  $\xi(\mathbf{x}, t)$  by

$$\begin{aligned} \eta &= \int_0^t p(\mathbf{x}, s) ds, & \zeta &= \int_0^t q(\mathbf{x}, s) ds, \\ \phi &= \int_0^t \theta_1(\mathbf{x}, s) ds, & \psi &= \int_0^t \theta_2(\mathbf{x}, s) ds, & \xi &= \int_0^t \theta_3(\mathbf{x}, s) ds. \end{aligned}$$

In terms of the variables  $\eta, \zeta, \phi, \psi$  and  $\xi$  we may integrate (5)<sub>2-6</sub> to derive the equations

$$\begin{aligned} \alpha p &= (k_{ij} \eta_{,j})_{,i} - \beta_{ij} u_{i,j} - \gamma(\eta - \zeta), \\ \beta q &= (m_{ij} \zeta_{,j})_{,i} - \gamma_{ij} u_{i,j} + \gamma(\eta - \zeta), \\ a_1 \theta_1 &= (\kappa_{ij} \phi_{,j})_{,i} - a_{ij}^{(1)} u_{i,j} - d_1(\phi - \psi) - d_2(\phi - \xi), \\ a_2 \theta_2 &= (\omega_{ij} \psi_{,j})_{,i} - a_{ij}^{(2)} u_{i,j} + d_1(\phi - \psi) - d_3(\psi - \xi), \\ a_3 \theta_3 &= (\pi_{ij} \xi_{,j})_{,i} - a_{ij}^{(3)} u_{i,j} + d_3(\psi - \xi) + d_2(\phi - \xi). \end{aligned} \tag{8}$$

We denote by  $\langle \cdot \rangle$  integration over  $\Omega$ , e.g.

$$\langle f \rangle = \int_{\Omega} f dx.$$

Then define the functional  $F$  by

$$\begin{aligned}
F(t) = & \langle \rho u_i \dot{u}_i \rangle + \int_0^t \langle k_{ij} \eta_j \eta_i \rangle ds + \int_0^t \langle m_{ij} \zeta_j \zeta_i \rangle ds \\
& + \int_0^t \langle \gamma (\eta - \zeta)^2 \rangle ds + \int_0^t \langle \kappa_{ij} \phi_j \phi_i \rangle ds \\
& + \int_0^t \langle \omega_{ij} \psi_j \psi_i \rangle ds + \int_0^t \langle \pi_{ij} \xi_j \xi_i \rangle ds \\
& + \int_0^t \langle d_1 (\phi - \psi)^2 \rangle ds + \int_0^t \langle d_2 (\phi - \xi)^2 \rangle ds \\
& + \int_0^t \langle d_3 (\psi - \xi)^2 \rangle ds.
\end{aligned} \tag{9}$$

Differentiate  $F$  to obtain

$$\begin{aligned}
F'(t) = & 2 \langle \rho u_i \dot{u}_i \rangle + 2 \int_0^t \langle k_{ij} \eta_j p_i \rangle ds + 2 \int_0^t \langle m_{ij} \zeta_j q_i \rangle ds \\
& + 2 \int_0^t \langle \gamma (\eta - \zeta)(p - q) \rangle ds + 2 \int_0^t \langle \kappa_{ij} \phi_j \theta_{1,i} \rangle ds \\
& + 2 \int_0^t \langle \omega_{ij} \psi_j \theta_{2,i} \rangle ds + 2 \int_0^t \langle \pi_{ij} \xi_j \theta_{3,i} \rangle ds \\
& + 2 \int_0^t \langle d_1 (\phi - \psi)(\theta_1 - \theta_2) \rangle ds + 2 \int_0^t \langle d_2 (\phi - \xi)(\theta_1 - \theta_3) \rangle ds + 2 \int_0^t \langle d_3 (\psi - \xi)(\theta_2 - \theta_3) \rangle ds.
\end{aligned} \tag{10}$$

After a further differentiation one finds

$$\begin{aligned}
F''(t) = & 2 \langle \rho \dot{u}_i \dot{u}_i \rangle + 2 \langle \rho u_i \ddot{u}_i \rangle + 2 \langle k_{ij} \eta_j p_i \rangle + 2 \langle m_{ij} \zeta_j q_i \rangle \\
& + 2 \langle \gamma (\eta - \zeta)(p - q) \rangle + 2 \langle \kappa_{ij} \phi_j \theta_{1,i} \rangle + 2 \langle \omega_{ij} \psi_j \theta_{2,i} \rangle \\
& + 2 \langle \pi_{ij} \xi_j \theta_{3,i} \rangle + 2 \langle d_1 (\phi - \psi)(\theta_1 - \theta_2) \rangle \\
& + 2 \langle d_2 (\phi - \xi)(\theta_1 - \theta_3) \rangle + 2 \langle d_3 (\psi - \xi)(\theta_2 - \theta_3) \rangle.
\end{aligned} \tag{11}$$

The next step is to multiply equation (5)<sub>1</sub> by  $u_i$  and integrate over  $\Omega$ . Then multiply each of equations (8)<sub>1</sub> to (8)<sub>5</sub> by  $p$ ,  $q$ ,  $\theta_1$ ,  $\theta_2$  and  $\theta_3$ , in turn, and integrate each over  $\Omega$  with integration by parts and use of the homogeneous boundary conditions. The resulting six equations are added to yield the following equation

$$\begin{aligned}
& \langle \rho u_i \ddot{u}_i \rangle + \langle k_{ij} \eta_j p_i \rangle + \langle m_{ij} \zeta_j q_i \rangle + \langle \kappa_{ij} \phi_j \theta_{1,i} \rangle \\
& \langle \omega_{ij} \psi_j \theta_{2,i} \rangle + \langle \pi_{ij} \xi_j \theta_{3,i} \rangle + \langle \gamma (\eta - \zeta)(p - q) \rangle \\
& + \langle d_1 (\phi - \psi)(\theta_1 - \theta_2) \rangle + \langle d_3 (\psi - \xi)(\theta_2 - \theta_3) \rangle \\
& + \langle d_2 (\phi - \xi)(\theta_1 - \theta_3) \rangle \\
= & - \langle a_{ijhk} u_{h,k} u_{i,j} \rangle - \langle \alpha p^2 \rangle - \langle \beta q^2 \rangle \\
& - \langle a_1 \theta_1^2 \rangle - \langle a_2 \theta_2^2 \rangle - \langle a_3 \theta_3^2 \rangle.
\end{aligned} \tag{12}$$

We now use equation (12) to substitute for the last ten terms in (11) to find

$$\begin{aligned}
F''(t) = & 2 \langle \rho \dot{u}_i \dot{u}_i \rangle - 2 \langle a_{ijhk} u_{h,k} u_{i,j} \rangle - 2 \langle \alpha p^2 \rangle - 2 \langle \beta q^2 \rangle \\
& - 2 \langle a_1 \theta_1^2 \rangle - 2 \langle a_2 \theta_2^2 \rangle - 2 \langle a_3 \theta_3^2 \rangle.
\end{aligned} \tag{13}$$

The next stage involves the energy equation for (5)–(7). Define the energy as

$$E(t) = \frac{1}{2} \langle \rho \dot{u}_i \dot{u}_i \rangle + \frac{1}{2} \langle a_{ijhk} u_{i,j} u_{h,k} \rangle + \frac{1}{2} \langle \alpha p^2 \rangle + \frac{1}{2} \langle \beta q^2 \rangle + \frac{1}{2} \langle a_1 \theta_1^2 \rangle + \frac{1}{2} \langle a_2 \theta_2^2 \rangle + \frac{1}{2} \langle a_3 \theta_3^2 \rangle . \quad (14)$$

To obtain the energy equation, one multiplies (5)<sub>1</sub> by  $\dot{u}_i$ , (5)<sub>2–6</sub> by  $p$ ,  $q$ ,  $\theta_1$ ,  $\theta_2$  and  $\theta_3$ , respectively, and integrates each over  $\Omega$ . Integration by parts, taking into account the homogeneous boundary conditions, and then integrating in time, readily yields

$$\begin{aligned} E(t) &+ \int_0^t \langle k_{ij} p_{,j} p_{,i} \rangle ds + \int_0^t \langle m_{ij} q_{,j} q_{,i} \rangle ds + \int_0^t \langle \gamma (p - q)^2 \rangle ds \\ &+ \int_0^t \langle \kappa_{ij} \theta_{1,j} \theta_{1,i} \rangle ds + \int_0^t \langle \omega_{ij} \theta_{2,j} \theta_{2,i} \rangle ds + \int_0^t \langle \pi_{ij} \theta_{3,j} \theta_{3,i} \rangle ds \\ &+ \int_0^t \langle d_1 (\theta_1 - \theta_2)^2 \rangle ds + \int_0^t \langle d_2 (\theta_1 - \theta_3)^2 \rangle ds + \int_0^t \langle d_3 (\theta_2 - \theta_3)^2 \rangle ds = E(0) . \end{aligned} \quad (15)$$

From (7) it follows that  $E(0) = 0$ .

Substitute in (13) for the last six terms using (14) and (15) to see that

$$\begin{aligned} F''(t) &= 4 \langle \rho \dot{u}_i \dot{u}_i \rangle - 4E(t) \\ &= 4 \langle \rho \dot{u}_i \dot{u}_i \rangle + 4 \int_0^t \langle k_{ij} p_{,j} p_{,i} \rangle ds + 4 \int_0^t \langle m_{ij} q_{,j} q_{,i} \rangle ds \\ &\quad + 4 \int_0^t \langle \kappa_{ij} \theta_{1,j} \theta_{1,i} \rangle ds \\ &\quad + 4 \int_0^t \langle \omega_{ij} \theta_{2,j} \theta_{2,i} \rangle ds + 4 \int_0^t \langle \pi_{ij} \theta_{3,j} \theta_{3,i} \rangle ds \\ &\quad + 4 \int_0^t \langle \gamma (p - q)^2 \rangle ds + 4 \int_0^t \langle d_1 (\theta_1 - \theta_2)^2 \rangle ds \\ &\quad + 4 \int_0^t \langle d_2 (\theta_1 - \theta_3)^2 \rangle ds + 4 \int_0^t \langle d_3 (\theta_2 - \theta_3)^2 \rangle ds . \end{aligned} \quad (16)$$

We now utilize (16), (9) and (10) to form the combination  $FF'' - (F')^2$  to see that

$$FF'' - (F')^2 = 4S^2 \geq 0, \quad (17)$$

where  $S^2$ , which is non-negative due to the Cauchy-Schwarz inequality, is given by

$$S^2 = AB - C^2,$$

where the terms  $A$ ,  $B$  and  $C$  have form

$$\begin{aligned}
A &= \langle \rho u_i \dot{u}_i \rangle + \int_0^t \langle k_{ij} \eta_j \eta_i \rangle ds + \int_0^t \langle m_{ij} \zeta_j \zeta_i \rangle ds \\
&+ \int_0^t \langle \kappa_{ij} \phi_j \phi_i \rangle ds + \int_0^t \langle \omega_{ij} \psi_j \psi_i \rangle ds + \int_0^t \langle \pi_{ij} \xi_j \xi_i \rangle ds \\
&+ \int_0^t \langle \gamma(\eta - \zeta)^2 \rangle ds + \int_0^t \langle d_1(\phi - \psi)^2 \rangle ds + \int_0^t \langle d_2(\psi - \xi)^2 \rangle ds + \int_0^t \langle d_3(\phi - \xi)^2 \rangle ds, \\
B &= \langle \rho \dot{u}_i \dot{u}_i \rangle + \int_0^t \langle k_{ij} p_j p_i \rangle ds + \int_0^t \langle m_{ij} q_j q_i \rangle ds \\
&+ \int_0^t \langle \kappa_{ij} \theta_{1j} \theta_{1i} \rangle ds + \int_0^t \langle \omega_{ij} \theta_{2j} \theta_{2i} \rangle ds + \int_0^t \langle \pi_{ij} \theta_{3j} \theta_{3i} \rangle ds \\
&+ \int_0^t \langle \gamma(p - q)^2 \rangle ds + \int_0^t \langle d_1(\theta_1 - \theta_2)^2 \rangle ds \\
&+ \int_0^t \langle d_2(\theta_1 - \theta_3)^2 \rangle ds + \int_0^t \langle d_3(\theta_2 - \theta_3)^2 \rangle ds, \\
C &= \langle \rho u_i \dot{u}_i \rangle + \int_0^t \langle k_{ij} \eta_j p_i \rangle ds + \int_0^t \langle m_{ij} \zeta_j q_i \rangle ds \\
&+ \int_0^t \langle \kappa_{ij} \phi_j \theta_{1i} \rangle ds + \int_0^t \langle \omega_{ij} \psi_j \theta_{2i} \rangle ds + \int_0^t \langle \pi_{ij} \xi_j \theta_{3i} \rangle ds \\
&+ \int_0^t \langle \gamma(\eta - \zeta)(p - q) \rangle ds + \int_0^t \langle d_1(\phi - \psi)(\theta_1 - \theta_2) \rangle ds \\
&+ \int_0^t \langle d_2(\phi - \xi)(\theta_1 - \theta_3) \rangle ds + \int_0^t \langle d_3(\psi - \xi)(\theta_2 - \theta_3) \rangle ds.
\end{aligned}$$

From (17) one now deduces  $F \equiv 0$  on  $[0, T]$ , see e.g. Straughan<sup>1</sup>, chapter 1. Roughly, one divides (17) by  $F^2$  to find  $(\log F)'' \geq 0$ , but care has to be taken since  $F(0) = 0$ . One has to use a contradiction argument involving  $F(\epsilon)$ , and then let  $\epsilon \rightarrow 0$ . Since  $F \equiv 0$  it follows from (9) that  $u_i \equiv 0$ . Next, we appeal to the energy equation (15) using (14), to conclude also that

$$p \equiv 0, \quad q \equiv 0, \quad \theta_1 \equiv 0, \quad \theta_2 \equiv 0, \quad \theta_3 \equiv 0,$$

on  $\Omega \times [0, T]$  and uniqueness follows, with only the symmetry conditions (4) on the elastic coefficients.

## 4 | DECAY

In double porosity elasticity theory several writers argue that for many applications one may dispense with the acceleration term in equation (1)<sub>1</sub>, see e.g. Scarpetta and Svanadze<sup>12</sup> and the references therein. In this case the system is referred to as a quasi-equilibrium system, cf. Scarpetta and Svanadze<sup>12</sup>, Svanadze<sup>5</sup>.

In this section we analyze the quasi-equilibrium version of equations (1) with zero source terms, and so omit the  $\rho \ddot{u}_i$  term. The boundary conditions adopted are those of (6), while the initial conditions are (3). The boundary-initial value problem thus



defined is denoted by  $\mathcal{M}$ . We retain the symmetries of (4), but now we require  $a_{ijhk}$  to be positive definite, i.e.

$$a_{ijhk}\varepsilon_{hk}\varepsilon_{ij} \geq a_0\varepsilon_{ij}\varepsilon_{ij}, \quad (18)$$

for all  $\varepsilon_{ij}$ , where  $a_0 > 0$  is a constant. In addition we impose the (realistic) bounds on the coefficients,

$$\beta_{ij}\beta_{ij}, \quad \gamma_{ij}\gamma_{ij}, \quad \gamma, \quad a_{ij}^{(1)}a_{ij}^{(1)}, \quad a_{ij}^{(2)}a_{ij}^{(2)}, \quad a_{ij}^{(3)}a_{ij}^{(3)} \leq M, \quad (19)$$

for some constant  $M$ , where (19) hold  $\forall \mathbf{x} \in \Omega$ .

Furthermore, for positive constants  $\alpha_0, \beta_0, \hat{\alpha}_1, \hat{\alpha}_2, \hat{\alpha}_3, \tilde{\alpha}_0, \tilde{\beta}_0, \tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3, k_1, m_1, \kappa_1, \omega_1, \pi_1$ , we require

$$\alpha_0 \leq \alpha \leq \tilde{\alpha}_0, \quad \beta_0 \leq \beta \leq \tilde{\beta}_0, \quad \hat{\alpha}_1 \leq \alpha_1 \leq \tilde{\alpha}_1, \quad \hat{\alpha}_2 \leq \alpha_2 \leq \tilde{\alpha}_2, \quad \hat{\alpha}_3 \leq \alpha_3 \leq \tilde{\alpha}_3, \quad (20)$$

and

$$\begin{aligned} k_{ij}\varepsilon_j\varepsilon_i &\geq k_1\varepsilon_i\varepsilon_i, & m_{ij}\varepsilon_j\varepsilon_i &\geq m_1\varepsilon_i\varepsilon_i, & \kappa_{ij}\varepsilon_j\varepsilon_i &\geq \kappa_1\varepsilon_i\varepsilon_i, \\ \omega_{ij}\varepsilon_j\varepsilon_i &\geq \omega_1\varepsilon_i\varepsilon_i, & \pi_{ij}\varepsilon_j\varepsilon_i &\geq \pi_1\varepsilon_i\varepsilon_i, \end{aligned} \quad (21)$$

$\forall \mathbf{x} \in \Omega$  and for all  $\varepsilon_i$ .

We shall establish that a solution to  $\mathcal{M}$  decays to zero in an appropriate manner. To do this we commence by multiplying the quasi-equilibrium version of (5)<sub>1</sub> by  $u_i$  and then integrate over  $\Omega$  and use the homogeneous boundary conditions to derive

$$\begin{aligned} \langle a_{ijhk}u_{h,k}u_{i,j} \rangle &= \langle \beta_{ij}pu_{i,j} \rangle + \langle \gamma_{ij}qu_{i,j} \rangle \\ &+ \langle a_{ij}^{(1)}\theta_1u_{i,j} \rangle + \langle a_{ij}^{(2)}\theta_2u_{i,j} \rangle + \langle a_{ij}^{(3)}\theta_3u_{i,j} \rangle. \end{aligned} \quad (22)$$

We next employ the Cauchy-Schwarz and arithmetic-geometric mean inequalities on the terms on the right of (22). After, using the bounds (19) and (20) one may readily deduce the inequality

$$\frac{1}{2} \langle a_{ijhk}u_{h,k}u_{i,j} \rangle \leq C^* \left[ \langle \alpha p^2 \rangle + \langle \beta q^2 \rangle + \langle a_1\theta_1^2 \rangle + \langle a_2\theta_2^2 \rangle + \langle a_3\theta_3^2 \rangle \right], \quad (23)$$

where

$$C^* = \frac{5MK}{2\hat{\alpha}_3a_0}$$

with

$$K = \frac{\hat{\alpha}_3}{\alpha_0} + \frac{\hat{\alpha}_3}{\beta_0} + \frac{\hat{\alpha}_3}{\hat{\alpha}_1} + \frac{\hat{\alpha}_3}{\hat{\alpha}_2} + 1.$$

The next step is to multiply the quasi-equilibrium version of (5)<sub>1</sub> by  $\dot{u}_i$  and integrate over  $\Omega$ , and then multiply (5)<sub>2</sub>–(5)<sub>6</sub> by  $p, q, \theta_1, \theta_2$  and  $\theta_3$ , respectively, and integrate each over  $\Omega$ . Denote by  $E_e$  and  $E_d$  the expressions

$$\begin{aligned} E_e &= \frac{1}{2} \langle a_{ijhk}u_{h,k}u_{i,j} \rangle, \\ E_d &= \frac{1}{2} \langle \alpha p^2 \rangle + \frac{1}{2} \langle \beta q^2 \rangle + \frac{1}{2} \langle a_1\theta_1^2 \rangle + \frac{1}{2} \langle a_2\theta_2^2 \rangle + \frac{1}{2} \langle a_3\theta_3^2 \rangle, \end{aligned}$$

corresponding to the elastic energy and to the (thermo-bidisperse) dissipative term. By adopting the procedure above one may show

$$\frac{d}{dt} (E_e + E_d) = -f(t), \quad (24)$$

where

$$\begin{aligned} f(t) &= \langle k_{ij}p_{,j}p_{,i} \rangle + \langle m_{ij}q_{,j}q_{,i} \rangle + \langle \kappa_{ij}\theta_{1,j}\theta_{1,i} \rangle \\ &+ \langle \omega_{ij}\theta_{2,j}\theta_{2,i} \rangle + \langle \pi_{ij}\theta_{3,j}\theta_{3,i} \rangle + \langle \gamma(p-q)^2 \rangle \\ &+ \langle d_1(\theta_1 - \theta_2)^2 \rangle + \langle d_2(\theta_1 - \theta_3)^2 \rangle + \langle d_3(\theta_2 - \theta_3)^2 \rangle. \end{aligned} \quad (25)$$

Upon integration, we see from (24) with  $E = E_e + E_d$

$$0 \leq E(t) = E(0) - \int_0^t f(s)ds. \quad (26)$$

Next, using Poincaré's inequality together with (21) and the definition of  $E$ , from (26) we may deduce

$$\begin{aligned} E_d(t) &\leq E(t) = E(0) - \int_0^t f(s) ds \\ &\leq E(0) - \lambda_1 \int_0^t (k_1 \langle p^2(s) \rangle + m_1 \langle q^2(s) \rangle + \kappa_1 \langle \theta_1^2(s) \rangle + \langle \omega_1 \langle \theta_2^2(s) \rangle + \langle \pi_1 \langle \theta_3^2(s) \rangle) ds, \end{aligned} \quad (27)$$

$\lambda_1$  being the first eigenvalue in the membrane problem for  $\Omega$ .

Therefore, using the upper bounds of (20), we may obtain

$$\begin{aligned} E_d(t) &\leq E(0) - \lambda_1 \int_0^t \left( \frac{k_1}{\tilde{\alpha}_0} \langle \alpha p^2(s) \rangle + \frac{m_1}{\tilde{\beta}_0} \langle \beta q^2(s) \rangle + \frac{\kappa_1}{\tilde{a}_1} \langle a_1 \theta_1^2(s) \rangle + \frac{\omega_1}{\tilde{a}_2} \langle a_2 \theta_2^2(s) \rangle + \frac{\pi_1}{\tilde{a}_3} \langle a_3 \theta_3^2(s) \rangle \right) ds \\ &\leq E(0) - k \int_0^t E_d(s) ds, \end{aligned} \quad (28)$$

where

$$k = 2\lambda_1 \min \left\{ \frac{k_1}{\tilde{\alpha}_0}, \frac{m_1}{\tilde{\beta}_0}, \frac{\kappa_1}{\tilde{a}_1}, \frac{\omega_1}{\tilde{a}_2}, \frac{\pi_1}{\tilde{a}_3} \right\}.$$

We integrate (27) with the aid of an integrating factor to find

$$\int_0^t E_d(s) ds \leq \frac{E(0)}{k} (1 - e^{-kt}), \quad \forall t > 0,$$

so

$$\int_0^\infty E_d(s) ds \leq \frac{E(0)}{k}. \quad (29)$$

However, from inequality (23) we see that

$$\frac{E_e}{2C^*} \leq E_d.$$

Hence, from (29) we may show

$$\int_0^\infty E(s) ds \leq \frac{1 + 2C^*}{k} E(0). \quad (30)$$

Therefore,  $E(t) \in L^1(0, \infty)$  and from (24)  $E'(t) \leq 0, \forall t > 0$ . Thus,

$$\lim_{t \rightarrow \infty} E(t) = 0. \quad (31)$$

Relation (31) shows that the solution to  $\mathcal{M}$  decays to zero in time in the measure  $E(t)$ .

## 5 | CONCLUSIONS

In this paper we have introduced a set of partial differential equations to model the evolutionary behaviour of an anisotropic elastic body which contains a double porosity structure under conditions of local thermal non-equilibrium. We have proved that a solution to the displacement boundary initial value problem is unique under the weak requirement of symmetry of the elastic coefficients. In addition, we have demonstrated that the solution to the equivalent problem for a quasi-equilibrium theory decays to zero in time in a precise sense, when there are no sources and external body force involved.

It is still an open problem to know the rate of decay, even in the quasi-equilibrium theory. It would also be very interesting to know properties of decay for the full theory along the lines of analogous results for classical linear thermoelasticity by Dafermos<sup>20</sup> and Lebeau and Zuazua<sup>21</sup>. Given that the decay rate even for thermoelasticity may be polynomial or exponential depending on the geometry, it is natural to expect something similar in the present more complicated model. In fact, in equation (1) there are two potential sources of damping. One is the analogous one from thermoelasticity involving the  $\beta_{ij}, \gamma_{ij}, a_{ij}^{(s)}$ ,

$s = 1, 2, 3$ , terms. However, a second source of damping could arise from the interaction terms between the macro and micro pressures, the terms involving  $\gamma$ , and between the temperatures, the  $d_\alpha$ ,  $\alpha = 1, 2, 3$ , terms. Since the  $\gamma$  terms involve the difference  $p - q$ , and  $d_\alpha$  terms involve  $\theta_\alpha - \theta_\beta$ , this has some resemblance to the situation for a mixture of two elastic solids where the dissipation involves the difference of velocities, a problem resolved by Dafermos<sup>22</sup>.

## Acknowledgment

Research performed under the auspices of G.N.F.M. - I.N.d.A.M. and partially supported by Italian M.I.U.R.. We should like to thank the anonymous referees for incisive remarks which have led to improvements in the paper.

## References

1. Straughan, B., *Mathematical aspects of multi-porosity continua*. Advances in Mechanics and Mathematics, vol. 38. New York: Springer; 2017.
2. Olusola, B. K., Yu, G., Aguilera, R., The use of electromagnetic mixing rules for petrophysical evaluation of dual-and triple-porosity reservoirs. *SPE Reservoir Evaluation & Engineering*. 2013;16(04):378–389.
3. Kim, J., Moridis, G. J., Numerical analysis of fracture propagation during hydraulic fracturing operations in shale gas systems. *Int J Rock Mech Min Sci*. 2015;76:127–137.
4. Svanadze, M., Scalia, A., Mathematical problems in the coupled linear theory of bone poroelasticity. *Comput. Math. Appl.*. 2013;66(9):1554–1566.
5. Svanadze, M., On the linear theory of double porosity thermoelasticity under local thermal non-equilibrium, manuscript.
6. Nield, D., Effects of local thermal nonequilibrium in steady convective processes in a saturated porous medium: forced convection in a channel. *J Porous Media*. 1998;1(2):181–186.
7. Rees, D. A. S., Bassom, A. P., Siddheshwar, P. G., Local thermal non-equilibrium effects arising from the injection of a hot fluid into a porous medium. *J. Fluid Mech.*. 2008;594:379–398.
8. Straughan, B., *Convection with local thermal non-equilibrium and microfluidic effects* Advances in Mechanics and Mathematics, vol. 32. New York: Springer; 2015.
9. Svanadze, M., Plane waves and boundary value problems in the theory of elasticity for solids with double porosity. *Acta Appl. Math.*. 2012;122:461–471.
10. Svanadze, M., Uniqueness theorems in the theory of thermoelasticity for solids with double porosity. *Meccanica*. 2014;49(9):2099–2108.
11. Ciarletta, M., Passarella, F., Svanadze, M., Plane waves and uniqueness theorems in the coupled linear theory of elasticity for solids with double porosity. *J. Elasticity*. 2014;114(1):55–68.
12. Scarpetta, E., Svanadze, M., Uniqueness theorems in the quasi-static theory of thermoelasticity for solids with double porosity. *J. Elasticity*. 2015;120(1):67–86.
13. Siratovich, P. A., Villeneuve, M. C., Cole, J. W., Kennedy, B. M., Bégué, F., Saturated heating and quenching of three crustal rocks and implications for thermal stimulation of permeability in geothermal reservoirs. *Int J Rock Mech Min Sci*. 2015;80:265–280.
14. Xinchun, S., Lakes, R. S., Stability of elastic material with negative stiffness and negative Poisson's ratio. *Phys. Status Solidi B*. 2007;244(3):1008–1026.
15. Ha, C. S., Hestekin, E., Li, J., Plesha, M. E., Lakes, R. S., Controllable thermal expansion of large magnitude in chiral negative Poisson's ratio lattices. *Phys. Status Solidi B*. 2015;252(7):1431–1434.

16. Chiriță, S., Ciarletta, M., Spatial estimates for the constrained anisotropic elastic cylinder. *J. Elasticity*. 2006;85(3):189–213.
17. Chiriță, S., Danescu, A., Ciarletta, M., On the strong ellipticity of the anisotropic linearly elastic materials. *J. Elasticity*. 2007;87(1):1–27.
18. Chiriță, S., Danescu, A., Strong ellipticity for tetragonal system in linearly elastic solids. *Int. J. Solids Struct.* 2008;45(17):4850–4859.
19. Chiriță, S., Ghiba, I.-D., Strong ellipticity and progressive waves in elastic materials with voids. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* 2010;466(2114):439–458.
20. Dafermos, C. M., On the existence and the asymptotic stability of solutions to the equations of linear thermoelasticity. *Arch. Rational Mech. Anal.* 1968;29(4):241–271.
21. Lebeau, G., Zuazua, E., Decay rates for the three-dimensional linear system of thermoelasticity. *Arch. Ration. Mech. Anal.* 1999;148(3):179–231.
22. Dafermos, C. M., Wave equations with weak damping. *SIAM J. Appl. Math.* 1970;18:759–767.

