



ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Exploiting phenotype diversity in a local animal genetic resource: Identification of a single nucleotide polymorphism associated with the tail shape phenotype in the autochthonous Casertana pig breed

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Exploiting phenotype diversity in a local animal genetic resource: Identification of a single nucleotide polymorphism associated with the tail shape phenotype in the autochthonous Casertana pig breed / Bertolini, Francesca; Schiavo, Giuseppina; Tinarelli, Silvia; Santoro, Laura; Utzeri, Valerio Joe; Dall'Olio, Stefania; Nanni Costa, Leonardo; Gallo, Maurizio; Fontanesi, Luca*. - In: LIVESTOCK SCIENCE. - ISSN 1871-1413. - ELETTRONICO. - 216:(2018), pp. 148-152. [10.1016/j.livsci.2018.08.007]

This version is available at: <https://hdl.handle.net/11585/661002> since: 2019-02-06

Published:

DOI: <http://doi.org/10.1016/j.livsci.2018.08.007>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

(Article begins on next page)

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

1 *This is the post-peer reviewed version of the following article: “Exploiting phenotype diversity in a*
2 *local animal genetic resource: identification of a single nucleotide polymorphism associated with*
3 *the tail shape phenotype in the autochthonous Casertana pig breed” by Francesca Bertolini,*
4 *Giuseppina Schiavo, Silvia Tinarelli, Laura Santoro, Valerio Joe Utzeri, Stefania Dall’Olio,*
5 *Leonardo Nanni Costa, Maurizio Gallo and Luca Fontanesi, which has been published in final form*
6 *at 10.1016/j.livsci.2018.08.007. This article may be used for non-commercial purposes in*
7 *accordance with Elsevier Terms and Conditions for Self-Archiving.”*
8

9 **Exploiting phenotype diversity in a local animal genetic resource: identification of a single**
10 **nucleotide polymorphism associated with the tail shape phenotype in the autochthonous**
11 **Casertana pig breed**

12

13 Francesca Bertolini^{1,2,*}, Giuseppina Schiavo^{1,*}, Silvia Tinarelli^{1,3}, Laura Santoro⁴, Valerio Joe
14 Utzeri¹, Stefania Dall'Olio¹, Leonardo Nanni Costa¹, Maurizio Gallo³ and Luca Fontanesi¹

15

16 ¹Department of Agricultural and Food Sciences, Division of Animal Sciences, University of Bologna,
17 Viale Fanin 46, 40127 Bologna, Italy

18 ²Department of Bio and Health Informatics, Technical University of Denmark, Kemitovet, 2800
19 Kgs. Lyngby, Denmark

20 ³Associazione Nazionale Allevatori Suini, Via L. Spallanzani 4, 00161 Roma, Italy

21 ⁴ConSDABI - National Focal Point Italiano FAO, Contrada Piano Cappelle, 82100 Benevento, Italy

22

23 Corresponding author: Luca Fontanesi, e-mail: luca.fontanesi@unibo.it

24

25 * These authors contributed equally to this work.

26

27 **Running title:** SNPs and tail shape phenotype in pigs

28 **Highlights**

- 29 • Casertana is an autochthonous pig genetic resource reared in Central-South of Italy.
- 30 • Tail shape phenotype variability in the breed was investigated in a GWAS.
- 31 • A single nucleotide polymorphism on porcine chromosome 12 was associated with this trait.
- 32 • This marker is close to the *SRY-box 9 (SOX9)* gene that is essential in skeletogenesis.

33 **Abstract**

34 Casertana is a local pig breed mainly raised in Central-South regions of Italy. Pigs of this breed
35 are considered the descendants of the ancient Neapolitan population that largely influenced the
36 constitution of the modern commercial pigs. The pigs of this breed are usually curly-tailed, like
37 several other domestic pig populations. However, Casertana population shows some variability for
38 this trait, including animals having straight tail as observed in wild boars. In this study, we run, for
39 the first time, a genome wide association study (GWAS) comparing the curly tailed (no. = 53) and
40 straight tailed (no. = 19) Casertana pigs to identify genomic regions associated with the tail shape
41 phenotype in *Sus scrofa*. All animals were genotyped with the Illumina PorcineSNP60 BeadChip v.2.
42 GEMMA software was used in the GWAS for which we were able to correct for stratification in the
43 analysed cohort. A single nucleotide polymorphism (rs81439488), located on porcine chromosome
44 12, was significantly associated with the investigated trait. This marker is close to the *SRY-box 9*
45 (*SOX9*) gene that encodes for a transcription factor that is required during sequential steps of the
46 chondrocyte differentiation pathway, notochord maintenance and skeletogenesis. As the shape of the
47 tail could be important in relation to the problem of tail biting in pigs, the obtained results might open
48 new perspectives for defining selection programs answering indirectly animal welfare issues. This
49 work demonstrated that autochthonous animal genetic resources might be used to disclose genetic
50 factors affecting peculiar traits by exploiting segregating phenotypes and genetic variability.

51

52 **Keywords:** Animal genetic resource; autochthonous breed; GWAS; morphological trait; SNP; *Sus*
53 *scrofa*.

54 **1. Introduction**

55 Conservation of animal genetic resources is mainly aimed to preserve genetic diversity and
56 associated inheritable phenotypes characterizing different populations that might be interesting for
57 current or future purposes, including potential use in breeding programs. These resources can be also
58 useful to understand biological mechanisms determining unique phenotypes derived by diversity in
59 selection pressures or as result of adaptation to environmental and production conditions (Leroy et
60 al., 2016).

61 Casertana pigs constitute a local breed mainly raised in Central-South regions of Italy. Pigs of
62 this breed are considered the descendants of the ancient Neapolitan pig population that largely
63 influenced the constitution of the modern commercial pig breeds through introgression of blood into
64 British pig populations during the 19th century (Porter, 1993). Neapolitan pigs were, in turn,
65 influenced by Asian blood in the late 18th century (Porter, 1993). Casertana is enlisted among the
66 endangered animal genetic resources as the herd book of this breed accounts for about 100 boars and
67 sows currently registered (ANAS, 2016). Animals are mainly raised in extensive or semi-extensive
68 production systems with possible contacts and crossbreeding with European wild boars that could
69 have contributed, at least in part, to shape their morphological characteristics. Casertana pigs have a
70 black or grey coat colour, wrinkled skin, forward ears, and usually a typical hairless phenotype. The
71 pigs of this breed are usually curly-tailed, like several other domestic pig populations. However,
72 Casertana population shows some variability for this trait, including animals having **straight** and wavy
73 tail as in a few other pig breeds and in wild boars.

74 Domestication in mammals has been a complex and continuous process associated with a series
75 of changes in the domesticated animals compared to the wild counterparts, derived by selective
76 breeding of animals showing favourable production and reproduction performances, and increased
77 docility that indirectly shaped the genome of domesticated populations (Wiener and Wilkinson, 2011;
78 Larson and Burger, 2013; Carneiro et al., 2014; Wang et al., 2014; Wilkins et al., 2014). Several
79 morphological features have been also directly or indirectly selected and, in most cases, fixed in

80 domesticated populations as result of the domestication process (Darwin, 1868). Coat colour is one
81 of the most common phenotypic traits that has been modified as result of reduced selective pressure
82 against colours with low fitness in the wild and of aesthetic preferences of the breeders, sometimes
83 associated with higher production performances (Clutton-Brock, 1999). Among several other
84 morphological characters, curliness of the tail and shape has been associated with domestication in
85 mammals (Trut et al., 2009).

86 The tail is considered an extension of the spinal column usually composed of specifically
87 shaped vertebrae. Spontaneous curly tail phenotypes in mice have been the matter of studies that
88 investigated the role of embryonic development in this morphological anomaly (Copp et al., 1988;
89 van Straaten and Copp, 2001; Ohnishi et al., 2017). Curly tail is also commonly observed in many
90 dog breeds. Vaysse et al. (2011) compared the genome of dog breeds having curly tails with that of
91 breeds with straight tails using single nucleotide polymorphisms (SNPs) chip data and identified a
92 genomic region on chromosome 1 significantly associated with these alternative tail shapes.

93 In pigs, few studies have been reported on the genetic factors affecting tail shape. A putative
94 recessive genetic defect known as kinky tail (or flexed or screw tail), derived by fused caudal
95 vertebrae associated in some cases with other defects, has been described in the mid of the last century
96 (Nordby, 1934; Donald, 1949; Brooksbank, 1958). It is not known if this defect could be, in some
97 way, related or not to the normal curling of the tail that is common in domestic pigs. This signature
98 of domestication, however, seems not fixed in all pig breeds (Porter, 1993) but no systematic study
99 has been conducted so far, probably because the difficulties in retrieving phenotype information due
100 to the usual practice of tail docking in most herds.

101 In this study, we took advantage from the variability of the shape of the tail that we recorded in
102 the Casertana pig population and run a genome wide association study (GWAS) comparing the
103 genome of curly-tailed and straight tailed animals to identify genomic regions associated with the tail
104 shape phenotype in *Sus scrofa*.

105

106 **2. Materials and methods**

107 **2.1. Animals**

108 A total of 101 Casertana pigs (of about 7 to 20 months old) from six different farms were
109 evaluated. Photographic records of each animal were obtained to capture information on the tail shape
110 in standardized restraining conditions (including a direct evaluation of the personnel on this
111 phenotype during this phase for the biological sampling) for all animals and after release (Figure 1).
112 Pigs were classified as follows: 53 (25 males and 28 females) showed the curly tail phenotype; 19
113 (five males and 14 females) showed the straight tail phenotype; 29 were not classified and excluded
114 from the study as tail docking, that was practised by the farmers as routine before weaning of the
115 piglets, prevented the recording of any tail phenotype.

116

117 **2.2. Genotyping**

118 Hairs (with roots) were collected from the investigated pigs. DNA extraction was carried out
119 using the Wizard® Genomic DNA Purification kit (Promega Corporation, Madison, WI, USA)
120 following the manufacturer's instructions. Genotyping of the extracted DNA was obtained with the
121 Illumina PorcineSNP60 BeadChip v.2 (Illumina, Inc., San Diego, CA, USA) that interrogated 61,565
122 SNPs. Single nucleotide polymorphisms were assigned to the Sscrofa11.1 genome version, as
123 previously described (Fontanesi et al., 2012). PLINK 1.9 software (Chang et al., 2015) was used to
124 filter SNPs and genotyping data using the following criteria already used in a similar study (Schiavo
125 et al., 2018): genotyping call rate >0.9, minor allele frequency >0.01 and Hardy-Weinberg
126 equilibrium $P > 0.001$.

127

128 **2.3. Data analysis and genome wide association**

129 To evaluate distance relationships among the animals of the investigated cohort,
130 multidimensional scaling (MDS) was obtained with the PLINK 1.9 software (Chang et al., 2015).
131 Genome wide association study was carried out by applying the univariate mixed model of GEMMA

132 (Zhou and Stephens, 2012) that can accommodate the centered relatedness matrix calculated from
133 SNP genotypes to correct for population stratification in a case and control analysis. The model also
134 included the farm and the sex as fixed effects. To be able to identify associated markers in this
135 experiment that included a low number of animals (derived by the fact that the analysed pigs were
136 almost a complete representation of the whole population of the Casertana breed) and that used a SNP
137 chip that might originally have an ascertain bias (as local breeds were not used for the selection of
138 the informative SNPs), the significant threshold was defined at the $P_{nominal\ value} < 5.00E-05$ level,
139 according to the Wellcome Trust Case Control Consortium (2007) and as also applied in several other
140 GWAS in livestock (e.g. Fontanesi et al., 2012; Sanchez et al., 2014). Genomic inflation factor (λ)
141 and quantile–quantile (Q–Q) plot were obtained with GenABEL (Aulchenko et al., 2007). Gene
142 annotation information was retrieved from the Sscrofa11.1 genome version available at the Ensembl
143 database (http://www.ensembl.org/Sus_scrofa/Info/Index), release 91.

144

145 **3. Results and discussion**

146 A recent phenotypic characterization of the endangered Casertana pig population that we
147 carried out noted several morphological differences among distinct animals of this autochthonous
148 breed (data not shown). For example, in addition to the hairless or hypotrichotic condition (that is the
149 characteristic phenotype of the Casertana animals), we already described the presence of haired pigs
150 in this population and this morphological variability was used for a GWAS that we have recently
151 reported (Schiavo et al. 2018). Despite a limited number of animals was included in that study, we
152 were able to identify genomic regions associated with the hairless phenotype, demonstrating that local
153 animal genetic resources can be used to genetically describe phenotypic variability of simple traits
154 (Schiavo et al., 2018).

155 Another morphological trait that is not fixed in this breed is the shape of the tail (Figure 1). Of
156 the animals for which we could record this phenotype, 26% (19 out of 72) showed a straight tail
157 without any curls, similarly to the usual shape of wild boars. This shape was clearly different from

158 the curly tails reported in the remaining investigated pigs (74%). There was no age effect and the two
159 groups included a comparable number of males and females so that sex did not explain the tail shape
160 phenotype. In addition, we could exclude the possible effect of the behavioral change of tail posture
161 on this phenotype (Zonderland et al., 2009). The recording system was based on standardized
162 conditions and subsequent photographic records of the animals confirmed, at least for the
163 photographic time point, their assignment to one or to the other group of tail shape phenotype. The
164 two groups were observed in animals from all six farms. Limited pedigree record prevented the
165 possibility to evaluate any potential founder effect.

166 A total of 36,533 autosomal SNPs, mapped to a unique position in the Sscrofa11.1 genome
167 version, was used for MDS. The obtained MDS plot showed some structures not well defined in the
168 analysed pigs that however did not clearly separate the curly and straight tailed Casertana pigs (Figure
169 2). A stratified sample could be a critical point in GWAS in a very small population where, to some
170 extent, all animals might be related. Figure S1 reports the genomic inflation factor (λ) and Q-Q plot
171 that did not show any biased test statistic distribution, suggesting that the investigated cohort was
172 corrected for a possible stratification effect.

173 Figure 3 reports the Manhattan plot obtained in this GWAS. One significant SNP ($P=2.3E-05$)
174 was identified on porcine chromosome 12 (SSC12). This marker indicated as ALGA0064877
175 (rs81439488 C/A) is located at position 10,301,075 of this chromosome. Allele A of this SNP was
176 more frequent in the straight tail group (0.17) than in the curly tailed group (0.03).

177 **The complete list of genes annotated around the ALGA0064877 marker (with known functions)**
178 **is reported in Table S1. Figure S2 report a screenshot of this SSC12 region retrieved from ENSEMBL**
179 **database with the annotated genes.** One of the closest gene in this desert chromosome region is the
180 *SRY-box 9 (SOX9)* gene (positions 8,641,629-8,647,764, encoded by the -1 strand; actually the closest
181 upstream gene to the significant SNP), that, according to its function, might be the most plausible
182 candidate gene, explaining the recorded phenotypic variability. It is well established that the
183 expression of this gene at the embryonal level marks the onset of cartilage differentiation (Wright et

184 al., 1995; Healy et al., 1996). *SOX9* encodes for a transcription factor that is required during sequential
185 steps of the chondrocyte differentiation pathway, notochord maintenance and skeletogenesis
186 (Akiyama et al., 2002; Barrionuevo et al., 2006; Montero et al., 2017). Continued expression of *Sox9*
187 in differentiated chondrocytes is essential for subsequent hypertrophy and sustains chondrocyte-
188 specific survival mechanisms (Ikegami et al., 2011). Although *SOX9* seems to be a master gene for
189 chondrocytes differentiation, the whole region surrounding the marker ALGA0064877 is downstream
190 closer to a few genes [i.e. potassium voltage-gated channel subfamily J member 2 (*KCNJ2*), 61169
191 bp of distance from the SNP; potassium voltage-gated channel subfamily J member 16 (*KCNJ16*),
192 147461 bp of distance; mitogen-activated protein kinase kinase 6 (*MAP2K6*), 570407 bp of distance].
193 Alterations in the genes of this region have been also observed to affect the Wnt pathway (Kurth et
194 al., 2009). Heterozygous mutations within and around human *SOX9* cause campomelic dysplasia that
195 is a malformation syndrome characterized by cartilage derived skeletal structure defects (Foster et
196 al., 1994; Wagner et al., 1994). These mutations, most of which reduce the level of expression of this
197 gene, are located upstream spanning a large region (from 50 kb to more than 1 Mb) in which
198 regulatory elements are present (Wunderle et al., 1998; Bagheri-Fam et al., 2006). Close upstream
199 mutations produce more severe defects whereas far upstream mutations cause mild defects (Pfeifer
200 et al., 1999; Velagaleti et al., 2005; Leipoldt et al., 2007). About 40 k SNPs, located between *SOX9*
201 and *MAP2K6*, are reported in the Sscrofa11.1 genome version and could be considered for future
202 studies to identify the causative mutation(s). Among them, 20 are also present in the PorcineSNP60
203 BeadChip but only ALGA0064877 reached the threshold of significance, probably due to the biased
204 chip design that might not be able to capture particular haplotype structures of this breed.

205 Based on these studies in other species it is tempting to suggest a possible regulatory mechanism
206 affecting *SOX9* expression in porcine developing chondrocytes that would, in turn, produce a mild
207 cartilage/skeletal effect determining the shape of the tail. This hypothesis might be worth of further
208 investigation starting from a precise characterization of the structure and morphology of the pig tail
209 with different shapes for which, at present, there is no detailed investigation. Our phenotype records

210 were based only on an external morphological evaluation of the shape of the tail. Furthermore,
211 analysis of gene expression of *SOX9* at different developmental stages should be also carried out to
212 evaluate the role of this gene in the phenotype observed in pigs.

213 The results we obtained might have broader impacts than those that would be limited to a simple
214 morphological characterization. The shape of the tail could be important in relation to the problem of
215 tail biting in pigs. Tail biting is a widespread behavioral vice with significant animal welfare
216 implications and economic losses in commercial pig farms (Bracke et al., 2004). A few studies have
217 established correlations between tail posture and tail biting incidence suggesting limited damages and
218 related welfare complications with behaviors of the pigs that tended to have a tail posture up that
219 those with tail posture down (Zonderland et al., 2009; Lahrmann et al., 2018). It would be interesting
220 to evaluate if pigs with genetically determined curly tails (as a possible adaptation derived by the
221 domestication process) are less affected by tail biting damages than pigs with straight tails.

222

223 **4. Conclusions**

224 This work demonstrated that autochthonous animal genetic resources, even constituted by very
225 small populations, might be used to disclose genetic factors affecting peculiar traits by exploiting
226 segregating phenotypes and genetic variability. To our knowledge, this is the first study that reported
227 a frequency distribution of the tail shape phenotype in a pig population. Our results indicated that this
228 morphological trait is associated with a marker close to an important gene involved in embryonic
229 development, opening other hypothesis, worth of further investigations. It will be important to
230 validate the results we obtained in this GWAS in other breeds and populations, including a more
231 precise anatomical characterization of this trait, to further extend the impact of the results reported in
232 Casertana pigs. It would be however first needed to know if diversity for this morphological
233 characteristic is common in commercial pig populations as at present, there is not information on this
234 aspect, mainly due to the usual practice of tail docking that prevents the recording of this phenotype.
235 Considering the potential relationship between tail shape and tail biting damages (that, however,

236 remains to be formally demonstrated), it could be possible to envisage practical applications of the
237 identified marker in selection programs aimed to respond to animal welfare issues. From this study it
238 emerges that conservation strategies of autochthonous pig genetic resources should take also into
239 account the preservation of phenotypic variability within populations. Our study represents one of the
240 few examples of exploitation of animal genetic resources to recover information that might have
241 potential impacts in commercial populations.

242

243 **Acknowledgements**

244 The Authors thank all farmers and personnel that contributed to collect phenotypes and samples.

245

246 **Funding**

247 This work has received funding from the Italian Ministry of Agriculture, Food and Forestry
248 (MiPAAF) under the project INNOVAGEN and from the European Union's Horizon 2020 research
249 and innovation programme under grant agreement No 634476 (Project acronym: TREASURE). The
250 content of this paper reflects only the authors' view and the European Union Agency is not
251 responsible for any use that may be made of the information it contains. This study was associated
252 with the PSRN program (2017-2019).

253

254 **Conflict of interest statement**

255 The authors declare that there is no conflict of interest regarding the publication of this paper.

256

257 **References**

258 Akiyama, H., Chaboissier, M.C., Martin, J.F., Schedl, A., de Crombrughe, B. 2002. The
259 transcription factor Sox9 has essential roles in successive steps of the chondrocyte
260 differentiation pathway and is required for expression of Sox5 and Sox6. *Genes Dev.* 16, 2813–
261 2828.

262 ANAS, 2016. *Registro Anagrafico*. <http://www.anas.it/>.

263 Aulchenko, Y.S., Ripke, S., Isaacs, A., van Duijn C.M., 2007. GenABEL: an R library for genome-
264 wide association analysis. *Bioinformatics* 23, 1294–1296.

265 Bagheri-Fam, S., Barrionuevo, F., Dohrmann, U., Günther, T., Schüle, R., Kemler, R., Mallo, M.,
266 Kanzler, B., Scherer, G. 2006. Long-range upstream and downstream enhancers control distinct
267 subsets of the complex spatiotemporal Sox9 expression pattern. *Dev. Biol.* 291, 382–397.

268 Barrionuevo, F., Taketo, M. M., Scherer, G., Kispert, A. 2006. Sox9 is required for notochord
269 maintenance in mice. *Dev. Biol.* 295, 128–140.

270 Bracke, M.B.M., Hulsege, B., Keeling, L., Blokhuis, H.J. 2004. Decision support system with
271 semantic model to assess the risk of tail biting in pigs. 1. Modelling. *Appl. Anim. Behav. Sci.*
272 87, 31–44.

273 Brooksbank, N.H. 1958. Congenital deformity of the tail in pigs. *Br. Vet. J.* 114, 50–55.

274 Carneiro, M., Rubin, C.J., Di Palma, F., Albert, F.W., Alföldi, J., Martinez Barrio, A., Pielberg, G.,
275 Rafati, N., Sayyab, S., Turner-Maier, J., Younis, S., Afonso, S., Aken, B., Alves, J.M., Barrell,
276 D., Bolet, G., Boucher, S., Burbano, H.A., Campos, R., Chang, J.L., Duranthon, V., Fontanesi,
277 L., Garreau, H., Heiman, D., Johnson, J., Mage, R.G., Peng, Z., Queney, G., Rogel-Gaillard,
278 C., Ruffier, M., Searle, S., Villafuerte, R., Xiong, A., Young, S., Forsberg-Nilsson, K., Good,
279 J.M., Lander, E.S., Ferrand, N., Lindblad-Toh, K., Andersson, L. 2014. Rabbit genome analysis
280 reveals a polygenic basis for phenotypic change during domestication. *Science* 345, 1074–
281 1079.

282 Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J. 2015. Second-generation
283 PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 4, 7.

284 Clutton-Brock, J. 1999. *A Natural History of Domesticated Mammals*. 2nd Edition, Cambridge, UK:
285 Cambridge University Press.

286 Copp, A. J., Brook, F. A., & Roberts, H. J. 1988. A cell-type-specific abnormality of cell proliferation
287 in mutant (curly tail) mouse embryos developing spinal neural tube defects. *Development* 104,
288 285–295.

289 Darwin, C. 1868. *The Variation of Animals and Plants under Domestication*. London, UK: John
290 Murray.

291 Donald, H. P., 1949. The inheritance of a tail abnormality associated with urogenital disorders in pigs.
292 *J. Agric. Sci.* 39, 164–173.

293 Fontanesi, L., Schiavo, G., Galimberti, G., Calò, D.G., Scotti, E., Martelli, P.L., Buttazzoni, L.,
294 Casadio, R., Russo, V. 2012. A genome wide association study for backfat thickness in Italian
295 Large White pigs highlights new regions affecting fat deposition including neuronal genes.
296 *BMC Genomics* 13, 583.

297 Foster, J.W., Dominguez-Steglich, M.A., Guioli, S., Kwok, C., Weller, P.A., Stevanović, M.,
298 Weissenbach, J., Mansour, S., Young, I.D., Goodfellow, P.N., et al. 1994. Campomelic
299 dysplasia and autosomal sex reversal caused by mutations in an SRY-related gene. *Nature* 372,
300 525–530.

301 Healy, C., Uwanogho, D., Sharpe, P.T. 1996. Expression of the chicken Sox9 gene marks the onset
302 of cartilage differentiation. *Ann. N. Y. Acad. Sci.* 785, 261–262.

303 Ikegami, D., Akiyama, H., Suzuki, A., Nakamura, T., Nakano, T., Yoshikawa, H., Tsumaki, N. 2011.
304 Sox9 sustains chondrocyte survival and hypertrophy in part through Pik3ca-Akt pathways.
305 *Development* 138, 1507–1519.

306 Kurth, I., Klopocki, E., Stricker, S., van Oosterwijk, J., Vanek, S., Altmann, J., Santos, H.G., van
307 Harssel, J.J., de Ravel, T., Wilkie, A.O., Gal, A., Mundlos, S. 2009 Duplications of
308 noncoding elements 5' of SOX9 are associated with brachydactyly-anonychia. *Nat. Genet.*
309 41, 862–863.

310 Lahrmann, H. P., Hansen, C. F., D'Eath, R., Busch, M. E., & Forkman, B. (2018). Tail posture
311 predicts tail biting outbreaks at pen level in weaner pigs. *Appl. Anim. Behav. Sci.* 200, 29–35.

312 Larson, G., Burger, J. 2013. A population genetics view of animal domestication. *Trends Genet.* 29,
313 197–205.

314 Leipoldt, M., Erdel, M., Bien-Willner, G.A., Smyk, M., Theurl, M., Yatsenko, S.A., Lupski, J.R.,
315 Lane, A.H., Shanske, A.L., Stankiewicz, P., Scherer, G. 2007. Two novel translocation
316 breakpoints upstream of SOX9 define borders of the proximal and distal breakpoint cluster
317 region in campomelic dysplasia. *Clin. Genet.* 71, 67–75.

318 Leroy, G., Besbes, B., Boettcher, P., Hoffmann, I., Capitan, A., Baumung, R. 2016. Rare phenotypes
319 in domestic animals: unique resources for multiple applications. *Anim. Genet.* 47, 141–153.

320 Montero, J.A., Lorda-Diez, C.I., Francisco-Morcillo, J., Chimal-Monroy, J., Garcia-Porrero, J.A.,
321 Hurle, J.M. 2017. Sox9 expression in Amniotes: Species-specific differences in the formation
322 of digits. *Front. Cell Dev. Biol.* 5, 23.

323 Nordby J.E. 1934. Kinky tail in swine. *J. Hered.* 25, 171–174.

324 Ohnishi, T., Miura, I., Ohba, H., Shimamoto, C., Iwayama, Y., Wakana, S., Yoshikawa, T. 2017. A
325 spontaneous and novel Pax3 mutant mouse that models Waardenburg syndrome and neural tube
326 defects. *Gene* 607, 16–22.

327 Pfeifer, D., Kist, R., Dewar, K., Devon, K., Lander, E.S., Birren, B., Korniszewski, L., Back, E.,
328 Scherer, G. 1999. Campomelic dysplasia translocation breakpoints are scattered over 1 Mb
329 proximal to SOX9: evidence for an extended control region. *Am. J. Hum. Genet.* 65, 111–124.

330 Porter, V. 1993. *Pigs: A Handbooks to the Breeds of the World*. Cornell University Press,

331 Sanchez M.P., Tribout T., Iannuccelli N., Bouffaud, M., Servin, B., Tenghe, A., Dehais, P., Muller,
332 N., Del Schneider, M.P., Mercat, M.J., Rogel-Gaillard, C., Milan, D., Bidanel, J.P., Gilbert, H.
333 2014. A genome-wide association study of production traits in a commercial population of
334 Large White pigs: evidence of haplotypes affecting meat quality. *Genet. Sel. Evol.* 46, 12.

335 Schiavo, G., Bertolini, F., Utzeri, V.J., Ribani, R., Geraci, C., Santoro, L., Óvilo, C., Fernández, A.I.,
336 Gallo, M., Fontanesi, L. 2018. Taking advantage from phenotype variability in a local animal

337 genetic resource: identification of genomic regions associated with the hairless phenotype in
338 Casertana pigs. *Anim. Genet.* 49, 321–325.

339 Trut, L., Oskina, I., Kharlamova, A. 2009. Animal evolution during domestication: the domesticated
340 fox as a model. *Bioessays* 31, 349–360.

341 van Straaten, H. W., Copp, A. J. 2001. Curly tail: a 50-year history of the mouse spina bifida model.
342 *Anat. Embryol.* 203, 225–238.

343 Vaysse, A., Ratnakumar, A., Derrien, T., Axelsson, E., Rosengren Pielberg, G., Sigurdsson, S., Fall,
344 T., Seppälä, E.H., Hansen, M.S., Lawley, C.T., Karlsson, E.K.; LUPA Consortium, Bannasch,
345 D., Vilà, C., Lohi, H., Galibert, F., Fredholm, M., Häggström, J., Hedhammar, A., André, C.,
346 Lindblad-Toh, K., Hitte, C., Webster, M.T. 2011. Identification of genomic regions associated
347 with phenotypic variation between dog breeds using selection mapping. *PLoS Genet.* 7,
348 e1002316.

349 Velagaleti, G.V., Bien-Willner, G.A., Northup, J.K., Lockhart, L.H., Hawkins, J.C., Jalal, S.M.,
350 Withers, M., Lupski, J.R., Stankiewicz, P. 2005. Position effects due to chromosome
351 breakpoints that map approximately 900 Kb upstream and approximately 1.3 Mb downstream
352 of SOX9 in two patients with campomelic dysplasia. *Am. J. Hum. Genet.* 76, 652–662.

353 Wagner, T., Wirth, J., Meyer, J., Zabel, B., Held, M., Zimmer, J., Pasantes, J., Bricarelli, F.D., Keutel,
354 J., Hustert, E., Wolf, U., Tommerup, N., Schempp, W., Scherer, G. 1994. Autosomal sex
355 reversal and campomelic dysplasia are caused by mutations in and around the SRY-related gene
356 SOX9. *Cell* 79, 1111–1120.

357 Wang, G.D., Xie, H.B., Peng, M.S., Irwin, D., Zhang, Y.P. 2014. Domestication genomics: evidence
358 from animals. *Annu. Rev. Anim. Biosci.* 2, 65–84.

359 Wiener, P., Wilkinson, S. 2011. Deciphering the genetic basis of animal domestication. *Proc. Biol.*
360 *Sci.* 278, 3161–3170.

361 Wilkins, A. S., Wrangham, R. W., Fitch, W. T. 2014. The “domestication syndrome” in mammals: a
362 unified explanation based on neural crest cell behavior and genetics. *Genetics* 197, 795–808.

363 Wright, E., Hargrave, M.R., Christiansen, J., Cooper, L., Kun, J., Evans, T., Gangadharan, U.,
364 Greenfield, A., Koopman, P. 1995. The Sry-related gene Sox9 is expressed during
365 chondrogenesis in mouse embryos. *Nat. Genet.* 9, 15–20.

366 Wunderle, V.M., Critcher, R., Hastie, N., Goodfellow, P.N., Schedl, A. 1988. Deletion of long-range
367 regulatory elements upstream of SOX9 causes campomelic dysplasia. *Proc. Natl. Acad. Sci*
368 *USA* 95, 10649–10654.

369 Zhou, X., Stephens, M. 2012. Genome-wide efficient mixed-model analysis for association studies.
370 *Nat. Genet.* 44, 821–824.

371 Zonderland, J.J., van Riel, J.W., Bracke, M.B.M., Kemp, B., den Hartog, L.A., Spoolder, H.A.M.,
372 2009. Tail posture predicts tail damage among weaned piglets. *Appl. Anim. Behav. Sci.* 121,
373 165–170.

374 **Figure 1.** Tail shape of Casertana pigs: a) curly tail; b) straight tail.

375

a)




b)



376


377

378 **Figure 2.** Multidimensional scaling (MDS) with represented the pigs (dots) included in this study
379 divided in the two groups of tail shape.



380

381 **Figure 3.** Manhattan plot obtained for the genome wide association study.




382

383 **Supplementary material**

384


385 **Figure S1.** Quantile–quantile (Q–Q) plot obtained from the genome wide association analysis.



386

387

388 **Figure S2.** Graphical representation of the annotated genes in the porcine chromosome 12 region
 389 around the ALGA0064877 (rs81439488 C/A; nucleotide position 10,301,075. The screenshot has
 390 been retrieved from the ENSEMBL database (<http://www.ensembl.org/index.html>; release 93, July
 391 2018).



392

393 **Table S1.** List of genes annotated in the Sscrofa11.1 genome version around the ALGA0064877
 394 (rs81439488 C/A; nucleotide position 10,301,075) marker (2 Mbp on both directions) on porcine
 395 chromosome 12.

Position	Gene symbol	Gene name	Role/function ¹
8,641,629-8,647,764	<i>SOX9</i>	SRY-box 9	The encode protein is involved in chondrocyte differentiation and, with steroidogenic factor 1, regulates transcription of the anti-Muellerian hormone (AMH) gene. Deficiencies lead to the skeletal malformation syndrome campomelic dysplasia
10,351,715-10,362,244	<i>KCNJ2</i>	potassium voltage-gated channel subfamily J member 2	The protein is an integral membrane protein and inward-rectifier type potassium channel. It probably participates in establishing action potential waveform and excitability of neuronal and muscle tissues. Mutations in this gene have been associated with Andersen syndrome in humans, which is characterized by periodic paralysis, cardiac arrhythmias, and dysmorphic features.
10,393,179-10,448,536	<i>KCNJ16</i>	potassium voltage-gated channel subfamily J member 16	Similar to the function of the previous gene product. It may act in fluid and pH balance regulation.
10,897,104-11,018,943	<i>MAP2K6</i>	mitogen-activated protein kinase kinase 6	It encodes for a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis.
11,018,687-11,152,913	<i>ABCA5</i>	ATP binding cassette subfamily A member 5	The encode protein is is a member of the superfamily of ATP-

			binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intracellular membranes. Its function is not well known.
11,169,809-11,211,801	<i>ABCA10</i> (ENSSSCG00000032844)	ATP-binding cassette sub-family A member 10	Similar to the previous gene.
11,225,408-11,286,018	<i>ABCA6</i>	ATP binding cassette subfamily A member 6	Similar to the previous gene.
11,295,697-11,359,520	<i>ABCA9</i>	ATP-binding cassette sub-family A member 9	Similar to the previous gene.
11,374,028-11,447,292	<i>ABCA8</i>	ATP-binding cassette sub-family A member 8	Similar to the previous gene.
11,607,233-11,657,737	<i>FAM20A</i>	FAM20A, golgi associated secretory pathway pseudokinase	This gene encodes a protein that is likely secreted and may function in hematopoiesis. A mutation at this locus has been associated with amelogenesis imperfecta and gingival hyperplasia syndrome in humans.
11,661,872-11,681,435	<i>PRKARIA</i>	protein kinase cAMP-dependent type I regulatory subunit alpha	This gene encodes for one of the regulatory subunits of the kinase holoenzyme. This protein was found to be a tissue-specific extinguisher that down-regulates the expression of several liver genes. Mutations in this gene cause Carney complex (CNC) in humans. Other functions have been inferred based on the protein structure.
11,715,274-11,752,377	<i>WIPI1</i>	WD repeat domain, phosphoinositide interacting 1	It encodes a WD40 repeat protein, which is a key components of many essential biologic functions, by regulating the assembly of multiprotein complexes.
11,826,578-11,827,867	ENSSSCG00000034083	Novel gene	Not known
11,857,792-11,875,313	<i>SLC16A6</i>	solute carrier family 16 member 6	Not well defined.
11,882,800-11,895,486	<i>AMZ2</i>	archaelysin family metalloproteinase 2	The encoded protein is a zinc metalloprotease that displays some activity against angiotensin-3. The encoded protein is inhibited by the aminopeptidase inhibitor amastatin, as well as by the general inhibitors o-phenanthroline and batimastat. Defects in this

			gene may be associated with lung tumorigenesis.
11,922,572-11,966,747	<i>GNA13</i>	G protein subunit alpha 13	Not well defined.
11,996,142-12,089,466	<i>RGS9</i>	regulator of G protein signaling 9	This gene encodes a member of the RGS family of GTPase activating proteins that function in various signaling pathways by accelerating the deactivation of G proteins. This protein is anchored to photoreceptor membranes in retinal cells and deactivates G proteins in the rod and cone phototransduction cascades. Mutations in this gene result in bradyopsia in humans.

396 ¹ Information has been adapted from the NCBI Gene database (<https://www.ncbi.nlm.nih.gov/gene/>; August 2018).

397

398