

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Protease-activated receptor 1 is implicated in irritable bowel syndrome mediators-induced signaling to thoracic human sensory neurons

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Protease-activated receptor 1 is implicated in irritable bowel syndrome mediators-induced signaling to thoracic human sensory neurons / Desormeaux, Cleo; Bautzova, Tereza; Garcia-Caraballo, Sonia; Rolland, Corinne; Barbaro, Maria Raffaella; Brierley, Stuart M.; Barbara, Giovanni; Vergnolle, Nathalie; Cenac, Nicolas. - In: PAIN. - ISSN 0304-3959. - ELETTRONICO. - 159:7(2018), pp. 1257-1267. [10.1097/j.pain.0000000000001208]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/674755> since: 2019-02-26

*Published:*

DOI: <http://doi.org/10.1097/j.pain.0000000000001208>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the accepted manuscript of:

Desormeaux C, Bautzova T, Garcia-Caraballo S, Rolland C, Barbaro MR, Brierley SM, Barbara G, Vergnolle N, Cenac N.

Protease-activated receptor 1 is implicated in irritable bowel syndrome mediators-induced signaling to thoracic human sensory neurons.

Pain. 2018 Jul;159(7):1257-1267.

Final peer reviewed version available at: <https://doi.org/10.1097/j.pain.0000000000001208>

#### Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

## Protease-Activated Receptor 1 is implicated in irritable bowel syndrome mediators-induced signaling to thoracic human sensory neurons

Cleo Desormeaux<sup>1</sup>, Tereza Bautzova<sup>1</sup>, Sonia Garcia-Caraballo<sup>2,3</sup>, Corinne Rolland<sup>1</sup>, Maria Raffaella Barbaro<sup>4</sup>, Stuart M. Brierley<sup>2,3</sup>, Giovanni Barbara<sup>4</sup>, Nathalie Vergnolle<sup>1,5\*</sup>§, Nicolas Cenac<sup>1§</sup>.

<sup>1</sup> IRSD, Université de Toulouse, INSERM, INRA, ENVT, UPS, Toulouse, France.

<sup>2</sup> Visceral Pain Group, Human Physiology, Flinders University, Bedford Park, South Australia, 5042, Australia

<sup>3</sup> Centre for Nutrition and Gastrointestinal Diseases, Discipline of Medicine, University of Adelaide, South Australian Health and Medical Research Institute (SAHMRI), North Terrace, Adelaide, South Australia 5000, Australia

<sup>4</sup> Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy.

<sup>5</sup> Department of Physiology and Pharmacology, Faculty of Medicine, University of Calgary, Calgary, Canada

§ Equal contribution

\* Correspondance: Dr. Nathalie Vergnolle, Inserm UMR-1220, Institut de Recherche en Santé Digestive, CS60039 CHU Purpan, 31024 Toulouse, Cedex-3, France. E-mail address: [nathalie.vergnolle@inserm.fr](mailto:nathalie.vergnolle@inserm.fr)

The authors have declared that no conflict of interest exists.

Word count: 3982

Abstract : words 243

Proteases and Protease-Activated Receptors (PARs) are major mediators involved in irritable bowel syndrome (IBS). Our objectives were to decipher the expression and functionality (calcium signaling) of PARs in human dorsal root ganglia (DRG) neurons, and to define mechanisms involved in human sensory neuron signaling by IBS patient mediators.

Human thoracic DRG were obtained from the national disease resource interchange. Expression of PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> was assessed by immunohistochemistry and RT-qPCR in whole DRG or in primary cultures of isolated neurons. Calcium signaling in response to PAR agonist peptides (PAR-AP), their inactive peptides (PAR-IP), thrombin (10u/ml), supernatants from colonic biopsies of IBS patients or healthy controls (HC), with or without PAR<sub>1</sub> or PAR<sub>4</sub> antagonist were studied in cultured human DRG neurons.

PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> were all expressed in human DRG, respectively in 20%, 40% and 40% of the sensory neurons. PAR<sub>1</sub>-AP increased intracellular calcium concentration in a dose-dependent manner. This increase was inhibited by PAR<sub>1</sub> antagonism. In contrast, PAR<sub>2</sub>-AP, PAR<sub>4</sub>-AP and PAR-IP did not cause calcium mobilization. PAR<sub>1</sub>-AP-induced calcium flux was significantly reduced by pre-incubation with PAR<sub>4</sub>-AP, but not with PAR<sub>2</sub>-AP. Thrombin increased calcium flux, which was inhibited by a PAR<sub>1</sub> antagonist and increased by a PAR<sub>4</sub> antagonist. Supernatants from colonic biopsies of IBS patients induced calcium flux in human sensory neurons compared to HC, this induction was reversed by a PAR<sub>1</sub> antagonist.

Taken together, our results highlight that PAR<sub>1</sub> antagonism should be investigated as a new therapeutic target for IBS symptoms.

**Keywords:** Proteases; PARs; Protease-Activated Receptors; Visceral pain; Inflammation; Irritable Bowel Syndrome; Visceral hypersensitivity; Thrombin; Human dorsal root ganglia neurons.

## Introduction

Irritable Bowel Syndrome (IBS) affects 11 to 20% of the Western population with a higher prevalence in women [14; 31]. IBS associates abdominal pain, diarrhea (IBS-D), constipation (IBS-C) or both (IBS-A, for alternate) [30]. Although IBS is a functional gastrointestinal disorder, not associated with gross structural or biochemical abnormalities [23], several recent studies indicate the presence of meaningful micro-organic changes [7]. One of the emerging ideas to explain the visceral pain associated with IBS is that sensory neurons innervating the colon are hyperexcitable in these patients [10; 11; 13]. However, because of the difficulties associated with human sensory neuron cultures, it has been difficult to evaluate the relevance of identified mediators in the context of human pathology.

Among the molecular targets explored to decipher neuronal hyperexcitability in IBS, several studies showed that proteases released by colonic biopsies of IBS patients were able to activate mouse and rat intestinal neurons *in vitro* and to induce somatic and visceral hypersensitivity *in vivo* [9; 13; 17; 38; 47]. Proteases are known to signal to mouse or rat sensory neurons through the activation of Protease-Activated Receptors (PARs) [19; 46; 53; 55], a family of G protein-coupled receptors that includes 4 members: PAR<sub>1</sub>, PAR<sub>2</sub>, PAR<sub>3</sub> and PAR<sub>4</sub> [40]. Only PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> seem to be able to signal through calcium mobilization and to exert a role in nociception and pain (PAR<sub>3</sub> has been considered more as a co-factor for other PAR activation[35] so far). PAR<sub>1</sub>, 2 and 4 are activated by the proteolytic cleavage of their N-terminal domain, which reveals a tethered ligand that binds and activates the receptors. The role of PARs in visceral inflammation and pain has been well studied in animal models[49; 50]. Both PAR<sub>1</sub> and PAR<sub>2</sub> agonists induce calcium mobilization in rodent sensory neurons[17; 21]. PAR<sub>2</sub> agonists induce pain in somatic and visceral models [51; 53], while PAR<sub>1</sub> and PAR<sub>4</sub> agonists attenuate nociception and pain symptoms in rodents [4-6; 32]. In mouse sensory neurons, the calcium mobilization induced by supernatants from colonic biopsies of IBS patients is dependent on supernatant's proteolytic activity and on PAR<sub>2</sub> expression in mouse primary afferents [17]. In contrast, PAR<sub>4</sub> activation inhibits PAR<sub>2</sub>-induced calcium mobilization and intrinsic excitability of colonic dorsal root ganglia (DRG) neurons, as well as overall pain [5; 6; 28]. However, in the context of human nociception and pain, the effects of PAR agonists and PAR signaling mechanisms are largely unknown.

Proteases show different specificity for the different PARs[52]. For instance, thrombin can activate PAR<sub>1</sub>, PAR<sub>3</sub>, PAR<sub>4</sub> and to a lower extent PAR<sub>2</sub>, trypsin can activate PAR<sub>2</sub>, PAR<sub>3</sub> and PAR<sub>4</sub> and at high concentrations PAR<sub>1</sub> [25; 33; 42]. In the context of IBS, tryptase and trypsin-3 are up-regulated, and are considered as possible endogenous agonists for PARs[9; 43].

Our study aimed at deciphering PAR signaling in human sensory neurons, determining whether proteases and PARs could play a role in human sensory neuron biology and nociceptive mechanisms. Further, we investigated human sensory neuron responses to IBS mediators in cultured human DRG neurons. Although not only neurons projecting from the digestive tract are present in our cultures, the present study could give insights in the response of human primary afferents to mediators present in IBS patient tissues.

## Methods

### Chemicals

Each agonist and inactive peptides of PARs, respectively PAR-AP and PAR-IP, were purchased from Genscript (France): PAR<sub>1</sub>-AP (TFFLR), PAR<sub>2</sub>-AP (SLIGKV), PAR<sub>4</sub>-AP (GYPGQV), PAR<sub>1</sub>-IP (RLLFT), PAR<sub>2</sub>-IP (LRGILS) and PAR<sub>4</sub>-IP (YAPGQV). Thrombin, PAR<sub>1</sub>-antagonist (SCH79797), PAR<sub>2</sub> antagonist (GB83) and PAR<sub>4</sub>-antagonist (ML354) were obtained from Tocris (Denver, USA).

### Patient biopsies and supernatant collection

Colon biopsy samples from 24 patients with IBS (8 IBS-D, 8 IBS-C, 8 IBS-A) and 5 healthy controls, HCs) undergoing colorectal cancer screening were collected during colonoscopy at the Department of Medical and Surgical Sciences of the University of Bologna, Italy (Supplementary Table 1, available online at <http://links.lww.com/PAIN/A553>). Rome III criteria were used for the diagnosis of IBS patients. Additional exclusion criteria were major abdominal surgery, celiac disease, asthma, allergic disorders, anti-inflammatory treatments, organic syndrome. Symptoms as bloating, pain and bowel habit changes in the last 12 months were also excluded from control group.

Supernatants from colonic biopsies were obtained following a previously validated and published method [8], with few modifications. Briefly, after removal, biopsies were immersed in plastic tubes containing 1 ml of Hepes-Krebs solution. After weighing the biopsies, supernatant volume was adjusted to incubate 15 mg of biopsies in 1 ml of buffer. Incubation was carried out in oxygenation at 37°C for 25 min. Samples were centrifuged at 200g for 10 min and supernatant collected and stored at –20°C until the assay.

#### **Human Dorsal Root Ganglia Neurons Isolation**

Experiments were conducted under the Institutional Review Board numbers IRB00003888, FWA00005831. Human DRG (thoracic position 11 and 12) were collected in Dulbecco's Modified Eagle's medium by the National Disease Resource Interchange (NDRI). Twenty DRG were obtained from 10 post-mortem donors (21-60 years old, 10 hours maximum post-mortem) with the following exclusion criteria: chemotherapy, drug abuse, infectious disease, neurodegenerative diseases and opioid medications. None of the donors had a reported history of colitis or inflammatory bowel disease, but no information was provided about possible IBS. DRG were cut in small pieces, rinsed in Hank's balanced salt solution (HBSS; Thermo Fisher, Villebon-sur-Yvette, France) and digested in L-Cystein (pH 7.4, Sigma Aldrich, Missouri, USA), Papain (27 µg/ml, Sigma Aldrich, Missouri, USA) for 20 minutes at 37°C, rinsed 2 minutes in Leibovitz's L15 Medium (Thermo Fisher Scientific, Waltham, Massachusetts, USA) containing 10 % of FBS. A second enzymatic dissociation was performed in 4 mg/ml dispase II (Sigma Aldrich, Missouri, USA), and 1 mg/ml collagenase (collagenase (type IV, Serlabo Technologies, France) for 15 minutes at 37°C, followed by mechanical dissociation. This step was repeated until complete dissociation of the DRG up to 4 times. Finally, neurons were plated in 8-wells Nunc™ Lab-Tek™ II CC2™ chamber slide system (Thermo Fisher Scientific, Waltham, Massachusetts, USA) and cultured for 7 days in complete Neurobasal-A medium (Thermo Fisher Scientific, Waltham, Massachusetts, USA) containing penicillin (100 µg/ml), streptomycin (100µg/ml), B27 (1X, Thermo Fisher Scientific, Waltham, Massachusetts, USA), L-glutamine (1X, Thermo Fisher Scientific, Waltham, Massachusetts, USA), and inhibitors of mitosis (Cytosine-B-arabinofuranoside 1.5µM, F-Uridine 10µM, Uridine, 10µM, Sigma Aldrich, Missouri, USA). Cytosine-B-arabinofuranoside was removed after 3 days of culture and the medium was changed every 2 days.

For RT-qPCR and single-cell PCR studies, human thoracolumbar DRG (T9-L1) were acquired from five (three females, two males) human adult organ donors ( $22.2 \pm 2.08$  years of age) during the removal of vital organs for transplantation. The harvested intact DRG were kept for quantitative-reverse-transcription-PCR (RT-qPCR) mRNA expression studies, while additional DRG were dissociated to allow individual DRG neurons to be isolated and studied with single-cell-reverse-transcription-PCR (RT-PCR).

### **Calcium imaging of human sensory neurons**

After washing with HBSS, cultured neurons were incubated for 1 hour (30 min at 37°C followed by 30 min at room temperature) in solution containing fluo-4 acetoxymethyl (AM) 1mM (Thermo Fisher Scientific, Waltham, Massachusetts, USA) reconstituted in 0.01% pluronic F-127 and 0.7% DMSO (Sigma Aldrich, Missouri, USA). The fluorescence was measured at 460-490 nm excitation and 515 nm emission in each well. Neurons were imaged using an inverted microscope (Zeiss, 10x-objective) and a CCD camera (Zeiss). Acquisition parameters were kept constant within each experiment. A kinetic of 80 recordings (one per second) was performed. From 0 to 5-sec basal fluorescence was determined, from 6 to 60-sec, neurons were exposed to the different molecules studied and finally to a KCl solution (50 mM), in order to discriminate neurons from glial cells. Neurons were identified by Image J software and variations in the fluorescence intensity of each neuron was measured. Results were expressed as  $\Delta F/F$ , representing the fluorescence intensity ratio between the highest measure during the 6 to 60 seconds period, and baseline measure.

In a first set of experiments, neurons were treated with individual PAR agonist peptides: PAR<sub>1</sub>-AP (1, 10, 50 or 100  $\mu$ M), PAR<sub>2</sub>-AP (100  $\mu$ M), PAR<sub>4</sub>-AP (100  $\mu$ M) or neurons were pre-incubated for 5 minutes with PAR<sub>2</sub>-AP (100  $\mu$ M) or PAR<sub>4</sub>-AP (100  $\mu$ M), before being exposed to PAR<sub>1</sub>-AP (100  $\mu$ M). For these experiments, the inactive peptides (PAR-IP, 100  $\mu$ M) and vehicle (HBSS) were used as control. In a second set of experiments, neurons were pre-incubated for 5 minutes with antagonist of PAR<sub>1</sub> (SCH79797, 10  $\mu$ M) or its vehicle (HBSS, 0.01% DMSO) and treated with PAR<sub>1</sub>-AP (100  $\mu$ M). In a third set of experiments, neurons were pretreated 5 minutes with the PAR<sub>1</sub> antagonist (SCH79797, 10  $\mu$ M), the PAR<sub>4</sub> antagonist (ML354, 10  $\mu$ M) or vehicle (HBSS, 0.01% DMSO) and were then treated with thrombin (10



U/ml). In a last set of experiments, neurons were pretreated 5 minutes with the PAR<sub>1</sub> antagonist (SCH79797, 10 $\mu$ M) or vehicle, before being exposed to supernatants from colonic biopsies of IBS-D, IBS-C, IBS-A patients or HCs.

### **Immunofluorescence in human Dorsal Root Ganglia**

Experiments were conducted under the Institutional Review Board numbers IRB00003888, and FWA00005831. Three Human DRG T11 and 3 DRG T12 (thoracic position 11 and 12) were cryoprotected in Tissue-Tek<sup>®</sup> optimum cutting temperature compound (Sakura Finetek, Netherlands) after their collection by the NDRI. Cryoprotected DRG were cut into 35  $\mu$ m sections in a cryostat (Leica CM1950; Nanterre, France) and mounted on Superfrost slides (Thermo Fisher Scientific, Villebon-sur-Yvette, France). Cultured DRG neurons were fixed with paraformaldehyde 4% during 20 min. Both slides and cultures were washed in Phosphate Buffered Saline (PBS), 0.5% Triton X-100, and 1% Bovine Serum Albumin solution (BSA, Sigma Aldrich, Missouri, USA) and were incubated overnight at 4°C with primary antibodies diluted at 1:100 for tissues and 1:500 for cultures and directed against PGP9.5, PAR<sub>1</sub>, PAR<sub>4</sub> (respectively, AB86808, AB111976, AB70400, Abcam, Cambridge, England) and PAR<sub>2</sub> (LSB2321, LifeSpan, Seattle, USA). After washing in PBS, slices or cultures were incubated with the appropriate secondary antibody conjugated to Alexa Fluor 488 or Alexa Fluor 555, they were washed, and mounted with ProLong Gold reagent containing 40,6-diamidino-2-phenylindole (DAPI, Molecular Probes). Controls for the specificity of the antibodies include incubation in the absence of secondary antibody, incubation in the presence of immunizing peptides and the use of PAR-deficient tissues or cells (not shown). Images were acquired using Zeiss LSM-710 confocal microscopes (Carl Zeiss MicroImaging, Jena, Germany) with 10x objective in the inverted configuration [20]. Quantification of labelling was determined using Image J software.

### **Quantitative-reverse-transcription-PCR (RT-qPCR):**

RNA was extracted from either whole human DRG or single human DRG neurons using RNA-isolation kits (PureLink<sup>™</sup> and Cells-to-CT<sup>™</sup>; Ambion). RT-qPCR was performed using human-specific Taqman primers for PAR<sub>1</sub>, PAR<sub>2</sub>, PAR<sub>4</sub> and GAPDH (Hs00169258\_m1, Hs00608346\_m1, Hs00765740\_m1, Hs01006385\_g1, Hs99999905\_m1). The comparative

cycle threshold method was used to quantify the abundance of target transcripts to reference genes.

#### **Single cell PCR:**

26 single dissociated DRG neurons were picked using a micromanipulator at 40x magnification. Cells were under a continuous slow flow of sterile and RNA/DNAse free PBS to reduce contamination. After a cell was picked, the glass capillary was broken into a tube containing 10ul of Lysis buffer and DNAse (TaqMan® Gene Expression Cells-to-CT™ Kit; Ambion). The whole content was used for cDNA synthesis (SuperScript® VILO™ cDNA Synthesis Kit, Thermo Fisher) and PAR<sub>1</sub>, PAR<sub>2</sub>, PAR<sub>4</sub> expression was measured using Taqman™ RT-qPCR for 50 cycles. For every coverslip, a bath control was taken and analyzed together with samples. TUBB3 (Hs00964962\_g1) expression served as positive control. One cell was excluded because no TUBB3 expression was present. Twenty-six cells were used to calculate frequency of presence of PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub>.

#### **Statistical Analysis**

Data are presented as means ± standard error of the mean (SEM). Analyses were performed using GraphPad Prism 5.0 software (GraphPad, San Diego, CA). Comparisons between-groups were performed by Mann-Whitney test or Wilcoxon matched pairs test. Multiple comparisons within groups were performed by Kruskal-Wallis test, followed by Dunn's post-test. Statistical significance was accepted at  $P < 0.05$ .

#### **Study approval**

The collection of biopsies from colonic patients was approved by the local Ethic Committee (64/2004/O/Sper and EM14/2006/O) and conducted in accordance with the Declaration of Helsinki. IBS patients and HCs gave their written and informed consent. Fixed and fresh human DRG were collected with the NDRI (reference: DCEN1 001), all human DRG trials were conducted following the opinion number 14-164 of the institutional review board (IRB00003888) of French institute of medical research and health.

## Results

### -PAR expression in human DRG neurons-

In whole human DRG, PAR mRNA expression was assessed by RT-qPCR analysis. The relative mRNA abundance of PAR<sub>2</sub> was the highest followed by PAR<sub>4</sub> and PAR<sub>1</sub> (Figure 1a). Then, to determine if PARs were expressed in neurons at the protein level, the percentage of neurons, identified by Pgp9.5 immunostaining and expressing each PAR was studied (Figure 1b). PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> were respectively expressed in 20%, 40% and 40% of neurons (Figure 1c). The percentage of PAR-positive neurons was expressed in function of the diameter of neurons. PAR<sub>1</sub> was preferentially expressed in 30 to 50  $\mu$ M diameter neurons, PAR<sub>2</sub> in neurons with a diameter < 30  $\mu$ M and PAR<sub>4</sub> in 30 to 50  $\mu$ M diameter neurons (Figure 1d).

PAR expression was then assessed in cultured human DRG neurons. Cultures were 90% pure for Pgp9.5 neuronal marker staining (Supplementary Figure 1, available online at <http://links.lww.com/PAIN/A553>). Single cell PCR experiments showed that PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> mRNA were respectively expressed in 20%, 25% and 17% of neurons (Figure 2a left panel). We found that 75% of PAR<sub>1</sub> expressing neurons also expressed PAR<sub>2</sub>, 100% of the PAR<sub>4</sub> expressing neurons also expressed PAR<sub>2</sub>, whilst 25% of the PAR<sub>1</sub> expressing neurons also expressed PAR<sub>4</sub> (Figure 2a right panel). PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> were respectively expressed at the protein level in 35%, 36% and 27% of neurons (Figure 2b and 2c). These results demonstrate that human DRG neurons in culture continue to express PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> and no major or significant difference was observed between the level of expression of PARs in whole DRGs and in cultured DRGs.

### -Effects of PAR agonists on calcium mobilization in human DRG neurons-

Calcium flux in response to specific synthetic agonist peptides of each PAR was quantified in human sensory neuron cultures. Only PAR<sub>1</sub>-activating peptide (PAR<sub>1</sub>-AP, 100- $\mu$ M) evoked a transient increase in [Ca<sup>2+</sup>]<sub>i</sub> that was maximal after 20 seconds and declined to baseline afterwards, compared to PAR<sub>2</sub>-AP and PAR<sub>4</sub>-AP (Figure 3a). This activation was characterized by an increased percentage of responding neurons (Figure 3b) and increased amplitude ( $\Delta F/F$ ) of response (Figure 3c), compared to its inactive peptide (PAR<sub>1</sub>-IP, 100- $\mu$ M). PAR<sub>1</sub>-AP-induced calcium mobilization was dose-dependent (Figure 3d) and abolished

by a PAR<sub>1</sub> antagonist (SCH79797, 10-μM) (Figure 3d). Agonists and inactive peptides of PAR<sub>2</sub> and PAR<sub>4</sub> (all at 100-μM) had no significant effect on calcium signaling in human cultured DRG (Figure 3a, 3b and 3c).

As PAR<sub>2</sub> and PAR<sub>4</sub> agonist peptides did not induce calcium mobilization, we tested their potential inhibitory role on PAR<sub>1</sub>-AP-induced calcium flux. Human DRG were pretreated with PAR<sub>2</sub>-AP or PAR<sub>4</sub>-AP and, 5 min after later, PAR<sub>1</sub>-AP was added (all agonists at 100 μM). The transient increase in [Ca<sup>2+</sup>]<sub>i</sub> induced by PAR<sub>1</sub>-AP was decreased by PAR<sub>4</sub>-AP but not by PAR<sub>2</sub>-AP pretreatment (Figure 3e). Both the percentage of responding neurons and the amplitude of response induced by a PAR<sub>1</sub> agonist were significantly reduced by PAR<sub>4</sub>-AP, but not by PAR<sub>2</sub>-AP pretreatment (Figure 3f and 3g).

#### **- Thrombin signals to human DRG neurons**

Considering the calcium mobilization responses of human sensory DRG neurons to PAR<sub>1</sub> and PAR<sub>4</sub> peptidic agonists, we investigated the effects of thrombin (10U/ml), a known endogenous activator of both PAR<sub>1</sub> and PAR<sub>4</sub>. The average amplitude of thrombin-induced calcium response in human DRG neurons was significantly increased compared to vehicle (Figure 4a). Pre-treatment with a PAR<sub>1</sub> antagonist (SCH79797, 10 μM) significantly reduced the effects of thrombin on the percentage of responding neurons. Pre-treatment with a PAR<sub>4</sub> antagonist (ML354, 10 μM) had no effect on this parameter (Figure 4b). Considering only neurons that responded to thrombin by mobilizing calcium (23% of all neurons), the amplitude of their response to thrombin was significantly increased in the presence of PAR<sub>4</sub> antagonist, but was not modified by a PAR<sub>1</sub> antagonist (Figure 4c). These results demonstrated that in human DRG neurons, thrombin activates both PAR<sub>1</sub> and PAR<sub>4</sub>, exerting opposite effects in terms of calcium mobilization.

#### **-IBS patient mediators mobilize calcium in human DRG neurons through a PAR<sub>1</sub>-dependent mechanism-**

IBS supernatants, but not supernatants from healthy control, evoked a transient increase in [Ca<sup>2+</sup>]<sub>i</sub> that was maximal after 30 seconds and declined to baseline afterwards (Figure 5a). Supernatants from colonic biopsies of IBS patients significantly increased the amplitude of the calcium flux response and the percentage of responding neurons compared to healthy control supernatants (Figure 5b and 5c). Neither the percentage of responding

neurons, nor the amplitude of the response of human neurons to IBS patient tissue biopsy supernatants correlated with abdominal pain scores or abdominal frequency scores (supplementary Figure 2, available online at <http://links.lww.com/PAIN/A553>). When considered by subgroups, only supernatants from IBS-A patients induced a significant increase in the number of responding neurons (Supplementary Figure 3, available online at <http://links.lww.com/PAIN/A553>). PAR<sub>1</sub> antagonist (SCH79797, 10 μM) pretreatment of human sensory neurons decreased the transient increase in [Ca<sup>2+</sup>]<sub>i</sub> induced by IBS supernatants (Figure 5d). The antagonist significantly reduced both the percentage of responding neurons to IBS supernatants (all IBS subgroups together) and the amplitude of their response (Figure 5e and 5f).

## Discussion

Since their discovery in rodent neurons [46; 53], PARs have been considered as new important therapeutic targets for the treatment of pain. A number of *in vivo* and *in vitro* studies have confirmed this potential role for PARs, particularly in the context of visceral pain and hypersensitivity [19; 49; 51; 52]. However, the relevance of considering PAR signaling as an important pathway for human pain has not been thoughtfully addressed thus far. Indeed, only one study performed in human subjects refers to a possible role for PAR<sub>2</sub> in pruritus[45], but no study has investigated the expression and functionality of PARs in human DRG neurons. Here, we provide evidences that PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> are all expressed in human sensory neurons. Furthermore, we showed that in culture, the expression of PARs is generally conserved and that the culture conditions we have defined for human DRG neurons can be used to investigate the functionality of human sensory neurons. Although our results provide new insights on human primary afferent signaling, the link to activation of pain pathways would clearly require further experiments.

Previous studies performed in rodent primary afferents have demonstrated the expression of the three functional PARs: PAR<sub>1</sub>, PAR<sub>2</sub> and PAR<sub>4</sub> [21; 46]. We confirmed here that those three receptors are also expressed in human primary afferent neurons both in cell bodies of whole DRG neurons and in cultures. In rodent sensory neurons, both PAR<sub>1</sub> and PAR<sub>2</sub> agonists induced calcium mobilization [15-17; 21]. In contrast, PAR<sub>4</sub> agonists did not

303 mobilize calcium, but decreased pro-nociceptive mediator-induced calcium signaling [6; 28].  
304 In human sensory neurons, we demonstrated that only PAR<sub>1</sub> agonist induced calcium  
305 mobilization. Like in rodents, in human DRG neurons, PAR<sub>4</sub> was not able to mobilize calcium,  
306 but decreased calcium mobilization induced by other agonists. Taken together, our results  
307 clearly demonstrated that both PAR<sub>1</sub> and PAR<sub>4</sub> were present and functional in human  
308 primary afferents, where they exerted opposing effects. The PAR peptide agonists used in  
309 the present study have been well characterized for their selectivity [26; 27]. The doses that  
310 were used for these peptide agonists in the present study are considered to be highly  
311 selective for their targeted receptors in cell culture assays. Therefore, we can reasonably  
312 think that the results obtained are truly representative of PAR selective activation. Indeed,  
313 this was confirmed for PAR<sub>1</sub> activation by incubation in the presence of the PAR<sub>1</sub> antagonist.  
314 At the concentrations used, both SCH79797 and ML354 antagonists were considered  
315 selective for inhibition of PAR<sub>1</sub> and PAR<sub>4</sub> respectively, compared to the inhibition of other  
316 PARs [1; 57]. However, PAR<sub>1</sub>-independent effects have also been described for the  
317 SCH79797 compound [22], and one cannot rule out that the effects of this antagonist alone  
318 cannot be due to MAPK inhibition as it has been shown in cell lines at similar concentrations  
319 [22]. The PAR<sub>4</sub> antagonist ML354 is potent only at micro-molar ranges, showing a reasonable  
320 selectivity for PAR<sub>4</sub>, but one cannot rule out at this concentration off-target binding for this  
321 antagonist [57].

322 Although DRG cultures were 90% pure for neurons, some glial cells or other  
323 supporting cells might still be present in our culture conditions. PARs are known to be  
324 expressed and functional in glial cells and to some extent, glial PAR activation might account  
325 for some of the calcium responses observed in our cultures.

326 Translating cellular signaling of sensory neurons into nociceptive response *in vivo* is  
327 complex, and the study of calcium signaling in primary afferents cannot fully reflect pain  
328 pathway activation. We found no correlation between pain severity scores and pain  
329 frequency scores with the percentage of responding neurons or with the amplitude of the  
330 response to IBS tissue supernatants. This could be due to the low number of samples we  
331 included in this study considering the precious nature of human DRG neuron cultures.  
332 Indeed, other animal studies suggest that calcium signals in sensory neurons are often  
333 associated with pro-nociceptive signals. Calcium mobilization and increasing excitability of

neurons reflected the sensitization of neurons associated to visceral hypersensitivity [12]. However, in rodents, while PAR<sub>1</sub> agonists mobilized calcium in primary afferents [21], they also increased nociceptive threshold and reduced inflammatory hyperalgesia [4; 32]. In the present study, we showed that PAR<sub>1</sub> specific activation mobilized calcium in human sensory neurons. The pro-nociceptive nature of this PAR<sub>1</sub> signal is supported by our subsequent observations demonstrating the involvement of PAR<sub>1</sub> activation in IBS patient biopsy supernatants-induced activation of human sensory neurons. Interestingly, in human submucosal and myenteric plexi, PAR<sub>1</sub>-AP (TFLLR), thrombin or supernatants of colonic biopsies from IBS patients were also able to induce spike discharges and calcium signaling. This reflected enteric neuronal excitability [29; 34]. This suggests that in humans, PAR<sub>1</sub> might be functional both in submucosal and myenteric neurons, as well as in primary afferents (per our results). In the context of IBS, this means that PAR<sub>1</sub> could contribute as our results suggested, to hyperexcitability of extrinsic sensory neurons leading to visceral hypersensitivity symptoms, but could also contribute to motor and secretory dysfunctions controlled by intrinsic enteric neurons. However, one has to be careful establishing a link between primary afferent response and activation of pain pathways, as in rodents, PAR<sub>1</sub> can be activated in primary afferents, but was shown to be analgesic. Therefore, the pro-nociceptive effects of PAR<sub>1</sub> agonists remain to be demonstrated in humans. Our study paves the way for studying in human clinical trials the potential that PAR<sub>1</sub> antagonists already developed for use in human could have at reducing pain and hyperalgesia.

In agreement with the results previously generated in mouse primary afferents [6; 28], we observed in human neurons as well, that PAR<sub>4</sub>-AP did not mobilize calcium, but significantly reduced calcium mobilization of stimulated sensory neurons. Here, we demonstrated for the first time an inhibitory effect for PAR<sub>4</sub> activation on human sensory pathways, suggesting that like in rodents [5; 6], PAR<sub>4</sub> activation could contribute at reducing pain and hypersensitivity. Taken together, our results highlight opposite effects for PAR<sub>1</sub> and PAR<sub>4</sub> activation in human primary sensory afferents. Considering that both receptors are activated by thrombin, although at different concentrations, it was important to investigate the overall effect of thrombin on human primary neuron calcium signaling. Thrombin is attracted to PAR<sub>1</sub> by a Hirudin-like site located at the N-terminal end of the receptor [56]. Thrombin binding facilitates the cleavage of PAR<sub>1</sub>, which thus does not require high



concentrations of thrombin for its activation (0.1 to 1u/ml). In contrast, PAR<sub>4</sub> has no hirudin-like site and higher concentrations of thrombin are requested to activate PAR<sub>4</sub> (10u/ml) [36]. We deliberately used a concentration of thrombin that would activate both PAR<sub>1</sub> and PAR<sub>4</sub> and observed that thrombin induced a PAR<sub>1</sub>-dependent calcium mobilization in human primary afferents (Figure 4 a,b). We also observed that the amplitude of the thrombin response in human sensory neurons was reduced by concomitant PAR<sub>4</sub> activation. Indeed, PAR<sub>4</sub> blockade enhanced the amplitude of thrombin-induced calcium response (Figure 4c). In keeping with these functional results, our single cell PCR data show that a population of human DRG neurons expressing PAR<sub>1</sub> also expresses PAR<sub>4</sub>. Interestingly, PAR<sub>4</sub> was strongly expressed in human DRG neurons. The magnitude of PAR<sub>4</sub> effect on PAR<sub>1</sub>- or thrombin-induced calcium signals in primary afferents was not as strong as it could have been expected considering the strong PAR<sub>4</sub> expression. In primary afferents, PAR<sub>4</sub> might have other function than counteracting the PAR<sub>1</sub>-induced signals.

In contrast to studies reporting calcium mobilization in rodent sensory neurons after stimulation with PAR<sub>2</sub>-APs, or in other human cell lines [37; 54], we observed that human sensory neurons did not mobilize calcium after exposure to PAR<sub>2</sub>-APs. Interestingly, PAR<sub>2</sub> expression in human sensory DRG neurons was clearly demonstrated both at the mRNA and protein levels (Figures 1 and 2). This result was quite surprising since numerous studies have demonstrated that human cells expressing PAR<sub>2</sub>, mobilize calcium after exposure to PAR<sub>2</sub> tethered ligand synthetic peptide [39]. The lack of calcium mobilization in human sensory neurons exposed to PAR<sub>2</sub> synthetic agonist does not mean that the receptor is silent or non-functional. Indeed, other signaling pathways that have been previously described for PAR<sub>2</sub>, such as pERK or cAMP signaling might be implicated in human sensory neurons. Such pathways would have to be investigated in future studies. Furthermore, in human submucosal neurons, PAR<sub>2</sub>-AP induces very weak calcium mobilization [34], while PAR<sub>2</sub> has been shown to be functional and potently activated by the Trypsin-3 protease in the same neurons [43]. This suggests that depending on the type of agonists that are tested (synthetic peptides or proteases), the cell response might be diverse and more or less potent. Similar findings have been reported for PAR pharmacology in a number of cells and tissues [39].



One crucial step in studying the relevance of PARs as therapeutic targets is the definition of protease profile and activities associated with pathological states. Indeed, the opposite roles for PAR<sub>1</sub> and PAR<sub>4</sub> we have defined here in human primary afferent signaling, suggest that depending on the proteases present and their concentration, pro- or anti-nociceptive signals could be prominent. Thrombin can activate both PAR<sub>1</sub> and PAR<sub>4</sub>, but as mentioned above, at different concentrations. Therefore, the concentration of active thrombin detected in colonic tissues from IBS patients could give an important indication on the activation status of PARs and the overall nociceptive signals. Cathepsin G activates PAR<sub>4</sub> [44], and disarms PAR<sub>1</sub> [41]. Here again, the presence of cathepsin G in tissues could modify nociceptive status and accordingly participate to pain relief. Thus, it is clear that protease profiling will be an important step towards our comprehension of nociceptive signaling to primary afferents. However, the results presented here clearly defined that PAR<sub>1</sub> activation on sensory neurons is involved in sensory response to mediators associated with IBS in humans, even though the proteases responsible for PAR<sub>1</sub> activation are not yet defined. Whether PAR<sub>1</sub> activation can potentiate transient receptor potential channels as it was demonstrated in rodent sensory neurons both for PAR<sub>1</sub> [48] and PAR<sub>2</sub> [2; 3; 15; 16; 18; 24] would still have to be confirmed in human primary afferents.

One major limitation of the present study though is that even if we have harvested thoracic DRGs containing neurons projecting from the colon, not all neurons present in the culture dishes are colonic projections. We cannot define whether colonic projections are responding to supernatants of IBS patients, or even whether colonic projections are indeed expressing the different PARs. In animal studies, retrograde labeling of projecting neurons is used to identify the origin of the neurons, but this is hardly applicable to human studies.

In conclusion, this study describes in human sensory neurons, the expression of PARs and their ability to generate calcium signaling. The results highlight the functionality of PAR<sub>1</sub> as an activator of calcium-dependent signaling and PAR<sub>4</sub> as an inhibitor of such signaling. Most importantly, mediators from IBS patient tissues signal to human primary afferent in a PAR<sub>1</sub>-dependent mechanism, illustrating the potential of PAR<sub>1</sub> antagonism as a new therapeutic option to treat symptoms associated with IBS.

## Authors participation

CD, TB, SG-C, SMB, and CR performed experiments and statistical analysis. MRG and GB provided technical and material supports. CD, NC, NV have drafted the manuscript. NC and NV designed and supervised the study and obtained funding.

## Grant support

This work was supported by the Agence Nationale de la Recherche (R12177BB to NV), the Region Midi-Pyrénées (to NV), by the European Research Council (ERC-2012-StG-20111109) (to NV). This work was funded in part by the National Health and Medical Research Council (NHMRC) of Australia Project Grant #1083480 (awarded to S.M.B). S.M.B is an NHMRC R.D Wright Biomedical Research Fellow.

## Acknowledgments

The authors acknowledge the microscope core facility (INSERM UMR1043, Toulouse), the National Diseases Research Interchange (NDRI) and Anabios Corporation (San Diego, USA) for collecting Human DRG. The authors declare no conflict of interest.

## References

- [1] Ahn HS, Foster C, Boykow G, Stamford A, Manna M, Graziano M. Inhibition of cellular action of thrombin by N3-cyclopropyl-7-[[4-(1-methylethyl)phenyl]methyl]-7H-pyrrolo[3, 2-f]quinazoline-1,3-diamine (SCH 79797), a nonpeptide thrombin receptor antagonist. *Biochem Pharmacol* 2000;60(10):1425-1434.
- [2] Amadesi S, Cottrell GS, Divino L, Chapman K, Grady EF, Bautista F, Karanjia R, Barajas-Lopez C, Vanner S, Vergnolle N, Bunnett NW. Protease-activated receptor 2 sensitizes TRPV1 by protein kinase Cepsilon- and A-dependent mechanisms in rats and mice. *J Physiol* 2006;575(Pt 2):555-571.
- [3] Amadesi S, Nie J, Vergnolle N, Cottrell GS, Grady EF, Trevisani M, Manni C, Geppetti P, McRoberts JA, Ennes H, Davis JB, Mayer EA, Bunnett NW. Protease-activated receptor 2 sensitizes the capsaicin receptor transient receptor potential vanilloid receptor 1 to induce hyperalgesia. *J Neurosci* 2004;24(18):4300-4312.
- [4] Asfaha S, Brussee V, Chapman K, Zochodne DW, Vergnolle N. Proteinase-activated receptor-1 agonists attenuate nociception in response to noxious stimuli. *Br J Pharmacol* 2002;135(5):1101-1106.
- [5] Asfaha S, Cenac N, Houle S, Altier C, Papez MD, Nguyen C, Steinhoff M, Chapman K, Zamponi GW, Vergnolle N. Protease-activated receptor-4: a novel mechanism of inflammatory pain modulation. *Br J Pharmacol* 2007;150(2):176-185.

- 458 [6] Auge C, Balz-Hara D, Steinhoff M, Vergnolle N, Cenac N. Protease-Activated Receptor-4 (PAR4): a  
459 role as inhibitor of visceral pain and hypersensitivity. *Neurogastroenterol& Motil* 2009;21(11):1189-  
460 e1107.
- 461 [7] Barbara G, Feinle-Bisset C, Ghoshal UC, Quigley EM, Santos J, Vanner S, Vergnolle N, Zoetendal  
462 EG. The Intestinal Microenvironment and Functional Gastrointestinal Disorders. *Gastroenterology*  
463 2016.
- 464 [8] Barbara G, Stanghellini V, De GR, Cremon C, Cottrell GS, Santini D, Pasquinelli G, Morselli-Labate  
465 AM, Grady EF, Bunnett NW, Collins SM, Corinaldesi R. Activated mast cells in proximity to colonic  
466 nerves correlate with abdominal pain in irritable bowel syndrome. *Gastroenterology*  
467 2004;126(3):693-702.
- 468 [9] Barbara G, Wang B, Stanghellini V, de Giorgio R, Cremon C, Di Nardo G, Trevisani M, Campi B,  
469 Geppetti P, Tonini M, Bunnett NW, Grundy D, Corinaldesi R. Mast cell-dependent excitation of  
470 visceral-nociceptive sensory neurons in irritable bowel syndrome. *Gastroenterology* 2007;132(1):26-  
471 37.
- 472 [10] Brierley S. Altered Ion Channel/Receptor Expression and Function in Extrinsic Sensory Neurons:  
473 The Cause of and Solution to Chronic Visceral Pain? *Adv Exp Med Biol* 2016;891:75-90.
- 474 [11] Brierley SM, Linden DR. Neuroplasticity and dysfunction after gastrointestinal inflammation. *Nat*  
475 *Rev Gastroenterol Hepatol* 2014;11(10):611-627.
- 476 [12] Buhner S, Braak B, Li Q, Kugler EM, Klooker T, Wouters M, Donovan J, Vignali S, Mazzuoli-Weber  
477 G, Grundy D, Boeckxstaens G, Schemann M. Neuronal activation by mucosal biopsy supernatants  
478 from irritable bowel syndrome patients is linked to visceral sensitivity. *Exp Physiol* 2014;99(10):1299-  
479 1311.
- 480 [13] Buhner S, Li Q, Vignali S, Barbara G, De Giorgio R, Stanghellini V, Cremon C, Zeller F, Langer R,  
481 Daniel H, Michel K, Schemann M. Activation of human enteric neurons by supernatants of colonic  
482 biopsy specimens from patients with irritable bowel syndrome. *Gastroenterology* 2009;137(4):1425-  
483 1434.
- 484 [14] Canavan C, West J, Card T. The epidemiology of irritable bowel syndrome. *Clin Epidemiol*  
485 2014;6:71-80.
- 486 [15] Cenac N, Altier C, Chapman K, Liedtke W, Zamponi G, Vergnolle N. Transient receptor potential  
487 vanilloid-4 has a major role in visceral hypersensitivity symptoms. *Gastroenterology* 2008;135(3):937-  
488 946, 946 e931-932.
- 489 [16] Cenac N, Altier C, Motta JP, d'Aldebert E, Galeano S, Zamponi GW, Vergnolle N. Potentiation of  
490 TRPV4 signalling by histamine and serotonin: an important mechanism for visceral hypersensitivity.  
491 *Gut* 2010;59(4):481-488.
- 492 [17] Cenac N, Andrews CN, Holzhausen M, Chapman K, Cottrell G, Andrade-Gordon P, Steinhoff M,  
493 Barbara G, Beck P, Bunnett NW, Sharkey KA, Ferraz JG, Shaffer E, Vergnolle N. Role for protease  
494 activity in visceral pain in irritable bowel syndrome. *J Clin Invest* 2007;117(3):636-647.
- 495 [18] Cenac N, Bautzova T, Le Faouder P, Veldhuis NA, Poole DP, Rolland C, Bertrand J, Liedtke W,  
496 Dubourdeau M, Bertrand-Michel J, Zecchi L, Stanghellini V, Bunnett NW, Barbara G, Vergnolle N.  
497 Quantification and Potential Functions of Endogenous Agonists of Transient Receptor Potential  
498 Channels in Patients With Irritable Bowel Syndrome. *Gastroenterology* 2015;149(2):433-444 e437.
- 499 [19] Coelho AM, Vergnolle N, Guiard B, Fioramonti J, Bueno L. Proteinases and proteinase-activated  
500 receptor 2: a possible role to promote visceral hyperalgesia in rats. *Gastroenterology* 2002;122  
501 1035-1047.
- 502 [20] D'Aldebert E, Cenac N, Rousset P, Martin L, Rolland C, Chapman K, Selves J, Alric L, Vinel JP,  
503 Vergnolle N. Transient receptor potential vanilloid 4 activated inflammatory signals by intestinal  
504 epithelial cells and colitis in mice. *Gastroenterology* 2011;140(1):275-285.
- 505 [21] de Garavilla L, Vergnolle N, Young SH, Ennes H, Steinhoff M, Ossovskaya VS, D'Andrea MR,  
506 Mayer EA, Wallace JL, Hollenberg MD, Andrade-Gordon P, Bunnett NW. Agonists of proteinase-  
507 activated receptor 1 induce plasma extravasation by a neurogenic mechanism. *Br J Pharmacol*  
508 2001;133(7):975-987.

- [22] Di Serio C, Pellerito S, Duarte M, Massi D, Naldini A, Cirino G, Prudovsky I, Santucci M, Geppetti P, Marchionni N, Masotti G, Tarantini F. Protease-activated receptor 1-selective antagonist SCH79797 inhibits cell proliferation and induces apoptosis by a protease-activated receptor 1-independent mechanism. *Basic Clin Pharmacol Toxicol* 2007;101(1):63-69.
- [23] Enck P, Aziz Q, Barbara G, Farmer AD, Fukudo S, Mayer EA, Niesler B, Quigley EM, Rajilic-Stojanovic M, Schemann M, Schwille-Kiuntke J, Simren M, Zipfel S, Spiller RC. Irritable bowel syndrome. *Nat Rev Dis Primers* 2016;2:16014.
- [24] Grant AD, Cottrell GS, Amadesi S, Trevisani M, Nicoletti P, Materazzi S, Altier C, Cenac N, Zamponi GW, Bautista-Cruz F, Lopez CB, Joseph EK, Levine JD, Liedtke W, Vanner S, Vergnolle N, Geppetti P, Bunnett NW. Protease-activated receptor 2 sensitizes the transient receptor potential vanilloid 4 ion channel to cause mechanical hyperalgesia in mice. *J Physiol* 2007;578(Pt 3):715-733.
- [25] Hollenberg MD, Mihara K, Polley D, Suen JY, Han A, Fairlie DP, Ramachandran R. Biased signalling and proteinase-activated receptors (PARs): targeting inflammatory disease. *Br J Pharmacol* 2014;171(5):1180-1194.
- [26] Hollenberg MD, Saifeddine M, Al-Ani B, Gui Y. Proteinase-activated receptor 4 (PAR4): action of PAR4-activating peptides in vascular and gastric tissue and lack of cross-reactivity with PAR1 and PAR2. *Can J Physiol Pharmacol* 1999;77(6):458-464.
- [27] Hollenberg MD, Saifeddine M, Sandhu S, Houle S, Vergnolle N. Proteinase-activated receptor-4: evaluation of tethered ligand-derived peptides as probes for receptor function and as inflammatory agonists in vivo. *Br J Pharmacol* 2004;143(4):443-454.
- [28] Karanjia R, Spreadbury I, Bautista-Cruz F, Tsang ME, Vanner S. Activation of protease-activated receptor-4 inhibits the intrinsic excitability of colonic dorsal root ganglia neurons. *Neurogastroenterol Motil* 2009;21(11):1218-1221.
- [29] Kugler EM, Mazzuoli G, Demir IE, Ceyhan GO, Zeller F, Schemann M. Activity of protease-activated receptors in primary cultured human myenteric neurons. *Front Neurosci* 2012;6:133.
- [30] Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130(5):1480-1491.
- [31] Lovell RM, Ford AC. Global prevalence of and risk factors for irritable bowel syndrome: a meta-analysis. *Clin Gastroenterol Hepatol* 2012;10(7):712-721 e714.
- [32] Martin L, Auge C, Boue J, Buresi MC, Chapman K, Asfaha S, Andrade-Gordon P, Steinhoff M, Cenac N, Dietrich G, Vergnolle N. Thrombin receptor: An endogenous inhibitor of inflammatory pain, activating opioid pathways. *Pain* 2009;146(1-2):121-129.
- [33] Mihara K, Ramachandran R, Saifeddine M, Hansen KK, Renaux B, Polley D, Gibson S, Vanderboor C, Hollenberg MD. Thrombin-Mediated Direct Activation of Proteinase-Activated Receptor-2: Another Target for Thrombin Signaling. *Mol Pharmacol* 2016;89(5):606-614.
- [34] Mueller K, Michel K, Krueger D, Demir IE, Ceyhan GO, Zeller F, Kreis ME, Schemann M. Activity of protease-activated receptors in the human submucous plexus. *Gastroenterology* 2011;141(6):2088-2097 e2081.
- [35] Nakanishi-Matsui M, Zheng YW, Sulciner DJ, Weiss EJ, Ludeman MJ, Coughlin SR. PAR3 is a cofactor for PAR4 activation by thrombin. *Nature* 2000;404(6778):609-613.
- [36] Nieman MT, Schmaier AH. Interaction of thrombin with PAR1 and PAR4 at the thrombin cleavage site. *Biochemistry* 2007;46(29):8603-8610.
- [37] Oikonomopoulou K, Hansen KK, Saifeddine M, Vergnolle N, Tea I, Diamandis EP, Hollenberg MD. Proteinase-mediated cell signalling: targeting proteinase-activated receptors (PARs) by kallikreins and more. *Biol Chem* 2006;387(6):677-685.
- [38] Ostertag D, Annahazi A, Krueger D, Michel K, Demir IE, Ceyhan GO, Zeller F, Schemann M. Trypsin potentiates enteric nerve activation by histamine and serotonin: Relevance for the effects of mucosal biopsy supernatants from irritable bowel syndrome patients. *Neurogastroenterol Motil* 2017;29(9).
- [39] Ramachandran R, Altier C, Oikonomopoulou K, Hollenberg MD. Proteinases, Their Extracellular Targets, and Inflammatory Signaling. *Pharmacol Rev* 2016;68(4):1110-1142.

- [40] Ramachandran R, Hollenberg MD. Proteinases and signalling: pathophysiological and therapeutic implications via PARs and more. *Br J Pharmacol* 2008;153 Suppl 1:S263-282.
- [41] Renesto P, Si-Tahar M, Moniatte M, Balloy V, Van Dorsselaer A, Pidard D, Chignard M. Specific inhibition of thrombin-induced cell activation by the neutrophil proteinases elastase, cathepsin G, proteinase 3: evidence for distinct cleavage sites within the aminoterminal domain of the thrombin receptor. *Blood* 1997;89:1944-1953.
- [42] Riewald M, Ruf W. Science review: role of coagulation protease cascades in sepsis. *Crit Care* 2003;7(2):123-129.
- [43] Rolland-Fourcade C, Denadai-Souza A, Cirillo C, Lopez C, Jaramillo JO, Desormeaux C, Cenac N, Motta JP, Larauche M, Tache Y, Berghe PV, Neunlist M, Coron E, Kirzin S, Portier G, Bonnet D, Alric L, Vanner S, Deraison C, Vergnolle N. Epithelial expression and function of trypsin-3 in irritable bowel syndrome. *Gut* 2017.
- [44] Sambrano GR, Huang W, Faruqi T, Mahrus S, Craik C, Coughlin SR. Cathepsin G activates protease-activated receptor-4 in human platelets. *JBiolChem* 2000;275(10):6819-6823.
- [45] Steinhoff M, Neisius U, Ikoma T. Proteinase-activated receptor-2 mediates itch: a novel pathway for pruritus in human skin *J Neurosci* 2003;In press
- [46] Steinhoff M, Vergnolle N, Young SH, Tognetto M, Amadesi S, Ennes HS, Trevisani M, Hollenberg MD, Wallace JL, Caughey GH, Mitchell SE, Williams LM, Geppetti P, Mayer EA, Bunnett NW. Agonists of proteinase-activated receptor 2 induce inflammation by a neurogenic mechanism. *Nat Med* 2000;6:151-158.
- [47] Valdez-Morales EE, Overington J, Guerrero-Alba R, Ochoa-Cortes F, Ibeakanma CO, Spreadbury I, Bunnett NW, Beyak M, Vanner SJ. Sensitization of peripheral sensory nerves by mediators from colonic biopsies of diarrhea-predominant irritable bowel syndrome patients: a role for PAR2. *Am J Gastroenterol* 2013;108(10):1634-1643.
- [48] Vellani V, Kinsey AM, Prandini M, Hechtfisher SC, Reeh P, Magherini PC, Giacomoni C, McNaughton PA. Protease activated receptors 1 and 4 sensitize TRPV1 in nociceptive neurones. *Mol Pain* 2010;6:61.
- [49] Vergnolle N. Modulation of visceral pain and inflammation by protease-activated receptors. *BrJ Pharmacol* 2004;141(8):1264-1274.
- [50] Vergnolle N. Postinflammatory visceral sensitivity and pain mechanisms. *Neurogastroenterol Motil* 2008;20 Suppl 1:73-80.
- [51] Vergnolle N. Protease-activated receptors as drug targets in inflammation and pain. *PharmacolTher* 2009;123(3):292-309.
- [52] Vergnolle N. Protease inhibition as new therapeutic strategy for GI diseases. *Gut* 2016;65(7):1215-1224.
- [53] Vergnolle N, Bunnett NW, Sharkey KA, Brussee V, Compton SJ, Grady EF, Cirino G, Gerard N, Basbaum AI, Andrade-Gordon P, Hollenberg MD, Wallace JL. Proteinase-activated receptor-2 and hyperalgesia: A novel pain pathway. *Nat Med* 2001;7(7):821-826.
- [54] Vergnolle N, Macnaughton WK, Al-Ani B, Saifeddine M, Wallace JL, Hollenberg MD. Proteinase-activated receptor 2 (PAR2)-activating peptides: identification of a receptor distinct from PAR2 that regulates intestinal transport. *Proc Natl Acad Sci U S A* 1998;95(13):7766-7771.
- [55] Vergnolle N, Wallace JL, Bunnett NW, Hollenberg MD. Protease-activated receptors in inflammation, neuronal signaling and pain. *Trends PharmacolSci* 2001;22(3):146-152.
- [56] Vu TK, Hung DT, Wheaton VI, Coughlin SR. Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 1991;64:1057-1068.
- [57] Young SE, Duvernay MT, Schulte ML, Nance KD, Melancon BJ, Engers J, Wood MR, Hamm HE, Lindsley CW. A Novel and Selective PAR4 Antagonist: ML354. *Probe Reports from the NIH Molecular Libraries Program*. Bethesda (MD), 2010.



## Figure legends

**Figure 1: Expression of PARs in whole human DRG.** (a) Relative expression of PAR<sub>1</sub> (white bar), PAR<sub>2</sub> (gray bar) and PAR<sub>4</sub> (black bar) mRNA in human DRG normalized with GAPDH expression; n=5 human DRG from 5 different donors. (b) Representative pictures of PAR<sub>1</sub> (blue), PAR<sub>2</sub> (red), PAR<sub>4</sub> (green) and PGP9.5 (white, arrows = neurons, arrow heads = nerves fibers) immunodetection on slices of human DRG; scale bar = 50  $\mu$ m. (c) Percentage of PAR expression on PGP9.5 positive cells in slices of T11 and T12 DRG from 3 donors. For PAR<sub>1</sub> expression 154 neurons were counted, 162 for PAR<sub>2</sub> and 89 for PAR<sub>4</sub>. (d) Percentage of PAR-immunoreactive neurons according to the diameter of neuronal perikarya. (c), (d), n = 3 x (T11 + T12). For each condition, an average of the percentage obtained on 3 pictures was performed.

## Figure 2: Expression of PARs in cultured human DRG neurons.

(a) Expression of PAR<sub>1</sub> (in white), PAR<sub>2</sub> (in gray) and PAR<sub>4</sub> (in black) mRNA transcripts by single-cell RT-qPCR (left panel) of human neurons (TUBB3 positive cells). Pie charts representation of the expression (dark color) or not (light color) of PAR<sub>1</sub> (in blue), PAR<sub>2</sub> (in red) and PAR<sub>4</sub> (in green) mRNA in human neurons (right panel). Each segment represents a single neuron. n = 26 neurons. (b) Representative pictures of PARs (in red) and PGP9.5 (in green) immunodetection in cultures of human DRG; scale bar = 50  $\mu$ m. (c) Percentage of PAR expression on PGP9.5-positive cells in human DRG neuron cultures from 5 different donors. Two wells for each condition were counted: 112 neurons for PAR<sub>1</sub> labelling, 134 for PAR<sub>2</sub> and 88 for PAR<sub>4</sub> were counted.

## Figure 3: Effects of PAR-AP in human sensory neurons.

(a) Representative trace of calcium flux experiment obtained in one well of human sensory neuron culture exposed to PAR<sub>1</sub>-AP, PAR<sub>2</sub>-AP or PAR<sub>4</sub>-AP (100  $\mu$ M each). Percentage of responding neurons (b) and amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; c) in human sensory neurons exposed to PAR agonist peptides (PAR-AP, 100  $\mu$ M, black bar) or inactive peptides (PAR-IP, 100  $\mu$ M, white bar). n=6 to 8 independent experiments of 3 wells per condition and 20-53 neurons per well. (d) Calcium flux amplitude of responding neurons exposed to PAR<sub>1</sub>-IP (100  $\mu$ M, white bar) or to increasing doses of PAR<sub>1</sub>-AP (1, 10, 50 and 100  $\mu$ M, black bar) pretreated or not with a PAR<sub>1</sub> antagonist (SCH79797, 10  $\mu$ M). n=4 to 5 independent experiments of 3 wells per condition and 36-62 neurons per well. (e)

Representative trace of calcium flux experiment obtained in one well of human sensory neurons culture exposed to PAR<sub>1</sub>-AP (100  $\mu$ M) and pre-incubated with PAR<sub>2</sub>-AP or PAR<sub>4</sub>-AP (100  $\mu$ M each). Percentage of responding neurons (**f**) and amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; **g**) in human sensory neurons exposed to PAR<sub>1</sub>-AP (100  $\mu$ M, all bars) and pretreated with PAR<sub>2</sub>-IP or PAR<sub>4</sub>-IP (100  $\mu$ M, white bar), PAR<sub>2</sub>-AP (100  $\mu$ M, gray bar) or PAR<sub>4</sub>-AP (100 $\mu$ M, black bar). n=3 to 4 independent experiments of 3 wells per condition and 30-58 neurons per well. Statistical analysis was performed using Kruskal-Wallis analysis of variance and subsequent Dunn's post hoc test. \* p<0.05, \*\* p<0.01, \*\*\* p<0.001, significantly different from the corresponding inactive-peptide groups; fff p < 0.001, significantly different from PAR<sub>1</sub>-AP (100 $\mu$ M).

**Figure 4: Effects of thrombin in human DRG neurons.**

Percentage of responding neurons (**a**) and amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; **b**) in human DRG neurons exposed to thrombin (10 U/mL, black bar) or its vehicle (HBSS, white bar). n=3 to 4 independent experiments of 3 wells per condition and 31-45 neurons per well. Statistical analysis was performed using Mann-Whitney test. \* p<0.05, \*\* p<0.01, significantly different from HBSS group. Percentage of responding neurons (**c**) and amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; **d**) in human sensory neurons exposed to thrombin (10 U/mL, all bars) and pretreated with PAR<sub>1</sub> antagonist (SCH79797, 10  $\mu$ M, gray bar), PAR<sub>4</sub> antagonist (ML354, 10  $\mu$ M, black bar) or their vehicle (HBSS, white bar). n=3 independent experiments of 3 wells per condition and 39-68 neurons per well. Statistical analysis was performed using Kruskal-Wallis analysis of variance and subsequent Dunn's post hoc test. \* p<0.05, \*\*\* p<0.001, significantly different from the corresponding inactive-peptide groups.


**Figure 5: Effects of supernatants from colonic biopsies of IBS patients or healthy controls on calcium mobilization in human DRG neurons.**

(**a**) Representative trace of calcium flux experiment obtained in one well of human sensory neuron culture exposed to supernatant of diarrhea-predominant IBS patient (hexagon) or to healthy control (HC, circle). Amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; **b**) in human sensory neurons and percentage of responding neurons (**c**) exposed to supernatants from colonic biopsies of IBS patients: constipation-predominant (-C, triangle), diarrhea-predominant (-D, hexagon), alternate (-A, square) or to supernatants from colonic biopsies of healthy control (HC, circle). (**d**) Representative trace of calcium flux experiment obtained in


676 one well of human sensory neurons culture exposed to supernatant of alternate-predominant  
677 IBS patient (square) and pretreated with PAR<sub>1</sub> antagonist (SCH79797, 10  $\mu$ M, gray square) or  
678 its vehicle (white square). Amplitude of intracellular calcium mobilization ( $\Delta F/F$ ; **e**) in human  
679 sensory neurons and percentage of responding neurons (**f**) exposed to supernatants from  
680 colonic biopsies of IBS patients: constipation-predominant (-C, triangle), diarrhea-  
681 predominant (-D, hexagon) or alternate (-A, square) and pretreated with PAR<sub>1</sub> antagonist  
682 (SCH79797, 10  $\mu$ M), or its vehicle. Data are represented as scattered dot plot with line at  
683 mean. Each symbol represents one patient. n=6 independent experiments of 3 wells per  
684 condition and 42-53 neurons per well. Statistical analysis was performed using Mann-  
685 Whitney test (**b** and **c**) or Wilcoxon matched pairs test (**e** and **f**). \* p<0.05, \*\* p<0.01,  
686 significantly different from HC (**b** and **c**) or from IBS group (**e** and **f**).




**Figure 1**




**Figure 2**



**Figure 3**



**Figure 4**



**Figure 5**

