

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The persistent homotopy type distance

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Frosini, P., Landi, C., Mémoli, F. (2019). The persistent homotopy type distance. *HOMOLOGY, HOMOTOPY AND APPLICATIONS*, 21(2), 231-259 [10.4310/HHA.2019.v21.n2.a13].

Availability:

This version is available at: <https://hdl.handle.net/11585/682656> since: 2021-02-28

Published:

DOI: <http://doi.org/10.4310/HHA.2019.v21.n2.a13>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Patrizio Frosini, Claudia Landi and Facundo Mémoli,
The persistence homotopy type distance,
Homology, Homotopy and Applications, vol. 21 (2019), n. 2, 231-259.
Preprint available at <https://arxiv.org/pdf/1702.07893.pdf>.

The final published version is available online at: <https://dx.doi.org/10.4310/HHA.2019.v21.n2.a13>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

THE PERSISTENT HOMOTOPY TYPE DISTANCE

PATRIZIO FROSINI, CLAUDIA LANDI AND FACUNDO MÉMOLI

Abstract

We introduce the *persistent homotopy type distance* d_{HT} to compare two real valued functions defined on possibly different homotopy equivalent topological spaces. The underlying idea in the definition of d_{HT} is to measure the minimal shift that is necessary to apply to one of the two functions in order that the sublevel sets of the two functions become homotopy equivalent. This distance is interesting in connection with persistent homology. Indeed, our main result states that d_{HT} still provides an upper bound for the bottleneck distance between the persistence diagrams of the intervening functions. Moreover, because homotopy equivalences are weaker than homeomorphisms, this implies a lifting of the standard stability results provided by the L^∞ distance and the natural pseudo-distance d_{NP} . From a different standpoint, we prove that d_{HT} extends the L^∞ distance and d_{NP} in two ways. First, we show that, appropriately restricting the category of objects to which d_{HT} applies, it can be made to coincide with the other two distances. Finally, we show that d_{HT} has an interpretation in terms of interleavings that naturally places it in the family of distances used in persistence theory.

Keywords: bottleneck distance between persistence diagrams, natural pseudo-distance, interleaving distance, stability, merge trees.

AMS Subject Classification: 55P10, 68U05, 18A23.

1. Introduction

Persistent homology has been developed as a theory to study topological properties of noisy or incomplete data, establishing itself as a fundamental tool for topological data analysis [15, 19, 7]. Persistent homology is characterized by an invariant called the persistence diagram (also known as the barcode) which summarizes both topological features of a dataset and their prominence. One of the reasons for the success of persistent homology is that persistence diagrams change continuously provided that

the input dataset also changes continuously. This is known as the Stability Theorem of Persistence [11]. Usually, (a) changes in persistence homology are measured via the bottleneck distance between persistence diagrams, (b) datasets are modeled as real valued functions defined on the same space, and (c) one uses the L^∞ distance between functions to quantify their changes:

Theorem 1 (Stability Theorem of Persistence [11]). *Let X be a compact polyhedron. Then, for all continuous tame functions $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}$, and all integers $k \geq 0$,*

$$d_B(D_k(\varphi_1), D_k(\varphi_2)) \leq \|\varphi_1 - \varphi_2\|_\infty. \quad (1.1)$$

Above, d_B stands for the bottleneck distance between persistence diagrams, and $D_k(\varphi)$ is the persistence diagram corresponding to the k -th homology of the sub-level set filtration of the function φ .

In order to lift the Stability Theorem of Persistence to the case when functions are defined on different, albeit homeomorphic spaces, one can resort to the natural pseudo-distance. If two continuous functions $\varphi_X : X \rightarrow \mathbb{R}$, $\varphi_Y : Y \rightarrow \mathbb{R}$ are given on homeomorphic spaces X and Y , then the natural pseudo-distance d_{NP} [14] between them is defined by

$$d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)) := \inf_h \|\varphi_1 - \varphi_2 \circ h\|_\infty \quad (1.2)$$

where $h : X \rightarrow Y$ varies among the morphisms of the category \mathbf{H}_0 whose objects are all topological spaces endowed with real-valued continuous functions, and morphisms are all homeomorphisms between topological spaces. If two continuous functions $\varphi_X : X \rightarrow \mathbb{R}$, $\varphi_Y : Y \rightarrow \mathbb{R}$ are given on non-homeomorphic spaces X and Y , then we set $d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)) := \infty$. Then, since persistence diagrams of sub-level set filtrations are invariant under reparameterization, one obtains an improvement of inequality (1.1) stated in Theorem 1:

$$d_B(D_k(\varphi_X), D_k(\varphi_Y)) \leq d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)). \quad (1.3)$$

However, the natural pseudo-distance is not suitable when we are interested in analyzing functions defined on non-homeomorphic topological spaces.

In this paper we construct a new extended pseudo-metric, called the *persistent homotopy type distance*, denoted d_{HT} , to quantify the distance between real-valued functions defined on different spaces which is meaningful when the spaces are at least homotopy equivalent. In plain words, the persistent homotopy type distance is a generalization of the natural pseudo-distance in that it uses homotopy equivalences instead of homeomorphisms, and these homotopy equivalences are required to satisfy a certain coherence property in terms of the sublevel sets of the given real valued functions. This allows us to use the persistent homotopy type distance to obtain a new and stronger stability theorem for persistent homology, which is the main contribution of our paper:

Theorem 2. *Let X and Y be compact polyhedra, and k be any non-negative integer. Let $\varphi_X : X \rightarrow \mathbb{R}$ and $\varphi_Y : Y \rightarrow \mathbb{R}$ be continuous functions. Then,*

$$d_B(D_k(\varphi_X), D_k(\varphi_Y)) \leq d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)).$$

We note that in the statement above d_{HT} becomes infinity when the underlying spaces above are not homotopy equivalent.

While lifting the stability theorem is the main motivation for introducing d_{HT} , this paper contains two further contributions as presented below. The first one concerns possible extensions of the basic definition of d_{HT} in Theorem 2 to more general settings in order to enhance its applicability, while the second one concerns the connections of d_{HT} to other well-known distances such as the interleaving distance.

Variations on the basic definition of d_{HT}

To prove Theorem 2, it is sufficient to take a basic definition of d_{HT} where functions φ_X and φ_Y are real-valued, and no restrictions are required on the possible homotopy equivalences apart from being sublevel set preserving up to a shift. This mirrors the early stage research about the natural pseudo-distance d_{NP} as defined in (1.2), where functions are real-valued as well, and homeomorphisms are generically morphisms of \mathbf{H}_0 . On the other hand, more recently the natural-pseudo distance d_{NP} has undergone mainly two variations in order to extend its range of applications. We now illustrate such variations for d_{NP} , with the goal in the following sections to achieve similar possibilities for d_{HT} .

The first variation we can consider in the basic definition of d_{NP} given in (1.2), is to extend the class of considered objects to compare from real-valued functions $\varphi_X : X \rightarrow \mathbb{R}$ to vector-valued functions $\varphi_X = (\varphi_i) : X \rightarrow \mathbb{R}^n$. In this case, we consider $\|\varphi_X\|_\infty := \sup_{x \in X} \max_{1 \leq i \leq n} |\varphi_i(x)|$. This generalization of d_{NP} to vector-valued functions is much valuable since it allows to lift the stability of the interleaving distance of multidimensional persistence modules [22] in much the same way as we lift the stability of the bottleneck distance in one-dimensional persistence.

The second variation we can consider in (1.2), is the restriction of the considered morphisms to use for the comparison from the category \mathbf{H}_0 (whose objects now we can take to be all topological spaces endowed with \mathbb{R}^n -valued continuous functions, and morphisms are all homeomorphisms between topological spaces) to any subcategory \mathbf{H} of \mathbf{H}_0 provided that \mathbf{H} is closed under inverse [2]: for any $\varphi_X : X \rightarrow \mathbb{R}^n$ and $\varphi_Y : Y \rightarrow \mathbb{R}^n$,

$$d_{\text{NP}}^{\mathbf{H}}((X, \varphi_X), (Y, \varphi_Y)) := \inf_{h \in \text{hom}_{\mathbf{H}}(X, Y)} \|\varphi_1 - \varphi_2 \circ h\|_\infty \quad (1.4)$$

if there is a homeomorphism from X to Y in \mathbf{H} , $d_{\text{NP}}^{\mathbf{H}}((X, \varphi_X), (Y, \varphi_Y)) = \infty$ otherwise. This variation is useful in applications of the natural pseudo-distance to cases when the desired invariance is not the one expressed by any homeomorphism [17, 18]. For example, two monotonic functions $\varphi, \psi : [0, 1] \rightarrow \mathbb{R}$ with the same set of extrema are equivalent under d_{NP} (and therefore equivalent for standard persistent homology) when every homeomorphism from $[0, 1]$ to $[0, 1]$ is accepted, whereas restricting the set of acceptable homeomorphisms permits distinguishing them. From a different perspective, since the group of all self-homeomorphisms of a topological space, even a compact one, is not itself compact, the possibility of restricting the set of homeomorphisms is also motivated by the desire of working with compact groups. This would be useful for obtaining interesting theorems e.g. good finite approximations of the considered groups.

Motivated by these earlier generalizations of the basic definition of the natural pseudo-distance, our purpose in the next sections will be to propose a definition for a persistent homotopy type distance that not only lifts the stability result, but also

behaves similarly to d_{NP} with respect to such generalizations.

The homotopy type distance as an interleaving distance

Starting with [9, 21], and more recently with [22, 4, 10], a unifying look at all the metrics usually used to state the stability theorems of persistence has been proposed in terms of interleaving distances. Interleavings apply between pairs of functors from the category of ordered reals to any other category. Interleavings are given by pairs of natural transformations between each one of the functors and a *shifted* version of the other functor.

The interleaving distance measures the smallest shift that allows the existence of an interleaving. Since many distances considered in topological data analysis can be formulated in terms of interleavings, it is natural to ask whether the same holds true for the persistent homotopy type distance. A further contribution of this paper is a positive answer to this question. Recent related work in this direction is that of Blumberg and Lesnick [3] where the authors construct the so called *homotopy interleaving distance* between \mathbb{R} -spaces and study its universality. Whereas the relationship (if any) between our homotopy type distance and that of [3] is currently unclear, in this paper we prove that, when restricted to merge trees, our homotopy type distance agrees with the interleaving distance between merge trees of Morozov et al. [23].

Organization of the paper

After introducing the persistent homotopy type distance in Section 2, we discuss its properties and give some examples. In Section 3, we prove that the bottleneck distance between persistence diagrams is upper-bounded by the persistent homotopy type distance (Theorem 2). In other words, we lift the Stability Theorem of Persistence (Theorem 1) to functions defined on different spaces provided that they are homotopy equivalent. In Section 4, we show that the interleaving distance between merge trees can be obtained as a special case of the persistence homotopy type distance. Then, Section 5 offers an interpretation of the persistent homotopy type distance as an interleaving distance more in general. Finally, a discussion section offers some thoughts on possible extensions.

2. Mathematical setting

For any integer $n \geq 1$, let us endow \mathbb{R}^n with the partial order \preceq defined by setting, for any $\alpha, \beta \in \mathbb{R}^n$, $\alpha = (\alpha_i) \preceq \beta = (\beta_i)$ whenever $\alpha_i \leq \beta_i$ for $i = 1, \dots, k$. When $\alpha \preceq \beta$ we also write $\beta \succeq \alpha$. For $\alpha = (\alpha_i) \in \mathbb{R}^n$, we set $\|\alpha\|_\infty = \max_{1 \leq i \leq k} |\alpha_i|$. Recall that for a function $\varphi_X : X \rightarrow \mathbb{R}^n$, we set $\|\varphi_X\|_\infty = \sup_{x \in X} \|\varphi_X(x)\|_\infty$. For $\alpha \in \mathbb{R}$, we denote by $\vec{\alpha}$ the diagonal element $(\alpha, \alpha, \dots, \alpha) \in \mathbb{R}^n$.

Let us consider the category \mathbf{S} such that: the objects of \mathbf{S} are all the pairs (X, φ_X) where X is a topological space and $\varphi_X : X \rightarrow \mathbb{R}^n$ is a vector-valued continuous function; the morphisms of \mathbf{S} from an object (X, φ_X) to another object (Y, φ_Y) are all the continuous maps $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$. The composition of morphisms is the usual composition of maps and identity morphisms are identity maps.

Definition 2.1. For every $\alpha \in \mathbb{R}^n$, with $\alpha \succeq \mathbf{0}$, the α -shift functor $(\cdot)_\alpha : \mathbf{S} \rightarrow \mathbf{S}$ is defined as follows: for every (X, φ_X) in $ob(\mathbf{S})$, $(X, \varphi_X)_\alpha = (X, \varphi'_X)$, where $\varphi'_X(x) =$

$\varphi_X(x) - \alpha$ for every $x \in X$; for every $f : (X, \varphi_X) \rightarrow (Y, \varphi_Y)$ in $\text{hom}(\mathbf{S})$, $(f)_\alpha = f$ regarded as a morphism between $(X, \varphi_X)_\alpha$ and $(Y, \varphi_Y)_\alpha$.

Instead of \mathbf{S} , we can restrict ourselves to any sub-category \mathbf{C} of \mathbf{S} provided that \mathbf{C} is closed with respect to the α -shift functor for any $\alpha \in \mathbb{R}^n$. From now on we assume one such \mathbf{C} is fixed.

Definition 2.2. Let $(X, \varphi_X), (Y, \varphi_Y)$ be objects in \mathbf{C} . Given $\alpha \succeq \mathbf{0}$ in \mathbb{R}^n , an α -map f with respect to (φ_X, φ_Y) is a morphism in \mathbf{C} from (X, φ_X) to $(Y, \varphi_Y)_\alpha$, that is a continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X + \alpha$. Furthermore, given two α -maps $f_1 : X \rightarrow Y$ and $f_2 : X \rightarrow Y$, an α -homotopy in \mathbf{C} between f_1 and f_2 with respect to the pair (φ_X, φ_Y) is a continuous map $H : X \times [0, 1] \rightarrow Y$ such that

1. $f_1 \equiv H(\cdot, 0)$;
2. $f_2 \equiv H(\cdot, 1)$;
3. $H(\cdot, t)$ is an α -map in \mathbf{C} with respect to the pair (φ_X, φ_Y) for every $t \in [0, 1]$.

If an α -homotopy between the α -maps f_1 and f_2 with respect to the pair (φ_X, φ_Y) exists, we say that f_1, f_2 are α -homotopic with respect to (φ_X, φ_Y) .

In plain words, an α -homotopy between α -maps is a homotopy that is a α -map at every instant.

Remark 2.1. Any map $f : X \rightarrow Y$ that induces a morphism $f : (X, \varphi_X) \rightarrow (Y, \varphi_Y)_\alpha$ with respect to (φ_X, φ_Y) , also induces a morphism $f : (X, \varphi_X) \rightarrow (Y, \varphi_Y)_\beta$, still with respect to (φ_X, φ_Y) , for every $\beta \succeq \alpha$. Therefore, an α -map can also be regarded as a β -map, for every $\beta \succeq \alpha$.

Remark 2.2. For any fixed $\alpha \succeq \mathbf{0}$, the α -homotopy with respect to (φ_X, φ_Y) is an equivalence relation on α -maps from (X, φ_X) to $(Y, \varphi_Y)_\alpha$.

We now introduce the relation of α -homotopy equivalence between objects of \mathbf{C} . In spite of its name, in general it is not going to be an equivalence relation.

Definition 2.3. For every $\alpha \succeq \mathbf{0}$ in \mathbb{R}^n and any two objects (X, φ_X) and (Y, φ_Y) in \mathbf{C} , we say that (X, φ_X) and (Y, φ_Y) are α -homotopy equivalent in \mathbf{C} if there exist α -maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ in \mathbf{C} , with respect to (φ_X, φ_Y) and (φ_Y, φ_X) respectively, such that the following properties hold:

- the map $g \circ f : X \rightarrow X$ is 2α -homotopic to id_X with respect to (φ_X, φ_X) ;
- the map $f \circ g : Y \rightarrow Y$ is 2α -homotopic to id_Y with respect to (φ_Y, φ_Y) .

If the two previous conditions hold, we say that g (resp. f) is an α -homotopy inverse of f (resp. g) with respect to (φ_X, φ_Y) (resp. (φ_Y, φ_X)), and that (f, g) constitutes a pair of α -homotopy equivalences in \mathbf{C} with respect to the pair (φ_X, φ_Y) .

We observe that in general the α -homotopy inverse of f with respect to the pair (φ_X, φ_Y) is not unique.

We are now ready to define the persistent homotopy type distance.

Definition 2.4. For any $\alpha \in \mathbb{R}$, set $\vec{\alpha} = (\alpha, \alpha, \dots, \alpha) \in \mathbb{R}^n$. For any $((X, \varphi_X), (Y, \varphi_Y))$ in $ob(\mathbf{C}) \times ob(\mathbf{C})$, we define the set

$$\Lambda^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) := \{\alpha \geq 0 : (X, \varphi_X) \text{ and } (Y, \varphi_Y) \text{ are } \vec{\alpha}\text{-homotopy equivalent in } \mathbf{C}\},$$

and the value

$$d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) := \inf \Lambda^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)),$$

where we use the convention that the infimum over the empty set is $+\infty$.

$d_{\text{HT}}^{\mathbf{C}}$ will be called the *persistent homotopy type (pseudo-)distance* on the category \mathbf{C} . When \mathbf{C} is taken to be the whole \mathbf{S} , sometimes we simply denote $d_{\text{HT}}^{\mathbf{C}}$ by d_{HT} .

Proposition 2.1. *Let (X, φ_X) and (Y, φ_Y) be two objects in \mathbf{S} such that φ_X and φ_Y are bounded, that is $\|\varphi_X\|_\infty, \|\varphi_Y\|_\infty < +\infty$. Then, $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) < \infty$ if and only if X and Y are homotopy equivalent.*

Proof. Let us assume that X and Y are homotopy equivalent. Then, there exist maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ and homotopies $H : X \times [0, 1] \rightarrow X$ and $G : Y \times [0, 1] \rightarrow Y$ between $g \circ f$ and id_X and between $f \circ g$ and id_Y , respectively. Since φ_X and φ_Y are bounded functions, the numbers $\alpha_X := \|\varphi_X - \varphi_Y \circ f\|_\infty$, $\alpha_Y := \|\varphi_X \circ g - \varphi_Y\|_\infty$, $\hat{\alpha}_X := \frac{1}{2} \max_{t \in [0, 1]} \|\varphi_X - \varphi_X \circ H(\cdot, t)\|_\infty$, and $\hat{\alpha}_Y := \frac{1}{2} \max_{t \in [0, 1]} \|\varphi_Y \circ G(\cdot, t) - \varphi_Y\|_\infty$ are all finite. Thus, $\alpha := \sup\{\alpha_X, \alpha_Y, \hat{\alpha}_X, \hat{\alpha}_Y\} < \infty$, and α belongs to $\Lambda^{\mathbf{S}}((X, \varphi_X), (Y, \varphi_Y))$ so that $d_{\text{HT}} \leq \alpha$. Conversely, if $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) < \infty$, there is some $\alpha \in \mathbb{R}$ such that (X, φ_X) and (Y, φ_Y) are $\vec{\alpha}$ -homotopy equivalent in \mathbf{S} with $\vec{\alpha} = (\alpha, \alpha, \dots, \alpha)$. Hence, X and Y are homotopy equivalent. \square

In this paper we will follow the convention that $\alpha + \infty = \infty + \alpha = \infty$ for every $\alpha \in \overline{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$.

We now prove that $d_{\text{HT}}^{\mathbf{C}}$ is an extended pseudo-metric. To this aim we use the following lemma.

Lemma 2.1. *Let $(X, \varphi_X), (Y, \varphi_Y), (Z, \varphi_Z)$ be three objects in \mathbf{C} . If (f_1, g_1) is a pair of α_1 -homotopy equivalences with respect to (φ_X, φ_Y) in \mathbf{C} and (f_2, g_2) is a pair of α_2 -homotopy equivalences with respect to (φ_Y, φ_Z) also in \mathbf{C} , then $(f_2 \circ f_1, g_1 \circ g_2)$ is a pair of $(\alpha_1 + \alpha_2)$ -homotopy equivalences with respect to the pair (φ_X, φ_Z) in \mathbf{C} .*

Proof. By definition, $f_2 \circ f_1$ (resp. $g_1 \circ g_2$) is an $(\alpha_1 + \alpha_2)$ -map with respect to (φ_X, φ_Z) (resp. (φ_Z, φ_X)). Again by definition, a $2\alpha_1$ -homotopy $H_1 : X \times [0, 1] \rightarrow X$ with respect to (φ_X, φ_X) from $g_1 \circ f_1$ to id_X and a $2\alpha_2$ -homotopy $H_2 : Y \times [0, 1] \rightarrow Y$ with respect to (φ_Y, φ_Y) from $g_2 \circ f_2$ to id_Y exist. Thus, we can define a map

$$\bar{H} : X \times [0, 1] \rightarrow X \text{ by setting } \bar{H}(x, t) := \begin{cases} g_1 \circ H_2(f_1(x), 2t), & \text{if } t \in [0, 1/2) \\ H_1(x, 2t - 1), & \text{if } t \in [1/2, 1] \end{cases}. \bar{H} \text{ is a}$$

$(2\alpha_1 + 2\alpha_2)$ -homotopy with respect to (φ_X, φ_X) from $g_1 \circ g_2 \circ f_2 \circ f_1$ to id_X . Similarly, we can define a $(2\alpha_1 + 2\alpha_2)$ -homotopy (φ_Z, φ_Z) from $f_2 \circ f_1 \circ g_1 \circ g_2$ to id_Z . This proves the claim. \square

Remark 2.3. The term *extended pseudo-metric* means that the function $d_{\text{HT}}^{\mathbf{C}}$ is a function defined on $ob(\mathbf{C}) \times ob(\mathbf{C})$ such that, for every $(X, \varphi_X), (Y, \varphi_Y) \in ob(\mathbf{C})$, it holds that

- (i) $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) \in [0, \infty]$,
- (ii) if $(X, \varphi_X) = (Y, \varphi_Y)$ then $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) = 0$,
- (iii) $d_{\text{HT}}^{\mathbf{C}}$ satisfies the symmetry property,
- (iv) $d_{\text{HT}}^{\mathbf{C}}$ satisfies the triangle inequality.

Proposition 2.2. *The function $d_{\text{HT}}^{\mathbf{C}} : \text{ob}(\mathbf{C}) \times \text{ob}(\mathbf{C}) \rightarrow \overline{\mathbb{R}}$ is an extended pseudo-metric.*

Proof. We check that $d_{\text{HT}}^{\mathbf{C}}$ satisfies the properties of an extended pseudo-metric.

1. By definition, $d_{\text{HT}}^{\mathbf{C}}$ cannot take negative values.
2. $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (X, \varphi_X)) = 0$ because the identity map of X , id_X , belongs to $\text{hom}(\mathbf{C})$ and $(\text{id}_X, \text{id}_X)$ is a pair of $\mathbf{0}$ -homotopy equivalences with respect to (φ_X, φ_X) .
3. The equality $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) = d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (X, \varphi_X))$ immediately follows from the symmetry of the definition of pairs of α -homotopy equivalences.
4. Let $\varphi_X : X \rightarrow \mathbb{R}^n$, $\varphi_Y : Y \rightarrow \mathbb{R}^n$, $\varphi_Z : Z \rightarrow \mathbb{R}^n$ be three objects in our category \mathbf{C} . Let $\vec{\alpha} = (\alpha, \alpha, \dots, \alpha) \in \mathbb{R}^n$ with $\alpha \geq 0$. If either (X, φ_X) is not $\vec{\alpha}$ -homotopy equivalent to (Y, φ_Y) , or (Y, φ_Y) is not $\vec{\alpha}$ -homotopy equivalent to (Z, φ_Z) for any $\alpha \in \mathbb{R}$, then the definition of $d_{\text{HT}}^{\mathbf{C}}$ implies that $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) + d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (Z, \varphi_Z)) = \infty$. In this case the inequality $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) + d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (Z, \varphi_Z)) \geq d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Z, \varphi_Z))$ is trivially satisfied.

Let us assume that (f_1, g_1) is a pair of $\vec{\alpha}_1$ -homotopy equivalences with respect to (φ_X, φ_Y) for some $\alpha_1 \geq 0$, and (f_2, g_2) is a pair of $\vec{\alpha}_2$ -homotopy equivalences with respect to (φ_Y, φ_Z) for some $\alpha_2 \geq 0$. By definition of $d_{\text{HT}}^{\mathbf{C}}$, we can assume that $\alpha_1 \leq d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) + \varepsilon$ and $\alpha_2 \leq d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (Z, \varphi_Z)) + \varepsilon$ for an arbitrarily small $\varepsilon > 0$. We know from Lemma 2.1 that $(f_2 \circ f_1, g_1 \circ g_2)$ is a pair of $(\vec{\alpha}_1 + \vec{\alpha}_2)$ -homotopy equivalences with respect to (φ_X, φ_Z) . It follows that

$$\begin{aligned} d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Z, \varphi_Z)) &\leq \alpha_1 + \alpha_2 \\ &\leq d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) + d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (Z, \varphi_Z)) + 2\varepsilon. \end{aligned}$$

By taking the limit for ε tending to 0, we obtain the triangle inequality

$$d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Z, \varphi_Z)) \leq d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) + d_{\text{HT}}^{\mathbf{C}}((Y, \varphi_Y), (Z, \varphi_Z)).$$

□

As we mentioned in the introduction, our definition of the persistent homotopy type distance is meant to be a generalization of the natural pseudo-distance. The next proposition gives a relationship between these two distances when we compare functions on two homeomorphic spaces, whereas Example 2.9 proves that we can read the natural pseudo-distance d_{NP} into the persistent homotopy type distance $d_{\text{HT}}^{\mathbf{C}}$ when we suitably restrict the underlying category.

Proposition 2.3. *Let (X, φ_X) and (Y, φ_Y) be objects in \mathbf{S} where X and Y are homeomorphic. Then,*

$$d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) \leq d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)).$$

In particular, if $X = Y$, and $(X, \varphi_1), (X, \varphi_2)$ are objects in \mathbf{S} , then

$$d_{\text{HT}}((X, \varphi_1), (X, \varphi_2)) \leq \|\varphi_1 - \varphi_2\|_{\infty}.$$

Proof. If $d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)) = \infty$ then there is nothing to prove. If not, then there exist $\alpha \geq 0$ and a homeomorphism $f : X \rightarrow Y$ such that $\|\varphi_X - \varphi_Y \circ f\|_\infty \leq \alpha$. Thus, f and f^{-1} are $\vec{\alpha}$ -maps, and $f^{-1} \circ f$ and $f \circ f^{-1}$ are $\mathbf{0}$ -homotopic, and hence $2\vec{\alpha}$ -homotopic, with $\vec{\alpha} = (\alpha, \alpha, \dots, \alpha)$ to id_X and id_Y , respectively, by constant homotopies. Therefore, (f, f^{-1}) is a pair of $\vec{\alpha}$ -homotopy equivalences for (φ_X, φ_Y) , implying that (X, φ_X) and (Y, φ_Y) are $\vec{\alpha}$ -homotopy equivalent. Thus, $d_{\text{HT}} \leq d_{\text{NP}}$. \square

In the next few sections we will compare the persistent homotopy type distance with other metrics widely used in the topological data analysis literature to measure the perturbations in the input functions. In particular, in Section 5 we will show that the persistent homotopy type distance can be represented as an interleaving type distance. For the moment, however, we proceed with the study of the persistent homotopy type distance using the current definition.

2.1. Examples

We now show how to compute the persistent homotopy type distance in some simple cases. We take $\mathbf{C} = \mathbf{S}$ and denote $d_{\text{HT}}^{\mathbf{S}}$ simply by d_{HT} . For the sake of simplicity, we take $n = 1$.

The first example pertains to the case when one can retract a given space X to a subset A in such a way that the function values do not increase.

Proposition 2.4. *Let (X, φ_X) be an object in \mathbf{S} and let A be a subspace of X such that there exists a deformation retract $F : X \times [0, 1] \rightarrow X$ of X onto A with the property that $\varphi_X(F(x, t)) \leq \varphi_X(x)$ for all $x \in X$ and $t \in [0, 1]$. Then,*

$$d_{\text{HT}}((X, \varphi_X), (A, \varphi_X|_A)) = 0.$$

Proof. The proof follows directly from the definition of the homotopy type distance. \square

As an immediate corollary, we obtain:

Example 2.1. Let X be a contractible space and let $z \in X$. For any fixed $c \in \mathbb{R}$, let $\varphi_c : X \rightarrow \mathbb{R}$ denote the constant function $\varphi_c(x) = c$ on X . Simply denote again by c the constant function equal to c on $\{z\}$. It holds that $d_{\text{HT}}((X, \varphi_c), (\{z\}, c)) = 0$.

In the next two examples, we show that d_{HT} may be different from d_{NP} even when the spaces are homeomorphic.

Example 2.2. Let X be the band obtained by gluing without any twist two opposite sides of a rectangle R , and Y the band obtained by gluing the same sides of R after applying a complete twist (i.e. a torsion of 2π radians). Assume that the glued sides have length equal to 2 and that X and Y are embedded into \mathbb{R}^3 as follows (see Figure 1):

$$X : \begin{cases} x(u, v) = 2 \cos u \\ y(u, v) = 2 \sin u \\ z(u, v) = v \end{cases} \quad Y : \begin{cases} x(u, v) = (2 + v \sin u) \cos u \\ y(u, v) = (2 + v \sin u) \sin u \\ z(u, v) = v \cos u \end{cases}$$

with $0 \leq u < 2\pi$ and $-1 \leq v \leq 1$. Moreover, define $\varphi_X : X \rightarrow \mathbb{R}$ by $\varphi_X(x, y, z) = z$, and similarly $\varphi_Y : Y \rightarrow \mathbb{R}$ by $\varphi_Y(x, y, z) = z$. This way the centerlines of both X




Figure 1: The twisted band Y used in Example 2.2. Colors are as follows: the red region corresponds to points with non-negative z -coordinate whereas the green region is the complement, that is, it corresponds to those points with negative z -coordinate.

and Y coincide with the curve $C = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 4, z = 0\}$ and φ_X takes values in $\{-1, 1\}$ at every boundary point of X , while φ_Y continuously varies in $[-1, 1]$ for every boundary point of Y . We claim that

$$d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) = 1, \quad (2.1)$$

and

$$d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)) = 2. \quad (2.2)$$

In order to establish (2.2) note that for any homeomorphism $f : X \rightarrow Y$ one has that $\|\varphi_X - \varphi_X \circ f\|_\infty = 2$ simply because f must take boundary points to boundary points. In order to establish (2.1), defining the retractions onto C given by $r_X : X \rightarrow C$ by $r_X(x, y, z) = (x, y, 0)$ and $r_Y : Y \rightarrow C$ by $r_Y((2 + v \sin u) \cos u, (2 + v \sin u) \sin u, v \cos u) = (2 \cos u, 2 \sin u, 0)$, and the inclusions $i_{C, X}$ and $i_{C, Y}$ of C into X and Y respectively, one immediately checks that $(i_{C, Y} \circ r_X, i_{C, X} \circ r_Y)$ is a pair of 1-homotopy equivalences with respect to (φ_X, φ_Y) . Thus, $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y))$ is at most 1. On the other hand, we claim that by Theorem 2 for $k = 1$ we have that $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) \geq 1$. Indeed, notice that $D_1(\varphi_X) = \{(-1, \infty)\}$ whereas $D_1(\varphi_Y) = \{(0, \infty)\}$, see Figure 1. It then follows that the bottleneck distance (see Section 3.1) between these diagrams satisfies $d_B(D_1(\varphi_X), D_1(\varphi_Y)) = 1$, which via Theorem 2 implies our claim.

Example 2.3 (Lens spaces). Let $X = Y$ be the disjoint union of the lens spaces $L(7, 1)$ and $L(7, 2)$. Define $\varphi_X : X \rightarrow \mathbb{R}$ by setting $\varphi_X|_{L(7, 1)} \equiv 0$ and $\varphi_X|_{L(7, 2)} \equiv 1$, and define $\varphi_Y : Y \rightarrow \mathbb{R}$ by setting $\varphi_Y|_{L(7, 1)} \equiv 1$ and $\varphi_Y|_{L(7, 2)} \equiv 0$. Then, $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y))$ is zero because $L(7, 1)$ and $L(7, 2)$ are homotopy equivalent but not homeomorphic [25], whereas

$$d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)) = 1.$$

Finally, we consider another example which shows that d_{NP} and d_{HT} can in fact be the same.

Example 2.4. Let M be any closed connected oriented manifold. Take $X = Y = M$ and $\varphi_Y = \varphi_c$, the constant function equal to c . In this case $d_{\text{NP}}(\varphi_X, \varphi_c) = \|\varphi_X - c\|_\infty$. We claim that in this case $d_{\text{HT}}(\varphi_X, \varphi_c) = d_{\text{NP}}(\varphi_X, \varphi_c)$. Assume that $\alpha \geq 0$ is such that there exists a pair (f, g) a pair of α -homotopy equivalences with respect to the pair (φ_X, φ_c) . We will prove that $\alpha \geq \|\varphi_X - c\|_\infty$. Note that from $c = \varphi_c(f(x)) \leq \alpha + \varphi_X(x)$ for $x \in M$ we have that $\alpha \geq c - \min \varphi_X$. Now, since $g \circ f$ and id_M are homotopic, their degrees are the same, therefore $|\deg(f)| = |\deg(g)| = 1$. Hence, both f and g are surjective. From the condition $\varphi_X(g(x)) \leq \alpha + \varphi_c(x) = \alpha + c$ for all $x \in M$ we obtain that $\alpha \geq \max \varphi_X - c$. Hence, $\alpha \geq \max(c - \min \varphi_X, \max \varphi_X - c) = \|\varphi_X - c\|_\infty$. The fact that $d_{\text{HT}} \leq d_{\text{NP}}$ (Proposition 2.3) yields the claim.

2.2. Comments on Definition 2.4

We now consider three questions that may naturally arise:

- What if we simplify the definition of α -homotopies given in Definition 2.2 by removing the condition about H being an α -map at each instant, requiring only it to be an α -map for $t = 0$ and $t = 1$? And, analogously, what if we remove the condition about an α -homotopy in a subcategory \mathbf{C} of \mathbf{S} to be a morphism in \mathbf{C} at each instant?
- Would it be possible to define $d_{\text{HT}}^{\mathbf{C}}$ via a minimum instead of an infimum?
- Is $d_{\text{HT}}^{\mathbf{C}}$ actually only an extended pseudo-metric or rather an extended metric?

We will answer this questions by means of examples, taking $n = 1$, i.e. real valued functions, for the sake of simplicity. Moreover, when possible, we take $\mathbf{C} = \mathbf{S}$, and in such cases we simply write d_{HT} instead of $d_{\text{HT}}^{\mathbf{S}}$.

As for the first issue, the definition of α -homotopy we have given may seem more complex than necessary. One could think of removing the condition about being an α -map at each instant, maintaining only the condition that it be an α -map for $t = 0$ and $t = 1$. Unfortunately, the new metric d^* that we would obtain from this simplified definition of α -homotopies would not give an upper bound for the bottleneck distance in persistent homology. In particular, the vanishing of d^* would not imply that the considered sublevel-set persistent homologies are the same. This is shown in the following example, proving that the analogue of Theorem 2 for d^* would not hold.




Figure 2: The functions considered in Example 2.5. In this case $d^*(\varphi, \psi) = 0$ but $d_{\text{HT}}([-1, 1], \varphi), ([-1, 1], \psi) > 0$.

Example 2.5. Define $\varphi, \psi : [-1, 1] \rightarrow \mathbb{R}$ by $\varphi(x) = 1 - |x|$ and $\psi(x) = (1 + x)/2$ (see Figure 2). We also define $f, g : [-1, 1] \rightarrow [-1, 1]$ by setting $f(x) = 1 - 2|x|$ and $g(x) = (x - 1)/2$. We have that $\psi \circ f(x) = \frac{1+(1-2|x|)}{2} = 1 - |x| = \varphi(x)$ and $\varphi \circ g(x) = 1 - \left|\frac{x-1}{2}\right| = 1 - \frac{1-x}{2} = \frac{1+x}{2} = \psi(x)$, so that the maps f and g are 0-maps with respect to (φ, ψ) and (ψ, φ) , respectively. Furthermore, $g \circ f(x) = \frac{(1-2|x|)-1}{2} = -|x|$ and $f \circ g(x) = 1 - 2\left|\frac{x-1}{2}\right| = 1 - 2\left(\frac{1-x}{2}\right) = x$. It follows that $f \circ g$ equals the identity, whereas $g \circ f$ is homotopic to the identity via the homotopy $H(x, t) := (t - 1)|x| + tx$. Note that $H(\cdot, 0)$ and $H(\cdot, 1)$ are 0-maps with respect to (φ, φ) . As a consequence, $d^*(\varphi, \psi) = 0$. Now we can observe that the sublevelset persistent homologies of φ and ψ are clearly different from each other, and hence the corresponding bottleneck distance is positive. This can be easily seen by checking that 1 is a homological critical value for φ , but not for ψ . Therefore, d^* is not an upper bound for the bottleneck distance. On the contrary, we shall prove in Section 3.1 that that property holds for d_{HT} (Theorem 2). This fact leads us to prefer the definition of α -homotopy, and consequently of d_{HT} , that we have presented.

Analogously, the following Example 2.6 shows the difference between asking an α -homotopy to be a morphism in \mathbf{C} rather than in \mathbf{S} at each instant.

Example 2.6. Denoting by $2\mathbb{Z}$ the even integers and by $2\mathbb{Z} + 1$ the odd integers, let $X = \{0\} \times \mathbb{R} \cup [0, 1] \times 2\mathbb{Z}$ and $Y = \{0\} \times \mathbb{R} \cup [0, 1] \times (2\mathbb{Z} + 1)$ be two subsets of \mathbb{R}^2 . Let $\varphi_X(s, t) = t$ and $\varphi_Y(s, t) = t$. Taking \mathbf{C} the subcategory of \mathbf{S} whose morphisms are homeomorphisms, it holds that $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) = 1$ whereas it holds that $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) = 0$. Notice that removing the request for an α -homotopy in \mathbf{C} to be a homeomorphism at each instant, we would obtain $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) = 0$ by using the 0-maps $f : X \rightarrow Y$ $f(s, t) = (s, t - 1)$ and $g : Y \rightarrow X$, $g(s, t) = (s, t - 1)$ shown in Figure 3.




Figure 3: The objects and maps considered in Example 2.6.

As for the second issue, the use of an infimum instead of a minimum is necessary, as the following example shows. The same example shows that d_{HT} is not an extended metric, but only an extended pseudo-metric, thus clarifying the third issue.

Example 2.7. Define $\varphi, \psi : [-2, 2] \rightarrow \mathbb{R}$ by setting $\varphi(x) = -|x|$, $\psi(x) = 0$ for $|x| \leq 1$ and $\psi(x) = 2 - 2|x|$ for $|x| > 1$ (see Figure 4). We claim that, for every small enough




Figure 4: The functions considered in Example 2.7. In this case the infimum of the values $\alpha > 0$ such that a pair of α -homotopy equivalences with respect to (φ, ψ) exists is 0. However, no pair of 0-homotopy equivalences with respect to (φ, ψ) exists.

$\alpha > 0$ in \mathbb{R} , we can find a pair of α -homotopy equivalences $(f_\alpha, f_\alpha^{-1})$ with respect to (φ, ψ) . As a consequence, $d_{\text{HT}}([\![-2, 2]\!], \varphi), ([\![-2, 2]\!], \psi) = 0$. In order to show our claim, let us take $0 < \alpha < 2$ and consider the homeomorphism $f_\alpha : [-2, 2] \rightarrow [-2, 2]$ defined by setting

$$f_\alpha(x) := \begin{cases} \frac{1}{2-\alpha}x + \frac{2\alpha-2}{2-\alpha}, & \text{if } -2 \leq x < -\alpha \\ \frac{1}{\alpha}x, & \text{if } -\alpha \leq x < \alpha \\ \frac{1}{2-\alpha}x - \frac{2\alpha-2}{2-\alpha}, & \text{if } \alpha \leq x \leq 2. \end{cases}$$

Observe that $f_\alpha([\![-2, -\alpha]\!]) = [\![-2, -1]\!]$, $f_\alpha([\![-\alpha, \alpha]\!]) = [\![-1, 1]\!]$ and $f_\alpha([\![\alpha, 2]\!]) = [\![1, 2]\!]$. It follows that

$$\psi \circ f_\alpha(x) := \begin{cases} 2 - 2 \left| \frac{x}{2-\alpha} + \frac{2\alpha-2}{2-\alpha} \right| = 2 + 2 \left(\frac{x}{2-\alpha} + \frac{2\alpha-2}{2-\alpha} \right) = 2 \frac{x+\alpha}{2-\alpha}, & \text{if } -2 \leq x < -\alpha \\ 0, & \text{if } -\alpha \leq x < \alpha \\ 2 - 2 \left| \frac{x}{2-\alpha} - \frac{2\alpha-2}{2-\alpha} \right| = 2 - 2 \left(\frac{x}{2-\alpha} - \frac{2\alpha-2}{2-\alpha} \right) = -2 \frac{x-\alpha}{2-\alpha}, & \text{if } \alpha \leq x \leq 2. \end{cases}$$

As a consequence

$$\varphi(x) - \psi \circ f_\alpha(x) := \begin{cases} -|x| - 2 \frac{x+\alpha}{2-\alpha} = x - 2 \frac{x+\alpha}{2-\alpha} = -\alpha \frac{2+x}{2-\alpha}, & \text{if } -2 \leq x < -\alpha \\ -|x|, & \text{if } -\alpha \leq x < \alpha \\ -|x| + 2 \frac{x-\alpha}{2-\alpha} = -x + 2 \frac{x-\alpha}{2-\alpha} = -\alpha \frac{2-x}{2-\alpha}, & \text{if } \alpha \leq x \leq 2. \end{cases}$$

We can easily check that $-\alpha \leq \varphi(x) - \psi \circ f_\alpha(x) \leq 0$ for every $x \in [-2, 2]$. It follows that $\|\varphi - \psi \circ f_\alpha\|_\infty \leq \alpha$. Hence, $(f_\alpha, f_\alpha^{-1})$ is a pair of α -homotopy equivalences with respect to (φ, ψ) . Given that α can be chosen arbitrarily close to 0, this implies that

$$d_{\text{HT}}([\![-2, 2]\!], \varphi), ([\![-2, 2]\!], \psi) = d_{\text{NP}}([\![-2, 2]\!], \varphi), ([\![-2, 2]\!], \psi) = 0.$$

We claim that no pair of 0-homotopy equivalences with respect to (φ, ψ) exists. We prove this by contradiction. Assume that a pair of 0-homotopy equivalences (f_0, g_0) with respect to (φ, ψ) exists. Then a homotopy $H_0 : [-2, 2] \times [0, 1] \rightarrow [-2, 2]$ exists, such that $H_0(x, 0) = g_0 \circ f_0(x)$, $H_0(x, 1) = x$ and $\varphi(H_0(x, t)) \leq \varphi(x)$ for every $x \in [-2, 2]$ and every $t \in [0, 1]$. It is easy to prove that $H_0(-2, t) = -2$ for every $t \in [0, 1]$:

$H_0(-2, 1) = -2$, $\varphi(H_0(-2, t)) \leq \varphi(-2)$ for every $t \in [0, 1]$, and -2 is a strict local minimum point for φ . Analogously, $H_0(2, t) = 2$ for every $t \in [0, 1]$. It follows that $g_0 \circ f_0(-2) = -2$ and $g_0 \circ f_0(2) = 2$.

Now, we observe that $f_0(\{-2, 2\}) \subseteq \{-2, 2\}$, because f_0 is a 0-map and $-2, 2$ are the only points where ψ takes a value that is not strictly greater than $\varphi(-2) = \varphi(2) = -2$. Analogously, $g_0(\{-2, 2\}) \subseteq \{-2, 2\}$. By possibly composing f_0 with the reflection $x \mapsto -x$, we can assume that $f_0(-2) = -2$. From the equality $g_0 \circ f_0(-2) = -2$, it follows that $g(-2) = -2$. We can now prove that $f_0([-2, 2]) \subseteq [-2, -1]$. Indeed, since $f_0(-2) = -2$, if $f_0([-2, 2])$ contained a point $\bar{x} > -1$, it should also contain an infinite number of points x where ψ takes the value 0. This contradicts the assumption that f_0 is a 0-map, because φ takes its maximum 0 only at the point 0. Furthermore, $g_0([-2, -1]) \subseteq [-2, 0]$. Indeed, since $g_0(-2) = -2$, if $g_0([-2, -1])$ contained a point $\bar{x} > 0$, there should exist a point $\bar{x} \in [-2, -1]$ such that $g(\bar{x}) = 0$, so that $\varphi(\bar{x}) = 0$. This contradicts the assumption that g_0 is a 0-map. In conclusion, we should have that $g_0 \circ f_0([-2, 2]) \subseteq [-2, 0]$, thus contradicting the fact that $g_0 \circ f_0(2) = 2$.

2.3. The importance of choosing a subcategory \mathbf{C} of \mathbf{S}

The main motivation for considering a subcategory \mathbf{C} of \mathbf{S} instead of only the category \mathbf{S} is to generalize the natural pseudo-distance, whose definition depends on the selection of a set of objects and a set of morphisms that may be respectively smaller than the set of all real-valued continuous functions and the set of all homeomorphisms (cf., e.g., Section 7.1 in [2], [6] and [18]). The following examples show that this choice is fruitful.

The first two examples show that the use of an appropriate subcategory \mathbf{C} of \mathbf{S} allows us to represent the L^∞ distance and the natural pseudo-distance d_{NP} as particular cases of $d_{\text{HT}}^{\mathbf{C}}$.

Example 2.8. For a fixed compact X , consider the category \mathbf{C} whose objects are given by the pairs (X, φ) where $\varphi : X \rightarrow \mathbb{R}^n$ is continuous, and such that between any two objects $(X, \varphi), (X, \varphi') \in \text{ob}(\mathbf{C})$ there is at most one morphism, $\text{id}_X : X \rightarrow X$, from (X, φ) to (X, φ') , and this happens provided that $\varphi' \preceq \varphi$. If φ' is not everywhere less than φ , then no morphism exists from (X, φ) to (X, φ') . By choosing this subcategory \mathbf{C} of \mathbf{S} we obtain that $d_{\text{HT}}^{\mathbf{C}}((X, \varphi), (X, \varphi')) = \|\varphi - \varphi'\|_\infty$.

Example 2.9. Let us set $n = 2m$ in the definition of the category \mathbf{S} . Take the category \mathbf{C} whose objects are the objects (X, φ_X) of \mathbf{S} and the morphisms from an object (X, φ_X) to another object (Y, φ_Y) are the homeomorphisms $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$. If $\psi_X : X \rightarrow \mathbb{R}^m$ is a continuous function, then

$$d_{\text{NP}}((X, \psi_X), (Y, \psi_Y)) = d_{\text{HT}}^{\mathbf{C}}((X, (\psi_X, -\psi_X)), (Y, (\psi_Y, -\psi_Y))).$$

Here the symbol $(\psi_X, -\psi_X)$ denotes the function $\varphi_X : X \rightarrow \mathbb{R}^n$ whose first m components define the function ψ_X , while the last m components define the function $-\psi_X$. Analogously for the symbol $(\psi_Y, -\psi_Y)$.

Let us prove the previous equality. In the case that X and Y are not homeomorphic, we have that $d_{\text{NP}}((X, \psi_X), (Y, \psi_Y)) = d_{\text{HT}}^{\mathbf{C}}((X, (\psi_X, -\psi_X)), (Y, (\psi_Y, -\psi_Y))) = \infty$. Therefore we can confine ourselves to assuming that X and Y are homeomorphic.

If there exist an $\alpha \geq 0$ and a homeomorphism $f : X \rightarrow Y$ such that $\|\psi_X - \psi_Y \circ f\|_\infty \leq \alpha$, then $\psi_Y \circ f \preceq \psi_X + \bar{\alpha}$ and $\psi_Y \circ f \succeq \psi_X - \bar{\alpha}$ (i.e. $-\psi_Y \circ f \preceq -\psi_X + \bar{\alpha}$),

and hence f is an $\vec{\alpha}$ -map in \mathbf{C} from $(X, (\psi_X, -\psi_X))$ to $(Y, (\psi_Y, -\psi_Y))$, with $\vec{\alpha} = (\alpha, \alpha, \dots, \alpha) \in \mathbb{R}^m$. Analogously, since $\|\psi_Y - \psi_X \circ f^{-1}\|_\infty = \|\psi_X - \psi_Y \circ f\|_\infty \leq \alpha$, f^{-1} is an $\vec{\alpha}$ -map in \mathbf{C} from $(Y, (\psi_Y, -\psi_Y))$ to $(X, (\psi_X, -\psi_X))$. Furthermore, $f^{-1} \circ f$ and $f \circ f^{-1}$ are $\mathbf{0}$ -homotopic, and hence $2\vec{\alpha}$ -homotopic to id_X and id_Y , respectively, by constant homotopies. Therefore, (f, f^{-1}) is a pair of $\vec{\alpha}$ -homotopy equivalences for $((\psi_X, -\psi_X), (\psi_Y, -\psi_Y))$, implying that $(X, (\psi_X, -\psi_X))$ and $(Y, (\psi_Y, -\psi_Y))$ are $(\vec{\alpha}, \vec{\alpha})$ -homotopy equivalent. Thus,

$$d_{\text{NP}}((X, \psi_X), (Y, \psi_Y)) \geq d_{\text{HT}}^{\mathbf{C}}((X, (\psi_X, -\psi_X)), (Y, (\psi_Y, -\psi_Y))).$$

If $(X, (\psi_X, -\psi_X))$ and $(Y, (\psi_Y, -\psi_Y))$ are $(\vec{\alpha}, \vec{\alpha})$ -homotopy equivalent with $\alpha \geq 0$, then there is a pair (f, g) of $(\vec{\alpha}, \vec{\alpha})$ -homotopy equivalences for $((\psi_X, -\psi_X), (\psi_Y, -\psi_Y))$. The definition of \mathbf{C} implies that f and g are two homeomorphisms. Moreover, $\psi_Y \circ f \preceq \psi_X + \vec{\alpha}$ and $-\psi_Y \circ f \preceq -\psi_X + \vec{\alpha}$ (i.e. $\psi_Y \circ f \succeq \psi_X - \vec{\alpha}$), and hence $\|\psi_X - \psi_Y \circ f\|_\infty \leq \alpha$.

Thus, $d_{\text{NP}}((X, \psi_X), (Y, \psi_Y)) \leq d_{\text{HT}}^{\mathbf{C}}((X, (\psi_X, -\psi_X)), (Y, (\psi_Y, -\psi_Y)))$.

Therefore, $d_{\text{NP}}((X, \psi_X), (Y, \psi_Y)) = d_{\text{HT}}^{\mathbf{C}}((X, (\psi_X, -\psi_X)), (Y, (\psi_Y, -\psi_Y)))$.

In other words, by taking a suitable subcategory \mathbf{C} of \mathbf{S} we can read the natural pseudo-distance d_{NP} into the persistent homotopy type distance $d_{\text{HT}}^{\mathbf{C}}$.

The following examples provide more insights into the different outcomes that can be obtained varying the category \mathbf{C} . In particular, Example 2.10 shows the effect of a restriction of both $\text{ob}(\mathbf{S})$ and $\text{hom}(\mathbf{S})$, while Example 2.11 illustrates the effect of a restriction of $\text{hom}(\mathbf{S})$.

Example 2.10. Let us imagine to be interested in comparing gray-level colorings of narrow strips (possibly segments), represented by pairs (R_ϵ, φ) where R_ϵ is the rectangle $[-1, 1] \times [0, \epsilon]$ for some $\epsilon \in [0, 1]$ and $\varphi : R_\epsilon \rightarrow \mathbb{R}$ takes values that depend only on the first coordinate (i.e. $\varphi(x, y_1) = \varphi(x, y_2)$ for every $(x, y_1), (x, y_2) \in R_\epsilon$). We will call each of these pairs a *strip coloring*. Let us also assume that we wish (i) to distinguish the generic strip coloring (R_ϵ, φ) from its horizontal reflection $(R_\epsilon, \hat{\varphi})$ with $\hat{\varphi}$ defined by setting $\hat{\varphi}(x, y) = \varphi(-x, y)$, and (ii) not distinguish the strip colorings (R_ϵ, φ) and $(R_0, \varphi|_{R_0})$ on the ground that the height of the strips is not important. Then, these requirements can be satisfied by considering the subcategory \mathbf{C} of \mathbf{S} whose objects are given by the previously defined strip colorings for all $\epsilon \in [0, 1]$, and whose morphisms between two strip colorings $(R_\epsilon, \varphi), (R_{\epsilon'}, \varphi') \in \text{ob}(\mathbf{C})$ are the continuous maps $f = (f_1, f_2) : R_\epsilon \rightarrow R_{\epsilon'}$ such that

1. f_1 depends only on the first coordinate;
2. $f_1(\cdot, 0)$ is a strictly increasing homeomorphism;
3. The inequality $\varphi' \circ f(x, y) \leq \varphi(x, y)$ holds for every $(x, y) \in R_\epsilon$.

Now, let us set $\varphi(x, y) = x$. If $f = (f_1, f_2) : R_\epsilon \rightarrow R_{\epsilon'}$ is an α -map with respect to $(\varphi, \hat{\varphi})$, then $\hat{\varphi}(f(x, y)) \leq \varphi(x, y) + \alpha$ for every $(x, y) \in R_\epsilon$. The definition of φ and $\hat{\varphi}$ implies that $-f_1(x, y) \leq x + \alpha$ for every $(x, y) \in R_\epsilon$. By setting $x = -1$, $y = 0$ and observing that $f_1(-1, 0) = -1$, we get $1 = -f_1(-1, 0) \leq -1 + \alpha$, i.e. $\alpha \geq 2$. It is immediate to check that $(\text{id}_{R_\epsilon}, \text{id}_{R_{\epsilon'}})$ is a pair of 2-homotopy equivalences with respect to $(\varphi, \hat{\varphi})$. Therefore $d_{\text{HT}}^{\mathbf{C}}((R_\epsilon, \varphi), (R_\epsilon, \hat{\varphi})) = 2$. Moreover, $d_{\text{HT}}^{\mathbf{C}}((R_\epsilon, \varphi), (R_0, \varphi|_{R_0})) = 0$, so that $d_{\text{HT}}^{\mathbf{C}}$ satisfies the two required properties (i) and (ii). It is also interesting to

observe that $d_{\text{NP}}((R_\epsilon, \varphi), (R_0, \varphi|_{R_0})) = \infty$ (because R_ϵ and R_0 are not homeomorphic) and $d_{\text{HT}}^{\mathbf{S}}((R_\epsilon, \varphi), (R_\epsilon, \hat{\varphi})) = 0$ (because the homeomorphism $f(x, y) = (-x, y)$ is a 0-homotopy equivalence with respect to the pair $(\varphi, \hat{\varphi})$), so that neither d_{NP} nor $d_{\text{HT}}^{\mathbf{S}}$ satisfy the two required properties (i) and (ii).

Example 2.11. Let \mathbf{C} be the category whose objects are pairs (X, φ) with $X = [-1, 1]$ and $\varphi : X \rightarrow \mathbb{R}$ a continuous function, and whose morphisms between two objects $(X, \varphi), (X, \varphi')$ are the non-decreasing continuous maps $f : X \rightarrow X$. Let us take the two functions $\varphi, \hat{\varphi} : X \rightarrow \mathbb{R}$ defined by setting $\varphi(x) = x$ and $\hat{\varphi}(x) = -x$. In the category \mathbf{C} , the map $f(x) \equiv 1$ is a 0-map with respect to the pair $(\varphi, \hat{\varphi})$, and the map $g(x) \equiv -1$ is a 0-map with respect to the pair $(\hat{\varphi}, \varphi)$. The function $H : [-1, 1] \times [0, 1] \rightarrow \mathbb{R}$, $H(x, t) = t(x + 1) - 1$ is a 0-homotopy between $g \circ f$ and id_X with respect to the pair (φ, φ) , and the function $H' : [-1, 1] \times [0, 1] \rightarrow \mathbb{R}$, $H'(x, t) = t(x - 1) + 1$ is a 0-homotopy between $f \circ g$ and id_X with respect to the pair $(\hat{\varphi}, \hat{\varphi})$. As a consequence, $d_{\text{HT}}^{\mathbf{C}}((X, \varphi), (X, \hat{\varphi})) = 0$. In other words, $d_{\text{HT}}^{\mathbf{C}}$ cannot distinguish φ from $\hat{\varphi}$. However, if we maintain the same objects and restrict the set of morphisms to the set of all increasing homeomorphisms from X to X , we obtain another subcategory \mathbf{C}' of \mathbf{S} such that $d_{\text{HT}}^{\mathbf{C}'}((X, \varphi), (X, \hat{\varphi})) > 0$. Therefore, different choices of the subcategory \mathbf{C} of \mathbf{S} can produce different pseudo-metrics in our model.

2.4. d_{HT} is the same in the topological, PL, and smooth categories

In this section we prove that d_{HT} is the same in the topological, PL, and smooth categories.

Proposition 2.5. *Let \mathbf{C} be the subcategory of \mathbf{S} such that: the objects of \mathbf{C} are all the pairs (X, φ_X) where X is a compact polyhedron, and $\varphi_X : X \rightarrow \mathbb{R}$ is a piecewise linear function; the morphisms of \mathbf{C} from an object (X, φ_X) to another object (Y, φ_Y) are all the piecewise linear maps $f : X \rightarrow Y$ such that $\varphi_Y \circ f \leq \varphi_X$. If (X, φ_X) and (Y, φ_Y) are two objects in \mathbf{C} , and so in \mathbf{S} , then $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) = d_{\text{HT}}^{\mathbf{S}}((X, \varphi_X), (Y, \varphi_Y))$.*

Proof. The inequality $d_{\text{HT}}^{\mathbf{C}}((X, \varphi_X), (Y, \varphi_Y)) \geq d_{\text{HT}}^{\mathbf{S}}((X, \varphi_X), (Y, \varphi_Y))$ holds because each morphism in \mathbf{C} is also a morphism in \mathbf{S} . To see that the converse inequality holds, let ω_X and ω_Y be the moduli of continuity of φ_X and φ_Y , respectively. Let (f, g) be a pair of α -homotopy equivalences in \mathbf{S} with respect to (φ_X, φ_Y) . Let K, L be simplicial complexes such that $X = |K|$ and $Y = |L|$. By the simplicial approximation theorem (cf., e.g., [16]), there exists $\varepsilon > 0$ and K^ε and L^ε subdivisions of K and L with $\text{mesh}(K^\varepsilon), \text{mesh}(L^\varepsilon) < \varepsilon$, respectively, and a PL map $f^\varepsilon : X \rightarrow Y$ such that, for any $x \in X$, each simplex of L^ε containing $f(x)$ contains also $f^\varepsilon(x)$. Since f^ε is homotopic to f via $F(x, t) = (1 - t)f^\varepsilon(x) + tf(x)$, and $\text{mesh}(L^\varepsilon) < \varepsilon$, it holds that f^ε is $\omega_Y(\varepsilon)$ -homotopic to f . Analogously, there exists a simplicial approximation g^ε of g that is $\omega_X(\varepsilon)$ -homotopic to g via $G(x, t) = (1 - t)g^\varepsilon(x) + tg(x)$. Notice that f^ε is an $(\alpha + \omega_Y(\varepsilon))$ -map with respect to (φ_X, φ_Y) and that g^ε is an $(\alpha + \omega_X(\varepsilon))$ -map with respect to (φ_Y, φ_X) . Thus, $S : X \times I \rightarrow X$ defined by $S(x, t) = G(F(x, t), t)$ is a $(\omega_X(\varepsilon) + \omega_Y(\varepsilon))$ -homotopy between $g^\varepsilon \circ f^\varepsilon$ and $g \circ f$. Because (f, g) is a pair of α -homotopy equivalences with respect to (φ_X, φ_Y) , $g \circ f$ is 2α -homotopic to id_X with respect to (φ_X, φ_X) . Thus, there is a $(\omega_X(\varepsilon) + \omega_Y(\varepsilon) + 2\alpha)$ -homotopy between $g^\varepsilon \circ f^\varepsilon$ and id_X with respect to (φ_X, φ_X) . Because any homotopy between continuous mappings can

likewise be approximated by a combinatorial version, possibly further subdividing K and L , we can approximate the $(\omega_X(\varepsilon) + \omega_Y(\varepsilon) + 2\alpha)$ -homotopy between $g^\varepsilon \circ f^\varepsilon$ and id_X by a $(2\omega_X(\varepsilon) + \omega_Y(\varepsilon) + 2\alpha)$ -homotopy between $g^\varepsilon \circ f^\varepsilon$ and id_X that is PL at each instant. Analogously, there is a $(\omega_X(\varepsilon) + 2\omega_Y(\varepsilon) + 2\alpha)$ -homotopy between $f^\varepsilon \circ g^\varepsilon$ and id_Y with respect to (φ_Y, φ_Y) that is PL at each instant. Hence, $(f^\varepsilon, g^\varepsilon)$ is a pair of $(\omega_X(\varepsilon) + \omega_Y(\varepsilon) + \alpha)$ -homotopy equivalences with respect to (φ_X, φ_Y) . As ε tends to 0, $\omega_X(\varepsilon)$ and $\omega_Y(\varepsilon)$ tend to 0. Hence, the claim. \square

Proposition 2.6. *Let \mathbf{D} be the subcategory of \mathbf{S} such that: the objects of \mathbf{D} are all the pairs (M, φ_M) where M is a smooth connected compact manifold, and $\varphi_M : M \rightarrow \mathbb{R}$ is a smooth function; the morphisms of \mathbf{D} from an object (M, φ_M) to another object (N, φ_N) are all the smooth maps $f : M \rightarrow N$ such that $\varphi_N \circ f \leq \varphi_M$. If (M, φ_M) and (N, φ_N) are two objects in \mathbf{D} , and hence in \mathbf{S} , then $d_{\text{HT}}^{\mathbf{D}}((M, \varphi_M), (N, \varphi_N)) = d_{\text{HT}}^{\mathbf{S}}((M, \varphi_M), (N, \varphi_N))$.*

Proof. It may be proved in much the same way as Proposition 2.5, using the fact that any continuous map $f : M \rightarrow N$ with M and N manifolds can be approximated by C^∞ -maps homotopic to f (cf. [20, Ch. 5, Lemma 1.5]). \square

3. Stability of persistent homology with respect to d_{HT}

In this section we establish some connections between the distance d_{HT} and persistent homology, in particular we lift the Stability Theorem of Persistence 1 via d_{HT} .

3.1. Preliminaries

3.1.1. Overview of persistence diagrams and the bottleneck distance

In persistent homology, for each $k = 0, 1, 2, \dots$ one seeks to summarize the topological information contained in the sequence of sublevel sets $\varphi_X^{-1}((-\infty, t])$ into a multiset $D_k(\varphi_X)$ of points of the extended plane called a *persistence diagram*: a birth–death pair (b, d) corresponding to a homological feature in degree k gives rise to a point $p = (b, d)$ in $D_k(\varphi_X)$; points are taken with multiplicity in order to take into account the presence of multiple features appearing and disappearing at the same sublevels.

There is a natural notion of distance, called the *bottleneck distance* d_{B} , that makes the set of all persistence diagrams into a metric space. The bottleneck distance between two persistence diagrams D_1, D_2 is

$$d_{\text{B}}(D_1, D_2) = \inf_M \max \left\{ \sup_{(p,q) \in M} \|p - q\|_\infty, \sup_{\substack{s \in D_1 \amalg D_2 \\ s \notin M(D_1) \amalg M^{-1}(D_2)}} \left| \frac{s_x - s_y}{2} \right| \right\},$$

where M varies among all the binary relations between D_1 and D_2 that are both right- and left-unique, i.e. partial matchings between D_1 and D_2 .

The following result, which for fixed X expresses the continuity of the assignment $\varphi \mapsto D_k(\varphi)$, is standard:

Theorem 1 (Stability Theorem of Persistence [11]). *Let X be a compact polyhedron. Then, for all continuous tame functions $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}$, and all integers $k \geq 0$,*

$$d_B(D_k(\varphi_1), D_k(\varphi_2)) \leq \|\varphi_1 - \varphi_2\|_\infty. \quad (1.1)$$

In the above statement, tameness refers to a certain regularity condition singling out functions φ for which the homology groups $H_k(\varphi^{-1}((-\infty, a]))$ are finite dimensional for all $a \in \mathbb{R}$, and in addition, the maps induced at homology level by the inclusions $\varphi^{-1}((-\infty, a - \varepsilon]) \hookrightarrow \varphi^{-1}((-\infty, a + \varepsilon])$ fail to be isomorphisms for $\varepsilon > 0$ small only at finitely many points.

One of the salient features of the above result is that it assumes the underlying space X to be fixed. Using our construction of the homotopy type distance we lift this result into a statement that applies to any pair (X, φ_X) and (Y, φ_Y) in the category \mathbf{C} satisfying minimal tameness conditions.

3.1.2. Persistence modules and interleavings

More recently, persistent homology has been revisited in terms of persistence modules and interleavings. The main references here are [9] and [10].

A persistent module (over \mathbb{R}) is by definition a directed sequence of vector spaces connected by linear maps $\{V_\delta \xrightarrow{v_{\delta, \delta'}} V_{\delta'}\}_{\delta \leq \delta'}$ such that $v_{\delta, \delta} = \text{id}$ for all $\delta \in \mathbb{R}$ and $v_{\delta', \delta''} \circ v_{\delta, \delta'} = v_{\delta, \delta''}$ for all $\delta \leq \delta' \leq \delta''$. It is said to be q -tame if the linear maps $v_{\delta, \delta'}$ with $\delta < \delta'$ have finite rank.

We now recall the notion of interleaving of persistence modules [9, §3.2]. Given two persistent modules $\{V_\delta \xrightarrow{v_{\delta, \delta'}} V_{\delta'}\}_{\delta \leq \delta'}$ and $\{W_\delta \xrightarrow{w_{\delta, \delta'}} W_{\delta'}\}_{\delta \leq \delta'}$, one says that they are $\alpha \geq 0$ interleaved if for each $\delta \geq 0$ there are maps $\phi_\delta : V_\delta \rightarrow W_{\delta+\alpha}$ and $\gamma_\delta : W_\delta \rightarrow V_{\delta+\alpha}$ such that the following four diagrams (3.1), (3.2), (3.3), and (3.4) commute for all $\delta, \delta' \in \mathbb{R}$ with $\delta \leq \delta'$:

$$\begin{array}{ccc} V_\delta & \xrightarrow{v_{\delta, \delta+2\alpha}} & V_{\delta+2\alpha} \\ & \searrow \phi_\delta & \nearrow \gamma_{\delta+\alpha} \\ & & W_{\delta+\alpha} \end{array} \quad (3.1)$$

$$\begin{array}{ccc} & & V_{\delta+\alpha} \\ & \nearrow \gamma_\delta & \searrow \phi_{\delta+\alpha} \\ W_\delta & \xrightarrow{w_{\delta, \delta+2\alpha}} & W_{\delta+2\alpha} \end{array} \quad (3.2)$$

$$\begin{array}{ccccc} V_\delta & \xrightarrow{v_{\delta, \delta'}} & V_{\delta'} & & \\ & \searrow \phi_\delta & & \searrow \phi_{\delta'} & \\ & & W_{\delta+\alpha} & \xrightarrow{w_{\delta+\alpha, \delta'+\alpha}} & W_{\delta'+\alpha} \end{array} \quad (3.3)$$

$$\begin{array}{ccc}
& & V_{\delta+\alpha} \xrightarrow{v_{\delta+\alpha, \delta'+\alpha}} V_{\delta'+\alpha} \\
& \nearrow \gamma_\delta & \\
W_\delta & \xrightarrow{w_{\delta, \delta'}} & W_{\delta'} \xrightarrow{\gamma_{\delta'}} \\
& &
\end{array} \tag{3.4}$$

In what follows, for each non-negative integer k , $H_k(\cdot)$ will denote the homology functor (with field coefficients). Given a pair (X, φ_X) and a non-negative integer k , we define the associated sublevelset persistence module

$$\mathfrak{P}_k^X := \{V_\delta \xrightarrow{v_{\delta, \delta'}} V_{\delta'}\}_{\delta \leq \delta'}$$

where:

- for each $\delta \in \mathbb{R}$, $V_\delta := H_k(X^\delta)$, where $X^\delta := \varphi_X^{-1}((-\infty, \delta])$.
- for each $\delta, \delta' \in \mathbb{R}$ with $\delta \leq \delta'$ the linear map $v_{\delta, \delta'} := H_k(\iota_{\delta, \delta'}^X : X^\delta \hookrightarrow X^{\delta'})$ induced by the natural inclusion $\iota_{\delta, \delta'}^X$ of X^δ into $X^{\delta'}$.

3.2. Lifting stability results via d_{HT}

In this section we obtain a lower bound for the homotopy type distance based on comparing persistence diagrams.

The context of the following theorem is that of the full subcategory \mathbf{C} of \mathbf{S} whose objects are compact polyhedra endowed with continuous real valued functions. Thus, the associated sublevelset persistence modules turn out to be q -tame [10], and one can still obtain persistence diagrams for sublevelset persistence without adding any extra tameness condition on the functions [8]. Moreover, because \mathbf{C} is a full subcategory of \mathbf{S} , $d_{\text{HT}}^{\mathbf{C}}$ coincides with $d_{\text{HT}}^{\mathbf{S}}$ restricted to objects of \mathbf{C} . Thus we can simply write d_{HT} for $d_{\text{HT}}^{\mathbf{C}}$.

We start proving that the persistence modules of α -homotopic pairs are α -interleaved.

Lemma 3.1. *Let (X, φ_X) and (Y, φ_Y) be two α -homotopy equivalent pairs in \mathbf{C} . Then, for every non-negative integer k , the persistence modules \mathfrak{P}_k^X and \mathfrak{P}_k^Y are α -interleaved.*

Proof. Fix a non-negative integer k , and write

$$\mathfrak{P}_k^X = \{V_\delta \xrightarrow{v_{\delta, \delta'}} V_{\delta'}\}_{\delta \leq \delta'} \text{ and } \mathfrak{P}_k^Y = \{W_\delta \xrightarrow{w_{\delta, \delta'}} W_{\delta'}\}_{\delta \leq \delta'}$$

where:

- for each $\delta \in \mathbb{R}$ $V_\delta := H_k(X^\delta)$ and $W_\delta := H_k(Y^\delta)$;
- for each $\delta, \delta' \in \mathbb{R}$ with $\delta \leq \delta'$ the maps $v_{\delta, \delta'} := H_k(\iota_{\delta, \delta'}^X)$ and $w_{\delta, \delta'} := H_k(\iota_{\delta, \delta'}^Y)$.

Let (f, g) be a pair of α -homotopy equivalences with respect to (φ_X, φ_Y) . For each $\delta \in \mathbb{R}$, let $f_\delta := f|_{X^\delta}$ and $g_\delta := g|_{Y^\delta}$ denote the restrictions of f and g to X^δ and Y^δ , respectively. Notice that from the fact that f and g are α -maps it follows that $\text{im}(f_\delta) \subset Y^{\delta+\alpha}$ and $\text{im}(g_\delta) \subset X^{\delta+\alpha}$ for each $\delta \in \mathbb{R}$.

In order to prove that \mathfrak{P}_k^X and \mathfrak{P}_k^Y are α -interleaved, for each $\delta \in \mathbb{R}$ we need to provide maps $\phi_\delta : V_\delta \rightarrow W_{\delta+\alpha}$ and $\gamma_\delta : W_\delta \rightarrow V_{\delta+\alpha}$ such that the four diagrams (3.1), (3.2), (3.3), and (3.4) commute for all $\delta, \delta' \in \mathbb{R}$ with $\delta \leq \delta'$.

In order to establish the commutativity of (3.1) consider the following diagram of topological spaces:

$$\begin{array}{ccc}
 X^\delta & \xrightarrow{\iota_{\delta, \delta+2\alpha}^X} & X^{\delta+2\alpha} \\
 & \searrow f_\delta & \nearrow g_{\delta+\alpha} \\
 & Y^{\delta+\alpha} &
 \end{array} \tag{3.5}$$

Notice that $g_{\delta+\alpha} \circ f_\delta : X^\delta \rightarrow X^{\delta+2\alpha}$ and the inclusion $\iota_{\delta, \delta+2\alpha}^X : X^\delta \hookrightarrow X^{\delta+2\alpha}$ are homotopic by hypothesis. Indeed, since a 2α -homotopy $H : X \times [0, 1] \rightarrow X$ between $g \circ f$ and id_X exists, the restriction of H to $X^\delta \times [0, 1]$ has its image contained in $X^{\delta+2\alpha}$, and is therefore a proper homotopy between the maps $g_{\delta+\alpha} \circ f_\delta$ and $\iota_{\delta, \delta+2\alpha}^X$. Thus $H_k(g_{\delta+\alpha} \circ f_\delta) = H_k(\iota_{\delta, \delta+2\alpha}^X)$. Applying the homology functor to diagram (3.5) therefore yields the commutative diagram (3.1). The commutativity of diagram (3.2) can be established in a similar way.

In order to establish the commutativity of (3.3) consider the following diagram of topological spaces:

$$\begin{array}{ccccc}
 X^\delta & \xrightarrow{\iota_{\delta, \delta'}^X} & X^{\delta'} & & \\
 & \searrow f_\delta & & \searrow f_{\delta'} & \\
 & & Y^{\delta+\alpha} & \xrightarrow{\iota_{\delta+\alpha, \delta'+\alpha}^Y} & Y^{\delta'+\alpha}
 \end{array} \tag{3.6}$$

We now verify that this diagram commutes so that the commutativity of (3.3) follows by applying the homology functor to (3.6). Indeed, pick any $x \in X^\delta$. Then

$$\begin{aligned}
 f_{\delta'}(\iota_{\delta, \delta'}^X(x)) &= f_{\delta'}(x) && \text{(Since } \iota_{\delta, \delta'}^X \text{ is the inclusion map)} \\
 &= f|_{X^{\delta'}}(x) && \text{(By definition of } f_{\delta'}) \\
 &= f|_{X^\delta}(x) && \text{(Since } x \in X^\delta \subseteq X^{\delta'}) \\
 &= f_\delta(x) && \text{(Definition of } f_\delta) \\
 &= \iota_{\delta+\alpha, \delta'+\alpha}^Y(f_\delta(x)). && \text{(Since } f_\delta(x) \in Y^{\delta+\alpha} \subseteq Y^{\delta'+\alpha})
 \end{aligned}$$

Since $x \in X^\delta$ was arbitrary it follows that $f_{\delta'+\alpha} \circ \iota_{\delta, \delta'}^X = \iota_{\delta+\alpha, \delta'+\alpha} \circ f_\delta$.

One can verify that (3.4) commutes using a similar argument. \square

Using Lemma 3.1, we now obtain the stability of persistence diagrams with respect to the persistent homotopy type distance.

Theorem 2. *Let X and Y be compact polyhedra, and k be any non-negative integer. Let $\varphi_X : X \rightarrow \mathbb{R}$ and $\varphi_Y : Y \rightarrow \mathbb{R}$ be continuous functions. Then,*

$$d_B(D_k(\varphi_X), D_k(\varphi_Y)) \leq d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)).$$

Proof. Under the assumption that X and Y are compact polyhedra, and $\varphi_X : X \rightarrow \mathbb{R}$ and $\varphi_Y : Y \rightarrow \mathbb{R}$ are continuous functions, the persistence modules \mathfrak{P}_k^X and \mathfrak{P}_k^Y are

q -tame by [8, Thm. 2.3]. By Lemma 3.1, if (X, φ_X) and (Y, φ_Y) are α -homotopy equivalent with respect to (φ_X, φ_Y) , then, the persistence modules \mathfrak{P}_k^X and \mathfrak{P}_k^Y are α -interleaved. Thus, the claim follows from the stability theorem for q -tame persistence modules over \mathbb{R} [10]. \square

4. The homotopy type distance for comparing merge trees

A merge tree is a structural descriptor used in shape analysis. For a continuous function $\varphi : X \rightarrow \mathbb{R}$ defined on a connected domain, the merge tree of φ encodes how the sublevel sets $\varphi^{-1}((-\infty, t])$ are connected for increasing values of $t \in \mathbb{R}$.

Following [23], it can be defined as follows. Consider the epigraph $\text{epi}(\varphi)$ of φ , that is the space

$$\text{epi}(\varphi) := \{(x, t) \in X \times \mathbb{R} : \varphi(x) \leq t\},$$

and the function $\bar{\varphi} : \text{epi}(\varphi) \rightarrow \mathbb{R}$ defined by $\bar{\varphi}(x, t) = t$. Consider the equivalence relation \sim on $\text{epi}(\varphi)$ defined by setting $(x, t) \sim (x', t')$ if and only if $t = t'$ and (x, t) and (x', t') belong to the same connected component of $\bar{\varphi}^{-1}(t)$. The merge tree of φ , denoted M_φ , is the quotient space $\text{epi}(\varphi)/\sim$. In other words, M_φ is the Reeb graph [24] of $\text{epi}(\varphi)$ with respect to $\bar{\varphi}$. M_φ is naturally endowed with the continuous function $\hat{\varphi} : M_\varphi \rightarrow \mathbb{R}$ defined by setting $\hat{\varphi}(p) := \bar{\varphi}(x)$ for any point x belonging to the equivalence class p .

Because M_φ is the Reeb graph of $\text{epi}(\varphi)$ with respect to $\bar{\varphi}$, the assumption that X is a compact polyhedron and φ is piecewise linear ensures that $\text{epi}(\varphi)/\sim$ is a (non-compact) polyhedron of dimension 1 [13]. In particular, M_φ turns out to be a tree with finitely many leaves. As such, merge trees can be compared using the persistent homotopy type distance. It is then interesting to study the persistent homotopy type distance on merge trees in relation to other distances that have been proposed for the same goal.

The manuscript [23] presents an interleaving distance between merge trees that is interesting because it satisfies a stability property with respect to perturbation of the function that defines the merge tree. The interleaving distance between merge trees is defined as follows. For $\varepsilon \geq 0$, define the ε -shift map $i_\varphi^\varepsilon : M_\varphi \rightarrow M_\varphi$ that sends, for every $t \in \mathbb{R}$, a connected component of $\bar{\varphi}^{-1}(t)$ to the connected component of $\bar{\varphi}^{-1}(t + \varepsilon)$ that contains it. In other words, i_φ^ε is the map induced by the inclusion of the sublevel sets of φ . Given two continuous functions $\varphi, \psi : X \rightarrow \mathbb{R}$, and $\varepsilon \geq 0$, consider the two ε -shift maps $i_\varphi^\varepsilon : M_\varphi \rightarrow M_\varphi$ and $i_\psi^\varepsilon : M_\psi \rightarrow M_\psi$. Two continuous maps $f^\varepsilon : M_\varphi \rightarrow M_\psi$ and $g^\varepsilon : M_\psi \rightarrow M_\varphi$ are said to be ε -compatible if the following diagrams commute

$$\begin{array}{ccc} M_\varphi & \xrightarrow{i_\varphi^{2\varepsilon}} & M_\varphi \\ & \searrow f^\varepsilon & \nearrow g^\varepsilon \\ & & M_\psi \end{array} \quad \begin{array}{ccc} & & M_\varphi \\ & \nearrow g^\varepsilon & \searrow f^\varepsilon \\ M_\psi & \xrightarrow{i_\psi^{2\varepsilon}} & M_\psi \end{array} \quad (4.1)$$

and, moreover, $\hat{\psi} \circ f^\varepsilon = \hat{\varphi} + \varepsilon$ and $\hat{\varphi} \circ g^\varepsilon = \hat{\psi} + \varepsilon$. In this setting, the interleaving distance, $d_I(M_\varphi, M_\psi)$ between two merge trees is the greatest lower bound on ε for

which there are ε -compatible maps [23]:

$$d_I(M_\varphi, M_\psi) = \inf\{\varepsilon \geq 0 : \exists \varepsilon\text{-compatible maps } f^\varepsilon : M_\varphi \rightarrow M_\psi, g^\varepsilon : M_\psi \rightarrow M_\varphi \}.$$

We prove that the persistent homotopy distance on merge trees coincides with such interleaving distance.

Proposition 4.1. *For every pair of piecewise linear functions $\varphi, \psi : X \rightarrow \mathbb{R}$ defined on a compact connected polyhedron X , it holds that*

$$d_I(M_\varphi, M_\psi) = d_{\text{HT}}((M_\varphi, \hat{\varphi}), (M_\psi, \hat{\psi})).$$




Figure 5: Elements of the proof of Proposition 4.1.

Proof. In order to see that $d_{\text{HT}} \leq d_I$, we will prove that, given $\varepsilon \geq 0$, every pair of ε -compatible maps between M_φ and M_ψ constitutes a pair of ε -homotopy equivalences. Pick $\varepsilon \geq 0$ and let $i_\varphi^\varepsilon : M_\varphi \rightarrow M_\varphi$ and $i_\psi^\varepsilon : M_\psi \rightarrow M_\psi$ be ε -shifts on M_φ and M_ψ , respectively. Assume that $f^\varepsilon : M_\varphi \rightarrow M_\psi$ and $g^\varepsilon : M_\psi \rightarrow M_\varphi$ are ε -compatible. Because $\hat{\psi} \circ f^\varepsilon = \hat{\varphi} + \varepsilon$ and $\hat{\varphi} \circ g^\varepsilon = \hat{\psi} + \varepsilon$, it follows that f^ε and g^ε are ε -maps. Let us see that $(f^\varepsilon, g^\varepsilon)$ is a pair of ε -homotopy equivalences with respect to $(\hat{\varphi}, \hat{\psi})$. To this end, we need to construct a 2ε -homotopy with respect to $(\hat{\varphi}, \hat{\varphi})$ (resp. $(\hat{\psi}, \hat{\psi})$) between the identity of M_φ (resp. M_ψ) and $g^\varepsilon \circ f^\varepsilon$ (resp. $f^\varepsilon \circ g^\varepsilon$). However, from the assumption that f^ε and g^ε are ε -compatible, we get $i_\varphi^{2\varepsilon} = g^\varepsilon \circ f^\varepsilon$ and $i_\psi^{2\varepsilon} = f^\varepsilon \circ g^\varepsilon$. Thus, equivalently, we need to construct a 2ε -homotopy H (resp. K) with respect to $(\hat{\varphi}, \hat{\varphi})$ (resp. $(\hat{\psi}, \hat{\psi})$) between the identity on M_φ (resp. M_ψ) and the 2ε -shift $i_\varphi^{2\varepsilon}$ (resp. $i_\psi^{2\varepsilon}$). We show how to construct H . Recall that M_φ is a tree, and that for every $p \in M_\varphi$ there are finitely many leaves $l_1, \dots, l_{n_p} \in M_\varphi$ which can be connected to p by a simple path increasing under $\hat{\varphi}$. For $i = 1, \dots, n_p$, set $\delta_i^p := (\hat{\varphi}(p) - \hat{\varphi}(l_i))/2\varepsilon \in \mathbb{R}$, and let $\gamma_i^p : [-\delta_i^p, +\infty) \rightarrow M_\varphi$ be the path increasing under $\hat{\varphi}$ with starting point l_i (i.e. $\hat{\varphi}(-\delta_i^p) = l_i$), passing through p , and parametrized by $\hat{\varphi} \circ \gamma_i^p(t) = \hat{\varphi}(p) + 2\varepsilon t$ with $t \in [-\delta_i^p, +\infty)$. Note that, as parameter t increases, the paths γ_i^p , with $i = 1, \dots, n_p$,

can only merge and not branch, and that $\gamma_i^p(0) = p$ for every i . See Figure 5 for notation and an illustration of intervening elements.

Setting $H : M_\varphi \times [0, 1] \rightarrow M_\varphi$, $(p, s) \mapsto \gamma_i^p(s)$ gives a well-defined set map because it does not depend on the choice of the leaf ℓ_i . Notice that $H(p, 0) = p$, $H(p, 1) = i^{2\varepsilon}(p)$, and $\hat{\varphi}(H(p, s)) = \hat{\varphi}(p) + 2\varepsilon s$ for all $p \in M_\varphi$ and $s \in [0, 1]$.

It remains to show that H is continuous. In the case when $\varepsilon = 0$, $H(p, s) = p$ for every $(p, s) \in M_\varphi \times [0, 1]$, and so H is the identity map of M_φ for every $s \in [0, 1]$, implying the continuity of H in that case. Now we pick $\varepsilon > 0$. Let q_0 belong to the image of H , that is $q_0 = H(p_0, s_0)$ for some $(p_0, s_0) \in M_\varphi \times [0, 1]$, and let V be an open set in M_φ containing q_0 . For every $\eta > 0$ and $q \in M_\varphi$, let $U(q, \eta)$ be the connected component of $\hat{\varphi}^{-1}((\hat{\varphi}(q) - \eta, \hat{\varphi}(q) + \eta))$ containing q . Let $\eta > 0$ be small enough such that $U(q_0, \eta)$ is contained in V . Set $J(s_0) = (s_0 - \frac{\eta}{4\varepsilon}, s_0 + \frac{\eta}{4\varepsilon}) \cap [0, 1]$. Note that $U(p_0, \eta/2) \times J(s_0)$ is an open neighborhood of (p_0, s_0) in $M_\varphi \times [0, 1]$. By construction, for every $(p, s) \in U(p_0, \eta/2) \times J(s_0)$, it holds that $\hat{\varphi}(H(p, s)) = \hat{\varphi}(\gamma_i^p(s)) = \hat{\varphi}(p) + 2\varepsilon s$. Moreover, $(p, s) \in U(p_0, \eta/2) \times J(s_0)$ implies that $\hat{\varphi}(p_0) - \eta/2 < \hat{\varphi}(p) < \hat{\varphi}(p_0) + \eta/2$ and $s_0 - \frac{\eta}{4\varepsilon} < s < s_0 + \frac{\eta}{4\varepsilon}$, so that

$$\hat{\varphi}(p_0) - \frac{\eta}{2} + 2\varepsilon \left(s_0 - \frac{\eta}{4\varepsilon} \right) < \hat{\varphi}(H(p, s)) < \hat{\varphi}(p_0) + \frac{\eta}{2} + 2\varepsilon \left(s_0 + \frac{\eta}{4\varepsilon} \right).$$

Because $q_0 = H(p_0, s_0)$ implies that $\hat{\varphi}(q_0) = \hat{\varphi}(p_0) + 2\varepsilon s_0$, we deduce that

$$\hat{\varphi}(q_0) - \eta < \hat{\varphi}(H(p, s)) < \hat{\varphi}(q_0) + \eta,$$

yielding $H(U(p_0, \eta/2) \times J(s_0)) \subseteq U(q_0, \eta) \subseteq V$. Indeed, we observe that $H(p, s)$ belongs to the connected component of $\hat{\varphi}^{-1}((\hat{\varphi}(q_0) - \eta, \hat{\varphi}(q_0) + \eta))$ containing q_0 . In other words, H is continuous.

So, H is a 2ε -homotopy between the identity on M_φ and $i_\varphi^{2\varepsilon}$. In a similar way, we can build a 2ε -homotopy K between the identity on M_ψ and $i_\psi^{2\varepsilon}$. So, we have finally proved that $(f^\varepsilon, g^\varepsilon)$ is a pair of ε -homotopy equivalences with respect to $(\hat{\varphi}, \hat{\psi})$.

Conversely, in order to see that $d_I \leq d_{HT}$, let $f : M_\varphi \rightarrow M_\psi$ and $g : M_\psi \rightarrow M_\varphi$ be such that (f, g) is a pair of ε -homotopy equivalences with respect to $(\hat{\varphi}, \hat{\psi})$ for some $\varepsilon \geq 0$. For every $p \in M_\varphi$, let

$$f^\varepsilon(p) := i_\psi^{\hat{\varphi}(p) + \varepsilon - \hat{\psi}(f(p))} \circ f(p).$$

Analogously, for every $q \in M_\psi$, let

$$g^\varepsilon(q) := i_\varphi^{\hat{\psi}(q) + \varepsilon - \hat{\varphi}(g(q))} \circ g(q).$$

In plain words, f^ε shifts the point $f(p)$ of M_ψ in the direction of increasing function values until it reaches the level $\hat{\varphi}(p) + \varepsilon$, and similarly g^ε shifts the point $g(q)$ of M_φ in the direction of increasing function values until it reaches the level $\hat{\psi}(q) + \varepsilon$. By construction, $\hat{\psi} \circ f^\varepsilon = \hat{\varphi} + \varepsilon$, $\hat{\varphi} \circ g^\varepsilon = \hat{\psi} + \varepsilon$, $f^\varepsilon \circ g^\varepsilon = i_\psi^{2\varepsilon}$ and $g^\varepsilon \circ f^\varepsilon = i_\varphi^{2\varepsilon}$. To prove that f^ε and g^ε are ε -compatible maps, it remains to check that they are continuous, which is implied by the structure of the tree M_φ . □

5. d_{HT} can be seen as an interleaving distance using categories

The goal of this section is to re-interpret some of the material contained in the previous sections in terms of interleavings. The advantage is that we obtain a unifying look at the distances we have encountered so far.

The theory of interleavings was initiated by Chazal et al. in [9], further developed by Lesnick in [22] to comprise also functional categories, and by Bubenik and Scott in [4] for functors from the category of ordered reals, and extended by Bubenik et al. in [5] to the case of functors from any preordered set.

5.1. Functorial definition of interleaving distance

For a given integer $n \geq 1$ we denote by \mathbf{R}^n the poset category with object set \mathbb{R}^n with a morphism between \mathbf{u} and \mathbf{v} in \mathbb{R}^n iff $\mathbf{u} \preceq \mathbf{v}$.

Let \mathbf{O} be an arbitrary category. We start with the general definition of interleaving distance between functors from \mathbf{R}^n to \mathbf{O} following [5].

Definition 5.1. Let $T : \mathbf{R}^n \rightarrow \mathbf{O}$ be a functor between \mathbf{R}^n and \mathbf{O} and $\varepsilon \succeq \mathbf{0}$. The ε -shift of T is the functor $T_\varepsilon : \mathbf{R}^n \rightarrow \mathbf{O}$ such that:

1. $T_\varepsilon(\mathbf{u}) = T(\mathbf{u} + \varepsilon)$;
2. $T_\varepsilon(\mathbf{u} \preceq \mathbf{v}) = T(\mathbf{u} + \varepsilon \preceq \mathbf{v} + \varepsilon)$.

Given two functors $T, T' : \mathbf{R}^n \rightarrow \mathbf{O}$, a natural transformation $\xi : T \Rightarrow T'$ between these functors consists of a morphism $\xi_{\mathbf{u}} : T(\mathbf{u}) \rightarrow T'(\mathbf{u})$ in \mathbf{O} for every $\mathbf{u} \in \mathbb{R}^n$, such that, for every $\mathbf{u} \preceq \mathbf{v} \in \mathbb{R}^n$, the following diagram commutes:

$$\begin{array}{ccc} T(\mathbf{u}) & \xrightarrow{\xi_{\mathbf{u}}} & T'(\mathbf{u}) \\ T(\mathbf{u} \preceq \mathbf{v}) \downarrow & & \downarrow T'(\mathbf{u} \preceq \mathbf{v}) \\ T(\mathbf{v}) & \xrightarrow{\xi_{\mathbf{v}}} & T'(\mathbf{v}) \end{array} \quad (5.1)$$

Given $\varepsilon \in \mathbb{R}$ by $\vec{\varepsilon} \in \mathbb{R}^n$ we denote the vector with all components equal to ε .

Definition 5.2. Given two functors $T, T' : \mathbf{R}^n \rightarrow \mathbf{O}$, and a real number $\varepsilon \geq 0$, T and T' are said to be ε -interleaved if there exist natural transformations $\xi : T \Rightarrow T'_\varepsilon$ and $\eta : T' \Rightarrow T_\varepsilon$ such that, for every $\mathbf{u} \in \mathbb{R}^n$, $\eta_{\mathbf{u} + \vec{\varepsilon}} \circ \xi_{\mathbf{u}} = T(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$ and $\xi_{\mathbf{u} + \vec{\varepsilon}} \circ \eta_{\mathbf{u}} = T'(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$. Moreover, in that case, the pair (ξ, η) is called an ε -interleaving between T and T' .

Definition 5.3. Given two functors $T, T' : \mathbf{R}^n \rightarrow \mathbf{O}$, the *interleaving distance* between T and T' is defined as

$$d_{\mathbf{I}}^{\mathbf{O}}(T, T') := \inf\{\varepsilon \geq 0 : T \text{ and } T' \text{ are } \varepsilon\text{-interleaved}\}$$

whenever T and T' are ε -interleaved for some real number $\varepsilon \geq 0$, while $d_{\mathbf{I}}^{\mathbf{O}}(T, T') = \infty$, otherwise.

Proposition 5.1 ([5]). *Let \mathbf{O} be an arbitrary category, and let $\mathbf{O}^{\mathbf{R}}$ be the category of functors from \mathbf{R} to \mathbf{O} . It holds that $d_{\mathbf{I}}^{\mathbf{O}}$ is an extended pseudo-metric on the objects of $\mathbf{O}^{\mathbf{R}}$.*

Next, we consider specific choices of the target category \mathbf{O} . When there is no risk of confusion, in order to avoid an overload of notation, we will avoid specifying the \mathbf{O} symbol in the notation $d_1^{\mathbf{O}}$.

5.1.1. The interleaving distance between \mathbb{R}^n -valued functions

Let $\mathbf{O} = \mathbf{Top}/\mathbb{R}_{\geq}^n$ be the category of continuous functions $\varphi_X : X \rightarrow \mathbb{R}^n$ as objects, and $\mathbf{0}$ -maps as morphisms: a morphism from φ_X to φ_Y is a continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$.

Proposition 5.2. *Every continuous function $\varphi_X : X \rightarrow \mathbb{R}^n$ defines a functor $T^{\varphi_X} : \mathbf{R}^n \rightarrow \mathbf{Top}/\mathbb{R}_{\geq}^n$ by setting*

- for every $\mathbf{u} \in \mathbb{R}^n$, $T^{\varphi_X}(\mathbf{u}) := \varphi_X - \mathbf{u}$;
- for every $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $\mathbf{u} \preceq \mathbf{v}$, $T^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) := \text{id}_X$.

Proof. For $\mathbf{u} \preceq \mathbf{v}$, $\text{id}_X : X \rightarrow X$ is a morphism between $\varphi_X - \mathbf{u}$ and $\varphi_X - \mathbf{v}$ because $(\varphi_X - \mathbf{v}) \circ \text{id}_X \preceq \varphi_X - \mathbf{u}$. Moreover, T^{φ_X} preserves identity and composition. \square

Let us now see what ε -shifts look like in this setting.

Proposition 5.3. *For every $\varepsilon \succeq \mathbf{0}$, and for every φ_X , the ε -shift of T^{φ_X} , $T_{\varepsilon}^{\varphi_X} : \mathbf{R}^n \rightarrow \mathbf{Top}/\mathbb{R}_{\geq}^n$, is equal to the functor $T^{\varphi_X - \varepsilon} : \mathbf{R}^n \rightarrow \mathbf{Top}/\mathbb{R}_{\geq}^n$.*

Proof. By Definition 5.1 and the definition of T^{φ_X} in Proposition 5.2, the ε -shift of T^{φ_X} is the functor $T_{\varepsilon}^{\varphi_X} : \mathbf{R}^n \rightarrow \mathbf{Top}/\mathbb{R}_{\geq}^n$ such that

1. for every $\mathbf{u} \in \mathbb{R}^n$, $T_{\varepsilon}^{\varphi_X}(\mathbf{u}) = T^{\varphi_X}(\mathbf{u} + \varepsilon) = \varphi_X - \mathbf{u} - \varepsilon = T^{\varphi_X - \varepsilon}(\mathbf{u})$;
2. for every $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $\mathbf{u} \preceq \mathbf{v}$, $T_{\varepsilon}^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = T^{\varphi_X}(\mathbf{u} + \varepsilon \preceq \mathbf{v} + \varepsilon) = \text{id}_X = T^{\varphi_X - \varepsilon}(\mathbf{u} \preceq \mathbf{v})$.

\square

The next lemma characterizes all the natural transformations between pairs of functors T^{φ_X} and T^{φ_Y} .

Lemma 5.1. *Let X, Y be topological spaces, and let $\varphi_X : X \rightarrow \mathbb{R}^n$, $\varphi_Y : Y \rightarrow \mathbb{R}^n$ be any two continuous functions. Then, every continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$ induces a natural transformation $\xi^f : T^{\varphi_X} \Rightarrow T^{\varphi_Y}$ such that, for every $\mathbf{u} \in \mathbb{R}^n$, $\xi^f(\mathbf{u})$ is equal to f . Reciprocally, to every natural transformation $\xi : T^{\varphi_X} \Rightarrow T^{\varphi_Y}$ corresponds a continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$ defined by $f = \xi(\mathbf{u})$ for any $\mathbf{u} \in \mathbb{R}^n$.*

Proof. Any continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$ induces a natural transformation $\xi^f : T^{\varphi_X} \Rightarrow T^{\varphi_Y}$ defined as follows: for every $\mathbf{u} \in \mathbb{R}^n$, $\xi^f(\mathbf{u}) := f$. Indeed, for every $\mathbf{u} \in \mathbb{R}^n$, f is a morphism between $\varphi_X - \mathbf{u}$ and $\varphi_Y - \mathbf{u}$ because $\varphi_Y \circ f \preceq \varphi_X$ implies $(\varphi_Y - \mathbf{u}) \circ f \preceq \varphi_X - \mathbf{u}$, proving that $\xi^f(\mathbf{u}) : T^{\varphi_X}(\mathbf{u}) \rightarrow T^{\varphi_Y}(\mathbf{u})$. Moreover, $f \circ \text{id}_X = \text{id}_Y \circ f$, proving that $\xi^f(\mathbf{u}) \circ T^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = T^{\varphi_Y}(\mathbf{u} \preceq \mathbf{v}) \circ \xi^f(\mathbf{v})$.

Reciprocally, assume that there is a family $\xi = \{\xi(\mathbf{u}) : T^{\varphi_X}(\mathbf{u}) \rightarrow T^{\varphi_Y}(\mathbf{u})\}_{\mathbf{u} \in \mathbb{R}^n}$ of continuous maps such that diagram (5.1) commutes for every $\mathbf{u} \preceq \mathbf{v} \in \mathbb{R}^n$:

$$\xi(\mathbf{u}) \circ T^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = T^{\varphi_Y}(\mathbf{u} \preceq \mathbf{v}) \circ \xi(\mathbf{v}).$$

Equivalently, for $\mathbf{u} \preceq \mathbf{v}$, $\xi(\mathbf{v}) \circ \text{id}_X = \text{id}_Y \circ \xi(\mathbf{u})$. Hence, $\xi(\mathbf{u}) = \xi(\mathbf{v})$ for every $\mathbf{u} \preceq \mathbf{v}$. Hence, it is sufficient to define $f := \xi(\mathbf{u})$ for one and hence all $\mathbf{u} \in \mathbb{R}^n$. \square

We can now show that the interleaving distance between functors T^{φ_X} and T^{φ_Y} in $\mathbf{Top}/\mathbb{R}_{\geq}^n$ coincides with the natural pseudo-distance.

Proposition 5.4. *Let X and Y be topological spaces. Denoting by d_{NP} the natural pseudo-distance, for any pair of continuous functions $\varphi_X : X \rightarrow \mathbb{R}^n$ and $\varphi_Y : Y \rightarrow \mathbb{R}^n$, it holds that*

$$d_{\mathbb{I}}(T^{\varphi_X}, T^{\varphi_Y}) = d_{\text{NP}}((X, \varphi_X), (Y, \varphi_Y)).$$

Proof. If X and Y are not homeomorphic, $d_{\text{NP}} = \infty$ trivially implying that $d_{\mathbb{I}} \leq d_{\text{NP}}$. Let $f : X \rightarrow Y$ be a homeomorphism such that $\|\varphi_X - \varphi_Y \circ f\|_{\infty} \leq \varepsilon$ for some $\varepsilon \geq 0$. It then holds that $(\varphi_Y - \vec{\varepsilon}) \circ f \preceq \varphi_X$ and $(\varphi_X - \vec{\varepsilon}) \circ f^{-1} \preceq \varphi_Y$. By Lemma 5.1, f induces a natural transformation ξ^f from T^{φ_X} to $T^{(\varphi_Y - \vec{\varepsilon})}$, as well as a natural transformation $\xi^{f^{-1}}$ from T^{φ_Y} to $T^{(\varphi_X - \vec{\varepsilon})}$. By Proposition 5.3, $T^{(\varphi_Y - \vec{\varepsilon})} = T_{\vec{\varepsilon}}^{\varphi_Y}$ and $T^{(\varphi_X - \vec{\varepsilon})} = T_{\vec{\varepsilon}}^{\varphi_X}$. Moreover, for every $\mathbf{u} \in \mathbb{R}^n$, $\xi^f(\mathbf{u}) = f$, and analogously $\xi^{f^{-1}}(\mathbf{u} + \vec{\varepsilon}) = f^{-1}$. Thus, we get $\xi^{f^{-1}}(\mathbf{u} + \vec{\varepsilon}) \circ \xi^f(\mathbf{u}) = f^{-1} \circ f = \text{id}_X = T^{\varphi_X}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$. Similarly, $\xi^f(\mathbf{u} + \vec{\varepsilon}) \circ \xi^{f^{-1}}(\mathbf{u}) = f \circ f^{-1} = \text{id}_Y = T^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$. Hence, the pair $(\xi^f, \xi^{f^{-1}})$ is an ε -interleaving between T^{φ_X} and T^{φ_Y} , proving that $d_{\mathbb{I}} \leq d_{\text{NP}}$ also in this case.

Let us now show that $d_{\text{NP}} \leq d_{\mathbb{I}}$. The claim is obvious if $d_{\mathbb{I}} = \infty$, so let us assume that T^{φ_X} and T^{φ_Y} are ε -interleaved for some $\varepsilon \geq 0$ by an ε -interleaving (ξ, η) with $\xi : T^{\varphi_X} \Rightarrow T_{\vec{\varepsilon}}^{\varphi_Y}$ and $\eta : T^{\varphi_Y} \Rightarrow T_{\vec{\varepsilon}}^{\varphi_X}$. By Definition 5.2, for every $\mathbf{u} \in \mathbb{R}^n$, $\eta(\mathbf{u} + \vec{\varepsilon}) \circ \xi(\mathbf{u}) = T^{\varphi_X}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$, and $\xi(\mathbf{u} + \vec{\varepsilon}) \circ \eta(\mathbf{u}) = T^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$. By Lemma 5.1 and Proposition 5.3, there are two continuous maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that, for each $\mathbf{u} \in \mathbb{R}^n$, $\xi(\mathbf{u}) = f$, $\eta(\mathbf{u}) = g$, $(\varphi_Y - \vec{\varepsilon}) \circ f \preceq \varphi_X$ and $(\varphi_X - \vec{\varepsilon}) \circ g \preceq \varphi_Y$. From $\eta(\mathbf{u} + \vec{\varepsilon}) \circ \xi(\mathbf{u}) = T^{\varphi_X}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$ and $\xi(\mathbf{u} + \vec{\varepsilon}) \circ \eta(\mathbf{u}) = T^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$, it follows that $g \circ f = \text{id}_X$ and $f \circ g = \text{id}_Y$. Hence, $g = f^{-1}$ and $\|\varphi_X - \varphi_Y \circ f\|_{\infty} \leq \varepsilon$. \square

5.1.2. The interleaving distance between functions up to homotopy

We now consider $\mathbf{O} = \mathbf{hTop}/\mathbb{R}_{\geq}^n$, the category with continuous functions $\varphi_X : X \rightarrow \mathbb{R}^n$ as objects, and \mathbf{O} -homotopy classes of \mathbf{O} -maps between X and Y as morphisms: for two objects φ_X, φ_Y in $\mathbf{hTop}/\mathbb{R}_{\geq}^n$, a morphism from φ_X to φ_Y is the \mathbf{O} -homotopy class with respect to (φ_X, φ_Y) of a \mathbf{O} -map $f : X \rightarrow Y$, that is, a continuous map between X and Y such that $\varphi_Y \circ f \preceq \varphi_X$, and is denoted by $[f]_{(\varphi_X, \varphi_Y)}$. In $\mathbf{hTop}/\mathbb{R}_{\geq}^n$, the composition of morphisms is defined as the \mathbf{O} -homotopy class of the composition of \mathbf{O} -maps: for $\varphi_X, \varphi_Y, \varphi_Z$ in $\mathbf{hTop}/\mathbb{R}_{\geq}^n$, and $f : X \rightarrow Y, g : Y \rightarrow Z$ being \mathbf{O} -maps with respect to (φ_X, φ_Y) and (φ_Y, φ_Z) , respectively, we set

$$[g]_{(\varphi_Y, \varphi_Z)} \circ [f]_{(\varphi_X, \varphi_Y)} = [g \circ f]_{(\varphi_X, \varphi_Z)}.$$

This is well defined because the composition does not depend on the representatives, and $\varphi_Z \circ g \leq \varphi_Y, \varphi_Y \circ f \preceq \varphi_X$ imply $\varphi_Z \circ g \circ f \preceq \varphi_X$. Composition is associative and, for any object (X, φ_X) , the \mathbf{O} -homotopy class of the identity id_X with respect to (φ_X, φ_X) is a morphism from (X, φ_X) to (X, φ_X) in $\mathbf{hTop}/\mathbb{R}_{\geq}^n$.

Proposition 5.5. *Every continuous function $\varphi_X : X \rightarrow \mathbb{R}^n$ defines a functor $hT^{\varphi_X} : \mathbf{R} \rightarrow \mathbf{hTop}/\mathbb{R}_{\geq}^n$ by setting*

- for every $\mathbf{u} \in \mathbb{R}^n$, $hT^{\varphi_X}(\mathbf{u}) := \varphi_X - \mathbf{u}$;

- for every $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $\mathbf{u} \preceq \mathbf{v}$, $hT^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) := [\text{id}_X]_{(\varphi_X - \mathbf{u}, \varphi_X - \mathbf{v})}$.

Proof. For $\mathbf{u} \preceq \mathbf{v}$, $[\text{id}_X]_{(\varphi_X - \mathbf{u}, \varphi_X - \mathbf{v})}$ is a morphism between $\varphi_X - \mathbf{u}$ and $\varphi_X - \mathbf{v}$ because $(\varphi_X - \mathbf{v}) \circ \text{id}_X \preceq \varphi_X - \mathbf{u}$. Moreover, hT^{φ_X} preserves the identity and the composition. \square

Let us now consider ε -shifts of functors hT^{φ_X} .

Proposition 5.6. *For every $\varepsilon \succeq \mathbf{0}$, and for every φ_X , the ε -shift of hT^{φ_X} , $hT_\varepsilon^{\varphi_X} : \mathbf{R}^n \rightarrow \mathbf{hTop}/\mathbb{R}_{\succeq}^n$, is equal to the functor $hT^{\varphi_X - \varepsilon} : \mathbf{R}^n \rightarrow \mathbf{hTop}/\mathbb{R}_{\succeq}^n$.*

Proof. By Definition 5.1 and the definition of hT^{φ_X} (Proposition 5.5), the ε -shift of hT^{φ_X} is the functor $hT_\varepsilon^{\varphi_X} : \mathbf{R}^n \rightarrow \mathbf{hTop}/\mathbb{R}_{\succeq}^n$ such that, for every $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $\mathbf{u} \preceq \mathbf{v}$, (i) $hT_\varepsilon^{\varphi_X}(\mathbf{u}) = hT^{\varphi_X}(\mathbf{u} + \varepsilon) = \varphi_X - \mathbf{u} - \varepsilon = hT^{\varphi_X - \varepsilon}(\mathbf{u})$; (ii) $hT_\varepsilon^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = hT^{\varphi_X}(\mathbf{u} + \varepsilon \preceq \mathbf{v} + \varepsilon) = [\text{id}_X]_{(\varphi_X - \mathbf{u} - \varepsilon, \varphi_X - \mathbf{v} - \varepsilon)} = hT^{\varphi_X - \varepsilon}(\mathbf{u} \preceq \mathbf{v})$. \square

The next lemma describes the natural transformations between pairs of functors hT^{φ_X} and hT^{φ_Y} .

Lemma 5.2. *Let X, Y be topological spaces, and let $\varphi_X : X \rightarrow \mathbb{R}^n$, $\varphi_Y : Y \rightarrow \mathbb{R}^n$ be any two continuous functions. Then, every continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$ induces a natural transformation $h\xi^f : hT^{\varphi_X} \Rightarrow hT^{\varphi_Y}$ such that, for every $\mathbf{u} \in \mathbb{R}^n$, $h\xi^f(\mathbf{u})$ is equal to $[f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})}$. Reciprocally, for every natural transformation $h\xi : hT^{\varphi_X} \Rightarrow hT^{\varphi_Y}$, with $h\xi(\mathbf{u}) = [f_{\mathbf{u}}]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})}$, and for every $\alpha \succeq \mathbf{0}$ in \mathbb{R}^n , $f_{\mathbf{u}}$ and $f_{\mathbf{u} + \alpha}$ are α -homotopic with respect to (φ_X, φ_Y) for every $\mathbf{u} \in \mathbb{R}^n$.*

Proof. Any continuous map $f : X \rightarrow Y$ such that $\varphi_Y \circ f \preceq \varphi_X$ induces a natural transformation $h\xi^f : hT^{\varphi_X} \Rightarrow hT^{\varphi_Y}$ defined as follows: for every $\mathbf{u} \in \mathbb{R}^n$, $h\xi^f(\mathbf{u}) := [f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})}$. Indeed, for every $\mathbf{u} \in \mathbb{R}^n$, f is a morphism between $\varphi_X - \mathbf{u}$ and $\varphi_Y - \mathbf{u}$ because $\varphi_Y \circ f \preceq \varphi_X$ implies $(\varphi_Y - \mathbf{u}) \circ f \preceq \varphi_X - \mathbf{u}$, proving that $h\xi^f(\mathbf{u}) : hT^{\varphi_X}(\mathbf{u}) \rightarrow hT^{\varphi_Y}(\mathbf{u})$. Moreover, we get that $[f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})} \circ [\text{id}_X]_{(\varphi_X - \mathbf{u}, \varphi_X - \mathbf{u})} = [\text{id}_Y]_{(\varphi_Y - \mathbf{u}, \varphi_Y - \mathbf{u})} \circ [f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})}$, giving $h\xi^f(\mathbf{u}) \circ hT^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = hT^{\varphi_Y}(\mathbf{u} \preceq \mathbf{v}) \circ h\xi^f(\mathbf{v})$.

Reciprocally, assume there is a family $h\xi = \{h\xi(\mathbf{u}) : hT^{\varphi_X}(\mathbf{u}) \rightarrow hT^{\varphi_Y}(\mathbf{u})\}_{\mathbf{u} \in \mathbb{R}^n}$ of $\mathbf{0}$ -homotopy classes of maps $f_{\mathbf{u}}$ with respect to $(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})$ such that diagram (5.1) commutes for every $\mathbf{u} \preceq \mathbf{v} \in \mathbb{R}^n$: $h\xi(\mathbf{v}) \circ hT^{\varphi_X}(\mathbf{u} \preceq \mathbf{v}) = hT^{\varphi_Y}(\mathbf{u} \preceq \mathbf{v}) \circ h\xi(\mathbf{u})$. Take $\alpha = \mathbf{v} - \mathbf{u}$. From

$$h\xi(\mathbf{u} + \alpha) \circ hT^{\varphi_X}(\mathbf{u} \preceq \mathbf{u} + \alpha) = hT^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + \alpha) \circ h\xi(\mathbf{u})$$

we deduce $[f_{\mathbf{u} + \alpha}]_{(\varphi_X - \mathbf{u} - \alpha, \varphi_Y - \mathbf{u} - \alpha)} \circ [\text{id}_X]_{(\varphi_X - \mathbf{u}, \varphi_X - \mathbf{u} - \alpha)} = [\text{id}_Y]_{(\varphi_Y - \mathbf{u}, \varphi_Y - \mathbf{u} - \alpha)} \circ [f_{\mathbf{u}}]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u})}$.

Equivalently, $[f_{\mathbf{u} + \alpha}]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u} - \alpha)} = [f_{\mathbf{u}}]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u} - \alpha)}$. In other words, there exists a homotopy $H : X \times I \rightarrow Y$ such that $H(\cdot, 0) = f_{\mathbf{u}}$, $H(\cdot, 1) = f_{\mathbf{u} + \alpha}$, and $\varphi_Y \circ H(\cdot, t) - \mathbf{u} - \alpha \preceq \varphi_X - \mathbf{u}$ for every $t \in I$. Hence $\varphi_Y \circ H(\cdot, t) \preceq \varphi_X + \alpha$ for every $t \in I$, yielding that $f_{\mathbf{u}}$ and $f_{\mathbf{u} + \alpha}$ are α -homotopic with respect to (φ_X, φ_Y) . \square

We can now show that the interleaving distance between functors hT^{φ_X} and hT^{φ_Y} in $\mathbf{hTop}/\mathbb{R}_{\succeq}^n$ coincides with the persistent homotopy type pseudo-distance.

Proposition 5.7. *Let X and Y be topological spaces. Denoting by d_{HT} the persistent homotopy type pseudo-distance, for every pair of continuous functions $\varphi_X : X \rightarrow \mathbb{R}^n$ and $\varphi_Y : Y \rightarrow \mathbb{R}^n$, it holds that*

$$d_1(hT^{\varphi_X}, hT^{\varphi_Y}) = d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)).$$

Proof. If X and Y are not homotopy equivalent, $d_{\text{HT}} = \infty$ trivially implying that $d_1 \leq d_{\text{HT}}$. Let $f : X \rightarrow Y$, $g : Y \rightarrow X$ form a pair (f, g) of ε -homotopy equivalences with respect to (φ_X, φ_Y) for some $\varepsilon \succeq \mathbf{0}$. It holds that $(\varphi_Y - \varepsilon) \circ f \preceq \varphi_X$ and $(\varphi_X - \varepsilon) \circ g \preceq \varphi_Y$. By Lemma 5.2, f induces the natural transformation $h\xi^f : hT^{\varphi_X} \Rightarrow hT^{(\varphi_Y - \varepsilon)}$ with $h\xi^f(\mathbf{u}) = [f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \varepsilon - \mathbf{u})}$, and g induces the natural transformation $h\xi^g : hT^{\varphi_Y} \Rightarrow hT^{(\varphi_X - \varepsilon)}$ with $h\xi^g(\mathbf{u}) = [g]_{(\varphi_Y - \mathbf{u}, \varphi_X - \varepsilon - \mathbf{u})}$. By Proposition 5.6, $hT^{(\varphi_Y - \varepsilon)} = hT_\varepsilon^{\varphi_Y}$ and $hT^{(\varphi_X - \varepsilon)} = hT_\varepsilon^{\varphi_X}$. Thus, we get

$$\begin{aligned} h\xi^g(\mathbf{u} + \varepsilon) \circ h\xi^f(\mathbf{u}) &= [g]_{(\varphi_Y - \varepsilon - \mathbf{u}, \varphi_X - 2\varepsilon - \mathbf{u})} \circ [f]_{(\varphi_X - \mathbf{u}, \varphi_Y - \varepsilon - \mathbf{u})} \\ &= [g \circ f]_{(\varphi_X - \mathbf{u}, \varphi_X - 2\varepsilon - \mathbf{u})} = [g \circ f]_{(\varphi_X, \varphi_X - 2\varepsilon)} \\ &= [\text{id}_X]_{(\varphi_X, \varphi_X - 2\varepsilon)} = hT^{\varphi_X}(\mathbf{u} \preceq \mathbf{u} + 2\varepsilon). \end{aligned}$$

Similarly, $h\xi^f(\mathbf{u} + \varepsilon) \circ h\xi^g(\mathbf{u}) = hT^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + 2\varepsilon)$. Hence, the pair $(h\xi^f, h\xi^g)$ is an $\|\varepsilon\|_\infty$ -interleaving between hT^{φ_X} and hT^{φ_Y} , proving that $d_1 \leq d_{\text{HT}}$ also in this case.

Let us now show that $d_{\text{HT}} \leq d_1$. The claim is obvious if $d_1 = \infty$, so let us assume that hT^{φ_X} and hT^{φ_Y} are ε -interleaved for some $\varepsilon \geq 0$ by an ε -interleaving $(h\xi, h\zeta)$. By Proposition 5.6, for any fixed \mathbf{u} arbitrarily chosen, there is a continuous $\mathbf{0}$ -map $f_{\mathbf{u}} : X \rightarrow Y$ with respect to $(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u} - \vec{\varepsilon})$ such that $[f_{\mathbf{u}}]_{(\varphi_X - \mathbf{u}, \varphi_Y - \mathbf{u} - \vec{\varepsilon})} = h\xi(\mathbf{u})$, and a continuous $\mathbf{0}$ -map $g_{\mathbf{u} + \vec{\varepsilon}} : Y \rightarrow X$ with respect to $(\varphi_Y - \mathbf{u} - \vec{\varepsilon}, \varphi_X - \mathbf{u} - 2\vec{\varepsilon})$ such that $[g_{\mathbf{u} + \vec{\varepsilon}}]_{(\varphi_Y - \mathbf{u} - \vec{\varepsilon}, \varphi_X - \mathbf{u} - 2\vec{\varepsilon})} = h\zeta(\mathbf{u} + 2\vec{\varepsilon})$. Note that $f_{\mathbf{u}}$ and $g_{\mathbf{u} + \vec{\varepsilon}}$ are also $\vec{\varepsilon}$ -maps with respect to (φ_X, φ_Y) and (φ_Y, φ_X) , respectively. By Definition 5.2, for every $\mathbf{u} \in \mathbb{R}^n$, $h\zeta(\mathbf{u} + \vec{\varepsilon}) \circ h\xi(\mathbf{u}) = hT^{\varphi_X}(\mathbf{u} \leq \mathbf{u} + 2\vec{\varepsilon})$, and $h\xi(\mathbf{u} + 2\vec{\varepsilon}) \circ h\zeta(\mathbf{u} + \vec{\varepsilon}) = hT^{\varphi_Y}(\mathbf{u} + \vec{\varepsilon} \preceq \mathbf{u} + 3\vec{\varepsilon})$. Hence, from $h\zeta(\mathbf{u} + \vec{\varepsilon}) \circ h\xi(\mathbf{u}) = hT^{\varphi_X}(\mathbf{u} \leq \mathbf{u} + 2\vec{\varepsilon})$ we deduce that $g_{\mathbf{u} + \vec{\varepsilon}} \circ f_{\mathbf{u}}$ is $\mathbf{0}$ -homotopic to id_X with respect to $(\varphi_X - \mathbf{u}, \varphi_X - \mathbf{u} - 2\vec{\varepsilon})$, that is $g_{\mathbf{u} + \vec{\varepsilon}} \circ f_{\mathbf{u}}$ is $2\vec{\varepsilon}$ -homotopic to id_X with respect to (φ_X, φ_X) . Analogously, from $h\xi(\mathbf{u} + \vec{\varepsilon}) \circ h\zeta(\mathbf{u}) = hT^{\varphi_Y}(\mathbf{u} \preceq \mathbf{u} + 2\vec{\varepsilon})$ we deduce that $f_{\mathbf{u} + \vec{\varepsilon}} \circ g_{\mathbf{u}}$ is $\mathbf{0}$ -homotopic to id_Y with respect to $(\varphi_Y - \mathbf{u}, \varphi_Y - \mathbf{u} - 2\vec{\varepsilon})$, that is $f_{\mathbf{u} + \vec{\varepsilon}} \circ g_{\mathbf{u}}$ is $2\vec{\varepsilon}$ -homotopic to id_Y with respect to (φ_Y, φ_Y) . On the other hand, by Lemma 5.2, $g_{\mathbf{u}}$ is $\vec{\varepsilon}$ -homotopic to $g_{\mathbf{u} + \vec{\varepsilon}}$ with respect to (φ_Y, φ_X) , and $f_{\mathbf{u} + \vec{\varepsilon}}$ is $\vec{\varepsilon}$ -homotopic to $f_{\mathbf{u}}$ with respect to (φ_X, φ_Y) . Thus, $f_{\mathbf{u} + \vec{\varepsilon}} \circ g_{\mathbf{u}}$ is $2\vec{\varepsilon}$ -homotopic to $f_{\mathbf{u}} \circ g_{\mathbf{u} + \varepsilon}$ with respect to (φ_Y, φ_Y) , implying that $f_{\mathbf{u}} \circ g_{\mathbf{u} + \varepsilon}$ is $2\vec{\varepsilon}$ -homotopic to id_Y with respect to (φ_Y, φ_Y) . In conclusion, for an arbitrarily chosen $\mathbf{u} \in \mathbb{R}^n$, $(f_{\mathbf{u}}, g_{\mathbf{u} + \vec{\varepsilon}})$ is a pair of $\vec{\varepsilon}$ -homotopy equivalences with respect to (φ_X, φ_Y) , thus proving that $d_{\text{HT}}((X, \varphi_X), (Y, \varphi_Y)) \leq \varepsilon$. \square

5.1.3. The interleaving distance between persistence modules

Let $\mathbf{Vect}_{\mathbb{F}}$ be the category of vector spaces over a fixed field \mathbb{F} . An n -dimensional persistence module can be viewed as a functor $\mathfrak{P} : \mathbf{R}^n \rightarrow \mathbf{Vect}_{\mathbb{F}}$. Replacing \mathbf{O} with $\mathbf{Vect}_{\mathbb{F}}$ in Definition 5.2, we obtain the interleaving distance on the category $\mathbf{Vect}_{\mathbb{F}}$ as usual.

A standard way in which one obtains n -dimensional persistence modules from topological spaces endowed with \mathbb{R}^n valued functions, is by composing the sublevelset

filtration functor with the homology functor: for k a non-negative integer, and for (X, φ_X) an object in $\mathbf{Top}/\mathbb{R}_{\leq}^n$, $\mathfrak{P}_k^X = H_k \circ T^{\varphi_X}$ is an object in $\mathbf{Vect}_{\mathbb{F}}^{\mathbf{R}^n}$.

An immediate remark is that given $k \in \mathbb{N}$ and any ε -interleaving (ζ, η) between objects (X, φ_X) and (Y, φ_Y) in $\mathbf{Top}/\mathbb{R}_{\leq}^n$, the pair $(H_k(\zeta), H_k(\eta))$ is an ε -interleaving between the n -dimensional persistence modules \mathfrak{P}_k^X and \mathfrak{P}_k^Y .

As a corollary of Propositions 5.7 and 5.4 we then obtain the following result linking the interleaving distances on the categories $\mathbf{Vect}_{\mathbb{F}}^{\mathbf{R}^n}$, $\mathbf{Top}/\mathbb{R}_{\leq}^n$, and $\mathbf{hTop}/\mathbb{R}_{\leq}^n$.

Corollary 5.1. *For every $(X, \varphi_X), (Y, \varphi_Y)$ with X, Y topological spaces, every $\varphi_X : X \rightarrow \mathbb{R}^n$, $\varphi_Y : Y \rightarrow \mathbb{R}^n$ continuous functions, and every non-negative integer k , it holds that*

$$d_I(\mathfrak{P}_k^X, \mathfrak{P}_k^Y) \leq d_I(hT^{\varphi_X}, hT^{\varphi_Y}) \leq d_I(T^{\varphi_X}, T^{\varphi_Y}).$$

Remark 5.1. Compare the leftmost inequality above (for $n = 1$) with Lemma 3.1.

6. Discussion

We have introduced the persistent homotopy type distance d_{HT} to quantify perturbations of functions defined on homotopy equivalent spaces. As a key consequence, we were able to lift the standard stability result of persistence diagrams of sublevel set filtrations of Cohen-Steiner, Edelsbrunner, and Harer [11] from the setting of functions defined on the *same* domain to the more general setting of functions defined on possibly different but homotopy equivalent domains.

Focusing, as we did, on sublevelset filtrations implies a certain asymmetry in the definition of persistent homotopy type distance in that only up-shifts of maps impact it, but not down-shifts, see Proposition 2.4 and Example 2.6.

This lack of symmetry is shared neither by the L^∞ distance nor the natural pseudo-distance. One interesting line of research is whether a certain modification of our definition of the persistent homotopy type distance would correct this asymmetry. Further, it would be interesting to investigate whether such modification implies a lift of the standard L^∞ stability results for extended persistence [12] and/or interlevel set persistence [1] to settings when functions are defined on possibly different homotopy equivalent domains.

Another interesting open line of research is uncovering the relationship, if any, between our homotopy type distance and the distances that appear in the recent work of Blumberg and Lesnick [3].

7. Acknowledgements

This work started during a visit of the third author to the first author at the University of Bologna in 2014 which was partially supported by INdAM-GNSAGA. The first two authors partially carried out this research within the activities of ARCES ‘‘E. De Castro’’, University of Bologna. The third author has been partially supported by NSF under grants DMS-1547357, CCF-1526513, and IIS-1422400. The authors thank Francesca Cagliari for her helpful advice about the categorical setting and Michael Lesnick for useful discussions that helped us to focus Section 5.

References

- [1] P. Bendich, H. Edelsbrunner, D. Morozov, and A. Patel. *Homology and robustness of level and interlevel sets*. Homology, Homotopy and Applications vol. 15, (2013) no. 1, 51–72.
- [2] S. Biasotti, L. De Floriani, B. Falcidieno, P. Frosini, D. Giorgi, C. Landi, L. Papaleo, M. Spagnuolo, *Describing shapes by geometrical-topological properties of real functions*, ACM Computing Surveys, vol. 40 (2008), n. 4, 12:1–12:87.
- [3] A. Blumberg and M. Lesnick. *Universality of the Homotopy Interleaving Distance*. ArXiv 2017, <https://arxiv.org/abs/1705.01690>.
- [4] P. Bubenik and J. A. Scott, *Categorification of persistent homology*, Discrete & Computational Geometry, vol. 51 (2014), n. 3, 600–627.
- [5] P. Bubenik, V. de Silva, and J. Scott. *Metrics for generalized persistence modules*. Foundations of Computational Mathematics vol. 15 (2015) n. 6, 1501–1531.
- [6] F. Cagliari, B. Di Fabio, C. Landi, *The natural pseudo-distance as a quotient pseudo-metric, and applications*, Forum Mathematicum, 27 (2013), n. 3, 1729–1742.
- [7] G. Carlsson, *Topology and data*, Bull. Amer. Math. Soc. (N.S.), vol. 46 (2009), n. 2, 255–308.
- [8] A. Cerri, B. Di Fabio, M. Ferri, P. Frosini, C. Landi, *Betti numbers in multi-dimensional persistent homology are stable functions*, Mathematical Methods in the Applied Sciences, vol. 36 (2013), 1543–1557.
- [9] F. Chazal, D. Cohen-Steiner, M. Glisse, L. Guibas, S. Oudot, *Proximity of persistence modules and their diagrams*, Proceedings of the twenty-fifth ACM Annual Symposium on Computational Geometry 2009, 237–246.
- [10] F. Chazal, V. de Silva, M. Glisse, S. Oudot, *The structure and stability of persistence modules*, SpringerBriefs in Mathematics, 2016.
- [11] D. Cohen-Steiner, David, H. Edelsbrunner, J. Harer, *Stability of persistence diagrams*, Discrete & Computational Geometry, vol. 37 (2007), n. 1, 103–120.
- [12] D. Cohen-Steiner, H. Edelsbrunner, and J. Harer. *Extending persistence using Poincaré and Lefschetz duality*. Foundations of Computational Mathematics vol. 9 (2009), n. 1, 79–103.
- [13] B. Di Fabio, C. Landi, *Reeb graphs of piecewise linear functions*, Proc. GbRPR 2017, LNCS 10310 (2017).
- [14] P. Donatini, P. Frosini, *Natural pseudodistances between closed manifolds*, Forum Mathematicum, vol. 16 (2004), n. 5, 695–715.
- [15] H. Edelsbrunner, D. Morozov, *Persistent homology: theory and practice*, Proceedings of the European Congress of Mathematics 2012, Eur. Math. Soc., Zürich, 31–50 (2014)
- [16] R. Fritsch, R. A. Piccinini, *Cellular Structures in Topology*, Cambridge University Press, 1990.
- [17] P. Frosini, *G-invariant persistent homology*, Math. Methods Appl. Sci., vol. 38 (2015), n. 6, 1190–1199.

- [18] P. Frosini, G. Jabłoński, *Combining persistent homology and invariance groups for shape comparison*, Discrete & Computational Geometry, vol. 5 (2016), n. 2, 373–409.
- [19] R. Ghrist, *Barcodes: the persistent topology of data*, Bull. Amer. Math. Soc. (N.S.), vol. 45 (2008), n. 1, 61–75.
- [20] M. W. Hirsch, *Differential Topology*, Springer New York, 1976.
- [21] M. Lesnick, *Multidimensional interleavings and applications to topological inference*, PhD. thesis, Stanford, 2012. <https://searchworks.stanford.edu/view/9697564>.
- [22] M. Lesnick, *The theory of the interleaving distance on multidimensional persistence modules*, Found. Comput. Math., vol. 15 (2015), n.3, 613–650.
- [23] D. Morozov, K. Beketayev, and G. H. Weber, *Interleaving distance between merge trees*, manuscript presented at TopoInVis 2013.
- [24] G. Reeb, *Sur les points singuliers d’une forme de Pfaff complètement intégrable ou d’une fonction numérique*, Comptes Rendus de L’Académie des Sciences 222 (1946), 847–849.
- [25] H. Seifert, W. Threlfall, *A textbook of topology*, Academic Press, 1980.

Patrizio Frosini `patrizio.frosini@unibo.it`

Patrizio Frosini, Department of Mathematics, University of Bologna, Italy

Claudia Landi `claudia.landi@unimore.it`

Dipartimento di Scienze e Metodi dell’Ingegneria, Università di Modena e Reggio Emilia Reggio Emilia, Italy

Facundo Mémoli `memoli@math.osu.edu`

Department of Mathematics, The Ohio State University, Columbus, Ohio, U.S.A.