

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Polyphenol-enriched spelt husk extracts improve growth and stress-related biochemical parameters under moderate salt stress in maize plants

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Ceccarini C., Antognoni F., Biondi S., Fraternale A., Verardo G., Gorassini A., et al. (2019). Polyphenolenriched spelt husk extracts improve growth and stress-related biochemical parameters under moderate salt stress in maize plants. PLANT PHYSIOLOGY AND BIOCHEMISTRY, 141, 95-104 [10.1016/j.plaphy.2019.05.016].

Availability:

This version is available at: https://hdl.handle.net/11585/691670 since: 2019-07-15

Published:

DOI: http://doi.org/10.1016/j.plaphy.2019.05.016

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

Accepted Manuscript

Polyphenol-enriched spelt husk extracts improve growth and stress-related biochemical parameters under moderate salt stress in maize plants

Ceccarini Chiara, Fabiana Antognoni, Stefania Biondi, Alessandra Fraternale, Giancarlo Verardo, Andrea Gorassini, Valeria Scoccianti

PII: S0981-9428(19)30204-9

DOI: https://doi.org/10.1016/j.plaphy.2019.05.016

Reference: PLAPHY 5710

To appear in: Plant Physiology and Biochemistry

Received Date: 20 February 2019

Revised Date: 26 April 2019

Accepted Date: 15 May 2019

Please cite this article as: C. Chiara, F. Antognoni, S. Biondi, A. Fraternale, G. Verardo, A. Gorassini, V. Scoccianti, Polyphenol-enriched spelt husk extracts improve growth and stress-related biochemical parameters under moderate salt stress in maize plants, *Plant Physiology et Biochemistry* (2019), doi: https://doi.org/10.1016/j.plaphy.2019.05.016.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.







Polyphenol-enriched spelt husk extracts improve growth and stress-related biochemical parameters under moderate salt stress in maize plants

Ceccarini Chiara¹, Fabiana Antognoni², Stefania Biondi³, Alessandra Fraternale¹, Giancarlo Verardo⁴, Andrea Gorassini⁵, Valeria Scoccianti¹

¹ Dipartimento di Scienze Biomolecolari, Università di Urbino Carlo Bo, 61029 Urbino, Italy

² Dipartimento di Scienze per la Qualità della Vita, Università di Bologna, 47921 Rimini, Italy

³Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università di Bologna, 40126 Bologna, Italy

⁴Dipartimento di Scienze Agroalimentari, Ambientali e Animali, Università di Udine, 33100 Udine, Italy

⁵Dipartimento di Studi Umanistici e del Patrimonio Culturale, Università di Udine, 33100 Udine, Italy

*Correspondence:

Prof. Fabiana Antognoni

Department for Life Quality Studies, Corso Augusto 237, University of Bologna, 47921, Rimini, Italy

e-mail: fabiana.antognoni@unibo.it

1

Abstract

Biostimulants improve yield, quality, and stress acclimation in crops. In this work, we tested the possibility of using phenolics-rich extracts from spelt (Triticum dicoccum L.) husks to attenuate the effects of salt stress (100-200 mM NaCl) in maize. Two methanolic extracts were prepared from the soluble-conjugated (SC), and the insoluble-bound (IB) phenolic acid fractions of the spelt husk, and their effects were investigated on several stress-associated biochemical parameters, such as proline, lipid peroxidation, H₂O₂, GSH levels, and ion content. Results show that SC and IB fractions of husk extracts behaved very differently, no doubt due to their greatly divergent chemical composition, as revealed by both GC-MS and HPLC analyses. The efficacy of treatments in mitigating salt stress was also dose- and timing-dependent. IB, even at the lower concentration tested, was able to recover the performance of stressed plants in terms of growth, photosynthetic pigments content, and levels of salt stress markers. Recovery of shoot growth to control levels and reduction of stress-induced proline accumulation occurred regardless of whether plants were pretreated or post-treated with IB, whereas only pre-treatment with the higher dose of IB was effective in mitigating oxidative stress. Although in some cases SC and even methanol alone exerted some positive effects, they could also be deleterious whereas IB never was. Overall, results indicate that a polyphenol-containing extract obtained from spelt by-products can behave as biostimulant in maize plants and can mitigate their response to salt stress, by acting on different biochemical targets.

Keywords: agricultural by-products, biostimulant, oxidative stress, phenolic acids, salt stress, *Triticum dicoccum*, *Zea mays*

Introduction

Salinity is one of the major abiotic stresses that negatively influence plant growth, biomass production, and crop yield. More than 800 million hectares worldwide are currently affected by salt and they are expected to increase even further because of global climate change (Mahajan and Tuteja, 2005). High concentrations of salt reduce water and nutrient uptake (Ashraf 2004), as well as chlorophyll content, and RuBisCO activity (Rady 2013; Kahrizi et al. 2012), leading to impaired plant growth and productivity. Salinity also imposes osmotic stress and ionic toxicity, and leads to oxidative stress. Consequently, antioxidant defense mechanisms play a fundamental role in mitigating salt stress (Zhu 2001; Sairam and Tyagi 2004).

The application of biostimulants has been proposed as an innovative agronomic practice, not only to ensure optimal nutrient uptake, crop yield, and quality (Chen et al. 2003; Schiavon et al, 2008, 2010; Ertani et al. 2009; Khan et al. 2009), but also to mitigate the detrimental effects of environmental stresses, including high salinity (Feitosa de Vasconcelos et al, 2009; Povero et al. 2016; Desoky et al. 2018). Due to the heterogeneous chemical composition of biostimulants, their biological activity is also highly variable (Ertani et al, 2013, 2016; du Jardin 2015). An interesting source of biostimulants is represented by agro-industrial residues, such as bark, straw, rice husks, etc. (Ertani et al., 2011a), which often still contain bioactive molecules (Schieber et al. 2001; Balasundram et al. 2005), able to modulate several physiological processes in plants. Phenolics are amongst these compounds. They represent a very large group of secondary metabolites, which are involved in many aspects of plant growth, structural support, and response to environmental stimuli (Caretto et al. 2015 and references therein); they also play a crucial role in response to variation in mineral supply (Clemens and Weber, 2016). The role of phenolic compounds in mediating stress responses is mainly due to their capacity to scavenge free radicals produced in stress-induced oxidative reactions (Cook and Samman 1996; Halliwell 2008; Sharma et al. 2012), thus protecting cell components from damage (Bulgari et al. 2017). In recent years, several studies have investigated the application in agriculture of food by-products enriched in phenols, since these secondary metabolites are particularly efficient in stimulating plant growth and in improving plantwater relationships (Ertani et al. 2016). Moreover, phenolics may exert a gibberellin-like activity (Savy et al. 2017) and interact with several other plant hormones, including auxins (Ertani et al. 2016).

Maize (*Zea mays* L.) is one of the world's most important cereal crops, classified as moderately sensitive to salinity, with wide intraspecific genetic variations in salt tolerance (Farooq et al., 2015). The salt tolerance of some maize genotypes has been explained by the maintenance and/or increased activity of antioxidant enzymes (De Azevedo Neto et al. 2006). The effects of biostimulants on growth and stress tolerance have also been investigated in maize. Maize seedlings treated with plant extracts derived from red grape, blueberry fruits, and hawthorn leaves showed increased root and leaf biomass, chlorophyll and sugar content, as well as phenolic acids compared to untreated plants (Ertani et al. 2016). Recently, a positive effect against drought stress was also observed in maize treated with an aqueous extract of sorghum leaves (Maqbool and Sadiq 2017).

In this work, an extract was prepared using a by-product of spelt (*Triticum dicoccum* L.) processing. Based on the assumption that this by-product may still contain bioactive polyphenols, the potential of this material as plant biostimulant under stress conditions was investigated. To this purpose, a pot trial was conducted to test the performance of methanolic extracts prepared from spelt husks in

mitigating the negative impact of salt stress in maize. Since methanol was used as solvent and previous reports demonstrate a role of this alcohol in modulating plant growth, development (Dorokhov et al. 2018) and the response to drought and salt stresses (Mirakori et al. 2009; Wei et al. 2015), the effect of methanol alone was investigated in parallel.

Materials and methods

Spelt husk extract preparation

Two phenolic acid-enriched extracts were prepared from husks of *Triticum dicoccum* L. (kindly provided by Terra Bio Soc. Coop, Schieti di Urbino, PU, Italy) according to the protocol described in Antognoni et al. (2017). Husk samples (1.5 g) were extracted with 30 mL acetone/methanol/water mixture (7:7:6, v/v/v) in an ultrasound bath at 30 °C for 30 min. The homogenate was centrifuged at 1,500 x g for 20 min and the pellet was re-extracted once with the same procedure. Pooled supernatants and pellets were used to prepare the soluble-conjugated (SC) and the insoluble-bound (IB) phenolic acid fractions, respectively. An aliquot (8 mL) of supernatant was mixed with 2 mL 10 M NaOH and hydrolyzed under nitrogen flow and constant stirring for 1 h. The solution was then acidified to pH 2 with 12 M HCl and extracted three times with an equal volume of a diethyl ether/ethyl acetate (1/1, v/v) mixture. The organic extracts were merged, brought to dryness in a rotary evaporator, and re-dissolved with 2 mL of methanol. A 0.5 g aliquot of the pellet was mixed with 40 mL of 2 M NaOH and hydrolyzed under nitrogen flow and constant stirring for 1 h. The sample was then centrifuged for 20 min at 1.500 x g; the supernatant was acidified to pH 2 and then subjected to the extraction procedure already described for the SC fraction.

Gas chromatography-mass spectrometry (GC-MS) analysis of extracts

An aliquot (*ca.* 1 mg) of each dry extract was derivatized with 200 μ L of bis-(trimethyl) trifluoroacetamide (BSTFA) including 1% trimethylchlorosilane (TMCS) (Sigma-Aldrich, Milan, Italy) and 200 μ L of pyridine (Sigma-Aldrich, Milan, Italy) for 2 h at room temperature (RT). The silylating reagent was removed under a nitrogen stream and the sample re-dissolved in 1 mL of *n*-heptane. The GC-MS analysis of the trimethylsilyl derivatives was carried out using a Trace GC Ultra coupled to an ion-trap mass spectrometer detector Polaris Q and equipped with a split/splitless injector (Thermo Scientific, Italy). The column was a Supelco SLB-5ms, 30 x 0.25 mm, 0.1 μ m film thickness (Supelco, Sigma-Aldrich, Milan, Italy). Column oven temperature was programmed as follows: 80 °C held for 2 min, up to 280 °C at 10 °C min⁻¹, held for 10 min, up to 300 °C at 10 °C min⁻¹. The

injector, transfer line, and ion source were held at 280, 280 and 200 °C, respectively. Split ratio was 1:10 and injection volume 1 μ L. The mass spectra were recorded under electron ionization (EI) conditions at 70 eV electron energy with a mass range from m/z 50 to 650. Relative percentage amount of each identified compound was expressed as percent peak area relative to total ion current GC-MS peak area without using the correction factor.

HPLC analysis of extracts

Twenty μ L of each extract were injected into an HPLC system (Jasco, Tokyo, Japan; PU-4180 pump, MD-4015 PDA detector, AS-4050 autosampler). The stationary phase was an Agilent (Santa Clara, CA, USA) Zorbax Eclipse Plus C18 reversed-phase column (100 mm x 3 mm I.D., particle size 3.5 μ m).The chromatographic method for the analysis of phenolic acids was adapted from Mattila et al. (2005). Gradient elution was carried out with a mixture of acidic phosphate buffer and acetonitrile flowing at 0.7 mL min⁻¹. The signals at 254, 280, and 329 nm were used for analyte quantitation. Identification and quantification were performed based on standard compounds (gallic, *p*-hydroxybenzoic, syringic, ferulic, *p*-coumaric, cinnamic, and caffeic acids). The recovery values in spiked samples ranged from 78.8 to 92.2% (RSD < 9.8%, n = 6).

The sum of all individual phenolic acid concentrations was calculated and used to express the total phenolics acid index (TPAI) for each extract.

Plant material

Seeds of *Zea mays* L. (var. FAO 700, kindly provided by Società Italiana Sementi, San Lazzaro di Savena, Bologna, Italy) were washed under running tap water for 30 min and then placed on wet filter paper in Petri dishes. They were germinated for two days in the dark at 25 °C. Seedlings were then transferred to plastic pots (12 x 8 x 7 cm) with drainage holes and containing Perlite soaked in Hoagland's nutrient solution (Hoagland and Arnon 1950). The pots were placed on trays to which half-strength Hoagland's solution was added when necessary in order to keep the Perlite fully wet. Plants were grown in a growth chamber with a photoperiod of 16/8 h day/night (400 µmol m⁻² s⁻¹ flux density supplied by fluorescent TL/D Aquarelle Philips lamps) at a temperature of 24 ± 2 °C. During the day, an additional irradiation (180 m⁻² s⁻¹ flux density from Lumatek HPS Grow lamps) was supplied for 5 h.

Salt treatment and husk extract application

After 8-10 days, the pots (four per treatment and each containing five plants) were transferred to trays containing Hoagland's solution added with either 0 (control), 50, 100, or 200 mM NaCl. The

solution in the trays was changed every two days to maintain the same salt concentration. Plants were harvested at 5, 12, and 19 days after the start of salt treatment.

Husk extracts (HEs) were added to the nutrient solution at two different concentrations, 0.1 or 1.0

mL L⁻¹, in the trays before (pre-treatment) or after (post-treatment) the addition of NaCl as follows:

- a) pre-treatment (PRE): IB, SC or methanol (solvent control) were added for two days before the beginning of salt treatment. Plants were then transferred to trays containing Hoagland's solution added with 0, 100 or 200 mM NaCl.
- b) post-treatment (POST): IB, SC or methanol (solvent control) were added two days after the start of salt treatment; plants were grown in the presence of extract or methanol plus NaCl until sampling.

Plants were harvested at 12 (100 mM NaCl) or 8 days (200 mM NaCl) from the start of salt treatment. Shoots and roots were weighed separately and then ground to a powder in liquid nitrogen; some samples were freeze-dried, while others were left frozen and kept at -80 °C until use. For dry weight (DW) determination, shoots and roots were dried in an oven at 80 °C for two days.

Photosynthetic pigment determination

Freeze-dried shoot samples (50 mg) were extracted in a chilled mortar with 80% (v/v) cold acetone (1:5 ratio) in dim light. The homogenate was centrifuged at 10,000 x g for 10 min at 4 °C and the extraction repeated once. The supernatants were pooled and absorbance determined spectrophotometrically (V-530 Jasco, Jasco Corporation, Tokyo, Japan) at 663 nm (chlorophyll *a*), 647 nm (chlorophyll *b*), and 470 nm (carotenoids and xanthophylls). Pigment concentrations were estimated based on specific absorbance coefficients (Lichtenthaler 1987) and are expressed as mg g⁻¹ DW.

Lipid peroxidation assay

The level of lipid peroxidation was measured in terms of malondialdehyde (MDA) production by the thiobarbituric acid (TBA) reaction method (Velikova et al. 2000). Absorbance of the supernatant was read at 532 nm. After subtracting the value for non-specific absorption at 600 nm, the concentration of the MDA-TBA complex was calculated from the extinction coefficient 155 $\text{mM}^{-1} \text{ cm}^{-1}$.

Hydrogen peroxide determination

Hydrogen peroxide (H_2O_2) levels were determined according to Velikova et al. (2000). Frozen shoot and root tissue (500 mg) was extracted on an ice-bath with 5 mL 0.1% (w/v) trichloroacetic

acid. The homogenate was centrifuged at 12,000 x g for 15 min. Then, an aliquot (0.5 mL) of the supernatant was added to a mixture of 0.5 mL potassium phosphate buffer (10 mM, pH 7.0) and 1 mL 1 M KI. After 10 min at RT, the absorbance of the samples was spectrophotometrically determined at 390 nm. The concentration of H_2O_2 was calculated from a standard curve.

Proline determination

Shoot and root proline concentrations were estimated following the method of Bates et al. (1973) with slight modifications. About 20 mg of freeze-dried plant material was crushed in 1.2 mL 3% sulphosalicylic acid and the homogenate centrifuged at 16,000 x g at RT for 20 min. A 0.5 mL aliquot of the supernatant was made up to 1 mL with water and to this 1 mL glacial acetic acid and 1 mL ninhydrin reagent [2.5% ninhydrin in glacial acetic acid-distilled water-85% orthophosphoric acid (6:3:1)] were added. The reaction mixtures were kept in a water bath at 90 °C for 1 h to develop the colour. Test tubes were then cooled in an ice-bath, and 3 mL toluene added to separate the chromophore. Absorbance of the toluene phase was read in a spectrophotometer at 546 nm, and proline concentration calculated by comparing sample absorbance with the standard proline curve.

Glutathione (GSH) measurement

GSH was determined as previously described by Brundu et al. (2016). Approximately 20-25 mg of freeze-dried shoots or roots were transferred to microcentrifuge tubes containing 200 μ L of precipitating solution (1.67 g glacial meta-phosphoric acid, 0.2 g disodium EDTA, and 30 g NaCl in 100 mL). Samples were homogenized on ice with a mortar and pestle, kept on ice for 10 min and then centrifuged at 12,000 × g for 10 min at 4 °C. Fifteen μ L 0.3 M Na₂HPO₄ were added to 60 μ L of extract followed immediately by 45 μ L 5,5'-dithiobis (2-nitrobenzoic acid) (DTNB) prepared as follows: 20 mg DTNB in 100 mL 1% (w/v) sodium citrate solution. The mixture was stirred for 1 min at RT, then left at RT for another 5 min and finally used for GSH determination by Reverse Phase-HPLC (Jasco LG-980-02, Jasco Europe S.R.L., Cremella, LC, Italy).

Na^+ and K^+ content determination

Freeze-dried samples were mineralized (MDS 2000, CEM, Italy) with a mixture of concentrated HNO_3/H_2O_2 (8+2 mL). All chemicals used in sample treatment were suprapure grade (HNO₃ 65%, H_2O_2 30%, Merck Suprapur, Darmstadt, Germany). Na⁺ and K⁺ concentrations were measured by inductively coupled plasma-optical emission spectrometry using a Perkin Elmer Optima 8000 ICP-OES Spectrometer (PerkinElmer Inc., Shelton, CT, USA). Ultrapure water (Milli-Q system, Millipore Corporation, USA) was used for all solutions. Standard solutions were prepared from stock solutions of ultrapure grade metals (PerkinElmer Pure Plus, 100 mg L⁻¹). The error of metal

determinations, based on variation in replicate analyses (n=2) on the same samples, was 10% or lower.

Results

Phytochemical characterization of HEs

The phytochemical composition of both IB and SC was investigated through HPLC-DAD and GC-MS analyses. As shown in Table 1, five phenolic acids were detected by HPLC-DAD in both extracts, i.e., two simple phenolics (*p*-hydroxybenzoic, and syringic acids) and three hydroxycinnamic acids (ferulic, *p*-coumaric, and caffeic). The most abundant were ferulic and *p*-coumaric acids, which together represented *ca*. 75% and 97% of total phenolic acids in SC and IB, respectively. In absolute terms, IB contained about 25-fold higher ferulic and *p*-coumaric acids compared to SC and more than six-fold higher caffeic acid; syringic acid was about two-fold higher in SC than IB (Table 1). In both extracts, *p*-hydroxybenzoic acid was detected at a similar concentration (Table 1). Total phenolic acid index was 17.71 μ g mL⁻¹ for SC and 713.82 μ g mL⁻¹ for IB (Table 1).

8

Phenolic acid	S	SC	IB			
	$\mu g m L^{-1}$	μg g DW ⁻¹	μg mL ⁻¹	μg g DW ⁻¹		
<i>p</i> -OH benzoic acid	1.72 ± 0.7 16.15 ± 6.6		2.77 ± 0.4	13.23 ±1.9		
Syringic acid	1.63 ± 0.7	15.36 ± 6.6	1.84 ± 0.3	8.79 ± 1.4		
<i>p</i> -Coumaric acid	7.22 ± 0.9	67.74 ± 8.4	343.37 ±12.1	1635.10 ± 57.6		
Caffeic acid	0.38 ± 0.05	3.61 ± 0.5	5.11 ± 0.2	24.34 ± 0.9		
Ferulic acid	6.76 ± 0.4	63.45 ± 3.8	360.73 ± 19.4	1717.76 ± 92.4		
Total phenolic acid index	17.71 ±1.33	166.31 ± 13.1	713.82±12.11	3399.22 ±108.9		

Table 1. HPLC-DAD analysis of phenolic acid composition of spelt husk extracts. Data are the means \pm S.E. of three independent extractions.

In addition, GC-MS analysis of IB revealed the presence of vanillin (0.7%), vanillic acid (1.6%) and a relatively higher percentage of *m*-coumaric (10.5%) and isoferulic acids (8.4%) (Table 2). GC-MS analysis of SC showed the presence of malic (14.2%), palmitic (27.6%), linoleic (16.8%), oleic (25.6%), and stearic (3.2%) acids accompanied by lesser amounts of 2-hydroxyglutaric acid (0.2%), *m*-anisic acid methyl ester (0.7%), campesterol (0.2%), stigmasterol (0.3%), and β -sitosterol (0.4%) (Table 2). Fatty acids, absent in IB, represented about 73% of all compounds identified in SC. Among phenolic acids, ferulic and *p*-coumaric acids were the major constituents in both extracts representing *ca*. 75% and 98% in SC and IB, respectively, in agreement with the results of the HPLC-DAD analysis. Vanillin and isoferulic acid were absent in SC.

	S	С	IB		
Compound	Mean (%) ^a	RSD ^b (%)	Mean (%) ^a	RSD ^b (%)	

Malic acid	14.2 ± 0.4	2.8	-	-
Vanillin	-	-	0.7 ± 0.01	3.2
2-Hydroxyglutaric acid	0.2 ± 0.01	2.0	-	-
<i>p</i> -Hydroxybenzoic acid	0.5 ± 0.01	3.6	0.2 ± 0.01	3.8
<i>m</i> -Anisic acid methyl ester	0.7 ± 0.01	1.2	-	-
Vanillic acid	1.8 ± 0.01	1.8	1.6 ± 0.1	4.4
<i>m</i> -Coumaric acid	0.7 ± 0.01	0.4	10.5 ± 0.1	0.7
Syringic acid	1.4 ± 0.1	4.0	0.4 ± 0.01	6.4
Isoferulic acid	-	-	8.4 ± 0.2	2.0
<i>p</i> -Coumaric acid	3.3 ± 0.1	1.7	35.2 ± 0.4	1.0
Palmitic acid	27.6 ± 1.2	4.2		-
Ferulic acid	3.0 ± 0.01	1.0	42.1 ± 0.1	0.3
Caffeic acid	0.1 ± 0.01	4.6	0.9 ± 0.1	5.9
Linoleic acid	16.8 ± 1.1	6.5	-	-
Oleic acid	25.6 ± 0.4	1.5	-	-
Stearic acid	3.2 ± 0.1	1.9	-	-
Campesterol	0.2 ± 0.01	1.6	-	-
Stigmasterol	0.3 ± 0.01	2.1	-	-
β-Sitosterol	0.4 ± 0.01	1.3	-	-

Table 2. GC-MS analysis of spelt husk extracts.

^aPercent peak area relative to total ion current GC-MS peak area \pm SD (n = 3).

^bRelative standard deviation.

Effects of salt treatment on maize plants

As revealed by the time-course analysis of growth performed with NaCl alone, shoot fresh weight (FW) and dry weight (DW) were negatively affected by 50, 100 and 200 mM NaCl starting on day 12, with the strongest inhibition in FW (58% relative to controls) at the highest concentration and longest exposure time; instead, root FW was less affected and DW not at all by all three salt concentrations (Supplementary Table 1).

		Shoots				Roots				
Time (days)	NaCl (mM)	FW (g plant ⁻¹)	DW (g plant ⁻¹)	RFW	RDW	FW (g plant ⁻¹)	DW (g plant ⁻¹)	RFW	RDW	
	0	0.689 ± 0.208^{a}	0.055 ± 0.017^{a}	100	100	0.572 ± 0.154^{a}	0.03 ± 0.017^{a}	100	100	
5	50	0.945±0.227 ^a	0.072 ± 0.017^{a}	137.2	130.9	0.655 ± 0.029^{a}	0.038 ± 0.017^{a}	114.5	115.2	
5	100	0.905 ± 0.216^{a}	0.076 ± 0.016^{a}	131.3	138.2	0.709 ± 0.089^{a}	0.036 ± 0.016^{a}	124.0	109.1	
	200	0.907 ± 0.282^{a}	0.083 ± 0.026^{a}	131.6	150.9	0.628 ± 0.178^{a}	0.044 ± 0.026^{a}	109.8	133.3	
	0	2.620 ± 0.263^{a}	0.206 ± 0.017^{a}	100	100	1.338 ± 0.252^{a}	0.077 ± 0.014^{a}	100	100	
12	50	1.968 ± 0.078^{b}	0.160 ± 0.022^{b}	75.1	77.7	1.144 ± 0.131^{b}	0.061 ± 0.016^{a}	85.5	79.2	
12	100	1.923 ± 0.220^{b}	0.158 ± 0.034^{b}	73.4	76.7	1.100 ± 0.137^{b}	$0.079{\pm}0.014^{a}$	82.2	102.3	
	200	1.229 ± 0.183^{c}	0.137±0.022 ^b	46.9	66.5	$0.823 \pm 0.126^{\circ}$	0.065 ± 0.006^{a}	61.5	84.4	
	0	4.350±0.39 ^a	0.350 ± 0.03^{a}	100	100	1.596 ± 0.25^{a}	0.100 ± 0.02^{a}	100	100	
10	50	4.220 ± 1.10^{a}	0.366 ± 0.06^{a}	97.0	105.7	1.560 ± 0.34^{a}	0.099 ± 0.05^{a}	97.7	99.0	
17	100	2.910 ± 0.55^{b}	0.281±0.06 ^{ab}	66.9	80.0	1.400 ± 0.30^{ab}	0.097 ± 0.02^{a}	87.7	97.0	
	200	2.820 ± 0.45^{b}	0.221 ± 0.05^{b}	41.8	62.9	1.130 ± 0.07^{b}	0.090 ± 0.01^{a}	70.8	90.0	

Supplementary Table 1. Fresh weight, dry weight, relative fresh weight (RFW), and relative dry weight (RDW) of maize plants after exposure for 5, 12, and 19 days to 0, 50, 100 or 200 mM NaCl. Data are means \pm S.D. (n=8). Different letters within a column and for each time point indicate significant differences (p<0.05).

As regards the stress-related biochemical parameters, all photosynthetic pigments showed a timeand dose-dependent reduction in the presence of NaCl (Supplementary Fig. 1).



Supplementary Fig. 1. Concentration of photosynthetic pigments after 5 (A), 12 (B), and 19 (C) days of exposure to 0, 50, 100 or 200 mM NaCl. Data are means \pm S.E. (n=3). Asterisks indicate significant differences (*P*<0.05) relative to controls (0 mM NaCl).

Proline, on the other hand, accumulated, also in a time- and dose-dependent manner, so that by day 19, the increase in shoot proline content was *ca*. 6-fold and 26-fold with 100 and 200 mM NaCl, respectively relative to controls and was equally dramatic in roots (Supplementary Fig. 2).



Supplementary Fig. 2. Proline concentration in shoots (A) and roots (B) after 5, 12, and 19 days of exposure to 0, 50, 100 or 200 mM NaCl. Data are means \pm S.E. (n=3). Asterisks indicate significant differences (*P*<0.05) relative to controls (0 mM NaCl) for each sampling time.

Salt also induced oxidative stress, as revealed by MDA production, which was significantly enhanced in shoots by all NaCl treatments; on day 19, it was *ca*. 40% and 70% higher than in controls with 100 and 200 mM NaCl, respectively (Supplementary Fig. 3 A). In roots, only the highest NaCl concentration significantly enhanced MDA levels (Supplementary Fig. 3 B). Likewise, both shoot and root H_2O_2 and GSH concentrations revealed a salt-induced response (Supplementary Fig. 3 C-F). Interestingly, at day 19, only 200 mM NaCl still exerted a significant effect in roots (Supplementary Fig. 3 D and F).





Supplementary Fig. 3. Malondialdehyde (A, B), H_2O_2 (C, D) and GSH (E, F) concentrations in shoots (A, C, E) and roots (B, D, F) after 5, 12, and 19 days of exposure to 0, 50, 100 or 200 mM NaCl. Data are means \pm S.E. (n=3). Asterisks indicate significant differences (*P*<0.05) relative to controls (0 mM NaCl) for each sampling time.

Effects of treatments with 100 mM NaCl and HEs

Pre-treatment with either 0.1 or 1.0 mL L⁻¹ IB led to a significant shoot growth recovery, similar for both concentrations, that went *ca.* 22-25% beyond non-saline controls and that was 73.6% (FW)and 77.0% (DW) higher than with NaCl alone (Fig. 1 A,C). Conversely, plants pre-treated with SC or methanol exhibited the same reduction in shoot FW and DW as those treated with NaCl alone, regardless of the dose applied. Roots also responded positively to pre-treatment with 1.0 mL L⁻¹ IB (PRE-IB 1.0) in terms of FW, but not DW (Fig.1 B,D). Improved shoot growth (FW and DW) relative to both control and saline conditions was also registered after post-treatment with IB (POST-IB) irrespective of IB concentration (Fig.1 A,C); a significant (46.2%) growth recovery, but only in terms of DW, was also observed after post-treatment with 1.0 mL L⁻¹ SC (POST-SC 1.0) (Fig. 1 C). Root growth (FW and DW) in the presence of 100 mM NaCl was likewise improved, even beyond control levels, by POST-IB 1.0 and, as regards DW, also by POST-SC 1.0 (Fig. 1 B,D). Methanol had no effect on the growth performance of maize plants.

CER C



Fig. 1. Shoot (A, C) and root (B, D) fresh and dry weights after 12 days of exposure to 100 mM NaCl in the presence of 0.1 or 1 ml L⁻¹ IB, SC, or methanol (MeOH) added two days before (PRE) or two days after (POST) the start of salt treatment. Data are means \pm S.E. (n=10). Different letters indicate significant differences (*P*<0.05).

Under saline conditions, both shoots and roots accumulated Na⁺, although the increase, relative to controls, was higher in the former (23-fold) than in the latter (*ca.* 10-fold). By contrast, K⁺ levels declined in both organs (25% and 47% in shoots and roots, respectively; Fig. 2 A,B). Na⁺ accumulation in shoots was not affected by IB treatments, but it was reduced to the same extent by PRE-SC 1.0 and PRE-MeOH 1.0 (Fig. 2 A). Na⁺ uptake in roots was most effectively reduced by PRE-MeOH 0.1 and POST-IB 1.0; conversely, it was enhanced by both doses of PRE-IB and POST-SC (Fig. 2 B). The decline in shoot K⁺ levels in salt-treated *vs* control plants was slightly, but significantly (p < 0.05), reverted by PRE-IB 0.1 (Fig. 2 A). In roots, all treatments with IB as well as PRE-MeOH 1.0 and POST-SC (both doses) increased K⁺ levels as compared with salt treatment alone (Fig. 2B). Salt stress caused a strong decline in K⁺/Na⁺ ratios in both shoots (from *ca.* 45 in controls to 1.5 with NaCl) and roots (from 11.5 to *ca.* 0.7 in control *vs* salt-treated plants).

Treatment with HEs or MeOH had no ameliorative effect, except POST-IB 1.0, which doubled the ratio in roots, relative to salt alone, by increasing K⁺ without increasing Na⁺ levels.



Fig. 2. Shoot (A) and root (B) sodium (Na⁺) and potassium (K⁺) concentrations after 12 days of exposure to 100 mM NaCl in the presence of 0.1 or 1 ml L⁻¹ IB, SC, or methanol (MeOH) added two days before (PRE) or two days after (POST) the start of salt treatment. Data are means \pm S.E. (n=3).

The salt-induced decline in Chl*a*, Chl*b* and their sum as well as that in total carotenoids was reverted only by pre-treatment with 0.1 or 1.0 mL L⁻¹ IB (Fig. 3 A). Post-treatment with IB, however, did not produce the same positive effects (Fig. 3 B). Both pre- and post-treatments with SC and MeOH had no effect or even exacerbated the negative response in terms of pigment concentrations (Fig. 3). PRE-IB (both doses) also decreased the Chl*a*/Chl*b* ratio, while other treatments either had no effect or increased it. Finally, the Σ carotenoid/ Σ chlorophyll ratio ranged from 0.14 to 0.18 and did not change notably in relation to the treatments (data not shown).



Fig. 3. Concentration of photosynthetic pigments and Chl*a*/Chl*b* ratios after 12 days of exposure to 100 mM NaCl in the presence of 0.1 or 1 ml L^{-1} IB, SC, or methanol (MeOH) added two days

before (A) or two days after (B) the start of salt treatment. Data are means \pm S.E. (n=3). Different letters indicate significant differences (*P*<0.05).

Upon treatment with IB (both doses), recovery of proline to control levels was observed, without differences between pre- and post-treatment; all other treatments resulted in no change relative to salt alone or even further accumulation (with methanol) of this salt stress-related compound (Fig. 4 A, B).



Fig. 4. Proline concentration in shoots (A) and roots (B) after 12 days of exposure to 100 mM NaCl in the presence of 0.1 or 1 ml L^{-1} IB, SC or methanol (MeOH) added two days before (PRE) or two

days after (POST) the start of salt treatment. Data are means \pm S.E. (n=3). Different letters indicate significant differences (*P*<0.05).

After pre-treatment with either dose of IB, MDA accumulation under saline conditions, indicative of oxidative stress, returned to control values in shoots, although the effect was not significantly different from that of PRE-MeOH 0.1 (Fig. 5 A). In roots, all pre-treatments reduced MDA accumulation as compared with NaCl alone, with PRE-IB 0.1 producing the strongest ameliorative effect (Fig. 5 B). POST-IB 0.1 also strongly contrasted the salt-induced MDA accumulation in both organs; in shoots, this beneficial effect was significantly stronger than with pre-treatment, while in roots pre-treatment was better than post-treatment. Post-treatment with the lower dose of SC and methanol exerted a similar stress-mitigating effect, but only in roots, so the effect was due to the solvent (Fig. 5 A, B).

Salt-induced accumulation of H_2O_2 in shoots and roots was slightly, but significantly, mitigated by all pre-treatments and to the same extent. In post-treatment, however, 0.1 mL L⁻¹ IB caused the strongest reduction in shoot H_2O_2 production, while other treatments had no or a negative effect (Fig. 5 C, D). In roots, POST-IB 0.1 was again the most effective treatment, while all other post-treatments (except 1.0 mL L⁻¹ MeOH) led to only a slight decline in H_2O_2 levels as compared with salt alone (Fig. 5 D).

Salt-induced GSH content in shoots was slightly, but significantly, decreased by PRE-IB 1.0, while neither SC nor methanol had any effect. When given after the start and for the entire duration of the salt treatment, none of the extracts contrasted the rise in shoot GSH concentration (Fig. 5 E). In roots, the response was quite different. In fact, a significant increase in GSH relative to NaCl was observed, both after pre- and post-treating plants with SC and methanol, while IB had no effect (Fig. 5 F).



Treatment

Fig. 5. Malondialdehyde (A, B), H_2O_2 (C, D) and GSH (E, F) concentrations in shoots (A, C, E) and roots (B, D, F) after 12 days of exposure to 100 mM NaCl in the presence of 0.1 or 1 ml L⁻¹ IB, SC or methanol (MeOH) added two days before (PRE) or two days after (POST) the start of salt treatment. Data are means \pm S.E. (n=3). Different letters indicate significant differences (*P*<0.05).

Effects of treatment with 200 mM NaCl and IB

The efficacy of the extract treatment that gave the best results with 100 mM NaCl (i.e., 1.0 mL L^{-1} IB) was tested with the higher dose of salt (200 mM NaCl) albeit for a shorter duration (8 days). As

shown in Table 3, both pre- and post-treatment with IB resulted in a growth stimulation of shoots and roots relative to the salt treatment both in terms of fresh and dry biomass. Pre-treatment with IB improved shoot growth to a greater extent than the post-treatment: *ca*. 53% (FW) and 78% (DW); in roots, FW and DW increments (relative to salt) were of the order of 50%. Although methanol also stimulated growth relative to salt, the increase was always lower than that of IB, so that net increases were around 30-40%. These positive effects on growth were not, however, accompanied by significant changes in the other parameters. Thus, the decline in photosynthetic pigments and the accumulation of proline, MDA, H_2O_2 , and GSH induced by 200 mM NaCl were not mitigated, either by IB or methanol (data not shown).

		Sh	oot		Root						
	Fresh weight variation (%)										
	1 ml l ⁻¹ IB 1 ml l ⁻¹ MeOH				1 ml	$\mathbf{I}^{1}\mathbf{I}\mathbf{B}$	1 ml l ⁻¹ MeOH				
VS	pre	post	pre	post	pre	post	pre	post			
NaCl	+ 53.1	+ 31.2	+ 25.2	+ 4.4	+ 50.9	+ 29.3	+ 10.4	- 5.7			
		Dry weight variation (%)									
VS	pre	post	pre	post	pre	post	pre	post			
NaCl	+ 77.8	+ 43.4	+ 44.4	+ 10.1	+ 47.5	+ 22.0	+ 16.9	- 5.1			

Table 3. Percent variation in shoot and root FW and DW relative to NaCl after 8 days of exposure to 200 mM NaCl and pre- or post-treatment with 1 mL L^{-1} IB or methanol.

Discussion

Present results indicate that the maize genotype used in this study was moderately sensitive to salt stress. Indeed, reductions in shoot and root growth (except root DW) of 20-33% were observed with 100 mM NaCl. This is in accord with a previous study, aimed at investigating the genetic variability of salt tolerance in maize, in which a reduction in growth from 20 to 80% was already evident at 100 mM NaCl in all tested accessions (Hoque et al., 2015). In the present study, the response to 100 mM salt was characterized by changes in several biochemical parameters, including photosynthetic pigments, proline and oxidative markers, such as MDA, and H_2O_2 . Salt stress also resulted in a lowering of the K⁺/Na⁺ ratio. Moreover, all growth, and biochemical changes observed under saline treatments were, in general, concentration- and time-dependent and more evident in

shoots than in roots, even though the amount of Na^+ accumulated in the two plant portions was similar in absolute terms.

IB and SC were tested for their possible role as biostimulants on maize plants subjected to salt stress. The responses appear to be differentially modulated by the two extracts, and chemical composition may account for these differences. Indeed, only IB, even at the lower concentration tested, was able to recover the performance of stressed plants in terms of growth, photosynthetic pigments content, and levels of salt stress markers, such as proline, lipid peroxidation products, and H₂O₂. Thus, IB can be regarded as a potential biostimulant able to mitigate salt stress. Present results also indicate that the biostimulant activity of IB, in salt-stressed maize plants, strictly depended on timing of treatment. Thus, whereas shoot growth recovery and reduction of proline accumulation occurred when plants were either pre-treated or post-treated with the extract, POST-IB 0.1 was most effective in mitigating oxidative stress. Conversely, the positive effect on photosynthetic pigments, i.e., recovery of Chla and b levels, only occurred when IB was applied to plants before salt stress; when applied two days after salt treatment, the stress-alleviating effect was no longer observed. Moreover, the decrease in Chla/b ratio induced by PRE-IB relative to control and salt-treated plants may be of particular relevance, since Chlb is favoured over Chla in exerting a protective function of the photosystems, due to the lower photo-oxidation rate of the former compared to the latter (de Souza et al., in press). A similar alleviating effect on chlorophyll content in NaCl-stressed bean plants was reported after foliar application Howladar et al. (2014) or seed pre-soaking with Moringa oleifera leaf extract (Rady et al. 2013). Ertani et al. (2016) also reported that Chla and Chlb were differentially affected by treatment with various plant extracts (blueberry, hawthorn, red grape skin). In Salicornia, adaptation to stress was revealed by the maintenance of a high ratio (0.12 to 0.14) of photo-protective pigments (i.e., carotenoids) against light-harvesting chlorophylls (de Souza et al. in press). In our study, neither salt alone nor salt combined with HEs had any effect on this ratio.

Proline accumulation under various types of abiotic stress (Scoccianti et al., 2016), including salinity, is a common response and overproduction of this compatible solute in transgenic plants confers a higher salt stress tolerance (Kishor et al. 1995). In addition to their osmoprotective role, organic osmolytes, such as proline, contribute to contrasting oxidative stress (Szabados and Savouré, 2010). Interestingly, IB, at both doses and times of application, reverted the salt-induced proline increase in shoots and roots to control levels, a strong indication in favor of its stress-mitigating effect.

Sodium and chloride are responsible for both osmotic stress and ion-specific toxicity that significantly reduce crop growth and yield. The best characterized mechanisms of tolerance involve

limiting Na⁺ uptake, excluding Na⁺ from leaves, and efficient vacuolar compartmentation of Na⁺ (Munns and Tester 2008). Soil salinity also causes ion imbalance by affecting, for example, the uptake of potassium (K⁺), which is an essential macronutrient in plants. Maintaining high K⁺/Na⁺ ratios is regarded as a major strategy for coping with salinity stress in salt-sensitive (glycophyte) species (e.g., cereals, such as barley) and this is often achieved by K⁺ retention rather than Na⁺ exclusion (Shabala and Pottosin 2014). The positive effect of moringa leaf extract on shoot K⁺ under high salinity was previously reported in wheat (Yasmeen et al., 2013). As regards ion homeostasis under salt stress, here we show that none of the treatments improved the K⁺/Na⁺ ratio in shoots, whereas in roots, post-treatment with the higher dose of IB was able to slightly ameliorate this parameter; the ratio increased due to enhanced K⁺ levels and no change in Na⁺ levels.

Salt stress is known to result in extensive lipid peroxidation, a parameter that has often been used as indicator of oxidative damage in membranes (Miller et al. 2010). Depending on timing and concentration, both HEs as well as methanol were able to reduce MDA production in salt-stressed roots. In shoots, however, post-treatment with the lower dose of IB was most effective in mitigating salt-induced lipid peroxidation as well as in reducing both shoot and root H_2O_2 concentration. In spite of the dose- and time-dependent ameliorative effect of IB on oxidative stress, growth recovery under saline conditions was observed when the extract was applied at either dose and both in pre- and post-treatment. This suggests that the positive effect of IB on growth was not simply the consequence of reduced oxidative stress, as also confirmed by the results of the experiment conducted using 200 mM NaCl, in which growth recovery was observed, without a substantial change in biochemical parameters.

It is worth noting that partial recovery from salt-induced oxidative stress was also observed with methanol alone, at the lower dose and in both organs. Some authors have investigated the biological functions of solvents used in plant experiments (Savvides et al. 2016), and a clear role of ethanol and methanol in influencing several plant responses has been reported. In *Arabidopsis thaliana* and rice plants, the application of exogenous ethanol enhanced salinity stress tolerance by regulating ROS-related genes and enhancing ROS detoxification (Nguyen et al. 2017); an increase in tolerance to chilling stress has also been reported in rice plants (Kato-Noguchi 2008). As regards methanol, its positive effects on growth and water use efficiency has been documented in various plant species (Behrouzyar et al. 2016), even though they strictly depend on exposure time, tissue morphology and, especially, application method. In tomato, Arabidopsis, and tobacco plants, foliar spraying with methanol enhanced plant growth under normal conditions, while root applications caused severe damage (Rowe et al. 1994; Ramirez et al. 2006). Nevertheless, it has to be underlined that the methanol concentrations (0.01-0.1%, v/v) applied by us to maize roots were much lower than those

used in the above-cited studies, and this can explain the, sometimes, ameliorative action or ineffectiveness of this solvent.

In plants, glutathione (G) is an essential component of the cellular antioxidant defense system. It is the substrate of glutathione peroxidase and glutathione-S-transferases, enzymes involved in the removal of ROS, and the ascorbate-G cycle is regarded as the principal means of superoxide and H₂O₂ removal. Reduced/oxidized forms of G (GSH/GSSG) influence the redox status of plant cells. Although increases in GSH levels have been measured in response to chilling, heat shock, and other forms of abiotic stress, the numerous publications on this topic (Tausz et al. 2004 and references therein) indicate that the GSH/GSSG ratio may change one way or another, suggesting that the role of G is particularly complex. In our study, maize plants exposed to 100 mM NaCl exhibited significantly higher GSH levels as compared with non-saline controls, with only a slight reduction in shoot FW/DW and no inhibition of root biomass, suggesting that GSH may have contributed to salt acclimation. The strongest increase in GSH levels, however, occurred with 200 mM NaCl, which significantly depressed plant biomass. Thus, under more severe conditions, although the plant activated this antioxidant response, it was not enough to ensure normal growth. The saltinduced increase in GSH concentrations could be interpreted as an "overcompensation" to keep G in its reduced, active form (Tausz et al. 2004) or an interference with cellular metabolism limiting the conversion of GSH to GSSG. Indeed, De Azevedo Neto et al. (2006) reported that G reductase activity in leaves of salt-stressed maize plants (both salt-tolerant and salt-sensitive) was greater than in control plants. Ruiz and Blumwald (2002) showed that synthesis of cysteine and GSH increased significantly when Brassica napus was exposed to salt stress. Similarly, the contents of GSH, MDA, O₂⁻⁻, and H₂O₂ were significantly increased in bean plants stressed with 200 mM NaCl (Latif and Mohamad 2016). Overall, these results suggest that the processes leading to the accumulation of GSH are salt stress-elicited. The pattern of changes in GSH levels in response to spelt HEs and methanol showed that, in shoots, the higher dose of IB reverted salt-induced GSH accumulation when plants were exposed to 100 mM NaCl. On the other hand, roots responded in a relevant manner only to SC and, especially, methanol, which, instead, caused a further increase in GSH levels. Thus, although GSH is regarded as having a positive (antioxidant) role under stress conditions, our results indicate that GSH increased with increasing stress (100 vs 200 mM NaCl), so that the effect of IB can be considered as stress-mitigating, while SC and methanol had the opposite effect.

Extract composition plays an outstanding role in determining the efficacy of biostimulants (Bulgari et al., 2015). The phytochemical investigation of the two spelt HEs revealed that their chemical composition was rather different, both in qualitative and quantitative terms. In particular, SC

contained, besides phenolic and hydroxycinnamic acids, also organic and fatty acids, which were absent in IB, while the latter had a much higher level of ferulic, p-coumaric, and caffeic acids compared to the former. Indeed, most hydroxycinnamic acids in spelt and einkorn (Triticum *monococcum* L.) grains are localized in cell walls where they are bound to structural components, such as cellulose, lignin, and arabinoxylans (Hidalgo and Brandolini 2014) and this might explain their higher levels in IB compared to SC. The greater amount of hydroxycinnamic acids, especially ferulic and p-coumaric acid, in IB could account for its higher efficacy, compared to SC, in counteracting the negative impact of salt stress. Besides the numerous functions of ferulic acid in plant growth and development (e.g., cell wall lignification), a major physiological role is likely to be its potent antioxidant capacity (Graf 1992). Thus, the higher antioxidant potential of IB, revealed by attenuated lipid peroxidation and H₂O₂ accumulation, may be due to its higher ferulic acid content and may have contributed to alleviate the toxicity generated by salt stress. Moreover, the higher fatty acid content of SC may render this extract more hydrophobic, thus less prone to interact with the root surface and be absorbed. Differences in the biostimulatory effects in maize plants of two phenol-containing extracts, deriving from lignin-rich biorefinery wastes, have been recently reported; both extracts were able to act as biostimulants, but at different concentration ranges and through diverse mechanisms, for example, via modification of the plant's hormonal balance (Savy et al. 2017). Other authors also investigated the potential biostimulant effect of polyphenol-enriched fractions derived from plant by-products. Maize plants supplied with two different doses of extracts obtained from dry apple and blueberry residues displayed a significant increase in root and leaf biomass and a higher content in macronutrients and proteins; extracts also exerted a positive impact on secondary metabolism associated with the synthesis of phenolic compounds (Ertani et al. 2011b). Phenolic acids also have allelopathic properties. Allelopathic water extracts (AWE) containing, among other compounds, phenolic acids, can improve tolerance to abiotic stresses when applied to crops, such as wheat (Farooq et al. 2018).

Finally, a growth-stimulating effect of IB was also observed in maize plants grown under non-saline conditions, confirming its potential as biostimulant. This effect could be accounted for by the phytohormone (gibberellin, auxin)-like activity reported for phenols (Ertani et al. 2016; Savy et al. 2017) and their ability to influence endogenous phytohormone levels (Einhellig, 2004).

In conclusion, we show here that polyphenol-containing methanolic extracts prepared from spelt husks have a growth-stimulating and stress-mitigating effect on maize plants by acting on different targets, including accumulation of compatible solutes, photosynthetic pigments, oxidative stress, and ion balance. The plethora of mechanisms activated by the extract can be related to its chemical complexity, as occurs with all biostimulants, with IB generally exerting a more positive action than

SC (Fig. 6). Work is in progress to investigate the effect of spelt HEs on plant polyphenol metabolism and regulation of genes involved in polyphenol biosynthesis. Experiments are also underway to test extracts prepared by using more environmentally compatible extraction procedures.

				PRE						POST			
		IB 0.1	IB 1.0	SC 0.1	SC 1.0	M 0.1	M 1.0	IB 0.1	IB 1.0	SC 0.1	SC 1.0	M 0.1	M 1.0
FW	S												
	R												
DW	S												
	R												
Chl <i>a</i>	S								Q				
Chl <i>b</i>	S												
Chla+b	S												
Cc+x	S												
proline	S												
	R												
MDA	S												
	R												
H_2O_2	S												
	R												
GSH	S						ĺ ́						
	R												

				$\boldsymbol{\boldsymbol{\zeta}}$
highes	t to lov	vest pos	sitive ef	fect
		~		
		T		

highest to lowest negative effect

Fig. 6 Graphical representation of the effects on growth and metabolic parameters of pre- or post-treatments with IB, SC or methanol in shoots (S) and roots (R) of maize plants grown under saline (100 mM NaCl) conditions. Darker and lighter colour shadings represent relatively higher and lower values, respectively.

CORTEN MARINES

Funding

This research was funded by Progetti Valorizzazione DiSB - Università di Urbino 2017/ 2018 to V.S. and by Ricerca Fondamentale Orientata - Università di Bologna to F.A. and S.B.

Declarations of interest: none

References

- Antognoni, F., Mandrioli, R., Bordoni, A., Di Nunzio, M., Viadel, B., Gallego, E., Villalba, M.P., Tomás-Cobos, L., Taneyo Saa, D.L., Gianotti, A., 2017. Integrated evaluation of the potential health benefits of einkorn-based breads. Nutrients 9, 1232-1248. https://doi.org/10.3390/nu9111232
- Ashraf, M., Harris, P.J.C., 2004. Potential biochemical indicators of salinity tolerance in plants. Plant Sci. 166, 3-16. https://doi.org/10.1016/j.plantsci.2003.10.024
- Ashraf, M.A., Akbar, A., Parveen, A., Rasheed, R., Hussain, I., Iqbal, M., 2018. Phenological application of selenium differentially improves growth, oxidative defense and ion homeostasis in maize under salinity stress. Plant Physiol. Biochem. 123, 268–280. https://doi.org/10.1016/j.plaphy.2017.12.023
- Balasundram, N., Ai, T.Y., Sambanthamurthi, R., Sundram, K., Samman, S., 2005. Antioxidant properties of palm fruit extracts. Asia Pac. J. Clin. Nutr. 14, 319–324. https://doi.org/10.1101/gr.174615.114
- Bates, L.S., Waldren, R.P., Teare, I.D., 1973. Rapid determination of free proline for water-stress studies. Plant Soil 39, 205–207. https://doi.org/10.1007/BF00018060
- Behrouzyar, E.K., Yarnia, M., 2016. Effects of methanol foliar application on nutrient content and RWC of sugar beet under water deficit stress. Bangladesh J. Bot. 45, 1069–1074.
- Brundu, S., Palma, L., Picceri, G.G., Ligi, D., Orlandi, C., Galluzzi, L., Chiarantini, L., Casabianca, A., Schiavano, G.F., Santi, M., Mannello, F., Green, K., Smietana, M., Magnani, M., Fraternale, A., 2016. Glutathione depletion is linked with Th2 polarization in mice with a retrovirus-induced immunodeficiency syndrome, murine AIDS: role of proglutathione molecules as immunotherapeutics. J. Virol. 90, 7118–30. https://doi.org/10.1021/nl034016+
- Bulgari, R., Cocetta, G., Trivellini, A., Vernieri, P., Ferrante, A., 2015. Biostimulants and crop responses: a review. Biol. Agric. Hortic. 31, 1-17. https://doi.org/10.1080/01448765.2014.964649
- Bulgari, R., Morgutti, S., Cocetta, G., Negrini, N., Farris, S., Calcante, A., Spinardi, A., Ferrari, E., Mignani, I., Oberti, R., Ferrante, A., 2017. Evaluation of borage extracts as potential biostimulant using a phenomic, agronomic, physiological, and biochemical approach. Front. Plant Sci. 8, 1–16. https://doi.org/10.3389/fpls.2017.00935

- Caretto, S., Linsalata, V., Colella, G., Mita, G., Lattanzio, V., 2015. Carbon fluxes between primary metabolism and phenolic pathway in plant tissues under stress. Int. J. Mol. Sci. 16, 26378-26394. https://doi.org/10.3390/ijms161125967
- Chen, S.-K., Edwards, C.A., Subler, S., 2003. The influence of two agricultural biostimulants on nitrogen transformations, microbial activity, and plant growth in soil microcosms. Soil Biol. Biochem. 35, 9–19. https://doi.org/10.1016/S0038-0717(02)00209-2
- Clemens, S., Weber, M., 2016. The essential role of coumarin secretion for Fe acquisition from alkaline soil. Plant Signal. Behav. 11:e1114197. doi: 10.1080/15592324.2015.1114197
- Cook, N.C., Samman, S., 1996. Flavonoids chemistry, metabolism, cardioprotective effects, and dietary sources. J. Nutr. Biochem. 7, 66-76. https://doi.org/10.1016/0955-2863(95)00168-9
- De Azevedo Neto, A.D., Prisco, J.T., Enéas-Filho, J., Abreu, C.E.B.D., Gomes-Filho, E., 2006. Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salttolerant and salt-sensitive maize genotypes. Environ. Exp. Bot. 56, 87–94. https://doi.org/10.1016/j.envexpbot.2005.01.008
- de Souza, M., Mendes, C., Doncato, K., Badiale-Furlong, E., Costa, C., 2018. Growth, phenolics, photosynthetic pigments, and antioxidant response of two new genotypes of sea asparagus (*Salicornia neei* Lag.) to salinity under greenhouse and field conditions. Agriculture 8, 115. https://doi.org/10.3390/agriculture8070115
- Desoky, E.S.M., Merwad, A.R.M., Rady, M.M., 2018. Natural biostimulants improve saline soil characteristics and salt stressed-sorghum performance. Commun. Soil Sci. Plant Anal. 49, 967–983. https://doi.org/10.1080/00103624.2018.1448861
- Dorokhov, Y.L., Sheshukova, E. V., Komarova, T. V., 2018. Methanol in plant life. Front. Plant Sci. 9, 1–6. https://doi.org/10.3389/fpls.2018.01623
- du Jardin, P., 2015. Plant biostimulants: definition, concept, main categories and regulation. Sci. Hortic. (Amsterdam) 196, 3-14. https://doi.org/10.1016/j.scienta.2015.09.021
- Einhelling F.A., 2004. Mode of allelochemical action of phenolic compounds. In: Macias, F.A., Galindo, J.C.G., Molinillo, J.M.G., Cutler, H.G. (Eds.) *Allelopathy Chemistry and Mode of Action of Allelochemicals* (pp. 217-238). CRC Press LLC, Boca Raton, Florida.
- Ertani, A., Nardi, S., Altissimo, A., 2013. Review: long-term research activity on the biostimulant properties of natural origin compounds. Acta Hortic. 1009, 181–187. https://doi.org/10.17660/ActaHortic. 2013.1009.22
- Ertani, A., Cavani, L., Pizzeghello, D., Brandellero, E., Altissimo, A., Ciavatta, C., Nardi, S., 2009. Biostimulant activity of two protein hydrolyzates in the growth and nitrogen metabolism of maize seedlings. J. Plant Nutr. Soil Sci. 172, 237–244. https://doi.org/10.1002/jpln.200800174
- Ertani, A., Francioso, O., Tugnoli, V., Righi, V., Nardi, S., 2011a. Effect of commercial lignosulfonate-humate on *Zea mays* L. metabolism. J. Agric. Food Chem. 59, 11940–11948. https://doi.org/10.1021/jf202473e

- Ertani, A., Pizzeghello, D., Francioso, O., Tinti, A., Nardi, S., 2016. Biological activity of vegetal extracts containing phenols on plant metabolism. Molecules 21, 1–14. https://doi.org/10.3390/molecules21020205
- Ertani, A., Schiavon, M., Altissimo, A., Franceschi, C., Nardi, S., 2011b. Phenol-containing organic substances stimulate phenylpropanoid metabolism in *Zea mays*. J. Plant Nutr. Soil Sci. 174, 496–503. https://doi.org/10.1002/jpln.201000075
- Farooq, M., Hussain, M., Wakeel, A., Siddique, K.H.M., 2015. Salt stress in maize: effects, resistance mechanisms, and management. A review. Agron. Sustain. Dev. 35, 461–481. https://doi.org/10.1007/s13593-015-0287-0.
- Feitosa de Vasconcelos, A.C.F. de, Zhang, X., Ervin, E.H., Kiehl, J. de C., 2009. Enzymatic antioxidant responses to biostimulants in maize and soybean subjected to drought. Sci. Agric. 66, 395–402. https://doi.org/10.1590/S0103-90162009000300015
- Graf, E., 1992. Antioxidant potential of ferulic acid. Free Radic. Biol. Med. 13, 435–448. https://doi.org/10.1016/0891-5849(92)90184-I
- Halliwell, B., 2008. Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and *in vivo* studies? Arch. Biochem. Biophys. 476, 107–112. https://doi.org/10.1016/j.abb.2008.01.028
- Hidalgo, A., Brandolini, A., 2014. Nutritional properties of einkorn wheat (*Triticum monococcum* L.). J. Sci. Food Agric. 94, 601-612. https://doi.org/10.1002/jsfa.6382
- Hoagland, D.R., Arnon, D.I., 1950. The water-culture method for growing plants without soil. Calif. Agr. Expt. Sta. Circ. 347, 1–32. https://doi.org/citeulike-article-id:9455435
- Hoque, M.M.I., Jun, Z., Guoying, W., Muhebbullah, M., Hoque, I., 2015. Evaluation of salinity tolerance in maize (*Zea mays* L.) genotypes at seedling stage. J. BioSci. Biotechnol 4, 39–49.
- Howladar, S.M., 2014. A novel *Moringa oleifera* leaf extract can mitigate the stress effects of salinity and cadmium in bean (*Phaseolus vulgaris* L.) plants. Ecotoxicol. Environ. Saf. 100, 69–75. https://doi.org/10.1016/j.ecoenv.2013.11.022
- Kahrizi, S., Sedighi, M., Sofalian, O., 2012. Effect of salt stress on proline and activity of antioxidant enzymes in ten durum wheat cultivars. Ann. Biol. Res 3, 3870–3874.
- Kato-Noguchi, H., 2008. Low temperature acclimation mediated by ethanol production is essential for chilling tolerance in rice roots. Plant Signal. Behav. 3, 202–203. https://doi.org/10.4161/psb.3.3.5542
- Kavi Kishor, P., Sangam, S., Amrutha, R.N., Laxmi, P.S., Naidu, K.R., Rao, K.R.S.S., Sreenath, R., Reddy, K.J., Theriappan, P., Sreenivasulu, N., 2005. Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress. Curr. Sci 88, 424–438.
- Khan, W., Rayirath, U.P., Subramanian, S., Jithesh, M.N., Rayorath, P., Hodges, D.M., Critchley, A.T., Craigie, J.S., Norrie, J., Prithiviraj, B., 2009. Seaweed extracts as biostimulants of plant

growth and development. J. Plant Growth Regul. 28, 386–399. https://doi.org/10.1007/s00344-009-9103-x

- Kiš, D., Jovičić, N., Matin, A., Kalambura, S., Vila, S., Guberac, S., 2017. Energy value of agricultural spelt residue (*Triticum spelta* L.) forgotten cultures. Teh. Vjesn. Tech. Gaz. 24, 369–373. https://doi.org/10.17559/TV-20170406124003
- Latif, H.H., Mohamed, H.I., 2016. Exogenous applications of moringa leaf extract effect on retrotransposon, ultrastructural and biochemical contents of common bean plants under environmental stresses. South African J. Bot. 106, 221–231. https://doi.org/10.1016/j.sajb.2016.07.010
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. Methods Enzymol. 148, 350–382. https://doi.org/10.1016/0076-6879(87)48036-1
- Mahajan, S., Tuteja, N., 2005. Cold, salinity and drought stresses: An overview. Arch. Biochem. Biophys. 444, 139–158. doi:10.1016/j.abb.2005.10.018
- Maqbool, N., Sadiq, R., 2017. Allelochemicals as growth stimulators for drought stressed maize. Am. J. Plant Sci. 08, 985–997. https://doi.org/10.4236/ajps.2017.85065
- Mattila, P., Pihlava, J.M., Hellstrőm, J., 2005. Contents of phenolic acids, alkyl- and alkenylresorcinols, and avenanthramides in commercial grain products. J. Agric. Food Chem. 53, 8290–8295. https://doi.org/10.1021/jf051437z
- Miller, G., Suzuki, N., Ciftci-Yilmaz, S., Mittler, R., 2010. Reactive oxygen species homeostasis and signalling during drought and salinity stresses. Plant, Cell Environ. 33, 453–467. https://doi.org/10.1111/j.1365-3040.2009.02041.x
- Mirakhori, M., Paknejad, F., Moradi, F., Ardakani, M., Zahedi, H., Nazeri, P., 2009. Effect of drought stress and methanol on yield and yield components of Soybean Max (L 17). Am. J. Biochem. Biotechnol. 5, 162–169. https://doi.org/10.3844/ajbbsp.2009.162.169
- Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. Annu. Rev. Plant Biol. 59, 651–681. https://doi.org/10.1146/annurev.arplant.59.032607.092911
- Nguyen, H.M., Sako, K., Matsui, A., Suzuki, Y., Mostofa, M.G., Ha, C. Van, Tanaka, M., Tran, L.-S.P., Habu, Y., Seki, M., 2017. Ethanol enhances high-salinity stress tolerance by detoxifying reactive oxygen species in *Arabidopsis thaliana* and Rice. Front. Plant Sci. 8, 1001. https://doi.org/10.3389/fpls.2017.01001
- Okoruwa, A., 1997. Utilization and processing of maize. IITA Res. Guid. 35, 29.
- Povero, G., Mejia, J.F., Di Tommaso, D., Piaggesi, A., Warrior, P., 2016. A systematic approach to discover and characterize natural plant biostimulants. Front. Plant Sci. 7, 1–9. https://doi.org/10.3389/fpls.2016.00435
- Rady, M.M., Varma C., B., Howladar, S.M., 2013. Common bean (*Phaseolus vulgaris* L.) seedlings overcome NaCl stress as a result of presoaking in *Moringa oleifera* leaf extract. Sci. Hortic. (Amsterdam) 162, 63–70. https://doi.org/10.1016/j.scienta.2013.07.046

- Ramírez, I., Dorta, F., Espinoza, V., Jiménez, E., Mercado, A., Peña-Cortés, H., 2006. Effects of foliar and root applications of methanol on the growth of Arabidopsis, tobacco, and tomato plants. J. Plant Growth Regul. 25, 30–44. https://doi.org/10.1007/s00344-005-0027-9
- Rowe, R.N., Farr, D.J., Richards, B.A.J., 1994. Effects of foliar and root applications of methanol or ethanol on the growth of tomato plants (*Lycopersicon esculentum* Mill). New Zeal. J. Crop Hortic. Sci. 22, 335–337. https://doi.org/10.1080/01140671.1994.9513842
- Ruiz, J.M., Blumwald, E., 2002. Salinity-induced glutathione synthesis in *Brassica napus*. Planta 214, 965–969. https://doi.org/10.1007/s00425-002-0748-y
- Sairam, R.K., Tyagi, A., 2004. Physiology and molecular biology of salinity stress tolerance in plants. Curr. Sci. 86, 407–421. https://doi.org/10.2307/24108735
- Savvides, A., Ali, S., Tester, M., Fotopoulos, V., 2016. Chemical priming of plants against multiple abiotic stresses: mission possible? Trends Plant Sci. 21, 329–340. https://doi.org/10.1016/j.tplants.2015.11.003
- Savy, D., Mazzei, P., Drosos, M., Cozzolino, V., Lama, L., Piccolo, A., 2017. Molecular characterization of extracts from biorefinery wastes and evaluation of their plant biostimulation. ACS Sustain. Chem. Eng. 5, 9023–9031. https://doi.org/10.1021/acssuschemeng.7b01928
- Schiavon, M., Ertani, A., Nardi, S., 2008. Effects of an alfalfa protein hydrolysate on the gene expression and activity of enzymes of the tricarboxylic acid (TCA) cycle and nitrogen metabolism in Zea mays L. J. Agric. Food Chem. 56, 11800–11808. https://doi.org/10.1021/jf802362g
- Schiavon, M., Pizzeghello, D., Muscolo, A., Vaccaro, S., Francioso, O., Nardi, S., 2010. High molecular size humic substances enhance phenylpropanoid metabolism in maize (*Zea mays* L.). J. Chem. Ecol. 36, 662–669. https://doi.org/10.1007/s10886-010-9790-6
- Schieber, A., Stintzing, F.C., Carle, R., 2001. By-products of plant food processing as a source of functional compounds – recent developments. Trends Food Sci. Technol. 12, 401–413. https://doi.org/10.1016/S0924-2244 (02)00012-2
- Scoccianti, V., Bucchini, A.E., Iacobucci, M., Ruiz, K.B., Biondi, S., 2016. Oxidative stress and antioxidant responses to increasing concentrations of trivalent chromium in the Andean crop species *Chenopodium quinoa* Willd. Ecotoxicol. Environ. Saf. 133, 25–35. https://doi.org/10.1016/j.ecoenv.2016.06.036
- Shabala, S., Pottosin, I., 2014. Regulation of potassium transport in plants under hostile conditions: Implications for abiotic and biotic stress tolerance. Physiol. Plant. 151, 257-279. https://doi.org/10.1111/ppl.12165
- Sharma, P., Jha, A.B., Dubey, R.S., Pessarakli, M., 2012. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. J. Bot. 2012, 1–26. https://doi.org/10.1155/2012/217037
- Szabados, L., Savouré, A., 2010. Proline: a multifunctional amino acid. Trends Plant Sci. 15, 89-97. https://doi.org/10.1016/j.tplants.2009.11.009

- Tausz, M., Šircelj, H., Grill, D., 2004. The glutathione system as a stress marker in plant ecophysiology: is a stress-response concept valid?. J. Exp. Bot. 55, 1955–1962. https://doi.org/10.1093/jxb/erh194
- Velikova, V., Yordanov, I., Edreva, A., 2000. Oxidative stress and some antioxidant systems in acid rain-treated bean plants protective role of exogenous polyamines. Plant Sci. 151, 59–66. https://doi.org/10.1016/S0168-9452(99)00197-1
- Wang, W., Vinocur, B., Altman, A., 2003. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. Planta 218, 1-14. https://doi.org/10.1007/s00425-003-1105-5
- Wei, P., Chen, D., Jing, R., Zhao, C., Yu, B., 2015. Ameliorative effects of foliar methanol spraying on salt injury to soybean seedlings differing in salt tolerance. Plant Growth Regul. 75, 133– 141. https://doi.org/10.1007/s10725-014-9938-4
- Yasmeen, A., Basra, S.M.A., Farooq, M., Rehman, H. ur, Hussain, N., Athar, H. ur R., 2013. Exogenous application of moringa leaf extract modulates the antioxidant enzyme system to improve wheat performance under saline conditions. Plant Growth Regul. 69, 225–233. https://doi.org/10.1007/s10725-012-9764-5
- Zhu, J.K., 2001. Plant salt tolerance. Trends Plant Sci. 6, 66-71. https://doi.org/10.1016/S1360-1385(00)01838-0

ACCEPTED MANUSCRIPT										
		Sh	oot		Root					
	Fresh weight variation (%)									
	1 ml L^{-1} IB 1 ml L^{-1} MeOH				1 ml L ⁻¹ IB		1 ml L ⁻¹ MeOH			
	Pre	Post	Pre	Post	Pre	Post	Pre	Post		
vs NaCl	+53.1	+31.2	+25.2	+4.4	+50.9	+29.3	+10.4	-5.7		
	Dry weight variation (%)									
	Pre	Post	Pre	Post	Pre	Post	Pre	Post		
vs NaCl	+77.8	+43.4	+44.4	+10.1	+47.5	+22.0	+16.9	-5.1		

Table 3. Percent variation in shoot and root FW and DW relative to NaCl after 8 days of exposure to 200 mM NaCl and pre- or post-treatment with 1 mL L^{-1} IB or methanol.

Two polyphenol-enriched extracts were prepared from spelt by-products.

Chemical composition of the two extracts was assessed by HPLC-DAD and GC-MS analysis.

Maize plants subjected to salt stress were treated with polyphenol-enriched extracts.

The efficacy of spelt husk extracts as biostimulant on maize plants was evaluated.

Authors' contributions to the manuscript Ceccarini et al.

V.S. and F.A. conceived and designed the experiments; C.C. and F.A. performed the experiments, collected and analyzed the data; A.F performed GSH measurements; G.V. and A.G. did the GC-MS analyses; V.S., F.A. and S.B. analyzed the data and drafted the manuscript. All authors read and approved the final manuscript.