



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

## ARCHIVIO ISTITUZIONALE DELLA RICERCA

### Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Pliocene colonization of the Mediterranean by Great White Shark inferred from fossil records, historical jaws, phylogeographic and divergence time analyses

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Agostino Leone, E.S. (2020). Pliocene colonization of the Mediterranean by Great White Shark inferred from fossil records, historical jaws, phylogeographic and divergence time analyses. JOURNAL OF BIOGEOGRAPHY, 47(5), 1119-1129 [10.1111/jbi.13794].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/712445> since: 2024-05-24

*Published:*

DOI: <http://doi.org/10.1111/jbi.13794>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

1 **Pliocene colonization of the Mediterranean by Great White Shark inferred from fossil records,**  
2 **historical jaws, phylogeographic and divergence time analyses**

3  
4 Agostino Leone<sup>1,\*</sup>, Gregory Neils Puncher<sup>2,3</sup>, Francesco Ferretti<sup>4,5</sup>, Emilio Sperone<sup>6</sup>, Sandro Tripepi<sup>6</sup>,  
5 Primo Micarelli<sup>7</sup>, Andrea Gambarelli<sup>8</sup>, Maurizio Sarà<sup>9</sup>, Marco Arculeo<sup>9</sup>, Giuliano Doria<sup>10</sup>, Fulvio  
6 Garibaldi<sup>11</sup>, Nicola Bressi<sup>12</sup>, Andrea Dall'Asta<sup>12</sup>, Daniela Minelli<sup>13</sup>, Elisabetta Cilli<sup>14</sup>, Stefano  
7 Vanni<sup>15</sup>, Fabrizio Serena<sup>16</sup>, Píndaro Díaz-Jaimes<sup>17</sup>, Guy Baele<sup>18</sup>, Alessia Cariani<sup>1</sup>, Fausto Tinti<sup>1</sup>

8 <sup>1</sup>Dept. Biological, Geological & Environmental Sciences (BiGeA), Laboratory of Genetics &  
9 Genomics of Marine Resources and Environment (GenoDREAM), University of Bologna, 48123,  
10 Ravenna, Italy

11 <sup>2</sup>Dept Biological Sciences, Canadian Rivers Institute, University of New Brunswick, Saint John,  
12 NB, E2L 4L5, Canada

13 <sup>3</sup>Genomics Laboratory, Maurice-Lamontagne Institute, Fisheries and Oceans Canada, Mont-Joli,  
14 QC, G5H 3Z4, Canada

15 <sup>4</sup>Hopkins Marine Station, Department of Biology, Stanford University, 120 Oceanview Boulevard,  
16 Pacific Grove, CA 93950, USA

17 <sup>5</sup>Department of Fish and Wildlife Conservation, Virginia Tech, 310 West Campus Drive, Blacksburg,  
18 VA 24060, USA

19 <sup>6</sup>DiBEST Department of Biology, Ecology and Earth Science, University of Calabria, 87036,  
20 Arcavacata di Rende, Italy

21 <sup>7</sup> Sharks Studies Center (C.S.S.), 58024, Massa Marittima (GR), Italy

22 <sup>8</sup>Museum of Zoology and Comparative Anatomy of Modena, University of Modena and Reggio  
23 Emilia, 41121, Modena, Italy

24 <sup>9</sup>Dipartimento di Scienze e Tecnologie Biologiche Chimiche e Farmaceutiche (STEBICEF),  
25 University of Palermo, 90123, Palermo, Italy

26 <sup>10</sup>Civic Museum of Natural History "Giacomo Doria", 16121, Genova, Italy

27 <sup>11</sup>Department of Earth Sciences, Environmental and Life, University of Genova, 16132, Genova, Italy

28 <sup>12</sup>Civic Museum of Natural History of Trieste, 34100, Trieste, Italy

29 <sup>13</sup>Museum of Comparative Anatomy, University of Bologna, 40126, Bologna, Italy

30 <sup>14</sup>Department of Cultural Heritage, University of Bologna, 48121, Ravenna, Italy

31 <sup>15</sup>Museum of Natural History of Firenze "La Specola", 50125, Firenze, Italy

32 <sup>16</sup>Institute for Biological Resources and Marine Biotechnology (IRBIM), National Research Council  
33 – CNR, 91026, Mazara del Vallo (TP), Italy

34 <sup>17</sup>Laboratorio de Genética de Organismos Acuáticos Instituto de Ciencias del Mar y Limnología,  
35 Universidad Nacional Autónoma de México, Apdo. Postal 70-305, 04510 Mexico, D.F., Mexico

36 <sup>18</sup> Department of Microbiology and Immunology, Rega Institute, KU Leuven, 3000 Leuven, Belgium

37 \*Corresponding author: agostino.leone@unibo.it

38

39

40 **Abstract**

41 **Aim**

42 Determine the evolutionary origin of the heretofore poorly characterized contemporary Great White  
43 Shark (GWS; *Carcharodon carcharias*) of the Mediterranean Sea, using phylogenetic and dispersal  
44 vicariance analyses to trace back its global paleo-migration pattern.

45 **Location** Mediterranean Sea

46 **Taxon** *Carcharodon carcharias*

47 **Methods** We have built the largest mtDNA control region (CR) sequence dataset for the  
48 Mediterranean GWS from referenced historical jaws spanning the 19th and 20th centuries.  
49 Mediterranean and global GWS CR sequences were analyzed for genetic diversity, phylogenetic  
50 relationships and divergence time. A Bayes factor approach was used to assess two scenarios of GWS  
51 lineage divergence and emergence of the Mediterranean GWS line using fossil records and paleo-  
52 geographical events for calibration of the molecular clock.

53 **Results** The results confirmed a closer evolutionary relationship between Mediterranean GWS and  
54 populations from Australia-New Zealand and the Northeastern Pacific coast rather than populations  
55 from South-African and Northwestern Atlantic. The Mediterranean GWS lineage showed the lowest  
56 genetic diversity at the global level, indicating its recent evolutionary origin. An evaluation of various  
57 divergence scenarios determined the Mediterranean GWS lineage most likely appeared some 3.23  
58 million years ago by way dispersal/vicariance from Australian/Pacific paleo-populations.

59 **Main conclusion** Based on the fossil records, phylogeographic patterns and divergence time, we  
60 revealed that the Mediterranean GWS population originated in the Pliocene following the Messinian  
61 Salinity Crisis. Colonization of the Mediterranean by GWS likely occurred via an eastward paleo-  
62 migration of Australian/eastern Pacific elements through the Central American Seaway, before the  
63 complete closure of the Isthmus of Panama. This Pliocene origin scenario contrasts with a previously  
64 proposed scenario in which Australian GWS colonized the Mediterranean via antipodean northward  
65 migration resulting from navigational errors from South Africa during Quaternary climatic  
66 oscillations.

67 **Keywords** Great White Shark, Mediterranean, historical DNA, divergence time, phylogeography,  
68 *Carcharodon carcharias*

## 70 **Background**

71 Large predatory shark populations in the Mediterranean Sea have declined dramatically over  
72 the last century (Ferretti, Myers, Serena, & Lotze, 2008). The loss of apex predators throughout the  
73 world's oceans over the past century is likely to have caused profound ecological alterations and  
74 potentially large-scale trophic cascades (Myers, Baum, Shepherd, Powers & Peterson, 2007; Ferretti,  
75 Worm, Britten, Heithaus, & Lotze, 2010). Among the species that witnessed the most precipitous  
76 declines is the Great White Shark (*Carcharodon carcharias*, L.1758, henceforth GWS; McPherson  
77 & Myers, 2009), which is currently listed as Critically Endangered in European seas (Nieto et al.,  
78 2015). GWS are widespread throughout the globe, aside from the polar regions, with hotspots of  
79 abundance located off the coasts of South Africa, Australia, New Zealand, Japan, North and South  
80 America, and in the Mediterranean (Compagno, 1984; Fergusson, 1996). Integrated ecological,  
81 genetic and tagging data have revealed natal homing and philopatric behavior of GWS, with  
82 extraordinary trans-oceanic migrations of both sexes between geographically distant populations in  
83 the Indian (Pardini et al., 2001; Bonfil et al., 2005; Blower, Pandolfi, Bruce, Gomez-Cabrera, &  
84 Ovenden, 2012) and North Pacific Ocean (Domeier & Nasby-Lucas, 2008; Jorgensen et al., 2010,  
85 2012). In the Mediterranean, GWS have long been observed and documented by the public, resource  
86 users and scientists, who have provided opportunistic occurrence records, from direct sightings,  
87 commercial fisheries, records of bite marks found on prey and museum specimens dating back to the  
88 early 19th century (Mancusi et al., 2002; De Maddalena, 2006; Sperone et al., 2012). In the past, free-  
89 swimming individuals and pairs were frequently observed in areas where large pelagic fisheries were  
90 intense (e.g. the Sicilian Channel, the Ligurian and Tyrrhenian seas; Fergusson, 1996; Storai, Vanni,  
91 Zuffa, & Biagi, 2005). A few records of GWS pups from Turkey and Tunisia, as well as juveniles in  
92 the Sicilian Channel, suggest that the Mediterranean may host GWS nursery areas (Fergusson, 1996;  
93 Storai, Mojetta, Zuffa, & Giulian, 2000; Kabasakal, & Gedikoğlu, 2008). However, the natural history  
94 of Mediterranean GWS still remains largely uncharacterized.

95 Efforts have been made in the past to resolve the phylogenetic relationships of the  
96 Mediterranean GWS population using mitochondrial DNA (mtDNA) sequences (Gubili et al., 2010;  
97 2015). Using up to five specimens from the Mediterranean, Gubili et al. (2010, 2015) concluded that  
98 the population is more closely related to populations in the Pacific Ocean (Australia, New Zealand  
99 and Northeastern Pacific) than to those from the western Indian Ocean (South-Africa) and  
100 northwestern Atlantic Ocean (Florida). Based on a nucleotide substitution rate between the two major  
101 lineages (Northeastern Pacific vs. North West Atlantic and Eastern Indian) calibrated by the  
102 formation of the Isthmus of Panama (3.5 Ma) and the Sunda-Sahul Shelves (5 Ma) respectively,  
103 Gubili et al. (2010) suggested that Mediterranean GWS are descendants of a few disoriented  
104 individuals who immigrated from Australia/New Zealand during the Pleistocene (348-565 ka) by an  
105 antipodean route along the western coast of Africa. A scenario of multiple relatively recent  
106 colonization events was also considered, based on the haplotype relationships that were generated  
107 using a few historical and contemporary Mediterranean specimens (Gubili et al., 2015).

108 Collection of fresh GWS specimens in the Mediterranean has proven to be difficult in recent  
109 decades, due to their precipitous decline in abundance. However, there is a great number of referenced

110 and dry-preserved GWS specimens in several Italian museums and private scientific archives, such  
111 as mounted skins, jaws, vertebrae and teeth collected from the Mediterranean during the last two  
112 centuries (Mancusi et al., 2002; De Maddalena, 2006). Recently developed ancient DNA (aDNA)  
113 techniques present a great opportunity for reconstructing the natural history of marine species using  
114 preserved historical materials (Hofreiter, Serre, Poinar, Kuch, & Pääbo, 2001; Riccioni et al., 2010).  
115 Unfortunately, most historical GWS specimens have been archived in sub-optimal conditions,  
116 thereby compromising their potential for DNA-based applications. Moreover, many collectors are  
117 reluctant to loan specimens for molecular study, due to their intrinsic, sentimental and market value.

118 By analyzing DNA extracted from preserved specimens of GWS caught during the last 195  
119 years, from eight Italian museums and private collections, we have been able to explore the  
120 evolutionary history of the Mediterranean GWS. Using effective and affordable aDNA techniques  
121 widely used to extract and genotype DNA from historical specimens of marine fish, we have  
122 generated a publicly available mtDNA sequence dataset from 18 Mediterranean GWS individuals.

123

## 124 **Methods**

125 Full details of the collected historical GWS specimens, sampling procedures, protocols for  
126 aDNA extraction, PCR amplification of Control Region (CR), sequencing and sequence analyses are  
127 provided in the in the Supporting Information (Supplementary Methods, Appendix S1; Figures S1-  
128 S3, Appendix S2; Tables S1-S3, Appendix S3).

### 129 *Analysis of fossil evidence*

130 The extensive catalogue of taxon-specific GWS fossils featured in the online and open access  
131 Paleobiology Database (<https://paleobiodb.org/#/>), and its associated R package ‘paleobioDB’  
132 (Varela et al., 2015), was used to create a distribution and stratigraphic map of global GWS fossils.  
133 The downloaded database was filtered manually to avoid the use of homonym extinct taxa and  
134 dubious records. Only reliable records specifically classified as *Carcharodon carcharias* and relative  
135 synonyms were retained (Table S4, Appendix S3). A detailed search of the literature with a focus on  
136 *C. carcharias* fossils from the Mediterranean Sea was carried out in order to retain reliable fossil  
137 records and exclude potential misidentification of fossil specimens (Table S5, Appendix S3) (Cigala-  
138 Fulgosi, 1990; Applegate, & Espinosa-Arrubarrena, 1996; Gottfried & Fordyce, 2001; Bianucci et  
139 al., 2002; Marsili, 2006; Marsili, 2008; Adnet, Balbino, Antunes, & Marin-Ferrer, 2009). The fossil  
140 records from the paleobioDB database were checked for correctness using the references and fossil  
141 collections associated to the paleobioDB codes (Table S4, Appendix S3). Records without  
142 descriptions of the fossil or without pictures, especially those from old references, as well as records  
143 with misidentified fossils (e.g. *Isurus* spp. or *Carcharodon* spp. identified as *Carcharodon*  
144 *carcharias*), were removed.

### 145 *Time of the Most Recent Common Ancestor (TMRCA) and estimation of evolutionary rate*

146 The divergence time analysis of the GWS lineages was carried out using Bayesian inference  
147 through Markov Chain Monte Carlo (MCMC) sampling as implemented in BEAST v1.10.0 (Suchard

148 et al., 2018). An initial analysis using calibration priors without sequence data was carried out to  
149 determine if calibration priors interacted unexpectedly and to assess if the data were informative  
150 (Fulton, & Strobeck, 2010). A relaxed molecular clock and a constant population size coalescent  
151 model were used to recover time-stamped phylogenies in BEAST. To ensure convergence of the  
152 posterior distributions, three independent MCMC analyses were run (20 million steps, sampled every  
153 1k generations, burn-in 50%). Convergence and effective sample sizes (ESS) were verified using  
154 Tracer v.1.7.0 (Rambaut, Drummond, Xie, Baele, & Suchard, 2018). A Maximum Clade Credibility  
155 (MCC) tree was summarized using TREEANNOTATOR V.1.10.0 and visualized in FIGTREE V.1.4.3  
156 (Rambaut, 2009). Since the molecular clock is sensitive to bias when a short fragment with fewer  
157 polymorphisms is used, the TMRCA analyses were carried out using the two CR sequence datasets  
158 of different length (516bp and 828bp), as previously used in the haplotype network analysis  
159 (Supplementary Methods in Appendix S1). The divergence time of the GWS lineages was estimated  
160 using two alternative combinations of calibration priors in BEAST v1.10.0 (Suchard et al., 2018).  
161 The alternative combinations of calibration priors were built by integrating the estimated age of the  
162 earliest GWS fossil records (Applegate & Espinosa-Arrubarrena, 1996) (Mean: 11.0 Ma; SD: 1.0)  
163 with two different secondary calibrations: a) the best dated fossil record of GWS in the Mediterranean  
164 Sea, dated back to the Pliocene (Cigala-Fulgosi, 1990; Bianucci et al., 2002) (Mean: 3.0 Ma; SD:  
165 0.30 Ma), and b) the molecular divergence previously estimated by Gubili et al. (2010) (Mean: 0.4  
166 Ma; SD: 0.15). Both combinations of calibration priors were implemented as normally distributed.  
167 Rather than using debated fossil records from the middle Miocene epoch (Gottfried & Fordyce, 2001),  
168 we used the divergence time between the GWS and the outgroup *Lamna nasus* (GenBank Acc. No.  
169 GU266755-GU266769) as an alternative first calibration. The divergence time of these two lineages  
170 has been estimated at around 46 Ma (Martin, 1996) (Mean: 46.0 Ma; SD: 1.0).

#### 171 *Marginal likelihood estimation and testing divergence time hypotheses*

172 After setting the first calibration for the earliest fossil attributed to the GWS, the two alternative  
173 secondary calibrations could be compared by estimating log marginal likelihoods using generalized  
174 stepping-stone (GSS) sampling, as implemented in BEAUti v1.10.0 and BEAST v1.10.0 (Suchard et  
175 al., 2018; Baele, Lemey, & Suchard, 2015). The log marginal likelihood values for the two different  
176 scenarios were first estimated using generalized stepping-stone sampling (100 stepping stones, 1  
177 million iterations, logging every 1,000 iterations). A total of 101 power posteriors, with one million  
178 iterations each, were sampled using MCMC for the GSS approach. In order to select the alternative  
179 calibrations that fits best with the principal timing information fixed for both scenarios, the log Bayes  
180 factor was calculated for both scenarios using the formula  $\log BF = \log \Pr(D | M1) - \log \Pr(D | M2)$ ,  
181 where  $\log \Pr(D | M1)$  is the log marginal likelihood for model 1, and  $\log \Pr(D | M2)$  is the log  
182 marginal likelihood for model 2.. This analysis was performed on both sequence datasets (i.e. 516bp  
183 and 828bp).

#### 184 *Reconstructing the historical biogeography*

185 Two approaches were implemented in order to reconstruct the historical biogeography of  
186 species: the Statistical Dispersal-Vicariance Analysis (S-DIVA; Yu, Harris, & He, 2010), which is a  
187 parsimony method of historical biogeography, and Dispersal-Extinction-Cladogenesis analysis  
188 (DEC; Ree, & Smith, 2008). An MCC estimated by using BEAST and TreeAnnotator on just  
189 haplotype sequences, and the specimen distribution through all biogeographical areas (A:  
190 Australia/New Zealand, AUS; B: Northeastern Pacific, NEP; C: Mediterranean, MED; D: South-  
191 Africa, SA; E: Northwestern Atlantic, NWA) was used to perform the S-DIVA and DEC analyses  
192 implemented in RASP v. 4.0 (Yu, Harris, Blair, & He, 2015).

193

## 194 **Results**

195 Partial CR sequences (515bp) from 18 GWS historical specimens were obtained and deposited  
196 in GenBank (Acc. No. MN718579-MN718596).

197 The multiple sequence alignment containing sequences from all 18 historical samples and four  
198 Mediterranean homologous modern sequences deposited in GenBank (HQ540294-HQ540296;  
199 JF715925; Table S3 in Appendix S3) showed an extremely low nucleotide diversity among the CR  
200 fragment (< 0.1%), with only one variable site at position 244, in which the specimen FICC02LI (See  
201 Table S1 in Appendix S3), with GenBank Acc. No. MN718587, showed a transition (A > G). A final  
202 alignment containing Mediterranean sequences merged with the homologous sequences from the  
203 other global populations (N = 99) resulted in a final dataset of 117 CR sequences of 516bp in which  
204 an indel at position 60 was revealed.

205 The global ML haplotype network revealed 31 haplotypes (Figure S4, Appendix S2) that were  
206 clustered in two main haplogroups differentiated by 28 mutations. As expected, based on the existing  
207 literature, the first haplogroup was formed by the individuals from the MED and the Pacific Ocean  
208 (AUS and NEP) and the second was composed of GWS from SA and the NWA. Three haplotypes of  
209 GWS collected in AUS (HQ414073, HQ414074 and AY026211; Table S3, Appendix S3) clustered  
210 in the latter haplogroup and these individuals are SA-like individuals that likely migrated across the  
211 Indian Ocean (Pardini et al., 2001; Blower et al., 2012). The ML haplotype network built with 99  
212 GWS sequences of 828bp revealed 68 haplotypes and a similar topology (Figure S4, Appendix S2)  
213 to the one reconstructed using the shorter sequence dataset, with 2 main haplogroups (AUS-NEP-  
214 MED vs SA-NWA).

215 The cross-plot for the haplotype and nucleotide diversity of the Mediterranean and global  
216 populations revealed that the MED and AUS populations have the lowest and highest values for both  
217 indices, respectively (Figure 1; Table S6, Appendix S3). The NEP and NWA populations showed  
218 high haplotype diversity and low nucleotide diversity. The SA population exhibited quite opposite  
219 positions in the plot, depending on the reference study used (Pardini et al. (2001) and O'Leary et al.  
220 (2015) both offered high values for both indices and Andreotti et al. (2016) provided low values.

221 No polymorphisms were detected among the four complete contemporary CR sequences  
222 isolated from Mediterranean GWS, demonstrating a low haplotype diversity among longer sequences  
223 as well, while the genetic diversity (e.g. nucleotide) is proportional to the length of the sequences.

224

225 *Marginal likelihood estimation and test for divergence time hypotheses*

226 The application of the log Bayes factor formula gave significant support to the Pliocene  
227 calibration scenario in every model tested (Table 1). Based on the 828bp dataset, the GWS  
228 Mediterranean population diverged from the Pacific populations approximately 3.23 Ma, a time that  
229 is congruent with the estimated closure of the Central American Seaway, CAS (about 3.5 Ma), after  
230 the formation of the Isthmus of Panama (O’Dea et al., 2016). Considering the “Pleistocene  
231 divergence” scenario, we obtained a slight mismatch between the posterior distribution generated  
232 from the data and posterior distribution generated from specified calibration using priors only. This  
233 may suggest that a second Pleistocene calibration could be conflicting with the data and that a  
234 “Pliocene divergence” scenario is preferable.

235 *Time of the Most Recent Common Ancestor (TMRCA) and estimation of evolutionary rate*

236 The analysis of fossil evidence suggested that GWS experienced a long evolutionary history  
237 (Tables S4 and S5, Appendix S3). The occurrence of GWS fossils in the Mediterranean area is high  
238 in the Pliocene, after the Messinian Salinity Crisis.

239 The CR relaxed substitution rate range inferred from the TMRCA analysis was estimated at  
240 0.38%-0.72% substitutions/site/my for the best model following the Bayes factor. Based on these  
241 estimates, the coalescence of the Mediterranean and Pacific GWS lineages was dated at 3.23 Ma for  
242 the alignment containing 828bp (Figure 2) and 1.81 Ma for the alignment of 516bp (Figure S5,  
243 Appendix S2).

244 *Historical biogeography reconstruction*

245 The analysis of the ancestral ranges of GWS, using both S-DIVA and DEC approaches,  
246 suggested that the Mediterranean population is potentially the result of a dispersal-vicariance  
247 scenario. The biogeographical reconstruction of the Mediterranean lineage estimated using S-DIVA  
248 resulted in two principal ancestral ranges, AC (AUS-MED) and ABC (AUS-NEP-MED), with  
249 probabilities (relative frequencies) for each range of 52.44% and 46.69% respectively. Two minor  
250 ancestral ranges were detected with very low probability: AB (AUS-NEP) and BC (NEP-MED), with  
251 probabilities of 0.52% and 0.35%, respectively. The DEC analysis resulted in concordant results with  
252 two ancestral ranges, AC (AUS-MED); BC (NEP-MED), with probabilities of 50.01% and 49.99%  
253 respectively, supporting that the origin of the Mediterranean population is likely Australia and the  
254 North Eastern Pacific. Both approaches suggest the following route of dispersal-vicariance:  
255 AC>CAB>C|AB. For both S-DIVA and DEC analyses, Australian and North Eastern Pacific lineages  
256 originated from a Pacific ancestral range AB (AUS-NEP) with a probability of 100%.

257

258 **Discussion**

259 Our phylogenetic analysis of contemporary and historical sequences indicates that the existing  
260 population of GWS in the Mediterranean Sea could be a relic of the Pacific GWS clade. This scenario  
261 would suggest that Pacific-born ancestors migrated eastward across the ocean, before the closure of  
262 the Central American Seaway. Following this, these itinerant migrants would have colonized the



263 North Atlantic Ocean before entering the Mediterranean Sea after the Messinian Salinity Crisis  
264 (MSC). They would have entered the Mediterranean during a period of ecological upheaval,  
265 following the MSC which caused the extinction of the local marine megafauna due to the total or  
266 partial drying up of the nearly landlocked sea (Garcia-Castellanos et al., 2009). This evolutionary  
267 pathway (Figure 3) is supported by Bayesian analyses of genetic diversity and divergence time  
268 estimates and it is coherent with the age of the main paleo-geographical events, paleo-climatic  
269 patterns and fossil records. This pathway represents an alternative to the antipodean dispersal  
270 hypothesis (Gubili et al., 2010) which suggested that a few Australian/New Zealand founder females  
271 that visited the Good Hope Cape area became confused by Pleistocene climatic oscillations, swam  
272 northward and found themselves in the Mediterranean Sea.

273 Our phylogenetic analysis of the CR sequences as well as the plot analysis of haplotype  
274 confirmed that the Mediterranean GWS have a closer evolutionary relationship with the  
275 Australian/New Zealand and Northeastern Pacific lineages than with the South African and  
276 Northwestern Atlantic lineage. These results are in agreement with previous studies (Gubili et al.,  
277 2010; Blower et al., 2012; Andreotti et al., 2016). The Mediterranean GWS exhibited a very shallow  
278 mtDNA genetic variation with only two haplotypes and extremely low genetic diversity. Among the  
279 modern samples collected from throughout the world that have been analyzed to date, the  
280 Mediterranean GWS have provided the lowest estimates of genetic diversity. This is likely an  
281 indication that the population is relatively young and has likely originated by a founder event by a  
282 limited number of mtDNA lineages or that the population has experienced a recent population  
283 bottleneck (Group 1 of Grant, & Bowen, 1998; Grant, & Waples, 2000). Within the  
284 Pacific/Mediterranean mitochondrial clade, the Australian/New Zealand GWS population was  
285 identified as the most ancestral with a long evolutionary/demographic history and divergent  
286 haplotypes that have accumulated over long periods of time (as indicated by the high haplotype and  
287 nucleotide diversity values; Group 3 of Grant, & Bowen, 1998; Grant, & Waples, 2000). An  
288 intermediate evolutionary position in the clade could be that of the Northeastern Pacific GWS  
289 population which possesses high haplotype diversity and low nucleotide diversity. This pattern is  
290 indicative of a population that has experienced a bottleneck event followed by rapid demographic  
291 growth and accumulation of mutations (Group 2 of Grant, & Bowen 1998; Grant, & Waples, 2000).  
292 Both S-DIVA and DEC analyses suggest that the Mediterranean GWS originated from a dispersal-  
293 vicariance event via eastward dispersal from a few Australian/Pacific individuals during the Pliocene.  
294 The founder event that gave origin to the modern Mediterranean GWS was dated by the TMRCA  
295 analysis, with high statistical support, to 3.23 Ma, during the Piacenzian Age (3.6-2.58 Ma). This is  
296 much earlier than the estimate provided by the antipodean dispersal hypothesis (348-565 ka;  
297 Calabrian, Pleistocene), which is based on a mtDNA substitution rate of 1.19-0.74% of divergence  
298 between lineages per million year (Gubili et al., 2010). These rates of evolutionary change were  
299 calibrated by Gubili et al. (2010) using estimates of vicariance events that separated GWS populations  
300 from the Northeastern Pacific and Northwestern Atlantic oceans (i.e. the rising of the Isthmus of  
301 Panama dated at 3.5 Ma; O'Dea et al., 2016) and the Western Pacific and Indian oceans (i.e. the rising  
302 of the Sunda-Sahul shelves dated at 5 Ma; Haq, Hardenbol, & Vail, 1987). However, all phylogenetic

303 analyses carried out so far (Gubili et al., 2010; Andreotti et al., 2016; present work) have revealed  
304 that the Northwestern Atlantic GWS are phylogenetically linked to the South African population but  
305 not to the Northeastern Pacific GWS. Therefore, the use of the vicariance event separating Atlantic  
306 and Pacific GWS might have led to an overestimation of the mutation rate and the subsequent time  
307 of divergence between Mediterranean and Pacific GWS at 348-565 ka during the Pleistocene.

308 The origination time of the species is still debated (see dedicated paragraph in Gottfried &  
309 Fordyce, 2001). Despite the fossil records from the Miocene are less common, this should not exclude  
310 a Miocene origin of the species. Moreover, the molecular clock analysis suggested a divergence  
311 between *Carcharodon* and *Isurus* dated back to 43 Ma (Martin, 1996), in contrast with a Pliocene  
312 origin of the *C. carcharias* from *I. hastalis*. However, a more dedicated study on the fossil history is  
313 needed to better clarify the origination time of the species.

314 Abundant fossil data suggests that GWS have inhabited the Mediterranean Sea since the early  
315 Pliocene, following the Messinian Salinity Crisis, with numerous specimens estimated to be between  
316 ~5 and ~2 Ma old, with peaks of abundance occurring during the Pliocene (Cigala-Fulgosi, 1990;  
317 Bianucci et al., 2002; Marsili, 2008; Adnet, Balbino, Antunes, & Marín-Ferrer, 2009). After these  
318 paleoclimatic phases, GWS could have colonized the Mediterranean Sea occupying the ecological  
319 niches left empty by other apex marine predators, such as the giant megatooth shark *Carcharocles*  
320 *megalodon*, which went extinct between 3.5 and 2.6 Ma (Pimiento, & Clements, 2014). Fossil  
321 evidence from Central America suggests that many species of sharks and marine mammals that were  
322 part of the region's faunal assemblage may have migrated across the CAS continuously before the  
323 formation of the Isthmus of Panama (Steeman et al., 2009; Pimiento et al., 2013; Velez-Juarbe, Wood,  
324 De Gracia, & Hendy, 2015). Upon the closure of the CAS, the newly formed Gulf Stream current  
325 could have facilitated a trans-Atlantic migration by way of the easterly current and subsequent  
326 eastward dispersion of nutrients and, consequently, food resources. There is evidence that intense  
327 fluctuations in the speed of the Gulf Stream occurred during the formation of the Isthmus of Panama,  
328 reaching a height during the late Miocene and early Pliocene (~6.1-4.8 Ma; Kaneps, 1979). The  
329 intensification of currents coincides with the end of the Messinian Salinity Crisis (5.33 Ma), which  
330 culminated with the Zanclean inflow in the Mediterranean Sea (Garcia-Castellanos et al., 2009). All  
331 of these oceanographic phenomena are concordant with our estimated divergence of the  
332 Mediterranean GWS population.

333 A potential earlier formation of the Isthmus of Panama (Bacon et al., 2015; Lessios et al., 2015;  
334 Montes et al., 2015; O'Dea et al., 2016) would not affect our results, since it is plausible that the  
335 Pacific lineage that gave rise to the MED population may already have been present in the North  
336 Atlantic before the closure of the CAS. Moreover, some researchers have suggested that the  
337 Caribbean Sea and Pacific Ocean may have remained intermittently connected by shallow waters  
338 other than the CAS (Jaramillo et al., 2017). The global phylogeography of contemporary GWS  
339 populations reveals a discontinuous distribution of the Pacific/Mediterranean GWS evolutionary  
340 lineage in the North Atlantic being interrupted by the recent colonization of the Northwestern Atlantic  
341 area by the Indian lineage. The phylogenetic analyses highlighted that the GWS of Northwestern  
342 Atlantic population does not exhibit a separated mtDNA cluster like the Pacific and Mediterranean

343 populations, and it form a unique lineage with the South-Africa (Figure S4, Appendix S2). This  
344 evidence is suggestive of a recent evolutionary history or a population bottleneck followed by  
345 population growth and accumulation of mutations.

346 A local extinction of the past Pacific GWS lineage in the North Atlantic Ocean and further  
347 replacement by components of the Indian lineage could explain such a phylogeographic discontinuity.  
348 An extensive analysis of the fossil records, highlighted an exceptional extinction rate of the marine  
349 fauna, including sharks, during the Pliocene epoch (Pimiento et al., 2017). Furthermore, the highest  
350 extinction rates occurred in the late Pliocene, between 3.8 and 2.4 Ma, which coincides with the  
351 Mediterranean population divergence time, and may have led to the local extinction of the ancient  
352 Atlantic population of GWS.

353 Additional genetic and tagging studies are required to determine if the Mediterranean GWS are  
354 ecologically and reproductively isolated from the adjacent populations occupying the North Atlantic  
355 Ocean. This issue is of high-priority for the development of efficient conservation actions and  
356 implementation of management strategies. Despite this knowledge gap, some information can be  
357 gleaned from the other populations in the Atlantic, as well as other species. For example, several other  
358 epipelagic sharks migrate throughout the temperate waters of the Atlantic and several are suspected  
359 of having nursery areas in the mid-Atlantic Ridge (Kohler, Turner, Hoey, Natanson, & Briggs, 2002;  
360 Stevens, 2010; Vandeperre et al., 2014; O'Leary et al., 2015). Elsewhere, in the Indian Ocean, GWS  
361 have been observed migrating between Australia and South Africa, while in the Pacific, GWS  
362 frequently migrate between the western coast of North America and Hawaii (Bonfil et al., 2005;  
363 Jorgensen et al., 2010, 2012; Blower et al., 2012). Clearly, the species is not averse to far reaching  
364 longitudinal movements. However, antipodean connections appear less frequently and evidence for  
365 them is limited to variations in the genetic code (O'Leary et al., 2015). The latitudinal distribution of  
366 GWS could be limited by thermal tolerance, prey availability, social structure and fidelity to nursery  
367 areas (Cliff, Dudley, & Davis, 1989; Curtis et al., 2014).

368 Our results suggest that the Mediterranean GWS have a more ancient origin than previously  
369 thought, and that this population is genetically disconnected from the adjacent Atlantic population.  
370 Due to historical and widespread declines of sharks in the Mediterranean Sea (Colloca et al., 2013;  
371 Ferretti, Osio, Jenkins, Rosenberg, & Lotze, 2013, Ferretti et al., 2008) as well as the ongoing over-  
372 exploitation of marine resources in the region, the current white shark population is critically  
373 endangered (Dulvy, Allen, Ralph, & Walls, 2016). This distinct genetic pool represents a valuable,  
374 albeit extremely vulnerable, component of the genetic diversity of a species that is endangered  
375 worldwide. Losing the Mediterranean GWS population would thus represent a significant blow to the  
376 global conservation of this species. Continued characterization of this population's ecology, spatial  
377 dynamics and population structure is paramount for effective management and restoration of this  
378 important top predator's ecological role in the region.

379

380 **Table 1.** Best model selection based on Bayes factors of tree topologies reconstructed with the earliest fossil occurrences using 828bp and 516 bp  
 381 mitochondrial control region sequences. Node constraints and calibration parameters on the phylogeographical tree of *C. carcharias* for the two  
 382 divergence models are also provided. logML\_GSS: log marginal likelihood from generalized stepping stone model, BF\_GSS: Bayes factors calculated  
 383 using the logML\_GSS. Models are ranked according to the logML values.  
 384

<b>Model</b>	<b>Dataset</b>	<b>Node1</b>	<b>Node2</b>	<b>logML_GSS</b>	<b>logBF_GSS</b>
MED Pliocene divergence	828bp	Ingroup Mean: 11 Ma, SD: 1.0 Ma	MED/Pacific Mean: 3.0 Ma, SD: 0.30 Ma	-2455.2796	0 (Best)
MED Pleistocene divergence	828bp	Ingroup; Mean: 11 Ma, SD: 1.0 Ma	MED/Pacific Mean: 0.4Ma, SD: 0.15 Ma	-2458.1264	2.8468
MED Pliocene divergence	516bp	Ingroup; Offset: 11 Ma, SD: 1.0 Ma	MED/Pacific Mean: 3.0 Ma, SD: 0.30 Ma	-1440.6171	0 (Best)
MED Pleistocene divergence	516bp	Ingroup; Offset: 11 Ma, SD: 1.0 Ma	MED/Pacific Mean: 0.4Ma, SD: 0.15 Ma	-1443.0069	2.3898

385

386 **Acknowledgements**

387 We are indebted to Alessandro Velonà, Adriana Villamor and Marco Stagioni for providing valuable  
388 technical support and advice. Clemente Ventrone and Gerlando Spagnolo provided samples from  
389 Favignana tuna trap.

390

391 **Funding**

392 AL, GNP, AC and FT were funded by the University of Bologna by RFO 2016-2018 and Canziani  
393 grants to FT. AL is also funded by Italian Ministry for University and Research by a PhD grant 2015-  
394 2017. All the other authors were funded by respective institutions. GB acknowledges support from  
395 the Interne Fondsen KU Leuven / Internal Funds KU Leuven under grant agreement C14/18/094.

396

397 **Data Availability Statement**

398 The sequence data obtained in the present study have been deposited to the GenBank database of the  
399 National Center for Biotechnology Information under accession numbers MN718579-MN718596.

400

401 **References**

402 Adnet, S., Balbino, A. C., Antunes, M. T., & Marín-Ferrer, J. M. (2009). New fossil teeth of the white  
403 shark (*Carcharodon carcharias*) from the early Pliocene of Spain. Implication for its paleoecology  
404 in the Mediterranean. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, 256(1), 7-  
405 16. <http://dx.doi.org/doi:10.1127/0077-7749/2009/0029>

406 Andreotti, S., von der Heyden, S., Henriques, R., Rutzen, M., Meyer, M., Oosthuizen, H., & Matthee,  
407 C. A. (2016) New insights into the evolutionary history of white sharks, *Carcharodon carcharias*.  
408 *Journal of Biogeography*, 43(2), 328-339. <https://doi.org/10.1111/jbi.12641>

409 Applegate, S. P., & Espinosa-Arrubarrena, L. (1996). The fossil history of *Carcharodon* and its  
410 possible ancestor, *Cretolamna*: a study in tooth identification. In A. P. Klimley & D. G. Ainley  
411 (Eds.), *Great white sharks: the biology of Carcharodon carcharias* (pp. 19-36). San Diego, CA:  
412 Academic Press.

413 Bacon, C. D., Silvestro, D., Jaramillo, C., Smith, B. T., Chakrabarty, P., Antonelli, A. (2015).  
414 Biological evidence supports an early and complex emergence of the Isthmus of Panama.  
415 *Proceedings of the National Academy of Sciences U.S.A.*, 112(19), 6110-6115. doi:  
416 <https://doi.org/10.1073/pnas.1423853112>

417 Baele, G., Lemey, P., & Suchard, M. A. (2015). Genealogical working distributions for Bayesian  
418 model testing with phylogenetic uncertainty. *Systematic Biology*, 65(2), 250-264.  
419 <https://doi.org/10.1093/sysbio/syv083>

420 Bianucci, G., Bisconti, M., Landini, W., Storai, T., Zuffa, M., Giuliani, S., & Mojetta, A. (2002).  
421 Trophic interaction between white shark, *Carcharodon carcharias*, and cetaceans: a comparison  
422 between Pliocene and recent data from Central Mediterranean Sea. In M. Vacchi, G. Lamesa, F.  
423 Serena, & B. Seret (Eds.), *Proceedings of the 4th European Elasmobranch Association Meeting 28-  
424 30 November 2000* (pp. 33-48). Livorno, Italy: ICRAM, ARPAT & SFI.

- 425 Blower, D. C., Pandolfi, J. M., Bruce, B. D., Gomez-Cabrera, M.D., & Ovenden, J. R. (2012)  
426 Population genetics of Australian white sharks reveals fine-scale spatial structure, transoceanic  
427 dispersal events and low effective population sizes. *Marine Ecology Progress Series*, 455, 229-244.  
428 [https://doi: 10.3354/meps09659](https://doi.org/10.3354/meps09659)
- 429 Bonfil, R., Meyer, M., Scholl, M. C., Johnson, R., O'Brien, S., Oosthuizen, H., ... Paterson, M.  
430 (2005). Transoceanic migration, spatial dynamics, and population linkages of white sharks. *Science*,  
431 310(5745), 100-103. [https://doi:10.1126/science.1114898](https://doi.org/10.1126/science.1114898)
- 432 Canese, S. P., Giusti, M., Salvati, E., Angiolillo, M., Cardinali, A., & Consalvo, I. (2011). *Final*  
433 *Report Project ISPRA PR0.3.452 Biodiversity of Sicily Channel*. Rome, Italy: Ministry of the  
434 Environment, Territorial and Sea Protection.
- 435 Cigala-Fulgosi, F. (1990). Predation (or possible scavenging) by a great white shark on an extinct  
436 species of bottlenosed dolphin in the Italian Pliocene. *Tertiary Research*, 12, 17-36.
- 437 Cliff, G., Dudley, S. F. J., & Davis, B. (1989). Sharks caught in the protective gill nets off Natal,  
438 South Africa. 2. The great white shark *Carcharodon carcharias* (Linnaeus). *South African Journal*  
439 *of Marine Science*, 8, 131-144. [https://doi:10.2989/02577618909504556](https://doi.org/10.2989/02577618909504556)
- 440 Colloca, F., Cardinale, M., Maynou, F., Giannoulaki, M., Scarcella, G., Jenko, K., ... Fiorentino, F.  
441 (2013). Rebuilding Mediterranean fisheries: a new paradigm for ecological sustainability. *Fish and*  
442 *Fisheries*, 14(1), 89-109. [https://doi:10.1111/j.1467-2979.2011.00453.x](https://doi.org/10.1111/j.1467-2979.2011.00453.x)
- 443 Compagno, L. J. V. (1984). FAO species catalogue. Vol. 4. Sharks of the world. An annotated and  
444 illustrated catalogue of shark species known to date. Part 1. Hexanchiformes to Lamniformes. *FAO*  
445 *Fish Synopsis*, 125, 1-249.
- 446 Curtis, T. H., McCandless, C. T., Carlson, J. K., Skomal, G. B., Kohler, N. E., Natanson, L. J., ...  
447 Pratt Jr., H. L. (2014) Seasonal Distribution and Historic Trends in Abundance of White Sharks,  
448 *Carcharodon carcharias*, in the Western North Atlantic Ocean. *PLOS ONE* 9(6), e99240.  
449 [doi:10.1371/journal.pone.0099240](https://doi.org/10.1371/journal.pone.0099240)
- 450 De Maddalena, A. (2006) A catalogue of great white sharks *Carcharodon carcharias* (Linnaeus,  
451 1758) preserved in European museums. *Journal of the National Museum, Natural History Series*,  
452 175(3-4), 109-125.
- 453 Domeier, M. L., Nasby-Lucas, N. (2008). Migration patterns of white sharks *Carcharodon*  
454 *carcharias* tagged at Guadalupe Island, Mexico, and identification of an eastern Pacific shared  
455 offshore foraging area. *Marine Ecology Progress Series*, 370, 221-237. [doi:](https://doi.org/10.3354/meps07628)  
456 <https://doi.org/10.3354/meps07628>
- 457 Dulvy, N. K., Allen, D. J., Ralph, G. M., & Walls, R. H. L. (2016). The conservation status of sharks,  
458 rays and Chimaeras in the Mediterranean Sea. IUCN, Malaga, Spain.
- 459 Fergusson, I. K. (1996) Distribution and autoecology of the white shark in the Eastern North Atlantic  
460 and the Mediterranean Sea. In A. P. Klimley & D. G. Ainley (Eds.), *Great white sharks: the biology*  
461 *of Carcharodon carcharias* (pp. 321-345). San Diego, CA: Academic Press.
- 462 Ferretti, F., Worm, B., Britten, G. L., Heithaus, M. R., & Lotze, H. K. (2010). Patterns and ecosystem  
463 consequences of shark declines in the ocean. *Ecology Letters*, 13(8), 1055-1071.  
464 [https://doi:10.1111/j.1461-0248.2010.01489.x](https://doi.org/10.1111/j.1461-0248.2010.01489.x)Ferretti, F., Myers, R. A., Serena, F., & Lotze, H. K.

- 465 (2008). Loss of large predatory sharks from the Mediterranean Sea. *Conservation Biology*, 22(4),  
466 952-964. <https://doi:10.1111/j.1523-1739.2008.00938.x>
- 467 Ferretti, F., Osio, G. C., Jenkins, C. J., Rosenberg, A. A., & Lotze, H. K. (2013) Long-term change  
468 in a meso-predator community in response to prolonged and heterogeneous human impact. *Scientific*  
469 *Reports*, 3, 1057. <https://doi:10.1038/srep01057>
- 470 Fulton, T. L., & Strobeck, C. (2010). Multiple fossil calibrations, nuclear loci and mitochondrial  
471 genomes provide new insight into biogeography and divergence timing for true seals (Phocidae,  
472 Pinnipedia). *Journal of Biogeography*, 37(5), 814-829. <https://doi:10.1111/j.1365-2699.2010.02271.x>
- 474 Garcia-Castellanos, D., Estrada, F., Jiménez-Munt, I., Gorini, C., Fernández, M., Vergés, J., & De  
475 Vicente, R. (2009). Catastrophic flood of the Mediterranean after the Messinian salinity crisis.  
476 *Nature*, 462, 778-781. <https://doi:10.1038/nature08555>
- 477 Gottfried, M. D., & Fordyce, R. E. (2001). An associated specimen of *Carcharodon angustidens*  
478 (Chondrichthyes, Lamnidae) from the late Oligocene of New Zealand, with comments on  
479 *Carcharodon* interrelationships. *Journal of Vertebrate Paleontology*, 21(4), 730-739.  
480 [https://doi:10.1671/0272-4634\(2001\)021\[0730:AASOCA\]2.0.CO;2](https://doi:10.1671/0272-4634(2001)021[0730:AASOCA]2.0.CO;2)
- 481 Grant, W. S., & Bowen, B. W. (1998). Shallow population histories in deep evolutionary lineages of  
482 marine fishes: insights from sardines and anchovies and lessons for conservation. *Journal of Heredity*,  
483 89(5), 415–426. <https://doi.org/10.1093/jhered/89.5.415>
- 484 Grant, W. S., & Waples, R. S. (2000). Spatial and temporal scales of genetic variability in marine and  
485 anadromous species: implications for fisheries oceanography. In P. J. Harrison & T. R. Parsons (Eds.),  
486 *Fisheries oceanography: an integrative approach to fisheries ecology and management* (pp. 63–93).  
487 Oxford, UK: Blackwell Science.
- 488 Gubili, C., Bilgin, R., Kalkan, E., Karhan, S. Ü., Jones, C. S., Sims, D. W., ... Noble, L. R. (2010).  
489 Antipodean white sharks on a Mediterranean walkabout? Historical dispersal leads to genetic  
490 discontinuity and an endangered anomalous population. *Proceedings of the Royal Society B:*  
491 *Biological Sciences*, 278(1712), 1679–1686. <https://doi:10.1098/rspb.2010.1856>
- 492 Gubili, G., Robinson, C. E. C., Cliff, G., Wintner, S. P., de Sabata, E., De Innocentiis, S., ... Jones,  
493 C. S. (2015). DNA from historical and trophy samples provides insights into white shark population  
494 origins and genetic diversity. *Endangered Species Research* 27(3), 233-241. <https://doi:10.3354/esr00665>
- 496 Haq, B. U., Hardenbol, J., & Vail, P. R. (1987). Chronology of fluctuating sea levels since the  
497 Triassic. *Science*, 235(4793), 1156-1167. <https://doi:10.1126/science.235.4793.1156>
- 498 Hasegawa, M., Kishino, H., & Yano, T. (1985). Dating of the human-ape splitting by a molecular  
499 clock of mitochondrial DNA. *Journal of Molecular Evolution*, 22(2), 160-174.
- 500 Hofreiter, M., Serre, D., Poinar, N. H., Kuch, M., & Pääbo, S. (2001). Ancient DNA. *Nature Reviews*  
501 *Genetics*, 2, 353-359. <https://doi:10.1038/35072071>
- 502 Jaramillo, C., Montes, C., Cardona, A., Silvestro, D., Antonelli, A., Bacon, C. D. (2017). Comment  
503 (1) on “Formation of the Isthmus of Panama” by O’Dea et al. *Science Advances*, 3(6), e1602321. doi:  
504 10.1126/sciadv.1602321

- 505 Jorgensen, S. J., Arnoldi, N. S., Estess, E. E., Chapple, T. K., Rückert, M., Anderson, S. D., & Block,  
506 B. A. (2012). Eating or meeting? Cluster analysis reveals intricacies of white shark (*Carcharodon*  
507 *carcharias*) migration and offshore behavior. *PLOS ONE* 7, e47819.  
508 <https://doi:10.1371/journal.pone.0047819>
- 509 Jorgensen, S. J., Reeb, C. A., Chapple, T. K., Anderson, S., Perle, C., Van Sommeran, S.R., ... Block,  
510 A. B. (2010). Philopatry and migration of Pacific white sharks. *Proceedings of the Royal Society B:*  
511 *Biological Sciences*, 277(1682), 679-688. <https://doi:10.1098/rspb.2009.1155>
- 512 Kabasakal, H., & Gedikoğlu, Ö. S. (2008). Two new-born great white sharks, *Carcharodon*  
513 *carcharias* (Linnaeus, 1758) (Lamniformes; Lamnidae) from Turkish waters of the north Aegean Sea.  
514 *Acta Adriatica*, 49, 125–135.
- 515 Kaneps, A. G. (1979). Gulf stream: velocity fluctuations during the late Cenozoic. *Science*,  
516 204(4390), 297-301. <https://doi:10.1126/science.204.4390.297>
- 517 Kohler, N. E., Turner, P. A., Hoey, J. J., Natanson, L. J., & Briggs, R. (2002). Tag and recapture data  
518 for three pelagic shark species: blue shark (*Prionace glauca*), shortfin mako (*Isurus oxyrinchus*), and  
519 porbeagle shark (*Lamna nasus*) in the North Atlantic Ocean. *ICCAT Collective Volumes of Scientific*  
520 *Papers*, 54, 1231-1260.
- 521 Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7 Molecular Evolutionary Genetics Analysis  
522 Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874.  
523 <https://doi.org/10.1093/molbev/msw054>
- 524 Lessios, H. A. (2015). Appearance of an early closure of the Isthmus of Panama is the product of  
525 biased inclusion of data in the metaanalysis. *Proceedings of the National Academy of Sciences USA*,  
526 112(43), E5765-E5765. doi: 10.1073/pnas.1514719112
- 527 Mancusi, C., Nicolosi, P., Arculeo, M., Barbagli, F., Carlini, R., Costantini, M., ... Vacchi, M. (2002).  
528 The presence of elasmobranchs in the collections of the main Italian natural history museums. In M.  
529 Vacchi, G. Lamesa, F. Serena, & B. Seret (Eds.), *Proceedings of the 4th European Elasmobranch*  
530 *Association Meeting 28-30 November 2000* (pp. 97-108). Livorno, Italy: ICRAM, ARPAT & SFI.
- 531 Marsili, S. (2008). Systematic, paleoecologic and paleobiogeographic analysis of the Plio-Pleistocene  
532 Mediterranean elasmobranch fauna. *Atti della Società Toscana di Scienze Naturali, Memorie, Serie*  
533 *A*, 113, 81-88.
- 534 Marsili, S. (2006). Analisi Sistematica, Paleoecologica e Paleobiogeografica della Selaciofauna plio-  
535 pleistocenica del Mediterraneo. University of Pisa. PhD thesis
- 536 Martin, A. P. (1996). Systematics of the Lamnidae and the Origination Time of *Carcharodon*  
537 *carcharias* Inferred from the Comparative Analysis of Mitochondrial DNA Sequences. In A. P.  
538 Klimley & D. G. Ainley (Eds.), *Great white sharks: the biology of Carcharodon carcharias* (pp. 49-  
539 53). San Diego, CA: Academic Press.
- 540 McPherson, J. M., & Myers, R. A. (2009). How to infer population trends in sparse data: examples  
541 with opportunistic sighting records for great white sharks. *Diversity and Distributions* 15(5), 880-  
542 890. <https://doi:10.1111/j.1472-4642.2009.00596.x>



- 543 Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J. C., Valencia, ... Niño, H. (2015). Middle  
544 Miocene closure of the Central American Seaway. *Science* 348(6231), 226–229. doi:  
545 10.1126/science.aaa2815
- 546 Myers, R. A., Baum, J. K., Shepherd, T. D., Powers, S. P., Peterson, C. H. (2007). Cascading Effects  
547 of the Loss of Apex Predatory Sharks from a Coastal Ocean. *Science*, 315(5820), 1846–1850. doi:  
548 10.1126/science.1138657
- 549 Nieto, A., Ralph, G. M., Comeros-Raynal, M. T., Kemp, J., García Criado, M., Allen, D. J., ...  
550 Williams, J. T. (2015) *European Red List of marine fishes*. Luxembourg, EU: Rosseels Printing.
- 551 O’Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L., ...  
552 Jackson, J. B. C. (2016). Formation of the Isthmus of Panama. *Science Advances*, 2(8), e1600883.  
553 <https://doi.org/10.1126/sciadv.1600883>
- 554 O’Leary, S. J., Feldheim, K. A., Fields, A. T., Natanson, L. J., Wintner, S., Hussey, N., ... Chapman,  
555 D. D. (2015). Genetic diversity of white sharks, *Carcharodon carcharias*, in the Northwest Atlantic  
556 and Southern Africa. *Journal of Heredity*, 106(3), 258–265. <https://doi.org/10.1093/jhered/esv001>
- 557 Pardini, A. T., Jones, C. S., Noble, L. R., Kreiser, B., Malcolm, H., Bruce, B. D., ... Martin, A. P.  
558 (2001). Sex-biased dispersal of great white sharks. *Nature*, 412(6843), 139–140.  
559 <http://doi.org/10.1038/35084125>
- 560 Pimiento, C., & Clements, C. F. (2014). When did *Carcharocles megalodon* become extinct? A new  
561 analysis of the fossil record. *PLOS ONE*, 9(10), e111086.  
562 <https://doi.org/10.1371/journal.pone.0111086>
- 563 Pimiento, C., Gonzalez-Barbad, G., Hendy, A. J. W., Jaramillo, C., MacFadden, B. J., Montes, C., ...  
564 Shippritt, M. (2013). Early Miocene chondrichthyans from the Culebra formation, Panama: a window  
565 into marine vertebrate faunas before closure the Central American Seaway. *Journal of South  
566 American Earth Sciences*, 42, 159–170. <https://doi.org/10.1016/j.jsames.2012.11.005>
- 567 Pimiento, C., Griffin, J. N., Clements, C. F., Silvestro, D., Varela, S., Uhen, M. D., & Jaramillo, C.  
568 (2017). The Pliocene marine megafauna extinction and its impact on functional diversity. *Nature  
569 Ecology & Evolution*, 1, 1100–1106. <https://doi.org/10.1038/s41559-017-0223-6>
- 570 Rambaut, A. (2009). FigTree v1.4.3. Retrieved from <http://tree.bio.ed.ac.uk/software/figtree/>
- 571 Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarisation  
572 in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, 67(5), 901–904.  
573 <https://doi.org/10.1093/sysbio/syy032>
- 574 Ree, R. H., & Smith, S. A. (2008). Maximum Likelihood inference of geographic range evolution by  
575 dispersal, local extinction, and cladogenesis. *Systematic Biology*, 57(1), 4–14.  
576 <https://doi.org/10.1080/10635150701883881>
- 577 Riccioni, G., Landi, M., Ferrara, G., Milano, I., Cariani, A., Zane, L., ... Tinti, F. (2010). Spatio  
578 temporal population structuring and genetic diversity retention in depleted Atlantic bluefin tuna of  
579 the Mediterranean Sea. *Proceedings of the National Academy of Sciences of the United States of  
580 America*, 107(5), 2102–2107. <https://doi.org/10.1073/pnas.0908281107>

- 581 Sperone, E., Parise, G., Leone, A., Milazzo, C., Circosta, V., Santoro, F., ... Tripepi, S. (2012).  
 582 Spatiotemporal patterns of distribution of large predatory sharks in Calabria (Central Mediterranean,  
 583 Southern Italy). *Acta Adriatica*, 53, 13-24.
- 584 Steeman, M. E., Hebsgaard, M. B., Fordyce, R. E., Ho, S. Y. W., Rabosky, D. L., Nielsen, R., ...  
 585 Willerslev, E. (2009). Radiation of Extant Cetaceans Driven by Restructuring of the Oceans.  
 586 *Systematic Biology*, 58(6), 573-585. <https://doi:10.1093/sysbio/syp060>
- 587 Stevens, J. D. (2010). Epipelagic Oceanic Elasmobranchs. In J. C. Carrier, J. A. Musick, & M. R.  
 588 Heithaus (Eds.), *Sharks and Their Relatives II: Biodiversity, Adaptive Physiology, and Conservation*  
 589 (pp. 3-37). Boca Raton, FL: CRC Press.
- 590 Storai, T., Mojetta, A., Zuffa, M., Giulian, S. (2000). Nuove segnalazioni di *Carcharodon carcharias*  
 591 (L.) nel Mediterraneo centrale. *Atti Società Toscana di Scienze Naturali*, 107, 139-142.
- 592 Storai, T., Vanni, S., Zuffa, M., & Biagi, V. (2005). Presenza di *Carcharodon carcharias* (Linnaeus,  
 593 1758) nelle acque toscane (Mar Ligure meridionale e Mar Tirreno settentrionale; Mediterraneo):  
 594 analisi e revisione delle segnalazioni (1839-2004). *Atti Società Toscana di Scienze Naturali*, 112,  
 595 153-166.
- 596 Suchard, M. A., Lemey, P., Baele, G., Ayres, D. L., Drummond, A. J., & Rambaut, A. (2018).  
 597 Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evolution*, 4(1),  
 598 vey016. <https://doi:10.1093/ve/vey016>
- 599 Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control  
 600 region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10(3),  
 601 512-526. <https://10.1093/oxfordjournals.molbev.a040023>
- 602 Vandeperre, F., Aires-da-Silva, A., Fontes, J., Santos, M., Serrão Santos, R., & Afonso, P. (2014).  
 603 Movements of blue sharks (*Prionace glauca*) across their life history. *PLOS ONE* 9(8), e103538.  
 604 <https://doi:10.1371/journal.pone.0103538>
- 605 Varela, S., González-Hernández, J., Sgarbi, L. F., Marshall, C., Uhen, M. D., Peters, S., &  
 606 McClennen, M. (2015). paleobioDB: an R package for downloading, visualizing and processing data  
 607 from the Paleobiology Database. *Ecography*, 38(4), 419–425. <https://doi:10.1111/ecog.01154>
- 608 Velez-Juarbe, J., Wood, A. R., De Gracia, C., & Hendy, A. J. W. (2015). Evolutionary patterns among  
 609 living and fossil kogiid sperm whales: evidence from the Neogene of Central America. *PLOS ONE*,  
 610 10(4), e0123909. <https://doi:10.1371/journal.pone.0123909>
- 611 Yu, Y., Harris, A. J., & He, X. (2010). S-DIVA (Statistical Dispersal-Vicariance Analysis): A tool  
 612 for inferring biogeographic histories. *Molecular Phylogenetics and Evolution*, 56(2), 848-850.  
 613 <https://doi.org/10.1016/j.ympev.2010.04.011>
- 614 Yu, Y., Harris, A. J., Blair, C., & He, X. J. (2015) RASP (Reconstruct Ancestral State in Phylogenies):  
 615 a tool for historical biogeography. *Molecular Phylogenetics and Evolution*, 87, 46-49.  
 616 <https://doi:10.1016/j.ympev.2015.03.008>

617

618 **Biosketch**

619

620 **Agostino Leone** is a marine molecular ecologist and population geneticist. Former PhD student at  
621 the University of Bologna. He has broad interests in researches related to marine conservation,  
622 ranging from biogeography to population genomics, investigating marine populations structure and  
623 their response to different pressures. AL, FT conceived the study; AL, GNP, EC, AC carried out  
624 molecular work and sequences analysis; AL, GNP, FF, PDJ, GB carried out statistical analyses; ES,  
625 ST, PM, AG, MS, MA, GD, FG, ADA, DM, SV, FS collected specimens; AL, GNP, MA, EC, FS,  
626 PDJ, GB, AC, FT drafted the manuscript.

627

## 628 **Supporting Information**

629

630 Additional Supporting Information may be found in the online version:

631 **Appendix S1** Supplementary methods for the tissue sampling, historical DNA extraction, species-  
632 specific primers design, DNA amplification, sequencing and genetic diversity analysis.

633 **Appendix S2** Supplementary figures

634 **Appendix S3** Supplementary tables




Figure 1. Plot of the haplotype and nucleotide diversity values (expressed as percentage values) of global populations of *Carcharodon carcharias* inferred using sequences from the mitochondrial control region AUS: Australia/New Zealand; NEP: Northeastern Pacific; MED: Mediterranean (516bp; n = 22); SA: South-Africa; NWA: Northwestern Atlantic. Asterisk indicates values reported by Andreotti et al. [29] for the South African population.

78x56mm (300 x 300 DPI)




Figure 2. Bayesian divergence time tree of populations of *Carcharodon carcharias* inferred using sequences from the mitochondrial control region (828bp). High posterior density (HPD 95%) values are featured as blue bars. Nodes with posterior values <0.5 are not shown. Abbreviations are provided in Figure 1.

153x71mm (300 x 300 DPI)




Figure 3. Global dispersal and Pacific/Mediterranean vicariance hypothesis for *Carcharodon carcharias*. Ancient great White Sharks from the Pacific Ocean, namely Australia, dispersed via two routes: westward to South African coasts (1, light grey dotted line) and eastward to the Northeastern Pacific Ocean (2, black dotted line). The Pacific Great White Sharks were free to move eastward to the Atlantic, and in the Pliocene to an ancient Mediterranean Sea after the Messinian Salinity Crisis (3-4). Past climatic oscillation due to the closure of the Central America Seaway and the formation of the Isthmus of Panama, could have caused a local extinction or an eastward mass migration of white shark from the North Atlantic Ocean, isolating the Mediterranean population from other ancestral populations. The North Atlantic was colonized, then, in relatively recent history, when the climate conditions became more suitable (5). Fossil records extrapolated from the Paleobiology Database are indicated with colored points (red: Miocene, blue: Pliocene, green: Pleistocene). The map was created using 'paleobioDB' package in R version 3.5.1 (Varela et al., 2015). Abbreviations are provided in Figure 1.

167x83mm (300 x 300 DPI)

**Supporting Information****Pliocene colonization of the Mediterranean by Great White Shark inferred from fossil records, historical jaws, phylogeographic and divergence time analyses**

Agostino Leone, Gregory Neils Puncher, Francesco Ferretti, Emilio Sperone, Sandro Tripepi, Primo Micarelli, Andrea Gambarelli, Maurizio Sarà, Marco Arculeo, Giuliano Doria, Fulvio Garibaldi, Nicola Bressi, Andrea Dall'Asta, Daniela Minelli, Elisabetta Cilli, Stefano Vanni, Fabrizio Serena, Píndaro Díaz-Jaimes, Guy Baele, Alessia Cariani, Fausto Tinti

**Appendix S1** Supplementary methods for the tissue sampling, historical DNA extraction, species-specific primers design, DNA amplification, sequencing and genetic diversity analysis.

**Supplementary methods***Sampling*

Tissue samples from 18 historical specimens identified as *Carcharodon carcharias*, captured in the Italian Seas from 1823 to the 1980s, were collected from museums and private archives (Figure S1, Appendix S2; Table S1, Appendix S3). Due to the cultural importance of the GWS museum specimens, sampling operations were carried out with utmost care to avoid extensive and unsightly damage (e.g. collecting tissue samples from the inner surface of jaws, internal dental pulp of teeth and dried skin debris). When jaws were available, the internal point of attachment between the lower hemi-arches was used for sampling, as it is concealed from public view and is the thickest part of the jaw. In this way, it was possible to drill deep into the jaw and avoid the use of surface materials that could act as a source of contaminants (Figure S2, Appendix S2). Exogenous DNA was removed by saturating sample surfaces with a 3.0% v/v sodium hypochlorite solution for approximately 10 minutes according to the protocol of Kemp and Smith (Kemp & Smith, 2005). All instruments were sterilized with bleach and UV irradiation between samplings. Holes of 5 mm in diameter were drilled into each jaw using an electric drill set to minimum speed to avoid damage to DNA due to thermal stress (Gibbon, Penny, Štrkalj, & Ruff, 2009). As the drill bit was removed from the cartilaginous jaws and vertebrae, all dust was carefully collected. The amounts of collected tissue ranged from 16 to 409.2 mg. The powdered tissue was then transported to a sterilized laboratory dedicated to aDNA analysis. The resulting holes left in specimens were filled with a low temperature restorative paste commonly used in anthropological studies (Figure S2, Appendix S2).

*Historical DNA extraction*

Extraction of DNA and polymerase chain reaction set-up were conducted in the “pre-PCR” clean-room of the Laboratory of Genetics & Genomics of Marine Resources and Environment (GenoDREAM) of the University of Bologna, dedicated to the analysis of degraded and low copy

number DNA. The most stringent criteria to minimize and detect exogenous contamination in aDNA analysis (Cooper & Poinar, 2000) were followed (e.g. the use of extraction and amplification blanks as negative controls in each reaction).

DNA was extracted from all samples using a protocol modified from Riccioni et al. (2010), whereby homogenized tissue powder was incubated in EDTA buffer (0.5M, pH 8.0) in a shaker overnight at room temperature. Samples were then precipitated and incubated in an extraction buffer (0.1M EDTA, 0.5% N-laurylsarcosine-Na) and Proteinase K (20 mg/mL) for an additional 24 hours at 44°C. After spinning, 250 µL supernatants were transferred to tubes containing 3.5 µL of 1 µg/µL Dextran Blue, 250 µL of 4M NH<sub>4</sub>-acetate, and 500 µL of 96% v/v ethanol. After precipitation, washing and drying, the genomic DNA was re-suspended in 50 µL of distilled sterile water and stored at -20°C.

#### *Design of primer pairs, PCR amplification and sequencing of historical DNA*

A fragment of the highly variable fragment of mitochondrial control region (CR) was targeted for analysis, since it is the most commonly used marker for reconstructing white shark phylogeography, and because there are available several sequences and haplotypes deposited in the public repositories (e.g. GenBank). Since historical DNA can be extensively fragmented (Allentoft et al. 2012), species-specific primers suitable for amplifying short fragments (167-221bp) of overlapping and adjacent DNA sequence fragments were developed (Figure S3, Appendix S2). Due to the intrinsic characteristic of ancient genetic material to be highly damaged, mainly for deamination processes, resulting in transitions from C to T and G to A (Hansen, Willerslev, Wiuf, Mourier, & Arctander, 2001), at least two independent amplifications were performed for each sample, in order to improve the detection of the damaged sites. Control region sequences/haplotypes of contemporary GWS specimens were retrieved from GenBank (Table S3, Appendix S3) and aligned with MEGA v.7.0.14 (Kumar, Stecher, & Tamura, 2016). using the ClustalW algorithm (Thompson, Higgins, & Gibson, 1994). From the alignment, five CR primer pairs (Table S2, Appendix S3) were designed with the software PRIMER3 v.4.0.0 (Untergrasser et al. 2012). These were subsequently tested *in silico* (Figure S3, Appendix S2) using AmplifX software, version 1.44 (©Nicolas Jullien 2004-2013; CNRS, Aix-Marseille Université, <http://crn2m.univ-mrs.fr/pub/amplifx-dist>). PCR conditions for all gene fragments consisted of 3 minutes of denaturation at 94°C, followed by 35 cycles of 30s at 94°C, 30s at 50°C, 30s at 72°C, and a final extension period of 7 minutes at 72°C. All PCR reactions were performed in a volume of 50 µL containing approximately 10-20ng of template DNA, 1X Tris-HCl, 200mM of each dNTP, 3mM MgCl<sub>2</sub>, 0.5µM of forward and reverse primers, and 1.25 units of *Taq* DNA Polymerase (Invitrogen). PCR amplicons were cycle-sequenced from both strands by a commercial sequence service provider (Macrogen Europe, Amsterdam, Netherlands).

#### *Sequence Analysis*

The mitochondrial Control Region (CR) partial sequences obtained from the historical samples were checked and edited using MEGA v.7.0.14 (Kumar, Stecher, & Tamura, 2016) and aligned with



homologous sequences deposited in the GenBank (Table S3, Appendix S3) using the ClustalW algorithm (Thompson, Higgins, & Gibson, 1994). The total number of haplotypes and haplotype and nucleotide diversities of the Mediterranean sequences (with associated standard deviations) were estimated using DnaSP v.5.1 (Librado, & Rosaz, 2009). The genetic diversity data of the global populations were taken from literature (see results). The relationship between haplotype and nucleotide diversity of each population was plotted using the R software package (R Core Team, 2008) to investigate GWS population diversity history. Phylogenetic relationships and haplotype genealogies were inferred using HapView (Salzburger, Ewing, & von Haeseler, 2011). Maximum likelihood (ML) clustering was constructed using the DNAML program in PHYLIP v.3.695 (Felsenstein, 2005), run in HapView. The best evolutionary model used in the phylogenetic analyses was inferred with JModelTest 2.1.1 (Darriba, Taboada, Doallo, & Posada, 2012), according to the Akaike Information Criteria (AIC; Akaike, 1974). Subsequently, the Hasegawa-Kishino-Yano model (Hasegawa, Kishino, & Yano, 1985) with the discrete Gamma distribution (0.8) and allowing for a proportion of invariant sites (0.4) to exist (HKY85+G+I) was selected as the best-fit model.

Since the historical Mediterranean sequence alignment (515bp) was shorter than most of the CR sequences deposited in the GenBank, a ML haplotype network was also reconstructed using a longer sequence alignment of 828bp based on haplotypes available in GenBank obtained from global contemporary GWS populations and specimens, including four Mediterranean sequences previously reported (Table S3, Appendix S3). A comparison between the topologies of the two haplotype networks permitted a test of the potential loss of informative sites in the shorter sequence alignment and the possible effects this might have on the reconstructed phylogenetic relationships.


### References of the supplementary methods

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19(6), 716–723. <https://doi:10.1109/TAC.1974.1100705>
- Allentoft, M. E., Collins, M., Harker, D., Haile, J., Oskam, C. L., Hale, M. L., ... Bunce, M., (2012). The half-life of DNA in bone: measuring decay kinetics in 158 dated fossils. *Proceedings of the Royal Society B: Biological Sciences*, 279(1748), 4724–4733. <https://doi:10.1098/rspb.2012.1745>
- Cooper, A., & Poinar, H. N. (2000). Ancient DNA: do it right or not at all. *Science*, 289(5482), 1139. <https://doi:10.1126/science.289.5482.1139b>
- Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, 9, 772. <https://doi:10.1038/nmeth.2109>
- Felsenstein, J. (2005). *PHYLIP (Phylogeny Inference Package) version 3.6*. University of Washington, Seattle. Retrieved from <http://evolution.genetics.washington.edu/phylip.html>.

- Gibbon, V. E., Penny, C. B., Štrkalj, G., & Ruff, P. (2009). Brief Communication: Minimally Invasive Bone Sampling Method for DNA Analysis. *American Journal of Physical Anthropology*, 139(4), 596–599. <https://doi:10.1002/ajpa.21048>
- Hansen, A. J., Willerslev, E., Wiuf, C., Mourier, T., & Arctander, P. (2001) Statistical Evidence for Miscoding Lesions in Ancient DNA Templates. *Molecular Biology and Evolution*, 18(2), 262–265. <https://doi:10.1093/oxfordjournals.molbev.a003800>.
- Hasegawa, M., Kishino, H., & Yano, T. (1985). Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution*, 22(2), 160–174.
- Kemp, B. M., & Smith, D. G. (2005). Use of bleach to eliminate contaminating DNA from the surface of bones and teeth. *Forensic Science International*, 154(1), 53–61. <https://doi:10.1016/j.forsciint.2004.11.017>
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7 Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, 33(7), 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Librado, P., & Rosaz, J. (2009). DNAsp v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25(11), 1451–1452. <https://doi:10.1093/bioinformatics/btp187>
- R Core Team. (2008). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Riccioni, G., Landi, M., Ferrara, G., Milano, I., Cariani, A., Zane, L., ... Tinti, F. (2010). Spatio temporal population structuring and genetic diversity retention in depleted Atlantic bluefin tuna of the Mediterranean Sea. *Proceedings of the National Academy of Sciences of the United States of America*, 107(5), 2102–2107. <https://doi:10.1073/pnas.0908281107>
- Salzburger, W., Ewing, G. B., & von Haeseler, A. (2011). The performance of phylogenetic algorithms in estimating haplotype genealogies with migration. *Molecular Ecology*, 20(9), 1952–1963. <https://dx.doi.org/10.1111/j.1365-294X.2011.05066.x>
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22(22), 4673–4680. <https://doi:10.1093/nar/22.22.4673>
- Untergrasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012 Primer3 - new capabilities and interfaces. *Nucl. Acids Res.* **40**, e115. <https://doi:10.1093/nar/gks596>

## Appendix S2 Supplementary figures


**Figure S1.** a) Jaws of the Great White Shark archived in the Museum of Comparative Anatomy of the University of Bologna. b, c) The cover and original illustrative note from Antonio Alessandrini (1854) “Catalogo degli Oggetti e Preparati più interessanti del Gabinetto D’Anatomia Comparata della Pontificia Università di Bologna dalla sua Fondazione all’ Ottobre del 1852”. The jaws were prepared from a GWS individual (TL = 473 cm) collected in the Adriatic Sea in 1827 and displayed to the public at the University of Bologna. d) Original cover from the publication of Ricciardi (1721) Pontificia Università di Bologna, Italy.




**Figure S2.** Illustration of the sampling procedures for the museum specimens of Great White Shark. a: Tooth pulp extraction from the crown. b-d: Drill sampling from dried jaw and restoration of the holes.




**Figure S3.** Results of the *in silico* PCR simulation with primer pairs designed for the amplification of the Control Region (CR) of *Carcharodon carcharias* (see Table S2, Appendix S3). The annealing positions of the primers with respect to the starting position of the gene on a reference mitochondrial genome (GenBank accession number NC\_022415) was shown in brackets. The length in base pairs (bp) of the amplified fragments are marked in yellow.



**Figure S4.** Maximum likelihood haplotype networks of worldwide *Carcharodon carcharias* population samples reconstructed using the 516bp (a) and the 828bp (b) CR sequence alignments, respectively. Acronyms are given in Table S2; Appendix S3. Numbers inside circles indicates the number of GWS individuals bearing the CR haplotype. Small blue dots indicate single nucleotide substitutions. Numbers outside the circles indicates the gross number of nucleotide substitutions separating the two haplogroups.



**Figure S5.** Bayesian divergence time tree of populations of *Carcharodon carcharias* inferred using the 516bp dataset of control region sequences. High posterior density (HPD 95%) in blue bars. Abbreviations are given in Table S2, Appendix S3.



### Appendix S3 Supplementary tables

**Table S1.** List of the 18 historical specimens of Mediterranean Great White Sharks, *Carcharodon carcharias* collected from museum and private archives.

Specimen code	GenBank Acc.No.	Geographic area	Sampling year/period	Tissue type*	Source §	Museum/Archive, Biological and Collecting Data
TRCC01AD	MN718579	Adriatic Sea	1906	V, T	1	Cat. No. 1182. Sex: female. Collecting site: Quarnero Gulf (Istria, Croatia). Collecting date: 29 May 1906. Donor: A. Morin
GECC01LI	MN718580	Ligurian Sea	1935	J	2	Cat. No. C.E. 32695. Sex: unknown. Collecting site: Riva Trigoso, Sestri Levante (Genoa, Italy). Collecting date: 03 July 1935
GECC02LI	MN718581	Ligurian Sea	1930s	J	2	Cat. No. C.E. 31916. Sex: unknown. Collecting site: unknown. Collecting date: 17 March 1933. Donor: E. Olivieri
GECC03LI	MN718582	Ligurian Sea	1958	J, T	2	No detailed data are available
LICC01LI	MN718583	Ligurian Sea	1950s	J	3	No detailed data are available
BOCC01AD	MN718584	Adriatic Sea	1823	J	4	Cat. No. 811, Catalogue Alessandrini. 1823 Sex: unknown. Collecting site: unknown. Collecting date: unknown. Additional info: mouth extension of 1.15 m
BOCC02AD	MN718585	Adriatic Sea	1827	J	4	Cat. No. ACP 114*; 1216 Catalogue Alessandrini. Sex: unknown. Collecting site: unknown. Collecting date: unknown. Additional info: mouth extension of 1.80 m
FICC01LI	MN718586	Ligurian Sea	1891	J	5	Cat. No. 6032; <i>Carcharodon rondeletii</i> M.H., 2775. Sex: female. Collecting site: Monterosso (Spezia, Italy). Collecting date: 10 December 1891. Additional info: Total length ~600 cm; Weight ~600 Kg. Donor: S.H. Giglioli.
FICC02LI	MN718587	Ligurian Sea	1879	V, T	5	Cat. No. 5983. Sex: unknown. Collecting site: Viareggio (Lucca, Italy). Collecting date: unknown.
MOCC01LI	MN718588	Ligurian Sea	1883	J, T	6	Cat. No. 50; <i>Carcharodon rondeletii</i> M.H. Sex: Male. Collecting site: Portofino (Genoa, Italy). Collecting date: January 1883.
PACC01TI	MN718589	Tyrrhenian Sea	1980s	J	7	No detailed data are available
PACC02TI	MN718590	Tyrrhenian Sea	1980s	J	7	No detailed data are available



PACC03TI	MN718591	Tyrrhenian Sea	1980s	J	7	No detailed data are available
FACC01TI	MN718592	Tyrrhenian Sea	1980s	V, M	8	Sex: unknown. Collecting site: Favignana (Trapani, Italy). Collecting date: unknown.
FACC02TI	MN718593	Tyrrhenian Sea	1980s	M	8	Sex: unknown. Collecting site: Favignana (Trapani, Italy). Collecting date: unknown.
FACC03TI	MN718594	Tyrrhenian Sea	1980s	V	8	Sex: unknown. Collecting site: Favignana (Trapani, Italy). Collecting date: unknown.
FACC04TI	MN718595	Tyrrhenian Sea	1980s	V	8	Sex: unknown. Collecting site: Favignana (Trapani, Italy). Collecting date: unknown.
FACC05TI	MN718596	Tyrrhenian Sea	1980s	M	8	Sex: unknown. Collecting site: Favignana (Trapani, Italy). Collecting date: unknown.

---

\* V: cartilage, vertebrae; T: osteodentine, tooth; J: cartilage, jaws; M; dried skeletal muscle

§ 1: Civic Museum of Natural History of Trieste; 2: Civic Museum of Natural History of Genova "Giacomo Doria"; 3: Regional Agency for Environmental Protection of Tuscany, ARPAT Livorno; 4: Museum of Comparative Anatomy, University of Bologna; 5: Museum of Natural History of Firenze "La Specola"; 6: University Museum of Natural History and Instrumentation of Modena; 7: Museum of Zoology of Palermo "P. Doderlein"; 8: Favignana Tuna Trap

**Table S2.** List of primer pairs (F: forward primer; R: reverse primer) designed for the PCR amplification and sequencing of the CR gene fragments from the historical DNA of Great White Shark.

---

Primer	5'>3' sequence
GWCRA1F:	TGACCTTCACCTAATGGTATCACA
GWCRA1R*:	AAGTCTCTGTGAGTGGAAGGAA
GWCRA2F:	CCTTCACCTAATGGTATCACACTC
GWCRA2R*:	AAGTCTCTGTGAGTGGAAGGAA
GWCRB1F:	TTCCTTCCACTCACAGAGACTT
GWCRB1R:	CAAGGACTGAAGTGTTACAAGCA
GWCRB2F:	TTTATTCCTTCCACTCACAGAGAC
GWCRB2R:	GACGGAAATGCTGTAAAGG

---

\* these two primers have identical sequences

**Table S3.** The mtDNA control region haplotypes and/or sequences of modern Great White Shark deposited and retrieved from the GenBank.

<b>Geographic Origin</b>	<b>Acronym</b>	<b>N</b>	<b>Reference</b>	<b>GenBank Accession Number</b>
Australia	AUS	14	[9]	HQ414073 - HQ414086
Australia	AUS	12		
New Zealand	AUS	4	[7]	AY026196 - AY026224
South Africa	SA	13		
Northeastern Pacific	NEP	20	[10]	GU002302 - GU002321
Florida	NWA	2		
Mediterranean	MED	3	[17]	HQ540294 - HQ540298
Mediterranean	MED	1	[27]	JF715925
Northwestern Atlantic	NWA	11		
South Africa	SA	15	[28]	KC511601 - KC511626
South Africa*	SA	4*	[29]	KP058665 - KP058902*

\*During the analyses, the 238 unique sequences from Andreotti et al. [29], were collapsed in the unique four haplotypes observed and added to the final dataset. AUS: Australia/New Zealand; NEP: Northeastern Pacific; MED: Mediterranean; SA: South-Africa; NWA: Northwestern Atlantic.

**Table S4.** Fossil data downloaded from PaleoDB (<https://paleobiodb.org/#/>) using the species-specific taxon “*Carcharodon carcharias*” and synonyms. References downloadable from the paleoDB database. The records, especially those from old references, without description of the fossil and without pictures, and those records with misidentified fossils (e.g. *Isurus* spp. or *Carcharodon* spp. identified as *C. carcharias*), were considered dubious and subsequently removed in order to create a fossil dataset of reliable *C. carcharias* records.

PaleobioDB N°	coll_no	PaleobioDB Epoch	early_age	late_age	Ref_no	longitude	latitude
465463	45458	Pliocene	7.246	5.333	12282	-115.175552	28.11472
465581	13079	Pliocene	7.246	5.333	12182	-74.719719	-15.5808
506112	51328	Pliocene	5.333	3.6	13672	141.603058	-38.3619
506149	51335	Pliocene	5.333	3.6	13672	141.944443	-37.7283
506450	46068	Pliocene	5.333	3.6	13672	147.966666	-37.8547
506468	51405	Pliocene	3.6	1.806	13672	138.609726	-34.8333
506557	51414	Pliocene	5.333	2.588	13672	148.083328	-39.9833
518397	52644	Pleistocene	1.806	0.781	14149	-52.326389	-32.3883
520948	28039	Pliocene	3.6	3	14399	174.28334	-39.5833
533364	20400	Pleistocene	1.8	0.3	1960	-82.5	27.7
558936	58089	Pleistocene	1.8	0.3	15601	-80.811386	27.83028
593653	52582	Pliocene	5.333	3.6	18094	-76.817497	35.35972
634589	68271	Pliocene	5.333	3.6	19640	-71.5	-30.3333
639047	55535	Miocene	7.246	5.333	19852	-70.841667	-27.0808
645725	69730	Miocene	11.62	7.246	23394	-70.87944	-27.1392
645727	69731	Pliocene	5.333	3.6	23394	-70.87944	-27.1392
668275	72085	Miocene	15.97	11.608	24392	90.666664	25.16667
732439	78614	Miocene	15.97	11.608	26436	92.73333	22.88333
789236	20646	Pleistocene	0.126	0.0117	28773	-118.199997	34
807796	88328	Pliocene	3.6	2.588	29650	10.888611	43.67222
899319	100174	Pliocene	3.6	2.588	34371	-70.534447	-23.3575
981089	117471	Pliocene	5.333	3.6	37795	-122.407997	37.288
984829	118104	Pliocene	5.333	3.6	38036	-0.676944	38.085
1087691	136597	Pliocene	5.333	3.6	43697	14.004444	32.7325
1192384	154111	Pliocene	5.333	2.588	49963	13.3	-8.75
1192410	154114	Miocene	15.97	3.6	49966	-70.833336	-27.1333

1192476	154117	Miocene	7.246	3.6	49968	-112.291389	27.36806
1192505	154118	Pliocene	5.333	2.588	49969	-109.616669	23.16667
1221552	159914	Pliocene	3.6	2.588	52184	138.600006	-34.9167
1227809	161451	Pliocene	5.333	2.588	52598	139.645554	35.4475
1228090	161503	Pleistocene	2.588	0.781	52618	136.96666	36.76389
1234295	162457	Miocene	11.608	5.333	52569	-81.650002	30.33
1360649	184953	Pliocene	3.6	2.588	37795	-122.407501	37.29
1360660	184959	Pliocene	3.6	2.588	37795	-122.404999	37.31
1360661	184960	Pliocene	5.333	3.6	37795	-122.407997	37.288
1374190	187584	Pleistocene	0.126	0.0117	62969	-88.183891	30.24965
1374545	55759	Pliocene	3.6	2.588	62984	9.998611	44.82667
1388536	190889	Pleistocene	0.126	0.0117	33088	-117.902222	33.65722
1406990	188304	Pleistocene	0.126	0.0117	66045	-80.31115	33.10401
1431917	100313	Pliocene	5.333	3.6	62247	18.42804	-33.6651

**Table S5.** Fossil data from the Mediterranean area. For museal fossil teeth catalogued from 1 to 87 see S. Marsili (2006).

<b>n</b>	<b>Location</b>	<b>Age</b>	<b>Reference</b>	<b>Stratigraphic References</b>
1a	Salsomaggiore Terme, Parma, Emilia-Romagna (Italy)	Pliocene	Bianucci et al. 2002	Bianucci et al. 1998
2a	Guardamar del Segura, Alicante (Spain)	Pliocene	Adnet et al. 2009	
		Pliocene		
1	Terreti, Reggio Calabria: Calabria(Italy)	sup./Pleistocene ?	Seguenza, 1901	Gaetani et al., 1986.
2	Nasiti e S.Agata, Reggio Calabria: Calabria(Italy)	Pleistocene inf.	Seguenza, 1901	Lombardo
3	Cetona (vicinanze), Siena: Toscana (Italy)	Pliocene	Principi, 1920	
4	Castiglione del Lago (a Ovest di), Perugia: Umbria (Italy)	Pliocene	Principi, 1920	
5	Città della Pieve, Perugia: Umbria (Italy)	Pliocene	Principi, 1920	
6	Imola (varie località): Emilia Romagna (Italy)	Pliocene	Collection Scarabelli (De Stefano, 1911)	
7	Imola (varie località): Emilia Romagna (Italy)	Pliocene	Collection Scarabelli (De Stefano, 1911)	
8	Castell'Arquato, Piacenza: Emilia Romagna (Italy).	Pliocene	De Stefano, 1912	
9	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Landini, 1977	
10	Punta Ristola, Capo di Leuca, Lecce: Puglia (Italy)	Pliocene middle	Menesini, 1968	
11	Torre del Lago, Lucca: Toscana (Italy)	Pleistocene	Ghelardoni, 1956	
12	Calanna, Reggio Calabria: Calabria (Italy)	Pliocene sup.	De Stefano, 1901	Barrier et al., 1986.
13	Castell'Arquato, Piacenza: Emilia Romagna (Italy).	Pliocene	Carraroli, 1897	
		Pliocene sup-		
14	Rometta, Messina: Sicilia (Italy)	Pleistocene inf.	Seguenza, 1900	For Violanti, 1989
15	Milazzo, Messina: Sicilia (Italy)	Pliocene sup ?	Seguenza, 1900	
16	Guardamar del segura, Alicante: Valezia (Spain)	Pliocene inf.	Mora Morote, 1996	
17	Ruvo del Monte, Potenza: Basilicata (Italy)	Pliocene	Bassani, 1901	
18	Taranto: Puglia (Italy)	Pleistocene	Bassani, 1905	
19	Volpedo, Alessandria: Piemonte (Italy)	Pliocene middle-sup	De Alessandri, 1895	Gabba, 82
20	Ruvo del Monte, Potenza: Basilicata (Italy)	Pliocene	Pasquale, 1903	
21	S.Agata , Reggio Calabria: Calabria(Italy)	Pleistocene inf.	Pasquale, 1903	Lombardo
22	Terreti, Reggio Calabria: Calabria (Italy)	Pliocene inf.-middle	Pasquale, 1903	Gaetani et al., 1986.
23	Reggio, Nasiti, Reggio Calabria: Calabria (Italy)	Pliocene	Pasquale, 1903	

24	Capo di Leuca, Lecce: Puglia (Italy)	Pliocene middle	Pasquale, 1903	Bossio et al., 2001
25	San Demetrio Corone, Cosenza: Calabria (Italy)	Pleistocene	Pasquale, 1903	
26	Tabiano, Piacenza: Emilia Romagna (Italy);.	Pliocene inf.	De Stefano, 1912	
27	Calanna, Reggio Calabria: Calabria (Italy)	Pliocene sup.	Pasquale, 1903	Barrier et al., 1986. Iannone et al., 1979; Cherubini et al., 1996; Pomar et al., 2001.
28	Matera: Basilicata (Italy)	Pliocene sup.- Pleistocene inf.	Pasquale, 1903	
29	Sestano e Medano, Siena: Toscana (Italy)	Pliocene inf.	Manganelli & Spadini, 2003	
30	San Quirico d'Orcia, Giustrigona, Terre Rosse e I Sodi, Siena: Toscana (Italy)	Pliocene middle	Manganelli & Spadini, 2003	
31	Allerona, Terni: Umbria (Italy)	Pliocene middle	Bellocchio et al., 1991	
32	Guardamar del segura, Alicante: Valezia (Spain)	Pliocene inf.	Mora Morote, 1996 De Stefano, 1910: Collection Lawley di Bologna	
33	Colline Toscane (Italy)	Pliocene	De Stefano, 1910: Collection Lawley di Bologna	
34	Colline Toscane (Italy)	Pliocene	Lawley di Bologna	
35	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection Lawley of Pisa	
36	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
37	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
38	Val d'Orcia: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze	
39	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
40	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
41	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
42	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
43	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
44	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
45		Pliocene	Collection fossil teeth of Firenze	
46	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
47	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
48	San Frediano, Pisa: Toscana (Italy)	Pliocene middle	Collection fossil teeth of Firenze	
49	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	
50	Colline Senesi: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze	
51		Pliocene	Collection fossil teeth of Firenze	
52	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze	

53	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
54	San Vivaldo, Firenze: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
55	San Quirico d'Orcia, Siena: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
56	Colline Senesi: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
57	Chiusi, Siena: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
58		Pliocene	Collection fossil teeth of Firenze
59	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
60	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
61	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
62	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
63	Siena: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
64	San Quirico d'Orcia, Siena: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
65	Bagni di Casciana, Pisa: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
66	Orciano, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
67	San Quirico d'Orcia, Siena: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
68	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
69	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection fossil teeth of Firenze
70	Santa Luce, Pisa: Toscana (Italy)	Pliocene	Collection fossil teeth of Firenze
71	Rione Castellana, Palermo: Sicilia (Italy)	Pleistocene	Gemellaro, 1913
72	Pradalbino (varie loc.), Bologna: Emilia Romagna (Italy)	Pliocene inf.-middle	Vinassa de Regny, 1899
73	Pradalbino (varie loc.), Bologna: Emilia Romagna (Italy)	Pliocene inf.-middle	Vinassa de Regny, 1900
74	Orciano; Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Bassani, 1901
75	San Quirico d'Orcia (dintorni), Siena: Toscana (Italy)	Pliocene inf.-middle	Simonelli, 1880
76	unknown locality	Pliocene	Accademia Fisiocritici of Siena
77	unknown locality	Pliocene	Accademia Fisiocritici of Siena
78	? Monte Follonico(1), Siena: Toscana (Italy)	Pliocene	Accademia Fisiocritici of Siena
79	Monte Follonico, Siena; ? Volterra(1), Pisa: Toscana (Italy)	Pliocene	Accademia Fisiocritici of Siena
80	Pod. Casabianca (Trequanda), Siena: Toscana (Italy)	Pliocene	Accademia Fisiocritici of Siena
81	Medane (Asciano), Siena: Toscana (Italy)	Pliocene	Accademia Fisiocritici of Siena
82	Volterra, Pisa: Toscana (Italy)	Pliocene inf.-middle	Collection Lawley of Pisa
83	Pontedera, Pisa: Toscana (Italy)	Pliocene	Collection Lawley of Pisa



84	Piacentino: Emilia Romagna (Italy)	Pliocene	Collection Lawley of Pisa
85	Terricciola, Pisa: Toscana (Italy)	Pliocene	Collection Lawley of Pisa
86	Val di Cecina: Toscana (Italy)	Pliocene	Collection Lawley of Pisa
87	Peccioli, Pisa: Toscana (Italy)	Pliocene	Collection Lawley of Pisa

**Table S6.** Control Region diversity of the Mediterranean and global populations of *Carcharodon carcharias*.

Population	Acronym	N	Nh	$h \pm SD$	$\pi \pm SD$	Reference
Mediterranean	MED	22*	3	0.091 $\pm$ 0.124	0.0002 $\pm$ 0.0007	This work
Australia-New Zealand	AUS	94	14	0.880 $\pm$ 0.015	0.0085 $\pm$ 0.0045	[Blower et al., 2012]
Northeastern Pacific	NEP	59	20	0.790 na	0.0013 $\pm$ 0.0009	[Jorgensen et al., 2010]
South Africa	SA	34	15	0.723 na	0.0059 na	[O'Leary et al. 2015]
South Africa	SA*	238	4	0.205 $\pm$ 0.033	0.0027 $\pm$ 0.0005	[Andreotti et al. 2016]
North West Atlantic	NWA	44	12	0.749 na	0.0030 na	[O'Leary et al. 2015]

Acronyms are given in Table S3.

N: number of individuals analysed; Nh: number of haplotypes;  $h$ : haplotype diversity;  $\pi$  nucleotide diversity; SD: Standard deviation

\* this sample included sequence records from GenBank (Acc. Num. JF715925, HQ540294, HQ540295, HQ540296).

**Table S6.** Best model selection based on Bayes factors using alternative primary calibration (estimated divergence between *Carcharodon carcharias* and *Lamna nasus*). logML\_GSS: log marginal likelihood from generalized stepping stone model, BF\_GSS: Bayes factors calculated using the logML\_GSS.

Model	Dataset	Node1	Node2	logML_GSS	logBF_GSS
MED Pliocene divergence	828bp	<i>C.carcharias/L.nasus</i> divergence	MED/Pacific		
		Mean: 46 Ma, SD: 1.0 Ma	Mean: 3.0 Ma, SD: 0.3 Ma	-3210.6566	0 (Best)
MED Pleistocene divergence	828bp	<i>C.carcharias/L.nasus</i> divergence	MED/Pacific		
		Mean: 46 Ma, SD: 1.0 Ma	Mean: 0.4Ma, SD: 0.15 Ma	-3215.5172	4.8606
MED Pliocene divergence	516bp	<i>C.carcharias/L.nasus</i> divergence	MED/Pacific		
		Offset: 46 Ma, SD: 1.0 Ma	Mean: 3.0 Ma, SD: 0.3 Ma	-1905,6997	0
MED Pleistocene divergence	516bp	<i>C.carcharias/L.nasus</i> divergence	MED/Pacific		
		Offset: 46 Ma, SD: 1.0 Ma	Mean: 0.4Ma, SD: 0.15 Ma	-1914,4034	8.7037



Great White Shark, Dyer Island, Gansbaai, South Africa, 2011, Photo: Agostino Leone

1286x965mm (72 x 72 DPI)



Lateral view of a Great White Shark historical jaw from Alessandrini Collection of the University of Bologna.  
Photo: Mr. Leonardo Piol



Frontal view of a Great White Shark historical jaw from Alessandrini Collection of the University of Bologna.  
Photo: Mr. Leonardo Piol