ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Normalized Difference Vegetation Index versus Dark Green Colour Index to estimate nitrogen status on bermudagrass hybrid and tall fescue

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:
Caturegli, L., Gaetani, M., Volterrani, M., Magni, S., Minelli, A., Baldi, A., et al. (2020). Normalized Difference Vegetation Index versus Dark Green Colour Index to estimate nitrogen status on bermudagrass hybrid and tall fescue. INTERNATIONAL JOURNAL OF REMOTE SENSING, 41(2), 455-470 [10.1080/01431161.2019.1641762].

Availability:
This version is available at: https://hdl.handle.net/11585/716429 since: 2021-11-25

Published:

DOI: http://doi.org/10.1080/01431161.2019.1641762

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/).
When citing, please refer to the published version.

(Article begins on next page)

11 October 2026

Title Normalized Difference Vegetation Index versus Dark Green Color Index to estimate nitrogen status on bermudagrass hybrid and tall fescue.

Short title: NDVI vs DGCI to estimate N on two turfgrass species

Authors: Lisa Caturegli[*1], Monica Gaetani[1], Marco Volterrani[1], Simone Magni[1], Alberto Minelli[2], Ada Baldi[3], Giada Brandani[3], Marco Mancini[3], Anna Lenzi[3], Simone Orlandini[3], Filippo Lulli[4], Claudia de Bertoldi[4], Marco Dubbini[5], Nicola Grossi[1]

[1] Department of Agriculture, Food and Environment, University of Pisa, Pisa, Italy,

[2] Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy

[3] Department of Agriculture, Food, Environment and Forestry, University of Florence, Florence, Italy

[4] Turf Europe Srl, Livorno, Italy

[5] Department of History, Cultures and Civilizations, University of Bologna, Bologna, Italy;

*Corresponding author Email: lisa.caturegli@gmail.com

Abbreviations:

DGCI Dark Green Color Index

GPS Global Positioning System

HSB Hue Saturation Brightness

NDVI Normalized Difference Vegetation Index

PA Precision Agriculture

PTM Precision Turfgrass Management

RGB Red Green Blue

UAS Unmanned Aerial Systems

UAV Unmanned Aerial Vehicle

Keywords:

Cynodon dactylon x transvaalensis, Schedonorus phoenix, Turfgrass, Color, Quality, Unmanned Aerial Vehicle.

Abstract

In recent years digital sensors have been successfully integrated on board Unmanned Aerial Vehicles (UAV) to assess crop vigor, vegetation coverage, and to quantify the “greenness” of foliage as indirect measurements of crop nitrogen status. The classical approach of precision agriculture has involved the use of multispectral sensors onboard UAV and the development of numerous vegetation indices associated with vegetation parameters, such as the mostly used Normalized Difference Vegetation Index (NDVI). However, the main negative issue when dealing with multi and hyper-spectral reflectance measuring tools is their high cost and complexity from the operational point of view. As a low-cost alternative, vegetation indices derived from Red Green Blue (RGB) cameras have been employed for remote sensing assessment, providing data on different stress conditions and species. Digital images record information as amounts of RGB light emitted for each pixel of the image; however, the intensity of red and blue will often alter how green an image appears. To simplify the interpretation of digital color data, recent studies have suggested converting RGB values to the more intuitive Hue, Saturation, and Brightness (HSB) color spectrum, and then into a single measure of dark green color, the Dark Green Color Index (DGCI). In this study NDVI acquired by a ground-based handheld crop sensor and by a multispectral camera mounted on board a UAV have been compared with DGCI calculated from images taken with a commercial digital camera on board a UAV, trying to quantify the color of turfgrass that had received different nitrogen (N) rates.

The objectives of the trial were to study an affordable easy-to-use tool evaluating the relationship among NDVI, DGCI and leaf nitrogen content on turfgrass.

Introduction

Nitrogen fertilization on turfgrasses is one of the factors that most influence physiological and aesthetic aspects (Volterrani et al. 2005; Perry and Davenport 2007; Samborski, Tremblay, and Fallon 2009; Caturegli et al., "Monitoring turfgrass", 2014; Caturegli et al., "Turfgrass spectral reflectance", 2014; Grossi et al. 2016). Thus, nitrogen (N) represents an important nutrient that contributes to maintain green color, density, recovery from drought diseases, and a general good turfgrass quality (Walters and Bingham 2007; Dordas 2008; Magni et al. 2014; Caturegli et al. 2016).

However, the excessive fertilization of N wastes fertilizers and leads to pollution of ground and surface water, not improving the quality of the turf (Bell and Xiong 2008; Bremer et al. 2011; Rhezali and Lahlali 2017). To avoid over-fertilization, site-specific nutrition management brought significant environmental and economic benefits (Huang et al. 2008). Indeed, a precise analysis of the plant nitrogen status is important to determine the amount of nitrogen fertilizer the plant really needs (Corwin and Lesch 2005; Li et al. 2015).

Previous studies have focused on implementation of indirect sensing tools (chlorophyll meters, reflectance measurements, color analysis) to try to obtain an almost optimal quality by reducing the N inputs and the loss N to a minimum (Rorie, Purcell, Mozaffari et al. 2011; Caturegli, Casucci et al. 2015; Caturegli, Grossi et al. 2015; Caturegli et al. 2016).

These concepts are the basis of Precision Agriculture (PA), which aims to obtain detailed site-specific information by mapping the variation in important soil and plant properties in order to allow better site-specific management. Inputs such as water, fertilizers and pesticides are applied only where, when and in the amount needed by plant (Caturegli et al., "Turfgrass spectral reflectance", 2014). Related to PA is Precision Turfgrass Management (PTM) that is useful to monitor pests, fertilization, salinity stress and irrigation deficiency on turfgrass (Carrow et al. 2010; Krum, Carrow, and Karnok 2010). The approach of PA implied the combined use of multispectral

78 sensors and vegetation indices associated with vegetation parameters (Trenholm, Carrow, and
79 Duncan 1999; Jiang and Carrow 2007; Vergara-Díaz et al. 2016). Thus, vegetation indices were
80 calculated by combining various reflectance bands of the spectrum and correlated with relevant
81 turfgrass canopy parameters. Among the indices, the Normalized Difference Vegetation Index
82 (NDVI) is the most widely used as reflectance-based plant stress indicator (Hansen and Schjoerring
83 2003; Johnsen et al. 2009; Aguilar et al. 2012; Barton 2012; Fensholt and Proud 2012; Rhezali and
84 Lahlali 2017). It is based on the relationship between the absorption of visible light and resilient
85 reflectance of near-infrared light to the chlorophyll in vegetation (Bell et al. 2004; Caturegli,
86 Casucci et al. 2015). The NDVI value ranges from -1 to 1, with higher values indicating greater
87 plant health, and correlates positively with turfgrass quality (Trenholm, Carrow, and Duncan 1999;
88 Fitz-Rodriguez and Choi 2002; Leinauer et al. 2014). This index is also influenced by differences in
89 species, environmental stresses, fertilization and pest injuries (Xiong et al. 2007; Bremer et al.
90 2011; Caturegli, Grossi et al. 2015). It can be obtained with hand-held ground-based instruments
91 (Graeff and Claupein 2003; Ma, Morrison, and Dwyer 1996) and aerial vehicle-mounted sensors
92 (Bausch and Duke 1996; Blackmer et al. 1996; Scharf and Lory 2009; Rorie, Purcell, Karcher et al.
93 2011). In recent years, digital sensors have been successfully integrated on board Unmanned Aerial
94 Vehicles (UAV) to assess crop vigor, vegetation coverage, and to quantify the “greenness” of
95 foliage as indirect measurements of crop N status (White et al. 2012; Andrade-Sanchez et al. 2014).
96 Furthermore, small commercial Unmanned Aerial Systems (UAS) (< 50 kg) (Laliberte and Rango
97 2011) have been available for PA for environmental and agricultural applications (Gupta et al.
98 2013; Zhang and Kovacs 2012; Caturegli et al. 2016). However, the main negative issue when it
99 comes to multi and hyper-spectral reflectance measuring tools is their high cost and complexity.
100 Vegetation indices derived from Red-Green-Blue (RGB) cameras have been employed for remote
101 sensing assessment, as a low-cost alternative (Vergara-Díaz et al. 2016). This method may provide
102 data on different stress conditions in different crops (Casadesús et al. 2007; Casadesus and Villegas
103 2014; Zhou et al. 2015) and turfgrass (Karcher and Richardson 2003; Karcher and Richardson

104 2013). Digital images are composed by pixels that record information as amounts of RGB light
105 emitted. However, the greenness of an image can be often altered by the intensity of red and blue.
106 To simplify the interpretation of data, Karcher and Richardson (2003) suggested converting RGB
107 values to the more intuitive Hue, Saturation, and Brightness (HSB) based on human perception of
108 color. Working with quality of turfgrass in response to N fertilizer, Karcher and Richardson (2003)
109 processed HSB values into a single measure of dark green color, the Dark Green Color Index
110 (DGCI).
111 This method proposed by (Karcher and Richardson 2003) may represent a proper alternative to the
112 spectroradiometric approaches that involves the use of NDVI from aerial platforms and from
113 ground-based measurements (Vergara-Díaz et al. 2016). To facilitate the DGCI acquisition,
114 recently, also a smartphone application called FieldScout GreenIndex+ Turf (Spectrum
115 Technologies, Inc., Aurora, IL, USA) (Spectrum Technologies, Inc. 2018) has been developed and
116 tested (O'Brien 2017; Xiang et al. 2017; Xiang et al. 2018) The application (APP) captures images
117 with a smartphone or tablet, calculates the DGCI, and shows a turfgrass quality visual rating
118 (Karcher and Richardson 2003).
119 The aim of this research was to study an affordable easy-to-use tool evaluating the relationship
120 among NDVI, DGCI and leaf nitrogen content on turfgrass. Trying to quantify the color of turfgrass
121 that had received different N rates, NDVI acquired by a ground-based handheld crop sensor and by
122 a multispectral camera mounted on board a UAV have been compared with DGCI calculated from
123 images taken with a commercial digital camera on board a UAV.
124

Materials and Methods

The trial was carried out in July 2017 in S. Piero a Grado, Pisa, at the Centre for Research on Turfgrass for the Environment and Sports (CeRTES) of the Department of Agriculture, Food and Environment of the University of Pisa (43°40'N, 10°19'E, 6 metres above sea level (m. a. s. l.).

The turfgrasses selected for the study were a mature turfgrass stands of the warm-season bermudagrass hybrid (*Cynodon dactylon* [L.] Pers. (Linnaeus Persoon) variety dactylon x *Cynodon transvaalensis* Burtt-Davy) cultivar (cv) 'Patriot' and the cool-season tall fescue (*Schedonorus phoenix* [Scop.] (Scopoli) Holub) cv 'Grande'.

The swards were all established on a calcaric fluvisoil (Coarse-silty, mixed, thermic, Typic Xerofluvents) with pH 7.8 and 18 g kg^{-1} of organic matter.

No fertilizer had been applied to the turfgrass before the trial started. In order to create a linear nitrogen gradient, on June 2017 fertilization was carried out applying ammonium sulphate (21-0-0) with a rotary spreader (ICL Specialty Fertilizers AccuPro 2000, Ipswich, UK).

The experimental designs were:

a) For tall fescue 8 nitrogen rates were applied, from 0 to 210 kg ha^{-1} of N with increases of 30 kg ha^{-1} (0 kg ha^{-1}, 30 kg ha^{-1}, 60 kg ha^{-1}, 90 kg ha^{-1}, 120 kg ha^{-1}, 150 kg ha^{-1}, 180 kg ha^{-1}, 210 kg ha^{-1} of N). The plot size was 3 m × 3 m, with 3 replications.

b) For bermudagrass hybrid, which tolerates higher doses of fertilizer, 11 nitrogen rates were applied, from 0 to 300 kg ha^{-1} of N with increases of 30 kg ha^{-1} (0 kg ha^{-1}, 30 kg ha^{-1}, 60 kg ha^{-1}, 90 kg ha^{-1}, 120 kg ha^{-1}, 150 kg ha^{-1}, 180 kg ha^{-1}, 210 kg ha^{-1}, 240 kg ha^{-1}, 270 kg ha^{-1}, 300 kg ha^{-1} of N). The plot size was 3 m × 3 m, with 3 replications.

Extreme N rates were applied in order to reach the nitrogen saturation level for both species, regardless of the agronomic drawbacks to the turfgrasses.

After the fertilization, an irrigation of 5 mm was applied. During the trial period a turf height of 2.0 cm was maintained by mowing with a walk-behind reel mower (John Deere 20SR7, Moline IL,

USA) with clippings removal. In the entire experimental area, in order to evaluate nitrogen fertilization as the only variability source, identical and maintenance practices were applied. Irrigation was applied as needed to avoid wilt, in order to maintain the soil moisture constant and equal in all areas. During the trial no weed or pest control was necessary.

On each of the two experimental areas proximity and remote sensed readings were acquired starting from the unfertilized control to the highest nitrogen rate in each plot.

The ground-based instrument used to acquire NDVI values was a Handheld Crop Sensor (HCS) (GreenSeeker, Model HSC-100, Trimble Navigation Unlimited, Sunnyvale, CA) while the remote sensed readings were collected with a UAV which was a VTOL (Vertical Take Off and Landing) DJI s900 hexacopter (DJI, Shenzhen, China) equipped with a digital commercial camera Sony Nex 5 (Sony, Surrey, United Kingdom) and a lightweight multispectral sensor MAIA S2 (SAL Engineering, Modena Italy; EOPTIS, Trento, Italy). Spectral measurements (proximity and aerial) were taken on 6 July 2017 between 11:30 AM (ante meridiem) and 1:30 PM (post meridiem) (local time), in complete absence of clouds. The weather parameters of July 2017 were as follows: average air temperature 25 °C, average relative humidity 60%; July average of the noon Photosynthetic Photon Flux Density 1,482 μmol m^{-2} s^{-1}; average wind speed 6 km h^{-1}. Each ground-based measurement was geo-referenced to sub-meter accuracy with a Global Positioning System (GPS) receiver Leica 1200 in Real Time Kinematic, in order to find the exact position on the UAV images and to compare data acquired with the two systems (Caturegli, Casucci et al. 2015; Caturegli, Grossi et al. 2015; Caturegli et al. 2016).

Ground-based measurements

Proximity sensed measurements of spectral reflectance were acquired with a HCS at a height of 110 cm from the ground, thus monitoring a surface of about 2,000 cm^2 ($\varnothing$ = 50 cm). The HCS has an active light source that makes readings unaffected by sunlight (Bell, Kruse, and Krum 2013). Reflectance was measured in the red region at 660 nm, and in the near infrared region of the

spectrum at 780 nm. The output is directly provided as NDVI, which is calculated using the equation:

$$NDVI = ((NIR) - R)/((NIR) + R) \quad (1)$$

where R is the reflectance in the red band and NIR is the reflectance in the near-infrared band.

In the same day and in the same area also the following parameters were studied:

- Color intensity (1 = very light green; 6 = acceptable green; 9 = very dark green): visual assessments (Morris and Shearman 2008);
- Turfgrass Quality: (1 = poor; 6 = acceptable; 9 = excellent): visual assessments (Morris and Shearman 2008);
- Total N content of leaves: samples of clippings were collected on each sampling area with a walk-behind reel mower from a surface of 0.5 m^2 (1.0 m × 0.5 m). Fresh clippings were put in a ventilated stove at 70 °C, dried to constant weight, and the total N was determined by the micro-Kjeldahl method (Bremner 1965);
- Plant water content (PWC): calculated as follows:

$$PWC\ (\%) = \frac{(FW) - (DW)}{(FW)} \times 100 \quad (2)$$

where FW is the leaf fresh weight and DW the leaf dry weight. Leaves were cut and quickly put into a plastic bag with hermetic closure. The bags were refrigerated and kept in the dark until arrival to the laboratory, where they have been weighed.

UAV flight and analysis of UAV derived imagery

The UAV system used for surveying was a DJI s900 hexacopter (Figure 1 (a)) with Global Navigation Satellite System (GNSS), with L1 code solution and a 3 axis accelerometer based stabilization system. The hexacopter was equipped with a digital commercial camera Sony Nex 5 (Sony, Surrey, United Kingdom) and a lightweight multispectral camera MAIA S2 (SAL Engineering, Modena Italy; EOPTIS, Trento, Italy) (Figure 1 (b)). The images were acquired at 90

200 m of altitude to guarantee a GSD (Ground Sample Distance) of less than 5 cm and a FOV (Field Of
201 View) of about 58 m × 43 m. The direction and altitude of the aircraft were controlled by the
202 rotation speed or by the direction of the propellers (Li et al. 2015). Real-time images, and other
203 information such as altitude and battery voltage, were transmitted to a ground monitor through a
204 radio link.
205 **Please insert Figure 1 near here**
206

UAS derived imagery NDVI

The UAS derived imagery NDVI was obtained using the UAV cited above equipped with a multispectral camera MAIA S2 (SAL Engineering, Modena Italy; EOPTIS, Trento, Italy), which features an array of nine sensors with 1.2 Megapixel resolution: specifically, one RGB color and eight monochrome sensors are available for analysis of the visible and near infra-red (VIS-NIR) spectrum from 390 nm to 950 nm, operating with a frame rate of 5 Hz per sensor. Each of the eight sensors is provided with a band-pass filter (Table 1). Global shutter technology is so such that all of the pixels in each sensor start to collect charge simultaneously, allowing images to be scanned in "one shot" for synchronized multiband measurements. The extremely fast exposure times of the nine global shutter complementary metal-oxide semiconductor (CMOS) sensors (up to 10^{-4} s) and the low travel speed (< 0.5 m s^{-1}) guarantee the absence of the blur effect. The images obtained were geometrically corrected with calibrated optics, and radiometrically corrected with the acquisition of the reflectance values of the incident light through a calibrated white panel. After the corrections, the 9 images for each shot were registered using the proprietary MAIA software based on photogrammetric method (Dubbini et al. 2017). Every pixel of the image contained coordinates and an NDVI value that was extracted using Quantum GIS (Geographic Information System) 2.18 software.

Please insert Table 1 near here

Dark Green Color Index (DGCI)

A common digital camera, Sony Nex 5 (Sony, Surrey, United Kingdom) was used to capture RGB images of the selected area. The Sony Nex-5 is a mirrorless interchangeable-lens camera, with the Advanced Photo System-Classic (APS-C) Exmor CMOS sensor and a maximum image resolution of 4,912 × 3,264 and a pixel size of 5 μm in both x and y directions (Remondino and Fraser 2006; Fryskowska et al. 2016). To reduce the effect of vibration during the flight and capture

clear images, the camera was mounted on a pan-tilt set which keeps the lens horizontal. In the same day as the ground NDVI readings and the NDVI by the multispectral camera, the digital camera recorded UAS derived imagery RGB images above the interested area, always in a zenithal plane. Images were taken with auto-focus, auto-white balance and an automatic exposure, and they were saved in Joint Photographic Experts Group (JPEG) format. Subsequently, images were analyzed with the open source Quantum GIS 2.18 software to extract the RGB values of the pixels where the NDVI values by the ground and by the multispectral camera were calculated. To simplify the interpretation of data, RGB values were converted into HSB values, using the method suggested by (Karcher and Richardson 2003), to finally calculate the DGCI. DGCI value is on a scale from 0 (very yellow) to 1 (dark green) (Rhezali and Lahlali 2017). DGCI was calculated as:

$$DGCI = [((Hue) - 60)/60 + (1 - (Saturation)) + (1 - (Brightness))]/3 \quad (3)$$

Statistical analysis

The correlations between the two different NDVI reading methods (ground-based sensing with a HCS and remote sensing with UAV) and DGCI were studied using CoStat software (CoHort, Monterey, CA, USA) and Pearson's correlation coefficients (r) were calculated in order to verify whether: (a) NDVI-ground data and NDVI-UAV were suitably correlated with DGCI obtained from RGB images captured by the digital camera on board a UAV; (b) UAV imagery with a low cost digital camera could be a diagnostic tool to identify variation in N status of turfgrass, comparable to a more expensive multispectral camera. Linear relationships were studied for the correlations showing statistically significant coefficients.

Results and discussion

Relationship between DGCI, NDVI and observed parameters

In bermudagrass hybrid, considering *r* among NDVI values obtained with the two different instruments (proximity sensed with the HCS GreenSeeker and remotely sensed with the multispectral camera MAIA mounted on board a UAV), and the measured parameters, the *r* values were highly significant. The *r* values ranged between 0.92 for PWC-NDVI of both the instruments and 0.97 for turfgrass quality-NDVI GreenSeeker. Comparing DGCI and all the measured parameters, the index was significantly correlated with color intensity, turfgrass quality and plant water content with *r* values ranging between 0.83 for color intensity and 0.84 for turfgrass quality and PWC (Table 2).

In tall fescue the correlations between NDVI obtained with GreenSeeker and with UAV and color intensity ($r = 0.96$ and $r = 0.95$) has showed higher *r* values than the same in bermudagrass hybrid *(*$r = 0.94$). Also PWC-NDVI (GreenSeeker and UAV) showed a degree of association significantly higher in tall fescue ($r = 0.98$) than bermudagrass hybrid ($r = 0.92$).

Furthermore, observing the correlations, the DGCI was highly correlated with all the measured parameters with *r* values ranging between 0.92 for DGCI-Quality and 0.98 for DGCI-PWC. These relationships were all significantly higher in tall fescue than bermudagrass hybrid (Table 2) also in the case of DGCI-turfgrass color ($r = 0.95$). Previous reports by Zhang and Kovacs (2012), and Leinauer et al. (2014) also indicated this trend of values between DGCI and turfgrass quality and turfgrass color. As in our study also in the report by Leinauer et al. (2014), the association between DGCI and turfgrass quality in tall fescue showed higher *r* values than the same association in bermudagrass hybrid. As for the turfgrass color, Zhang and Kovacs (2012) also studied the relationship between visual color rating and DGCI, with higher Pearson correlation coefficient in tall fescue than bermudagrass hybrid. Previous reports by Karcher and Richardson (2003) also

277 confirm that visual ratings can be used to separate treatment effects on turf color. Frequently raters
278 ranked the turf plots similarly although differences in color existed. Therefore, visual color rating
279 remains a valid evaluation tool if data are not compared across raters. However, the accuracy of
280 DGCI, as demonstrated in previous studies, enables researchers to record reflected turfgrass color
281 on a standardized scale rather than using arbitrary rating values.

282 **Please insert Table 2 near here**

283

Relationship between DGCI and NDVI

Both in bermudagrass hybrid (Figure 2 (a)) and tall fescue (Figure 2 (b)) DCGI significantly related to the average NDVI values measured with a HCS (GreenSeeker) and with the multispectral camera MAIA mounted on board a UAV, although data have been collected by instruments that measure at different heights and spatial resolutions. In fact DGCI has been collected only with RGB camera mounted on board a UAV, while NDVI has been measured by a multispectral camera on board a UAV and also by a ground based HCS.

As shown in the Figure 2, DGCI values were linearly associated with NDVI, as also demonstrated by Leinauer et al. 2014. In Figure 2 (a) bermudagrass hybrid has performed a higher degree of association in NDVI GreenSeeker-DGCI (r = 0.91) than with UAV (r = 0.85), while in the case of tall fescue the degree of association was statistically the same (Figure 2 (b)). Comparing the two species, it was interesting to note that in tall fescue the correlation coefficients (Table 2) between both NDVI (GreenSeeker and UAV) and DGCI were higher than in Bermudagrass (Table 2; Figures 2 (a) - (b)).

Please insert Figure 2 near here

Relationship between DGCI and clipping nitrogen content

Figure 3 showed the linear relationship between DGCI and clipping nitrogen content percentage in bermudagrass hybrid (a) and tall fescue (b) and it was of interest to note that the coefficients were high for both the species. However, DGCI in tall fescue showed a higher degree of association with clipping N content (r = 0.95), than in bermudagrass hybrid (r = 0.86) (Figures 3 (a) - (b)). DGCI values were linearly associated with clipping nitrogen content, as also demonstrated in other crops (Rorie, Purcell, Mozaffari et al. 2011; Vergara-Díaz et al. 2016). Thus, DGCI values could predict the average nitrogen concentrations of tall fescue and bermudagrass hybrid clippings in different plots and with different application rates.

The close association between DGCI and leaf nitrogen therefore provided an additional tool for the assessment of leaf nitrogen content. Our research was consistent with previous work by Karcher and Richardson (2003) who found that DGCI values were able to differentiate among turfgrass cultivars receiving various N treatments.

Please insert Figure 3 near here

Conclusions

DGCI values were highly correlated with the nitrogen clipping content and NDVI with a highly significant degree of association. The results suggested that UAS derived imagery RGB photography by UAVs had a great potential in supporting decisions. Thus, DGCI could be a promising remote-sensing tool for mapping the crop nitrogen status or NDVI at large scale with high precision and low cost (Li et al. 2015). This method could be used by farmers operating in large-scale farms to precisely manage the application of fertilizers, although the farmers especially in the developing and underdeveloped counties, they do not have enough knowledge to operate the UAV and manage the technology. As turfgrass, especially in the most developed countries, this method could allow golf course superintendents and turf management specialists to make critical decisions in real time without high up-front costs. Differences in camera quality and settings and lighting conditions could affect DGCI and limit their utility in diagnosing N deficiencies. Furthermore, disease, water status, nutritional deficiencies other than N, or different uniformity, texture and growth habit may affect greenness regardless of N status as suggested also by Rorie, Purcell, Karcher et al. 2011 and by Leinauer et al. 2014. More research is required on this technology and on the Smartphone APP FieldScout GreenIndex+ Turf (Spectrum Technologies, Inc., Aurora, IL, USA) (Spectrum Technologies, Inc. 2018) to study and overcome possible discrepancies between the APP and the Smartphone camera. Although the accuracy of a Smartphone camera is not comparable to a digital camera, the precision of a Smartphone camera could still help to detect minor changes in turf greenness over time and-or relative to other areas of the golf course or sports field. In fact, if the imagery was conveyed quickly to the user, a broader usage of this technology could allow golf course superintendents and turf management specialists to make critical decisions in real time without high up-front costs, in small areas. To use efficiently this technology on large scale, DGCI could be use directly on board an UAV and could serve as an indicator of N deficiency on turfgrass, thus increasing turfgrass nitrogen fertilization efficiency.

344 Indeed, applications installed in drones could be good solutions for farmers or golf course
345 superintendents and turf management specialists so they can adopt and benefit from DGCI
346 technology.

347 Acknowledgements

348 Authors wish to thank Mar. Project Srl, Geographike Srl, Geomind Srl and E.T.G. Srl staff for
349 the cooperation in research.
350
351 The present research was carried out within the Project "TurfUp_Remote monitoring and
352 management of turfgrass" co-financed under Tuscany ERDF ROP 2014-2020 Research
353 development and innovation Call 2.
354

References

Aguilar, C., J. C. Zinnert, M. J. Polo, and D. R. Young. 2012. "NDVI as an indicator for changes in water availability to woody vegetation". *Ecological Indicators* 23: 290–300.

Andrade-Sanchez, P., M. A. Gore, J. T. Heun, K. R. Thorp, A. E. Carmo-Silva, A. N. French, M. E. Salvucci, and J. White. 2014. "Development and evaluation of a field-based high-throughput phenotyping platform". Functional Plant Biology 41 (1): 68-79.

Barton, C. V. 2012. "Advances in remote sensing of plant stress". *Plant and Soil*; 354: 41–44.

Bausch, W. C., and H. R Duke. 1996. "Remote sensing of plant nitrogen status in corn". *Transactions of the ASAE* 39: 1869‑1875.

Bell, G. E., and X. Xiong. 2008. "The history, role, and potential of optical sensing for practical turf management". In P. Pessaraki (Ed.), *Handbook of turfgrass management and physiology* (pp. 641–658). Boca Raton, FL: CRC Press.

Bell, G. E., B. M. Howell, G. V. Johnson, W. R. Raun, J. B. Solie, and M. L. Stone. 2004. "Optical sensing of turfgrass chlorophyll content and tissue nitrogen". *HortScience* 39(5): 1130–1132.

Bell, G. E., J. K. Kruse, and J. M Krum. 2013. "The evolution of spectral sensing and advances in precision turfgrass management". In *Turfgrass: Biology, use and management,* edited by J. C. Stier, B. P. Horgan, and S. A. Bonos, 1151–1188. Madison, WI: American Society of Agronomy.

Blackmer, T. M., J. S. Schepers, G. E. Varvel, and G. E. Meyer. 1996 "Analysis of aerial photography for nitrogen stress within corn fields". *Agronomy Journal* 88 (5): 729-733.

Bremer, D. J., H. Lee, K. Su, and S. J. Keeley. 2011, "Relationships between normalized difference vegetation index and visual quality in cool-season turfgrass: II. Factors affecting NDVI and its component reflectances". *Crop Science* 51(5), 2219-2227.

379 Bremner, J. M. 1965 “Total Nitrogen”. In *Methods of Soil Analysis*, Part 2. Agron. Monogr. no.
380 9, edited by C. A. Black, 1149–1178. Madison, WI: American Society of Agronomy.
381 Carrow, R. N., J. M. Krum, I. Flitcroft, and V. Cline. 2010. “Precision turfgrass management:
382 challenges and field applications for mapping turfgrass soil and stress”. *Precision*
383 *Agriculture* 11: 115–134.
384 Casadesus, J., and D. Villegas. 2014. “Conventional digital cameras as a tool for assessing leaf
385 area index and biomass for cereal breeding”. *Journal of integrative plant biology* 56 (1): 7-
386 14.
387 Casadesús, J., Y. Kaya, J. Bort, M. M. Nachit, J. L. Araus, S. Amor, G. Ferrazzano, et al. 2007.
388 “Using vegetation indices derived from conventional digital cameras as selection criteria for
389 wheat breeding in water-limited environments”. *Annals of Applied Biology* 150 (2): 227-
390 236.
391 Caturegli L., F. Lulli, L. Foschi, L. Guglielminetti, E. Bonari, and M. Volterrani. 2014.
392 “Monitoring turfgrass species and cultivars by spectral reflectance”. *European Journal of*
393 *Horticultural Science* 79 (3): 97–107.
394 Caturegli L., F. Lulli, L. Foschi, L. Guglielminetti, E. Bonari, and M. Volterrani. 2014.
395 “Turfgrass spectral reflectance: simulating satellite monitoring of spectral signatures of main
396 C3 and C4 species”. *Precision. Agriculture.* 6: 297–310.
397 Caturegli L., M. Corniglia, M. Gaetani, N. Grossi, S. Magni, M. Migliazzi, L. Angelini, M. et
398 al. 2016. “Unmanned Aerial Vehicle to estimate nitrogen status of turfgrasses”. *PLOS ONE*.
399 11(6), e0158268. doi:10.1371/journal.pone.0158268.
400 Caturegli, L., M. Casucci, F. Lulli, N. Grossi, M. Gaetani, S. Magni, E. Bonari, and M.
401 Volterrani. 2015. “GeoEye-1 satellite versus groundbased multispectral data for estimating
402 nitrogen status of turfgrasses”. *International Journal of Remote Sensing* 36: 2238–2251.
403 Caturegli, L., N. Grossi, M. Saltari, M. Gaetani, S. Magni, A. E. Nikolopoulou, E. Bonari, and
404 M. Volterrani. 2015. “Spectral reflectance of tall fescue (Festuca Arundinacea Schreb.)

405 under different irrigation and nitrogen conditions". *Agriculture and Agricultural Science*
406 *Procedia* 4: 59–67.
407 Corwin, D. L., and S. M Lesch. 2005. "Apparent soil electrical conductivity measurements in
408 agriculture". *Computers and electronics in agriculture* 46(1), 11-43.
409 doi:10.1016/j.compag.2004.10.005.
410 Dordas, C. 2008. "Role of nutrients in controlling plant diseases in sustainable agriculture: A
411 review". *Agronomy for Sustainable Development* 28: 33–46. doi:10.1051/agro:2007051..
412 Dubbini, M. , A. Pezzuolo, M. D. Giglio, M. Gattelli, L. Curzio, D. Covi, T. Yezekyan, and F.
413 Marinello. 2017. "Last generation instrument for agriculture multispectral data collection".
414 *Agricultural Engineering International: CIGR Journal* 19 (1): 87–93.
415 Fensholt, R., and S. R. Proud. 2012. "Evaluation of earth observation based global long term
416 vegetation trends-Comparing GIMMS and MODIS global NDVI time series". *Remote*
417 *Sensing of Environment* 119: 131–147.
418 Fitz-Rodriguez, E., and C. Y. Choi. 2002. "Monitoring turfgrass quality using multispectral
419 radiometry". *Transactions of the ASAE* 45 (3):865‑867.
420 Fryskowska, A., M. Kedzierski, A. Grochala, and A. Braula. 2016. "Calibration of low cost
421 RGB and NIR UAV cameras". *International Archives of Photogrammetry, Remote Sensing*
422 *and Spatial Information Sciences* 41: 817-821.
423 Graeff, S., and W. Claupein. 2003. "Quantifying nitrogen status of corn (Zea mays L.) in the
424 field by reflectance measurements". *European Journal of Agronomy* 19 (4): 611–618.
425 Grossi N., M. Fontanelli, E. Garramone, A. Peruzzi, M. Raffaelli, M. Pirchio, L. Martelloni, et
426 al. 2016. "Autonomous mower saves energy and improves quality of tall fescue
427 lawn". *HortTechnology*, 26 (6): 825-830.
428 Gupta, S. G., M. M. Ghonge, and P. M. Jawandhiya. 2013. "Review of unmanned aircraft
429 system (UAS)". *International Journal of Advanced Research in Computer Engineering &*
430 *Technology* 2 (4): 1646–1658.

Hansen, P. M., and J. K Schjoerring. 2003. “Reflectance measurement of canopy biomass and nitrogen status in wheat crops using normalized difference vegetation indices and partial least squares regression”. *Remote Sensing of Environment* 86: 542–553.

Huang, J, F. He, K. Cui, R. J. Buresh, B. Xu, W. Gong, and S. Peng. 2008. “Determination of optimal nitrogen rate for rice varieties using a chlorophyll meter”. *Field Crops Research* 105: 70–80.

Jiang, Y, and R. N. Carrow. 2007. “Broadband spectral reflectance models of turfgrass species and cultivars to drought stress”. *Crop Science* 47: 1611–1618.

Johnsen, A. R., B. P. Horgan, B. S. Hulke, and V. Cline. 2009. “Evaluation of remote sensing to measure plant stress in Creeping Bentgrass (L.) fairways”. *Crop Science* 49: 2261–2274.

Karcher, D. E., and M. D. Richardson. 2003. “Quantifying turfgrass color using digital image analysis”. *Crop Science* 43: 943-951.

Karcher, D. E., and M. D. Richardson. 2013. “Digital image analysis in turfgrass research”. In *Turfgrass: Biology, use and management,* edited by J. C. Stier, B. P. Horgan, and S. A. Bonos, 1133–1149. Madison, WI: American Society of Agronomy.

Krum, J. M., R. N. Carrow, and K. Karnok. 2010. “Spatial mapping of complex turfgrass sites: Site-specific management units and protocols”. *Crop Science* 50: 301–315

Laliberte, A. S., and A. Rango. 2011. “Image processing and classification procedures for analysis of sub-decimeter imagery acquired with an unmanned aircraft over arid rangelands”. *GIScience & Remote Sensing* 48 (1): 4–23.

Leinauer, B., D. M. VanLeeuwen, M. Serena, M. Schiavon, and E. Sevostianova. 2014. “Digital image analysis and spectral reflectance to determine turfgrass quality”. *Agronomy Journal* 106 (5): 1787-1794.

Li, J., F. Zhang, X. Qian, Y. Zhu, and G. Shen. 2015. “Quantification of rice canopy nitrogen balance index with digital imagery from unmanned aerial vehicle”. *Remote Sensing Letters* 6: 183–189.

457 Ma, B. L., M. J. Morrison, and L. M Dwyer. 1996. "Canopy light reflectance and field
458 greenness to assess nitrogen fertilization and yield of maize". *Agronomy Journal* 88: 915–
459 920.
460 Magni S., M. Gaetani, N. Grossi, L. Caturegli, S. La Bella, C. Leto, G. Virga, T. Tuttolomondo,
461 F. Lulli and M. Volterrani. 2014. "Bermudagrass adaptation in the Mediterranean climate:
462 phenotypic traits of 44 accessions". *Advances in Horticultural Science* 28 (1): 29-34.
463 Morris, K. N., and R. C. Shearman. 2008. "NTEP turfgrass evaluation guidelines". In NTEP
464 turfgrass evaluation workshop, 1–5. Beltsville, MD. Available:
465 http://www.ntep.org/cooperator.html. Accessed August 2018.
466 O'Brien, D. 2017. "New technologies for evaluating putting green surface characteristics". Ph.D
467 dissertation, University of Arkansas, AR, USA. Available from
468 https://scholarworks.uark.edu/etd/2614/
469 Perry, E. M., and J. R. Davenport. 2007. "Spectral and spatial differences in response of
470 vegetation indices to nitrogen treatments on apple". *Computer and Electronics in*
471 *Agriculture* 59 (1-2): 56–65. doi:10.1016/j.compag.2007.05.002.
472 Quantum GIS 2.18 software. Accessed July, 2018. https://www.qgis.org
473 Remondino, F., and C. Fraser. 2006. "Digital camera calibration methods: considerations and
474 comparisons". *International Archives of Photogrammetry, Remote Sensing and Spatial*
475 *Information Sciences* 36 (5): 266-272.
476 Rhezali, A., and R. Lahlali. 2017. "Nitrogen (N) mineral nutrition and imaging sensors for
477 determining N status and requirements of maize". *Journal of Imaging* 3(4): 51.
478 Rorie, R. L., L. C. Purcell, D. E. Karcher, and C. A. King. 2011. "The assessment of leaf
479 nitrogen in corn from digital images". *Crop Science* 51 (5): 2174-2180.
480 Rorie, R. L., L. C. Purcell, M. Mozaffari, D. E. Karcher, C. A. King, M. C. Marsh, and D. E.
481 Longer. 2011. "Association of "greenness" in corn with yield and leaf nitrogen
482 concentration". *Agronomy Journal* 103(2): 529-535.

SAL engineering MAIA S2 (Modena Italy; EOPTIS, Trento, Italy). Accessed July, 2018. https://www.salengineering.it/public/en/p/maia.asp

Samborski, S. M., N. Tremblay, and E. Fallon. 2009. "Strategies to make use of plant sensors-based diagnostic information for nitrogen recommendations". *Agronomy Journal* 101 (4): 800–816.

Scharf, PC, and J. A. Lory. 2009. "Calibrating reflectance measurements to predict optimal sidedress nitrogen rate for corn". *Agronomy Journal* 101 (3): 615-625.

Spectrum Technologies, Inc. 2018. Product manual, Item no. 2910TA, 2910T. Available:http://www. specmeters.com/assets/1/22/2910TA_2910T_GreenIndex__Turf1.pdf Accessed 11 Sept. 2018.

Trenholm, L. E., R. N. Carrow, and R. R. Duncan. 1999. Relationship of multispectral radiometry data to qualitative data in turfgrass research. *Crop Science* 39: 763–769.

Vergara-Díaz, O., M. A. Zaman-Allah, B. Masuka, A. Hornero, P. Zarco-Tejada, B. M. Prasanna, J. E. Cairns and J. L. Araus. 2016. "A novel remote sensing approach for prediction of maize yield under different conditions of nitrogen fertilization". *Frontiers in Plant Science* 7:666. doi: 10.3389/fpls.2016.00666.

Volterrani, M., N. Grossi, L. Foschi, and S. Miele. 2005. "Effects of nitrogen nutrition on bermudagrass spectral reflectance". *International Turfgrass Society Research Journal* 10: 1005–1014.

Walters, D. R., and I. J. Bingham. 2007. "Influence of nutrition on disease development caused by fungal pathogens: implications for plant disease control." *Annals of Applied Biology* 151 (3): 307–324. doi:10.1111/aab.2007.151.issue-3.

White, J. W., P. Andrade-Sanchez, M. A. Gore, K. F. Bronson, T. A. Coffelt, M. M. Conley, K. A. Feldmann, et al. 2012. "Field-based phenomics for plant genetics research". *Field Crops Research 133*: 101-112.

508 Xiang, M., J. Q. Moss, D. L. Martin, and Y. Wu. 2018. “The salinity tolerance of seeded-type
509 common bermudagrass cultivars and experimental selections”. *HortTechnology* 28(3): 276-
510 283.
511 Xiang, M., J. Q. Moss, D. L. Martin, K. Su, B. L. Dunn, and Y. Wu. 2017. “Evaluating the
512 salinity tolerance of clonal-type bermudagrass cultivars and an experimental selection”.
513 *HortScience* 52 (1): 185-191.
514 Xiong, X., G. E. Bell, J. B. Solie, M. W. Smith, and B. Martin. 2007. “Bermudagrass seasonal
515 responses to nitrogen fertilization and irrigation detected using optical sensing”. *Crop*
516 *Science* 47 (4): 1603–1610.
517 Zhang, C., and J. M. Kovacs. 2012. “The application of small unmanned aerial systems for
518 precision agriculture: a review”. *Precision Agriculture* 13: 693–712.
519 Zhou, B., A. Elazab, J. Bort, O. Vergara, M. D. Serret, and J. L. Araus. 2015. “Low-cost
520 assessment of wheat resistance to yellow rust through conventional RGB
521 images”. *Computers and electronics in agriculture* 116: 20-29.
522

523 **Captions**

524 **Table 1.** Instrument monochrome sensors with relative band-pass filters of the multispectral camera
525 MAIA.

526 **Table 2.** Pearson product-moment correlation coefficients (*r*) among clipping nitrogen content,
527 color intensity, turfgrass quality, plant water content (PWC), NDVI measured with a handheld crop
528 sensor (GreenSeeker) and NDVI measured with multispectral camera mounted on an unmanned
529 aerial vehicle (UAV) and dark green color index (DGCI) on a) bermudagrass hybrid; b) tall fescue.
530 For each species correlation coefficients are calculated across all entries.
531 All values are significant at the 0.010 level, except for DGCI color intensity, quality and PWC for
532 bermudagrass hybrid and DGCI quality for tall fescue, whose values are significant at the 0.001
533 level.

534 **Figure 1.** (a) UAV during flight operations (6 July 2017; Pisa, Italy; 43°40'N, 10° 19'E, 6 m. a. s.
535 l.); (b) The multispectral camera MAIA mounted on the UAV.

536 **Figure 2.** Linear relationship between NDVI measured with a handheld crop sensor (GreenSeeker)
537 and NDVI measured with a multispectral camera mounted on UAV and DGCI on (a) bermudagrass
538 hybrid*;* and (b) tall fescue. Values represented the 3 replications.

539 **Figure 3.** Linear relationship between DGCI and the clipping nitrogen content (%) on (a)
540 bermudagrass hybrid*;* and (b) tall fescue. Values represented the average of 3 replications.

541

542

Tables

Table 1. Instrument monochrome sensors with relative band-pass filters of the multispectral camera MAIA.

Wavelength (nm)		
Start	Central	Stop
395.0	422.5	450.0
455.0	487.5	520.0
525.0	550.0	575.0
580.0	602.5	625.0
630.0	660.0	690.0
705.0	725.0	745.0
750.0	785.0	820.0
825.0	887.5	950.0

Table 2. Pearson product-moment correlation coefficients (*r*) among clipping nitrogen content, color intensity, turfgrass quality, plant water content (PWC), NDVI measured with a handheld crop sensor (GreenSeeker) and NDVI measured with multispectral camera mounted on an unmanned aerial vehicle (UAV) and dark green color index (DGCI) on a) bermudagrass hybrid; b) tall fescue. For each species correlation coefficients are calculated across all entries.

All values are significant at the 0.010 level, except for DGCI color intensity, quality and PWC for bermudagrass hybrid and DGCI quality for tall fescue, whose values are significant at the 0.001 level.

r	Color intensity	Quality	PWC	NDVI GreenSeeker	NDVI UAV	DGCI
a) Bermudagrass hybrid						
N clipping (%)	0.97	0.97	0.95	0.94	0.92	0.86
Color intensity (1-9)	N/A	0.94	0.99	0.94	0.94	0.83
Quality (1-9)	N/A	N/A	0.97	0.97	0.94	0.84
PWC (%)	N/A	N/A	N/A	0.92	0.92	0.84
NDVI GreenSeeker (780,660)	N/A	N/A	N/A	N/A	0.96	0.91
NDVI UAV (830,660)	N/A	N/A	N/A	N/A	N/A	0.85
b) Tall fescue						
N clipping (%)	0.99	0.99	0.99	0.95	0.94	0.95
Color intensity (1-9)	N/A	0.99	0.99	0.96	0.95	0.95
Quality (1-9)	N/A	N/A	0.98	0.94	0.93	0.92
PWC (%)	N/A	N/A	N/A	0.98	0.98	0.98
NDVI GreenSeeker (780,660)	N/A	N/A	N/A	N/A	0.99	0.95
NDVI UAV (830,660)	N/A	N/A	N/A	N/A	N/A	0.96