



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Replacement reactions and deformation by dissolution and precipitation processes in amphibolites

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Replacement reactions and deformation by dissolution and precipitation processes in amphibolites / Giuntoli F.; Menegon L.; Warren C.J.. - In: JOURNAL OF METAMORPHIC GEOLOGY. - ISSN 1525-1314. - ELETTRONICO. - 36:9(2018), pp. 1263-1286. [10.1111/jmg.12445]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/719234> since: 2020-01-30

*Published:*

DOI: <http://doi.org/10.1111/jmg.12445>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of

Giuntoli F.; Menegon L.; Warren C. J.: Replacement reactions and deformation by dissolution and precipitation processes in amphibolites. JOURNAL OF METAMORPHIC GEOLOGY 36. 1525-1314

DOI: 10.1111/jmg.12445

The final published version is available online at: <http://dx.doi.org/10.1111/jmg.12445>

#### Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

# Replacement reactions and deformation by dissolution and precipitation processes in amphibolites

Francesco Giuntoli<sup>1</sup>

Luca Menegon<sup>1</sup> Clare J. Warren<sup>2</sup>

<sup>1</sup>School of Geography, Earth and Environmental Sciences, Plymouth University, Plymouth, UK

<sup>2</sup>School of Environment, Earth and Ecosystem Sciences, Faculty of Science, Technology, Engineering and Mathematics, The Open University, Milton Keynes, UK

## Correspondence

Francesco Giuntoli, School of Geography, Earth and Environmental Sciences, Plymouth University, Plymouth PL4 8AA, UK.

Email: francesco.giuntoli@gmail.com

## Funding information

Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung, Grant/Award Number: P2BEP2 168722; FP7 People: Marie Curie Actions, Grant/Award Number: PCIG13 GA 2013 618289; Early Postdoc.Mobility, Grant/Award Number: P2BEP2 168722; FP7 Marie Curie Career Integration Grant, Grant/Award Number: PCIG13 GA 2013 618289

Handling Editor: Michael Brown

## Abstract

The deformation of the middle to lower crust in collisional settings occurs via deformation mechanisms that vary with rock composition, fluid content, pressure, and temperature. These mechanisms are responsible for the accommodation of large tectonic transport distances during nappe stacking and exhumation. Here, we show that fracturing and fluid flow triggered coupled dissolution precipitation and dissolution precipitation creep processes, which were responsible for the formation of a mylonitic microstructure in amphibolites. This fabric is developed over a crustal thickness >500 m in the Lower Seve Nappe (Scandinavian Caledonides). Amphibolites display a mylonitic foliation that wraps around albite porphyroclasts appearing dark in panchromatic cathodoluminescence (CL). The albite porphyroclasts were dissected and fragmented by fractures preferentially developed along the (001) cleavage planes and display lobate edges with embayments and peninsular features. Two albite/oligoclase generations, bright in CL, resorbed and overgrew the porphyroclasts, sealing the fractures. Electron backscattered diffraction shows that the two albite/oligoclase generations grew both pseudomorphically and topotaxially at the expense of the albite porphyroclasts and epitaxially around them. These two albite/oligoclase generations also grew as neoblasts elongated parallel to the mylonitic foliation. The amphibole crystals experienced a similar microstructural evolution, as evidenced by corroded ferrohornblende cores surrounded by ferrotschermakite rims that preserve the same crystallographic orientation of the cores. Misorientation maps highlight how misorientations in amphibole are related to displacement along fractures perpendicular to its *c*-axis. No crystal plasticity is observed in either mineral species. Plagioclase and amphibole display a crystallographic preferred orientation that is the result of topotaxial growth on parental grains and nucleation of new grains with a similar crystallographic orientation. Amphibole and plagioclase thermobarometry constrains the mylonitic foliation development to the epidote amphibolite facies (~600°C, 0.75–0.97 GPa). Our results demonstrate that at middle to lower crustal levels, the presence of H<sub>2</sub>O-rich fluid at grain boundaries facilitates replacement reactions by coupled dissolution precipitation and favours deformation by dissolution precipitation creep over dislocation creep in plagioclase and amphibole.

## KEYWORDS

Caledonides, dissolution precipitation processes, electron backscatter diffraction, replacement reactions, X ray mapping

## 1 | INTRODUCTION

The thermomechanical properties of the middle to lower crust exert a fundamental control on the structure of orogenic belts and on the amount and style of shortening during continental collision (e.g., Jackson, Austrheim, McKenzie, & Priestley, 2004; Mouthereau, Watts, & Burov, 2013). In particular, how strain is distributed vertically and horizontally in orogenic belts is one of the more important questions in crustal dynamics, and one that can be addressed by investigating the deformation mechanisms associated with the accumulation of hundreds of kilometre tectonic transport distances along thrust faults during mountain building processes (e.g., Fousseis & Handy, 2008; Gilotti, 1989; Mouthereau, Lacombe, & Vergés, 2012; Northrup, 1996; Royden, 1996).

In particular, the deformation processes and rheology of mafic shear zones are the subject of considerable debate because their main mineral constituents (e.g., plagioclase, amphibole, clinopyroxene) are expected to be rheologically strong at middle to lower crustal conditions (e.g., Bürgmann & Dresen, 2008). Thus, the weakening of mafic assemblages along major thrust faults developed at middle to lower crustal conditions seems to critically depend on the occurrence of metamorphic reactions, which can result in the formation of rheologically weaker phases or in the formation of fine-grained material able to deform by grain size sensitive creep or in both (e.g., Brander, Svahnberg, & Piazzolo, 2012; Okudaira, Shigematsu, Harigane, & Yoshida, 2017; Rutter & Brodie, 1992). Furthermore, the presence of H<sub>2</sub>O-rich fluid at the grain boundary typically facilitates dissolution and precipitation processes, which have been identified as the main deformation mechanisms in different mid-crustal lithologies and up to high temperature and high pressure (HP) conditions (Carmichael, 1969; Giuntoli, Lanari, & Engi, 2018; Gratier, Dysthe, & Renard, 2013; Imon, Okudaira, & Fujimoto, 2002; Imon, Okudaira, & Kanagawa, 2004; McAleer et al., 2017; Menegon, Pennacchioni, & Spiess, 2008; Mukai, Austrheim, Putnis, & Putnis, 2014; Putnis, 2009; Rutter, 1983; Stokes, Wintsch, & Southworth, 2012; Wassmann & Stöckhert, 2013; Wassmann, Stöckhert, & Trepmann, 2011; Wintsch & Yi, 2002). Two main dissolution and precipitation processes can be distinguished: coupled dissolution precipitation and dissolution precipitation creep. Coupled dissolution precipitation results in the pseudomorphic (maintaining the size and shape of the pre-existing phase) and topotaxial (using the orientation of the pre-existing phase) replacement of a parent phase by a product phase from a reaction interface (e.g., Putnis, 2002; Ruiz-Agudo, Putnis, & Putnis, 2014). Dissolution precipitation creep includes the transport of the chemical constituents from the dissolution sites, of locally high normal stress, to the precipitation site, of locally low

normal stress (e.g., Imon et al., 2002, 2004; Mukai et al., 2014; Wassmann et al., 2011). Both processes invariably require the presence of a fluid.


Porosity is another essential requirement for the operation of dissolution and precipitation processes, as it provides the necessary fluid pathways. Porosity may develop during replacement reactions due to molar volume differences between the dissolved and the precipitated minerals (Engvik, Putnis, Gerald, & Austrheim, 2008) or from dehydration reactions and associated fracturing (Plumper, John, Podladchikov, Vrijmoed, & Scambelluri, 2017). Other porosity-generating mechanisms in metamorphic environments include dilatancy at grain boundaries (Tullis, Yund, & Farver, 1996), fracturing (e.g., Brander et al., 2012), and creep cavitation in fine-grained ultramylonites deforming by grain size sensitive creep (Fousseis, Regenauer-Lieb, Liu, Hough, & De Carlo, 2009; Menegon, Fousseis, Stünitz, & Xiao, 2015).

Here, we show that mineral reactions and deformation in amphibolites occurred mainly by coupled dissolution precipitation and dissolution precipitation creep at epidote amphibolite facies conditions, and that fracturing was the most efficient porosity-generating mechanism assisting deformation. Metamorphism and deformation in the amphibolites resulted in the development of a >500 m thick mylonitic foliation during Caledonian nappe thrusting.

## 2 | GEOLOGICAL SETTING

The Scandinavian Caledonides developed due to the closure of the Iapetus Ocean in the Ordovician and the subsequent subduction and continent collision of the Baltica plate below the Laurentia plate in the Silurian to early Devonian (e.g., Gee, Fossen, Henriksen, & Higgins, 2008; Roberts, 2003; Roberts & Gee, 1985; Stephens, 1988). In the Scandinavian Caledonides, tectonic units were transported up to 400 km to the east (Gayer, Rice, Roberts, Townsend, & Welbon, 1987; Gee, 1975; Gee, Juhlin, Pascal, & Robinson, 2010; Rice & Anderson, 2016; Roberts & Gee, 1985) as a result of the collision, creating a nappe stack of several allochthons units on top of Autochthons Baltic Shield (Figure 1a,b). After emplacement, the nappe stack was folded into north-trending synforms and anti-forms, possibly related to the crustal extension and normal faulting occurring during the latest orogenic phases (Bergman & Sjöström, 1997). By virtue of their deep erosional level, the internal parts of the Scandinavian Caledonides expose middle and lower crustal sections involved in subduction exhumation history and nappe stacking.

The Scandinavian Caledonides are subdivided, from top to bottom, into an Uppermost Allochthon, an Upper Allochthon, a Middle Allochthon, and a Lower Allochthon



**FIGURE 1** Geological setting of the Scandinavian Caledonides. (a) Tectonic map with inferred paleogeography of the nappes (modified after Gee et al., 2010). (b) Cross section marked in (a) with vertical exaggeration of 5× and approximate location of the COSC 1 borehole (modified after Gee et al., 2010). (c) Detail of the study area with location of the COSC 1 drilling site (modified after Strömberg et al., 1984)

based on tectonostratigraphy (Figure 1; Gee & Sturt, 1985; Strand & Kulling, 1972; Strömberg et al., 1984). The Middle Allochthon, the target of this study, includes several basement units and associated metasedimentary rocks representing the outermost Baltica margin and possibly including units derived from an ocean continent transition zone (e.g., Andréasson, 1994; Gee, Janák, Majka, Robinson, & van Roermund, 2013; Gee et al., 2008; Roberts, 2003; Stephens, 1988). The upper tectonic unit of the Middle Allochthon is the Seve Nappe Complex (SNC; e.g., Sjöström, 1983), which, in the central Scandes, outcrops over a N S distance of ~1,000 km and an W E distance of ~200 km (Figure 1; Andréasson, 1994).

In the Jämtland region, the SNC can be further subdivided into Lower, Middle, and Upper Seve Nappe by the presence of internal thrust sheets (Zachrisson & Sjöstrand, 1990). The Lower Seve Nappe is mainly composed of micaschists, quartzites, and metapsammites with gneisses, metabasics, and with minor peridotites and serpentinites (Figure 1c). The Middle Seve Nappe is composed of

similar lithotypes, but overprinted by a pervasive migmatization. Several parts of the Lower and Middle Seve preserve evidence of HP to ultrahigh pressure (UHP) metamorphism (summary in figure 4 of Klonowska, Janák, Majka, Froitzheim, & Kościńska, 2016) spanning from ~1.1 GPa and 600°C up to 4 GPa and 800°C, within the stability field of coesite and diamond (Brueckner & van Roermund, 2004; Gilio, Clos, & van Roermund, 2015; Janák, van Roermund, Majka, & Gee, 2013; Klonowska et al., 2016, 2017; Majka et al., 2014; Van Roermund, 1985, 1989). The HP-UHP metamorphism is the manifestation of the Ordovician subduction of the SNC (Brueckner & Van Roermund, 2007; Ladenberger et al., 2013; Root & Corfu, 2012). It is worth noting that, to date, no evidence of (U)HP metamorphism has been recorded in the Lower Seve Nappe in the central Jämtland. In the Middle Seve Nappe, the granulite and amphibolite facies metamorphism appears to postdate the HP-UHP stage and produced partial melting at 442–436 Ma (Ladenberger et al., 2013). A recent study has related the amphibolite facies metamorphic

“stage” to the exhumation and lateral extrusion of the SNC (Grimmer, Glodny, Drüppel, Greiling, & Kontny, 2015). In the Lower Seve Nappe, a pervasive amphibolite facies foliation overprints the (U)HP fabric where present; where not present, it represents the main metamorphic fabric. In the Åreskutan area, the amphibolite facies metamorphic stage was constrained at 550°C and 0.2–0.5 GPa (Arnbom, 1980).

The “Collisional Orogeny in the Scandinavian Caledonides (COSC-1; IGSN ICDP5054EHW1001)” borehole (Lorenz et al., 2015; see location in Figure 1b,c) is located in the central Jämtland region, near Åre (Sweden). The drill core provides an almost complete section (recovery rate higher than 99%) through the Lower Seve Nappe. In detail, the core comprises alternating layers of felsic gneisses, calc-silicates, and amphibolites displaying narrow (mm–cm) and localized shear zones from the surface down to 1,700 m (Hedin et al., 2016). The rocks show strongly deformed fabrics from 1,700 to 2,500 m (the end of the core; Lorenz et al., 2015), with the development of mylonitic fabrics. The lowermost portion of the core is composed of strongly deformed metasedimentary rocks. Acoustic televiewer data indicate that the foliation is generally shallow and trends N–S, but there are remarkable exceptions, where the foliation has dips to the E or W with angles >50°, related to recumbent folds and boudinage (Wenning et al., 2017).

## 3 | METHODS

### 3.1 | Scanning electron microscopy

All the scanning electron microscopy (SEM) analyses were performed on carbon-coated polished thin sections cut perpendicular to the foliation and parallel to the stretching lineation of the sample. Backscattered electron (BSE) and cathodoluminescence (CL) analyses were performed at the Open University (UK), using an FEI Quanta 200 three-dimensional SEM on carbon-coated thin sections. Analyses were conducted under high vacuum, using an accelerating voltage of 10 kV, a beam current of 3.3 nA, a working distance of 13 mm, and an electron source provided by a tungsten filament. The panchromatic CL detector used is a Centaurus Deben with a photo multiplier tube (Hamamatsu R316) characterized by sensitivity in the range of 400–1,200 nm.

Electron backscattered diffraction (EBSD) analyses were conducted with a Jeol-7001FEG SEM at the Electron Microscopy Centre, Plymouth University (UK). EBSD patterns were acquired with a 70° tilted sample geometry, 20 kV accelerating voltage, 18–23 mm working distance, and 1.3–1.7 µm step size. Diffraction patterns were automatically indexed using AZtec (Oxford Instruments). The indexing match units used for the analysed phases were

“anorthite” (Laue group-1) for albite and oligoclase and “hornblende” (Laue group 2/m) for tschermakite ferrotschermakite. Both match units were taken from the American Mineralogist database. Raw maps were processed with HKL Channel 5 (Oxford Instruments), using the noise reduction procedure tested by Prior, Wheeler, Peruzzo, Spiess, and Storey (2002). Wild spikes were removed, and unindexed points were replaced by the average orientation of the neighbour points. Grains smaller than three times the step size were not considered in the analysis. The mean angular deviation values were 0.3 for amphibole and 0.4–0.6 for plagioclase; the raw indexing rate ranged between 90 and 95%. Crystallographic directions were plotted on pole figures (upper and lower hemisphere of the stereographic projection), with X parallel to the stretching lineation and Z parallel to the pole of the mylonitic foliation. The grain orientation spread maps (GOS maps) were calculated as the average misorientation between every pixel in the grain and the grain's average orientation.

### 3.2 | Electron probe microanalyser

Electron probe microanalyser (EPMA) analyses were conducted at the Open University (UK), using a Cameca SX100 connected to five spectrometers. Wavelength dispersive spectrometers were used for both spot analyses and X-ray maps, the latter acquired following the procedure of Lanari et al. (2013). Spot analyses were acquired first for each mineral, before performing the X-ray maps on the same area. Spot analyses were performed with 20 KeV accelerating voltage, 20 nA specimen current, and 2 µm beam diameter. Ten oxide compositions were measured, using natural standards: K-feldspar (SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, K<sub>2</sub>O), bustamite (CaO, MnO), haematite (FeO), forsterite (MgO), jadeite (Na<sub>2</sub>O), rutile (TiO<sub>2</sub>), apatite (P<sub>2</sub>O<sub>5</sub>). A ZAF matrix correction routine was applied; uncertainty on major element concentrations was <1%. X-ray maps were acquired with 15 KeV accelerating voltage, 100 nA specimen current, dwell times of 70–100 ms, and step size of 5 µm. Ten elements (Si, Ti, Al, Fe, Mn, Mg, Na, Ca, K, and P) were measured at the specific wavelength in two series. Intensity X-ray maps were standardized to concentration maps of oxide weight percentage using spot analyses as internal standard. X-ray maps were processed using XMapTools 2.2.1 (Lanari et al., 2014).

### 3.3 | Geothermobarometry

#### 3.3.1 | Amphibole–plagioclase thermobarometry

Temperature was estimated using the Holland and Blundy (1994) geothermometer, which is based on element

exchange between amphibole and plagioclase pairs in equilibrium. The calibration reaction edenite+albite=richterite+anorthite was constrained from experimental and natural data for silica-saturated and silica-rich igneous and metamorphic rocks in the range of 0.1–1.5 GPa and 400–1,000°C.

Pressure was estimated using two geobarometers: Bhadra and Bhattacharya (2007) and Anderson and Smith (1995). The former is based on element distribution between amphibole and plagioclase pairs in equilibrium. Experimental data were conducted on silica-saturated assemblages in the  $P$ – $T$  range of 0.1–1.5 GPa and 650–950°C, and on the reaction tremolite+tschermakite+2 albite = 2 pargasite+8 quartz. The latter is based on the increase of Al in hornblende with increasing pressure and is calibrated on experimental data at 675 and 760°C, accounting for the effects of temperature and  $f_{O_2}$ . Pressure and temperature were calculated using the Plagioclase Hornblende Thermobarometry spreadsheet (Anderson, Barth, Wooden, & Mazdab, 2008) and the spreadsheet of Wallis, Phillips, and Lloyd (2014) (the latter was used to derive pressure from the calibration of Bhadra & Bhattacharya, 2007).

### 3.3.2 | Chlorite+Quartz+H<sub>2</sub>O thermometry


Chlorite+Quartz+H<sub>2</sub>O thermometry was performed using the program CHLMICA<sub>EQUI</sub> (Lanari, 2012). The crystallization temperature of chlorite and the  $X_{Fe^{3+}}$  were computed at a fixed pressure of 0.7 GPa in the range of 100–550°C from the convergence of four equilibria involving five chlorite end-members, quartz, and H<sub>2</sub>O (Lanari et al., 2012; Vidal, Lanari, Munoz, Bourdelle, & De Andrade, 2016) using standard state properties and solid solution models of Vidal, Parra, and Vieillard (2005) and Vidal et al. (2006).

## 4 | RESULTS


### 4.1 | Petrography and microstructure

In the middle and lower portions of the COSC-1 borehole (~1,500–2,300 m depth), amphibolites show a mylonitic foliation. The studied sample (International Geological Sample Number: ICDP5054EX8E601) was extracted from a depth of 2,206.97 m from the surface and is similar to several amphibolite samples present between ~1,600 and ~2,300 m depth. In this sample, the mylonitic foliation is subhorizontal, it wraps around plagioclase porphyrocrysts and is defined by amphibole, plagioclase (with smaller grain size, details in the following), chlorite, quartz, epidote, and ilmenite (Figure 2). Quartz and calcite occur mainly in veins subparallel to the mylonitic foliation or as fine grains dispersed in the mylonitic foliation, typically along the phase boundaries between plagioclase and amphibole.

Plagioclase is present as zoned porphyrocrysts (up to 0.5–1 cm in size) wrapped by the mylonitic foliation, and as finer (up to hundreds of  $\mu\text{m}$  in size) albite/oligoclase grains elongated parallel to the mylonitic foliation (aspect ratios up to ~5; mean ~2). Plagioclase porphyrocrysts have albite cores that appear turbid due to abundant fine-grained (up to 30  $\mu\text{m}$  in size) inclusions of ilmenite, rutile, and epidote. Ilmenite crystals are also included with bigger grain size (100  $\mu\text{m}$  in size); quartz is present as inclusions with variable grain size (tens to hundreds of  $\mu\text{m}$  in size; Figures 3 and 4). The inclusions locally define a rotated internal foliation (upper plagioclase porphyrocryst in Figure 3a). These cores are porphyroclastic and are cut by fracture systems and some are dismembered parallel to the foliation (see details in Section 4.2). The cores are always surrounded by albite/oligoclase rims. Pores are evident in the albite/oligoclase rims: they are few microns in size and



**FIGURE 2** Amphibolite with Pl porphyroclast displaying a mylonitic foliation defined by Amp, Pl, Chl, Qz, Ep, and Ilm. Cal and Qz ribbons, presumably representing transposed veins, are parallel to the foliation. Both dextral (clasts close to Figure 3 rectangle, marked by the white arrow) and sinistral sense of shear ( $C'$  planes, red dashed line) are visible. (a) plane polarized light; (b) crossed polarized light. Mineral abbreviation from Whitney and Evans (2010)




occur in trails parallel to fracture systems that dissect the plagioclase core (Figure 4). Several mineral inclusions also occur along such trails, ranging from a few microns to tens of microns in size. These are muscovite, epidote, and calcite, with rare Ba-rich muscovite (Figure 4). Such inclusions confer a turbid aspect to plagioclase rim in plane-polarized light micrographs (Figure 3a; Table 1).

Amphibole is pleochroic, with absorption colours ranging from light brown to dark green blue; in some bigger crystals absorption colours highlight a paler core and a darker rim. Amphibole displays the *C*-axis parallel to the stretching lineation and a maximum grain size >1 mm. Chlorite has a pleochroism varying from light yellow to light green, a negative optical sign and grain size similar to



**FIGURE 3** More detailed areas of sections shown in Figure 2. (a) Ab porphyroclasts with dark trails of inclusions of Ep and Cal few  $\mu\text{m}$  in size. The Ab porphyroclasts are wrapped by the mylonitic foliation defined by Amp, Olig, and Chl. The white rectangle indicates the site of the EBSD map shown in Figure 7 (plane polarized light). (b) X ray map showing Pl, Amp, Chl, and Ep that crystallize in the pressure shadows of Ab porphyroclasts. (c) BSE image showing zoned Pl with dark cores (P11) and brighter rims (P12 and P13). Bright inclusions inside Pl are Ilm crystals. (d) CL image highlighting the difference between Ab cores (dark, P11) rimmed by Pl with higher An content (bright, P12 and P13). This bright Pl crystallizes also in the pressure shadows and in the fractures that dissect the Ab cores. (e) Standardized X ray map of the X anorthite ( $X_{\text{An}}$ ) in the Pl. Note the fractures in P11 sealed by P13 and sheared off fragments of P11 rimmed by P12 and P13 (see text and Figure 5 for distinction). (f) Standardized X ray map of Amp crystals displaying relic cores higher in Mg# (Amp1) and rims lower in Mg# lengthened as the main foliation (Amp2). A minor but consistent shift towards higher Mg# is visible from the top to the bottom of the picture, probably related to an analytical artefact




**FIGURE 4** Details of the microstructures shown in Figure 3c. BSE images. (a) Dark P11, highlighted by the dashed yellow line, with lobate edges and peninsular features rimmed by P12 and P13. (b) Iso oriented Ilm and Rt inclusions in P11. Note the two perpendicular trails of pores in P12 and P13. (c) Detail of the pore trails and associated Ms and Cal inclusions. (d) Close up of (c) highlighting pores, abundant Ms inclusions and rare Ba rich Ms

the amphibole. Asymmetric pressure shadows around plagioclase porphyroclasts indicate both dextral and sinistral sense of shear. In these pressure shadows amphibole, albite/oligoclase grains and chlorite crystallize intergrown and elongated parallel to the mylonitic foliation.  $C'$  planes display a dominant sinistral sense of shear and are defined by the same minerals found along the mylonitic foliation (Figures 2 and 3a,b). Opaque minerals include mostly ilmenite, up to 1 mm in size, with minor magnetite and pyrite.


Ilmenite crystals are elongate and lie parallel to the mylonitic foliation.

## 4.2 | Microstructure and chemistry of plagioclase

Backscattered electron images highlight two plagioclase generations: dark cores with an albitic composition (P11) are surrounded by brighter plagioclase rims with higher anorthite



(d) Diagram Parameters ( $Ca_B \geq 1.50$ ;  $(Na+K)_A < 0.50$ ;  $Ca_A < 0.50$ )



**FIGURE 5** Mineral chemistry of Pl, Amp, and Chl. The subdivision in generations was made based on microtextures. (a) Pl groups based on the  $X_{Ab}$  and  $X_{An}$  content: PI1  $X_{An}$  0 0.05; PI2  $X_{An}$  0.05 0.13; PI3  $X_{An}$  0.13 0.25. Compare with Figure 3e. (b) Amp groups based on the Mg# and Si apfu content: Amp1 Mg# 0.5 0.44, Si apfu 7 6.5; Amp2 Mg# 0.44 0.36, Si apfu 6.5 6.1. Compare with Figure 3f. (c) Compositional map of Chl Mg# for 0% Fe<sup>3+</sup>. The crystals display homogeneous compositions (Chl1 Mg# 0.59 0.56), except at grain boundaries and along C' band where lower values are present (Chl2 Mg# 0.55 0.52). (d) Diagram of classification of calcic amphibole plotted with the average chemical compositions of Amp1 and Amp2 (from Leake et al., 1997)

content (PI2 and PI3, as defined based on chemistry in the next paragraph; Figures 3c and 4). Furthermore, the cores are cut by fractures filled with plagioclase with the same composition as the plagioclase rims. The distinction of these plagioclase generations is more evident in the CL images, where plagioclase cores appear moderately luminescent and are surrounded by bright plagioclase rims (Figure 3d). The fractures dissecting the cores are as bright as the plagioclase rims. The plagioclase cores (PI1) are locally fragmented and display lobate edges and embayments that are typically surrounded by bright plagioclase (Figures 3d, 7a and 8b). In some areas, the plagioclase rims can be further subdivided

into two generations based on the brightness of their CL response: a brighter plagioclase generally surrounded by a darker one (PI2 and PI3, respectively).

Compositional maps of plagioclase display a perfect match with the CL images (Figure 3e). The maps highlight three plagioclase compositions, numbered from the older to the younger: PI1 ( $X_{An}$  0 0.05), PI2 ( $X_{An}$  0.13 0.25), PI3 ( $X_{An}$  0.05 0.13; Figures 5a and 7b; Table 2). The plagioclase core (PI1) has an albite composition and is overgrown by the two albite/oligoclase rims (PI2 and PI3). PI2 is present only locally and is overgrown by PI3, as shown in the top right corner of Figure 3e. Furthermore, PI3 seals the

**TABLE 1** Metamorphic and deformation evolution of sample ICDP5054EX8E601

Minerals	Premain foliation	Main foliation	C' planes
Amp	-----	=====	
Pl	<u>Ab core</u>	Ab/Olig Rims	
Chl		=====	-----
Qz	=====	=====	-----
Cal	-----	=====	-----
Ep	=====	-----	
Ilm	=====	=====	

fractures inside P11. P11 forms porphyroclasts wrapped by the mylonitic foliation, whereas P12 and P13 occur in the pressure shadows of P11.

**TABLE 2** Representative average composition analysis (wt%) of P1 and Amp

	P1			Amp	
	Core (P11)	Rim1 (P12)	Rim2 (P13)	Core (Amp1)	Rim (Amp2)
SiO <sub>2</sub>	68.21	62.95	64.99	43.52	41.62
TiO <sub>2</sub>	0.01	0.01	0.01	0.22	0.35
Al <sub>2</sub> O <sub>3</sub>	19.08	24.37	21.55	13.21	15.42
FeO	0.03	0.03	0.06	18.41	18.56
MnO	0.00	0.00	0.00	0.17	0.17
MgO	0.00	0.00	0.00	8.34	7.37
CaO	0.11	4.30	2.30	11.42	10.72
Na <sub>2</sub> O	12.34	8.93	10.47	1.72	2.02
K <sub>2</sub> O	0.04	0.05	0.05	0.21	0.35
Sum	99.83	100.64	99.43	97.22	96.57
	<b>Formulae based on 8 O</b>			<b>Formulae based on 23 anhydrous O</b>	
Si	2.99	2.76	2.88	6.49	6.26
Ti				0.02	0.04
Al	0.99	1.26	1.12	2.33	2.73
Fe <sup>3+</sup>				0.41	0.49
Fe <sup>2+</sup>				1.89	1.85
Mn				0.02	0.02
Mg				1.86	1.65
Ca	0.01	0.20	0.11	1.83	1.73
Na	1.05	0.76	0.90	0.50	0.59
K	0.00	0.00	0.00	0.04	0.07
Sum	5.04	4.99	5.01	15.38	15.42
Mg#				0.45	0.41
X <sub>An</sub>	0.01	0.21	0.11		
X <sub>Ab</sub>	0.99	0.79	0.89		

### 4.3 | Microstructure and chemistry of amphibole and chlorite

Compositional maps of amphibole crystals display corroded cores, higher in Mg# ( $Mg\# = Mg/(Fe^{2+} + Mg)$ ; Amp1: Mg# 0.5 0.44, Si apfu 7 6.5), and rims, lower in Mg# (Amp2: Mg# 0.44 0.36, Si apfu 6.5 6.1; Figures 3f and 5b; Table 2). Based on the classification of Leake et al. (1997), the Amp1 is at the compositional boundary between tschermakite ferrotschermakite magnesiohornblende ferrohornblende; Amp2 is a ferrotschermakite (Figure 5c). There are local exceptions to this zoning pattern, in which the two peripheral areas have higher Mg# (Figure 9c). Amp2 and P13 grains displaying mutual intergrowths and elongation parallel to the foliation (Figure 3c f).

Chlorite is characterized by a homogeneous Mg# (Mg# 0.59 0.56, Chl1), except along grain boundaries and the C' planes, where some lower values are visible (Mg# 0.55 0.52, Chl2; Figure 5c; Table 3). Chl1 grains occur as intergrown with P13 and Amp2 along the mylonitic.

### 4.4 | Amp-P1 thermobarometry

Thermometric estimates for the P11 and Amp1 pair yield 400°C; thermobarometric estimates for the P12 and Amp1 pair and the Amp2 and P13 pair yield 615°C and 0.97 GPa

**TABLE 3** Representative average composition analysis (wt%) of Chl

Chl	Chl1-High Mg#		Chl2-Low Mg#	
	0% XFe <sup>3+</sup>	30% XFe <sup>3+</sup>	0% XFe <sup>3+</sup>	30% XFe <sup>3+</sup>
SiO <sub>2</sub>	25.70	25.70	26.52	26.69
Al <sub>2</sub> O <sub>3</sub>	22.33	22.34	21.70	21.69
FeO	22.00	21.98	22.60	22.41
Fe <sub>2</sub> O <sub>3</sub>		0.02		0.21
MnO	0.15	0.15	0.15	0.14
MgO	16.43	16.43	15.37	15.42
CaO	0.02	0.02	0.03	0.03
Na <sub>2</sub> O	0.01	0.01	0.01	0.02
K <sub>2</sub> O	0.00	0.00	0.01	0.01
Sum	86.63	86.65	86.38	86.62
	<b>Formulae based on 14 anhydrous O</b>			
Si	2.69	2.63	2.78	2.74
Al	2.75	2.70	2.68	2.62
Mg	2.56	2.51	2.41	2.36
Fe <sup>3+</sup>	0.00	0.57	0.00	0.58
Fe <sup>2+</sup>	1.92	1.32	1.98	1.35
Sum	9.92	9.73	9.85	9.65
Mg#	0.57	0.66	0.55	0.64

**TABLE 4** Results of amphibole plagioclase geothermobarometry computed from the values of Table 2

Amp-Pl couples	Thermometer HB (°C)	Barometer	
		BB (GPa)	AS (GPa)
Amp1 P11	392		
Amp1 P12	615	0.97	0.87
Amp2 P13	605	0.74	1.09

*Note.* Thermometer abbreviation: HB: Holland and Blundy (1994). Barometer abbreviations: BB: Bhadra and Bhattacharya (2007); AS: Anderson and Smith (1995). The favoured results are highlighted in bold (see Section 5.1 for details).

and 605°C and 0.74 GPa, respectively ( $\pm 50^\circ\text{C} \pm 0.2$  GPa, Table 4; see Section 5.1 for discussion on the plagioclase and amphibole growth zones that we consider to be in equilibrium).

#### 4.5 | Chlorite+Quartz+H<sub>2</sub>O thermometry

Chlorite+Quartz+H<sub>2</sub>O thermometry was performed on the two different chlorite groups, Chl1 and Chl2 (Figure 5c), at a fixed pressure of 0.7 GPa. The Chl1 displays two peaks in the T-frequency diagram (Figure 6a): one close to the temperature limit of the thermometer (550–450°C) and one at lower temperature (400–250°C). Chl2 yields a range of 200–350°C, with a peak at ~250°C (Figure 6b), with only one grain yielding a higher value (~500°C).

#### 4.6 | EBSD analysis

Figures 7 and 8 show the results of EBSD analysis of two microstructural domains (see Figures 2 and 3 for the location of EBSD maps): domain 1 includes a plagioclase porphyrocryst with a core of P11 composition rimmed by P12 and P13 compositions. The porphyrocryst P11 is cut by a network of intracrystalline microfractures oriented NW–SE and filled with P12–P13. Domain 1 also includes clusters of amphibole and plagioclase grains around the plagioclase porphyrocryst (Figure 7). Domain 2 contains an


aggregate of amphibole grains, mostly elongate parallel to the foliation, and several grains of plagioclase showing a core-and-rim zoning in CL images (Figure 8).

The comparison between EBSD maps, CL images, and compositional maps shows that the P11 porphyroclasts, which EBSD identifies as one individual grain (i.e., it does not contain high-angle boundaries with misorientation  $>10^\circ$ ), actually include areas with P12 and P13 compositions, which maintain the original crystal shape and crystallographic orientation of P11 (Figures 7 and 8). The internal distortion of the compositionally zoned porphyrocrysts, as evaluated with the GOS, is rather low (GOS values lower than 2°: Figures 7c and 8e). Low-angle boundaries (misorientations  $2-10^\circ$ ) are preferentially distributed along the microfractures. The analysis of local misorientations within P11 cores indicates that P12 and P13 areas have misorientations of up to 2° with respect to the P11 cores (Figure 7d). Higher misorientations (up to 3°) are found in the plagioclase porphyrocryst of the domain 2 (Figure 8d).


In domain 1, the compositionally zoned plagioclase porphyrocryst is surrounded by P12–P13 grains that range in size between 30 and 200  $\mu\text{m}$ , delimited by high-angle boundaries with misorientation  $>10^\circ$  from the porphyrocryst. The P12 and P13 grains have mean GOS values lower than 1°, with one maximum value of 4°.

Amphibole generally shows GOS values lower than 2°, with a few maxima of up to 7° (Figures 7e, 8g, and 9d). Misorientations progressively increase towards fractures perpendicular to the crystal elongation and, locally, towards very few low-angle boundaries oriented subparallel to the crystal elongation (Figures 7f, 8h, and 9f). The misorientation profile in Figure 9e highlights a jump in misorientation of up to 7° across the low-angle boundary.

The crystallographic orientation of the new grains of P12 and P13 displays a variable degree of overlap with the one of the P11 core that they overgrow (Figures 10a,b and 11a,b). In some cases, all the new grains inherit the crystallographic orientation of the P11 core (Figure 10a,b), whereas in other cases, the overlap is more limited and the










**FIGURE 6** Chlorite+Quartz+H<sub>2</sub>O thermometry results calculated at a pressure of 0.7 GPa and a range between 0 and 50% of Fe<sup>3+</sup>




**FIGURE 7** Details of the microstructure shown in Figure 3 (referred to as domain 1 in the text). (a) CL image of a dark P11 porphyroblast with fractures and rims containing the bright P12. (b) Standardized X ray map of the  $X_{An}$  content showing the overlap between the bright CL areas and the P12-P13 compositions. (c) EBSD GOS map superposed to the band contrast (BC) map of the same area shown in (a). White lines: low angle boundaries  $2-10^\circ$ . Black lines: high angle boundaries  $>10^\circ$ . Light blue lines: Twin boundaries in P1. (d) EBSD texture component map (TCM) of the P11 porphyroblast, showing the misorientation from the reference point marked by the red cross. White lines: low angle boundaries  $2-10^\circ$ . Black lines: high angle boundaries  $>10^\circ$ . Light blue lines: Twin boundaries in P1. (e) EBSD GOS of Amp from the area shown in (a). White lines: low angle boundaries  $2-10^\circ$ . Black lines: high angle boundaries  $>10^\circ$ . (f) EBSD TCM of Amp, showing the misorientation from the reference point marked by the red cross. White lines: low angle boundaries  $2-10^\circ$ . Black lines: high angle boundaries  $>10^\circ$ .

new grains show a significant dispersion of crystallographic directions with respect to the P11 cores (Figure 11). The crystallographic inheritance is more evident if we consider the crystallographic orientation of the P12-P13 grains in direct contact with the P11 core (Figures 10a and 11b). The

P12-P13 grains dispersed in the mylonitic matrix show a wide range of crystallographic orientations, which do not necessarily overlap with those of the P11 cores included in the map (Figures 10b and 11a). EBSD analysis also shows that the most common fracture set in the P11 porphyroclasts



**FIGURE 8** Details of Figure 2 (referred to as domain 2 in the text). (a) Light microscopy microstructure of the site analysed with EBSD. (b) CL image of the site shown in (a). Dark P11 porphyroclasts are overgrown by bright P12 and P13. (c) EBSD phases map of the site shown in (a). Note the epitaxial growth of P12 and P13 on P11, as indicated by the lack of high angle boundaries separating P11 from P12 and P13, and the very few low angle boundaries in the P11 porphyroclast. (d) EBSD TCM showing the misorientation from reference point (red cross) in P11 porphyroclasts. Note the correlation between higher misorientation and P12 P13 overgrowth on the right hand side of the reference point. (e) EBSD GOS map suggesting that the Pl crystals are very low internal strain. (f) EBSD TCM showing the misorientation from reference point of another P11 porphyroclast. (g) EBSD GOS map of Amp. (h) EBSD TCM showing the misorientation from reference point in an Amp crystal elongate parallel to the foliation



**FIGURE 9** Details of Figure 2. (a) Sigmoidal grain of Amp surrounded by Chl along the foliation wrapping around Pl porphyroclasts. Several brittle fractures perpendicular to the Amp elongation are visible (plane polarized light). The white rectangle encompasses the site of the EBSD maps shown in (d) and (f). (b) X ray map of the mineral phases in (a). (c) Standardized X ray map showing the variation in Mg# of Amp grains (scale bar on the right hand side). Amp displays cores higher in Mg# (Amp1) and rims lower in Mg# (Amp2) elongate parallel to the foliation (compare with Figure 3f). (d) EBSD GOS map of Amp. The black line is the trace of the misorientation profile a b shown in (e). See (a) for the location of the map. (e) Misorientation profile a b drawn across a low angle boundary. (f) EBSD TCM showing the misorientation from the reference point marked by a red cross

is parallel to the (001) plane, which is a perfect cleavage plane in plagioclase (Figure 11a).

Amphibole has a strong CPO, with the (100) and [001] subparallel to the foliation and to the stretching lineation, respectively (Figures 10c and 11c). The misorientation angle distribution of amphibole displays the strongest peaks between 2 and 25° and ~50 and 180°, with higher values for the correlated pairs (Figure 11d). The misorientation axes of amphibole, plotted in crystal coordinates, show maxima around the c-axis for misorientations up to 30°, which, in sample coordinates, is oriented subparallel to the stretching lineation of the mylonite (Figure 11d).


## 5 | DISCUSSION

### 5.1 | *P-T* conditions of metamorphism and deformation

The Anderson and Smith (1995) geobarometer calibration was based on rocks that equilibrated in the presence of

melt. There is no evidence of melt in our sample; therefore, we favour the pressure results from Bhadra and Bhattacharya (2007; Table 4). However, both results are kept as the Plagioclase Hornblende Thermobarometry spreadsheet (Anderson et al., 2008) that were used to compute *P-T* utilizes a convergence between the results of Anderson and Smith (1995) geobarometer and Holland and Blundy (1994) geothermometer. As a test, the Bhadra and Bhattacharya (2007) pressure results were used as input for the Holland and Blundy (1994) geothermometer and coincident temperature results were obtained, within error.

The oldest metamorphic stage preserved in the sample is represented by the albite cores (P11) with their oriented inclusions of epidote, ilmenite, and rutile. In particular, rutile inclusions suggest relatively HP, but the coexistence with albite crystals would constrain the maximum pressure of this metamorphic stage below the albite breakdown reaction (albite=jadeite+quartz; Newton & Smith, 1967). However, the exact conditions of this first metamorphic stage were not constrained in this study.



**FIGURE 10** Pole figures of the crystallographic orientation data of Pl (colour coding as in the grain size maps) and Amp; same site as maps shown in Figure 8. X is the extensional instantaneous stretching axis, Z is the pole of foliation, stereographic projections, lower hemisphere if not specified; U: upper hemisphere, L: lower hemisphere. (a) Subset of the Pl porphyroclast and adjacent grains sharing the same crystallographic orientation, with the exception of a few data points. The Pl displays a CPO with the (100) and (001) maxima approximately at 45° from X. The (010) maximum is perpendicular to X. a: 189029 Pl data points. (b) Subset of the Pl crystals of the matrix (PI2 and PI3) not adjacent to the porphyroclast. A weak CPO similar to Figure 11a is visible. 34676 Pl data points. (c) Amp displaying a CPO with the (100) and [001] perpendicular and parallel to X, respectively. 671 Amp data points (one point per grain).  $n$  = number of grains. Half width 10° and cluster size 5°, maximum value is given. Contouring is 1

The microstructural relationships between Pl1 and Amp1 are equivocal; therefore, it is difficult to prove that they grew in equilibrium. Their compositions suggest that had


they grown in equilibrium, they would have crystallized at 400°C (Table 4). These results, however, are far away from the lower limits of the calibration of Bhadra and



Bhattacharya (2007; 650–950°C) and Anderson and Smith (1995; 675–760°C): for this reason, pressures were not computed for the Amp1-P11 pairs. Furthermore, P11 is out of the compositional range of the Bhadra and Bhattacharya (2007) equilibration (table 10 of that contribution).

Within uncertainty, the temperature results of Amp1-P12 and Amp2-P13 pairs lie just within the lower limits of the two calibrations. The  $P$ – $T$  estimates from the Amp1-P12 and Amp2-P13 pairs are coincident within error ( $\pm 0.2$  GPa and  $\pm 50^\circ\text{C}$ ). This is due to the subtle chemical differences existing between these two generations. Thermobarometric calculations suggest that P12 may have grown synchronously with Amp1 at 615°C and 0.97 GPa. The crystallization of P12 marks the first stage of the mylonitic foliation development, as evidenced by the presence of P12 in the pressure shadows of P11 (Figures 3, 5a and 7). The mylonitic foliation continued to develop under conditions of  $\sim 600^\circ\text{C}$  and 0.75 GPa (Amp2-P13 pair), as supported by the microtextural observation of Amp2, P13, and Ch11 intergrowths within the P11 pressure shadows (Section 4.1). This temperature result is coincident, within error, with the Chlorite+Quartz+H<sub>2</sub>O thermometry results for Ch11 (550°C; higher temperature peak in Figure 6a).

The  $X_{\text{An}}$  increase from P11 to P12 and P13 could reflect the following reaction from Apter and Liou (1983) for a pressure of 0.7 GPa:



The abundant epidote inclusions in P11 and its corroded aspect in the matrix of the sample would support this hypothesis. P12 likely crystallized at  $>600^\circ\text{C}$  in the (high- $T$ ) epidote amphibolite facies field (Apter & Liou, 1983; Liou, Kuniyoshi, & Ito, 1974; Miyashiro, 1968; Winkler, 1980), and P13 at lower temperatures, as the associated Amp2 and Ch11 limit the maximum temperature to 550–600°C (Figures 5 and 6a). A final retrograde stage is recorded by the Ch12 compositions suggesting a range of 350–200°C (using the Chlorite+Quartz+H<sub>2</sub>O thermometry, Figure 6b), and by the overprinted Ch11 values yielding the low- $T$  peak of Figure 6a. Indeed, Ch12 appears to be associated with the development of  $C'$  shear bands overprinting the mylonitic foliation.

## 5.2 | The origin of luminescence in plagioclase

Backscattered electron and CL images provide important insight into feldspar textures (e.g., Lee, Parsons, Edwards, & Martin, 2007; Parsons & Lee, 2009; Parsons, Steele, Lee, & Magee, 2008). Plagioclase luminescence has been related to many causes (summarized in Götze, 2012): the






Mn<sup>2+</sup>, Ti, Fe<sup>3+</sup> content (Götze, Habermann, Kempe, Neuser, & Richter, 1999; Mariano & King, 1975), trace and REE elements (Götze, Habermann, Neuser, & Richter, 1999; Mariano & King, 1975), lattice defects (e.g., Al–O–Al bridge; Finch & Klein, 1999), and/or mineral inclusions (Smith & Stenstrom, 1965). The CL signal is most commonly linked to Ti concentrations (Lee et al., 2007; Parsons et al., 2008), with a contribution from Fe (Lee et al., 2007). Furthermore, Parsons et al. (2008) has noted that CL emissions appear to be linked to Ca zoning, but they did not find a direct link between Ca concentration (or other trace element concentrations) and CL intensity.

In our samples, TiO<sub>2</sub> appears uniform across all Pl generations (at 0.01 wt%; Table 2). This is also the case for the MnO, with a scatter between 0.002 and 0.006 wt%. The FeO (Fe total) in P11 and P12 is the same (0.03 wt%; Table 2), but increases in P13 (0.06 wt%; Table 2). The CL signal in our samples appears to be related to the Ca/Na ratio (with maybe some contribution from Fe): the higher the ratio, the brighter the growth zone, as visible from the perfect match between the SEM-CL and EPMA compositional maps (Figures 3 and 7), even though we cannot rule out a contribution from trace and REE elements or lattice defects. Luminescence induced by calcite inclusions in P12 and P13 can be excluded, as the calcite veins in the sample do not luminesce.


## 5.3 | Replacement reactions by coupled dissolution–precipitation and deformation by dissolution–precipitation creep in plagioclase and amphibole

Our analysis highlights a strong correlation between CL images and both BSE images and EPMA compositional maps of plagioclase. Textural and chemical features similar to those reported in this study are recognized in several amphibolites present over  $>500$  m of the COSC-1 core (from  $\sim 1,600$  to  $\sim 2,300$  m depth), implying that the deformation mechanisms and the mineral replacement reactions discussed below are important for the development of the middle to lower crustal thrusts and associated tectonic transport during Caledonian nappe stacking.

The plagioclase porphyrocrysts deformation history is summarized in Figure 12a. During the first stage P11 grew including a foliation defined by quartz, epidote, ilmenite, and rutile. Subsequently, fractures developed along the (001) cleavage planes (Ague, 1988; Brander et al., 2012; Brown & Macaudière, 1984; McLaren & Pryer, 2001) enhanced fluid infiltration that triggered replacement reactions occurring via coupled dissolution of P11 and precipitation of P12 (Brander et al., 2012; Marti, Stünitz, Heilbronner, Plümper, & Drury, 2017) in equilibrium with the changed  $P$ – $T$  conditions of 0.9 GPa and 615°C. These



**FIGURE 11** Pole figures of the crystallographic orientation data of Pl (colour coding as in the grain size maps) and Amp; same site as maps shown in Figure 7. (a) The Pl displays a CPO with the (100) and (001) at low to medium angles to X. Note how the maximum of the values coincides with the porphyroclast values, in red. 357353 Pl data points. (b) Subset of the Pl porphyroclast and adjacent grains; some of the latter display a similar crystallographic orientation as the porphyroclast. 238333 Pl data points (c) Pole figures of Amp displaying a strong CPO with the (100) and [001] perpendicular and parallel to X, respectively. 140 Amp data points (one point per grain).  $n$  = number of grains. Half width  $10^\circ$  and cluster size  $5^\circ$ , maximum value is given. Contouring is 2. (d) Histogram of distribution of misorientation angles and misorientation axes of amphibole plotted in crystal coordinates. 3295 Amp data points. Dashed line: forbidden zone limit



**FIGURE 12** Idealized sketch of the deformation history of Pl and Amp, colour coding as in the compositional maps. (a) 1: P11 porphyrocrysts grew including a foliation marked by Ep, Qz, and Ilm. 2: Fracturing of the P11 porphyrocrysts, mostly imposed along the (001) plane. 3: Fluid infiltration triggered mineral replacement by coupled dissolution precipitation with topotaxial and pseudomorphic growth of P12 and, successively, P13 on P11. Replacement occurred mostly along and in proximity of the fractures as well as at the edges of the crystals. P13 nucleated also as newly grown grains in the matrix. (b) 1: Amp1 grew with a CPO. 2: Replacement of Amp1 by Amp2 by coupled dissolution precipitation; topotaxial growth is suggested by the same CPO shared by the two generations. 3: Development of misorientation due to displacement along the fractures and along the cleavage planes

replacement processes continued precipitating P13, which rimmed P11 and P12 and sealed the fractures in P11. The slightly different chemistry of P13 compared to P12 and the microstructural observation that P13 overgrows P12 suggest that these two growth zones were closely related in time or crystallization reaction (Section 4.2).

The growth zones of P12 and P13 inherit the crystallographic orientation of the P11 cores, with maximum misorientation of  $5^\circ$  (Figure 7c,d). Thus, the precipitation of P12 and P13 on P11 is an example of pseudomorphic and topotaxial growth, as often observed during coupled dissolution precipitation processes (Engvik et al., 2008; Hövelmann, Putnis, Geisler, Schmidt, & Golla-Schindler, 2010; Plümper et al., 2017; Putnis & Putnis, 2007; Spruzeniece, Piazzolo, &

Maynard-Casely, 2017). The small (generally  $<2^\circ$ ) and only local difference in crystallographic orientation between P11 and P12-P13 presumably results from the slightly different unit cell parameters between albite and oligoclase. The few low-angle boundaries associated with P12-P13 growth zones are typically found along intracrystalline fractures (Figure 7d), and we interpret them as evidence of slightly rotated fractured fragments of P11 that re-equilibrated to P12-P13 compositions via coupled dissolution precipitation processes. Pore trails occur in the plagioclase porphyrocrysts parallel to the two fracture systems, together with several micrometric mineral inclusions resulting in a turbid aspect of the plagioclase (Figures 3a,b and 4). Those trails probably represented the main pathways of fluid circulation during

coupled P11 dissolution and P12 and P13 precipitation. These mineral inclusions may represent the signature of transient porosity during plagioclase replacement reactions (e.g., Plümper & Putnis, 2009; Plümper et al., 2017; Putnis, 2015; Walker, Lee, & Parsons, 1995).

Crystallographic continuity indicative of epitaxial overgrowth is also observed between plagioclase porphyrocrysts and P12 and P13 grains around them, but to different extents (compare Figure 10a with Figure 11b). Thus, the crystallographic orientation of new P12 and P13 grains may be inherited from the P11 parent grain due to epitaxy, as described for several minerals deforming by dissolution precipitation creep at different crustal levels (Engvik et al., 2008; Imon et al., 2002, 2004; Jiang, Prior, & Wheeler, 2000; Mukai et al., 2014; Spruzeniec et al., 2017; Wassmann & Stöckhert, 2012; Wassmann et al., 2011). The crystallographic continuity seems to decrease away from the plagioclase porphyrocrysts (Figures 10b and 11a), presumably reflecting the heterogeneous nucleation of P12 and P13 neoblasts in the surrounding matrix together with Amp2 and Chl1. Additionally, these neoblasts may have also undergone grain-boundary sliding during deformation, given their fine grain size, which can further disperse their inherited crystallographic orientation (Okudaira et al., 2017).

Thus, we conclude that metamorphism of plagioclase at ~600°C and 0.75–0.9 GPa occurred by coupled dissolution precipitation processes with pseudomorphic and topotaxial replacement of P11 by P12 and P13, and deformation was accommodated by dissolution precipitation creep with nucleation of P12–P13 grains around the plagioclase porphyrocrysts (epitaxial) and in the surrounding matrix. It is worth noting that the replacement of P11 porphyroclasts by coupled dissolution precipitation processes generally occurred concentrically (Figure 7). Consequently, at least part of this replacement process outlasted the deformation. The similar chemistry of P13 replacing P11 and the P13 neoblasts in the surrounding matrix suggest that there was no major change in metamorphic conditions throughout the timing of growth of P13.

No significant contribution of crystal plasticity was observed, although deformation occurred at  $P$ – $T$  conditions at which crystal plasticity in plagioclase is expected to occur (e.g., Gerald & Stünitz, 1993; Pearce, Wheeler, & Prior, 2011). The porosity necessary to maintain fluid transport during coupled dissolution precipitation was generated mostly by fracturing, as shown by the common occurrence of P12–P13 growth zones along fractures parallel to the (001) perfect cleavage planes. Finally, the sharp chemical transition (across ~1–5  $\mu\text{m}$ ) between all the plagioclase generations, visible from the EPMA and CL maps (Figures 3 and 7), suggest that chemical equilibration did not occur by solid-state diffusion (e.g., Hövelmann et al., 2010).

The textural features of amphibole suggest a similar deformation history to plagioclase (Figure 12b). Amp1 (high Mg#) grains are preserved mostly as relict cores and display embayments, lobate edges, and truncated chemical zoning patterns (Figures 3f and 9c; Bukovská, Wirth, & Morales, 2015; Gratier et al., 2013; Hyppolito, García-Casco, Juliani, Meira, & Hall, 2014; Passchier & Trouw, 1996; Rutter, 1983; Stokes et al., 2012; Wassmann & Stöckhert, 2013; Wintsch & Yi, 2002). These textural features suggest that Amp1 underwent coupled dissolution and Amp2 precipitated on Amp1. As for plagioclase, the growth of Amp2 on Amp1 was pseudomorphic and topotaxial (Figure 9d–f). Amp2 grew also as smaller neoblasts (maximum few hundreds of  $\mu\text{m}$  in size), elongated parallel to the mylonitic foliation, and preferentially elongated parallel to their  $c$ -axis, due to dissolution precipitation creep. These crystals do not display Amp1 cores in the compositional maps (Figures 3f and 5b).

The CPO and shape-preferred orientation of the amphibole can be acquired via different mechanisms: dissolution precipitation creep (Bons & den Brok, 2000; Imon et al., 2004; Pearce et al., 2011), oriented grain growth and passive rotation after growth (Berger & Stünitz, 1996; Kanagawa, Shimano, & Hiroi, 2008), and/or diffusion creep (Getsinger & Hirth, 2014). In the studied sample, the CPO displayed by Amp2 is mostly inherited due to the pseudomorphic and topotaxial growth on Amp1. The small misorientations (<3°) evident in the EBSD map (Figure 9f) are attributed to fractures that developed perpendicular to the crystal elongation (Figure 9a). Some amphibole crystals display more complex zoning (e.g., central crystal in Figure 9c), probably due to a preferential replacement of the central part of the crystal (e.g., Hyppolito et al., 2014). The CPO of Amp1 grains was presumably formed via oriented grain growth during an earlier deformation event. Moreover, as presented in Section 4.6, the misorientation axes of amphibole show maxima around the  $c$ -axis, which is oriented subparallel to the stretching lineation of the mylonite (Figure 11d). This geometry is not consistent with dislocation creep on the prism  $\langle c \rangle$  slips system of amphibole, as the misorientation axis cannot coincide with the Burgers vector (e.g., Kruse, Stünitz, & Kunze, 2001; Lloyd, Farmer, & Mainprice, 1997). Instead, we interpret the cluster of misorientation axes around  $\langle c \rangle$  as evidence of topotaxial growth of elongated amphibole grains that preferentially share their  $c$ -axis. As the data come from all the grains included in the map (and not only from the rims of Amp2), we think that oriented growth was the dominant deformation mechanisms of amphibole throughout the deformation history.

In the studied sample, amphibole, like plagioclase, shows no evidence of deformation via crystal plasticity, such as high intracrystalline misorientations, misorientation

bands, subgrains. Instead, it appears to have deformed by fracturing and coupled dissolution precipitation, as also suggested in other studies (Berger & Stünitz, 1996; Brodie & Rutter, 1985; Lafrance & Vernon, 1993; Nyman, Law, & Smelik, 1992; Pearce et al., 2011). Crystal plasticity is potentially a more effective deformation mechanism at higher temperatures (e.g., Skrotzki, 1992).

It is worth noting that the synkinematic reaction that produced Pl<sub>2</sub>-Pl<sub>3</sub> and Amp<sub>2</sub> (reaction 1) could have been a dehydration reaction. If this was the case, the aqueous fluid necessary to sustain coupled dissolution precipitation processes did not necessarily infiltrate from an external source, but may have been released internally. The role of dehydration reactions in the rheological evolution of crustal rocks has received little attention so far, as reaction weakening is commonly associated with hydration reactions during retrogression (e.g., Gueydan, Leroy, Jolivet, & Agard, 2003). However, dehydration reactions that release fluids at grain boundaries can also potentially result in weakening and strain localization during burial and nappe stacking if the released fluids facilitate the activation of coupled dissolution precipitation creep and of diffusion creep.

## 6 | CONCLUSIONS

Middle to lower crustal mylonites from the COSC-1 drill core (Lower Seve Nappe) were investigated with EPMA compositional maps, CL images, and EBSD maps to constrain the mechanism(s) responsible for their formation. The data suggest that fracturing, coupled dissolution precipitation, and dissolution precipitation creep were responsible for the development of the mylonitic fabric in amphibolites at conditions of ~600°C, 0.75–0.97 GPa, in the epidote amphibolite facies, over a thickness >500 m. No evidence of deformation via crystal plasticity is present in either plagioclase or amphibole in the analysed sample, even though deformation occurred at *P-T* conditions at which plagioclase is expected to deform by dislocation creep. The presence of H<sub>2</sub>O-rich fluid at the grain boundaries appears to have enhanced replacement reactions and facilitated dissolution and precipitation processes, which in turn considerably decreased the strength of this middle to lower crustal shear zone. Importantly, replacement reactions of plagioclase only occurred by coupled dissolution precipitation at grain boundaries and along fractures and were otherwise sluggish.

Our study shows that crystallographic preferred orientation in plagioclase and amphibole can be inherited from parental grains due to pseudomorphic and topotaxial growth during coupled dissolution precipitation during deformation. Thus, care must be taken when considering

CPOs in deformed rocks as evidence of deformation by dislocation creep.

The development of a mylonitic fabric by coupled dissolution precipitation and dissolution precipitation creep in amphibolites over a thickness >500 in the Lower Seve Nappe suggests that the strength of amphibolites can be significantly low in the presence of grain-boundary aqueous fluid during nappe thrusting in the middle to lower crust.

## ACKNOWLEDGEMENTS

S. Hammond and D. Johnson are warmly thanked for the help with EPMA and SEM analyses, respectively. The staff at Plymouth University Electron Microscopy Centre is acknowledged for the support during EBSD analysis. B. Almqvist is acknowledged for his support in acquiring the sample. We acknowledge constructive comments and suggestions from M.A. Pearce, R. Wintsch, and two anonymous referees that improved the quality of the manuscript. We are grateful to M. Brown for editorial handling. This work was funded by an Early Postdoc.Mobility funding scheme (project number: P2BEP2 168722) granted by Swiss National Science Foundation. L.M. acknowledges financial support from a FP7 Marie Curie Career Integration Grant (grant agreement PCIG13-GA-2013-618289). The COSC-1 drilling operations were supported by the International Continental Scientific Drilling Program (ICDP) and the Swedish Research Council (Vetenskapsrådet), and the Swedish national research infrastructure for scientific drilling “Riksriggen”.

## ORCID

Francesco Giuntoli <http://orcid.org/0000-0003-1999-5725>

## REFERENCES

- Ague, D. M. (1988). Universal stage measurements and transmission electron microscope observations of fractured plagioclase. *Journal of Structural Geology*, 10(7), 701–705. [https://doi.org/10.1016/0191-8141\(88\)90077-6](https://doi.org/10.1016/0191-8141(88)90077-6)
- Anderson, J. L., Barth, A. P., Wooden, J. L., & Mazdab, F. (2008). Thermometers and thermobarometers in granitic systems. *Reviews in Mineralogy and Geochemistry*, 69(1), 121–142. <https://doi.org/10.2138/rmg.2008.69.4>
- Anderson, J. L., & Smith, D. R. (1995). The effects of temperature and fO<sub>2</sub> on the Al in hornblende barometer. *American Mineralogist*, 80(5–6), 549–559. <https://doi.org/10.2138/am.1995.5.614>
- Andréasson, P. (1994). The Baltoscandian margin in Neoproterozoic early Palaeozoic times. Some constraints on terrane derivation and accretion in the Arctic Scandinavian Caledonides. *Tectonophysics*, 231(1–3), 1–32. [https://doi.org/10.1016/0040-1951\(94\)90118-X](https://doi.org/10.1016/0040-1951(94)90118-X)

- Apted, M. J., & Liou, J. (1983). Phase relations among greenschist, epidote amphibolite, and amphibolite in a basaltic system. *American Journal of Science*, 283, 328–354.
- Arnbom, J. O. (1980). Metamorphism of the Seve Nappes at Åreskutan, Swedish Caledonides. *Geologiska Föreningen i Stockholm Förhandlingar*, 102(4), 359–371. <https://doi.org/10.1080/11035898009454493>
- Berger, A., & Stünitz, H. (1996). Deformation mechanisms and reaction of hornblende: Examples from the Bergell tonalite (Central Alps). *Tectonophysics*, 257(2), 149–174. [https://doi.org/10.1016/0040-1951\(95\)00125-5](https://doi.org/10.1016/0040-1951(95)00125-5)
- Bergman, S., & Sjöström, H. (1997). Accretion and lateral extension in an orogenic wedge: Evidence from a segment of the Seve Kōlli terrane boundary, central Scandinavian Caledonides. *Journal of Structural Geology*, 19(8), 1073–1091. [https://doi.org/doi:10.1016/S0191-8141\(97\)00028-X](https://doi.org/doi:10.1016/S0191-8141(97)00028-X)
- Bhadra, S., & Bhattacharya, A. (2007). The barometer tremolite+ tschermakite+ 2 albite = 2 pargasite+ 8 quartz: Constraints from experimental data at unit silica activity, with application to garnet free natural assemblages. *American Mineralogist*, 92(4), 491–502. <https://doi.org/10.2138/am.2007.2067>
- Bons, P. D., & den Brok, B. (2000). Crystallographic preferred orientation development by dissolution precipitation creep. *Journal of Structural Geology*, 22(11), 1713–1722. [https://doi.org/10.1016/S0191-8141\(00\)00075-4](https://doi.org/10.1016/S0191-8141(00)00075-4)
- Brander, L., Svahnberg, H., & Piazzolo, S. (2012). Brittle plastic deformation in initially dry rocks at fluid present conditions: Transient behaviour of feldspar at mid crustal levels. *Contributions to Mineralogy and Petrology*, 163(3), 403–425. <https://doi.org/10.1007/s00410-011-0677-5>
- Brodie, K., & Rutter, E. (1985). On the relationship between deformation and metamorphism, with special reference to the behavior of basic rocks. In A. B. Thompson, & D. C. Rubie (Eds.), *Metamorphic reactions: kinetics, textures, and deformation* (pp. 138–179). New York, NY: Springer. <https://doi.org/10.1007/978-1-4612-5066-1>
- Brown, W. L., & Macaudière, J. (1984). Microfracturing in relation to atomic structure of plagioclase from a deformed meta anorthosite. *Journal of Structural Geology*, 6(5), 579–586. [https://doi.org/10.1016/0191-8141\(84\)90067-1](https://doi.org/10.1016/0191-8141(84)90067-1)
- Brueckner, H. K., & van Roermund, H. L. M. (2004). Dunk tectonics: A multiple subduction/eduction model for the evolution of the Scandinavian Caledonides. *Tectonics*, 23(2), TC2004. <https://doi.org/10.1029/2003TC001502>
- Brueckner, H. K., & Van Roermund, H. L. (2007). Concurrent HP metamorphism on both margins of Iapetus: Ordovician ages for eclogites and garnet pyroxenites from the Seve Nappe Complex, Swedish Caledonides. *Journal of the Geological Society*, 164(1), 117–128. <https://doi.org/10.1144/0016-76492005-139>
- Bukovská, Z., Wirth, R., & Morales, L. F. (2015). Pressure solution in rocks: Focused ion beam/transmission electron microscopy study on orthogneiss from South Armorican Shear Zone, France. *Contributions to Mineralogy and Petrology*, 170(3), 31. <https://doi.org/10.1007/s00410-015-1186-8>
- Bürgmann, R., & Dresen, G. (2008). Rheology of the lower crust and upper mantle: Evidence from rock mechanics, geodesy, and field observations. *Annual Review of Earth and Planetary Sciences*, 36, 531–567
- Carmichael, D. M. (1969). On the mechanism of prograde metamorphic reactions in quartz bearing pelitic rocks. *Contributions to Mineralogy and Petrology*, 20(3), 244–267. <https://doi.org/10.1007/BF00377479>
- Engvik, A. K., Putnis, A., Gerald, J. D. F., & Austrheim, H. (2008). Albitization of granitic rocks: The mechanism of replacement of oligoclase by albite. *The Canadian Mineralogist*, 46(6), 1401–1415. <https://doi.org/10.3749/canmin.46.6.1401>
- Finch, A. A., & Klein, J. (1999). The causes and petrological significance of cathodoluminescence emissions from alkali feldspars. *Contributions to Mineralogy and Petrology*, 135(2–3), 234–243. <https://doi.org/10.1007/s004100050509>
- Fussey, F., & Handy, M. (2008). Micromechanisms of shear zone propagation at the brittle–viscous transition. *Journal of Structural Geology*, 30(10), 1242–1253. <https://doi.org/10.1016/j.jsg.2008.06.005>
- Fussey, F., Regenauer-Lieb, K., Liu, J., Hough, R., & De Carlo, F. (2009). Creep cavitation can establish a dynamic granular fluid pump in ductile shear zones. *Nature*, 459(7249), 974–977. <https://doi.org/10.1038/nature08051>
- Gayer, R., Rice, A., Roberts, D., Townsend, C., & Welbon, A. (1987). Restoration of the Caledonian Baltoscandian margin from balanced cross sections: The problem of excess continental crust. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 78(3), 197–217. <https://doi.org/10.1017/S026359330001110X>
- Gee, D. (1975). A tectonic model for the central part of the Scandinavian Caledonides. *American Journal of Science*, 275(A), 468–515.
- Gee, D., Fossen, H., Henriksen, N., & Higgins, A. K. (2008). From the early Paleozoic platforms of Baltica and Laurentia to the Caledonide Orogen of Scandinavia and Greenland. *Episodes*, 31(1), 44–51.
- Gee, D., Janák, M., Majka, J., Robinson, P., & van Roermund, H. (2013). Subduction along and within the Baltoscandian margin during closing of the Iapetus Ocean and Baltica–Laurentia collision. *Lithosphere*, 5(2), 169–178. <https://doi.org/10.1130/L220.1>
- Gee, D., Juhlin, C., Pascal, C., & Robinson, P. (2010). Collisional orogeny in the Scandinavian Caledonides (COSC). *GFF*, 132(1), 29–44. <https://doi.org/10.1080/11035891003759188>
- Gee, D., & Sturt, B. (1985). *The Caledonide orogen: Scandinavia and related areas (Vol. 2)*. Chichester, UK: Wiley.
- Gerald, J. F., & Stünitz, H. (1993). Deformation of granitoids at low metamorphic grade. I: Reactions and grain size reduction. *Tectonophysics*, 221(3), 269–297. [https://doi.org/10.1016/0040-1951\(93\)90163-E](https://doi.org/10.1016/0040-1951(93)90163-E)
- Getsinger, A., & Hirth, G. (2014). Amphibole fabric formation during diffusion creep and the rheology of shear zones. *Geology*, 42(6), 535–538. <https://doi.org/10.1130/G35327.1>
- Gilio, M., Clos, F., & van Roermund, H. L. (2015). The Friningen garnet peridotite (central Swedish Caledonides). A good example of the characteristic PTt path of a cold mantle wedge garnet peridotite. *Lithos*, 230, 1–16. <https://doi.org/10.1016/j.lithos.2015.05.003>
- Gilotti, J. A. (1989). Reaction progress during mylonitization of basalitic dikes along the Särvi thrust, Swedish Caledonides. *Contributions to Mineralogy and Petrology*, 101(1), 30–45. <https://doi.org/10.1007/BF00387199>
- Giuntoli, F., Lanari, P., & Engi, M. (2018). Deeply subducted continental fragments – Part 1: Fracturing, dissolution precipitation, and diffusion processes recorded by garnet textures of the central Sesia Zone (western Italian Alps). *Solid Earth*, 9(1), 167–189. <https://doi.org/10.5194/se-9-167-2018>
- Götze, J. (2012). Application of cathodoluminescence microscopy and spectroscopy in geosciences. *Microscopy and Microanalysis*, 18(6), 1270–1284. <https://doi.org/10.1017/S1431927612001122>

- Götze, J., Habermann, D., Kempe, U., Neuser, R. D., & Richter, D. K. (1999a). Cathodoluminescence microscopy and spectroscopy of plagioclases from lunar soil. *American Mineralogist*, *84*(7–8), 1027–1032. <https://doi.org/10.2138/am.1999.7.804>
- Götze, J., Habermann, D., Neuser, R. D., & Richter, D. K. (1999b). High resolution spectrometric analysis of rare earth elements activated cathodoluminescence in feldspar minerals. *Chemical Geology*, *153*(1), 81–91. [https://doi.org/10.1016/S0009-2541\(98\)00153-3](https://doi.org/10.1016/S0009-2541(98)00153-3)
- Gratier, J. P., Dysthe, D. K., & Renard, F. (2013). The role of pressure solution creep in the ductility of the Earth's upper crust. *Advances in Geophysics*, *54*, 47–179. <https://doi.org/10.1016/B978-0-12-380940-7.00002-0>
- Grimmer, J. C., Glodny, J., Drüppel, K., Greiling, R. O., & Kontny, A. (2015). Early to mid Silurian extrusion wedge tectonics in the central Scandinavian Caledonides. *Geology*, *43*(4), 347–350. <https://doi.org/10.1130/G36433.1>
- Gueydan, F., Leroy, Y. M., Jolivet, L., & Agard, P. (2003). Analysis of continental midcrustal strain localization induced by microfracturing and reaction softening. *Journal of Geophysical Research: Solid Earth*, *108*(B2), 2064. <https://doi.org/10.1029/2001JB000611>
- Hedin, P., Almqvist, B., Berthet, T., Juhlin, C., Buske, S., Simon, H., ... Alm, P. G. (2016). 3D reflection seismic imaging at the 2.5 km deep COSC 1 scientific borehole, central Scandinavian Caledonides. *Tectonophysics*, *689*, 40–55. <https://doi.org/10.1016/j.tecto.2015.12.013>
- Holland, T., & Blundy, J. (1994). Non ideal interactions in calcic amphiboles and their bearing on amphibole plagioclase thermometry. *Contributions to Mineralogy and Petrology*, *116*(4), 433–447. <https://doi.org/10.1007/BF00310910>
- Hövelmann, J., Putnis, A., Geisler, T., Schmidt, B. C., & Golla-Schindler, U. (2010). The replacement of plagioclase feldspars by albite: Observations from hydrothermal experiments. *Contributions to Mineralogy and Petrology*, *159*(1), 43–59. <https://doi.org/10.1007/s00410-009-0415-4>
- Hypolito, T., García Casco, A., Juliani, C., Meira, V., & Hall, C. (2014). Late Paleozoic onset of subduction and exhumation at the western margin of Gondwana (Chilena Terrane): Counterclockwise P–T paths and timing of metamorphism of deep seated garnet mica schist and amphibolite of Punta Sirena, Coastal Accretionary Complex, central Chile (34°S). *Lithos*, *206*, 409–434. <https://doi.org/10.1016/j.lithos.2014.07.023>
- Imon, R., Okudaira, T., & Fujimoto, A. (2002). Dissolution and precipitation processes in deformed amphibolites: An example from the ductile shear zone of the Ryoke metamorphic belt, SW Japan. *Journal of Metamorphic Geology*, *20*(3), 297–308. <https://doi.org/10.1046/j.1525-1314.2002.00367.x>
- Imon, R., Okudaira, T., & Kanagawa, K. (2004). Development of shape and lattice preferred orientations of amphibole grains during initial cataclastic deformation and subsequent deformation by dissolution precipitation creep in amphibolites from the Ryoke metamorphic belt, SW Japan. *Journal of Structural Geology*, *26*(5), 793–805. <https://doi.org/10.1016/j.jsg.2003.09.004>
- Jackson, J. A., Austrheim, H., McKenzie, D., & Priestley, K. (2004). Metastability, mechanical strength, and the support of mountain belts. *Geology*, *32*(7), 625–628. <https://doi.org/10.1130/G20397.1>
- Janák, M., van Roermund, H., Majka, J., & Gee, D. (2013). UHP metamorphism recorded by kyanite bearing eclogite in the Seve Nappe Complex of northern Jämtland, Swedish Caledonides. *Gondwana Research*, *23*(3), 865–879. <https://doi.org/10.1016/j.gr.2012.06.012>
- Jiang, Z., Prior, D. J., & Wheeler, J. (2000). Albite crystallographic preferred orientation and grain misorientation distribution in a low grade mylonite: Implications for granular flow. *Journal of Structural Geology*, *22*(11), 1663–1674. [https://doi.org/10.1016/S0191-8141\(00\)00079-1](https://doi.org/10.1016/S0191-8141(00)00079-1)
- Kanagawa, K., Shimano, H., & Hiroi, Y. (2008). Mylonitic deformation of gabbro in the lower crust: A case study from the Panke-nushi gabbro in the Hidaka metamorphic belt of central Hokkaido, Japan. *Journal of Structural Geology*, *30*(9), 1150–1166. <https://doi.org/10.1016/j.jsg.2008.05.007>
- Klonowska, I., Janák, M., Majka, J., Froitzheim, N., & Kościńska, K. (2016). Eclogite and garnet pyroxenite from Stor Jougdan, Seve Nappe Complex, Sweden: Implications for UHP metamorphism of allochthons in the Scandinavian Caledonides. *Journal of Metamorphic Geology*, *34*(2), 103–119. <https://doi.org/10.1111/jmg.12173>
- Klonowska, I., Janák, M., Majka, J., Petřík, I., Froitzheim, N., Gee, D., & Sasínková, V. (2017). Microdiamond in Åreskutan confirms regional UHP metamorphism in the Seve Nappe Complex of the Scandinavian Caledonides. *Journal of Metamorphic Geology*, *35*, 541–564. <https://doi.org/10.1111/jmg.12244>
- Kruse, R., Stünitz, H., & Kunze, K. (2001). Dynamic recrystallization processes in plagioclase porphyroclasts. *Journal of Structural Geology*, *23*(11), 1781–1802. [https://doi.org/10.1016/S0191-8141\(01\)00030-X](https://doi.org/10.1016/S0191-8141(01)00030-X)
- Ladenberger, A., Be'eri Shlevin, Y., Claesson, S., Gee, D. G., Majka, J., & Romanova, I. V. (2013). Tectonometamorphic evolution of the Åreskutan Nappe Caledonian history revealed by SIMS U–Pb zircon geochronology. *Geological Society, London, Special Publications*, *390*, SP390.310.
- Lafrance, B., & Vernon, R. (1993). Mass transfer and microfracturing in gabbroic mylonites of the Guadalupe igneous complex, California. In J. N. Boland, & J. D. Fitz Gerald (Eds.), *Defects and processes in the solid state: Geoscience applications: The McLaren volume (Developments in Petrology, Vol. 4)* (pp. 151–167). Amsterdam, The Netherlands: Elsevier Science.
- Lanari, P. (2012). *Micro cartographie P–T e dans les roches métamorphiques. Applications aux Alpes et à l'Himalaya*. Doctorate PhD, Grenoble, France.
- Lanari, P., Guillot, S., Schwartz, S., Vidal, O., Tricart, P., Riel, N., & Beyssac, O. (2012). Diachronous evolution of the alpine continental subduction wedge: Evidence from P–T estimates in the Briançonnais Zone houillère (France–Western Alps). *Journal of Geodynamics*, *56–57*, 39–54. <https://doi.org/10.1016/j.jog.2011.09.006>
- Lanari, P., Riel, N., Guillot, S., Vidal, O., Schwartz, S., Pêcher, A., & Hattori, K. H. (2013). Deciphering high pressure metamorphism in collisional context using microprobe mapping methods: Application to the Stak eclogitic massif (northwest Himalaya). *Geology*, *41*(2), 111–114. <https://doi.org/10.1130/G33523.1>
- Lanari, P., Vidal, O., Lewin, E., Dubacq, B., De Andrade, V., & Schwartz, S. (2014). XMapTools a Matlab© based graphic user interface for microprobe quantified image processing. *Computers and Geosciences*, *62*, 227–240. <https://doi.org/10.1016/j.cageo.2013.08.010>
- Leake, B. E., Woolley, A. R., Arps, C. E., Birch, W. D., Gilbert, M. C., Grice, J. D., ... Krivovichev, V. G. (1997). Report.

- Nomenclature of amphiboles: Report of the subcommittee on amphiboles of the international mineralogical association commission on new minerals and mineral names. *Mineralogical Magazine*, 61(2), 295–321. <https://doi.org/10.1180/minmag.1997.061.405.13>
- Lee, M. R., Parsons, I., Edwards, P. R., & Martin, R. W. (2007). Identification of cathodoluminescence activators in zoned alkali feldspars by hyperspectral imaging and electron probe microanalysis. *American Mineralogist*, 92(2–3), 243–253. <https://doi.org/10.2138/am.2007.2160>
- Liou, J., Kuniyoshi, S., & Ito, K. (1974). Experimental studies of the phase relations between greenschist and amphibolite in a basaltic system. *American Journal of Science*, 274(6), 613–632. <https://doi.org/10.2475/ajs.274.6.613>
- Lloyd, G. E., Farmer, A. B., & Mainprice, D. (1997). Misorientation analysis and the formation and orientation of subgrain and grain boundaries. *Tectonophysics*, 279(1–4), 55–78. [https://doi.org/10.1016/S0040-1951\(97\)00115-7](https://doi.org/10.1016/S0040-1951(97)00115-7)
- Lorenz, H., Rosberg, J. E., Juhlin, C., Bjelm, L., Almqvist, B. S. G., Berthet, T., ... Pascal, C. (2015). COSC 1 drilling of a subduction related allochthon in the Palaeozoic Caledonide orogen of Scandinavia. *Scientific Drilling*, 19, 1. <https://doi.org/10.5194/sd-19-1-2015>
- Majka, J., Rosén, Å., Janák, M., Froitzheim, N., Klonowska, I., Maneck, M., ... Yoshida, K. (2014). Microdiamond discovered in the Seve Nappe (Scandinavian Caledonides) and its exhumation by the “vacuum cleaner” mechanism. *Geology*, 42(12), 1107–1110. <https://doi.org/10.1130/G36108.1>
- Mariano, A., & King, P. (1975). Europium activated cathodoluminescence in minerals. *Geochimica et Cosmochimica Acta*, 39(5), 649–660. [https://doi.org/10.1016/0016-7037\(75\)90008-3](https://doi.org/10.1016/0016-7037(75)90008-3)
- Marti, S., Stünitz, H., Heilbronner, R., Plümper, O., & Drury, M. (2017). Experimental investigation of the brittle viscous transition in mafic rocks: Interplay between fracturing, reaction, and viscous deformation. *Journal of Structural Geology*, 105, 62–79. <https://doi.org/10.1016/j.jsg.2017.10.011>
- McAleer, R., Bish, D., Kunk, M., Sicard, K., Valley, P., Walsh, G., ... Wintsch, R. (2017). Reaction softening by dissolution precipitation creep in a retrograde greenschist facies ductile shear zone, New Hampshire, USA. *Journal of Metamorphic Geology*, 35(1), 95–119. <https://doi.org/10.1111/jmg.12222>
- McLaren, A., & Pryer, L. (2001). Microstructural investigation of the interaction and interdependence of cataclastic and plastic mechanisms in feldspar crystals deformed in the semi brittle field. *Tectonophysics*, 335(1), 1–15. [https://doi.org/10.1016/S0040-1951\(01\)00042-7](https://doi.org/10.1016/S0040-1951(01)00042-7)
- Menegon, L., Fousseis, F., Stünitz, H., & Xiao, X. (2015). Creep cavitation bands control porosity and fluid flow in lower crustal shear zones. *Geology*, 43(3), 227–230. <https://doi.org/10.1130/G36307.1>
- Menegon, L., Pennacchioni, G., & Spiess, R. (2008). Dissolution precipitation creep of K feldspar in mid crustal granite mylonites. *Journal of Structural Geology*, 30(5), 565–579. <https://doi.org/10.1016/j.jsg.2008.02.001>
- Miyashiro, A. (1968). Metamorphism of mafic rocks. *Basalts*, 2, 799–834.
- Mouthereau, F., Lacombe, O., & Vergés, J. (2012). Building the Zagros collisional orogen: Timing, strain distribution and the dynamics of Arabia/Eurasia plate convergence. *Tectonophysics*, 532, 27–60. <https://doi.org/10.1016/j.tecto.2012.01.022>
- Mouthereau, F., Watts, A. B., & Burov, E. (2013). Structure of orogenic belts controlled by lithosphere age. *Nature Geoscience*, 6(9), 785–789. <https://doi.org/10.1038/ngeo1902> <http://www.nature.com/ngeo/journal/v6/n9/abs/ngeo1902.html#supplementary-information>
- Mukai, H., Austrheim, H., Putnis, C. V., & Putnis, A. (2014). Textural evolution of plagioclase feldspar across a shear zone: Implications for deformation mechanism and rock strength. *Journal of Petrology*, 55(8), 1457–1477. <https://doi.org/10.1093/petrology/egu030>
- Newton, R. C., & Smith, J. (1967). Investigations concerning the breakdown of albite at depth in the earth. *The Journal of Geology*, 75(3), 268–286. <https://doi.org/10.1086/627255>
- Northrup, C. (1996). Structural expressions and tectonic implications of general noncoaxial flow in the midcrust of a collisional orogen: The northern Scandinavian Caledonides. *Tectonics*, 15(2), 490–505. <https://doi.org/10.1029/95TC02951>
- Nyman, M. W., Law, R. D., & Smelik, E. A. (1992). Cataclastic deformation mechanism for the development of core mantle structures in amphibole. *Geology*, 20(5), 455–458. [https://doi.org/10.1130/0091-7613\(1992\)020<455:CDMFTD>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<455:CDMFTD>2.3.CO;2)
- Okudaira, T., Shigematsu, N., Harigane, Y., & Yoshida, K. (2017). Grain size reduction due to fracturing and subsequent grain size sensitive creep in a lower crustal shear zone in the presence of a CO<sub>2</sub> bearing fluid. *Journal of Structural Geology*, 95, 171–187. <https://doi.org/10.1016/j.jsg.2016.11.001>
- Parsons, I., & Lee, M. R. (2009). Mutual replacement reactions in alkali feldspars I: Microtextures and mechanisms. *Contributions to Mineralogy and Petrology*, 157(5), 641. <https://doi.org/10.1007/s00410-008-0355-4>
- Parsons, I., Steele, D. A., Lee, M. R., & Magee, C. W. (2008). Titanium as a cathodoluminescence activator in alkali feldspars. *American Mineralogist*, 93(5–6), 875–879. <https://doi.org/10.2138/am.2008.2711>
- Passchier, C., & Trouw, R. (1996). *Microtectonics*. Berlin, Germany: Springer Verlag, 289 pp.
- Pearce, M. A., Wheeler, J., & Prior, D. J. (2011). Relative strength of mafic and felsic rocks during amphibolite facies metamorphism and deformation. *Journal of Structural Geology*, 33(4), 662–675. <https://doi.org/10.1016/j.jsg.2011.01.002>
- Plümper, O., Botan, A., Los, C., Liu, Y., Malthe Sørensens, A., & Jamtveit, B. (2017). Fluid driven metamorphism of the continental crust governed by nanoscale fluid flow. *Nature Geoscience*, 10, 685. <https://doi.org/10.1038/ngeo3009> <https://www.nature.com/articles/ngeo3009#supplementary-information>
- Plümper, O., John, T., Podladchikov, Y. Y., Vrijmoed, J. C., & Scambelluri, M. (2017). Fluid escape from subduction zones controlled by channel forming reactive porosity. *Nature Geoscience*, 10(2), 150–156. <https://doi.org/10.1038/ngeo2865> <http://www.nature.com/ngeo/journal/v10/n2/abs/ngeo2865.html#supplementary-information>
- Plümper, O., & Putnis, A. (2009). The complex hydrothermal history of granitic rocks: Multiple feldspar replacement reactions under subsolidus conditions. *Journal of Petrology*, 50(5), 967–987. <https://doi.org/10.1093/petrology/egp028>
- Prior, D. J., Wheeler, J., Peruzzo, L., Spiess, R., & Storey, C. (2002). Some garnet microstructures: An illustration of the potential of orientation maps and misorientation analysis in microstructural studies. *Journal of Structural Geology*, 24(6), 999–1011. [https://doi.org/10.1016/S0191-8141\(01\)00087-6](https://doi.org/10.1016/S0191-8141(01)00087-6)



- Putnis, A. (2002). Mineral replacement reactions: From macroscopic observations to microscopic mechanisms. *Mineralogical Magazine*, 66(5), 689–708. <https://doi.org/10.1180/0026461026650056>
- Putnis, A. (2009). Mineral replacement reactions. *Reviews in Mineralogy and Geochemistry*, 70(1), 87–124. <https://doi.org/10.2138/rmg.2009.70.3>
- Putnis, A. (2015). Transient porosity resulting from fluid mineral interaction and its consequences. *Reviews in Mineralogy and Geochemistry*, 80, 1–23. <https://doi.org/10.2138/rmg.2015.80.01>
- Putnis, A., & Putnis, C. V. (2007). The mechanism of reequilibration of solids in the presence of a fluid phase. *Journal of Solid State Chemistry*, 180(5), 1783–1786. <https://doi.org/10.1016/j.jssc.2007.03.023>
- Rice, A. H. N., & Anderson, M. W. (2016). Restoration of the external Scandinavian Caledonides. *Geological Magazine*, 153, 1136–1165.
- Roberts, D. (2003). The Scandinavian Caledonides: Event chronology, palaeogeographic settings and likely modern analogues. *Tectonophysics*, 365(1), 283–299. [https://doi.org/10.1016/S00401951\(03\)00026X](https://doi.org/10.1016/S00401951(03)00026X)
- Roberts, D., & Gee, D. (1985). An introduction to the structure of the Scandinavian Caledonides. In D. G. Gee, & B. A. Sturt (Eds.), *The Caledonide orogen Scandinavia and related areas*, vol. 1 (pp. 55–68). Chichester, UK: Wiley.
- Root, D., & Corfu, F. (2012). U–Pb geochronology of two discrete Ordovician high pressure metamorphic events in the Seve Nappe Complex, Scandinavian Caledonides. *Contributions to Mineralogy and Petrology*, 163(5), 769–788. <https://doi.org/10.1007/s0041001106980>
- Royden, L. (1996). Coupling and decoupling of crust and mantle in convergent orogens: Implications for strain partitioning in the crust. *Journal of Geophysical Research: Solid Earth*, 101(B8), 17679–17705. <https://doi.org/10.1029/96JB00951>
- Ruiz Agudo, E., Putnis, C., & Putnis, A. (2014). Coupled dissolution and precipitation at mineral fluid interfaces. *Chemical Geology*, 383, 132–146. <https://doi.org/10.1016/j.chemgeo.2014.06.007>
- Rutter, E. (1983). Pressure solution in nature, theory and experiment. *Journal of the Geological Society*, 140(5), 725–740. <https://doi.org/10.1144/gsjgs.140.5.0725>
- Rutter, E., & Brodie, K. (1992). Rheology of the lower crust. In D. M. Fountain, R. Arculus, & R. W. Kay (Eds.), *Continental lower crust* (pp. 201–267). Amsterdam, The Netherlands: Elsevier.
- Sjöström, H. (1983). The Seve–Köli Nappe Complex of the Handölen–Storlien–Essandsjøen area, Scandinavian Caledonides. *Geologiska Föreningen i Stockholm Förhandlingar*, 105(2), 93–117. <https://doi.org/10.1080/11035898309454553>
- Skrotzki, W. (1992). Defect structure and deformation mechanisms in naturally deformed hornblende. *Physica Status Solidi (A)*, 131(2), 605–624. [https://doi.org/10.1002/\(ISSN\)1521396X](https://doi.org/10.1002/(ISSN)1521396X)
- Smith, J., & Stenstrom, R. (1965). Electron excited luminescence as a petrologic tool. *The Journal of Geology*, 73(4), 627–635. <https://doi.org/10.1086/627098>
- Spruzeniece, L., Piazzolo, S., & Maynard Casely, H. E. (2017). Deformation resembling microstructure created by fluid mediated dissolution precipitation reactions. *Nature Communications*, 8, 14032. <https://doi.org/10.1038/ncomms14032>
- Stephens, M. B. (1988). The Scandinavian Caledonides: A complexity of collisions. *Geology Today*, 4(1), 20–26. <https://doi.org/10.1111/j.13652451.1988.tb00537.x>
- Stokes, M., Wintsch, R., & Southworth, C. (2012). Deformation of amphibolites via dissolution precipitation creep in the middle and lower crust. *Journal of Metamorphic Geology*, 30(7), 723–737. <https://doi.org/10.1111/j.15251314.2012.00989.x>
- Strand, T., & Kulling, O. (1972). *Scandinavian caledonides*. Chichester, UK: John Wiley & Sons.
- Strömberg, A., Karis, L., Zachrisson, E., Sjöstrand, T., Skoglund, R., Lundegårdh, P., ... Kornfält, K. (1984). Berggrundskarta över Jämtlands län utom förutvarande Fjällsjö kommun, scale 1: 200 000. Geological Survey of Sweden, Ca, 53.
- Tullis, J., Yund, R., & Farver, J. (1996). Deformation enhanced fluid distribution in feldspar aggregates and implications for ductile shear zones. *Geology*, 24(1), 63–66. [https://doi.org/10.1130/00917613\(1996\)024<063:DEFDIF>2.3.CO;2](https://doi.org/10.1130/00917613(1996)024<063:DEFDIF>2.3.CO;2)
- Van Roermund, H. (1985). Eclogites of the Seve nappe, central Scandinavian Caledonides. In D. G. Gee, & B. A. Sturt (Eds.), *The Caledonide Orogen Scandinavia and related areas* (pp. 873–886). Chichester, UK: John Wiley & Sons
- Van Roermund, H. (1989). High pressure ultramafic rocks from the allochthonous nappes of the Swedish Caledonides. In R. A. Grayer (Ed.), *The Caledonide Geology of Scandinavia* (pp. 205–219). London: Graham and Trotman. <https://doi.org/10.1007/9789400925496>
- Vidal, O., De Andrade, V., Lewin, E., Munoz, M., Parra, T., & Pascarelli, S. (2006). P–T deformation Fe<sup>3+</sup>/Fe<sup>2+</sup> mapping at the thin section scale and comparison with XANES mapping: Application to a garnet bearing metapelite from the Sambagawa metamorphic belt (Japan). *Journal of Metamorphic Geology*, 24(7), 669–683. <https://doi.org/10.1111/j.15251314.2006.00661.x>
- Vidal, O., Lanari, P., Munoz, M., Bourdelle, F., & De Andrade, V. (2016). Deciphering temperature, pressure and oxygen activity conditions of chlorite formation. *Clay Minerals*, 51(4), 615–633. <https://doi.org/10.1180/claymin.2016.051.4.06>
- Vidal, O., Parra, T., & Vieillard, P. (2005). Thermodynamic properties of the Tschermak solid solution in Fe chlorite: Application to natural examples and possible role of oxidation. *American Mineralogist*, 90(2–3), 347–358. <https://doi.org/10.2138/am.2005.1554>
- Walker, F. D. L., Lee, M. R., & Parsons, I. (1995). Micropores and micropore texture in alkali feldspars: Geochemical and geophysical implications. *Mineralogical Magazine*, 59(3), 505–534. <https://doi.org/10.1180/minmag.1995.059.396.12>
- Wallis, D., Phillips, R., & Lloyd, G. (2014). Evolution of the Eastern Karakoram Metamorphic Complex, Ladakh, NW India, and its relationship to magmatism and regional tectonics. *Tectonophysics*, 626, 41–52. <https://doi.org/10.1016/j.tecto.2014.03.023>
- Wassmann, S., & Stöckhert, B. (2012). Matrix deformation mechanisms in HP–LT tectonic mélanges: microstructural record of jadeite blueschist from the Franciscan Complex, California. *Tectonophysics*, 568, 135–153. <https://doi.org/10.1016/j.tecto.2012.01.009>
- Wassmann, S., & Stöckhert, B. (2013). Rheology of the plate interface: dissolution precipitation creep in high pressure metamorphic rocks. *Tectonophysics*, 608, 1–29. <https://doi.org/10.1016/j.tecto.2013.09.030>
- Wassmann, S., Stöckhert, B., & Trepman, C. A. (2011). Dissolution precipitation creep versus crystalline plasticity in high pressure metamorphic serpentinites. *Geological Society, London, Special Publications*, 360(1), 129–149. <https://doi.org/10.1144/SP360.8>
- Wenning, Q. C., Berthet, T., Ask, M., Zappone, A., Rosberg, J. E., & Almqvist, B. S. (2017). Image log analysis of in situ stress orientation, breakout growth, and natural geologic structures to 2.5 km

- depth in central Scandinavian Caledonides: Results from the COSC 1 borehole. *Journal of Geophysical Research: Solid Earth*, 122, 3999–4019
- Whitney, D. L., & Evans, B. W. (2010). Abbreviations for names of rock forming minerals. *American Mineralogist*, 95(1), 185–187. <https://doi.org/10.2138/am.2010.3371>
- Winkler, W. (1980). *Petrogenesis of metamorphic rocks* (5th edn). New York, NY: Springer Verlag, 348pp.
- Wintsch, R., & Yi, K. (2002). Dissolution and replacement creep: A significant deformation mechanism in mid crustal rocks. *Journal of Structural Geology*, 24(6), 1179–1193. [https://doi.org/10.1016/S0191-8141\(01\)00100-6](https://doi.org/10.1016/S0191-8141(01)00100-6)
- Zachrisson, E., & Sjöstrand, T. (1990). Bedrock Map 22E Frostviken. Sveriges Geologiska Undersökning (SGU) Ai 44, scale 1: 50,000.

**How to cite this article:** Giuntoli F, Menegon L, Warren CJ. Replacement reactions and deformation by dissolution and precipitation processes in amphibolites. *J Metamorph Geol.* 2018;36:1263–1286. <https://doi.org/10.1111/jmg.12445>