

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

ROCK1 is associated with non-syndromic cleft palate

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version: Palmieri A., Scapoli L., Carrozzo M., Cura F., Morselli P.G., Pannuto L., et al. (2020). ROCK1 is associated with non-syndromic cleft palate. JOURNAL OF ORAL PATHOLOGY & MEDICINE, 49, 164-168 [10.1111/jop.12973].

Availability: This version is available at: https://hdl.handle.net/11585/723036 since: 2020-02-07

Published:

DOI: http://doi.org/10.1111/jop.12973

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)



DR LUCA SCAPOLI (Orcid ID : 0000-0003-4006-9910)

Article type : Original Article

ROCK1 is associated with non-syndromic cleft palate

Annalisa Palmieri^{1,\$}, Luca Scapoli^{1,\$}, Marco Carrozzo², Francesca Cura³, Paolo Giovanni Morselli^{1,4}, Lucia Pannuto⁴, Nayereh Nouri^{5,6}, Francesco Carinci³, Dorina Lauritano⁷, Marcella Martinelli¹

¹Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Bologna, 40126 Bologna, Italy

²School of Dental Sciences, University of Newcastle upon Tyne, Newcastle upon Tyne NE2 4BW, United Kingdom

³Department of Morphology, Surgery and Experimental Medicine, University of Ferrara, 44121 Ferrara, Italy

⁴Plastic Surgery Unit, University Hospital of Bologna Sant'Orsola Malpighi Polyclinic, 40138 Bologna, Italy

⁵Craniofacial and Cleft Research Center, Isfahan University of Medical Sciences, 81745-319 Isfahan, Iran

⁶Department of Genetics and Molecular Biology, Medical School, Isfahan University of Medical Sciences, 81745-319 Isfahan

⁷Department of Medicine and Surgery, Centre of Neuroscience of Milan, University of Milano-Bicocca, 20126 Milan, Italy

^{\$} Palmieri and Scapoli should be considered joint first author.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi: 10.1111/JOP.12973</u>

This article is protected by copyright. All rights reserved

Running title: ROCK1 role in non-syndromic cleft palate

Correspondence:

Luca Scapoli, Dept. of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Via Belmeloro 8 - 40126 Bologna, Italy

Tel. +39.051.2094106 - Fax. +39.051.2094110 Email: luca.scapoli2@unibo.it

ABSTRACT

Background Craniofacial morphogenesis is the result of an intricate multistep network of tightly controlled spatial and temporal signalling that involves several molecules and transcription factors organised into highly coordinated pathways. Any alteration in even one step of this delicate process can lead to congenital malformations such as cleft palate. One of the first steps in embryonal orofacial development is the migration of cells from the neural crests to the branchial arches. Next, the cells have to proliferate, differentiate, move and connect to each other in order to correctly form the palate. Cell contraction, promoted by the interaction of non-muscle myosin II and actin A, is a crucial step in morphogenesis and is regulated by ROCK1 protein.

Methods A family-based association study was carried out in order to verify whether or not genetic variants of *ROCK1* were associated with non-syndromic cleft palate (nsCP). Two cohorts from Italy and Iran, a total of 189 nsCP cases, and their parents were enrolled.

Results The rs35996865-G allele was under-transmitted in cases of nsCP [P = 0.006, odds ratio (OR) = 0.63 (95% C.I. 0.45–0.88)].

Conclusion This investigation reveals for the first time data supporting a role for *ROCK1* in nsCP aetiology.

KEYWORDS

eleft palate, orofacial malformation, ROCK1, association analysis

INTRODUCTION

During human organogenesis, neural crest cells (NCCs), a population of cells with migratory capacity, migrate from the dorsal surface of the neural tube to different regions of the embryo, where they differentiate into a broad range of cell types.¹ In vertebrates, the cranial NCC-derived mesenchyme contributes considerably to the development of the face and pharyngeal arches.² Orofacial development is a complex and delicate process, and craniofacial morphogenesis and palate formation occur through a well-orchestrated series of actions - including growth, expansion and fusion of the prominences - which is tightly controlled by the spatial and temporal signalling network.³ Correct formation of the secondary palate is conditioned by the ability of cells to interact with neighbouring cells and adapt their reaction to the changes occurring in the extracellular matrix leading to proliferation, migration, rearrangement, adhesion and fusion of the palatal shelves.^{4,5}

Aberration in any step of morphogenesis could lead to inborn malformation. In particular, the lack of fusion of embryonic palatal shelves between the 6th and 8th weeks results in cleft palate, a condition that affects about 1 to 25 per 10,000 newborns.⁶ Cleft palate is one of the commonest congenital defects. It may occur as a cleft in the palate alone (nsCP), affecting only the secondary palate involving the hard palate and/or soft palate, and sometimes being limited to one cleft uvula.^{7,8} In other cases, cleft palate can arise together with cleft lip, potentially involving the alveolar process and the primary palate (cleft lip with or without cleft palate, CL/P).⁹ Despite their apparent similarity, however, there is epidemiological evidence to support nsCP and CL/P being different malformations with distinct but potentially overlapping aetiology.¹⁰

A variety of signalling pathways are involved in palate development, and several mutations in developmental genes have been found to contribute to nsCP.¹¹⁻¹³ During morphogenetic movement, the cytoskeleton plays a crucial role in correctly allocating cells along the midline where the fusion of palatal shelves takes place.¹⁴ Proliferation, migration and adhesion of cells in the branchial processes involves the myosin 9 gene (*MYH9*), one of the three class II myosins expressed in non-muscle cells to exert contractile force.^{15,16} In a previous investigation, our group provided evidence supporting *MYH9* as a predisposing factor for CL/P.¹⁷ Its involvement in orofacial clefts, including nsCP, has been confirmed in different populations.¹⁸⁻²⁰

Crucial for the contractile ability expressed by cells is the interaction of activated non-muscle myosin II with actin.²¹ Activation of myosin is brought about by Rho-associated coiled-coil-containing protein kinase 1 (ROCK1). Rho kinases can act both directly, through phosphorylation

of regulatory myosin II light chain, and indirectly, by inhibiting the activity of myosin phosphatase.²²

Interestingly, transgenic mice embryos expressing a dominant-negative form of Rho kinase (Rock) developed severe NCC-related defects, including severe craniofacial clefting affecting the nose, palate and lip.²³ These mutant embryos showed disruption of the actin cytoskeleton, associated with NCC cell death in the craniofacial region. A recent investigation demonstrated that cell motility is crucial for normal palate development and that functional integrity of a contractility pathway, that includes both the *Myh9* gene product and its activator ROCK1, is needed.¹⁴

The two isoforms of the main downstream effectors of the Rho family, *ROCK1* and *ROCK2*, share 65% amino acid sequence homology, and both modulate cytoskeleton organization by phosphorylating different substrates, mainly myosin light chain and myosin phosphatase.²⁴ Nevertheless, they have distinct functions. In fact, while ROCK2 activity facilitates phagocytosis, ROCK1 is mainly responsible for stress fiber association and focal adhesion formation.²⁵ Since it is ROCK1, rather than ROCK2 that could play a role in the morphogenetic events that lead to the palate formation, we set out to conduct a family-based association study of *ROCK1* polymorphisms. A sample of 136 unrelated nsCP probands from Italy and their parents, as well as a second cohort of 53 triads from Iran were investigated in order to assess for genetic heterogeneity among populations of different ethnicities. This study represents the first attempt to evaluate a potential role for *ROCK1* in nsCP onset to date.

This article is protected by copyright. All rights reserved

MATERIALS AND METHODS

Study subjects

This case-parent association study was performed on two cohorts: a set of 136 unrelated trios of Italian ancestry, and a set of 53 trios of Fars ethnicity (from the central region of Iran). Triads were defined by a nsCP proband, free of any other congenital anomalies, and his/her parents. Guidelines were shared by research teams to ensure that samples from the two groups were collected uniformly. A clinical team ascertained the non-syndromic status of all probands. Exclusion criteria were exposure to known risk factors like phenytoin and warfarin, and tobacco smoking or alcohol consumption during pregnancy. Positive family history of cleft was defined as having at least one relative (second, third, or fourth degree) affected by nsCP. The study was approved by the ethics committees of Ferrara Province (Comitato Etico Unico della Provincia di Ferrara) (Study n° 130795) and Sant'Orsola-Malpighi General Hospital (project n° 14020) in Italy, and the Ethics and Research Committee of Isfahan University of Medical Sciences (process n° 189087) in Iran. The study complied with the Helsinki Declaration's Ethical Principles for Medical Research Involving Human Subjects. Prior informed written consent was obtained from all participants.

Genotyping

Four tag single nucleotide polymorphisms (SNPs) were selected by the Tagger tool using Haploview v.4.2 software.²⁶ The four tagSNPs (Table 2) captured 100% of the *ROCK1* alleles, with a minor allele frequency (MAF) > 0.05 and a mean max $r^2 = 0.951$.

DNA was extracted from peripheral whole blood using the GenEluteTM Blood Genomic DNA Kit (Sigma), and then quantified with a Nanodrop spectrophotometer (Thermo Fisher Scientific). Genotyping was performed using KASPTM genotyping technology (LGC group), a competitive allele-specific PCR, with the ABI PRISM 7500 thermalcycler (Applied Biosystems).

Statistical analyses

Statistical analysis to check for Hardy-Weinberg equilibrium in parents and family-based association was performed as before.²⁷ Briefly, the transmission disequilibrium test was performed using PLINK software v1.07 on a Windows PC.^{28,29} The association level was quantified by means of the odds ratio (OR), i.e. the M1/M2 ratio, where M1 and M2 were the number of times the allele was transmitted and not transmitted from heterozygous parents, respectively.³⁰ Multiple testing correction was performed by the "set-based test" procedure of PLINK. The method, suited

for candidate gene studies, uses permutation to determine empirical P values of association between the disease and the sets of independent SNPs, which are recognized considering the observed levels of linkage disequilibrium between the SNPs.

This article is protected by copyright. All rights reserved

RESULTS

The study included a total of 189 patient-parents triads of two different ethnicities. A description of the enrolled sample study was reported in Table 1.

Four tagSNPs at the *ROCK1* locus were investigated (Table 2). Genotype frequencies were distributed according to Hardy-Weinberg equilibrium law, and no Mendelian errors were detected. A family-based association study was performed to determine whether *ROCK1* polymorphisms were involved in nsCP. The analysis was performed for each population (Italian and Iranian samples), as well as on the entire data set, by the transmission disequilibrium test (Table 3). In the set pertaining to the entire sample, evidence of the association between rs35996865 alleles and the occurrence of nsCP emerged; indeed, the rs35996865-G allele was under-transmitted in nsCP cases [P = 0.006, odds ratio (OR) = 0.63 (95% C.I. 0.45-0.88)]. The multiple-test correction method adopted in the study indicates that this association is significant at a level of 0.05; indeed, the permutation set-based tests produced a corrected P of 0.013. Considering the two populations individually, the effect size appeared higher among Iranians [OR = 0.52 (95% C.I. 0.27–1.02)] than in Italians [OR = 0.70 (95% C.I. 0.48–1.03)], although the association between rs35996865 and nsCP was not significant in either the Iranian (P value = 0.052) or the Italian sample (P value = 0.069).

The haplotype association analysis was conducted with a combination of all four markers. Table 4 shows *ROCK1* haplotype association data for the sample as a whole. The haplotype rs288980-T|rs17202375-A|rs2127958-T|rs35996865-G was significantly under-transmitted to nsCP probands in both the entire sample (*P* value = 0.01) and the Italian group (*P* value = 0.018).

DISCUSSION

As *ROCK1*, which codes for a kinase that regulates myosin II activity and its interaction with actin filaments, is among the genes coding for proteins responsible for the reorganization of the actin cytoskeleton, cell shape regulation, attachment, and motility, we hypothesised that it may be implicated in the nsCP phenotype. Indeed, our association analysis supported the involvement of *ROCK1* in nsCP aetiology in the entire sample of 189 trios (P = 0.006). The association between nsCP and *ROCK1* was confirmed (P = 0.013) after corrections for multiple testing, and by haplotype association analysis, which detected significant under-transmission of a specific *ROCK1* haplotype in affected children (P = 0.01). The association analysis was repeated on the sample stratified by ethnicity, i.e. separating the 136 trios of Italian ancestry from the 53 trios of Fars ethnicity (Iran). Basing on the ORs obtained, the rs35996865-G allele showed association levels for the two populations similar to those observed for the whole sample. Because of the small sample size, however, the statistical power of the association was limited, yielding non-significant P values, albeit approaching the threshold level. That being said, a significant association with a specific haplotype (rs288980-T|rs17202375-A|rs2127958-T|rs35996865-G) was observed for the Italian population.

This investigation has yielded evidence of a significant association between *ROCK1* alleles and nsCP. Specifically, the rs35996865-G allele appeared to be under-transmitted in both Italian and Iranian populations. The associated SNP rs35996865 maps in the *ROCK1* promoter region, about 500 bp upstream of the transcription start site. However, it is unknown whether this polymorphism can alter the expression level of the investigated gene. Indeed, the SNP could be in linkage disequilibrium with another causative polymorphism, or a mutation that could affect either the promoter or coding sequence of *ROCK1*.

A genome-wide association analysis of a multiethnic sample, searching for common gene variants associated with nsCP risk, discovered a genome-wide significant association with a missense variant in the transcription factor *GRHL3*.³¹ No association with polymorphisms in *ROCK1* locus was found in that investigation. However, genetic heterogeneity may account for different association results. Also of note, the same genome wide investigation failed to replicate a previously reported association with *FAF1*.³² Nevertheless, the association data collected in this investigation support a role for *ROCK1* in nsCP aetiology. Cell motility appears to be determinant in normal palate development. Functional studies in mice indicate that cell extrusion from the medial epithelial seam appears a key step for palatal shelves fusion.¹⁴ Particularly, at molecular

level, this process requires the activation of a contractility pathway that includes Rock1 and its downstream effector NMHCIIA, the *Myh9* gene product.¹⁴ Considering that the product of *ROCK1*, by modulating the interaction between actin and non-muscle myosin, plays a role in the cell dynamics needed for palate development, the results of this investigation suggest the need to further explore the role of *ROCK1* in nsCP in independent sample studies.

Acknowledgments

This work was supported in part by the Association Interethnos-Interplast Italy (to Marcella Martinelli) and the University of Ferrara (FIR2017 to Francesco Carinci).

Conflict of interest

The authors declare no competing interests.

Author contributions

A Palmieri, L Scapoli and M Martinelli designed the study, as well as writing and revising the manuscript. PG Morselli, L Pannuto, N Nouri, F Carinci and D Lauritano collected biological samples and acquired clinical data. A Plamieri, F Cura and N Nouri performed the genetic analysis and screened the scientific literature. L Scapoli performed the statistical analysis. A Palmieri, L Scapoli and M Martinelli interpreted the data. M Carrozzo supervised the manuscript. All authors took part in manuscript preparation and gave final approval of the version to be submitted.

Accepted

REFERENCES

- 1. Etchevers HC, Dupin E, Le Douarin NM. The diverse neural crest: from embryology to human pathology. *Development*. 2019;146(5).
- 2. Frisdal A, Trainor PA. Development and evolution of the pharyngeal apparatus. *Wiley interdisciplinary reviews Developmental biology*. 2014;3(6):403-418.
- 3. Minoux M, Rijli FM. Molecular mechanisms of cranial neural crest cell migration and patterning in craniofacial development. *Development*. 2010;137(16):2605-2621.
- 4. Tarr JT, Lambi AG, Bradley JP, Barbe MF, Popoff SN. Development of Normal and Cleft Palate: A Central Role for Connective Tissue Growth Factor (CTGF)/CCN2. *Journal of developmental biology*. 2018;6(3).
- Lan Y, Xu J, Jiang R. Cellular and Molecular Mechanisms of Palatogenesis. *Current topics in developmental biology*. 2015;115:59-84.
- Burg ML, Chai Y, Yao CA, Magee W, 3rd, Figueiredo JC. Epidemiology, Etiology, and
 Treatment of Isolated Cleft Palate. *Front Physiol.* 2016;7:67.
- Lewin ML, Croft CB, Shprintzen RJ. Velopharyngeal insufficiency due to hypoplasia of the musculus uvulae and occult submucous cleft palate. *Plastic and reconstructive surgery*. 1980;65(5):585-591.
- 8. Gosain AK, Conley SF, Marks S, Larson DL. Submucous cleft palate: diagnostic methods and outcomes of surgical treatment. *Plastic and reconstructive surgery*. 1996;97(7):1497-1509.
- 9. Mossey PA, Little J, Munger RG, Dixon MJ, Shaw WC. Cleft lip and palate. *Lancet*. 2009;374(9703):1773-1785.
- 10. Leslie EJ, Marazita ML. Genetics of cleft lip and cleft palate. *American journal of medical genetics Part C, Seminars in medical genetics*. 2013;163C(4):246-258.
- Nakajima A, C FS, Gulka AOD, Hanai JI. TGF-beta Signaling and the Epithelial-Mesenchymal Transition during Palatal Fusion. *International journal of molecular sciences*. 2018;19(11).
- 12. Beaty TH, Marazita ML, Leslie EJ. Genetic factors influencing risk to orofacial clefts: today's challenges and tomorrow's opportunities. *F1000Res*. 2016;5:2800.
- Li C, Lan Y, Jiang R. Molecular and Cellular Mechanisms of Palate Development. *Journal of dental research*. 2017;96(11):1184-1191.

- 14. Kim S, Lewis AE, Singh V, Ma X, Adelstein R, Bush JO. Convergence and extrusion are required for normal fusion of the mammalian secondary palate. *PLoS biology*. 2015;13(4):e1002122.
- 15. Vicente-Manzanares M, Ma X, Adelstein RS, Horwitz AR. Non-muscle myosin II takes centre stage in cell adhesion and migration. *Nature reviews Molecular cell biology*. 2009;10(11):778-790.
- 16. Shutova MS, Svitkina TM. Common and Specific Functions of Nonmuscle Myosin II Paralogs in Cells. *Biochemistry Biokhimiia*. 2018;83(12):1459-1468.
- 17. Martinelli M, Di Stazio M, Scapoli L, et al. Cleft lip with or without cleft palate: implication of the heavy chain of non-muscle myosin IIA. *J Med Genet*. 2007;44(6):387-392.
- 18. Birnbaum S, Reutter H, Mende M, et al. Further evidence for the involvement of MYH9 in the etiology of non-syndromic cleft lip with or without cleft palate. *European journal of oral sciences*. 2009;117(2):200-203.
- Wang Y, Li D, Xu Y, et al. Functional Effects of SNPs in MYH9 and Risks of Nonsyndromic Orofacial Clefts. *Journal of dental research*. 2018;97(4):388-394.
- 20. Peng HH, Chang NC, Chen KT, et al. Nonsynonymous variants in MYH9 and ABCA4 are the most frequent risk loci associated with nonsyndromic orofacial cleft in Taiwanese population. *BMC medical genetics*. 2016;17(1):59.
- 21. Landsverk ML, Epstein HF. Genetic analysis of myosin II assembly and organization in model organisms. *Cell Mol Life Sci.* 2005;62(19-20):2270-2282.
- 22. Matsumura F, Totsukawa G, Yamakita Y, Yamashiro S. Role of myosin light chain phosphorylation in the regulation of cytokinesis. *Cell Struct Funct*. 2001;26(6):639-644.
- 23. Phillips HM, Papoutsi T, Soenen H, Ybot-Gonzalez P, Henderson DJ, Chaudhry B. Neural crest cell survival is dependent on Rho kinase and is required for development of the mid face in mouse embryos. *PLoS One*. 2012;7(5):e37685.
- 24. Amano M, Nakayama M, Kaibuchi K. Rho-kinase/ROCK: A key regulator of the cytoskeleton and cell polarity. *Cytoskeleton*. 2010;67(9):545-554.
- 25. Yoneda A, Multhaupt HA, Couchman JR. The Rho kinases I and II regulate different aspects of myosin II activity. *J Cell Biol.* 2005;170(3):443-453.
- 26. Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*. 2005;21(2):263-265.

- 27. Cura F, Palmieri A, Girardi A, et al. Possible effect of SNAIL family transcriptional repressor
 1 polymorphisms in non-syndromic cleft lip with or without cleft palate. *Clin Oral Investig.* 2018;22(7):2535-2541.
- 28. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics*. 2007;81(3):559-575.
- 29. Spielman RS, McGinnis RE, Ewens WJ. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *American journal of human genetics*. 1993;52(3):506-516.
- 30. Kazeem GR, Farrall M. Integrating case-control and TDT studies. *Ann Hum Genet*. 2005;69(Pt 3):329-335.
- 31. Leslie EJ, Liu H, Carlson JC, et al. A Genome-wide Association Study of Nonsyndromic Cleft Palate Identifies an Etiologic Missense Variant in GRHL3. *American journal of human genetics*. 2016;98(4):744-754.
- 32. Ghassibe-Sabbagh M, Desmyter L, Langenberg T, et al. FAF1, a gene that is disrupted in cleft palate and has conserved function in zebrafish. *American journal of human genetics*. 2011;88(2):150-161.

1		ITALIAN COHORT	IRANIAN COHORT		
	Sex				
	Male	57	25		
	Female	79	28		
	Parental consanguinity				
	YES	0	21		
	NO	136	32		
	History of familial cleft				
	YES	8	8		
	NO	128	45		
	Type of cleft				
	Hard, soft and uvula	73	27		
	Soft and uvula	61	23		
	Uvula	2	0		
	Submucous	0	3		

TABLE 1 Characteristics of the two nsCP cohorts

SNIP ID	CHPa	BDp	A 110100C	MAF ^d	MAF ^d
SINI ID	CIIK	DI	Alleles	ITALY	IRAN
rs288980	18	18609580	C/T	0.42	0.50
rs17202375	18	18621273	A/G	0.14	0.07
rs2127958	18	18653610	T/C	0.48	0.41
rs35996865	18	18692344	T/G	0.28	0.22

TABLE 2 Gene polymorphisms and observed allele frequency

^achromosome

^bchromosome position GRCh37/hg19

^cmajor allele first

^dminor allele frequency

This article is protected by copyright. All rights reserved

Q	ENTIRE SAMPLE	SNP ID	Alleles ^a	Tb	Uc	<i>P</i> value	OR (95% CI)
		rs288980	C/T	66	73	0.553	0.90 (0.65, 1.26)
		rs17202375	A/G	39	36	0.729	1.08 (0.69, 1.70)
		rs2127958	C/T	77	67	0.405	1.15 (0.83, 1.60)
		rs35996865	T/G	55	88	0.006	0.63 (0.45, 0.88)
	ITALIAN COHORT	SNP ID	Alleles ^a	T ^b	U°	<i>P</i> value	OR (95% CI)
		rs288980	C/T	45	58	0.200	0.78 (0.53, 1.15)
		rs17202375	A/G	35	30	0.535	1.17 (0.72, 1.9)
		rs2127958	C/T	57	49	0.437	1.16 (0.79, 1.70)
		rs35996865	T/G	45	64	0.069	0.70 (0.48, 1.03)
	IRANIAN COHORT	SNP ID	Alleles ^a	Tb	Uc	<i>P</i> value	OR (95% CI)
		rs288980	C/T	22	22	1.000	1.00 (0.55, 1.81)
	2	rs17202375	A/G	5	7	0.564	0.71 (0.23, 2.25)
		rs2127958	C/T	25	22	0.662	1.14 (0.64, 2.02)
		rs35996865	T/G	13	25	0.052	0.52 (0.27, 1.02)

TABLE 3 Family-based association analysis of ROCK1 polymorphisms in nsCP

^amajor allele first

^bnumber of times the minor allele was transmitted from heterozygous parents ^cnumber of times the minor allele was untransmitted from heterozygous parents Italic font is used for *P* values < 0.05

		1 51	5			
Q	ENTIRE SAMPLE	HAPLOTYPE ^a	T ^b	Uc	CHISQ	P value
		C A C G	15.0	16.4	0.06	0.806
		T A T G	34.2	59.3	6.71	0.010
		C G C T	29.8	23.4	0.78	0.374
		C A C T	63.3	54.2	0.70	0.403
	5	T A T T	64.7	47.6	2.61	0.106
		C A T T	28.6	35.6	0.76	0.382
	ITALIAN COHORT	HAPLOTYPE ^a	T ^b	Uc	CHISQ	P value
		C A C G	13.7	10.0	0.59	0.444
		T A T G	25.4	45.4	5.61	0.018
	4	C G C T	26.0	18.8	1.14	0.287
		C A C T	46.0	41.1	0.27	0.604
		T A T T	45.2	36.6	0.91	0.340
	5	C A T T	22.2	26.6	0.40	0.528
	IRANIAN COHORT	HAPLOTYPE ^a	T ^b	Uc	CHISQ	P value
		C A C G	2.9	6.9	1.64	0.200
		T A T G	9.1	14.1	1.08	0.300
		C G C T	3.5	5.5	0.44	0.505
		C A C T	21.6	13.6	1.80	0.180
		T A T T	20.8	15.9	0.67	0.414
		C A T T	9.1	10.0	0.05	0.828

TABLE 4 Haplotype association analysis at ROCK1

^acombination of alleles of the SNPs rs288980|rs17202375|rs2127958|rs35996865 ^bnumber of times the minor allele was transmitted from heterozygous parents ^cnumber of times the minor allele was untransmitted from heterozygous parents Italic font is used for P values < 0.05Ì