

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

An innovative approach for the assessment of mood disturbances in patients with eating disorders

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Tecuta, L., Fava, G.A., Tomba, E. (2020). An innovative approach for the assessment of mood disturbances in patients with eating disorders. *CNS SPECTRUMS*, 25(1), 71-78 [10.1017/S1092852919000798].

Availability:

This version is available at: <https://hdl.handle.net/11585/739435> since: 2020-02-28

Published:

DOI: <http://doi.org/10.1017/S1092852919000798>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

**An Innovative Approach for the Assessment of Mood Disturbances in Patients with Eating
Disorders**

Lucia Tecuta¹, Giovanni Andrea Fava^{1,2}, Elena Tomba¹

¹ Department of Psychology, University of Bologna, Bologna, Italy

² Department of Psychiatry, State University of New York at Buffalo, Buffalo, NY

Corresponding Author:

Giovanni A. Fava, MD

Department of Psychology,

University of Bologna,

Viale Berti Pichat, 5

40127 Bologna, Italy,

phone: +39-051-2091339 fax: +39-051-243086

E-mail: giovanniandrea.fava@unibo.it

Acknowledgements

The authors would like to thank the therapists and the collaborators of the Eating Disorder Clinics “Centro Gruber” (Bologna, Italy) and "Residenza Gruber" (Bologna, Italy), in particular Dr. Donatella Ballardini, MD, Dr. Romana Schumann, Psy.D. and Dr. Anna Franco, PsyD for assistance in participant recruitment and data gathering.

Abstract

Objective: Assessment of mood in Eating disorders (EDs) has important clinical implications, but the current standard psychiatric classification (DSM-5) has limitations. The aim of the current study is to broaden the evaluation of depressive symptomatology by providing a comprehensive and innovative assessment approach in EDs through instruments that capture clinical phenomena of demoralization, subclinical distress, and psychological well-being.

Methods: Seventy-nine patients who met diagnostic criteria for EDs of the Diagnostic and Statistical Manual of Mental Disorders- Fifth edition (DSM-5) were evaluated for depressive symptoms through Paykel's Clinical Interview for Depression, the Structured Clinical Interview for DSM 5 for major depressive episode and persistent depressive disorder, and the Diagnostic Criteria for Psychosomatic Research (DCPR) interview for demoralization. Further, self-report inventories encompassing psychological well-being and distress were used.

Results: Guilt, abnormal reactivity to social environment, and depressed mood were the most common depressive symptoms in the sample. DSM-defined depressive disorders were found in 55.7% of patients. The DCPR-demoralization criteria identified an additional 20.3% of the sample that would have been undetected with DSM criteria. Both DSM and DCPR diagnostic categories were associated with compromised psychological well-being and distress. Demoralization, unlike depression, was not associated with the severity of ED symptomatology.

Conclusion: The findings indicate that a standard psychiatric approach, DSM-5-based, captures only a narrow part of the spectrum of mood disturbances affecting patients with EDs. A broadened clinimetric assessment unravels the presence of demoralization and yields clinical distinctions that may entail prognostic and therapeutic differences among patients who would be otherwise simply labeled as depressed.

Key words: demoralization, depression, eating disorders, psychological well-being

INTRODUCTION

Depression is a frequent complication of eating disorders (EDs) and eating disturbances are common manifestations of depressive illness. The nature of the relationship, however, has been a source of controversy. A shared etiology postulates a common set of risk factors leading to the development of both EDs and depression¹. Mood disorder onset might precede, follow or develop simultaneously to the ED², suggesting the need of specifically evaluating the individual case.

Diagnosis of depression in EDs constitutes a difficult task. Not surprisingly, comorbidity rates of major depressive disorder (MDD) display wide fluctuations, from 40 to 80%^{2,3} and lack predictive value as to response to antidepressant therapy⁴. Overlapping symptomatology, such as excessive weight loss, over-eating, sleep disturbance, fatigue, irritability, concentrating difficulties, and poor memory^{5,6} may account for inflated depression diagnoses in this clinical population^{7,8}.

It has been suggested that exclusive reliance on conventional diagnostic classification systems may not provide sufficient clinical information and assessment may benefit from additional sources of information⁹. One source derives from expanding collection of symptoms to clinical manifestations that frequently occur in the longitudinal development of mood disorder¹⁰. The Clinical Interview for Depression^{11,12} is uniquely suited for capturing such manifestations. Another important element of the clinical process comes from the concept of demoralization^{13,14}, a feeling state characterized by the perception of being unable to cope with some pressing problems and/or of lack of adequate support from others¹⁵. Demoralization, seldom investigated in eating disorders¹⁶ may co-occur with major depression or be independent¹⁵ and is associated with adverse health outcomes and poor quality of life. Finally, a neglected area in assessment is psychological well-being, despite the availability of validated instruments and its growing importance in establishing resilience. Dimensions of positive functioning were found to affect the complex balance between positive and negative affects both in mood¹⁷ and eating^{18,19} disorders.

The aim of the current study was to broaden the evaluation of depressive symptomatology in EDs with instruments that capture clinical phenomena such as demoralization and subclinical distress, and to examine their associations with dimensional measures of psychological well-being. More specifically, the study aimed to provide a comprehensive and innovative assessment of mood in EDs, capturing not only traditional psychiatric disturbances such as depression but also subclinical manifestations and psychological states such as demoralization. We hypothesize that a subset of ED patients exhibit demoralization syndrome in absence of major depression and vice versa. In addition, we hypothesize that demoralization is associated with significant distress and impaired positive functioning.

Methods

Participants

Consecutively recruited patients (n=81) who met Diagnostic and Statistical Manual of Mental Disorders 5 (DSM-5) criteria for EDs²⁰, anorexia nervosa (AN), bulimia nervosa (BN), binge-eating disorder (BED), and other specified feeding or eating disorder (OSFED), were recruited from a specialized ED treatment centers, Centro Gruber and Residenza Gruber (Bologna, Italy) before commencing treatment. ED diagnoses were established at intake by the consensus of a psychiatrist and a clinical psychologist independently using the Structured Clinical Interview for DSM 5 (SCID)²⁰. With the exception of two patients who refused to participate, all invited patients took part in the study (n=79).

Ethical review committees of the Centro Gruber and Residenza Gruber in Bologna, Italy approved the study and all patients provided written informed consent after the procedures were explained to them.

Measures and Clinical Variables

The evaluation was performed during routine assessment visits. Participants underwent detailed clinical interviews by a trained clinical psychologist and completed several self-rating

questionnaires for the assessment of distress and psychological well-being. Data were collected between April 2016 and October 2017.

1) Depressive disorder diagnoses were obtained using the **Structured Clinical Interview for DSM 5 (SCID)**²⁰ for depressive disorders. For a diagnosis of major depression, patients had to exhibit 5 out of 8 symptoms one of which was depressed mood or loss of interest or pleasure. The criterion of significant weight gain or loss or change in appetite was excluded as eating disorder patients exhibit changes in weight and appetite in accordance with ED disorder diagnosis. For a diagnosis of persistent depressive disorder, in addition to depressed mood for most of the day, for more days than not for two years, the patients had to exhibit at least two additional symptoms, with the exception of poor appetite or overeating.

2) Depressive symptoms were assessed with the change version of the **Clinical Interview for Depression- 20 item** interview^{11,12} a dimensional observer-rated assessment instrument which consists of an expanded version of the Hamilton Rating Scale for Depression²¹. The interview covers 20 symptom areas. In this modified version, two items concerning appetite and weight gain/loss (items 12 and 13) were omitted due to the potentially confounding aspects of eating disorder-related symptomatology. Each item is rated on a 1–7 point scale, with 1 indicating absence of symptoms and 7 severe incapacitating manifestations. A score of 3 or above in the individual items was considered the cut-off for presence of the symptom. The scale encompasses a wide range of symptoms (such as irritability and phobic anxiety) compared to other scales and is particularly suitable to assess subclinical symptoms of mood disorders^{10,12,22}. One item concerning reactivity to social environment, selected from the full version of the CID, was added to the 18 items.

3) Demoralization diagnosis was obtained using **The revised Structured Interview for the Diagnostic Criteria for Psychosomatic Research (DCPR)-Demoralization Criteria**¹⁴. Diagnoses were formulated independently of DSM diagnostic findings. Items of the interview for DCPR are scored through a yes/no response format. The structured interview has demonstrated high inter-rater

reliability and Cohen's kappa for demoralization was found to be 0.90²³. The revised DCPR criteria¹⁴ allow differentiation of two expressions of demoralization: helplessness (the individual maintains the capacity to react, but lacks adequate support) and hopelessness (when the individual feels he/she alone is responsible for the situation and there is nothing he /she or anyone else can do to overcome the problem).

4) Self-report depressive symptoms were assessed with the **Beck Depression Inventory II** (BDI-II)²⁴ a 21 item questionnaire. A total score ranging 0-63 indicates depression severity with higher total scores indicate more severe depressive symptoms. Composite scales of cognitive and somatic-affective symptoms were calculated²⁵.

5) Positive functioning was evaluated with the **Psychological Well-being Scales-PWB**^{26,27}, an 84 item self-rated questionnaire that covers 6 inter-related areas of psychological well-being which allow the development of optimal functioning: autonomy, environmental mastery, personal growth, positive relations with others, purpose in life and self-acceptance. Items are constructed on a six-point 1–6 Likert scale, yielding six subscale scores ranging from 14 to 84. Subscale scores range from 0-98 with higher scores indicate greater psychological well-being in specific dimensions.

6) Eating disorder symptomatology was assessed with the **Eating Attitudes Test-40**^{28,29}, a 40 item screening measure identifying behaviors and cognitive patterns of eating disorders. Items are constructed on a 0–3 four-point Likert scale, yielding three subscale scores for dimensions of dieting, body, and food preoccupation, oral control, and a total score ranging from 0 to 120. Higher scores indicate greater eating disorder psychopathology.

In addition to administering these clinical scales, **Body mass index (BMI)**, **illness duration in months** and **type of antidepressant therapy** were collected from the medical records.

Data Analysis

Descriptive analyses were run for frequency of CID-rated specific depression symptoms and frequencies of demoralization and depressive illness (persistent depressive disorder and MDD) in the total sample. Univariate analyses of variance using the general linear model were performed to test for associations between DSM 5 Depressive Disorders and the DCPR-based classification of demoralization and average scores on dimensional psychological measures after controlling for illness duration. DSM 5 and DCPR-based diagnoses were examined separately. For all tests performed, the significance level was set at .05, two-tailed. In view of the exploratory nature of the investigation, adjustment for multiple testing was not performed. Age, educational level and BMI were not significantly correlated with any outcome variable and were therefore excluded from analyses³⁰.

Results

ED patients characteristics

The patient response rate was high with 97.53% (n=79) of ED outpatients out of 81 agreeing to participate (see Table 1 for descriptive socio-demographic and clinical data). Data on specific depression symptoms through CID interview was available for 72 patients. The 79 eating disorder patients were all female with mean age 28.83 ± 11.25 years, range 15-58 years and mean educational years 14.44 ± 3.15 . About half, 53.2% (n=42) were outpatients and the remaining 46.8% were inpatients (n=37). Outpatients and inpatients did not differ significantly in main socio-demographic characteristics, that is age, education or in BMI. They differed significantly in illness duration ($p=.04$) with inpatients reporting longer length of illness (11.69 ± 9.02 years) compared to outpatients (7.49 ± 8.96 years). Almost a third (n=22, 27.8%) of patients were currently on antidepressants, the most common being selective serotonin-reuptake inhibitors (n=18). Inpatient and outpatient groups did not differ significantly in severity (BDI and CID total scores) of depression. Diagnostic subgroups (AN, BN, BED, OSFED) also did not differ in severity of depression. See *Table 1*.

Frequency of depressive symptoms in ED patients

The most common CID depressive symptoms in the ED sample (n=72) were feelings of guilt, environmental reactivity, depressed mood, and low energy or fatigue, which were present in about two-thirds of the sample. Please see Table 2 for frequencies of these and of other depressive symptoms .

Relationship of DSM depression diagnoses and DCPR demoralization

Diagnoses are displayed in Figure 1. Nineteen patients (24.0%) were without any mood-related comorbidity (i.e. unaffected). A fifth of patients reported only demoralization (20.3%), 12 with helpless demoralization and four cases of hopeless demoralization. In terms of DSM-defined diagnoses, comorbid persistent depressive disorder was reported by 16.5% of the patients, while major depressive disorder was the most prevalent (39.2%). Demoralization overlapped partially with both persistent depressive disorder and MDD. None of the unaffected ED patients and about half (n=16) of the comorbid MDD group were on antidepressant medications. Three demoralized and four ED patients with persistent depressive disorder were also on antidepressants.

Associations between DSM 5 TR Depressive Disorders and dimensional psychological variables

According to univariate analyses of variance comparisons, comorbid depressive disorder (MDD or persistent depressive disorder) was associated with significantly greater distress in terms of depressive symptoms in BDI-II as well as in EAT-eating-related pathology, including oral control, food and bulimic worry, and dietary restraint. In terms of psychological well-being, presence of a depressive illness was associated with significantly worse functioning in PWB dimensions of environmental mastery, positive relations with others, purpose in life, and self-acceptance. Table 3 displays the comparisons between ED patients with and without comorbid DSM depressive illness.

Associations between DCPR demoralization and dimensional psychological variables

Univariate analyses of variance yielded some significant associations between diagnoses formulated according to the DCPR-based demoralization and dimensional psychological variables. Demoralization diagnosis in ED patients was also associated with significantly greater distress in terms of BDI-II scores but not in terms of EAT-eating-related pathology, in which no significant differences between demoralized and non-demoralized ED patients emerged. Nonetheless, in PWB dimensions of psychological well-being, occurrence of demoralization was associated with significantly worse functioning in environmental mastery, positive relations with others, purpose in life, and self-acceptance. Table 3 illustrates the comparisons between demoralized and non-demoralized ED patients.

Discussion

Despite some limitations of the current study, namely the small sample size and cross-sectional design, the joint use of a comprehensive clinical interview for depression^{11,12} with the DSM diagnostic criteria²⁰ and demoralization and psychological well-being assessment has yielded important clinical insights into mood disturbances in EDs.

First, diagnostic criteria place particular emphasis on a specific set of symptoms within a certain symptomatology. Such priority, however, does not necessarily apply to a setting of comorbidity, where other symptoms may be prominent and characteristic. The innovativeness of the applied assessment approach resides in the use of the clinimetric approach. Clinimetrics refers to clinically relevant information frequently ignored by traditional psychiatric approaches such as patterns of symptoms, severity of illness, effects of comorbid conditions, timing of phenomena, rate of progression of illness, functional capacity and other aspects such as positive functioning. The use of a macro-analytic evaluation using clinician-rated scales such as the CID and DCPR, followed by the use of a micro-analysis of specific symptoms through self-rating scales allows to capture subclinical symptoms, which would have been otherwise undetected⁹. Using the CID, the most common depression symptoms in EDs were depressed mood, feelings of guilt and abnormal reactivity to social environment. Guilt has been

found to be exceedingly common in ED patients, not globally but in particular in relation to eating and eating behaviors^{31,32} and in relation to body shame³³. Moreover, it has been found to persist throughout the recovery process from the disorder³¹. Reactivity to social environment refers to the changes in mood and symptomatology, as a result of environmental circumstances, either improvement or worsening. It has been found to characterize cyclothymia³⁴ and to be prevalent in the prodromal phase of bulimia nervosa compared to unaffected controls³⁵. Recent studies on emotional reactivity in EDs³⁶ suggest that such reactivity in AN and BN might be related to social situations in which patients are pressured to consume high-calorie food, evoking states of anxiety and depression³⁷.

Anxiety, whether generalized, somatic or phobic, was also found to be very common. Studies have shown that anxiety symptomatology is often comorbid and may precede EDs³⁸. More specifically eating disorders have been found to be associated with body anxiety, eating and food preoccupations, pre-meal anxiety³⁹ and avoidance behaviors relating to food, body and interpersonal situations^{40,41}. Indeed recent research explores the efficacy of exposure therapy and response prevention in the treatment of anorexia nervosa⁴².

Figure 1 illustrates the complexity of mood assessment in eating disorders. Using DCPR criteria, 20% of patients met the criteria for demoralization. Such patients would not be identified using DSM criteria only. The percentage of demoralization cases is in line with varying prevalence rates in the medical and psychiatric setting¹⁵. In EDs, a previous study which had not taken into account a possible overlap with major depression and dysthymic disorder, had reported higher rates¹⁶. Interestingly, cases of demoralization in absence of depressive disorder were mostly of the helpless subtype.

Associations of DCPR-demoralization with dimensional psychological measures provide further support to its validity and utility. Consistent with previous studies^{15,43}, demoralization was found to be associated with lower psychological well-being and greater distress. Moreover, the

same number of significant associations were found between demoralization and depressive illness and psychological variables, with the exception of ED-related symptoms, which were not associated with demoralization. Findings lends support to the hypothesis that DCPR-demoralization might be suitable for classifying psychological distress in EDs that is not confounded by the ED symptoms themselves. Indeed demoralization does not seem to depend on illness type or severity, affecting a wide range of psychiatric and medical illnesses alike¹⁵.

Subjective incompetence (a feeling of being trapped or blocked because of a sense of inability to plan or start actions toward goals) is a major component of demoralization^{44,45}. Such feelings of inadequacy and low self-efficacy have been previously documented in EDs^{46,47}. Individuals who perceive themselves as incompetent are uncertain and indecisive as to their directions, display high reactivity to environmental stimuli and low psychological well-being. Not surprisingly, patients with eating disorders were found to present with very high rates of drop-out⁴⁸.

In both standard assessment and treatment approaches, most studies focus on pathological symptomatology and its reduction, as well as modifications of physical and behavioral aspects, ignoring gains in positive aspects such as quality of life and psychological well-being^{18,19,49}. The pursuit of euthymia, defined as how the individual adjusts the psychological dimensions of well-being to changing needs may thus become one of the targets of treatment. Such positive functioning characteristics have been found to be persistently compromised in various psychiatric illnesses including eating disorders¹⁸ and their impairments are correlated with increased vulnerability to future adversity and may thus be a viable psychotherapeutic target⁵⁰⁻⁵³.

The high comorbidity rates found in the current study between eating disorders and depression are in line with those in the literature, ranging from 40 to 80%, as well as high comorbidity rates with dysthymic disorder which surpass those found in the general population^{2,3,4}. In the current study, depressive disorders were associated with worse overall functioning both in terms of psychological distress and well-being, a result that is in line with the literature. Depressed ED patients exhibit greater dietary restriction, body dissatisfaction and worse quality of life⁵⁴,

social⁵⁵ and global functioning compared to unaffected ED patients^{56,57,58}. Moreover, it is well-documented that eating disorder severity is significantly associated with severity of the depression⁵⁹.

As expected, an overlap between the demoralization and depressive illness was found, as in other medical psychiatric populations¹⁵.

Clinical guidelines for the screening of depression in EDs recommend the Beck Depression Inventory-II and the Hamilton Depression Rating Scale^{60,61}. However, such standardized screening measures mainly based on DSM criteria might not be sufficient in such a complex clinical population with frequent medical complications.

Conclusion

The presence of demoralization syndrome in eating disorders is undoubtedly relevant to treatment and recovery from EDs. Hopelessness and helplessness⁶² and poor self-efficacy^{63,64} have been identified by ED patients themselves as barriers to change and recovery in qualitative studies. In chronic AN patients, the recovery process may be hindered by feelings of hopelessness and "feeling stuck"⁶⁵. Indeed, demoralization has been found to affect response to psychotherapy^{66,67}.

Moreover, the clinical utility of depression diagnoses in ED populations is called into question by treatment trials. Depression comorbidity in EDs has an unclear role in treatment response with inconsistent results across ED categories^{68,69}. For instance, inconclusive or mixed findings have been reported for FDA approved fluoxetine in BN^{70,71,72} with lack of improvement in AN⁷³. Antidepressants in ED patients were devoid of impact on likelihood and persistence of recovery from MDD in a longitudinal study⁴.

Fava, Rafanelli and Tomba⁹ have advocated that exclusive reliance on diagnostic criteria has impoverished the clinical process in psychiatry. Customary clinical taxonomy in psychiatry does not include clinical distinctions that demarcate major prognostic and therapeutic differences among patients who otherwise seem to be deceptively similar since they share the same psychiatric

diagnosis⁹. This investigation has illustrated how a broader perspective in evaluation of mood disturbances in EDs may lead to an individualized assessment of the complex balance between euthymia, dysthymia and eating behavior which may entail important treatment implications. Further studies, using a comprehensive clinimetric approach⁷⁴ with larger samples and a longitudinal design, are needed.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Disclosure

Lucia Tecuta, Giovanni Andrea Fava, and Elena Tomba have nothing to disclose.

References

1. Puccio F, Fuller-Tyszkiewicz M, Ong D, Krug I. A systematic review and meta-analysis on the longitudinal relationship between eating pathology and depression. *Int J Eat Disord.* 2016;49(5):439-54
2. Godart N, Radon L, Curt F, Duclos J, Perderau F, Lang F, Venisse JL, Halfon O, Bizouard P, Loas G, Corcos M, Jeammet, Flament M. Mood disorders in eating disorder patients: prevalence and chronology of onset. *J Affect Disord.* 2015;185:115-122
3. Godart NT, Perdereau F, Rein Z, Berthoz S, Wallier J, Jeammet P, Flament M. Comorbidity studies of eating disorders and mood disorders. Critical review of the literature. *J Affect Disord.* 2007;97(1-3):37–49
4. Mischoulon D, Eddy KT, Keshaviah A, Dinescu D, Ross SL, Kass AE, Franko DL, Herzog DB. Depression and Eating Disorders: Treatment and Course. *J Affect Disord.* 2011;130(3):470-477
5. Meehan KG, Loeb KL, Roberto CA, Attia E. Mood change during weight restoration in patients with anorexia nervosa. *Int J Eat Disord.* 2006;39:587–589
6. Mattar L, Huas C, Duclos J, Apfel A, Godart N. Relationship between malnutrition and depression or anxiety in Anorexia Nervosa: A critical review of the literature. *J Affect Disord.* 2010;132(3):311-8
7. Garfinkel PE, Garner DM. *Anorexia Nervosa: A Multidimensional Perspective*. New York, NY: Brunner/Mazel Inc; 1982
8. Casper R. Depression in eating disorders. *Depress Anxiety* 1998;8:96-104
9. Fava GA, Rafanelli C, Tomba E. The clinical process in psychiatry: a clinimetric approach. *J Clin Psychiatry* 2012; 73(2):177-184
10. Fava GA. Subclinical symptoms in mood disorders. *Psychol Med.* 1999;29(1):47–61
11. Paykel ES. The Clinical Interview for Depression: development, reliability and validity. *J Affect Disord.* 1985;9(1):85–96
12. Guidi J, Fava GA, Bech P, Paykel E. The Clinical Interview for Depression: a comprehensive

- review of studies and clinimetric properties. *Psychother Psychosom.* 2011;80(1):10–27
13. Fava GA, Freyberger HJ, Bech P, Christodoulou G, Sensky T, Wise TN. Diagnostic criteria for use in psychosomatic research. *Psychother Psychosom.* 1995;63(1):1-8
 14. Fava GA, Cosci F, Sonino N. Current psychosomatic practice. *Psychother Psychosom.* 2017;86(1):13-30
 15. Tecuta L, Tomba E, Grandi S, Fava GA. Demoralization and its clinical characterization. *Psychol Med.* 2015;45(4):673-91
 16. Abbate-Daga G, Delsedime N, Nicotra, B, Giovannone, C, Marola E, Amianto F, Fassino S. Psychosomatic syndromes and anorexia nervosa. *BMC Psychiatry.* 2013;13:14
 17. Fava GA, Cosci F, Guidi J, Tomba E. Well-being therapy in depression: new insights into the role of psychological well-being in the clinical process. *Depress Anxiety.* 2017;34(9):801-808
 18. Tomba E, Offidani E, Tecuta L, Schmann R, Ballardini D. Psychological Well-being in out-patients with eating disorders: a controlled study. *Int J Eat Disord.* 2014;47(3):252-258
 19. Tomba E, Tecuta L, Schumann R, Ballardini D. Does psychological well-being change following treatment? An exploratory study on outpatients with eating disorders. *Compr Psychiatry.* 2017;74:61-69
 20. First MB, Williams JBW, Karg RS, Spitzer RL. *Structured Clinical Interview for DSM-5 Disorders, Clinician Version (SCID-5-CV)*. Arlington, VA: American Psychiatric Association; 2015
 21. Hamilton M. Development of a rating scale for primary depressive illness. *Brit J Soc Clin Psychol.* 1967(4);6:278–296
 22. Bech P. Fifty years with the Hamilton Scales for Anxiety and Depression. *Psychother Psychosom.* 2009;78(4):202–211
 23. Galeazzi GM, Ferrari S, Mackinnon A, Rigatelli M. Inter-rater reliability, prevalence and relation to ICD-10 diagnoses of the Diagnostic Criteria for Psychosomatic Research in consultation-liaison psychiatry patients. *Psychosom.* 2004;45(5):386-393

24. Beck AT, Steer, RA, Brown GK. *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation; 1996
25. Wang YP, Gorenstein C. Psychometric properties of the Beck Depression Inventory-II: a comprehensive review. *Rev Bras Psiquiatr*. 2013;35(4):416-31
26. Ryff CD. Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *J Pers Soc Psychol*. 1989;57(6):1069-1081
27. Ryff CD. Psychological well-being revisited: advances in the science and practice of eudaimonia. *Psychother Psychosom*. 2014;83(1):10-28
28. Garner DM, Garfinkel PE. The Eating Attitude Test: An index of the symptoms of anorexia nervosa. *Psychol Med*. 1979;9(2):273-279
29. Garfinkel PE, Newman A. The eating attitudes test: twenty-five years later. *Eat Weight Disord*. 2001;6(1):1-24
30. Miller GA, Chapman JP. Misunderstanding analysis of covariance. *J Abnorm Psychol*. 2001;110(1):40-48
31. Frank ES. Shame and guilt in eating disorders. *Am J Orthopsychiatry*. 1991;61(2):303-306
32. Petry N, Vasconcelos FAG, Costa LDCE. Feelings and perceptions of women recovering from anorexia nervosa regarding their eating behavior. *Cad Saude Publica*. 2017;33(9):e00048716
33. Burney J, Irwin HJ. Shame and guilt in women with eating-disorder symptomatology. *J Clin Psychol*. 2000;56(1):51-61
34. Tomba E, Rafanelli C, Grandi S, Guidi J, Fava GA. Clinical configuration of cyclothymic disturbances. *J Affect Disord*. 2012;139(3):244-249
35. Raffi AR, Rondini M, Grandi S, Fava GA. Life events and prodromal symptoms in bulimia nervosa. *Psychol Med*. 2000;30(3):727-731
36. Tapajóz P, de Sampaio F, Soneira S, Aulicino A, Harris P, Allegri RF. Emotional reactivity to social stimuli in patients with eating disorders. *Psychiatry Res*. 2015;229(3):887-94
37. Gutiérrez-Maldonado J, Ferrer-García M, Caqueo-Úrizar A, Letosa-Porta A. Assessment of


- emotional reactivity produced by exposure to virtual environments in patients with eating disorders. *Cyberpsychol Beh.* 2006;9(5):507-513
38. Swinbourne JM, Touyz SW. The co-morbidity of eating disorders and anxiety disorders: a review. *Eur Eat Disord Rev.* 2007;15(4):253-274
39. Steinglass J, Albano AM, Simpson HB, Carpenter K, Schebendach J, Attia E. Fear of food as a treatment target: exposure and response prevention for anorexia nervosa in an open series. *Int J Eat Disord.* 2012;45(4):615-621
40. Arkell J, Robinson P. A pilot case series using qualitative and quantitative methods: biological, psychological and social outcome in severe and enduring eating disorder (anorexia nervosa). *Int J Eat Disord.* 2008;41(7):650-6
41. Bamford BH, Attoe C, Mountford VA, Morgan JF, Sly R. Body checking and avoidance in low weight and weight restored individuals with anorexia nervosa and non-clinical females. *Eat Behav.* 2014;15(1):5-8
42. Steinglass JE, Albano AM, Simpson HB, Wang Y, Zou J, Attia E, Walsh BT. Confronting fear using exposure and response prevention for anorexia nervosa: A randomized controlled pilot study. *Int J Eat Disord.* 2014;47(2):174-80
43. Grandi S, Sirri L, Tossani E, Fava GA. Psychological characterization of demoralization in the setting of heart transplantation. *J Clin Psychiatry.* 2011;72(5):648-654
44. de Figueiredo JM. Depression and demoralization: phenomenologic differences and research perspectives. *Compr Psychiatry.* 1993;34(5):308-311
45. de Figueiredo JM. Distress, demoralization and psychopathology: diagnostic boundaries. *Eur J Psychiatry.* 2013;27(1):61-73
46. Garner DM. *EDI2—Eating Disorders Inventory 2. Professional manual.* Odessa, FL: Psychological Assessment Resources; 1991
47. Surgenor LJ, Maguire S, Russell J, Touyz S. Self-liking and self-competence: relationship to symptoms of anorexia nervosa. *Eur Eat Disord Rev.* 2007;15(2):139-145

48. Fassino S, Pierò A, Tomba E, Abbate-Daga G. Factors associated with dropout from treatment for eating disorders: a comprehensive literature review. *BMC Psychiatry*. 2009;9:67
49. Ackard DM, Richter S, Egan A, Engel S, Cronemeyer CL. The meaning of (quality of) life in patients with eating disorders: a comparison of generic and disease-specific measures across diagnosis and outcome. *Int J Eat Disord*. 2014;47(3):259–267
50. Jahoda M. *Current concepts of positive mental health*. New York, NY: Basic Books; 1958.
51. Fava GA, Bech P. The concept of euthymia. *Psychother Psychosom*. 2016;85(1):1-5
52. Tomba E, Tecuta L. Well-being therapy in a patient with anorexia nervosa. *Psychother Psychosom*. 2016;85:369-70
53. Fava GA. *Well-being therapy. Treatment Manual and Clinical Applications*. Basel: Karger; 2016.
54. Padierna A, Quintana J, Arostegui, I, Gonzalez N, Horcajo M. The health-related quality of life in eating disorders. *Qual Life Res*. 2000;9(6):667-674
- Hatsukami DK, Mitchell JE, Eckert ED, Pyle RL. Characteristics of patients with bulimia only, bulimia with affective disorder, and bulimia with substance abuse problems. *Addict Behav*. 1986;11(4):399-406
56. Bulik CM, Sullivan PF, Carter FA, Joyce PR. Lifetime anxiety disorders in women with bulimia nervosa. *Compr Psychiatry*. 1996;37(5):368-374
57. Hughes EK, Goldschmidt AB, Labuschagne Z, Loeb KL, Sawyer SM, Le Grange D. Eating disorders with and without comorbid depression and anxiety: Similarities and differences in a clinical sample of children and adolescents. *Eur Eat Disord Rev*. 2013;21(5):386-394
58. Brand-Gothelf A, Leor S, Apter A, Fennig S. The impact of comorbid depressive and anxiety disorders on severity of anorexia nervosa in adolescent girls. *J Nerv Ment Dis*. 2014;202(10):759-762
59. Herpertz-Dahlmann B, Remschmidt H. Depression in anorexia nervosa at follow-up. *Int J Eat Disord*. 1993;14(2):163-169

60. Anderson DA, Lundgren JD, Shapiro JR, Paulosky CA. Assessment of eating disorders: review and recommendations for clinical use. *Behav Modif.* 2004; 28(6):63-82
61. Bowers WA, Haedt-Matt AA. Psychological Assessment of Eating Disorders. In: Preedy VR, Watson RR, Colin RM, eds. *Handbook of Behavior, Food and Nutrition*. New York, NY: Springer; 2011: 3425-3435
62. Waller G. The myths of motivation. Time for a fresh look at some received wisdom in the eating disorders? *Int J Eat Disord.* 2012;45(1):1-16
63. Wade TD, Wilksch SM, Paxton SJ, Byrne SM, Austin SB. How perfectionism and ineffectiveness influence growth of eating disorder risk in young adolescent girls. *Beh Res Ther.* 2015;66:56-63
64. Dawson L, Rhodes P, Touyz S. “Doing the Impossible”: the process of recovery from chronic anorexia nervosa. *Qual Health Res.* 2014;24(4):494-505
65. Nordbø RHS, Espeset EMS, Gulliksen KS, Skårderud F, Geller J, Holte A. Reluctance to recover in anorexia nervosa. *Eur Eat Disord Rev.* 2012;20(1):60-67
66. Frank JD. *Persuasion and healing*. Baltimore: Johns Hopkins University Press; 1961
67. Tomba E, Tecuta L, Guidi J, Grandi S, Rafanelli C. Demoralization and response to psychotherapy: a pilot study comparing the sequential combination of cognitive-behavioral therapy and well-being therapy with clinical management in cyclothymic disorder. *Psychother Psychosom.* 2016;85(1):56-57
68. Agras WS, Walsh BT, Fairburn CG, Wilson GT, Kraemer HC. A multicenter comparison of cognitive-behavioral therapy and interpersonal psychotherapy for bulimia nervosa. *Arch Gen Psychiatry.* 2000;57(5):459–466
69. Collin P, Power K, Karatzias T, Grierson D, Yellowlees A. The effectiveness of and predictors of response to inpatient treatment of anorexia nervosa. *Eur Eat Disord Rev.* 2010;18(6):464–474
70. Fluoxetine Bulimia Nervosa Collaborative Study Group. Fluoxetine in the Treatment of Bulimia Nervosa. *Arch Gen Psychiatry.* 1992;49(2):139–147

71. Walsh BT, Wilson GT, Loeb KL, Devlin MJ, Pike KM, Roose SP, Fleiss J, Waternaux C. Medication and psychotherapy in the treatment of bulimia nervosa. *Am J Psychiatry*. 1997;154(4): 523–531
72. Romano SJ, Halmi KA, Sarkar NP, Koke SC, Lee JS. A placebo-controlled study of fluoxetine in continued treatment of bulimia nervosa after successful acute fluoxetine treatment. *Am J Psychiatry*. 2002;159(1):96–102
73. Walsh BT, Kaplan AS, Attia E, Olmsted M, Parides M, Carter JC, Pike KM, Devlin MJ, Woodside B, Roberto CA, Rockert W. Fluoxetine after weight restoration in anorexia nervosa: A randomized controlled trial. *JAMA*. 2006;295(22): 2605–2612
74. Fava GA, Carrozzino D, Lindberg L, Tomba E. The clinimetric approach to psychological assessment. *Psychother Psychosom*. 2018; 87(6):321-326

Figure 1. Prevalence of demoralization subtypes, persistent depressive disorder and major depression in ED patients



Notes. DCPR, Diagnostic Criteria for Psychosomatic Research; ED, eating disorder, MDD, major depressive disorder

Table 1. Socio-demographic characteristics of ED patient sample (N= 79)

Variable	Total ED sample (n=79)	Outpatients (n=42)	Inpatients (n=37)	p
Age	28.83±11.25	28.64±12.38	29.06± 9.943	.873 ⁺
Education (years)	14.44±3.15	14.40± 3.03	14.48± 3.355	.912 ⁺
Marital status (% single)	74.7	84.6	74.3	.087 [*]
Occupation (% employed or student)	75.3	92.3	51.35	.001[*]
BMI (kg/m ²)				
AN (n)	(42): 15.26±1.76	(17): 15.63±1.69	(25): 15.00±1.79	.260 [*]
BN (n)	(13):22.60±5.29	(7): 24.05±6.91	(6): 20.90±1.87	.350 [*]
BED (n)	(13): 35.53±9.95	(10): 32.73±9.77	(3):43.92±4.69	.092 [*]
OSFED (n)	(11): 21.26±8.61	(8): 18.30±3.65	(3): 33.11±14.93	.392 [*]
Illness duration (years)	9.45±9.17	7.49±8.96	11.69±9.021	.04⁺
Antidepressant use within group (%)	27.8	22.2	38.9	.125 [*]

Note. AN, anorexia nervosa; BED, binge-eating disorders; BN, bulimia nervosa; OSFED, Other-specified feeding or eating disorder

^{*}Pearson Chi-squared

⁺T-test for independent samples

Table 2. Frequency of depression symptoms (CID items) in ED patients (n=72)

CID ITEM	Symptom frequency (n)	Symptom frequency (%)
Guilt	57	79.2
Environmental reactivity	56	77.8
Depressed mood	55	76.4
Energy and fatigue	51	70.8
Generalized anxiety	46	63.9
Work and interests	42	58.3
Somatic anxiety	41	56.9
Pessimism	38	52.8
Delayed insomnia	33	45.8
Phobic anxiety	29	40.3
Suicidal tendencies	25	34.7
Phobic avoidance	23	31.9
Irritability	23	31.9
Early insomnia	22	30.6
Depressed appearance	22	30.6
Panic attacks	14	19.4
Psychomotor retardation	14	19.4
Agitation	7	9.7
Hostility	4	5.6

Note: CID, Clinical Interview for Depression

Table 3. Associations of DSM-5 Depressive Disorders and DCPR-based demoralization with dimensional psychological measures (N=79)

VARIABLE	Depressive disorder (+) N=44	Depressive disorder (-) N=35	F (1,72)	P	Partial eta squared	Demoralization (+) N=51	Demoralization (-) N=28	F (1,72)	P	Partial eta squared
	Estimated marginal mean (S.E.)	Estimated marginal mean (S.E.)				Estimated marginal mean (S.E.)	Estimated marginal mean (S.E.)			
BDI-II Cognitive subscale	17.667 (1.092)	9.764 (1.124)	25.122	<.0001	.273	11.641 (.808)	5.234 (1.044)	22.062	<.0001	.253
BDI-II Somatic-affective subscale	12.297 (.856)	6.086 (.856)	26.145	<.0001	.287	17.272 (1.018)	8.345 (1.309)	26.883	<.0001	.286
EAT oral control	12.493 (1.326)	7.071 (1.520)	7.211	.009	.092	10.533 (1.333)	9.439 (1.848)	.217	.643	.003
EAT food - bulimic worry	12.232 (.938)	6.383 (1.075)	16.777	<.0001	.191	10.648 (.984)	7.957 (1.364)	2.409	.125	.033
EAT dietary restraint	27.497 (2.281)	13.347 (2.614)	16.597	<.0001	.189	23.717 (2.388)	17.061 (3.312)	2.499	.118	.034
PWB autonomy	47.035 (2.356)	54.367 (2.499)	4.557	.037	.066	46.595(2.095)	58.097 (3.005)	9.230	.003	.124
PWB environmental mastery	55.965 (1.772)	64.767 (1.772)	12.318	.001	.161	42.342 (1.612)	56.549 (2.312)	23.779	<.0001	.268
PWB personal growth	57.388 (1.888)	64.063 (2.002)	5.880	.018	.083	58.228 (1.750)	65.033 (2.510)	4.631	.035	.067
PWB positive relations	50.666 (2.184)	63.564 (2.317)	16.398	<.0001	.201	51.021(1.924)	67.915 (2.758)	23.625	<.0001	.267
PWB purpose in life	47.537 (1.779)	55.614 (1.887)	9.694	.003	.130	47.425 (1.562)	58.994 (2.240)	16.799	<.0001	.205
PWB self-acceptance	46.150 (2.356)	35.227 (2.221)	11.375	.001	.149	34.705 (1.890)	51.447 (2.711)	24.025	<.0001	.270

Note: (+) Disorder or syndrome present, (-) Disorder or syndrome absent; BDI-II, Beck Depression Inventory II; EAT, Eating Attitudes Test; PWB, Psychological Well-Being scales