

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Low Degree Morphisms of $E(5, 10)$ -Generalized Verma Modules

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Low Degree Morphisms of $E(5, 10)$ -Generalized Verma Modules / Cantarini N.; Caselli F.. - In: ALGEBRAS AND REPRESENTATION THEORY. - ISSN 1386-923X. - ELETTRONICO. - 23:6(2020), pp. 2131-2165. [10.1007/s10468-019-09925-0]

Availability:

This version is available at: <https://hdl.handle.net/11585/746275> since: 2020-03-03

Published:

DOI: <http://doi.org/10.1007/s10468-019-09925-0>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Cantarini, N., Caselli, F. Low Degree Morphisms of $E(5, 10)$ -Generalized Verma Modules. *Algebr Represent Theor* 23, 2131–2165 (2020)

The final published version is available online at
<https://dx.doi.org/10.1007/s10468-019-09925-0>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

LOW DEGREE MORPHISMS OF $E(5, 10)$ -GENERALIZED VERMA MODULES

NICOLETTA CANTARINI AND FABRIZIO CASELLI*

ABSTRACT. In this paper we face the study of the representations of the exceptional Lie superalgebra $E(5, 10)$. We recall the construction of generalized Verma modules and give a combinatorial description of the restriction to \mathfrak{sl}_5 of the Verma module induced by the trivial representation. We use this description to classify morphisms between Verma modules of degree one, two and three proving in these cases a conjecture given by Rudakov [8]. A key tool is the notion of dual morphism between Verma modules.

1. INTRODUCTION

Infinite dimensional linearly compact simple Lie superalgebras over the complex numbers were classified by Victor Kac in 1998 [3]. A complete list, up to isomorphisms, consists of ten infinite series and five exceptions, denoted by $E(1, 6)$, $E(3, 6)$, $E(3, 8)$, $E(5, 10)$ and $E(4, 4)$. See also [1, 9, 10, 11] for the genesis of these superalgebras. Some years later Kac and Rudakov initiated the study of the representations of these algebras [4, 5, 7, 6] developing a general theory of Verma modules that we briefly recall.

Let $L = \bigoplus_{j \in \mathbb{Z}} L_j$ be a \mathbb{Z} -graded Lie superalgebra, let $L_- = \bigoplus_{j < 0} L_j$, $L_+ = \bigoplus_{j > 0} L_j$ and $L_{\geq 0} = L_0 \oplus L_+$. We denote by $U(L)$ the universal enveloping algebra of L . If F is an irreducible L_0 -module we define

$$M(F) = U(L) \otimes_{U(L_{\geq 0})} F$$

where we extend the action of L_0 to $L_{\geq 0}$ by letting L_+ act trivially on F . We call $M(F)$ a minimal generalized Verma module associated to F . If $M(F)$ is not irreducible we say that it is degenerate.

In [4, 5, 7, 6], a complete description of the degenerate Verma modules for $E(3, 6)$ and $E(3, 8)$ is given, as well as of their unique irreducible quotients. In [6] some basic ideas and constructions are settled also for $E(5, 10)$. In this case Kac and Rudakov conjecture a complete list of L_0 -modules which give rise to the degenerate Verma modules (see Conjecture 4.6).

In 2010 Rudakov tackled the proof of the conjecture through the study of morphisms between Verma modules. The existence of a degenerate Verma module is indeed strictly related to the existence of such morphisms of positive degree (see Proposition 3.5). In [8] Rudakov classified morphisms of degree one and gave some examples of morphisms of degree at most 5. He also conjectured that there exists no morphism of higher degree and that

2010 *Mathematics Subject Classification.* 17B15, 17B25 (primary), 17B65, 17B70 (secondary).

Key words and phrases. Lie superalgebras, Verma modules, Singular vectors.

*Corresponding author

his list exhausts all the examples. A more general family of modules, possibly induced from infinite-dimensional \mathfrak{sl}_5 -modules, had been studied in [2], where some of Rudakov's examples in degree one and two had been obtained through the use of the computer.

In this paper we study morphisms between generalized Verma modules and to this aim we analyze the structure of the universal enveloping algebra $U_- = U(L_-)$ as an L_0 -module. This analysis has its own interest and provides an explicit combinatorial description of the action of L_0 . This description is the main ingredient in our study of morphisms, together with a systematic use of the dominance order of the weights of the L_0 -modules. Our main result is the proof of Rudakov's conjecture in degree two and three (see Theorems 9.8, 10.15). A useful observation that we made is that if there exists a morphism $\varphi : M(V) \rightarrow M(W)$ between generalized Verma modules of degree d , then there exists a dual morphism $\psi : M(W^*) \rightarrow M(V^*)$ of the same degree. This duality is here proved in low degree for the purpose of this work but it holds in a much wider context as a consequence of the fact that the conformal dual of a Verma module is itself a Verma module. This will be shown in a forthcoming paper.

The paper is organized as follows: in Section 2 we recall the basic definitions and fix the notation. Section 3 is dedicated to Verma modules. Here we characterize degenerate Verma modules in terms of singular vectors and morphisms. In Section 4, following [8], we give examples of morphisms of degree one, two and three. Section 5 contains our first main result on the structure of U_- as an L_0 -module: we construct an explicit basis of U_- and describe its combinatorial properties. Section 6 is dedicated to the analysis of the dominance order of the weights of the basis elements of U_- . In Section 7 we develop the idea of dual morphism between generalized Verma modules and establish sufficient conditions for the existence of such a morphism (see Remark 7.2). Finally, Sections 8, 9 and 10 contain the classification of morphisms of degree one, two and three, respectively.

We thank Victor Kac for useful discussions.

2. PRELIMINARIES

We let $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ be the set of non-negative integers and for $n \in \mathbb{N}$ we set $[n] = \{i \in \mathbb{N} \mid 1 \leq i \leq n\}$.

If P is a proposition we let $\chi_P = 1$ if P is true and $\chi_P = 0$ if P is false.

We consider the simple, linearly compact Lie superalgebra of exceptional type $L = E(5, 10)$ whose even and odd parts are as follows: $L_{\bar{0}}$ consists of zero-divergence vector fields in five (even) indeterminates x_1, \dots, x_5 , i.e.,

$$L_{\bar{0}} = S_5 = \left\{ X = \sum_{i=1}^5 f_i \partial_i \mid f_i \in \mathbb{C}[[x_1, \dots, x_5]], \operatorname{div}(X) = 0 \right\},$$

where $\partial_i = \partial_{x_i}$, and $L_{\bar{1}} = \Omega_{cl}^2$ consists of closed two-forms in the five indeterminates x_1, \dots, x_5 . The bracket between a vector field and a form is given by the Lie derivative and for $f, g \in \mathbb{C}[[x_1, \dots, x_5]]$ we have

$$[f dx_i \wedge dx_j, g dx_k \wedge dx_l] = \varepsilon_{ijkl} f g \partial_{t_{ijkl}}$$

where, for $i, j, k, l \in [5]$, ε_{ijkl} and t_{ijkl} are defined as follows: if $|\{i, j, k, l\}| = 4$ we let $t_{ijkl} \in [5]$ be such that $|\{i, j, k, l, t_{ijkl}\}| = 5$ and ε_{ijkl} be the sign of the permutation (i, j, k, l, t_{ijkl}) . If $|\{i, j, k, l\}| < 4$ we let $t_{ijkl} = 1$ (this choice will be irrelevant) and $\varepsilon_{ijkl} = 0$.

From now on we shall denote $dx_i \wedge dx_j$ simply by d_{ij} .

The Lie superalgebra L has a consistent, irreducible, transitive \mathbb{Z} -grading of depth 2 where, for $k \in \mathbb{N}$,

$$\begin{aligned} L_{2k-2} &= \langle f\partial_i \mid i = 1, \dots, 5, f \in \mathbb{C}[[x_1, \dots, x_5]]_k \rangle \cap S_5 \\ L_{2k-1} &= \langle fd_{ij} \mid i, j = 1, \dots, 5, f \in \mathbb{C}[[x_1, \dots, x_5]]_k \rangle \cap \Omega_{cl}^2 \end{aligned}$$

where by $\mathbb{C}[[x_1, \dots, x_5]]_k$ we denote the homogeneous component of $\mathbb{C}[[x_1, \dots, x_5]]$ of degree k .

Note that $L_0 \cong \mathfrak{sl}_5$, $L_{-2} \cong (\mathbb{C}^5)^*$, $L_{-1} \cong \wedge^2 \mathbb{C}^5$ as L_0 -modules (where \mathbb{C}^5 denotes the standard \mathfrak{sl}_5 -module). We set $L_- = L_{-2} \oplus L_{-1}$, $L_+ = \oplus_{j>0} L_j$ and $L_{\geq 0} = L_0 \oplus L_+$. We denote by U (resp. U_-) the universal enveloping algebra of L (resp. L_-). Note that U_- is an L_0 -module with respect to the adjoint action: for $x \in L_0$ and $u \in U_-$,

$$x.u = [x, u] = xu - ux.$$

We also point out that the \mathbb{Z} -grading of L induces a \mathbb{Z} -grading on the enveloping algebra U_- . It is customary, though, to invert the sign of the degrees hence getting a grading over \mathbb{N} . Note that the homogeneous component $(U_-)_d$ of degree d of U_- under this grading is an L_0 -submodule. Section 3 will be dedicated to the study of these homogeneous components.

We fix the Borel subalgebra $\langle x_i \partial_j, h_{ij} = x_i \partial_i - x_j \partial_j \mid i < j \rangle$ of L_0 and we consider the usual base of the corresponding root system given by $\{\alpha_{12}, \dots, \alpha_{45}\}$. We let Λ be the weight lattice of \mathfrak{sl}_5 and we express all weights of \mathfrak{sl}_5 using their coordinates with respect to the fundamental weights $\omega_{12}, \omega_{23}, \omega_{34}, \omega_{45}$, i.e., for $\lambda \in \Lambda$ we write $\lambda = (\lambda_{12}, \dots, \lambda_{45})$ for some $\lambda_{i+1} \in \mathbb{Z}$ to mean $\lambda = \lambda_{12}\omega_{12} + \dots + \lambda_{45}\omega_{45}$.

For $i < j$ we denote as usual

$$\alpha_{ij} = \sum_{k=i}^{j-1} \alpha_{k,k+1}$$

and $\alpha_{ji} = -\alpha_{ij}$. For notational convenience we also let $\alpha_{ii} = 0$. Viewed as elements in the weight lattice we have

$$\alpha_{12} = (2, -1, 0, 0), \alpha_{23} = (-1, 2, -1, 0), \alpha_{34} = (0, -1, 2, -1), \alpha_{45} = (0, 0, -1, 2).$$

If $\lambda \in \Lambda$ is a weight, we use the following convention: for all $1 \leq i < j \leq 5$ we let

$$\lambda_{ij} = \sum_{k=i}^{j-1} \lambda_{k,k+1}.$$

If V is a \mathfrak{sl}_5 -module and $v \in V$ is a weight vector we denote by $\lambda(v)$ the weight of v and by $\lambda_{ij}(v) = (\lambda(v))_{ij}$.

If $\lambda = (a, b, c, d) \in \Lambda$ is a dominant weight, i.e. $a, b, c, d \geq 0$, let us denote by $F(\lambda) = F(a, b, c, d)$ the irreducible \mathfrak{sl}_5 -module of highest weight λ . In this paper we always think of $F(a, b, c, d)$ as the irreducible submodule of

$$\text{Sym}^a(\mathbb{C}^5) \otimes \text{Sym}^b(\wedge^2(\mathbb{C}^5)) \otimes \text{Sym}^c(\wedge^2(\mathbb{C}^5)^*) \otimes \text{Sym}^d((\mathbb{C}^5)^*)$$

generated by the highest weight vector $x_1^a x_{12}^b x_{45}^{*c} x_5^{*d}$ where $\{x_1, \dots, x_5\}$ denotes the standard basis of \mathbb{C}^5 , $x_{ij} = x_i \wedge x_j$, and x_i^* and x_{ij}^* are the corresponding dual basis elements. Besides, for a weight $\lambda = (a, b, c, d)$ we let $\lambda^* = (d, c, b, a)$, so that $F(\lambda)^* \cong F(\lambda^*)$.

Notice that $L_1 \cong F(1, 1, 0, 0)$ and that $x_5 d_{45}$ is a lowest weight vector in L_1 . Moreover, for $j \geq 1$, we have $L_j = L_1^j$.

3. GENERALIZED VERMA MODULES AND MORPHISMS

We recall the definition of generalized Verma modules introduced in [4]. For the reader's convenience we also sketch some proofs of basic results. Given an L_0 -module V we extend it to an $L_{\geq 0}$ -module by letting L_+ act trivially, and define

$$M(V) = U \otimes_{U(L_{\geq 0})} V.$$

Note that $M(V)$ has a L -module structure by multiplication on the left, and is called the (generalized) Verma module associated to V . We also observe that $M(V) \cong U_- \otimes_{\mathbb{C}} V$ as \mathbb{C} -vector spaces.

If V is finite-dimensional and irreducible, then $M(V)$ is called a minimal Verma module. We denote by $M(\lambda)$ the minimal Verma module $M(F(\lambda))$. A minimal Verma module is said to be non-degenerate if it is irreducible and degenerate if it is not irreducible.

Definition 3.1. We say that an element $w \in M(V)$ is homogeneous of degree d if $w \in (U_-)_d \otimes V$.

Definition 3.2. A vector $w \in M(V)$ is called a singular vector if it satisfies the following conditions:

- (i) $x_i \partial_{i+1} w = 0$ for every $i = 1, \dots, 4$;
- (ii) $zw = 0$ for every $z \in L_1$;
- (iii) w does not lie in V .

We observe that the homogeneous components of positive degree of a singular vector are singular vectors. The same holds for its weight components. From now on we will thus assume that a singular vector is a homogeneous weight vector unless otherwise specified. Notice that if condition (i) is satisfied then condition (ii) holds if $x_5 d_{45} w = 0$ since $x_5 d_{45}$ is a lowest weight vector in L_1 .

Proposition 3.3. *A minimal Verma module $M(V)$ is degenerate if and only if it contains a singular vector.*

Proof. Let $w \in M(V)$ be a singular vector. We may assume that w is homogeneous of degree $d > 0$. Hence the singular vector w generates a submodule of $M(V)$ which is proper since it is contained in $\oplus_{k \geq d} (U_-)_k \otimes V$.

On the other hand, if $M(V)$ is degenerate let us consider a proper non-zero submodule W of $M(V)$. Let $z \in W$ be a non-zero vector. By repeatedly applying L_1 to z if necessary we can find a non-zero element $w \in W$ such that $L_1 w = 0$, since the action of L_1 lowers the degree of the homogeneous components of z by 1. We observe that L_1 vanishes on the L_0 -module generated by w . Any highest weight vector in such a module is a singular vector. \square

Degenerate Verma modules can also be described in terms of morphisms. A linear map $\varphi : M(V) \rightarrow M(W)$ can always be associated to an element $\Phi \in U_- \otimes \text{Hom}(V, W)$ as follows: for $u \in U_-$ and $v \in V$ we let

$$\varphi(u \otimes v) = u\Phi(v)$$

where, if $\Phi = \sum_i u_i \otimes \theta_i$ with $u_i \in U_-$, $\theta_i \in \text{Hom}(V, W)$, we let $\Phi(v) = \sum_i u_i \otimes \theta_i(v)$. We will say that φ (or Φ) is a morphism of degree d if $u_i \in (U_-)_d$ for every i .

The following proposition characterizes morphisms between Verma modules.

Proposition 3.4. [8] *Let $\varphi : M(V) \rightarrow M(W)$ be the linear map associated with the element $\Phi \in U_- \otimes \text{Hom}(V, W)$. Then φ is a morphism of L -modules if and only if the following conditions hold:*

- (a) $L_0 \cdot \Phi = 0$;
- (b) $t\Phi(v) = 0$ for every $t \in L_1$ and for every $v \in V$.

We observe that if $M(V)$ is a minimal Verma module and condition (a) holds it is enough to verify condition (b) for an element t generating L_1 as an L_0 -module and for v a highest weight vector in V .

Proposition 3.5. *Let $M(\mu)$ be a minimal Verma module. Then the following are equivalent:*

- (a) $M(\mu)$ is degenerate;
- (b) $M(\mu)$ contains a singular vector;
- (c) there exists a minimal Verma module $M(\lambda)$ and a morphism $\varphi : M(\lambda) \rightarrow M(\mu)$ of positive degree.

Proof. We already know that condition (a) is equivalent to condition (b) by Proposition 3.3. Assume condition (c) holds: if $s \in F(\lambda)$ is a highest weight vector, then $\varphi(1 \otimes s)$ is a singular vector in $M(\mu)$.

On the other hand, if w is a singular vector in $M(\mu)$, we can define $\varphi : M(\lambda(w)) \rightarrow M(\mu)$ as the unique morphism of L -modules such that $\varphi(1 \otimes s) = w$, s being a highest weight vector in $M(\lambda(w))$. \square

Remark 3.6. Let $\varphi : M(V) \rightarrow M(W)$ be a linear map of degree d associated to an element $\Phi \in U_- \otimes \text{Hom}(V, W)$ that satisfies condition (a) of Proposition 3.4. Then there exists an L_0 -morphism $\psi : (U_-)_d^* \rightarrow \text{Hom}(V, W)$ such that $\Phi = \sum_i u_i \otimes \psi(u_i^*)$ where $\{u_i, i \in I\}$ is any basis of $(U_-)_d$ and $\{u_i^*, i \in I\}$ is the corresponding dual basis.

Definition 3.7. Let $M(\mu)$ be a minimal Verma module and let $\pi : M(\mu) \rightarrow U_- \otimes F(\mu)_\mu$ be the natural projection, $F(\mu)_\mu$ being the weight space of $F(\mu)$ of weight μ . Given a singular vector $w \in M(\mu)$ we call $\pi(w)$ the leading term of w .

Proposition 3.8. *If w is a singular vector in $M(\mu)$ then:*

- (i) $\pi(w) \neq 0$;
- (ii) if two singular vectors in $M(\mu)$ have the same leading term then they coincide.

Proof. If w is a weight vector homogeneous of degree d then we can write $w = \sum_i u_i \otimes v_i$ for some basis $\{u_i\}$ of $(U_-)_d$ consisting of weight vectors and $v_i \in F(\mu)_{\lambda_i}$ for some weight λ_i . Let λ_{i_0} be maximal in the dominance order such that $v_{i_0} \neq 0$. Then v_{i_0} is a highest weight vector in $F(\mu)$. Indeed, for $r < s$ we have:

$$0 = x_r \partial_s w = \sum_i [x_r \partial_s, u_i] \otimes v_i + \sum_i u_i \otimes x_r \partial_s v_i.$$

By the maximality of λ_{i_0} it follows that $x_r \partial_s v_{i_0} = 0$. (ii) follows from (i). \square

4. EXAMPLES

In this section we give some examples of singular vectors and the corresponding morphisms of Verma modules. These were described in [8]. We will need the following technical result.

Lemma 4.1. *Let $\varphi : M(\lambda) \rightarrow M(W)$ be a morphism of Verma modules of degree one associated to $\Phi = \sum_{i < j} d_{ij} \otimes \theta_{ij}$ and let s be a highest weight vector in $F(\lambda)$. Let \tilde{W} be an L_0 -module containing W and let $\tilde{\theta}_{ij} \in \text{Hom}(F(\lambda), \tilde{W})$ be such that the map $(U_-)_1^* \rightarrow \text{Hom}(F(\lambda), \tilde{W})$ given by $d_{ij}^* \mapsto \tilde{\theta}_{ij}$ is well defined and L_0 -equivariant. Then $\tilde{\theta}_{ij}(s) = \theta_{ij}(s)$ implies $\tilde{\theta}_{ij}(v) = \theta_{ij}(v)$ for all $v \in F(\lambda)$.*

Proof. It is enough to show that if $\tilde{\theta}_{ij}(v) = \theta_{ij}(v)$ for some $v \in F(\lambda)$ and all $i \neq j$, then $\theta_{ij}(x_h \partial_k \cdot v) = \tilde{\theta}_{ij}(x_h \partial_k \cdot v)$ for all $i \neq j$ and $h \neq k$. We have:

$$\begin{aligned} \tilde{\theta}_{ij}(x_h \partial_k \cdot v) &= x_h \partial_k (\tilde{\theta}_{ij}(v)) - (x_h \partial_k \cdot \tilde{\theta}_{ij})(v) = x_h \partial_k (\tilde{\theta}_{ij}(v)) + \delta_{hi} \tilde{\theta}_{kj}(v) + \delta_{hj} \tilde{\theta}_{ik}(v) \\ &= x_h \partial_k (\theta_{ij}(v)) + \delta_{hi} \theta_{kj}(v) + \delta_{hj} \theta_{ik}(v) = \theta_{ij}(x_h \partial_k \cdot v) \end{aligned}$$

where we used Remark 3.6 in order to write the action of L_0 on the θ_{ij} 's. Namely, we have:

$$x_h \partial_k \cdot \theta_{ij} = -\delta_{hi} \theta_{kj} - \delta_{hj} \theta_{ik}$$

where if $r > s$, $\theta_{rs} = -\theta_{sr}$. □

Example 4.2. Let us consider the Verma module $M(m, n, 0, 0)$. We first observe that $d_{12} \otimes x_1^m x_{12}^n$ is a singular vector in $M(m, n, 0, 0)$. Indeed, for $i = 1, \dots, 4$,

$$x_i \partial_{i+1} d_{12} \otimes x_1^m x_{12}^n = 0;$$

besides,

$$x_5 d_{45} d_{12} \otimes x_1^m x_{12}^n = x_5 \partial_3 \otimes x_1^m x_{12}^n = 0.$$

By Proposition 3.5 we can define a morphism of Verma modules $\nabla_A : M(m, n+1, 0, 0) \rightarrow M(m, n, 0, 0)$ by setting $\nabla_A(1 \otimes s) = d_{12} \otimes x_1^m x_{12}^n$. By Lemma 4.1 used with $\tilde{W} = \text{Sym}^m(\mathbb{C}^5) \otimes \text{Sym}^n(\wedge^2 \mathbb{C}^5)$ we have that ∇_A is associated to:

$$\sum_{i < j} d_{ij} \otimes \frac{\partial}{\partial x_{ij}} \in U_- \otimes \text{Hom}(F(m, n+1, 0, 0), F(m, n, 0, 0)).$$

Example 4.3. Let us consider the Verma module $M(m, 0, 0, n+1)$. One can check that $\sum_{j=2}^5 d_{1j} \otimes x_1^m x_j^*(x_5^*)^n$ is a singular vector in $M(m, 0, 0, n+1)$, with leading term $d_{15} \otimes x_1^m (x_5^*)^{n+1}$. By Remark 3.5 we can define a morphism of Verma modules $\nabla_B : M(m+1, 0, 0, n) \rightarrow M(m, 0, 0, n+1)$ by setting $\nabla_B(1 \otimes s) = \sum_{j=2}^5 d_{1j} \otimes x_1^m x_j^*(x_5^*)^n$. By Lemma 4.1, we have that ∇_B is associated to

$$\sum_{i < j} d_{ij} \otimes (x_i^* \partial_j - x_j^* \partial_i).$$

Example 4.4. We shall now exhibit a singular vector in $M(0, 0, m+1, n)$. To this aim it is convenient to think of $F(0, 0, m+1, n)$ as the dual L_0 -module $F(n, m+1, 0, 0)^*$. We shall later investigate the role of duality between Verma modules in Section 7, where we will show, in particular, that the morphism we are going to construct can be seen in a certain sense as the dual of the morphism ∇_A defined in Example 4.2.

Let us observe that the vector $\sum_{i<j} d_{ij} \otimes x_{ij}^*(x_{45}^*)^m(x_5^*)^n$ is a singular vector in $M(F(n, m+1, 0, 0)^*)$ (with leading term $d_{45} \otimes (x_{45}^*)^{m+1}(x_5^*)^n$). Indeed, one immediately checks that $x_k \partial_{k+1}(\sum_{i<j} d_{ij} \otimes x_{ij}^*(x_{45}^*)^m(x_5^*)^n) = 0$ for every $k = 1, \dots, 4$. Besides, we have:

$$\begin{aligned} x_5 d_{45} \left(\sum_{i<j} d_{ij} \otimes x_{ij}^*(x_{45}^*)^m(x_5^*)^n \right) \\ = x_5 \partial_3 x_{12}^*(x_{45}^*)^m(x_5^*)^n - x_5 \partial_2 x_{13}^*(x_{45}^*)^m(x_5^*)^n + x_5 \partial_1 x_{23}^*(x_{45}^*)^m(x_5^*)^n \\ = m(x_{45}^*)^{m-1}(x_5^*)^n(x_{12}^*x_{34}^* + x_{13}^*x_{42}^* + x_{14}^*x_{23}^*) - n(x_{45}^*)^m(x_5^*)^{n-1}(x_{12}^*x_3^* + x_{23}^*x_1^* + x_{31}^*x_2^*) = 0. \end{aligned}$$

Notice that, in fact,

$$x_{ab}^*x_{cd}^* + x_{ac}^*x_{db}^* + x_{ad}^*x_{bc}^* = 0$$

and

$$x_{ab}^*x_c^* + x_{bc}^*x_a^* + x_{ca}^*x_b^* = 0$$

in $F(n, m+1, 0, 0)^*$ for all $a, b, c, d \in [5]$, as one can check by applying these elements to the highest weight vector $x_1^n x_{12}^{m+1}$ in $F(n, m+1, 0, 0)$ and using the L_0 -action.

By Remark 3.5 we can thus define a morphism of Verma modules $\nabla_C : M(0, 0, m, n) \rightarrow M(F(n, m+1, 0, 0)^*)$ by setting $\nabla_C(1 \otimes s) = \sum_{i<j} d_{ij} \otimes x_{ij}^*(x_{45}^*)^m(x_5^*)^n$. Once again, Lemma 4.1 implies that the morphism ∇_C is associated to

$$\sum_{i<j} d_{ij} \otimes x_{ij}^*.$$

Examples 4.2, 4.3 and 4.4 imply the following result.

Proposition 4.5. *Let $m, n \geq 0$. Then $M(m, n, 0, 0)$, $M(m, 0, 0, n)$ and $M(0, 0, m, n)$ are degenerate Verma modules.*

Kac and Rudakov proposed the following conjecture [6]:

Conjecture 4.6. *Let $a, b, c, d \geq 0$ be such that $M(a, b, c, d)$ is a degenerate Verma module. Then $a = b = 0$ or $b = c = 0$ or $c = d = 0$.*

By Proposition 3.5 a possible strategy to prove Conjecture 4.6 is to construct all possible morphisms between minimal Verma modules. One of the main results of this paper is a complete classification of such morphisms of degree at most 3.

Example 4.7. The following are nonzero morphisms of degree 2:

- $\nabla_B \nabla_A : M(m, 1, 0, 0) \rightarrow M(m-1, 0, 0, 1)$;
- $\nabla_C \nabla_B : M(1, 0, 0, n) \rightarrow M(0, 0, 1, n+1)$;
- $\nabla_C \nabla_A : M(0, 1, 0, 0) \rightarrow M(0, 0, 1, 0)$;

Indeed,

$$\nabla_B \nabla_A(1 \otimes x_1^m x_{12}) = \nabla_B(d_{12} \otimes x_1^m) = -m \sum_{j>1} d_{12} d_{1j} \otimes x_1^{m-1} x_j^* \neq 0$$

$$\nabla_C \nabla_B(1 \otimes x_1(x_5^*)^n) = \sum_{j>1} \sum_{h<k} d_{1j} d_{hk} \otimes x_{hk}^* x_j^*(x_5^*)^n \neq 0$$

$$\nabla_C \nabla_A(1 \otimes x_{12}) = \sum_{i<j} d_{12} d_{ij} \otimes x_{ij}^* \neq 0.$$

We observe that the leading terms of these singular vectors are $d_{12}d_{15} \otimes x_1^{m-1}x_5^*$, $d_{15}d_{45} \otimes x_{45}^*(x_5^*)^{n+1}$ and $d_{12}d_{45} \otimes x_{45}^*$, respectively. (We also observe that the other compositions $\nabla_A\nabla_B$, $\nabla_A\nabla_C$, $\nabla_B\nabla_C$ are not defined). Moreover, one can also verify that $\nabla_A^2 = \nabla_B^2 = \nabla_C^2 = 0$ whenever they are defined: this will also be a consequence of the general treatment of morphisms of degree 2 in Section 9.

Example 4.8.

$$\nabla_C\nabla_B\nabla_A : M(1, 1, 0, 0) \rightarrow M(0, 0, 1, 1)$$

is a nonzero morphism of degree 3. We have that $\nabla_C\nabla_B\nabla_A(x_1x_{12}) = \sum_{j>1, k<l} d_{12}d_{1j}d_{kl} \otimes x_j^*x_{kl}^*$ is a singular vector in $M(0, 0, 1, 1)$ with leading term $d_{12}d_{15}d_{45} \otimes x_{45}^*x_5^*$.

We will prove that the morphisms described in this section are all possible morphisms between minimal Verma modules of degree at most 3.

5. STRUCTURE OF U_-

In order to classify morphisms between generalized Verma modules of a given degree we need to better understand the structure of U_- as an L_0 -module. The main result of this section is the construction of an explicit linear basis of U_- which realizes its structure of L_0 -module in a combinatorial way.

We recall that $(U_-)_d$ denotes the homogeneous component of U_- of degree d . We let

$$\mathcal{I}_d = \{I = (I_1, \dots, I_d) : I_l = (i_l, j_l) \text{ with } 1 \leq i_l, j_l \leq 5 \text{ for every } l = 1, \dots, d\}.$$

If $I = (I_1, \dots, I_d) \in \mathcal{I}_d$ we let $d_I = d_{I_1} \cdots d_{I_d} \in (U_-)_d$, with $d_{I_l} = d_{i_l j_l}$.

We set $[5]^k = \{(t_1, \dots, t_k) \mid t_i \in [5]\}$ and for $T = (t_1, \dots, t_k) \in [5]^k$ we let $\partial_T = \partial_{t_1} \cdots \partial_{t_k}$.

We have that $(U_-)_d$ is spanned by all elements of the form d_I as I varies in \mathcal{I}_d . One can also consider the following filtration of subspaces of $(U_-)_d$: for all $k \leq d/2$ we let

$$(U_-)_{d,k} = \text{Span}\{\partial_T d_I : T \in [5]^k, I \in \mathcal{I}_{d-2k}\}.$$

We have the following chain of inclusions

$$(U_-)_d = (U_-)_{d,0} \supseteq (U_-)_{d,1} \supseteq (U_-)_{d,2} \supseteq \cdots$$

We observe that for all $k \leq d/2$ the subspace $(U_-)_{d,k}$ is also an L_0 -submodule of $(U_-)_d$ and so we have the following isomorphism of L_0 -modules

$$(U_-)_d \cong \bigoplus_{k \leq d/2} (U_-)_{d,k} / (U_-)_{d,k+1},$$

where we let $(U_-)_{d,k} = 0$ if $k > d/2$. For example, we have

$$(U_-)_5 \cong \frac{(U_-)_{5,0}}{(U_-)_{5,1}} \oplus \frac{(U_-)_{5,1}}{(U_-)_{5,2}} \oplus (U_-)_{5,2}.$$

Moreover, one can check that there is an isomorphism of L_0 -modules $\psi : (U_-)_{d,k} / (U_-)_{d,k+1} \rightarrow \text{Sym}^k(\mathbb{C}^{5*}) \otimes \wedge^{d-2k}(\wedge^2 \mathbb{C}^5)$: this isomorphism is simply given by extending multiplicatively the following formulas

$$\psi(\partial_i) = x_i^*, \quad \psi(d_{ij}) = x_{ij}.$$

and so we have that

$$(U_-)_d \cong \bigoplus_{k \leq d/2} \text{Sym}^k(\mathbb{C}^{5*}) \otimes \wedge^{d-2k}(\wedge^2 \mathbb{C}^5)$$

as L_0 -modules. The main goal of this section is to explicitly construct such isomorphism.

We need some further technical notation. If $1 \leq i, j \leq 5$ we let $\overline{(i, j)} = (j, i)$. There is a natural action of B_d , the Weyl group of type B and rank d , on \mathcal{I}_d that can be described in the following way. If $w = (\eta_1 \sigma_1, \dots, \eta_d \sigma_d) \in B_d$, where $\sigma = (\sigma_1, \dots, \sigma_d)$ is a permutation of $[d]$ and $\eta_j = \pm 1$ for all $j \in [d]$, we let

$$w(I) = J$$

where

$$J_j = \begin{cases} I_{\sigma_j} & \text{if } \eta_j = 1 \\ \overline{I_{\sigma_j}} & \text{if } \eta_j = -1. \end{cases}$$

The fact that this is a B_d -action is an easy verification and is left to the reader.

We let \mathcal{S}_d be the set of subsets of $[d]$ of cardinality 2, so that $|\mathcal{S}_d| = \binom{d}{2}$.

Note that elements in \mathcal{I}_d are ordered tuples of ordered pairs, while elements in \mathcal{S}_d are unordered tuples of unordered pairs.

If $\{k, l\} \in \mathcal{S}_d$ and $I \in \mathcal{I}_d$ we let $t_{I_k, I_l} = t_{i_k, j_k, i_l, j_l}$ and $\varepsilon_{I_k, I_l} = \varepsilon_{i_k, j_k, i_l, j_l}$ (see Section 2).

Note that the definitions of t_{I_k, I_l} and ε_{I_k, I_l} do not depend on the order of k and l but only on the set $\{k, l\}$. We also let

$$D_{\{k, l\}}(I) = \frac{1}{2}(-1)^{l+k} \varepsilon_{I_k, I_l} \partial_{t_{I_k, I_l}} \in (U_-)_2.$$

For example, if $I = ((1, 2), (2, 3), (3, 5)) \in \mathcal{I}_3$ then $D_{\{1, 3\}}(I) = \frac{1}{2}(-1)^4 \varepsilon_{12354} \partial_4 = -\frac{1}{2} \partial_4$.

Definition 5.1. A subset S of \mathcal{S}_d is *self-intersection free* if its elements are pairwise disjoint.

For example $S = \{\{1, 3\}, \{2, 5\}, \{4, 7\}\}$ is self-intersection free while $\{\{1, 3\}, \{2, 5\}, \{3, 7\}\}$ is not. We denote by SIF_d the set of self-intersection free subsets of \mathcal{S}_d .

Definition 5.2. Let $\{k, l\}, \{h, m\} \in \mathcal{S}_d$ be disjoint. We say that $\{k, l\}$ and $\{h, m\}$ *cross* if exactly one element in $\{k, l\}$ is between h and m . If $S \in \text{SIF}_d$ we let the crossing number $c(S)$ of S be the number of pairs of elements in S that cross.

For example, if $S = \{\{1, 3\}, \{2, 5\}, \{4, 7\}\}$ then $\{1, 3\}$ and $\{2, 5\}$ cross, $\{1, 3\}$ and $\{4, 7\}$ do not cross, and $\{2, 5\}$ and $\{4, 7\}$ cross, so the crossing number of S is $c(S) = 2$ (see Figure 1 for a graphical interpretation).

Definition 5.3. Let $S = \{S_1, \dots, S_r\} \in \text{SIF}_d$. We let

$$D_S(I) = \prod_{j=1}^r D_{S_j}(I) \in (U_-)_{2r}$$

if $r \geq 2$ and $D_\emptyset(I) = 1$ (note that the order of multiplication is irrelevant as the elements $D_{S_j}(I)$ commute among themselves).

Definition 5.4. For $I = (I_1, \dots, I_d) \in \mathcal{I}_d$ and $S = \{S_1, \dots, S_r\} \in \text{SIF}_d$ we let $C_S(I) \in \mathcal{I}_{d-2r}$ be obtained from I by removing all I_j such that $j \in S_k$ for some $k \in [r]$.

For example, if $d = 7$ and $S = \{\{1, 4\}, \{2, 7\}\}$ then $C_S(I) = (I_3, I_5, I_6)$. We are now ready to give the main definition of this section.




FIGURE 1. A graphical interpretation of the crossing number

Definition 5.5. For all $I \in \mathcal{I}_d$ we let

$$\omega_I = \sum_{S \in \text{SIF}_d} (-1)^{c(S)} D_S(I) d_{C_S(I)} \in (U_-)_d.$$

For example, if $I = (21, 13, 45, 25) \in \mathcal{I}_4$ we have

- $D_\emptyset(I) = 1$;
- $D_{\{1,3\}}(I) = -\frac{1}{2}\partial_3$;
- $D_{\{2,3\}}(I) = +\frac{1}{2}\partial_2$;
- $D_{\{2,4\}}(I) = +\frac{1}{2}\partial_4$;
- $D_{\{1,3\},\{2,4\}}(I) = D_{\{1,3\}}(I)D_{\{2,4\}}(I) = -\frac{1}{4}\partial_3\partial_4$

and all other $D_S(I)$ vanish. We also have, $c(\{\{1, 3\}, \{2, 4\}\}) = 1$ so

$$\omega_I = d_I - \frac{1}{2}\partial_3 d_{13} d_{25} + \frac{1}{2}\partial_2 d_{21} d_{25} + \frac{1}{2}\partial_4 d_{21} d_{45} + \frac{1}{4}\partial_3 \partial_4.$$

Proposition 5.6. For all $I \in \mathcal{I}_d$ and all $g \in B_d$ we have

$$\omega_{g(I)} = (-1)^{\ell(g)} \omega_I,$$

where $\ell(g)$ is the length of g with respect to the Coxeter generators $\{s_0, s_1, s_2, \dots, s_{d-1}\}$, with $s_0 = (-1, 2, 3, \dots, d)$ and s_1, \dots, s_{d-1} the usual simple transpositions.

Proof. It is enough to verify the statement for $g \in \{s_0, \dots, s_{d-1}\}$. If $g = s_0$ we have, for all k, l , $1 \leq k, l \leq d$:

- $\varepsilon_{s_0(I)_k, s_0(I)_l} = (-1)^{\chi_{1 \in \{k, l\}}} \varepsilon_{I_k, I_l}$;
- $t_{s_0(I)_k, s_0(I)_l} = t_{I_k, I_l}$;

hence $D_S(s_0(I)) = (-1)^{\chi_{1 \in S}} D_S(I)$ while $d_{C_S(s_0(I))} = (-1)^{\chi_{1 \notin S}} d_{C_S(I)}$, and therefore we have

$$\begin{aligned}\omega_{s_0(I)} &= \sum_{S \in SIF_d} (-1)^{c(S)} D_S(s_0(I)) d_{C_S(s_0(I))} \\ &= \sum_{S \in SIF_d} (-1)^{c(S)} (-1)^{\chi_{1 \in S}} D_S(I) (-1)^{\chi_{1 \notin S}} d_{C_S(I)} \\ &= -\omega_I.\end{aligned}$$

Now let $h \in \{1, \dots, d-1\}$ and, for notational convenience, let $\sigma = s_h$. We have:

- $\varepsilon_{\sigma(I)_k, \sigma(I)_l} = \varepsilon_{I_{\sigma(k)}, I_{\sigma(l)}}$;
- $t_{\sigma(I)_k, \sigma(I)_l} = t_{I_{\sigma(k)}, I_{\sigma(l)}}$;
- $(-1)^{k+l} = (-1)^{\sigma(k)+\sigma(l)+\chi_{h \in \{k,l\}}+\chi_{h+1 \in \{k,l\}}}$

hence $D_S(\sigma(I)) = (-1)^{\chi_{h \in S} + \chi_{h+1 \in S}} D_{\sigma(S)}(I)$, where “ $h \in S$ ” means that h belongs to some element of S . We also observe that

$$(-1)^{c(S)} = (-1)^{c(\sigma(S))} (-1)^{\chi_{h \in S, h+1 \in S, \{h, h+1\} \notin S}}$$

i.e. the parity of the crossing number of S is opposite to the parity of the crossing number of $\sigma(S)$ precisely if h and $h+1$ belong to two distinct elements of S . Moreover we observe that $d_{C_S(\sigma(I))} = d_{C_{\sigma(S)}(I)}$ if h or $h+1$ belong to S . If $h, h+1$ do not belong to S we have

$$d_{C_S(\sigma(I))} = -d_{C_{\sigma(S)}(I)} - 2D_{\{h, h+1\}}(I) d_{C_{\sigma(\tilde{S})}(I)}$$

where \tilde{S} is obtained from S by adding the pair $\{h, h+1\}$. We are now ready to compute $\omega_{\sigma(I)}$. We have

$$\begin{aligned}\omega_{\sigma(I)} &= \sum_{S \in SIF_d} (-1)^{c(S)} D_S(\sigma(I)) d_{C_S(\sigma(I))} \\ &= \sum_{S \ni h \text{ or } S \ni h+1 \text{ but } S \not\ni \{h, h+1\}} (-1)^{c(S)} D_S(\sigma(I)) d_{C_S(\sigma(I))} \\ &\quad + \sum_{S \not\ni h, h+1} \left((-1)^{c(S)} D_S(\sigma(I)) d_{C_S(s_0(I))} + (-1)^{c(\tilde{S})} D_{\tilde{S}}(\sigma(I)) d_{C_{\tilde{S}}(\sigma(I))} \right) \\ &= \sum_{S \ni h \text{ or } S \ni h+1 \text{ but } S \not\ni \{h, h+1\}} (-1)^{\chi_{h \in S, h+1 \in S}} (-1)^{c(\sigma(S))} (-1)^{\chi_{h \in S} + \chi_{h+1 \in S}} D_{\sigma(S)}(I) d_{C_{\sigma(S)}(I)} \\ &\quad + \sum_{S \not\ni h, h+1} \left((-1)^{c(\sigma(S))} D_{\sigma(S)}(I) (-d_{C_{\sigma(S)}(I)} - 2D_{\{h, h+1\}}(I) d_{C_{\sigma(\tilde{S})}(I)}) + (-1)^{c(\sigma(\tilde{S}))} D_{\sigma(\tilde{S})}(I) d_{C_{\sigma(\tilde{S})}(I)} \right) \\ &= - \sum_{S \ni h \text{ or } S \ni h+1 \text{ but } S \not\ni \{h, h+1\}} (-1)^{c(\sigma(S))} D_{\sigma(S)}(I) d_{C_{\sigma(S)}(I)} - \sum_{S \not\ni h, h+1} (-1)^{c(\sigma(S))} D_{\sigma(S)}(I) d_{C_{\sigma(S)}(I)} \\ &\quad + \sum_{S \not\ni h, h+1} \left((-2)(-1)^{c(\sigma(S))} D_{\sigma(S)}(I) D_{\{h, h+1\}}(I) d_{C_{\sigma(\tilde{S})}(I)} + (-1)^{c(\sigma(\tilde{S}))} D_{\sigma(\tilde{S})}(I) d_{C_{\sigma(\tilde{S})}(I)} \right) \\ &= - \sum_{S \in SIF_d} (-1)^{c(\sigma(S))} D_{\sigma(S)}(I) d_{C_{\sigma(S)}(I)} \\ &= -\omega_I\end{aligned}$$

where we used that $D_{\sigma(S)}(I)D_{h,h+1}(I) = D_{\sigma(\tilde{S})}(I)$ and $(-1)^{c(\sigma(S))} = (-1)^{c(\sigma(\tilde{S}))}$. \square

Corollary 5.7. *If $I = (I_1, \dots, I_d)$ is such that $I_j = I_k$ for some $j < k$, then $\omega_I = 0$; if $I_j = \bar{I}_k$ for some $j \leq k$, then $\omega_I = 0$.*

Now we want to study the action of L_0 on the elements ω_I . If $I = (I_1, \dots, I_d)$ and r appears once in I_b for some b we let $I^{b,s,r}$ be the sequence obtained from I by substituting the letter r in I_b by s . We want to prove the following

Theorem 5.8. *Let $I \in \mathcal{I}_d$ and $r, s \in [5]$, $r \neq s$. Assume that the letter r appears in I_1, \dots, I_c , once in each pair, and does not appear in I_{c+1}, \dots, I_d . Then*

$$x_s \partial_r \omega_I = \sum_{b=1}^c \omega_{I^{b,s,r}}.$$

Proof. For notational convenience, since r and s are fixed in this proof, we simply let $I^b = I^{b,s,r}$ for all $1 \leq b \leq c$. We start by calculating the left-hand side. We have

$$x_s \partial_r \omega_I = x_s \partial_r \cdot \sum_S (-1)^{c(S)} D_S(I) d_{C_S(I)}.$$

Now we observe that $x_s \partial_r \cdot D_{\{k,l\}}(I)$ is non zero if and only if I_k and I_l have the four indices distinct from s , hence k and l cannot be both less than or equal to c or both strictly greater than c . We then assume that $k \leq c$ and $l > c$; in this case we have

$$x_s \partial_r \cdot D_{\{k,l\}}(S) = x_s \partial_r \cdot \left(\frac{1}{2} (-1)^{k+l} \varepsilon_{I_k, I_l} \partial_{t_{I_k, I_l}} \right) = \frac{1}{2} (-1)^{k+l+1} \varepsilon_{I_k, I_l} \partial_r.$$

So we have

$$\begin{aligned} x_s \partial_r \omega_I &= \sum_{k \leq c < l, s \notin I_k, s \notin I_l} \frac{1}{2} (-1)^{k+l+1} \varepsilon_{I_k, I_l} \partial_r \sum_{S \not\ni k, l} (-1)^{c(S)} D_S(I) d_{C_{S \cup \{k, l\}}(I)} + \\ &+ \sum_S (-1)^{c(S)} D_S(I) \sum_{b \leq c, b \notin S} d_{C_S(I^b)}. \end{aligned}$$

Now we compute the right-hand side:

$$\sum_{b \leq c} \omega_{I^b} = \sum_{b \leq c} \sum_S (-1)^{c(S)} D_S(I^b) d_{C_S(I^b)}.$$

Now we observe that if $b \notin S$ we have $D_S(I^b) = D_S(I)$ and so we reduce to prove the following:

$$\sum_{k \leq c < l, s \notin I_k, s \notin I_l} \frac{1}{2} (-1)^{k+l+1} \varepsilon_{I_k, I_l} \partial_r \sum_{S \not\ni k, l} (-1)^{c(S)} D_S(I) d_{C_{S \cup \{k, l\}}(I)} = \sum_{S, b: b \leq c, b \in S} (-1)^{c(S)} D_S(I^b) d_{C_S(I^b)}$$

We notice that if $\{b, b'\} \in S$ with both $b, b' \leq c$ then $D_S(I^b) = -D_S(I^{b'})$ hence we reduce to prove that

$$\begin{aligned} \sum_{k \leq c < l, s \notin I_k, s \notin I_l} \frac{1}{2} (-1)^{k+l+1} \varepsilon_{I_k, I_l} \partial_r \sum_{S \not\ni k, l} (-1)^{c(S)} D_S(I) d_{C_{S \cup \{k, l\}}(I)} &= \\ &= \sum_{b \leq c < l} \sum_{S: S \not\ni b, l} (-1)^{c(S)} D_{\{b, l\}}(I^b) D_S(I^b) d_{C_{S \cup \{b, l\}}(I^b)}. \end{aligned}$$

Finally, in order to prove this last equation we observe that if $b \leq c < l$ then $D_{\{b,l\}}(I^b)$ is nonzero only if $s \notin I_b, I_l$, that in this case $\varepsilon_{(I^b)_b, (I^b)_l} = -\varepsilon_{I_b, I_l}$, that $D_{\{b,l\}}(I^b) = -\frac{1}{2}(-1)^{b+l}\varepsilon_{I_b, I_l}\partial_r$ and that $d_{\mathcal{C}_{S \cup \{b,l\}}(I^b)} = d_{\mathcal{C}_{S \cup \{b,l\}}(I)}$. The proof is complete. \square

If $I = (I_1, \dots, I_d)$ with $I_k = (i_k, j_k)$ we let

$$D_{s \rightarrow r}(\omega_I) = \delta_{r, i_1} \omega_{((s, j_1), I_2, \dots, I_d)} + \delta_{r, j_1} \omega_{((i_1, s), I_2, \dots, I_d)} + \delta_{r, i_2} \omega_{(I_1, (s, j_2), I_3, \dots, I_d)} + \dots + \delta_{r, j_d} \omega_{(I_1, \dots, I_{d-1}, (i_d, s))}.$$

Corollary 5.9. *Let $I = (I_1, \dots, I_d)$ be arbitrary. Then*

$$x_s \partial_r \omega_I = D_{s \rightarrow r}(\omega_I).$$

Proof. If there exists k such that $i_k = j_k$ then $\omega_I = 0$ and clearly also $D_{s \rightarrow r}(\omega_I) = 0$ since all summands in the definition above vanish except possibly two of them which cancel out. If such k does not exist let $w \in B_d$ be such that $J = w(I)$ satisfies the following property: there exists $0 \leq c \leq d$ such that r appears in J_1, \dots, J_c and does not appear in J_{c+1}, \dots, J_d . By Theorem 5.8 we know that the result holds for J hence the result follows since $D_{s \rightarrow r}$ commutes with the action of B_d (we leave this to the reader). \square

Corollary 5.10. *The map*

$$\varphi : \bigoplus_k \text{Sym}^k(\mathbb{C}^{5*}) \otimes \wedge^{d-2k}(\wedge^2 \mathbb{C}^5) \rightarrow (U_-)_d$$

given by

$$\varphi(x_{t_1}^* \cdots x_{t_k}^* \otimes x_{i_1 j_1} \wedge \cdots \wedge x_{i_{d-2k} j_{d-2k}}) = \partial_{t_1} \cdots \partial_{t_k} \omega_{(i_1, j_1), \dots, (i_{d-2k}, j_{d-2k})}$$

for all $k \leq d/2$ and $t_1, \dots, t_k, i_1, j_1, \dots, i_{d-2k}, j_{d-2k} \in [5]$ is an isomorphism of L_0 -modules, hence the set

$$\bigcup_{k \leq d/2} \{\partial_T \omega_I \mid T = (t_1, \dots, t_k) \in [5]^k, t_1 < \cdots < t_k, I \in \mathcal{I}_{d-2k}/B_{d-2k}\}$$

is a basis of $(U_-)_d$.

6. PROPERTIES OF THE DOMINANCE ORDER

In this section we establish simple combinatorial criteria to determine whether the weights of vectors in U_- and $(U_-)^*$ are comparable.

Remark 6.1. If $\varphi : M(V) \rightarrow M(W)$ is a linear map of degree d which satisfies condition (a) of Proposition 3.4 let $\psi : (U_-)_d^* \rightarrow \text{Hom}(V, W)$ be as in Remark 3.6. By Corollary 5.10 we can identify $(U_-)_d^*$ with $\bigoplus_k \text{Sym}^k(\mathbb{C}^5) \otimes \wedge^{d-2k}(\wedge^2(\mathbb{C}^5)^*)$ and we let for all $T = (t_1, \dots, t_k) \in [5]^k$ and $I = (I_1, \dots, I_{d-2k}) \in \mathcal{I}_{d-2k}$ with $I_h = (i_h, j_h)$,

$$\theta_I^T = \psi(x_{t_1} \cdots x_{t_k} \otimes x_{i_1 j_1}^* \wedge \cdots \wedge x_{i_{d-2k} j_{d-2k}}^*).$$

We observe that $\theta_{g(I)}^T = (-1)^{\ell(g)} \theta_I^T$ for every $g \in B_{d-2k}$ hence $\partial_T \omega_I \otimes \theta_I^T$ is invariant with respect to the action of B_{d-2k} on I . We can thus write

$$\Phi = \sum_{\substack{T=(t_1, \dots, t_k): \\ 1 \leq t_1 \leq \dots \leq t_k \leq 5}} \sum_{I \in \mathcal{I}_{d-2k}/B_{d-2k}} \partial_T \omega_I \otimes \theta_I^T.$$

Moreover, we have:

$$x_s \partial_r \cdot \theta_I^T = x_s \partial_r \cdot \theta_{I_1, \dots, I_{d-2k}}^{t_1, \dots, t_k} = \sum_{h=1}^k \Delta_{s \rightarrow r}^h \theta_I^T - \sum_{l=1}^{d-2k} D_{r \rightarrow s}^l \theta_I^T$$

where $\Delta_{s \rightarrow r}^h(\theta_I^T) = \delta_{r, t_h} \theta_I^{t_1, \dots, t_{h-1}, s, t_{h+1}, \dots, t_k}$ and

$$D_{r \rightarrow s}^l(\theta_I^T) = \delta_{i_l, s} \theta_{I_1, \dots, I_{l-1}, (r, j_l), I_{l+1}, \dots, I_{d-2k}}^T + \delta_{s, j_l} \theta_{I_1, \dots, I_{l-1}, (i_l, r), I_{l+1}, \dots, I_{d-2k}}^T.$$

We now study the dominance order on the weights of the elements d_I , ω_I and θ_I^T . This will turn out to play a fundamental role in the study of morphisms of Verma modules.

We observe that d_{kl} is a weight vector for L_0 . Indeed we have:

$$[h_{ij}, d_{kl}] = (\delta_{i,k} + \delta_{i,l} - \delta_{j,k} - \delta_{j,l}) d_{k,l}$$

and so $\lambda_{ij}(d_{kl})$ is the number of occurrences of i minus the number of occurrences of j in $\{k, l\}$. If $I = (i_1, \dots, i_d)$ is a sequence of integers and we let

$$m_k(I) = |\{s \in [d] : i_s = k\}|$$

be the multiplicity of k in I , we have

$$\lambda_{ij}(d_{kl}) = m_i(k, l) - m_j(k, l).$$

More generally, if $I = \{i_1, j_1, \dots, i_d, j_d\}$ and $d_I = d_{i_1 j_1} \cdots d_{i_d j_d}$ we have

$$\lambda_{ij}(d_I) = m_i(I) - m_j(I).$$

In order to understand when the weights of d_I and d_K are comparable in the dominance order, we first observe that the weight of d_I does not depend on the order of its entries. If $I = (i_1, \dots, i_{2d})$ we let $I_o = (i'_1, \dots, i'_{2d})$ be the non decreasing reordering of I . We write $I \leq K$ if $i'_1 \leq k'_1, \dots, i'_{2d} \leq k'_{2d}$ and $I < K$ if $I \leq K$ and at least one of the previous inequalities is strict (notice that this is different that requiring $I \neq K$).

Proposition 6.2. *For all $I, K \in \mathcal{I}_d$ we have $\lambda(d_I) \geq \lambda(d_K)$ if and only if $I \leq K$.*

Proof. We can assume that $I = (i_1, \dots, i_{2d})$ and $K = (k_1, \dots, k_{2d})$ are such that $I = I_o$ and $K = K_o$. We express the difference of the weights as a linear combination of roots. First assume that all entries of I and K coincide except in position r and that $i_r = h$ and $k_r = h + 1$. We have $m_l(I) = m_l(K)$ for all $l \neq h, h + 1$, $m_h(I) = m_h(K) + 1$ and $m_{h+1}(I) = m_{h+1}(K) - 1$. Therefore $\lambda_{l, l+1}(d_I) = \lambda_{l, l+1}(d_K)$ for all $l \neq h - 1, h, h + 1$, $\lambda_{h-1, h}(d_I) = \lambda_{h-1, h}(d_K) + 1$, (if $h \neq 1$), $\lambda_{h, h+1}(d_I) = \lambda_{h, h+1}(d_K) - 2$ and $\lambda_{h+1, h+2}(d_I) = \lambda_{h+1, h+2}(d_K) + 1$ (if $h \neq 4$). Therefore

$$\lambda(d_I) - \lambda(d_K) = \alpha_{h, h+1}.$$

From this we can deduce that

$$\lambda(d_I) - \lambda(d_K) = \alpha_{i_1, k_1} + \alpha_{i_2, k_2} + \cdots + \alpha_{i_{2d}, k_{2d}}.$$

In particular, if $i_1 \leq k_1, \dots, i_{2d} \leq k_{2d}$ then $\lambda(d_I) \geq \lambda(d_K)$. Now we assume that the inequalities $i_1 \leq k_1, \dots, i_{2d} \leq k_{2d}$ are not all satisfied and we let r be minimum such that $i_r > k_r$. If we express $\lambda(d_I) - \lambda(d_K)$ as a linear combination of the simple roots then $\alpha_{k_r, k_{r+1}}$ necessarily appears with a negative coefficient and we are done. \square

Corollary 6.3. *For all $I, K \in \mathcal{I}_d$ and all $T, R \in [5]^k$ we have:*

- (i) $\lambda(\theta_I^T) \leq \lambda(\theta_K^T)$ if and only if $I \leq K$;

(ii) $\lambda(\theta_I^T) \geq \lambda(\theta_I^R)$ if and only if $T \leq R$.

Proof. In order to prove (i) it is sufficient to notice that $\lambda(\theta_I^T) = -(\lambda(\partial_T \omega_I)) = -\lambda(\partial_T) - \lambda(d_I)$ and then use Proposition 6.2.

In order to prove (ii) it is convenient to introduce the following notation. For $t \in [5]$ we let $t^{(1)} < t^{(2)} < t^{(3)} < t^{(4)}$ such that $\{t, t^{(1)}, t^{(2)}, t^{(3)}, t^{(4)}\} = [5]$ and, for $T = (t_1, \dots, t_k) \in [5]^k$, $T^c = (t_1^{(1)}t_1^{(2)}, t_1^{(3)}t_1^{(4)}, \dots, t_k^{(1)}t_k^{(2)}, t_k^{(3)}t_k^{(4)}) \in \mathcal{I}_{2k}$. Then it is enough to notice that $\lambda(\partial_T) = \lambda(d_{T^c})$ and that $T \leq R$ if and only if $T^c \geq R^c$. Then one can use (i). \square

7. DUALITY

Consider a morphism $\varphi : M(V) \rightarrow M(W)$ of generalized Verma modules of degree d associated to an element $\Phi \in (U_-)_d \otimes \text{Hom}(V, W)$. We ask the natural question: does it exist a “related” morphism $\psi : M(W^*) \rightarrow M(V^*)$ of the same degree d ? The first natural candidate to look at is the following: if $\Phi = \sum_i u_i \otimes \theta_i$, where $\{u_i \mid i \in I\}$ is any basis of $(U_-)_d$ and $\theta_i \in \text{Hom}(V, W)$ then we can consider the linear map $\psi : M(W^*) \rightarrow M(V^*)$ associated to $\Psi = \sum_i u_i \otimes \theta_i^*$, where, for all $\theta \in \text{Hom}(V, W)$ we denote by $\theta^* \in \text{Hom}(W^*, V^*)$ the pull-back of θ given by $\theta^*(f) = f \circ \theta$ for all $f \in W^*$. One can easily check that the map ψ does not depend on the chosen basis $\{u_i \mid i \in I\}$ of $(U_-)_d$. It turns out that for $d = 1$ the map ψ is also a morphism of L -modules, but this is not the case in general if the degree d is at least 2.

In this section we develop some tools which will allow us to construct a morphism of L -modules $\psi : M(W^*) \rightarrow M(V^*)$ starting from a morphism $\varphi : M(V) \rightarrow M(W)$ of degree at most 3 and we conjecture that our construction provides such morphism in all degrees.

The main result that we will need is the following.

Proposition 7.1. *Let $\theta_1, \dots, \theta_r, \sigma_1, \dots, \sigma_s \in \text{Hom}(V, W)$ for some L_0 -modules V, W , and let $z_1, \dots, z_t \in L_0$. Let $a_i, b_{j,k} \in \mathbb{C}$ be such that*

$$\sum_i a_i \theta_i(v) + \sum_{j,k} b_{j,k} (z_k \cdot (\sigma_j(v)) + \sigma_j(z_k \cdot v)) = 0 \in W$$

for all $v \in V$. Then

$$\sum_i a_i \theta_i^*(f) + \sum_{j,k} b_{j,k} (z_k \cdot ((-\sigma_j^*)(f)) + (-\sigma_j^*)(z_k \cdot f)) = 0 \in V^*$$

for all $f \in W^*$.

Proof. For all $v \in V$ we have

$$\begin{aligned}
& \left(\sum_i a_i \theta_i^*(f) + \sum_{j,k} b_{j,k} (z_k \cdot ((-\sigma_j^*)(f)) + (-\sigma_j^*)(z_k \cdot f)) \right) (v) \\
&= \sum_i a_i f(\theta_i(v)) + \sum_{j,k} b_{j,k} (\sigma_j^*(f))(z_k \cdot v) + (z_k \cdot f)(-\sigma_j(v)) \\
&= \sum_i a_i f(\theta_i(v)) + \sum_{j,k} b_{j,k} (f(\sigma_j(z_k \cdot v)) + f(z_k \cdot (\sigma_j(v)))) \\
&= f \left(\sum_i a_i \theta_i(v) + \sum_{j,k} b_{j,k} (\sigma_j(z_k \cdot v) + z_k \cdot (\sigma_j(v))) \right) \\
&= 0.
\end{aligned}$$

□

Remark 7.2. We will use Proposition 7.1 also in the following equivalent formulation: let $\theta_1, \dots, \theta_r, \sigma_1, \dots, \sigma_s \in \text{Hom}(V, W)$ for some L_0 -modules V, W and $z_1, \dots, z_t \in L_0$. Let $a_i, b_{j,k} \in \mathbb{C}$ be such that

$$\sum_i a_i \theta_i(v) + \sum_{j,k} b_{j,k} (2z_k \cdot (\sigma_j(v)) - (z_k \cdot \sigma_j)(v)) = 0 \in W$$

for all $v \in V$. Then

$$\sum_i a_i \theta_i^*(f) + \sum_{j,k} b_{j,k} (2z_k \cdot ((-\sigma_j^*)(f)) - (z_k \cdot (-\sigma_j^*))(f)) = 0 \in V^*$$

for all $f \in W^*$.

Conjecture 7.3. Let $\varphi : M(V) \rightarrow M(W)$ be a morphism of degree d associated to $\Phi := \sum_{T,I} \partial_T \omega_I \otimes \theta_I^T$ for some $\theta_I^T \in \text{Hom}(V, W)$. Then the linear map $\psi : M(W^*) \rightarrow M(V^*)$ associated to $\Psi := \sum_{T,I} \partial_T \omega_I \otimes (-1)^{\ell(T)} (\theta_I^T)^*$ is also a morphism of Verma modules, where if $T \in [5]^k$, we let $\ell(T) = k$.

In the following sections we will verify Conjecture 7.3 for morphisms of degree at most 3 as a straightforward application of Proposition 7.1.

Definition 7.4. Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules. The weight $\mu - \lambda$ is called the leading weight of φ .

The reason of the terminology in the previous definition is motivated by the following observation.

Remark 7.5. Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules of leading weight ν . If φ is associated to $\Phi = \sum_i u_i \otimes \theta_i$, where $\{u_i \mid i \in I\}$ is a basis of $(U_-)_d$ consisting of weight vectors, let θ_{i_0} be of maximal weight such that $\theta_{i_0}(s) \neq 0$ for a highest weight vector $s \in F(\lambda)$. Then $\theta_{i_0}(s)$ is a highest weight vector in $F(\mu)$ and so the weight of θ_{i_0} is the leading weight of φ . Therefore if φ has leading weight ν the leading term of the singular vector $\varphi(1 \otimes s)$ is

$$\sum_{i: \lambda(\theta_i) = \nu} u_i \otimes \theta_i(s).$$

We also say that $\theta \in \text{Hom}(V, W)$ has the leading weight of φ if $\theta(s) \neq 0$ and the weight of θ is ν . A general strategy to study a morphism $\varphi : M(V) \rightarrow M(W)$ is to understand elements $\theta \in \text{Hom}(V, W)$ which have the leading weight of φ ; in particular we will often show that there is no such morphism by showing that there is no $\theta \in \text{Hom}(V, W)$ that may possibly have the leading weight of a morphism.

Whenever Conjecture 7.3 holds the next result allows us to simplify the classification of morphisms.

Remark 7.6. Let $\varphi : M(V) \rightarrow M(W)$ and $\psi : M(W^*) \rightarrow M(V^*)$ be morphisms of Verma modules and let $\nu = (a, b, c, d)$ be the leading weight of φ . Then the leading weight of ψ is $-\nu^* = -(d, c, b, a)$.

8. MORPHISMS OF DEGREE ONE

In this section we classify morphisms of degree one between generalized Verma modules, slightly simplifying Rudakov's argument [8].

We let $C(a, b, c)$ be the set of cyclic permutations of a, b, c , i.e., $C(a, b, c) = \{(a, b, c), (b, c, a), (c, a, b)\}$.

Theorem 8.1. *Let $\varphi : M(V) \rightarrow M(W)$ be a linear map of degree one associated to*

$$\Phi = \sum_{I \in \mathcal{I}_1/B_1} \omega_I \otimes \theta_I$$

such that $L_0 \cdot \Phi = 0$. Then φ is a morphism of Verma modules if and only if for all distinct $a, b, c, p \in [5]$ and for all $v \in V$ we have

$$(1) \quad \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}(v)) = 0.$$

Proof. By Proposition 3.4 it is enough to check when $x_p d_{pq} \Phi(v) = 0$ for all $p, q \in [5]$. For notational convenience we let $Q = (p, q)$ and $\{a, b, c, p, q\} = [5]$. We have:

$$\begin{aligned} x_p d_Q \Phi(v) &= x_p d_Q \sum_{I \in \mathcal{I}_1/B_1} \omega_I \otimes \theta_I(v) = x_p d_Q \sum_{I \in \mathcal{I}_1/B_1} d_I \otimes \theta_I(v) \\ &= \sum_{I \in \mathcal{I}_1/B_1} \varepsilon_{Q, I} x_p \partial_{t_{Q, I}} \cdot (\theta_I(v)) = \varepsilon_{pqabc} \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}(v)). \end{aligned}$$

□

Remark 8.2. We point out that Equation (1) satisfies the hypotheses of Proposition 7.1 since in this case

$$x_p \partial_\gamma \cdot (\theta_{\alpha\beta}(v)) = \theta_{\alpha\beta}(x_p \partial_\gamma \cdot v)$$

hence we can write

$$x_p \partial_\gamma \cdot (\theta_{\alpha\beta}(v)) = \frac{1}{2} (x_p \partial_\gamma \cdot (\theta_{\alpha\beta}(v)) + \theta_{\alpha\beta}(x_p \partial_\gamma \cdot v)).$$

Conjecture 7.3 then holds in degree one. This will be also confirmed by Theorem 8.4.

Proposition 8.3. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules of degree one and let θ_{hk} have the leading weight of φ . Then if $i < j$ are distinct from h, k we have*

$$\mu_{ij} = -\chi_{i < h < j} - \chi_{i < k < j}.$$

Proof. Consider Equation (1) with $p = j$, $c = i$, $a = h$, $b = k$ and $v = s$ a highest weight vector in $F(\lambda)$:

$$x_j \partial_i.(\theta_{hk}(s)) + x_j \partial_k.(\theta_{ih}(s)) + x_j \partial_h.(\theta_{ki}(s)) = 0.$$

Now we apply $x_i \partial_j$ to this equation. We have

$$h_{ij}.(\theta_{hk}(s)) - \chi_{i < k < j} \theta_{kh}(s) - \chi_{i < h < j} \theta_{kh}(s) = 0$$

and the result follows. \square

Theorem 8.4. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules of degree one. Then one of the following occurs:*

- $\lambda = (m, n+1, 0, 0)$, $\mu = (m, n, 0, 0)$ for some $m, n \geq 0$ and, up to a scalar, $\varphi = \nabla_A$.
- $\lambda = (m+1, 0, 0, n)$, $\mu = (m, 0, 0, n+1)$ for some $m, n \geq 0$ and, up to a scalar, $\varphi = \nabla_B$.
- $\lambda = (0, 0, m, n)$, $\mu = (0, 0, m+1, n)$ for some $m, n \geq 0$ and, up to a scalar, $\varphi = \nabla_C$.

Proof. Let θ_{hk} have the leading weight of φ . By Proposition 8.3 we have that if $(h, k) \neq (1, 2), (1, 5), (4, 5)$ we can find i, j such that $\mu_{ij} < 0$, a contradiction. Proposition 8.3 also provides

- $\mu_{3,5} = 0$ if $(h, k) = (1, 2)$;
- $\mu_{2,4} = 0$ if $(h, k) = (1, 5)$;
- $\mu_{1,3} = 0$ if $(h, k) = (4, 5)$,

and the rest follows using Lemma 4.1 and Proposition 3.8 recalling that $\lambda(\theta_{hk}) = -\lambda(d_{hk})$. \square

9. MORPHISMS OF DEGREE 2

In this section we provide a complete classification of morphisms between Verma modules of degree 2. We will make use of the following preliminary result which holds in a much wider generality. Here and in what follows we denote by (p, q, a, b, c) any permutation of $[5]$ and we set $Q = (p, q)$.

Lemma 9.1. *Suppose that $\Phi = \sum_{T, I} \partial_T \omega_I \otimes \theta_I^T$ defines a morphism of Verma modules $\varphi : M(V) \rightarrow M(W)$. Then for all $t_1 \dots t_h \in [5]$, $I_1, \dots, I_k \in \mathcal{I}_1$ and $v \in V$ we have*

$$\begin{aligned} \sum_{I, J_1, \dots, J_r \in \mathcal{I}_1} \varepsilon_{Q, I} x_p \partial_{t_{Q, I}} d_{J_1} \cdots d_{J_r} \otimes \theta_{I_1, \dots, I_k, I, J_1, \dots, J_r}^{t_1, \dots, t_h}(v) &= 2 \sum_{\substack{(\alpha, \beta, \gamma) \in C(a, b, c) \\ H_1, \dots, H_r \in \mathcal{I}_1}} \varepsilon_{pqabc} d_{H_1} \cdots d_{H_r} \otimes \\ &\left(\theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(x_p \partial_\gamma.v) + \sum_{s=1}^h \Delta_{p \rightarrow \gamma}^s \theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(v) - \sum_{s=1}^k D_{\gamma \rightarrow p}^s \theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(v) \right) \end{aligned}$$

Proof. Using the definitions of $D_{a \rightarrow b}^h$, of $\Delta_{a \rightarrow b}^h$, of $\theta_{I_1, \dots, I_k}^{t_1, \dots, t_h}$ and of the action of L_0 on the latter elements, we have

$$\begin{aligned} \sum_{I, J_1, \dots, J_r} \varepsilon_{Q, I} x_p \partial_{t_{Q, I}} d_{J_1} \cdots d_{J_r} \otimes \theta_{I_1, \dots, I_k, I, J_1, \dots, J_r}^{t_1, \dots, t_h}(v) = -2 \sum_{\substack{(\alpha, \beta, \gamma) \in C(a, b, c) \\ H_1, \dots, H_r \in \mathcal{I}_1}} \varepsilon_{pqabc} d_{H_1} \cdots d_{H_r} \otimes \\ \left((x_p \partial_\gamma \cdot \theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h})(v) - \sum_{s=1}^h \Delta_{p \rightarrow \gamma}^s \theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(v) + \sum_{s=1}^k D_{\gamma \rightarrow p}^s \theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(v) \right. \\ \left. - x_p \partial_\gamma (\theta_{I_1, \dots, I_k, \alpha\beta, H_1, \dots, H_r}^{t_1, \dots, t_h}(v)) \right) \end{aligned}$$

from which the thesis follows. \square

We are now ready to state the following characterization result.

Theorem 9.2. *Let $\varphi : M(V) \rightarrow M(W)$ be a linear map of degree 2 associated to*

$$\Phi = \sum_{(I, J) \in \mathcal{I}_2/B_2} \omega_{I, J} \otimes \theta_{I, J} + \sum_{t=1}^5 \partial_t \otimes \theta^t$$

such that $x \cdot \Phi = 0$ for all $x \in L_0$. Then φ is a morphism of Verma modules if and only if for all $K \in \mathcal{I}_1$ and all $v \in V$ we have

$$-\chi_{(K \in B_1 Q)} \theta^p(v) + \frac{1}{2} \varepsilon_{pqabc} \sum_{(\alpha\beta\gamma) \in C(a, b, c)} \left(-((x_p \partial_\gamma) \cdot \theta_{\alpha\beta, K})(v) + 2x_p \partial_\gamma \cdot (\theta_{\alpha\beta, K}(v)) \right) = 0$$

Proof. By Proposition 3.4 we have that φ is a morphism of Verma modules if and only if

$$x_p d_Q \left(\sum_{(I, J) \in \mathcal{I}_2/B_2} \omega_{I, J} \otimes \theta_{I, J}(v) + \sum_t \partial_t \otimes \theta^t(v) \right) = 0$$

for all $v \in V$. It is convenient for us to consider the first sum running over all $(I, J) \in \mathcal{I}_2$ and so we have

$$\begin{aligned} x_p d_Q \left(\frac{1}{8} \sum_{(I, J) \in \mathcal{I}_2} \omega_{I, J} \otimes \theta_{I, J}(v) + \sum_t \partial_t \otimes \theta^t(v) \right) \\ (2) \quad = x_p d_Q \left(\frac{1}{8} \sum_{I, J} (d_I d_J - \frac{1}{2} \varepsilon_{I, J} \partial_{t_{I, J}}) \otimes \theta_{I, J}(v) + \sum_t \partial_t \otimes \theta^t(v) \right). \end{aligned}$$

We split Equation (2) into three parts:

In the first part of Equation (2) we have, using Lemma 9.1,

$$\begin{aligned}
x_p d_Q \sum_{I,J} d_I d_J \otimes \theta_{I,J}(v) &= \sum_{I,J} \left(\varepsilon_{Q,I}(x_p \partial_{t_{Q,I}}) d_J - \varepsilon_{Q,J} d_I (x_p \partial_{t_{Q,J}}) \right) \otimes \theta_{I,J}(v) \\
&= 2 \sum_H d_H \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \theta_{\alpha\beta,H}(x_p \partial_\gamma \cdot v) \\
&\quad - 2 \sum_I d_I \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} (\theta_{I,\alpha\beta}(x_p \partial_\gamma \cdot v) - D_{\gamma \rightarrow p}^1 \theta_{I,\alpha\beta}(v)) \\
&= 4 \sum_H d_H \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} (\theta_{\alpha\beta,H}(x_p \partial_\gamma \cdot v) + \frac{1}{2} (x_p \partial_\gamma \cdot \theta_{\alpha\beta,H})(v)) \\
&= 4 \sum_H d_H \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} (x_p \partial_\gamma \cdot (\theta_{\alpha\beta,H}(v)) - \frac{1}{2} (x_p \partial_\gamma \cdot \theta_{\alpha\beta,H})(v))
\end{aligned}$$

where the sums run over $I, J, H \in \mathcal{I}_1$ and $(\alpha, \beta, \gamma) \in C(a, b, c)$.

In the second part of Equation (2) we have

$$\sum_{I,J} \frac{1}{2} \varepsilon_{I,J} \partial_{t_{I,J}} \otimes \theta_{I,J}(v) = 0$$

since the term indexed by (I, J) cancels the term indexed by (J, I) .

In the third part of Equation (2) we have:

$$\sum_t x_p d_Q \partial_t \otimes \theta^t(v) = -d_Q \otimes \theta^p(v).$$

Putting the three parts together Equation (2) becomes

$$\begin{aligned}
&x_p d_Q \left(\frac{1}{8} \sum_{I,J \in \mathcal{I}_1} \omega_{I,J} \otimes \theta_{I,J}(v) + \sum_t \partial_t \otimes \theta^t(v) \right) \\
&= \sum_{K \in \mathcal{I}_1/B_1} d_K \otimes \left(-\chi_{(K \in B_1 Q)} \theta^p(v) + \varepsilon_{pqabc} \sum_{(\alpha\beta\gamma) \in C(a,b,c)} -\frac{1}{2} (x_p \partial_\gamma \cdot \theta_{\alpha\beta,K})(v) + x_p \partial_\gamma \cdot (\theta_{\alpha\beta,K}(v)) \right)
\end{aligned}$$

and the result follows. \square

We deduce that Conjecture 7.3 holds for morphisms of degree 2 and in particular we have the following duality result for degree 2 morphisms.

Corollary 9.3. *Let $\varphi : M(V) \rightarrow M(W)$ be a morphism of Verma modules of degree 2 associated to*

$$\Phi = \sum_{(I,J) \in \mathcal{I}_2/B_2} \omega_{I,J} \otimes \theta_{I,J} + \sum_t \partial_t \otimes \theta^t.$$

Then the linear map $\psi : M(W^) \rightarrow M(V^*)$ associated to*

$$\Psi = \sum_{(I,J) \in \mathcal{I}_2/B_2} \omega_{I,J} \otimes \theta_{I,J}^* + \sum_t \partial_t \otimes (-\theta^t)^*$$

is also a morphism of Verma modules.

Proof. This is an immediate consequence of Remark 7.2 and Theorem 9.2. \square

Corollary 9.4. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules and $s \in F(\lambda)$ a highest weight vector. Then for all $K \in \mathcal{I}_1$ we have*

$$2\chi_{K \in B_1 Q} \varepsilon_{pqabc} \theta^p(s) + \sum_{(\alpha\beta\gamma) \in C(abc)} \left((-1)^{\chi_{p>\gamma}} (x_p \partial_\gamma \cdot \theta_{\alpha\beta, K})(s) + 2\chi_{p>\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta, K}(s)) \right) = 0$$

Proof. This result immediately follows from Theorem 9.2 by observing that if $p < \gamma$ then $x_p \partial_\gamma \cdot s = 0$. \square

In the following results we fix a morphism $\varphi : M(\lambda) \rightarrow M(\mu)$ of Verma modules of degree 2 associated to $\Phi = \sum \omega_{I,J} \otimes \theta_{I,J} + \sum \partial_t \otimes \theta^t$ and we exploit Corollary 9.4 to obtain some constraints on the weights λ and μ . The next result is analogous to Proposition 8.3.

Proposition 9.5. *Let $h, k, l, m \in [5]$ be such that $\theta_{hk,lm}$ has the leading weight of φ . Let $1 \leq i < j \leq 5$ be such that $j \neq h, k, l, m$ and $i \neq h, k$. Then*

$$\mu_{ij} = -\chi_{i<h<j} - \chi_{i<k<j}.$$

Proof. By Corollary 9.4 used with $a = i$, $b = h$, $c = k$, $p = j$ and $K = (l, m)$, observing that $x_j \partial_\gamma \cdot \theta_{\alpha\beta, K} = 0$ for all $(\alpha, \beta, \gamma) \in C(i, h, k)$, we obtain the following relation

$$x_j \partial_i \cdot (\theta_{hk,lm}(s)) + \chi_{h<j} x_j \partial_h \cdot (\theta_{ki,lm}(s)) + \chi_{k<j} x_j \partial_k \cdot (\theta_{ih,kl}(s)) = 0.$$

Applying $x_i \partial_j$ to this equation we have

$$\begin{aligned} h_{ij} \cdot (\theta_{hk,lm}(s)) + \chi_{h<j} (x_i \partial_h \cdot (\theta_{ki,lm}(s)) - x_j \partial_h \cdot (\theta_{kj,lm}(s))) \\ + \chi_{k<j} (x_i \partial_k \cdot (\theta_{ih,lm}(s)) - x_j \partial_k \cdot (\theta_{jh,lm}(s))) = 0 \end{aligned}$$

Since $\theta_{hk,lm}$ has the leading weight of φ , if $h < j$ we necessarily have $\theta_{kj,lm}(s) = 0$, by Corollary 6.3. Similarly, if $k < j$, we have $\theta_{jh,lm}(s) = 0$. Therefore the previous equation becomes

$$h_{ij} \cdot (\theta_{hk,lm}(s)) + \chi_{h<j} x_i \partial_h \cdot (\theta_{ki,lm}(s)) + \chi_{k<j} x_i \partial_k \cdot (\theta_{ih,lm}(s)) = 0$$

Again, if $i > h$, we have $\theta_{ki,lm}(s) = 0$ and otherwise we have $x_i \partial_h \cdot (\theta_{ki,lm}(s)) = -\theta_{kh,lm}(s)$ and similarly for the other term, and so we have

$$h_{ij} \cdot (\theta_{hk,lm}(s)) - \chi_{h<j} \chi_{i<h} \theta_{kh,lm}(s) - \chi_{k<j} \chi_{i<k} \theta_{kh,lm}(s) = 0$$

i.e.,

$$h_{ij} \cdot (\theta_{hk,lm}(s)) = -(\chi_{i<h<j} + \chi_{i<k<j}) \theta_{hk,lm}(s).$$

\square

Proposition 9.6. *Let $i, h, k, l, m \in [5]$, with i, h, k, m distinct and $i < m$, be such that $\theta_{hk,lm}$ has the leading weight of φ . Then*

$$\begin{aligned} h_{im} \cdot (\theta_{hk,lm}(s)) = \\ \left(\frac{1}{2} - \chi_{i<h<m} - \chi_{i<k<m} \right) \theta_{hk,lm}(s) - \varepsilon_{mlhki} \theta^i(s) - \frac{1}{2} \left((-1)^{\chi_{h<m}} \theta_{hl,km}(s) + (-1)^{\chi_{k<m}} \theta_{hm,kl}(s) \right). \end{aligned}$$

Proof. We consider Corollary 9.4 with $a = h$, $b = k$, $c = i$, $p = m$ and $K = (l, m)$. We observe that

$$\varepsilon_{pqabc} \chi_{K \in B_1 Q} = \varepsilon_{mqhki} \chi_{l=q} = \varepsilon_{mlhki}$$

and so we obtain

$$\begin{aligned} \varepsilon_{mlhki}\theta^m(s) + \frac{1}{2}\left((-1)^{\chi_{h<m}}\theta_{ki,hl}(s) + (-1)^{\chi_{k<m}}\theta_{ih,kl}(s) - \theta_{hk,il}(s)\right) \\ + \chi_{h<m}x_m\partial_{h\cdot}(\theta_{ki,lm}(s)) + \chi_{k<m}x_m\partial_{k\cdot}(\theta_{ih,lm}(s)) + x_m\partial_{i\cdot}(\theta_{hk,lm}(s)) = 0 \end{aligned}$$

We apply $x_i\partial_m$ to this equation and we obtain

$$\begin{aligned} \varepsilon_{mlhki}\theta^i(s) - \frac{1}{2}\left((-1)^{\chi_{h<m}}\theta_{km,hl}(s) + (-1)^{\chi_{k<m}}\theta_{mh,kl}(s) + \theta_{hk,ml}(s)\right) \\ - \chi_{i<h<m}\theta_{kh,lm}(s) - \chi_{i<k<m}\theta_{kh,lm}(s) + h_{im\cdot}(\theta_{hk,lm}(s)) = 0 \end{aligned}$$

and the result follows. \square

Proposition 9.7. *Let $h, k, m, i \in [5]$ be distinct, $i < m$, be such that $\theta_{hk,hm}$ has the leading weight of $\varphi : M(\lambda) \rightarrow M(\mu)$. Then*

$$\mu_{i,m} = \chi_{k<m} - \chi_{i<h<m} - \chi_{i<k<m}$$

and

$$\lambda_{i,m} = \chi_{k<m} - \chi_{i<h<m} - \chi_{i<k<m} - 1.$$

Proof. We use Proposition 9.6 with $l = h$ and deduce

$$\begin{aligned} h_{im\cdot}(\theta_{hk,hm}(s)) &= \left(\frac{1}{2} - \chi_{i<h<m} - \chi_{i<k<m}\right)\theta_{hk,hm}(s) - \frac{1}{2}(-1)^{\chi_{k<m}}\theta_{hm,kh} \\ &= \left(\frac{1}{2} - \frac{1}{2}(-1)^{\chi_{k<m}} - \chi_{i<h<m} - \chi_{i<k<m}\right)\theta_{hk,hm}(s) \\ &= (\chi_{k<m} - \chi_{i<h<m} - \chi_{i<k<m})\theta_{hk,hm}(s). \end{aligned}$$

and the first part of the statement follows. The second part is an easy consequence since

$$\lambda_{i,m}(\theta_{hk,hm}) = 1.$$

\square

Theorem 9.8. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of degree 2. Then one of the following occurs:*

- (1) $\lambda = (1, 0, 0, n)$, $\mu = (0, 0, 1, n+1)$ for some $n \geq 0$ and, up to a scalar, $\varphi = \nabla_C \nabla_B$;
- (2) $\lambda = (n+1, 1, 0, 0)$, $\mu = (n, 0, 0, 1)$ for some $n \geq 0$ and, up to a scalar, $\varphi = \nabla_B \nabla_A$;
- (3) $\lambda = (0, 1, 0, 0)$, $\mu = (0, 0, 1, 0)$, and, up to a scalar, $\varphi = \nabla_C \nabla_A$.

Proof. We first make the following observation that will allow us to simplify several arguments. If $\nu \in \Lambda$ is any weight, by Corollary 9.3, if the statement holds for all morphisms of leading weight ν then it holds also for all morphisms of leading weight $-\nu^*$.

We let s be a highest weight vector of $F(\lambda)$ and we suppose that $\theta_{hk,lm}$ has the leading weight of φ . Let us first assume $|\{h, k, l, m\}| = 3$ i.e., without loss of generality, $h = l$.

By Corollary 9.4 with $K = (p, a)$ we have:

$$\begin{aligned} (3) \quad & -((-1)^{\chi_{b<p}} + (-1)^{\chi_{c<p}})\theta_{ab,ca}(s) \\ & + 2\chi_{a<p}x_p\partial_{a\cdot}(\theta_{bc,pa}(s)) + 2\chi_{b<p}x_p\partial_{b\cdot}(\theta_{ca,pa}(s)) + 2\chi_{c<p}x_p\partial_{c\cdot}(\theta_{ab,pa}(s)) = 0. \end{aligned}$$

Using this equation with $a = h$, $b = k$, $c = m$, since $\theta_{hk, hm}$ has the leading weight of φ , we immediately obtain

$$((-1)^{\chi_{k < p}} + (-1)^{\chi_{m < p}})\theta_{hk, hm}(s) = 0.$$

In particular, if we can choose p such that $p > k, m$ or $p < k, m$ we have $\theta_{hk, hm}(s) = 0$, a contradiction. So we reduce to study the following cases: (a) $k = 1, m = 5$; (b) $k = 2, m = 5, h = 1$; (c) $k = 1, m = 4, h = 5$.

- (a) By duality, since $\lambda(\theta_{21, 25}) = -(\lambda(\theta_{41, 45}))^*$, it is enough to consider only the cases $h = 2, 3$; we have, by Proposition 9.5,

$$\mu_{14} = -\chi_{1 < h < 4} - \chi_{1 < 5 < 4} = -1,$$

a contradiction.

- (b) In this case we have, by Proposition 9.5

$$\mu_{23} = -\chi_{2 < 1 < 3} - \chi_{2 < 5 < 3} = 0$$

and by Proposition 9.7 we have

$$\mu_{35} = \chi_{2 < 5} - \chi_{3 < 1 < 5} - \chi_{3 < 2 < 5} = 1.$$

Since the leading weight of φ is $\lambda(\theta_{12, 15}) = (-1, -1, 0, 1)$ we conclude that $\mu = (n, 0, 0, 1)$ for some $n \geq 0$ and so $\lambda = (n + 1, 1, 0, 0)$. The leading term of the singular vector $\varphi(1 \otimes s)$ is $\omega_{12, 15} \otimes \theta_{12, 15}(s) = d_{12}d_{15} \otimes \theta_{12, 15}(s)$ hence, up to a scalar, $\varphi = \nabla_B \nabla_A$ by Proposition 3.8.

- (c) Since $\lambda(\theta_{51, 54}) = -\lambda(\theta_{12, 15})^*$ this follows from case (b) and we obtain in this case the morphism $\nabla_C \nabla_B$.

This concludes the study of all possible $\theta_{hk, lm}$ having the leading weight of φ with h, k, l, m not distinct.

In order to deal with the case where h, k, l, m are distinct we let p be different from h, k, l, m . If $p = 4, 5$ we apply Proposition 9.5 with $i = 1$ and $j = p$ and we get that $\mu_{1p} < 0$ hence $\theta_{hk, lm}$ does not have the leading weight of φ . By Corollary 9.4 we also have $\theta^p(s) = 0$ and so also θ^p can not have the leading weight of φ .

For $p = 1$ we have $\lambda(\theta^1) = -\lambda(\theta^5)^*$ and if $p = 2$ we have $\lambda(\theta^2) = -\lambda(\theta^4)^*$ and so these cases follows from the previous discussion by Corollary 9.3.

For $p = 3$ Proposition 9.5 with $i = 1$, $j = 3$ shows that $\theta_{14, 25}$ and $\theta_{15, 24}$ cannot have the leading weight of φ , i.e. $\theta_{14, 25}(s) = \theta_{15, 24}(s) = 0$, and that if $\theta_{12, 45}$ has leading weight then $\mu_{1, 3} = 0$. Besides, by Corollary 9.4, $\theta_{12, 45}(s) = 2\theta^3(s)$. By Proposition 9.6 we immediately get

$$h_{35} \cdot (\theta_{12, 45}(s)) = \theta_{12, 45}(s)$$

and so $\mu_{3, 5} = 1$. Since the leading weight is $\lambda(\theta_{12, 45}) = (0, -1, 1, 0)$ we conclude that $\mu = (0, 0, 1, 0)$ and so $\lambda = (0, 1, 0, 0)$. The leading term of $\varphi(1 \otimes s)$ is

$$\omega_{12, 45} \otimes \theta_{12, 45}(s) + \partial_3 \otimes \theta^3(s) = d_{12}d_{45} \otimes \theta^3(s)$$

hence, up to a scalar, $\varphi = \nabla_C \nabla_A$ by Proposition 3.8.

□

10. MORPHISMS OF DEGREE 3

This section is dedicated to the study of morphisms of Verma modules of degree three. We consider a linear map $\varphi : M(\lambda) \rightarrow M(\mu)$ of degree three associated to

$$\Phi = \sum_{I \in \mathcal{I}_3/B_3} \omega_I \otimes \theta_I + \sum_{t \in [5], I \in \mathcal{I}_1/B_1} \partial_t \omega_I \otimes \theta_I^t.$$

As in the case of morphisms of degree one and two, our goal is to establish necessary and sufficient conditions to ensure that φ is a morphism of Verma modules.

Lemma 10.1. *If $x \cdot \Phi = 0$ for every $x \in L_0$, then the following relation holds for every $v \in F(\lambda)$:*

$$\sum_{I \in \mathcal{I}_3} \omega_I \otimes \theta_I(v) = \sum_{I \in \mathcal{I}_3} d_I \otimes \theta_I(v).$$

Proof. Indeed we have

$$\begin{aligned} \sum_{I \in \mathcal{I}_3} \omega_I \otimes \theta_I(v) &= \sum_{I \in \mathcal{I}_3} d_I \otimes \theta_I(v) \\ &+ \sum_{I_1, I_2, I_3} \left(-\frac{1}{2} \varepsilon_{I_1, I_2} \partial_{t_{I_1, I_2}} d_{I_3} + \frac{1}{2} \varepsilon_{I_1, I_3} \partial_{t_{I_1, I_3}} d_{I_2} - \frac{1}{2} \varepsilon_{I_2, I_3} \partial_{t_{I_2, I_3}} d_{I_1} \right) \otimes \theta_{I_1, I_2, I_3}(v) \end{aligned}$$

and the last sum vanishes since the coefficients of $\theta_{I_1, I_2, I_3}(v)$ and $\theta_{I_3, I_2, I_1}(v)$ coincide. \square

Theorem 10.2. *Let us assume that $x \cdot \Phi = 0$ for every $x \in L_0$. Then φ is a morphism of Verma modules if and only if for every $H, L \in \mathcal{I}_1$, every permutation (p, q, a, b, c) of $[5]$ and every $v \in F(\lambda)$, the following equations hold:*

$$(4) \quad \chi_{L \in B_1 Q} \theta_H^p(v) + \frac{1}{2} \varepsilon_{pqabc} \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} \left(- (x_p \partial_\gamma \cdot \theta_{\alpha\beta, H, L})(v) + 2x_p \partial_\gamma \cdot (\theta_{\alpha\beta, H, L}(v)) \right) = 0$$

$$(5) \quad \frac{1}{4} \theta_{ab, bc, cq}(v) + \frac{1}{4} \theta_{ac, cb, bq}(v) + \frac{1}{2} \varepsilon_{pqabc} \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} \left(- (x_p \partial_\gamma \cdot \theta_{\alpha\beta}^a)(v) + 2x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^a(v)) \right) = 0$$

$$(6) \quad \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^p(v)) = 0$$

$$(7) \quad \varepsilon_{pqabc} \sum_{(\alpha, \beta, \gamma) \in C(a, b, c)} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^q(v)) - \frac{1}{2} \theta_{ab, bc, ca}(v) = 0.$$

Proof. By Proposition 3.4 we need to compute $x_p d_Q \Phi(v)$ for $v \in F(\lambda)$. We compute the different summands separately. Using Lemma 10.1 and Lemma 9.1 we have

$$\begin{aligned}
x_p d_Q \sum_{(I,J,K) \in \mathcal{I}_3/B_3} \omega_{I,J,K} \otimes \theta_{I,J,K}(v) &= \frac{1}{48} \sum_{I,J,K \in \mathcal{I}_3} \omega_{I,J,K} \otimes \theta_{I,J,K}(v) \\
&= \frac{1}{48} x_p d_Q \sum_{I,J,K} d_I d_J d_K \otimes \theta_{I,J,K}(v) \\
&= \frac{1}{48} \sum_{I,J,K} (\varepsilon_{Q,I} x_p \partial_{t_{Q,I}} d_J d_K - d_I \varepsilon_{Q,J} x_p \partial_{t_{Q,J}} d_K + d_I d_J \varepsilon_{Q,K} x_p \partial_{t_{Q,K}}) \otimes \theta_{I,J,K}(v) \\
&= \frac{1}{48} \sum_{H,L} d_H d_L \otimes 2\varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(D_{\gamma \rightarrow p}^2 \theta_{\alpha\beta,H,L}(v) + 2D_{\gamma \rightarrow p}^3 \theta_{\alpha\beta,H,L}(v) + 3x_p \partial_{\gamma \cdot} (\theta_{\alpha\beta,H,L}(v)) \right),
\end{aligned}$$

where the sums run over $I, J, K \in \mathcal{I}_1$ and $(\alpha, \beta, \gamma) \in C(a, b, c)$.

Recalling that $d_H d_L = \omega_{H,L} + \frac{1}{2} \varepsilon_{H,L} \partial_{t_{H,L}}$ we have:

$$\begin{aligned}
x_p d_Q \sum_{(I,J,K) \in \mathcal{I}_3/B_3} \omega_{I,J,K} \otimes \theta_{I,J,K}(v) &= \frac{1}{48} \sum_{H,L} \omega_{H,L} \otimes 2\varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(D_{\gamma \rightarrow p}^2 \theta_{\alpha\beta,H,L}(v) + 2D_{\gamma \rightarrow p}^3 \theta_{\alpha\beta,H,L}(v) + 3x_p \partial_{\gamma \cdot} (\theta_{\alpha\beta,H,L}(v)) \right) \\
&\quad + \frac{1}{48} \sum_{H,L} \partial_{t_{H,L}} \otimes \varepsilon_{H,L} \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(D_{\gamma \rightarrow p}^2 \theta_{\alpha\beta,H,L}(v) + 2D_{\gamma \rightarrow p}^3 \theta_{\alpha\beta,H,L}(v) + 3x_p \partial_{\gamma \cdot} (\theta_{\alpha\beta,H,L}(v)) \right) \\
&= \frac{1}{48} \sum_{(H,L) \in \mathcal{I}_2/B_2} \omega_{H,L} \otimes 2\varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(12D_{\gamma \rightarrow p}^2 \theta_{\alpha\beta,H,L}(v) + 12D_{\gamma \rightarrow p}^3 \theta_{\alpha\beta,H,L}(v) + 24x_p \partial_{\gamma \cdot} (\theta_{\alpha\beta,H,L}(v)) \right) \\
&\quad + \frac{1}{48} \sum_{(H,L) \in \mathcal{I}_2/B_2} \partial_{t_{H,L}} \otimes \varepsilon_{H,L} \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(-4D_{\gamma \rightarrow p}^2 \theta_{\alpha\beta,H,L}(v) + 4D_{\gamma \rightarrow p}^3 \theta_{\alpha\beta,H,L}(v) \right) \\
&= \sum_{(H,L) \in \mathcal{I}_2/B_2} \omega_{H,L} \otimes \frac{1}{2} \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(-(x_p \partial_{\gamma \cdot} \theta_{\alpha\beta,H,L})(v) + 2x_p \partial_{\gamma \cdot} (\theta_{\alpha\beta,H,L}(v)) \right) \\
&\quad + \partial_q \otimes -\frac{1}{2} \theta_{ab,bc,ca}(v) + \sum_{\alpha\beta\gamma} \partial_\alpha \otimes \frac{1}{4} (\theta_{\alpha\beta,\beta\gamma,\gamma q}(v) + \theta_{\alpha\gamma,\gamma\beta,\beta q}(v)).
\end{aligned}$$

We also need the following computation

$$\begin{aligned}
x_p d_Q \sum_{t \in [5]} \sum_{I \in \mathcal{I}_1/B_1} \partial_t \omega_I \otimes \theta_I^t(v) &= -\frac{1}{2} \sum_{I \in \mathcal{I}_1} d_Q d_I \otimes \theta_I^p(v) + \frac{1}{2} \sum_t \partial_t x_p d_Q \sum_I d_I \otimes \theta_I^t(v) \\
&= -\frac{1}{2} \sum_I d_Q d_I \otimes \theta_I^p(v) + \sum_t \partial_t \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^t(v)) \\
&= \frac{1}{2} \sum_I (\omega_{I,Q} - \frac{1}{2} \varepsilon_{Q,I} \partial_{t_{Q,I}}) \otimes \theta_I^p(v) + \sum_t \partial_t \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^t(v)) \\
&= \sum_{I \in \mathcal{I}_1/B_1} (\omega_{I,Q} \otimes \theta_I^p(v) - \partial_{t_{Q,I}} \otimes \frac{1}{2} \varepsilon_{Q,I} \theta_I^p(v)) + \sum_t \partial_t \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^t(v)).
\end{aligned}$$

Now we can use these two relations and compute

$$\begin{aligned}
x_p d_Q \Phi(v) &= x_p d_Q \left(\sum_{(I,J,K) \in \mathcal{I}_3/B_3} \omega_{I,J,K} \otimes \theta_{I,J,K}(v) + \sum_{t \in [5]} \sum_{I \in \mathcal{I}_1/B_1} \partial_t \omega_I \otimes \theta_I^t(v) \right) \\
&= \sum_{(H,L) \in \mathcal{I}_2/B_2} \omega_{H,L} \otimes \frac{1}{2} \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(- (x_p \partial_\gamma \cdot \theta_{\alpha\beta,H,L}(v)) + 2x_p \partial_\gamma \cdot (\theta_{\alpha\beta,H,L}(v)) \right) \\
&\quad + \partial_q \otimes -\frac{1}{2} \theta_{ab,bc,ca}(v) + \sum_{\alpha\beta\gamma} \partial_\alpha \otimes \frac{1}{4} (\theta_{\alpha\beta,\beta\gamma,\gamma q}(v) + \theta_{\alpha\gamma,\gamma\beta,\beta q}(v)) \\
&\quad + \sum_{I \in \mathcal{I}_1/B_1} (\omega_{I,Q} \otimes \theta_I^p(v) - \partial_{t_{Q,I}} \otimes \frac{1}{2} \varepsilon_{Q,I} \theta_I^p(v)) + \sum_t \partial_t \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^t(v)) \\
&= \sum_{(H,L) \in \mathcal{I}_2/B_2} \omega_{H,L} \otimes \left(\chi_{L \in B_1 Q} \theta_H^p(v) + \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(-\frac{1}{2} (x_p \partial_\gamma \cdot \theta_{\alpha\beta,H,L}(v)) + x_p \partial_\gamma \cdot (\theta_{\alpha\beta,H,L}(v)) \right) \right) \\
&\quad + \partial_p \otimes \varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^p(v)) + \partial_q \otimes \left(\varepsilon_{pqabc} \sum_{\alpha\beta\gamma} x_p \partial_\gamma \cdot (\theta_{\alpha\beta}^q(v)) - \frac{1}{2} \theta_{ab,bc,ca}(v) \right) \\
&\quad + \sum_{\alpha\beta\gamma} \partial_\alpha \otimes \left(\frac{1}{4} \theta_{\alpha\beta,\beta\gamma,\gamma q}(v) + \frac{1}{4} \theta_{\alpha\gamma,\gamma\beta,\beta q}(v) + \varepsilon_{pqabc} \left(-\frac{1}{2} \theta_{\beta\gamma}^p(v) + x_p \partial_c \cdot (\theta_{ab}^\alpha(v)) \right. \right. \\
&\quad \left. \left. + x_p \partial_b \cdot (\theta_{ca}^\alpha(v)) + x_p \partial_a \cdot (\theta_{bc}^\alpha(v)) \right) \right).
\end{aligned}$$

This completes the proof of Equations (4), (6) and (7). In order to deduce Equation (5) we consider the coefficient of ∂_a in the previous equation (the coefficients of ∂_b and ∂_c provide

equivalent conditions) and we have

$$\begin{aligned}
& \frac{1}{4}\theta_{ab,bc,cq}(v) + \frac{1}{4}\theta_{ac,cb,bq}(v) + \varepsilon_{pqabc} \left(-\frac{1}{2}\theta_{bc}^p(v) + x_p\partial_c.(\theta_{ab}^a(v)) + x_p\partial_b.(\theta_{ca}^a(v)) + x_p\partial_a.(\theta_{bc}^a(v)) \right) \\
&= \frac{1}{4}\theta_{ab,bc,cq}(v) + \frac{1}{4}\theta_{ac,cb,bq}(v) + \varepsilon_{pqabc} \left(-\frac{1}{2}((x_p\partial_a.\theta_{bc}^a(v)) + (x_p\partial_b.\theta_{ca}^a(v)) + (x_p\partial_c.\theta_{ab}^a(v))) \right. \\
&\quad \left. + x_p\partial_c.(\theta_{ab}^a(v)) + x_p\partial_b.(\theta_{ca}^a(v)) + x_p\partial_a.(\theta_{bc}^a(v)) \right) \\
&= \frac{1}{4}\theta_{ab,bc,cq}(v) + \frac{1}{4}\theta_{ac,cb,bq}(v) + \frac{1}{2}\varepsilon_{pqabc} \sum_{\alpha\beta\gamma} \left(- (x_p\partial_\gamma.\theta_{\alpha\beta}^a(v)) + 2x_p\partial_\gamma.(\theta_{\alpha\beta}^a(v)) \right).
\end{aligned}$$

□

Corollary 10.3. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of Verma modules of degree 3 associated to*

$$\Phi = \sum_{I \in \mathcal{I}_3/B_3} \omega_I \otimes \theta_I + \sum_{t \in [5]} \sum_{I \in \mathcal{I}_1/B_1} \partial_t \omega_I \otimes \theta_I^t.$$

Then the linear map $\psi : M(\mu^) \rightarrow M(\lambda^*)$ associated to*

$$\Psi = \sum_{I \in \mathcal{I}_3/B_3} \omega_I \otimes \theta_I^* + \sum_{t \in [5]} \sum_{I \in \mathcal{I}_1/B_1} \partial_t \omega_I \otimes (-\theta_I^t)^*$$

is also a morphism of Verma modules.

Proof. This is an immediate consequence of Remark 7.2 and Theorem 10.2. □

If we consider Equation (4) on a highest weight vector $s \in F(\lambda)$ (and we multiply it by $2\varepsilon_{pqabc}$) we obtain the following equation:

$$(8) \quad 2\varepsilon_{pqabc}\chi_{L \in B_1 Q}\theta_H^p(s) + \sum_{\alpha\beta\gamma} ((-1)^{\chi_{p>\gamma}}(x_p\partial_\gamma.\theta_{\alpha\beta,H,L}(s)) + 2\chi_{p>\gamma}x_p\partial_\gamma.(\theta_{\alpha\beta,H,L}(s))) = 0.$$

Remark 10.4. If $x_p\partial_c.\theta_{ab,H,L}$ has the leading weight of φ then $\chi_{p>\gamma}x_p\partial_\gamma.(\theta_{\alpha\beta,H,L}(s)) = 0$ for all $(\alpha, \beta, \gamma) \in C(a, b, c)$ and so we obtain the following

$$(9) \quad 2\varepsilon_{pqabc}\chi_{L \in B_1 Q}\theta_H^p(s) + \sum_{\alpha\beta\gamma} (-1)^{\chi_{p>\gamma}}(x_p\partial_\gamma.\theta_{\alpha\beta,H,L}(s)) = 0.$$

This equation has several immediate consequences.

Lemma 10.5. *If $a, b, c, d \in [5]$ are distinct then $\theta_{ab,ac,ad}$ does not have the leading weight of φ .*

Proof. Without loss of generality we can assume that the fifth element p is either bigger than both b and c or smaller than both b and c . Otherwise we can rename b, c, d accordingly. Remark 10.4 applies with $H = (a, p)$, $q = d$ and $L = (a, d)$ so we have

$$(-1)^{\chi_{p>c}}\theta_{ab,ac,ad}(s) + (-1)^{\chi_{p>b}}\theta_{ab,ac,ad}(s) = 0.$$

□

Lemma 10.6. *If $a, b, c \in [5]$ are distinct then $\theta_{ab,bc,ca}$ does not have the leading weight of φ .*

Proof. Without loss of generality we can choose p such that p is either bigger than both a and c or smaller than both a and c . Remark 10.4 applies with $H = (b, p)$ and $L = (c, a)$ so we have

$$(-1)^{\chi_{p>c}} \theta_{ab,bc,ca}(s) + (-1)^{\chi_{p>a}} \theta_{ab,bc,ca}(s) = 0.$$

□

Lemma 10.7. *If $x, y, z, w \in [5]$ are distinct and $\theta_{xy,zw,xw}$ has the leading weight of φ , then $\theta_{xy,zw,xw} = \theta_{12,45,kl}$ for some $k, l \in \{1, 2, 4, 5\}$.*

Proof. Let us first assume that $\{x, y, z, w\} \neq \{1, 2, 4, 5\}$. This assumption ensures that we can assume that the fifth element p is either bigger or smaller than both y and w (otherwise exchange the roles of x, z and y, w). Use Remark 10.4 with $a = x$, $b = y$, $c = w$, $q = z$, $H = (x, p)$, $L = (z, w)$. Then we have:

$$(-1)^{\chi_{p>y}} \theta_{xy,xw,zw}(s) + (-1)^{\chi_{p>w}} \theta_{xy,xw,zw}(s) = 0.$$

Now let $\{x, y, z, w\} = \{1, 2, 4, 5\}$. If either $\{y, w\} = \{1, 2\}$ or $\{y, w\} = \{4, 5\}$ then we can use the same argument as above.

Now let $\{y, z\} = \{4, 5\}$ so that $\theta_{1y,2z,12}$ has the leading weight of φ . Equation (8) with $a = 1$, $b = 2$, $q = 3$, $c = y$, $p = z$, $H = (2, p)$ and $L = (1, 2)$ gives

$$x_z \partial_2.(\theta_{1y,2z,12}(s)) = 0$$

hence if we apply $x_2 \partial_z$ we get $h_{2z}.(\theta_{1y,2z,12}(s)) = 0$ which implies in particular that $\mu_{34} = 0$. Since $\lambda_{34}(\theta_{1y,2z,12}) = 1$ this contradicts the dominance of λ . The thesis follows. □

Lemma 10.8. *The elements $\theta_{12,45,14}$, $\theta_{12,45,25}$ and $\theta_{12,45,24}$ do not have the leading weight of φ .*

Proof. Use Equation (8) with $a = 1$, $b = 2$, $c = 4$, $q = 3$ and $p = 5$, $H = (4, 5)$ and $L = (1, 2)$. We obtain

$$(10) \quad \theta_{24,41,12}(s) + \theta_{41,42,12}(s) + 2x_5 \partial_1.(\theta_{24,45,12}(s)) + 2x_5 \partial_2.(\theta_{41,45,12}(s)) = 0.$$

Assume $\theta_{12,45,14}$ has the leading weight of φ . Then $\theta_{24,45,12}(s) = 0$ and we apply $x_2 \partial_5$ to Equation (10) to obtain

$$-\theta_{54,41,12}(s) - \theta_{24,41,15}(s) - \theta_{41,45,12}(s) - \theta_{41,42,15}(s) + 2h_{25}.(\theta_{41,45,12}(s)) = 0.$$

But by Lemma 10.7 we have $\theta_{24,41,15}(s) = 0$ and so we have

$$-2\theta_{41,45,12}(s) + 2h_{25}.(\theta_{41,45,12}(s)) = 0.$$

It follows that $\lambda_{25}(\theta_{41,45,12}(s)) = 1$ and so $\lambda_{34}(\theta_{41,45,12}(s)) \leq 1$ and, since $\lambda_{34}(\theta_{41,45,12}) = 2$ this would imply $\lambda_{34}(s) \leq -1$, a contradiction.

By Corollary 10.3 the element $\theta_{12,45,25}$ does not have the leading weight of φ since $\lambda(\theta_{12,45,25}) = -\lambda(\theta_{12,45,14})^*$.

Now we assume that $\theta_{12,45,24}$ has the leading weight of φ . We apply $x_1 \partial_5$ to Equation (10) to obtain

$$-\theta_{24,45,12}(s) - \theta_{24,41,52}(s) - \theta_{45,42,12} - \theta_{41,42,52} + 2h_{15}.(\theta_{24,45,12}(s)) + 2x_1 \partial_2.(\theta_{41,45,12}(s)) = 0.$$

Lemma 10.7 ensures $\theta_{24,41,52}(s) = 0$ and so we obtain

$$-2\theta_{24,45,12}(s) + 2h_{15}.(\theta_{24,45,12}(s) - 2\theta_{42,45,12}(s)) = 0$$

and we conclude

$$h_{15} \cdot (\theta_{24,45,12}(s)) = 0.$$

We obtain a contradiction with the same argument used in the other case. \square

Lemma 10.9. *Assume that $\theta_{12,15,45}$ has the leading weight of φ . Then $\lambda = (1, 1, 0, 0)$, $\mu = (0, 0, 1, 1)$ and $\varphi = \nabla_C \nabla_B \nabla_A$ (up to a scalar).*

Proof. Use Equation (8) with $a = 1$, $b = 2$, $c = 4$, $q = 3$, $p = 5$, $H = (1, 5)$ and $L = (4, 5)$. We obtain

$$\theta_{12,14,45}(s) + \theta_{24,15,41}(s) + \theta_{41,12,45}(s) + \theta_{41,15,42}(s) + 2x_5 \partial_4 \cdot (\theta_{12,15,45}(s)) = 0$$

since $\theta_{24,14,45}(s) = \theta_{41,15,45}(s) = 0$. Applying $x_4 \partial_5$ we get

$$-\theta_{12,15,45}(s) - \theta_{25,15,41}(s) - \theta_{51,12,45}(s) - \theta_{41,15,52}(s) + 2h_{45} \cdot (\theta_{12,15,45}(s)) = 0.$$

By Lemma 10.7 we have $\theta_{25,15,41}(s) = 0$ and so we obtain

$$-2\theta_{12,15,45}(s) + 2h_{45} \cdot (\theta_{12,15,45}(s)) = 0$$

and so

$$\lambda_{45}(\theta_{12,15,45}(s)) = 1.$$

Now we consider Equation (8) with $a = 1$, $b = 3$, $c = 5$, $q = 2$, $p = 4$, $H = (1, 2)$ and $L = (4, 5)$. We obtain

$$\theta_{35,12,15}(s) + \theta_{51,12,35}(s) + 2x_4 \partial_3 \cdot (\theta_{51,12,45}(s)) = 0.$$

Applying $x_3 \partial_4$ to this equation we have

$$-\theta_{45,12,15}(s) - \theta_{51,12,45}(s) + 2h_{34} \cdot (\theta_{51,12,45}(s)) = 0$$

and from this we get $\lambda_{34}(\theta_{12,15,45}(s)) = 1$.

Finally, we use again Equation (8) with $a = 1$, $b = 4$, $c = 5$, $q = 2$, $p = 3$, $H = (1, 2)$, $L = (1, 5)$ which gives $2x_3 \partial_1 \cdot (\theta_{45,12,15}(s)) = 0$, hence

$$\lambda_{13}(\theta_{12,15,45}(s)) = 0$$

proving that $\mu = (0, 0, 1, 1)$. It follows that $\lambda = (1, 1, 0, 0)$ since $\lambda(\theta_{12,15,45}) = (-1, -1, 1, 1)$.

By Remark 10.4 we have $-2\theta_{15}^3(s) - \theta_{12,15,45}(s) = 0$ hence the leading term of the singular vector $\varphi(1 \otimes s)$ is $\omega_{12,15,45} \otimes \theta_{12,15,45}(s) + \partial_3 d_{15} \otimes \theta_{15}^3(s) = d_{12} d_{15} d_{45} \otimes \theta_{12,15,45}(s)$. It follows that $\varphi = \nabla_C \nabla_B \nabla_A$ due to Proposition 3.8. \square

In the next result, for notational convenience, for all $a, b \in [5]$ we let $(-1)^{a < b} = (-1)^{\chi_{a < b}}$.

Proposition 10.10. *Let $\{x, y, z, w, t\} = [5]$ and let s be a highest weight vector in $F(\lambda)$. Assume that $\theta_{xy,xz,wt}$ has the leading weight of φ . Then the following equations hold:*

$$(11) \quad -2\varepsilon_{xyzwt} \theta_{xy}^y(s) + (-1)^{y < t} \theta_{xz,xt,yw}(s) + (-1)^{y < t} \theta_{xz,xy,tw}(s) + (-1)^{y < z} \theta_{xz,xt,yw}(s) \\ + (-1)^{y < z} \theta_{xy,xt,zw}(s) + (-1)^{y < x} \theta_{xy,xw,zt}(s) = 0$$

$$(12) \quad 2\varepsilon_{xyzwt} \theta_{xy}^y(s) + (-1)^{y < t} \theta_{xw,xt,yz}(s) + (-1)^{y < t} \theta_{xw,xy,tz}(s) + (-1)^{y < w} \theta_{xw,xt,yz}(s) \\ + (-1)^{y < w} \theta_{xy,xt,wz}(s) + (-1)^{y < x} \theta_{xy,xz,wt}(s) = 0$$

$$(13) \quad -2\varepsilon_{xyzwt}\theta_{xy}^y(s) + (-1)^{y<z}\theta_{xw,xz,yt}(s) + (-1)^{y<z}\theta_{xw,xy,zt}(s) + (-1)^{y<w}\theta_{xw,xz,yt}(s) \\ + (-1)^{y<w}\theta_{xy,xz,wt}(s) + (-1)^{y<x}\theta_{xy,xt,wz}(s) = 0$$

$$(14) \quad 2\varepsilon_{xyzwt}\theta_{xz}^z(s) + (-1)^{z<t}\theta_{xy,xt,zw}(s) + (-1)^{z<t}\theta_{xy,xz,tw}(s) + (-1)^{z<y}\theta_{xy,xt,zw}(s) \\ + (-1)^{z<y}\theta_{xz,xt,yw}(s) + (-1)^{z<x}\theta_{xz,xw,yt}(s) = 0$$

$$(15) \quad -2\varepsilon_{xyzwt}\theta_{xz}^z(s) + (-1)^{z<t}\theta_{xw,xt,zy}(s) + (-1)^{z<t}\theta_{xw,xz,ty}(s) + (-1)^{z<w}\theta_{xw,xt,zy}(s) \\ + (-1)^{z<w}\theta_{xz,xt,wy}(s) + (-1)^{z<x}\theta_{xz,xy,wt}(s) = 0$$

$$(16) \quad 2\varepsilon_{xyzwt}\theta_{xz}^z(s) + (-1)^{z<y}\theta_{xw,xy,zt}(s) + (-1)^{z<y}\theta_{xw,xz,yt}(s) + (-1)^{z<w}\theta_{xw,xy,zt}(s) \\ + (-1)^{z<w}\theta_{xz,xy,wt}(s) + (-1)^{z<x}\theta_{xz,xt,wy}(s) = 0$$

$$(17) \quad 2\varepsilon_{xyzwt}\theta_{xt}^t(s) + (-1)^{t<y}\theta_{xz,xy,tw}(s) + (-1)^{t<y}\theta_{xz,xt,yw}(s) + (-1)^{t<z}\theta_{xz,xy,tw}(s) \\ + (-1)^{t<z}\theta_{xt,xy,zw}(s) + (-1)^{t<x}\theta_{xt,xw,zy}(s) = 0$$

$$(18) \quad -2\varepsilon_{xyzwt}\theta_{xt}^t(s) + (-1)^{t<w}\theta_{xz,xw,ty}(s) + (-1)^{t<w}\theta_{xz,xt,wy}(s) + (-1)^{t<z}\theta_{xz,xw,ty}(s) \\ + (-1)^{t<z}\theta_{xt,xw,zy}(s) + (-1)^{t<x}\theta_{xt,xy,zw}(s) = 0$$

$$(19) \quad 2\varepsilon_{xyzwt}\theta_{xt}^t(s) + (-1)^{t<w}\theta_{xy,xw,tz}(s) + (-1)^{t<w}\theta_{xy,xt,wz}(s) + (-1)^{t<y}\theta_{xy,xw,tz}(s) \\ + (-1)^{t<y}\theta_{xt,xw,yz}(s) + (-1)^{t<x}\theta_{xt,xz,yw}(s) = 0$$

$$(20) \quad -2\varepsilon_{xyzwt}\theta_{xw}^w(s) + (-1)^{w<t}\theta_{xy,xt,wz}(s) + (-1)^{w<t}\theta_{xy,xw,tz}(s) + (-1)^{w<y}\theta_{xy,xt,wz}(s) \\ + (-1)^{w<y}\theta_{xw,xt,yz}(s) + (-1)^{w<x}\theta_{xw,xz,yt}(s) = 0$$

$$(21) \quad 2\varepsilon_{xyzwt}\theta_{xw}^w(s) + (-1)^{w<t}\theta_{xz,xt,wy}(s) + (-1)^{w<t}\theta_{xz,xw,ty}(s) + (-1)^{w<z}\theta_{xz,xt,wy}(s) \\ + (-1)^{w<z}\theta_{xw,xt,zy}(s) + (-1)^{w<x}\theta_{xw,xy,zt}(s) = 0$$

$$(22) \quad -2\varepsilon_{xyzwt}\theta_{xw}^w(s) + (-1)^{w<y}\theta_{xz,xy,wt}(s) + (-1)^{w<y}\theta_{xz,xw,yt}(s) + (-1)^{w<z}\theta_{xz,xy,wt}(s) \\ + (-1)^{w<z}\theta_{xw,xy,zt}(s) + (-1)^{w<x}\theta_{xw,xt,zy}(s) = 0$$

Proof. We use Remark 10.4 twelve times with $L = Q = (p, q)$ any ordered pair in $\{y, z, w, t\}$ and $H = (x, p)$ to obtain the stated equations. More precisely we get Equation (11) with $p = y, q = w$; Equation (12) with $p = y, q = z$; Equation (13) with $p = y, q = t$; Equation (14) with $p = z, q = w$; Equation (15) with $p = z, q = y$; Equation (16) with $p = z, q = t$; Equation (17) with $p = t, q = w$; Equation (18) with $p = t, q = y$; Equation (19) with $p = t, q = z$; Equation (20) with $p = w, q = z$; Equation (21) with $p = w, q = y$; Equation (22) with $p = w, q = t$. \square

Proposition 10.10 provides 12 linear equations in the ten unknown $\theta_{xy,xz,wt}(s) = f_{wt}$, $\theta_{xy,xw,zt}(s) = f_{zt}$, $\theta_{xy,xt,zw}(s) = f_{zw}$, $\theta_{xz,xw,yt}(s) = f_{yt}$, $\theta_{xz,xt,yw}(s) = f_{yw}$, $\theta_{xw,xt,yz}(s) = f_{yz}$, $\varepsilon_{xyzwt}\theta_{xy}^y(s) = b_y$, $\varepsilon_{xyzwt}\theta_{xz}^z(s) = b_z$, $\varepsilon_{xyzwt}\theta_{xw}^w(s) = b_w$, $\varepsilon_{xyzwt}\theta_{xt}^t(s) = b_t$. We are now interested in the study of the weights $\lambda_{i,j}(\theta_{xy,xz,wt}(s))$.

Proposition 10.11. *Let $\{p, q, a, b, c\} = [5]$ with $c < p$, let s be a highest weight vector in $F(\lambda)$, $H, L \in \mathcal{I}_1$ and assume that $\theta_{ab,H,L}$ has the leading weight of φ . Then we have*

$$\begin{aligned} 2h_{cp} \cdot (\theta_{ab,H,L}(s)) = \\ - 2\varepsilon_{pqabc} \chi_{L \in B_1 Q} (x_c \partial_p \cdot \theta_H^p)(s) + (x_c \partial_p \cdot (x_p \partial_c \theta_{ab,H,L}))(s) + (-1)^{\chi_{p < b}} (x_c \partial_p \cdot (x_p \partial_b \cdot \theta_{ca,H,L}))(s) \\ + (-1)^{\chi_{p < a}} (x_c \partial_p \cdot (x_p \partial_a \cdot \theta_{bc,H,L}))(s) - 2\chi_{c < b < p} (x_c \partial_b \cdot \theta_{ca,H,L})(s) - 2\chi_{c < a < p} (x_c \partial_a \cdot \theta_{bc,H,L})(s) \end{aligned}$$

Proof. Equation (8) is equivalent to the following

$$\begin{aligned} 2\varepsilon_{pqabc} \chi_{L \in B_1 Q} \theta_H^p(s) - (x_p \partial_c \cdot \theta_{ab,H,L})(s) + (-1)^{\chi_{p > b}} (x_p \partial_b \cdot \theta_{ca,H,L})(s) + (-1)^{\chi_{p > a}} (x_p \partial_a \cdot \theta_{bc,H,L})(s) \\ + 2x_p \partial_c \cdot (\theta_{ab,H,L}(s)) + 2\chi_{p > b} x_p \partial_b \cdot (\theta_{ca,H,L}(s)) + 2\chi_{p > a} x_p \partial_a \cdot (\theta_{bc,H,L}(s)) = 0. \end{aligned}$$

We apply $x_c \partial_p$ to this equation and we obtain

$$\begin{aligned} 2\varepsilon_{pqabc} \chi_{L \in B_1 Q} x_c \partial_p \cdot (\theta_H^p(s)) - (x_c \partial_p \cdot (x_p \partial_c \cdot \theta_{ab,H,L}))(s) + (-1)^{\chi_{p > b}} (x_c \partial_p \cdot (x_p \partial_b \cdot \theta_{ca,H,L}))(s) \\ + (-1)^{\chi_{p > a}} (x_c \partial_p \cdot (x_p \partial_a \cdot \theta_{bc,H,L}))(s) + 2h_{cp} \cdot (\theta_{ab,H,L}(s)) + 2\chi_{c < b < p} (x_c \partial_b \cdot \theta_{ca,H,L})(s) \\ + 2\chi_{c < a < p} (x_c \partial_a \cdot \theta_{bc,H,L})(s) = 0. \end{aligned}$$

The result follows. \square

Corollary 10.12. *Let $\{x, y, z, w, t\} = [5]$ and assume that $\theta_{xy,xz,wt}$ has the leading weight of φ . Then we have*

if $z < w$,

$$(23) \quad 2h_{zw} \cdot f_{wt} = 2(b_w - b_z) + f_{zt} + f_{wt} + (-1)^{\chi_{w < x}} (f_{yw} + f_{yz}) - 2\chi_{z < x < w} f_{wt};$$

if $y < z$,

$$(24) \quad \begin{aligned} 2h_{yz} \cdot f_{wt} = 2(b_y - b_z) + (-1)^{\chi_{t < z}} (-f_{zw} - f_{yw}) + (-1)^{\chi_{w < z}} (f_{zt} + f_{yt}) \\ - 2\chi_{y < t < z} (f_{wt} + f_{yw}) - 2\chi_{y < w < z} (f_{wt} - f_{yt}) \end{aligned}$$

if $w < t$,

$$(25) \quad \begin{aligned} 2h_{wt} \cdot f_{wt} = (-1)^{\chi_{y < t}} (f_{yw} + f_{yt}) + (-1)^{\chi_{x < t}} (f_{yt} + f_{yw}) \\ - 2\chi_{w < y < t} (f_{wt} - f_{yt}) - 2\chi_{w < x < t} (f_{wt} - f_{yw}). \end{aligned}$$

if $w < z$,

$$(26) \quad 2h_{wz} \cdot f_{wt} = f_{wt} + f_{zt} + (-1)^{\chi_{z < y}} (f_{wt} + f_{zt}) + 2\chi_{w < y < z} (-f_{wt} + f_{yt}) + 2\chi_{w < x < z} (-f_{wt} + f_{yw})$$

if $x < y$,

$$(27) \quad \begin{aligned} 2h_{xy} \cdot f_{wt} = ((-1)^{\chi_{y < t}} + (-1)^{\chi_{y < w}}) (-f_{yw} + f_{yt} + f_{yz}) \\ - 2\chi_{x < t < y} (f_{wt} + f_{yt} - f_{zt}) - 2\chi_{x < w < y} (f_{wt} + f_{yw} - f_{yz}) \end{aligned}$$

Proof. The statement follows from Proposition 10.11 with the following choices:

- (1) $a = x, b = y, c = z, p = w, q = t, H = (x, z), L = (w, t).$
- (2) $c = y, p = z, a = w, b = t, q = x, H = (x, y), L = (z, x).$
- (3) $c = w, p = t, a = x, b = y, q = z, H = (x, z), L = (w, t).$
- (4) $c = w, p = z, a = x, b = y, q = t, H = (x, z), L = (w, t).$
- (5) $c = x, p = y, a = w, b = t, q = z, H = (x, y), L = (x, z).$

\square

Proposition 10.13. *Let s be a highest weight vector in $F(\lambda)$. For $c < p$ we have*

$$4h_{cp}(\theta_{ab}^a(s)) = (-4\chi_{c<b<p} - 4\chi_{c<a<p})\theta_{ab}^a(s) + (2 - 4\chi_{c<a})\theta_{bc}^c(s) + (-2 + 4\chi_{p<a})\theta_{bp}^p(s) \\ + \varepsilon_{pqabc}(\theta_{ab,bp,cq}(s) + \theta_{ab,bc,pq}(s) + \theta_{ap,cb,bq}(s) + \theta_{ac,pb,bq}(s))$$

Proof. We start from Equation (5):

$$(28) \quad \theta_{ab,bc,cq}(s) + \theta_{ac,cb,bq}(s) + \varepsilon_{pqabc}(-2\theta_{bc}^p(s) + 4x_p\partial_{c\cdot}(\theta_{ab}^a(s)) + 4x_p\partial_{b\cdot}(\theta_{ca}^a(s)) + 4x_p\partial_{a\cdot}(\theta_{bc}^a(s))) = 0.$$

We want to apply $x_c\partial_p$ to this equation and so we do the following two preliminary calculations:

$$\begin{aligned} x_c\partial_p(x_p\partial_{b\cdot}(\theta_{ca}^a(s))) &= \chi_{c<b}x_c\partial_p(x_p\partial_{b\cdot}(\theta_{ca}^a(s))) \\ &= \chi_{c<b}x_c\partial_{b\cdot}(\theta_{ca}^a(s)) + \chi_{c<b}x_p\partial_{b\cdot}(x_c\partial_p(\theta_{ca}^a(s))) \\ &= -\chi_{c<b}\theta_{ba}^a(s) - \chi_{c<b}x_p\partial_{b\cdot}(\theta_{pa}^a(s)) \\ &= \chi_{c<b}\theta_{ab}^a(s) + \chi_{c<b}\chi_{p<b}\theta_{ba}^a(s) \\ &= \chi_{c<b}(1 - \chi_{p<b})\theta_{ab}^a(s) \\ &= \chi_{c<b<p}\theta_{ab}^a(s) \end{aligned}$$

$$\begin{aligned} x_c\partial_p(x_p\partial_{a\cdot}(\theta_{bc}^a(s))) &= \chi_{c<a}x_c\partial_p(x_p\partial_{a\cdot}(\theta_{bc}^a(s))) \\ &= \chi_{c<a}x_c\partial_{a\cdot}(\theta_{bc}^a(s)) + \chi_{c<a}x_p\partial_{a\cdot}(x_c\partial_p(\theta_{bc}^a(s))) \\ &= \chi_{c<a}(\theta_{bc}^c(s) - \theta_{ba}^a(s)) - \chi_{c<a}x_p\partial_{a\cdot}(\theta_{bp}^a(s)) \\ &= \chi_{c<a}\theta_{bc}^c(s) + \chi_{c<a}\theta_{ab}^a(s) - \chi_{c<a}\chi_{p<a}(\theta_{bp}^p(s) - \theta_{ba}^a(s)) \\ &= \chi_{c<a}\theta_{bc}^c(s) - \chi_{p<a}\theta_{bp}^p(s) + \chi_{c<a<p}\theta_{ab}^a(s) \end{aligned}$$

Therefore, if we apply $x_c\partial_p$ to Equation (28), using the previous computations, we obtain

$$\begin{aligned} -\theta_{ab,bp,cq}(s) - \theta_{ab,bc,pq}(s) - \theta_{ap,cb,bq}(s) - \theta_{ac,pb,bq}(s) + \varepsilon_{pqabc}(-2\theta_{bc}^c(s) + 2\theta_{bp}^p(s) \\ + 4h_{cp}(\theta_{ab}^a(s)) + 4\chi_{c<b<p}\theta_{ab}^a(s) + 4\chi_{c<a}\theta_{bc}^c(s) - 4\chi_{p<a}\theta_{bp}^p(s) + 4\chi_{c<a<p}\theta_{ab}^a(s)) = 0 \end{aligned}$$

hence we get the statement. \square

Proposition 10.14. *Let $\{h, k, l, m, n\} = [5]$. Then $\theta_{hk,hl,mn}$ and θ_{hk}^k do not have the leading weight of φ .*

Proof. We first assume $h = 1$ and we let $x = 1, y = 2, z = 3, w = 4, t = 5$. We use notation introduced after the proof of Proposition 10.10 and we observe that, up to a sign, $\theta_{1k,1l,mn}(s) \in \{f_{23}, f_{24}, f_{25}, f_{34}, f_{35}, f_{45}\}$ and $\theta_{1k}^k(s) \in \{b_2, b_3, b_4, b_5\}$. We solve the linear system provided by Proposition 10.10 and we have:

- $f_{35} = -f_{45} = -f_{34}$
- $f_{24} = -f_{25} = -f_{23}$
- $2b_2 = -3f_{34} + 2f_{23}$
- $2b_3 = 2b_5 = 2b_4 = -f_{34}$.

We use Proposition 10.13 with $a = 4$, $b = 1$, $c = 2$, $p = 3$, $q = 5$ and we obtain

$$\begin{aligned} h_{23}.b_4 &= \frac{1}{2}b_2 - \frac{1}{2}b_3 + \frac{1}{4}(f_{25} + f_{35} + f_{34} + f_{24}) \\ &= \frac{1}{4}(-3f_{34} + 2f_{23}) + \frac{1}{4}f_{34} + \frac{1}{4}(f_{23} - f_{34} + f_{34} - f_{23}) \\ &= -\frac{1}{2}f_{34} + \frac{1}{2}f_{23} \end{aligned}$$

therefore

$$h_{23}.f_{34} = f_{34} - f_{23}.$$

Now we use Equation (24):

$$2h_{23}.f_{45} = 2(b_2 - b_3) - f_{34} - f_{24} + (f_{35} + f_{25})$$

i.e.

$$2h_{23}.f_{34} = -3f_{34} + 2f_{23} + f_{34} - f_{34} + f_{23} - f_{34} + f_{23} = -4f_{34} + 4f_{23}$$

or

$$h_{23}.f_{34} = -2f_{34} + 2f_{23}.$$

Comparing this with the previous equation we obtain $f_{34} = f_{23}$.

Now we use Equation (27):

$$2h_{12}.f_{45} = 2f_{24} - 2f_{25} - 2f_{23}$$

i.e.

$$2h_{12}.f_{34} = -2f_{23} - 2f_{23} - 2f_{23} = -6f_{34}$$

This implies that $f_{34} = f_{23} = 0$. It follows that $\theta_{1k,1l,mn}(s) = 0$ and $\theta_{1k}^k(s) = 0$.

Now let $h = 2$ and $x = 2$, $y = 1$, $z = 3$, $w = 4$, $t = 5$. Similarly as above we have, up to a sign, $\theta_{2k,2l,mn}(s) \in \{f_{13}, f_{14}, f_{15}, f_{34}, f_{35}, f_{45}\}$ and $\theta_{2k}^k(s) \in \{b_1, b_3, b_4, b_5\}$. We solve the linear system provided by Proposition 10.10 and we have:

- $f_{35} = -f_{45} = -f_{34}$
- $f_{14} = -f_{15} = -f_{13}$
- $2b_1 = -f_{34} + 2f_{13}$
- $2b_3 = 2b_4 = 2b_5 = -f_{34}$

We use Proposition 10.13 with $a = 4$, $b = 2$, $c = 1$, $p = 5$, $q = 3$ and we obtain:

$$h_{15}.b_4 = \frac{1}{2}f_{34} + \frac{1}{2}f_{13}$$

i.e.,

$$h_{15}.f_{34} = -f_{34} - f_{13}$$

Now we use Equations (23), (24), (25) and we obtain:

$$h_{15}.f_{34} = 2f_{13} - f_{34}$$

It follows that:

$$2f_{13} - f_{34} = -f_{34} - f_{13}$$

i.e., $f_{13} = 0$, hence $h_{15}.f_{34} = -f_{34}$ which implies $f_{34} = 0$. It follows that $\theta_{2k,2l,mn}(s) = 0$ and $\theta_{2k}^k = 0$.

Now let $h = 3$ and $x = 3, y = 1, z = 2, w = 4, t = 5$. Similarly as above we have, up to a sign, $\theta_{3k,3l,mn}(s) \in \{f_{12}, f_{14}, f_{15}, f_{24}, f_{25}, f_{45}\}$ and $\theta_{3k}^k(s) \in \{b_1, b_2, b_4, b_5\}$. We solve the linear system provided by Proposition 10.10 and we have:

- $f_{15} = f_{24} = -f_{25} = -f_{14}$
- $f_{45} = -2b_4 = -2b_5 = -2f_{14} - f_{12}$
- $2b_1 = 2b_2 = f_{12}$

We use Proposition 10.13 with $a = 2, b = 3, c = 1, p = 5, q = 4$ and we obtain:

$$-h_{15}(b_2) = \frac{1}{2}f_{12} - \frac{1}{2}f_{14}$$

i.e.,

$$h_{15}(f_{12}) = f_{14} - f_{12}$$

Now we use Equations (23), (24), (25) and we obtain:

$$h_{15}(f_{45}) = 3f_{14} + f_{12}.$$

It follows that:

$$h_{15}.f_{14} = -\frac{1}{2}h_{15}.(f_{45} + f_{12}) = -2f_{14}$$

hence $f_{14} = 0$ and $h_{15}.f_{12} = -f_{12}$ from which it follows that $f_{12} = 0$. We conclude that $\theta_{3k,3l,mn}(s) = 0$ and $\theta_{3k}^k(s) = 0$.

If $h = 4, 5$ the result follows from Corollary 10.3. □

Now we can summarize the classification of morphisms of degree 3 in the next result.

Theorem 10.15. *Let $\varphi : M(\lambda) \rightarrow M(\mu)$ be a morphism of degree 3. Then $\lambda = (1, 1, 0, 0)$, $\mu = (0, 0, 1, 1)$ and up to a scalar $\varphi = \nabla_C \nabla_B \nabla_C$.*

Proof. This follows from Lemmas 10.5, 10.6, 10.7, 10.8, 10.9 and Proposition 10.14. □

REFERENCES

- [1] S.-J. CHENG, V. G. KAC *A new $N = 6$ superconformal algebra*, Comm. Math. Phys. **186** (1997), 219–231.
- [2] P. GROZMAN, D. LEITES, I. SHCHEPOCHKINA, *Invariant differential operators on supermanifolds and The Standard Model*, In: M. Olshanetsky, A. Vainstein (eds.) Multiple facets of quantization and supersymmetry. Michael Marinov Memorial Volume, World Sci. Publishing, River Edge, NJ (2002) 508–555;
- [3] V. G. KAC *Classification of infinite-dimensional simple linearly compact Lie superalgebras*, Adv. Math. **139** (1998), 1–55.
- [4] V. G. KAC, A. RUDAKOV *Representations of the exceptional Lie superalgebra $E(3,6)$. I. Degeneracy condition*, Transformation Groups **7** (2002), 67–86.
- [5] V. G. KAC, A. RUDAKOV *Representations of the exceptional Lie superalgebra $E(3,6)$. II. Four series of degenerate modules*, Comm. Math. Phys. **222** (2001), 611–661.
- [6] V. G. KAC, A. RUDAKOV *Complexes of modules over the exceptional Lie superalgebras $E(3,8)$ and $E(5,10)$* , Int. Math. Res. Not. **19** (2002), 1007–1025.
- [7] V. G. KAC, A. RUDAKOV *Representations of the exceptional Lie superalgebra $E(3,6)$. III. Classification of singular vectors*, J. Algebra Appl. **4** (2005), 15–57.
- [8] A. RUDAKOV *Morphisms of Verma modules over exceptional Lie superalgebra $E(5,10)$* , arXiv 1003.1369v1 (2010), 1–12.
- [9] I. SHCHEPOCHKINA *New exceptional simple Lie superalgebras*, C. R. Acad. Bulgare Sci. **36** (1983), no. 3, 313–314.
- [10] I. SHCHEPOCHKINA *The five exceptional simple Lie superalgebras of vector fields*, Funktsional Anal. i Prilozhen **33** (1999), **3**, 59–72, 96; transl. in Funct. Anal. Appl. **33** (1999), **3**, 208–219 (2000).

- [11] I. SHCHEPOCHKINA *The five exceptional simple Lie superalgebras of vector fields and their fourteen regradings*, Represent. Theory **3** (1999), 373–415.

FABRIZIO CASELLI AND NICOLETTA CANTARINI, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA,
PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY
Email address: `fabrizio.caselli@unibo.it`
Email address: `nicoletta.cantarini@unibo.it`