

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Imprints of volcanic, erosional, depositional, tectonic and mass-wasting processes in the morphology of Santa Maria insular shelf (Azores).

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Ricchi, A., Quartau, R., Ramalho, R.S., Romagnoli, C., Casalbore, D., Zhao, Z. (2020). Imprints of volcanic, erosional, depositional, tectonic and mass-wasting processes in the morphology of Santa Maria insular shelf (Azores). MARINE GEOLOGY, 424, 1-16 [10.1016/j.margeo.2020.106163].

Availability: This version is available at: https://hdl.handle.net/11585/752571 since: 2020-03-20

Published:

DOI: http://doi.org/10.1016/j.margeo.2020.106163

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Ricchi, A., Quartau, R., Ramalho, R. S., Romagnoli, C., Casalbore, D., & Zhao, Z. (2020). Imprints of volcanic, erosional, depositional, tectonic and mass-wasting processes in the morphology of Santa Maria insular shelf (Azores). Marine Geology, 424

The final published version is available online at https://dx.doi.org/10.1016/j.margeo.2020.106163

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<u>https://cris.unibo.it/</u>)

When citing, please refer to the published version.

# Title: Imprints of volcanic, erosional, depositional, tectonic and mass-wasting processes in the morphology of Santa Maria insular shelf.

Alessandro Ricchi<sup>a</sup>, Rui Quartau<sup>b,c</sup>, Ricardo S. Ramalho<sup>c,d,e,f</sup>, Claudia Romagnoli<sup>a</sup>, Daniele Casalbore<sup>g,h</sup>, Zhongwei Zhao<sup>i</sup>

<sup>a</sup>University of Bologna, Dip. Scienze Biologiche, Geologiche e Ambientali, Bologna, Italy <sup>b</sup>Instituto Hidrográfico, Divisão de Geologia Marinha, Lisboa, Portugal <sup>c</sup>Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal <sup>d</sup>Departamento de Geologia, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal <sup>e</sup>School of Earth Sciences, University of Bristol, Bristol, UK

<sup>f</sup>Lamont-Doherty Earth Observatory at Columbia University, Palisades, NY, USA <sup>g</sup>Istituto di Geologia Ambientale e Geoingegneria, Consiglio Nazionale delle Ricerche, Area della Ricerca, Roma, Italy

<sup>h</sup>Sapienza Università di Roma, Dipartimento Scienze della Terra, Roma, Italy <sup>i</sup>Department of Earth and Environmental Sciences, University of Manchester, Williamson Building, Oxford Road, Manchester M13 9PL, UK.

Corresponding author: Rui Quartau (rui.quartau@hidrografico.pt)

Highlights: (85 characters)

- The wider northern shelf suggests that the island might be older than 6 Ma
- The shelf is mostly sediment-deprived, except in fault-controlled basins
- Depositional sequences in island shelves are hard to preserve due to wave exposure
- Mass wasting at the shelf edge is driven by tectonics and sediment accumulation

- 1 Abstract
- 2

3 The present-day morphology of volcanic island shelves is the result of several factors, namely the 4 original shape and structure of the volcanic edifice, the nature and age of the materials that compose its flanks, and the erosional, volcanic, tectonic and sedimentary processes that acted to 5 6 form these shelves. By combining detailed seafloor mapping with onshore studies it is possible to 7 gain a unique insight onto the evolution of volcanic island edifices and the processes that helped 8 to shape their shelves. Here we look at Santa Maria Island in the Azores Archipelago, which is a 6 9 Ma old ocean-island volcano surrounded by a 135 km<sup>2</sup> shelf, characterized by an history of 10 subsidence since its formation c. 6 Ma ago, followed by uplift from 3.5 Ma onwards. By analyzing the geomorphic parameters of the insular shelf (such as the shelf width and the erosive edge 11 12 depth), the island's history of vertical movements, and by relating the morphology of the outcrops 13 found on the shelf with those onshore, we contribute to a better knowledge of the evolutionary history of Santa Maria with particular focus on its earlier stages of growth. We infer that the outer 14 part of the northern shelf is carved on units older than the oldest volcanic sequences exposed on 15 16 land; the western and southern shelves are almost entirely carved in the products of the oldest 17 shield volcano (Anjos Volcanic Complex), whilst the eastern shelf is the youngest in formation, being related to units of the Pico Alto volcanic complex. We also discuss the distribution of 18 19 sediments on the shelf. High wave energy concomitant with sea level drops contributed to the 20 erosion and transport of older sequences offshore, which were ultimately lost to the slopes of the 21 island, resulting in a shelf that is sediment-stripped for the most part. Presently, wide shelves 22 further contribute to wave-energy dissipation, resulting in a significantly diminished sediment 23 production. The shelf edges around Santa Maria exhibit several erosive scars, interpreted as the head of a submarine drainage network carved on the slopes of the island. On the western, 24

southern and eastern shelves, these mass-wasting features are mostly controlled by tectonics since the extension of faults and lineaments can be followed from onshore to the shelf edge. On the northern shelf mass-wasting appears to be mostly correlated with presence of unstable sediments on the shelf edge that triggered landsliding. Overall, this work shows how detailed mapping of the shelves surrounding old volcanic islands can still provide important complementary information about the evolution of these islands, which could not be acquired solely by the study of their subaerial and deeper submarine parts.

# 1 **1. Introduction**

2 The subaerial and submarine morphology of volcanic islands results from the interplay between 3 volcanic, tectonic, erosive and depositional processes acting at different time-scales (Ramalho et 4 al., 2013; Casalbore et al., 2015; Lebas et al., 2018; Quartau et al., 2018a). These processes and their influence on the geomorphology are very well known for the subaerial portions of volcanic 5 6 islands (e.g., Thouret, 1999). However, volcanic islands are mostly developed beneath the sea; 7 hence the study of their submarine areas can provide much more information than the subaerial 8 parts, which constitute only the "tip of iceberg" (e.g., Quartau and Mitchell, 2013; Quartau et al., 9 2015a). Most part of submarine studies was mainly addressed to reconstruct large-scale instability processes that commonly affect the flanks of these edifices (Moore et al., 1989; Deplus et al., 10 2001; Masson et al., 2002; Mitchell et al., 2002; Oehler et al., 2008; Casalbore et al., 2011; 11 12 Montanaro and Beget, 2011). On the contrary, less attention has been paid on the impact of 13 volcanism, erosion, tectonics, time-variant sediment supply and climatic/oceanographic conditions on the surrounding shelves and the link with the geology onshore. Only in the last decade, a few 14 number of nearshore marine surveys performed with high-resolution multibeam and seismic 15 reflection systems revealed the importance of insular shelf studies to obtain a more detailed 16 picture of the geological history of volcanic islands (Quartau et al., 2012; Romagnoli, 2013; 17 Quartau et al., 2014; Quartau et al., 2015b; 2016; Casalbore et al., 2018; Ricchi et al., 2018; 18 19 Romagnoli et al., 2018). Insular shelves surrounding volcanic islands are, in fact, representative of 20 their original subaerial extension during glacial maxima, thus playing a potential role for palaeomorphological and relative age reconstructions as well as for evaluating the vertical mobility of 21 such areas (Lucchi et al., 2019; Quartau et al., 2016, 2018b; Ricchi et al., 2018). Geomorphic 22 23 parameters such as the shelf edge depth and shelf width, as well as the morphology of the 24 submerged lava flows or rocky outcrops, provide useful insights concerning the original

25 morphology of the volcanic edifices, and the occurrence of (older or younger than expected) 26 volcanic activity not recorded onshore (Quartau et al., 2014; 2015b; Ramalho et al., in press; Romagnoli et al, 2018). For instance, shelf width can be related with shelf age (beginning of 27 incision) and as such may be used to infer reveal relative ages of undated coastal sectors (Quartau 28 et al., 2010; Romagnoli et al, 2018). In a similar fashion, coastal sectors formed by volcanism with 29 30 similar ages but different orientations to waves generally exhibit different shelf widths and may 31 thus reveal that wave regimes have probably not changed through time (Llanes et al., 2009; 32 Mitchell et al, 2003; Quartau et al., 2010). Erosional shelf breaks which are now deeper than the lowest sea-level stands, may also be related with subsidence of the volcanic edifices and thus rates 33 of vertical movements can also be estimated from the depth of the shelf break (Lucchi et al., 2019; 34 35 Quartau et al., 2014, 2016, 2018b), in a similar way that constructional slopes breaks were used in 36 Hawaii to estimate subsidence and relative age constraints of volcanic edifices (Taylor, 2019). To date, however, only a few studies have discussed sediment deposition on volcanic island shelves 37 and the preservation of their depositional sequences (Ávila et al., 2008; Casalbore et al., 2018; 38 39 Meireles, 2013; Quartau et al., 2012). Moreover, all the existing shelf studies have focused on relatively young islands (~1 Ma or less). At Santa Maria, however, there there is the opportunity of 40 41 looking into a much older island (~6 Ma) and test whether the approaches and interpretation 42 taken from younger islands could also apply to older edifices. In addition, Santa Maria's shelf is up 43 to now, one of the best surveyed insular shelves in the world, with a cover of multibeam bathymetry of 1 m resolution and a grid of high-resolution seismic profiles not spaced more than 44 45 250 m. This opportunity provides a unique and more detailed, careful analysis on the factors that 46 impact shelf evolution and sedimentary dynamics in an energetic environment such as the Azores. In this study, we combined high-resolution geophysical dataset with detailed onshore 47 geological field studies performed on Santa Maria Island (Azores Archipelago) to better 48

49 understand the connection between the onshore and the offshore domain. This allows 50 unprecedented inferences on the contribution of tectonics, wave regime, volcanism, erosion and 51 sediment supply in shaping the insular shelf. Moreover, through the analysis and comparison of 52 the shelf width, shelf edge depth and morphology for each sector of the island, a better 53 understanding of the evolution of Santa Maria is derived.

54

# 55 **2. Geological setting**

56

## 57 2.1 The Azores Archipelago

The Azores Archipelago is located in the North Atlantic Ocean, on both sides of the Mid-Atlantic 58 59 Ridge (MAR in Fig. 1), which separates the westernmost group of islands from the central and 60 easternmost groups. Volcanism here possibly results from the interaction between the triple 61 junction of the Eurasian, Nubian and American lithospheric plates and a hotspot (Gente et al., 2003). The complex interaction between the seafloor spreading forces driven by the Mid Atlantic 62 Ridge and the right-transtensional movements associated with the Nubian/Eurasian plate's 63 64 boundary dominates the area (Lourenço et al., 1998; Hipólito et al., 2013; Marques et al., 2013; 65 Madeira et al., 2015). Santa Maria sits at the SE corner of a roughly triangular shaped area located between the Nubian/European plate boundary (the active Terceira Rift, TR in Fig. 1) and the 66 67 presently inactive East Azores Fracture Zone (EAFZ in Fig. 1), most likely already outside the influence of the Nubian/European plate boundary (Miranda et al., 2018). 68

Being under an oceanic regime, the coastlines of the Azores Archipelago are frequently subjected
to high wave energy. The North Atlantic Oscillation and Eastern Atlantic atmospheric circulation
bring extreme storms (wave heights exceeding 16 m and peak periods exceeding 18 s) affecting
the Azores at least once every seven years (Andrade et al., 2008; Zhao et al., 2019). For instance,

- the recent Hurricane Lorenzo impacted the Azores in October 2019, with wave heights yielding 15
  m that violently destroyed the port of Lajes on Flores Island, western Azores (Hatton, 2019).
- 75

#### 76 2.2 Geology of Santa Maria Island

Santa Maria Island rises from the seafloor at around -2500 m and it is considered the oldest 77 78 island of the Azores Archipelago, since it emerged from the sea c. 6 Ma ago (Ramalho et al., 2017). 79 The eastern (leeward) side of the island exhibits a rugged morphology, culminating in the Pico Alto 80 high (587 m in elevation, Fig. 2). In this sector, the coastline is made of up to 250-300 m high plunging cliffs, occasionally displaying wave-cut notches at varying elevations. On the contrary, the 81 western (windward) side has a typical staircase morphology due to the occurrence of a number of 82 83 raised marine terraces from 7 m to 200 m in elevation, some of which still exhibit remains of 84 coastal sediments (Ramalho et al., 2017). Offshore, Santa Maria is surrounded by a wide insular shelf, in particular along the northern coastal sector (up to 8 km wide), in contrasts with narrower 85 shelves at the western, southern and eastern sides (~2 km). Five sets of submerged marine 86 terraces have been recognized on the shelf, reflecting the interplay between the island's vertical 87 movements and sea-level fluctuations (Ricchi et al., 2018). 88

89 The volcanic history of Santa Maria is the result of the activity of different overlapping 90 volcanoes, interrupted by relative long periods of quiescence and erosion (Serralheiro et al., 1987; 91 Serralheiro and Madeira, 1993; Ramalho et al., 2017). The volcanism of the Cabrestantes and 92 Porto Formations on the westernmost side of the island (Fig. 2), emplaced by surtseyan and 93 strombolian activity, witnesses the first emergence of the island and transition to a subaerial 94 volcanic edifice c. 6 Ma ago. The products of the Anjos Volcanic Complex are widespread in the 95 central and western portions of the island, representing the remnants of a large shield volcano 96 formed between 5.8 Ma and 5.3 Ma (Fig. 2) (Sibrant et al., 2015; Ramalho et al., 2017). The

97 following 1 Ma was mostly characterized by marine erosion that caused the partial dismantling, 98 and possibly submersion of the previously formed shield volcano. A collapse of the eastern flank of the Anjos volcanic edifice has also been proposed by Sibrant et al (2015). Simultaneously with 99 marine sedimentation on top of the eroded volcano, which probably constituted a wide shoal with 100 101 some residual subaerial reliefs, low-volume submarine volcanism took place, mainly focused on 102 the eastern portion of the island (Ramalho et al., 2017). This culminated in the emplacement of 103 the Touril Volcano-sedimentary Complex between 5.3 Ma and 4.1 Ma (Fig. 2). During this period 104 the island experienced a subsidence trend (c. 100 m/Myr on average) that lasted until 3.5 Ma as attested by the numerous passage zones of Pico Alto and Anjos lava deltas (which indicate 105 increasingly higher relative paleo sea levels towards the top of the sequence) and the 106 107 transgressive sequences of Touril complex showing a deepening-up sequence (Ramalho et al., 108 2017). Widespread volcanic activity resumed c. 4.1 Ma, promoting the re-emergence of the island edifice. The products of the Pico Alto volcanic complex (Fig. 2) correspond to the lateral growth of 109 the edifice due to lava delta progradation over the Touril marine sediments and the eroded 110 remains of the Anjos edifice. Subsidence reversed into uplift at 3.5 Ma (estimated at c. 59 m/Myr 111 112 from 3.5 Ma to 2.14 Ma and 42 m/Myr from 2.14 Ma to present), as attested by the position of 113 the passage zone of Pico Alto's lava deltas (up to 200 m), and the occurrence of marine terraces on 114 the western side of the island and offshore (Ramalho et al., 2017; Ricchi et al., 2018). Although the 115 last stage of the island evolution is considered mainly erosional, from 3.2 to 2.8 Ma a low volume 116 of magmatic and hydromagmatic activity led to the formation of the cones and products of the 117 last post-erosional phase, roughly corresponding to the Feteiras Formation of Serralheiro et al. 118 (1987) or the Young Parasitic Cones of Sibrant et al. (2015) (Fig. 2).

119 Tectonics is expressed onshore by the main NW–SE, N-S and NE–SW-trending faults (Fig. 2, 120 Madeira et al., 2015) that partly influenced the development of the subaerial terraces and

submerged terraces on the shelf (Ricchi et al., 2018). Moreover, Zbyszewski and Ferreira (1960)
and Serralheiro et al. (1987) reported a large set of N 045° and N 150° striking dikes located mostly
in the SW and NE portion of the island (Fig. 2), being likely contemporaneous to the main building
stages of the island.

Santa Maria is the driest island in the Azores, with the low western half exhibiting between 670 and 800 mm/ year of precipitation and the more rugged and taller eastern side reaching higher values, between 1100-1700 mm/year (Fernandes, 2004).

128

## 129 **2.3 Sedimentary dynamics at reefless Volcanic Ocean Islands**

130

131 On reefless volcanic islands the production of sediments reaching the coast largely depends on 132 volcanic activity (for instance explosive volcanism produces much more clastic materials than 133 effusive volcanism), the occurrence of a well-developed subaerial drainage system, wet climate 134 stimulating the subaerial erosion, and high wave energy promoting coastal erosion and sediment redistribution (Krastel et al., 2001; Ramalho et al., 2013; Quartau et al., 2018a). In particular, on 135 136 small and relatively dry reefless volcanic islands (like Santa Maria), sediment is produced mainly 137 through the erosion of coastal cliffs by wave processes because rivers do not have enough area to develop and the low precipitation imprints a weak fluvial erosion regime (Quartau et al., 2012). 138 139 Thus, on these islands, sediments that result from cliff erosion accumulate in the inner shelf and 140 are partly transported offshore during storms. Several studies showed, in fact, that storms are able to produce storm-induced downwelling currents that transport sediments from shallow to deeper 141 142 areas of the shelf (Field and Roy, 1984; Tsutsui et al., 1987; Hernández-Molina et al., 2000; Chiocci 143 and Romagnoli, 2004; Meireles et al., 2013). These sediments tend to form clinoform deposits, as 144 recognized on other volcanic islands of the Azores (Quartau et al., 2012; 2014; 2015b) and in the

Mediterranean Sea (where they are named "submerged or submarine depositional terraces",
Chiocci and Romagnoli, 2004; Casalbore et al., 2017; Romagnoli et al., 2020).

147

## 148 **3. Material and methods**

149

#### 150 **3.1 Shelf data acquisition and processing**

151 The shelf around Santa Maria Island has been surveyed onboard the R/V Arguipélago from 24<sup>th</sup> 152 August to 15<sup>th</sup> September 2016 in the scope of PLATMAR project. The survey (PLATMAR 1/2016) included the acquisition of multibeam bathymetry and single-channel seismic profiles (boomer) 153 between c. -20 m and c. -250 m (Figs. 3 and 4). Further details of the geometry acquisition and 154 equipment were reported by Ricchi et al. (2018). The subaerial DTM was generated from a 1/5000 155 scale digital altimetric database with a resolution of 5 m. An integrated analysis of the multibeam 156 157 bathymetry and subaerial topography has been mainly used to study the morphology of the shelf by mapping the volcanic, structural, erosional and depositional features and comparing with those 158 of the corresponding onshore sectors (Figs. 4, 5a, 6a, 7a, 8a and 1ESM). The shelf width and the 159 160 erosive edge depth values in each sector were determined from the seismic profiles collected perpendicular to the coastlines (Fig. 3) and reported in graphs in order to constrain its variability 161 along the shelf (Figs. 5a, 6a, 7a and 8a). The erosive shelf edge was defined as the point, in the 162 163 rocky basement of the seismic profiles, where there is a significant increase in gradients to the slope of the island (e.g., Fig. 2ESM). Shelf width was only measured where erosive shelf edge and 164 coastlines stretches were roughly parallel to each other, whilst shelf edge depth was measured 165 166 along the entire shelf. Since mass wasting of the shelf edge can cause a landward shift of that feature and shallow its depth, we always took into account in our interpretation areas without 167 scars, which is considered to represent the original depth at which waves cut the island subaerial 168

flanks. The analysis of seismic reflection profiles allowed mapping the distribution and thickness of the unconsolidated sediments lying on the shelf of Santa Maria Island (Fig. 9 and Table 1). In each seismic profile we picked two reflectors corresponding to the seafloor and, where present, the bottom of the sedimentary cover, respectively. For time/depth conversion we used a sound velocity of 1500 m/s for the water column and 1800 m/s for the sediments. Sound velocity is based on data of Hamilton and Bachman (1982) for coarse sands, which in the Azorean shelves are the dominant grain-size (Quartau et al., 2012; Quartau et al., 2015b).

176

### 177 **3.2 Wave regime characterization**

Using the model outputs of the European Centre for Medium range Weather Forecasts 178 (ECMWF) 40-yr Re-Analysis (ERA-40) (Uppala et al., 2005) and ECMWF Interim Re-Analysis (ERA-179 180 Interim) (Dee et al., 2011), a rose diagram for Santa Maria offshore significant wave heights (and 181 their associated percentages and originating directions was produced (Figure 10, Table 1ESM). These wave properties were extracted from a six hourly significant wave height H<sub>s</sub> database and 182 averaged for the sea area adjacent to Santa Maria Island (36.875°N - 37.125°N, 24.9375°W -183 184 25.25°W) for the period September 1957 to August 2019 and represent the base for discussing the 185 the wave regime in the different shelf sectors.

186

# 187 **4. Results**

188

# **4.1** Morphology of the insular shelf around Santa Maria Island

## 190 4.1.1 Northern shelf sector

This sector is located offshore the northern coast of the island from Baía do Mar da Barca to
Ponta do Norte (A to B in Fig. 5a). The roughly ENE-WSW oriented coastline has a strongly irregular
1

193 profile characterized by embayments and promontories. From Ponta do Norte to Ponta dos Frades, sea cliffs are steep, up to 200 m high, and were cut in the products of the Anjos Volcanic 194 Complex at the base, overlapped by those of Touril Volcano-sedimentary and Pico Alto Volcanic 195 196 Complex (see Fig. 2), forming a tabular sequence, dipping very gently to the eastern quadrant. 197 From Ponta dos Frades to Baía do Mar da Barca, cliffs are entirely carved in the products of the 198 Anjos Volcanic Complex and their height progressively decreases westwards to ~20-30 m. The 199 entire shelf has a roughly trapezoidal shape and shows a different morphology between the 200 eastern and the western sides, similarly to the asymmetric topography of the island. To the east of Ponta dos Frades the shelf width varies between 6 km and 8 km, while to west is narrower being c. 201 4 km wide (not considering the sector where the submarine cone of Baixa do Ambrósio is located, 202 203 Figs. 5a and 5b). The maximum depth of the erosive shelf edge is c. 170 m on the northwestern 204 portion (off Baía do Mar da Barca, Figs. 5a and profile aa' in Fig. 2ESM), then it slowly decreases to 205 156 m to the north and increases again up to 172 m in the northeast. Depth values as low as c. 115 206 m can be found, corresponding to scars affecting the shelf edge.

To the west of Ponta dos Frades, rocky outcrops have been recognized down to a depth of 80-207 208 100 m, being mostly located in the inner to middle shelf. These are cut by sharp linear 209 escarpments, ranging in heights from c.10 m to c.30 m in three compartmented blocks (Fig. 5a). 210 These lineaments, 2 to 2.7 km long, can be followed from the subaerial island into shallow water 211 and down to -100 m, where they apparently disappear in an area characterized by an overall 212 smooth morphology. Offshore continuation of these lineaments is possibly represented by a sharp 213 escarpment that dissects a rocky outcrop located to the south of Baixa do Ambrósio at c.-130/-170 214 m (probable extension of AjF in Fig. 5a) and by a narrow gully head, located SE of Baixa do 215 Ambrósio at -135 m (probable extension of CF in Fig. 5a). To the east, another lineament (RF in Fig. 216 5a) can be also apparently followed offshore in a NNW-SSE linear escarpment c. 2 km long, down

217 to -50 m. These major lineaments are in clear connection with recognized fault 218 planes/escarpments on land, as they are more evident in the case of the Aeroporto and Anjos 219 Faults (AF and AjF in Fig. 5a and profile cc' in Fig. 3ESM).

To the east of Ponta dos Frades, the entire shelf is dominated by rocky outcrops with some sub-220 221 conical or elongated morphological highs. A total of 26 sub-conical structures were recognized 222 from c. -70 m to c. -120 m on this shelf sector (Fig. 5a). These morphological features show a 223 random arrangement on the shelf and exhibit different sizes and morphologies. Four types of sub-224 conical landforms were identified in this sector: a) small conical, circular or elongated, positive relief (numbers 1–9 in Fig. 5a); b) larger, more irregular and concentric morphologies with positive 225 relief and sometimes a flat or depressed top, and with a basal diameter from c. 0.25 km to c. 1 km 226 227 and heights comprehended between c. 20 m and c. 40 m (numbers 10-18 in Fig. 5a); c) large 228 roughly-circular features with a wide central depression, ranging from 800 m to 1.2 km in width and up to 30 m in height (numbers 19-23 in Fig. 5a); and d) small round-to-elliptical ring-like 229 features, up to c. 600 m in diameter and 15 m high, with an internal depressed area covered by 230 sediments (numbers 24 and 25 in Fig. 5a). A large elliptical elevation (Baixa do Ambrósio, 1.3 km in 231 232 diameter and about 130 m high) also stands prominently isolated on the NW portion of this shelf 233 sector. This is a popular recreational diving site and submarine photos of this area abound. These 234 show that this morphology is a pillow-lava cone whose flanks are cut by marine terraces located at 235 different depths (Ricchi et al., 2018). In addition to the conical morphologies, a prominent NNW-236 SSE rocky ridge can be seen in the central-eastern part of the shelf, extending from the coastline 237 to the shelf edge, almost exactly in line with the Pico Alto ridge onshore (see arrow in Fig. 5a and 238 compare it to Fig. 2). Along this submarine ridge, several N-S linear positive reliefs are present also 239 in line with dykes exposed at the coast (see Fig. 4 and 1ESM), in between Lagoinhas and Ponta do 240 Pesqueiro Alto (Fig. 2) (Zbyszewski and Ferreira, 1960; Madeira, 1986; Madeira et al., 2015).

241 In terms of surface area, the sedimentary cover amounts almost the same as the sediment-free rocky surfaces, i.e., around 50% each (see Table 1). A total of 0.515 km<sup>3</sup> of sediments are 242 deposited on this sector and the ratio volume of sediments/shelf surface area is 0.007. Sediments 243 are mostly restricted to the outer part of shelf in the NW and in-between the rocky outcrops in the 244 245 NE (Fig. 9). Deposits are normally very thin (mostly between 0 and 2 m), being locally thicker (up to 246 14 m) on local depocenters bordered by sub-vertical escarpments (e.g., the NW-SE oriented small 247 basin located offshore Baia da Cré, and the ENE-WSW oriented basin offshore Ponta do Norte -248 see Fig. 9 and seismic profiles bb' and cc' in Fig. 3ESM).

249

#### 250 4.1.2 Western shelf sector

251 This sector is located offshore the western coast of the island from Baía do Mar da Barca to 252 Ilhéu da Vila (Fig. 6a). The coastline is NNW-SSE oriented with sub-vertical sea cliffs up to 75 m high, entirely carved in the subaerial westward-dipping lava flows and pyroclastic succession of the 253 254 Anjos Volcanic Complex (Fig. 2). The shelf runs parallel to the coastline and is generally rocky down to c. -50/-70 m. Offshore Campo Grande a NW-SE strip of sedimentary seafloor (Fig. 6a and 9) 255 256 around 1.7 km long and c. 150 m wide separates the shelf in two parts characterized by different 257 widths (c.1.2-1.6 km to the north, c.2 km to the south). In this sector the erosive shelf edge is generally at -100/-110 m along the northern portion, whilst in the southernmost part its depth 258 259 increases to -135/-147 m (Figs. 6a, 6b and profile dd' in Fig. 4ESM). However, between offshore 260 Campo Grande and Ponta do Poção, the shelf edge has been indented, shallowing its depth up to -261 50 m.

The lineament located offshore Campo Grande exhibits onshore an arc-shaped escarpment dissected by a set of WNW-ESE and WSW-ENE structures and the Campo Grande fault (CgF in Fig. 6a, Madeira et al., 2015). In this whole area, the rocky outcrops seen in the high-resolution

bathymetry are frequently cut by these WNW-ESE and WSW-ENE lineaments (see figure ESM1),
which are parallel to the dike network seen on the adjacent coast (Zbyszewsky and Ferreira, 1960;
Madeira, 1986; Madeira et al. 2015). In a similar fashion, a few well-pronounced NNW-SSE to NNESSW lineaments can be seen cutting the rocky shelf in this area, running approximately parallel to
shore and to the Aeroporto Fault further inland (Fig. 4, AF in Fig. 6a and Fig. 1ESM).

The surface of the seafloor in this sector corresponds to ~75% of rocky surfaces and 25% of sedimentary cover (Table 1). It has total of 0.037 km<sup>3</sup> of sediments and the ratio volume of sediments/shelf surface area is 0.003. Sediments are therefore restricted to a few scattered patches at -100/-120 m. These are also present inside the WNW-ESE oriented lineament offshore Campo Grande, with thicknesses less than 5 m, which increase up to c. 10 m at the shelf edge (Fig. 9).

276

#### 277 4.1.3 Southern shelf sector

This sector is located offshore the southern coast of the island from Ilhéu da Vila to Ponta do 278 Castelo (Fig. 7a). In the western-most portion to Ponta da Marvão, the coastline is very irregular in 279 280 plan-view, with cliffs mostly formed by SW-dipping lava flows and pyroclastic products of the 281 Anjos Volcanic complex (Fig. 2). From Ponta da Marvão to Ponta do Malbusca, the coastline exhibits an arcuate shape due to the occurrence of a 6 km-wide bay (Praia Formosa Bay, Fig. 7a), 282 283 bounded by up to 200 m-high, sub-vertical sea cliffs. These are cut in a sequence comprising the Anjos Volcanic Complex lava flows at the base, overlapped by marine sediments and 284 submarine/subaerial volcanic products of both the Touril and the Pico Alto Volcanic Complexes 285 286 (Fig. 2). From Ponta da Malbusca to Ponta do Castelo the coastline is made of steep cliffs, carved 287 on the Touril and Pico Alto sequences (Fig. 2).

288 The southern shelf has an irregular morphology, with its edge showing a configuration that roughly mimics the coastline west of Ponta da Malbusca. The westernmost portion of the shelf is 289 around 2 km wide, up to the eastern end of Praia Formosa Bay, after which it is reduced to c. 1 km 290 offshore Ponta da Malbusca. Eastwards it enlarges again, increasing to up to 2.6 km in width 291 292 offshore Ponta do Castelo, at the SE edge of the sector (Figs. 7a and 7b). The depth of the erosive 293 shelf edge follows a similar trend, i.e., deeper where the shelf is wider and shallower when it is 294 narrower. Between south of Ilhéu da Vila to Praia Formosa Bay it is 80/90 m (although can reach 295 120 m in places), while it is in the range of 50/90 m at Ponta da Malbusca. Then, the shelf edge depth progressively increases to a maximum value of 142 m offshore Ponta do Castelo (Figs. 7a, 7b 296 and profile ee' in Fig. 5ESM). 297

298 A dense network of NNW-SSE and NNE-SSW trending lineaments, 0.5 km to 2 km long and with 299 escarpments up to 15 m high, dissects the shelf from the coastline down to its edge, creating a succession of rocky structural high interspersed by sediment-covered depressed areas (Figs. 7a, 9 300 and seismic profile ff' in Fig. 6ESM). The most evident features are located offshore Vila do Porto, 301 302 Praia Formosa Bay and Ponta da Malbusca, also having a clear morphological expression onshore 303 (Fig. 7a). The lineaments that compartmentalize the shelf show an orientation similar to the dike 304 network exposed all along this stretch of coastline (see Zbyszewsky and Ferreira, 1960 and Madeira, 1986), (Fig. 7a) and, in some cases, can be followed to the outer shelf. The western Fault 305 306 of Malbusca (FM in Fig. 7a) also continues offshore, as expressed by a small NNE-SSW linear escarpment. 307

The distribution of sedimentary cover and rocky surfaces is very similar to the northern sector, around 50% each (Table 1). This shelf sector has a total of 0.221 km<sup>3</sup> of sediments and a ratio volume of sediments/shelf surface area of 0.012. The distribution of the sediments and their thickness is highly variable being from <6 m (offshore Ilhéu da Vila) up to 15-20 m (offshore Praia

Formosa Bay, Fig. 9, seismic profile ff' in Fig. 6ESM and seismic profile gg' in Fig. 7ESM). Offshore Praia Formosa and west of Ponta da Malbusca the sediments cover most of the shelf, whilst offshore Ilhéu da Vila and from Ponta do Malbusca to Ponta do Castelo the sediments cover only the outer shelf with a gradually decreasing thickness towards the shelf edge (Fig. 9).

316

# 317 4.1.4 Eastern shelf sector

318 This sector is located offshore the eastern coast of the island from Ponta do Castelo to Ponta do 319 Norte. The coast is very irregular in plan-view, being characterized by several embayments and promontories, with steep cliffs carved in the eastward-prograding lava-delta sequences of the Pico 320 Alto Volcanic Complex (Fig. 2). The NNW-SSE oriented shelf runs parallel to the coastline and is c. 321 322 0.6-2.4 km wide, being wider offshore Ponta do Castelete and São Lourenço bay (Figs 8a and 8b). 323 The erosive shelf edge depth varies mostly between 80 m and 100 m between Ponta do Norte and the shelf edge embayment offshore Pontinha, whilst south of Potinha it is much shallower, being 324 between 50 m and 100 m (Figs. 8a and 8b). Rocky outcrops are frequently extended down to the 325 shelf edge, where in places they form eastward-dipping ramps, with landward lips up to few 326 meters high (Fig. 8a and seismic profile hh' in Fig. 8ESM). Apart from some small-scale 327 328 indentations affecting the shelf edge, the eastern sector exhibits only an evident embayment that 329 dissects the shelf at c. -60- -80 m offshore Pontinha, in front of a concave cliff on the coast (Fig. 330 8a).

The distribution between the sedimentary cover and rocky surfaces is very similar to the northern and southern sectors, with 54% corresponding to sediments and 46% corresponding to rocky surfaces. This sector has a total of 0.187 km<sup>3</sup> of sediments and a ratio volume of sediments/shelf surface area of 0.015. Along the eastern shelf sector the sediments fill some depressions bordered by rocky outcrops and cover the shelf down to c. -70 m, showing a thickness

variable from 2 to 15 m (Fig. 9). The most extensive sediment accumulation, lying offshore
Pontinha is a depression between rocky outcrops (Figs. 9 and 9ESM).

338

# **4.2 Wave regime around Santa Maria Island**

The ERA outputs of Santa Maria wave properties suggest that the prevailing swells approach Santa Maria Island from the northwest (35%) and west (22%), with average significant wave heights ( $H_s$ ) of 2.14 m and 2.44 m respectively (Fig. 10 and Table 1). Waves from north and northeast are also frequent (respectively 19% and 12%) but lower (respectively  $H_s$  of 1.92 m and 1.83 m). The remaining directions are insignificant (< 5% each), however waves from the southwest can still reach 2.45 m, but only during 5% of the year.

346

# 347 **5. Discussion**

348

# 349 **5.1 Sediment distribution on the shelf**

350 In principle the shelf sectors surrounding the eastern part of Santa Maria should be filled with 351 more sediment because cliffs are higher and streams are more abundant. In what concerns 352 accommodation space the western, southern and eastern shelf sectors have similar areas, hence 353 similar space to accommodate deposits (see Table 1). However, the western shelf is mostly sediment-free, with deposits only on the outer shelf and very thin (c. 2 m thick, Fig. 9). The ratio 354 volume of sediments to area of the shelf in this sector is the lowest (0.003) and around one quart 355 356 to one fifth of the southern and eastern sectors (respectively 0.012 and 0.015) The southern and eastern shelf sectors exhibit then a wider sedimentary cover (around 50% of the seafloor), but 357 mostly located in the middle to outer shelf. The northern sector shows a similar sediment 358 359 distribution pattern to the southern and eastern sectors, although being much wider. The deposits

360 are very thin (c. 2m) and are mostly on the middle to outer shelf. The fact that nearshore areas of the four sectors are mostly sediment-stripped, despite that most sediments should be produced 361 362 from cliff erosion is somehow puzzling. This sediment distribution pattern is also seen in other 363 islands of the Azores archipelago such as Faial and Pico and related with high wave energy of this archipelago (Quartau et al., 2012; Quartau et al., 2015b). At Santa Maria this pattern is 364 365 exacerbated on the western sector relatively to the southern and eastern sectors because this 366 sector is more exposed to the dominant waves, hence transport of sandy sediments to deeper areas is higher and these are probably lost to the slope. In addition, this sector has a poorly 367 developed subaerial drainage and the cliffs are smaller, so fewer sediments accumulate nearshore. 368 Sediments of the southern and eastern shelf sectors, being on the leeward side of the island are 369 370 less easily transported offshore and consequently may rest at shallower depths (from 35/40 m 371 downwards). The presence of preserved small terraces and wave-cut notches attributed to the last interglacial (and older interglacials) along the SE, E and NE shores (Ramalho et al., 2017) also 372 suggests that wave influence is also lower in these sectors, otherwise these morphologies would 373 have been obliterated receding cliff erosion. In addition, these sectors are partially or entirely 374 375 adjacent to the higher and rainier side of the island, with a more evolved subaerial drainage, which 376 presumably delivers higher quantities of sediments that then accumulate nearshore. The northern sector is also exposed to the dominant waves as the western sector. However, given that this 377 378 sector is much wider, sediments are probably not transported much further than the middle to 379 outer shelf and thus are not entirely lost to the slope, leading to a higher sediment accumulation when compared with the western sector. Nevertheless, due to such a large accommodation space, 380 381 the ratio volume of sediments to area of the shelf in this sector is still the second lowest (0.014) 382 after the western sector.

The shelf around Santa Maria lacks well-developed sedimentary bodies as those recognized on other islands of the Azores and the Mediterranean Sea where some portions of the shelf are covered by up to 30-40 m thick volcaniclastic sequences (Chiocci and Romagnoli, 2004; Quartau et al., 2012; Casalbore et al., 2017; Romagnoli et al., 2020). Given that Santa Maria is much older and eroded for a longer period of time should not the shelf have thicker deposits? We believe that the thin sediment cover offshore Santa Maria can be explained by three main reasons:

389

 Increasing wave attenuation with shelf width. As shelves became wider, not only the space for sediment accommodation increases, but the energy reaching the coastline decreases due to wave attenuation over wider and shallower shelves (Quartau et al., 2010).
 Consequently, the lower sediment production by cliff retreat and the higher accommodation space explains the thin cover around Santa Maria shelf.

2. Santa Maria Island is a low and small island, with less precipitation and lower rates of stream erosion when compared with other Azorean Islands (which exhibit rainfall regimes frequently above 2000 mm/year, AEM & IMP, 2012). Thus, sediment production is likely dominated by cliff erosion. On the other hand given that island is mostly bound by low cliffs, their erosion produces less sediments than in taller islands.

Preservation issues, i.e., the weaker potential for preservation of depositional sequences on
 the shelves of reefless volcanic islands that are subjected to highly-energetic wave regimes.
 The Azores archipelago is considered an energetic environment in terms of winds and wave
 action (Andrade et al., 2008; Quartau et al., 2012; Rusu and Guedes Soares, 2012). In these
 settings, sediments tend to be transported offshore. Furthermore, as sea level falls, most of
 the sediments are transported below the shelf edge, leaving the shelves completely
 stripped of sediments. This cycle is repeated every glacio-eustatic oscillation, preventing the

407 accumulation of several depositional sequences, as seen on continental shelves (Ávila et al,
408 2008; Quartau et al., 2012).

409

The preservation issues also apply to other volcanic ocean islands subjected to high wave energy regimes. However, on the case of Santa Maria, the two previous reasons are more relevant and probably are enough to explain why Santa Maria lacks thicker sedimentary deposits in its shelf.

414

## 415 **5.2 Mass wasting processes**

416

417 Hampton et al. (1996) and Locat and Lee (2002) suggested several mechanisms for submarine 418 landslide triggering. These may include oversteepening, seismic and storm-wave loading and the incohesive nature of the sediments. The subaerial and submarine flanks of volcanic islands are 419 prone to be affected by large-scale sector collapses, as attested by both collapse scars and 420 submarine deposits seen in archipelagos such as the Hawaii (Moore et al., 1994), Canaries 421 (Masson et al., 2002), Cape Verde (Masson et al., 2008), Aeolian (Romagnoli et al., 2009a; 422 423 Romagnoli et al., 2009b), Aleutians (Montanaro and Beget, 2011), Azores (Costa et al., 2015; 424 Quartau et al., 2015b) and Madeira (Quartau et al., 2018a). Large-scale flank collapses have been 425 proposed for Santa Maria by Sibrant et al. (2015) but at least the younger collapse proposed by these authors is at odds with onshore field studies documented by Ramalho et al. (2017), and 426 submarine evidence for these collapses is absent. However, small scale mass wasting features 427 428 were identified at the edge of the insular shelf surrounding Santa Maria Island, and these are 429 mostly concentrated on the northern and southern sectors. The largest scars are located along the 430 NE shelf edge, between the areas offshore Ponta do Norte and Baía da Cré, and exhibit a wide

431 semicircular shape (Fig. 5a). Several gully heads, with an elongated shape, are also found at the southern shelf edge, between the areas offshore Vila do Porto and Praia Formosa (Fig. 7a). These 432 433 morphological differences might be related to the processes responsible for the formations of the 434 scars. Along the edge of the NE shelf sector, the scars are apparently controlled by the distribution 435 of the eroded cones and the rocky outcrops on the shelf (Fig. 5a). The areas between the 436 cones/outcrops are probably composed of more friable materials, thus representing a possible 437 lithological weakness, and a preferential pathway for sediment reworking. Despite the fact that nowadays the wide northern shelf prevents any direct stream discharge onto the outer shelf and 438 439 the upper submarine slope beyond the shelf break, during lowstands sedimentary flows probably were able to reach these areas, carving the head of those "proto-gullies". 440

441 Cones are absent along the southern shelf sector, where the elongated morphology of the scars 442 seem to reflect instead NE-SW striking structural weakness corridors, related to the faults that 443 extend to the south of the island and affect the shelf inducing localized erosion (Fig. 7a). The same 444 structural control might be inferred for the formation of the narrow scar located at the shelf edge offshore Ponta dos Frades, in the northern shelf sector (Fig. 5a), being probably the continuation 445 446 of the faults recognized onshore (Madeira et al., 2015) and on the inner part of the shelf (Ricchi et 447 al., 2018). Similar features can be found on the NE, southern and eastern shelf sectors where likely fault-controlled basins coincide with the scars of shelf-edge landslide (Fig. 9 and profile b-b' in Fig. 448 449 3ESM, profile f-f' in Fig. 6ESM and profile i-i' in Fig. 9ESM).

In the case of Santa Maria, it appears that on the southern and in some places on the eastern shelves and northern shelf these mass-wasting processes are mostly controlled by tectonics, because scars appear to be the extension of onshore tectonic features. However, on the northern shelf sector, tectonics appears to play a smaller role. On these sites it is more likely that the accumulation of incohesive sediments near the shelf edge (with high gradients) might trigger

455 those landslides as suggested by Casalbore et al. (2015) in Terceira Island, Quartau et al. (2012) in Faial Island, Quartau et al. (2018a) and Santos et al. (2019) in Madeira archipelago, promoting 456 shelf edge retreat. A quite similar setting is found in other shelves around Italian islands, such as 457 Vulcano (Romagnoli et al., 2013), Lipari (Casalbore et al., 2016a), and Ventotene islands (Casalbore 458 459 et al., 2016b), evidencing how the shelf edge is susceptible to landsliding processes. Most of these 460 islands are volcanic active, thus the production of volcanogenic sediment is higher favoring the 461 widespread occurrence of small-scale mass wasting, especially where thick volcanogenic aprons 462 are present. On the western side of Santa Maria, probably due to the scarcity of sediments and faults there are no significant scars at the shelf edge. 463

464

# 465 **5.3** New insights on the evolution of Santa Maria Island from submarine morphologies

466

Here we discuss the evolution of Santa Maria Island, based on the analysis of the morphologic parameters of the shelf (erosive shelf edge depth, shelf width, and type of submarine morphologies, Figures 5-8 and Table 1).

470

#### 471 **5.3.1** Insights from the shelf width

Generally, on volcanic islands a correspondence between shelf width and age of volcanic products outcropping onland is found (Menard, 1983). The width/age relationship is associated to the fact that the shelf width increases with increasing time as shorelines retreat with prolonged exposure to marine erosion (Quartau et al., 2010, 2012, 2014; Romagnoli, 2013; Romagnoli et al., 2018).

477 At Santa Maria this relationship can be better observed between the two width/age extremes, 478 the northern shelf (up to c. 8.7 km), likely carved in the c. 5.8 Ma-old Anjos volcanic products, and

479 the eastern shelf (2.4 km), carved in the c. 4.1-3.5 Ma old products of the Pico Alto volcanic Complex (Figs. 2 and 4). However, the western shelf shows a dramatically lower width (up to c. 2.3) 480 km) with respect the northern one (up to c. 8.7 km), despite both sectors are carved in the Anjos 481 volcanic products and are similarly exposed to the same dominant waves. Remarkably, the 482 483 western sector shelf is generally narrower than the southern one (2.6 km, also carved in the Anjos 484 volcanic units) and with similar width to the eastern sector (2.4 km). This apparent incongruence 485 can be explained by the paleogeography of the island in its earlier stages of growth. The western 486 subaerial side of the island is interpreted as an uplifted shelf that, at the onset of the uplift (3.5 Ma), was entirely submerged (Fig. 11) as suggested by presently exposed subaerial marine 487 terraces (Ramalho et al., 2017; Ricchi et al., 2018). The width of the western shelf, if measured 488 from the paleo-coastline attributed to 3.5 Ma BP is between 5.3 km and 8.5 km (Fig. 11). This 489 490 explains why the western shelf, today is not wider than the leeward and younger southeastern 491 and eastern shelves. However, it does not explain why the northern is presently much wider than 492 the western shelf. Given that both northern and western shelves are exposed to similar wave 493 energy they should have similar widths (Figs. 5 and 6 and Table 1ESM).

494

495 **5.3.2** Insights from the shelf edge depth

The analysis of the shelf edge depth might also help us in our understanding (Figs. 5-8). The erosive shelf edge depth is related to sea level coeval to the shelf formation; therefore this morphological feature can be used as a marker of palaeo sea levels, and thus be used to assess subsequent vertical movements affecting the volcanic edifices (Quartau et al. 2014; 2016, Romagnoli et al., 2018, Lucchi et al., 2019). Since glacial eustatic sea level never reached depths below 125 m in the last 3.5 Ma (Bintanja and van de Wal, 2008; de Boer et al., 2010), so any part of the shelf that is presently located below this depth should have experienced net subsidence

503 (Quartau et al., 2014; Quartau et al., 2015b; 2016; Quartau et al., 2018b; Romagnoli et al., 2018). Given that Santa Maria was affected by early subsidence prior to a reversal into an uplift trend at 504 ~3.5 Ma (Ramalho et al., 2017), the portions of the shelf exhibiting the deepest erosive edges can 505 be considered the oldest since they underwent the largest submergence after formation. 506 507 According to the history of vertical movements reconstructed for Santa Maria (Ramalho et al., 508 2017; Ricchi et al., 2018), the northern shelf sector, being the deepest of all (erosive shelf edge 509 depth down to -170 m, Figs. 5a and 5b), can be thus considered the oldest in formation. Deep (>-510 125 m) shelf edges have been also identified offshore the western sector of the island. However, in this shelf sector the maximum edge depth is 154 m (Figs. 5 and 6), indicating that it has suffered 511 less subsidence than the northern shelf. Being likely carved in the products of the Anjos Volcanic 512 513 Complex (Fig. 2) this shelf sector might be younger than the northern portion of the shelf.

Along the southern shelf sector, the shelf exhibits a more complex morphology and the shelf edge depth has a strongly irregular trend mostly around 80/100 m and being locally at 50/60 m (Fig. 7). The deeper shelf edge offshore Ponta do Castelo (147 m) is in agreement with the inferred subsidence processes experienced by the island and the relatively old age of this shelf sector. It is comparable to the maximum depth along the western sector (154 m, Fig. 6), suggesting that the southeastern portion of Santa Maria could be as old as the western portion of the island if they had been exposed to the same vertical movements.

The eastern shelf shows shelf edge depths not deeper than 100 m, contrary to the remaining sectors that show depths below 125 m. This is because when Pico Alto volcanism ended, the island was on a turning point in terms of vertical movements (see Fig. 6 of Ramalho et al., 2017). It was the beginning of the uplift trend, so the erosional shelf edge of this sector never experienced subsidence and is now always well above 125 m.

#### 527 **5.3.3 Insights from submarine morphologies**

## 528 5.3.3.1 Northern shelf sector

The morphology of the NW inner shelf shows, from Baía do Mar da Barca to Baía da Cré, a ~3 km wide belt of flat rocky outcrops that is very similar to those that can be seen on the western shelf (Figs. 5a and 6a). This suggests that, most likely, this area was carved on the Anjos Volcanic Complex. We believe that an earlier part of the island, probably older than the products of the Anjos Volcanic Complex, lies offshore those rocky outcrops on the northern shelf. This limit cannot be followed eastwards due to the profusion of eroded prominent reliefs existing in this portion of the shelf.

From Baía da Cré to the east, all rocky features are interpreted as volcanic cones or other type 536 of eruptive centers, albeit being of different age and having experienced different degrees and 537 538 styles of differential erosion. Type (a) described in section 4.1.1 are interpreted either as volcanic 539 necks exposed by differential erosion, or also as pillow mounds. Type (b), with their concentric 540 nature, are interpreted as old, well-consolidated strombolian cinder-and-scoria cones exposed and truncated by marine erosion. It is unlikely these are recent surtseyan cones, given that surtseyan 541 542 cones are more friable and typically erode to a flat platform (notwithstanding some concentric 543 features), as the recent surveys of Surtla, Jólnir and Syrtlingur (Surtsey, Iceland; see Romagnoli & Jacobson, 2015) suggest. These cones are probably contemporary of either the Porto Formation or 544 545 the Anjos Volcanic Sequence, given their inferred subaerial nature and the fact that the only time 546 relative sea level was low enough to allow their formation was during the period between 5.8 and 547 5.3 Ma (see Figs. 6 and 7 of Ramalho et al. 2017). In a similar fashion, type (c) cones are 548 interpreted as subaerial cones buried by effusive sequences, which are more erosion-resistant 549 than the softer inner core of the cinder cones. Given this, and the fact that these cones were 550 buried by effusive sequences that are in continuity with the Anjos Volcanic sequence onshore, and

given that they do not exhibit any of the prominent concentric morphologies typical of type (b) cones, these morphologies are interpreted to correspond to eroded surtseyan cones of the Cabrestantes Formation. Finally, type (d) cones, given their very pristine ring-like morphologies, with very low degrees of erosion, are interpreted as very young volcanic cones, perhaps younger than any structure found onshore.

The NNW-SSE submarine rocky ridge (Fig. 5a) that extends offshore Ilhéu das Lagoínhas is in clear alignment with both the dykes that cross-cut the Anjos Volcanic Complex along the adjacent coastline and the Pico Alto's ridge. This suggests long-lived fissural volcanism roughly oriented N-S in this part of the island, both during the Anjos and Pico Alto's volcanic phases. This inference is also supported by the presence of elongated morphologies interpreted as dykes exposed by differential erosion (Figs. 4 and 1ESM).

562 The pillow-lava cone of Baixa do Ambrósio at the NW edge of the northern shelf (Fig. 5a) is not directly associated to any of the volcanic complexes making up Santa Maria Island. The lack of 563 dated samples prevented us to make any inference regarding its earlier growth. The depth of its 564 top (c. 50 m) suggests that probably it is younger with respect the rest of the northern shelf 565 566 sector; otherwise it would be a more flat-topped cone. On its flanks, Ricchi et al. (2018) recognized 567 a set of submerged marine terraces from 70/80 m to 120/140 m. The shallowest terraces are considered to be the oldest and were likely carved at least since 866 ka (Ricchi et al., 2018), 568 569 suggesting that the cone is older than this age.

570 We therefore infer that all these eroded cones are covering a "proto-Santa Maria Island" on 571 which the outer northern shelf was carved. This shelf is much wider than the western shelf and the 572 shelf edge deeper. The occurrence of an older volcanic edifice offshore the northern portion of the 573 island had already been proposed by Ramalho et al. (2017), based on the wide northern shelf. But

now, an improved insight can be taken from the interpretation of the deeper shelf edge and highresolution images of the submarine morphologies.

576

577 5.3.3.2 Western shelf sector

The morphology of this sector shows a ~1-2 km wide belt of flat rocky outcrops that is very similar to those that can be seen on the NW inner shelf (Figs. 5a and 6a). This suggests that, most likely, this area was carved on Anjos Volcanic Complex.

581

582 5.3.3.3 Southern shelf sector

This shelf sector shows a ~0.7-2.3 km wide belt of flat rocky outcrops (Fig. 7a) that are very similar to those on the NW inner shelf and western shelf (Figs. 5a and 6a). These rocky outcrops are separated by sedimentary bottoms that correspond to fault controlled basins. The similarity with the NW inner shelf and western shelf suggests that, most likely, this area was also carved on Anjos Volcanic Complex.

588

589 5.3.3.4 Eastern shelf sector

590 The association of the eastern shelf sector to the Pico Alto products is supported by the outer shelf morphology, which fin-like outward dipping rocky surfaces (Fig. 8 and seismic profile hh' in 591 592 Fig. 8ESM), interpreted as preserved substructural surfaces of the foreset units of Gilbert-type lava-fed deltas. Effectively, these morphologies are in structural continuity with the eastward-593 dipping foreset units of the lava-fed delta sequences that crop out all along the eastern coast of 594 595 the island (e.g. see Figs. 3G and 3H of Ramalho et al. 2017), which extend from the passage zone 596 (between subaerial topset and submarine foreset units) currently located up to c. 130 m in 597 elevation to present-day sea level (Ramalho et al., 2017). These observations therefore suggest

that the eastern submarine upper slope of Santa Maria corresponds to the outer submarine slopes
of former Gilbert-type lava-fed deltas, i.e. are constructional morphologies contemporaneous of
the eastward expansion of the Pico Alto volcanic edifice, prior to uplift and erosion of the shelf.

# 602 6. Conclusions

603

The morphologic analysis of the shelf surrounding Santa Maria Island complemented with the known geological history of the subaerial island allowed us to improve our understanding of the island's evolution and the processes which have shaped its shallow submerged portions.

The shelves surrounding Santa Maria are mostly sediment-starved. Around 50% of the shelf is 607 rocky and sediment thickness is on average less than 2 m. A few depocenters exist, however, with 608 609 thicknesses up to 14 m, but these are clearly fault-controlled basins. The lack of thicker and older 610 submarine deposits that should have accumulated by erosion of the subaerial flanks of the island are related with the high wave energy in the Azores and the cyclicity of sea-level changes. High 611 wave energy tends to transport sediments offshore and during sea level falls, sediments that 612 613 normally accumulate on the shelf are swept offshore until they are lost to the slopes as they cross the shelf edge. In the present-day, and since shelves are wide, waves tend to attenuate their 614 energy, eroding the cliffs with less efficiency. In addition, it has lower cliffs and rains less, when 615 616 compared to other studied islands, so fewer sediments are expected to be delivered from the island to the coastline. Thus, not only fewer sediments are produced, but also larger 617 accommodation spaces (wider shelves) preclude the existence of thicker submarine deposits, as 618 found on other studied volcanic islands. 619

620 Small mass-wasting features were found on the shelf edge of Santa Maria. These are mostly 621 restricted to the northern and southern shelves, with only one evident scar on the eastern shelf.

However, different processes appear to be responsible for such features. On the southern and eastern shelf sectors, is mostly tectonics that controls the location of mass-wasting morphologies, with faults extending offshore and reaching the shelf break. On the northern shelf, the predisposing factor is mostly the accumulation of unconsolidated sediments near the edge.

626 Based on the analysis of the shelf geomorphic features (shelf width, erosive shelf edge depth 627 and morphology) and the correlation with volcanic sequences outcropping on the island, we were 628 able to improve our understanding of Santa Maria geological evolution. We suggest that given the 629 wide extension of the northern shelf, its outer part belongs to an older and undated volcanic sequence of Santa Maria that predated the Cabrestantes and Porto Formations. The mid-to inner 630 northern shelf, as well as the western and southern shelves were probably carved mainly on the 631 products of Anjos Volcanic Complex, whilst the eastern shelf results from erosion of the products 632 633 of Pico Alto Volcanic Complex. The present-day narrow width of the western shelf, when compared with the southern and eastern shelves, which are more protected to wave erosion, can 634 635 be explained by the uplift mechanism that raised part of the western shelf, originally ~5.3-8.5 km wide, leaving now a ~2 km wide shelf. 636

This work shows that, despite the fact that Santa Maria Island is much older than islands addressed in previous shelf studies, the various geological processes acting on the shelf can still leave significant morphological imprints that provide us important knowledge about the evolutionary history of this island.

641

# 642 **7. Acknowledgments**

The acquisition of multibeam bathymetry and seismic reflection profiles were funded by Fundação
 para a Ciência e a Tecnologia (FCT) through the PLATMAR project (*Development of volcanic island shelves: insights from Sta. Maria Island and implications on hazard assessment, habitat mapping*

and marine aggregates management, PTDC/GEO-GEO/0051/2014). The authors are grateful to the
crew of R/V Arquipélago and Pedro Afonso for all their help in the preparation and execution of
the geophysical survey. R. Ramalho acknowledges its "Investigador FCT" contract IF/01641/2015
funded by FCT. This study has been developed in the framework of the PhD thesis "Insular shelves
as a tool for reconstructing the evolution of volcanic islands" (A. Ricchi, University of Bologna).

## 652 **8. References**

653

- Agencia Estatal de Meteorología & Instituto de Meteorologia de Portugal, 2012. Climate atlas of
  the archipelagos of the Canary Islands, Madeira and the Azores. p. 78.
- Andrade, C., Trigo, R.M., Freitas, M.C., Gallego, M.C., Borges, P., Ramos, A.M., 2008. Comparing
- 657 historic records of storm frequency and the North Atlantic Oscillation (NAO) chronology for the

Azores region. The Holocene 18, 745-754.

- Ávila, S.P., Madeira, P., Silva, C.M.d., Cachão, M., Quartau, R., Landau, B., Martins, A.M.d.F., 2008.
  Local disappearance of bivalves in the Azores during the last glaciation. Journal of Quaternary
  Science 23, 777-785.
- 662 Bintanja, R., van de Wal, R.S.W., 2008. North American ice-sheet dynamics and the onset of
- 663 100,000-year glacial cycles. Nature 454, 869-872.
- 664 Casalbore, D., Bosman, A., Romagnoli, C., Di Filippo, M., & Chiocci, F.L., 2016a. Morphology of
- Lipari offshore (Southern Tyrrhenian Sea). Journal of Maps 12, 77-86.
- 666 Casalbore, D., Bosman, A., Martorelli, E., Sposato, A., Chiocci, F.L., 2016b. Mass wasting features
- on the submarine flanks of Ventotene volcanic edifice (Tyrrhenian Sea, Italy): In Submarine Mass

- Movements and Their Consequences, (Eds Krastel et al), Advances in Natural and Technological
  Hazards Research, 37, 285-293, DOI 101007/978-3-319-00972-8 25
- 670 Casalbore, D., Falese, F., Martorelli, E., Romagnoli, C., Chiocci, F.L., 2017. Submarine depositional
- 671 terraces in the Tyrrhenian Sea as a proxy for paleo-sea level reconstruction: Problems and
- 672 perspective. Quatern. Int. 439, 169-180.
- 673 Casalbore, D., Romagnoli, C., Adami, C., Bosman, A., Falese, F., Ricchi, A., Chiocci, F., 2018.
- 674 Submarine Depositional Terraces at Salina Island (Southern Tyrrhenian Sea) and Implications on
- the Late-Quaternary Evolution of the Insular Shelf. Geosciences 8, 20.
- 676 Casalbore, D., Romagnoli, C., Bosman, A., Chiocci, F.L., 2011. Potential tsunamigenic landslides at
- 677 Stromboli Volcano (Italy): Insight from marine DEM analysis. Geomorphology 126, 42-50.
- 678 Casalbore, D., Romagnoli, C., Pimentel, A., Quartau, R., Casas, D., Ercilla, G., Hipólito, A., Sposato,
- A., Chiocci, F.L., 2015. Volcanic, tectonic and mass-wasting processes offshore Terceira Island
- 680 (Azores) revealed by high-resolution seafloor mapping. Bull. Volc. 77, 1-19.
- 681 Chiocci, F.L., Romagnoli, C., 2004. Submerged depositional terraces in the Aeolian Islands (Sicily),
- in: Chiocci, F.L., D'Angelo, S., Romagnoli, C. (Eds.), Atlas of submerged depositional terraces along
- the Italian coasts. Memorie Descrittive della Carta Geologica d'Italia. APAT, Roma, pp. 81-114.
- 684 Costa, A.C.G., Hildenbrand, A., Marques, F.O., Sibrant, A.L.R., Santos de Campos, A., 2015.
- 685 Catastrophic flank collapses and slumping in Pico Island during the last 130 kyr (Pico-Faial ridge,
- Azores Triple Junction). J. Volcanol. Geotherm. Res. 302, 33-46.

de Boer, B., Van De Wal, R.S.W., Bintanja, R., Lourens, L.J., Tuenter, E., 2010. Cenozoic global icevolume and temperature simulations with 1-D ice-sheet models forced by benthic 18O records.
Ann. Glaciol. 51, 23-33.

690 Dee, D. P., S. Uppala, A. Simmons, P. Berrisford, P. Poli, S. Kobayashi, U. Andrae, M. Balmaseda, G.

Balsamo, and P. Bauer., 2011. The ERA-Interim reanalysis: Configuration and performance of the

data assimilation system. Q. J. Roy. Meteor. Soc., 137 (656), 553-597.

693 Deplus, C., Le Friant, A., Boudon, G., Komorowski, J.-C., Villemant, B., Harford, C.L., Segoufin, J.,

694 Cheminée, J.-L., 2001. Submarine evidence for large-scale debris avalanches in the Lesser Antilles

695 Arc. Earth Planet. Sci. Lett. 192, 145-157.

696 Fernandes, J., 2004. Caracterização Climática das ilhas de S Miguel e Santa Maria com base no

697 modelo CIELO. BSc thesis. Dept. de Ciências Agrárias. Universidade dos Açores, Angra do
698 Heroísmo, p. 210.

699 Field, M.E., Roy, P.S., 1984. Offshore transport and sand-body formation; evidence from a steep,

high energy shoreface, southeastern Australia. J. Sediment. Res. 54, 1292-1302.

701 Gente, P., Dyment, J., Maia, M., Goslin, J., 2003. Interaction between the Mid-Atlantic Ridge and

the Azores hot spot during the last 85 Myr: Emplacement and rifting of the hot spot-derived

703 plateaus. Geochem. Geophys. Geosyst. 4, 8514.

Hampton, M.A., Lee, H.J., Locat, J., 1996. Submarine landslides. Rev. Geophys. 34, 33-59.

Hatton, B. 2019. Hurricane Lorenzo batters mid-Atlantic Azores Islands. ABC News. Associated
Press. Retrieved February 9, 2020.

- 707 URL:https://abcnews.go.com/International/wireStory/category-hurricane-batters-mid-atlantic-708 azores-islands-65997843.
- 709 Hernández-Molina, F.J., Fernández-Salas, L.M., Lobo, F., Somoza, L., Díaz-del-Río, V., Alveirinho
- Dias, J.M., 2000. The infralittoral prograding wedge: a new large-scale progradational sedimentary
- body in shallow marine environments. Geo-Mar. Lett. 20, 109-117.
- Hipólito, A., Madeira, J., Carmo, R., Gaspar, J.L., 2013. Neotectonics of Graciosa island (Azores): a
  contribution to seismic hazard assessment of a volcanic area in a complex geodynamic setting.
  Ann. Geophys. 56, 1-18.
- Krastel, S., Schmincke, H.-U., Jacobs, C.L., 2001. Formation of submarine canyons on the flanks of
  the Canary Islands. Geo-Mar. Lett. 20, 160-167.
- Lebas, E., Friant, A.L., Deplus, C., Voogd, B., 2018. Understanding the Evolution of an Oceanic
- 718 Intraplate Volcano From Seismic Reflection Data: A New Model for La Réunion, Indian Ocean. J.

719 Geophys. Res. Solid Earth 123, 1035-1059.

- Llanes, P., Herrera, R., Gómez, M., Muñoz, A., Acosta, J., Uchupi, E., Smith, D., 2009. Geological
- evolution of the volcanic island La Gomera, Canary Islands, from analysis of its geomorphology.

722 Mar. Geol. 264, 123-139.

- Locat, J., Lee, H.J., 2002. Submarine landslides: advances and challenges. Can. Geotech. J. 39, 193212.
- Lourenço, N., Miranda, J.M., Luís, J.F., Ribeiro, A., Mendes Victor, L.A., Madeira, J., Needham, H.D.,
- 1998. Morpho-tectonic analysis of the Azores Volcanic Plateau from a new bathymetric
- 727 compilation of the area. Mar. Geophys. Res. 20, 141-156.
  - 1

Lucchi, F., Ricchi, A., Romagnoli, C., Casalbore, D., Quartau, R., 2019. Late Quaternary paleo sea
level geomorphological markers of opposite vertical movements at Salina volcanic island (Aeolian
Arc). Earth Surf. Proc. Land., , https://doi.org/10.1002/esp.4651.

Madeira, J., 1986. Geologia estrutural e enquadramento geotectónico da ilha de Santa Maria, MSc
thesis. Dept. de Geologia. Universidade de Lisboa, Lisboa, p. 107.

733 Madeira, J., Brum da Silveira, A., Hipólito, A., Carmo, R., 2015. Active tectonics in the central and

eastern Azores islands along the Eurasia–Nubia boundary: a review, in: Gaspar, J.L., Guest, J.E.,

Duncan, A.M., Barriga, F.J.A.S., Chester, D.K. (Eds.), Geological Society, London, Memoirs, 44, pp.
15-32.

Marques, F.O., Catalão, J.C., DeMets, C., Costa, A.C.G., Hildenbrand, A., 2013. GPS and tectonic
evidence for a diffuse plate boundary at the Azores Triple Junction. Earth Planet. Sci. Lett. 381,
177-187.

Masson, D.G., Le Bas, T.P., Grevemeyer, I., Weinrebe, W., 2008. Flank collapse and large-scale
landsliding in the Cape Verde Islands, off West Africa. Geochem. Geophys. Geosyst. 9,

742 doi:10.1029/2008GC001983.

743 Masson, D.G., Watts, A.B., Gee, M.J.R., Urgeles, R., Mitchell, N.C., Le Bas, T.P., Canals, M., 2002.

Slope failures on the flanks of the western Canary Islands. Earth-Sci. Rev. 57, 1-35.

745 Mazuel, A., Sisavath, E., Babonneau, N., Jorry, S.J., Bachèlery, P., Delacourt, C., 2016. Turbidity

current activity along the flanks of a volcanic edifice: The Mafate volcaniclastic complex, La

747 Réunion Island, Indian Ocean. Sed. Geol. 335, 34-50.

- 748 Meireles, R., Quartau, R., Ramalho, R.S., Rebelo, A.C., Madeira, J., Zanon, V., Ávila, S.P., 2013.
- 749 Depositional processes on oceanic island shelves evidence from storm-generated Neogene
- deposits from the mid-North Atlantic. Sedimentology 60, 1769-1785.
- 751 Menard, H.W., 1983. Insular erosion, isostasy, and subsidence. Science 220, 913-918.
- 752 Miranda, J.M., Luís, J., Lourenço, N., 2018. The geophysical architecture of the Azores from
- magnetic data, in: Kueppers, U., Beier, C. (Eds.), Volcanoes of the Azores. Springer-Verlag Berlin
  Heidelberg, pp. 89-100.
- Mitchell, N.C., Dade, W.B., Masson, D.G., 2003. Erosion of the submarine flanks of the Canary
  Islands. J. Geophys. Res. 108, 3-1 3-11.
- 757 Mitchell, N.C., Masson, D.G., Watts, A.B., Gee, M.J.R., Urgeles, R., 2002. The morphology of the
- flanks of volcanic ocean islands: A comparative study of the Canary and Hawaiian hotspot islands.

759 J. Volcanol. Geotherm. Res. 115, 83-107.

- 760 Montanaro, C., Beget, J., 2011. Volcano collapse along the Aleutian Ridge (western Aleutian Arc).
- 761 Nat. Hazards Earth Syst. Sci. 11, 715-730.
- Moore, J.G., Clague, D.A., Holcomb, R.T., Lipman, P.W., Normark, W.R., Torresan, M.E., 1989.
- Prodigious submarine landslides on the Hawaiian Ridge. J. Geophys. Res.: Solid Earth 94, 1746517484.
- Moore, J.G., Normark, W.R., Holcomb, R.T., 1994. Giant Hawaiian Landslides. Annu. Rev. Earth Pl.
  Sc. 22, 119-144.
- Oehler, J.-F., Lénat, J.-F., Labazuy, P., 2008. Growth and collapse of the Reunion Island volcanoes.
  Bull. Volc. 70, 717-742.

769 Quartau, R., Hipólito, A., Mitchell, N.C., Gaspar, J.L., Brandão, F., 2015a. Comment on

"Construction and destruction of a volcanic island developed inside an oceanic rift: Graciosa

771 Island, Terceira Rift, Azores" by Sibrant et al. (2014) and proposal of a new model for Graciosa

geological evolution [J. Volcanol. Geotherm. Res. 284 (2014) 32-45]. J. Volcanol. Geotherm. Res.

773 303, 146-156.

Quartau, R., Hipólito, A., Romagnoli, C., Casalbore, D., Madeira, J., Tempera, F., Roque, C., Chiocci,
F.L., 2014. The morphology of insular shelves as a key for understanding the geological evolution
of volcanic islands: Insights from Terceira Island (Azores). Geochem. Geophys. Geosyst. 15, 1801–
1826.

Quartau, R., Madeira, J., Mitchell, N.C., Tempera, F., Silva, P.F., Brandão, F., 2015b. The insular
shelves of the Faial-Pico Ridge: a morphological record of its geologic evolution (Azores
archipelago). Geochem. Geophys. Geosyst. 16, 1401–1420.

781 Quartau, R., Madeira, J., Mitchell, N.C., Tempera, F., Silva, P.F., Brandão, F., 2016. Reply to

comment by Marques et al. on "The insular shelves of the Faial-Pico Ridge (Azores archipelago): A

morphological record of its evolution". Geochem. Geophys. Geosyst. 17, 633-641.

48]. J. Volcanol. Geotherm. Res. 255, 124-126.

Quartau, R., Mitchell, N.C., 2013. Comment on "Reconstructing the architectural evolution of volcanic islands from combined K/Ar, morphologic, tectonic, and magnetic data: The Faial Island example (Azores)" by Hildenbrand et al. (2012) [J. Volcanol. Geotherm. Res. 241-242 (2012) 39-

Quartau, R., Ramalho, R.S., Madeira, J., Santos, R., Rodrigues, A., Roque, C., Carrara, G., Brum da
 Silveira, A., 2018a. Gravitational, erosional and depositional processes on volcanic ocean islands:

1

790 Insights from the submarine morphology of Madeira archipelago. Earth Planet. Sci. Lett. 482, 288-791 299.

792 Quartau, R., Tempera, F., Mitchell, N.C., Pinheiro, L.M., Duarte, H., Brito, P.O., Bates, R., Monteiro,

J.H., 2012. Morphology of the Faial Island shelf (Azores): The interplay between volcanic,

rosional, depositional, tectonic and mass-wasting processes. Geochem. Geophys. Geosyst., 13,

795 Q04012, doi:10.1029/2011GC003987.

796 Quartau, R., Trenhaile, A.S., Mitchell, N.C., Tempera, F., 2010. Development of volcanic insular

shelves: Insights from observations and modelling of Faial Island in the Azores Archipelago. Mar.

798 Geol. 275, 66-83.

Quartau, R., Trenhaile, A.S., Ramalho, R.S., Mitchell, N.C., 2018b. The role of subsidence in shelf
widening around ocean island volcanoes: Insights from observed morphology and modeling. Earth
Planet. Sci. Lett. 498, 408-417.

802 Ramalho, R.S., Helffrich, G., Madeira, J., Cosca, M., Thomas, C., Quartau, R., Hipólito, A., Rovere,

A., Hearty, P.J., Ávila, S.P., 2017. Emergence and evolution of Santa Maria Island (Azores)—The

conundrum of uplifted islands revisited. Geol. Soc. Am. Bull. 129, 372-390.

805 Ramalho, R.S., Quartau, R., Hóskuldsson, A., Madeira, J., Ventura da Cruz, J., Rodrigues, A., in

806 press. Evidence for late Pleistocene volcanism at Santa Maria Island, Azores? J Volcanol Geoth Res.

Ramalho, R.S., Quartau, R., Trenhaile, A.S., Mitchell, N.C., Woodroffe, C.D., Ávila, S.P., 2013.

808 Coastal evolution on volcanic oceanic islands: A complex interplay between volcanism, erosion,

sedimentation, sea-level change and biogenic production. Earth-Sci. Rev. 127, 140-170.

Ricchi, A., Quartau, R., Ramalho, R.S., Romagnoli, C., Casalbore, D., Ventura da Cruz, J., Fradique,
C., Vinhas, A., 2018. Marine terraces development on reefless volcanic islands: new insights from
high-resolution marine geophysical data offshore Santa Maria Island (Azores Archipelago). Mar.
Geol. 406, 42-56.

- 814 Romagnoli, C., 2013. Characteristics and morphological evolution of the Aeolian volcanoes from
- the study of submarine portions, in: Lucchi, F., Peccerillo, A., Keller, J., Tranne, C.A., Rossi, P.L.

816 (Eds.), Geological Society, London, Memoirs, 37, pp. 13-26.

817 Romagnoli, C., Belvisi, V., Innangi, S., Di Martino, G., Tonielli, R., 2020. New insights on the

evolution of the Linosa volcano (Sicily Channel) from the study of its submarine portions. MarineGeology 419, 106060.

Romagnoli C., Casalbore D., Bosman A., Braga R., Chiocci F.L. (2013). Submarine structure of
Vulcano Volcano (Aeolian Islands) revealed by high-resolution bathymetry and seismo-acoustic

data. Marine Geology 338, 30–45.

823 Romagnoli, C., Casalbore, D., Chiocci, F.L., Bosman, A., 2009a. Offshore evidence of large-scale

lateral collapses on the eastern flank of Stromboli, Italy, due to structurally-controlled, bilateral
flank instability. Mar. Geol. 262, 1-13.

826 Romagnoli, C., Casalbore, D., Ricchi, A., Lucchi, F., Quartau, R., Bosman, A., Tranne, C.A., Chiocci,

827 F.L., 2018. Morpho-bathymetric and seismo-stratigraphic analysis of the insular shelf of Salina

828 (Aeolian archipelago) to unveil its Late-Quaternary geological evolution. Mar. Geol. 395, 133-151.

829 Romagnoli, C., Kokelaar, P., Casalbore, D., Chiocci, F.L., 2009b. Lateral collapses and active

sedimentary processes on the northwestern flank of Stromboli volcano, Italy. Mar. Geol. 265, 101-

831 119.

- Rusu, L., Guedes Soares, C., 2012. Wave energy assessments in the Azores islands. Renew. Energy
  45, 183-196.
- 834 Serralheiro, A., Alves, C.A.M., Forjaz, V.H., Rodrigues, B., 1987. Carta Vulcanológica dos Açores.
- 835 Ilha de Santa Maria na (folhas 1 e 2), scale 1:15,000. Serviço Regional de Protecção Civil dos
- Açores, Universidade dos Açores and Centro de Vulcanologia, Ponta Delgada.
- Serralheiro, A., Madeira, J., 1993. Stratigraphy and geochronology of Santa Maria Island (Azores).
  Açoreana 7, 575-592.
- 839 Sibrant, A.L.R., Hildenbrand, A., Marques, F.O., Costa, A.C.G., 2015. Volcano-tectonic evolution of
- 840 the Santa Maria Island (Azores): Implications for paleostress evolution at the western Eurasia-
- Nubia plate boundary. J. Volcanol. Geotherm. Res. 291, 49-62.
- Taylor, B., 2019. Shoreline slope breaks revise understanding of Hawaiian shield volcanoes
- evolution. Geochem. Geophys. Geosyst. 20, 4025-4045.
- Thouret, J.C., 1999. Volcanic geomorphology-an overview. Earth-Sci. Rev. 47, 95-131.
- Tsutsui, B., Campbell, J.F., Coulbourn, W.T., 1987. Storm-generated, episodic sediment movements
  off Kahe Point, Oahu, Hawaii. Mar. Geol. 76, 281-299.
- Uppala, S. M., P. Kållberg, A. Simmons, U. Andrae, V. d. Bechtold, M. Fiorino, J. Gibson, J. Haseler,
- A. Hernandez, and G. Kelly., 2005. The ERA-40 re-analysis. Q. J. Roy. Meteor. Soc. 131 (612):29613012.
- Zbyszewski, G., Ferreira, O.V., 1960. Carta Geológica de Portugal—Ilha de Santa Maria (Açores),
- scale 1:50,000. Serviços Geológicos de Portugal, Lisboa.
  - 1

- Zhao, Z., Mitchell, N.C., Quartau, R., Tempera, F. and Bricheno, L., 2019. Submarine Platform
- 853 Development by Erosion of a Surtseyan Cone at Capelinhos, Faial Island, Azores. Earth Surf. Proc.
- Land. 44(15), 2982-3006.

855

856

858 Fig. 1. Shaded relief map illustrating the geodynamic setting of the islands (western, central and eastern groups) in the Azores Archipelago (islands in white, Santa Maria Island in yellow) and how 859 they straddle the triple junction between the North American (NA), Eurasian (EU) and Nubian (NU) 860 lithospheric plates. MAR: Mid-Atlantic Ridge; PAR: Princess Alice Rift; EAFZ: Eastern Azores 861 Fracture Zone; TR: Terceira Rift; GF: Gloria Fault. Grey shading represents the deformation zone of 862 863 Terceira rift and dashed lines of the inactive Princess Alice Rift. The upper-right inset shows the 864 geographical setting of the Azores Archipelago within the Northern Atlantic Ocean. The bathymetry is derived from the EMODNET web portal (http://portal.emodnet-bathymetry.eu/) 865 and tectonic structures adapted from Miranda et al. (2018). 866

867

Fig. 2. (a) Geological map of Santa Maria Island (modified after Ramalho et al., 2017), and (b)
WNW–ESE oriented cross-section. Underlying digital elevation model (DEM) is generated from a
1/5000 scale digital altimetric database from Secretaria Regional do Turismo e Transportes of the
Azores Government. Note that the Porto Formation is not visible on the map/section at this scale.

872

Fig. 3. Dataset used for this study, showing the seismic reflection profiles acquired during the PLATMAR 1/2016 cruise. The multibeam survey covered the area shown by the seismic grid. Blue dots represent location of sound speed profiles collected during the multibeam survey. Bathymetry shown in this figure (~200m resolution) was derived from the EMODNET web portal (http:// portal.emodnet-bathymetry.eu); topography is the same as the one shown in Figure 2.

878

Fig. 4. Shaded relief map of the multibeam bathymetry. Topography is the same as the one shownin Figure 2.

881

Fig. 5. (a) Shaded relief map of the northern shelf sector showing the erosive shelf edge, headwalls scars, extension of tectonic features and volcanic cones. The seismic profile aa' is shown in Fig. 2ESM and seismic profiles bb' and cc' are shown in Fig. 3ESM. (b) Variation of shelf width and erosive shelf edge depth along the northern sector from west to east (from A to B). Shelf width was only measured on profiles perpendicular to the north coast.

887

Fig. 6. (a) Shaded relief map of the western shelf sector showing the erosive shelf edge, headwalls scars, and extension of tectonic features. The seismic profile dd' is shown in Fig. 4ESM (b) Variation of shelf width and erosive shelf edge depth along the western sector from north to south (from C to D).

892

Fig. 7. (a) Shaded relief map of the southern shelf sector showing the erosive shelf edge, headwalls scars, and extension of tectonic features. The seismic profiles ee', ff' and gg' are shown respectively in Figs. 5ESM, 6ESM and 7ESM (b) Variation of shelf width and erosive shelf edge depth along the southern sector from west to east (from E to F).

897

Fig. 8. (a) Shaded relief map of the eastern shelf sector showing the erosive shelf edge, headwalls scars, and extension of tectonic features. The seismic profiles hh' and ii' are shown in Figs. 8ESM and 9ESM, respectively. (b) Variation of shelf width and erosive shelf edge depth along the eastern sector from north to south (from G to H).

902

Fig. 9. Thickness map (in metres) of the unconsolidated sediments of the shelf surrounding SantaMaria Island.

905

- 906 Fig. 10. Offshore significant wave heights in the Santa Maria area derived from ERA 40 and ERA
- 907 Interim (Uppala et al., 2005; Dee et al., 2011)
- 908
- 909 Fig. 11. Reconstruction of the likely extension of the western shelf of Santa Maria Island 3.5 Ma
- ago, before being partially raised. Color scale of topography is the same as Figure 3.

1 Table 1 – Extension of the rocky and sedimentary seafloor areas (in km<sup>2</sup> and %) of each shelf sector; measured volume of

	Are	a (km²)		ې) Area	%)	Volume of sediments (km <sup>3</sup> )	Volume sediments/Total shelf area	
 Sectors	Sediments	Rocks	Total	Sediments	Rocks			
Northern	36.820	36.240	73.060	50.397	49.603	0.515	0.007	
Western	3.300	9.900	13.200	25.000	75.000	0.037	0.003	
Southern	9.450	9.580	19.030	49.658	50.342	0.221	0.012	
Eastern	6.530	5.670	12.200	53.525	46.475	0.187	0.015	

2 sediments in each sector; and ratio of volume of sediments of each shelf sector versus its total area.

3

3	Fig. 1ESM. High resolution (1200 dpi) shaded relief map of the multibeam bathymetry. Topography
4	is the same as the one shown in Figure 2.
5	
6	Fig. 2ESM. Boomer seismic profile aa' showing the shelf morphology off the NW sector of Santa
7	Maria. Location of the seismic profile in Figs. 5a and 9.
8	
9	Fig. 3ESM. Boomer seismic profiles bb' and cc' showing the shelf morphology controlled by faults
10	off the N and NE sector of Santa Maria. Location of the seismic profiles in Fig. 5a and 9.
11	
12	Fig. 4ESM. Boomer seismic profile dd' showing the shelf morphology off the western sector of
13	Santa Maria. Location of the seismic profile in Fig. 6a.
14	
15	Fig. 5ESM. Boomer seismic profile ee' showing the shelf morphology off the southern sector of
16	Santa Maria. Location of the seismic profile in Fig. 7a.
17	
18	Fig. 6ESM. Boomer seismic profile ff' showing the shelf morphology controlled by faults off the
19	southern sector of Santa Maria. Location of the seismic profile in Fig. 7a.
20	
21	Fig. 7ESM. Boomer seismic profile gg' showing the shelf morphology off the southern sector of
22	Santa Maria. Location of the seismic profile in Fig. 7a.
23	

- Fig. 8ESM. Boomer seismic profiles hh' showing the shelf morphology off the eastern sector of
- 25 Santa Maria. Location of the seismic profiles in Fig. 8a.
- 26
- Fig. 9ESM. Boomer seismic profiles ii' showing the the shelf morphology controlled by faults off
- the eastern sector of Santa Maria. Location of the seismic profiles in Fig. 8a.

# 1 Table 1ESM - Six hourly significant wave height Hs for the quadrant directions and respective frequencies derived from ERA

Wave			Sum of percent	Average							
originating											
direction	0 - 1 m	1 -2 m	2 - 3 m	3 - 4 m	4 - 5 m	5 - 6 m	6 - 7 m	7 - 8 m	>= 8 m	inequency (70)	<i>m<sub>s</sub></i> (m)
Ν	0.99471	10.86626	5.20769	1.40490	0.32027	0.06699	0.02207	0.00590	0.00020	18.88899	1.922017
NE	0.56254	7.96483	2.96023	0.74180	0.15010	0.01982	0.00332	0.00111	0.00000	12.40375	1.830305
E	0.15145	2.15108	0.86000	0.29517	0.06035	0.00528	0.00000	0.00000	0.00000	3.52333	1.890301
SE	0.05794	0.67366	0.40322	0.13622	0.02203	0.00299	0.00000	0.00000	0.00000	1.29606	2.017559
S	0.09557	0.72673	0.59120	0.27056	0.05839	0.01118	0.00000	0.00000	0.00000	1.75363	2.204198
SW	0.19097	1.88260	1.55349	0.89456	0.34652	0.12226	0.02182	0.00577	0.00098	5.01897	2.453569
W	0.77025	8.88117	6.84695	3.62394	1.44261	0.54584	0.18249	0.06727	0.03071	22.39123	2.436865
NW	1.48884	17.00499	10.67882	3.78117	1.24923	0.35548	0.11542	0.03660	0.01347	34.72402	2.142374

# 2 40 and ERA Interim for the period between 1957 and 2019 (Uppala et al., 2005; Dee et al., 2011).

3







































a)

