

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Pastorino, P., Elia, A.C., Caldaroni, B., Menconi, V., Abete, M.C., Brizio, P., et al. (2020). Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps). *SCIENCE OF THE TOTAL ENVIRONMENT*, 715, 1-11 [10.1016/j.scitotenv.2020.136946].

Availability:

This version is available at: <https://hdl.handle.net/11585/758233> since: 2020-05-08

Published:

DOI: <http://doi.org/10.1016/j.scitotenv.2020.136946>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Manuscript Number: STOTEN-D-19-17682R2

Title: Oxidative stress ecology in brook trout (*Salvelinus fontinalis*)
from a high-mountain lake (Cottian Alps)

Article Type: Research Paper

Keywords: Alpine lakes, extreme ecosystems, oxidative stress biomarkers,
trace elements

Corresponding Author: Dr. Paolo Pastorino,

Corresponding Author's Institution: Istituto Zooprofilattico Sperimentale
del Piemonte, Liguria e Valle d'Aosta

First Author: Paolo Pastorino

Order of Authors: Paolo Pastorino; Antonia Concetta Elia; Barbara
Caldaroni; Vasco Menconi; Maria Cesarina Abete; Paola Brizio; Marco
Bertoli; Annalisa Zaccaroni; Magara Gabriele; Ambrosius Josef Martin
Dörr; Elisabetta Pizzul; Marino Prearo

Abstract: High-mountain lakes are pristine ecosystems characterized by extreme environmental conditions. The atmospheric transport of pollutants from lowlands may add further stress to organisms inhabiting these environments. We investigated the environmental stress pressure on brook trout (*Salvelinus fontinalis*) from a high-mountain lake in the Cottian Alps (Piedmont, northwest Italy). To do this, males and females of brook trout were sampled from Balma Lake in summer (August) and autumn (October) 2017 in order to assess the influence of trace elements accumulation and environmental parameters (physicochemical parameters and nutrient characteristics of water) on oxidative stress biomarkers. Bioaccumulation of Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se, and Zn and metallothionein levels were measured in muscle tissue of males and females. Liver, gills, kidney, and spleen tissue samples were analyzed for superoxide dismutase, catalase, total glutathione peroxidase, selenium-dependent glutathione peroxidase, glutathione reductase, and glutathione S-transferase activity. Analysis of environmental parameters showed changes in biomarker levels with seasonal variations. Water temperature was significantly higher in summer than autumn (Wilcoxon test; $p = 0.0078$), while pH was significantly higher in autumn than in summer (Wilcoxon test; $p = 0.0078$). Sex-related differences in oxidative stress biomarkers in tissues were unremarkable, whereas seasonal variability of oxidative stress biomarkers was observed, with major differences occurred for liver in summer and for gills, kidney, spleen and muscle in autumn. Positive correlations between environmental parameters and biomarkers were noted. Major fluctuations in water temperature, pH, Cu, Pb and Hg produced changes in biomarker levels; however, increased food intake during the ice-free season was probably the main factor that influenced changes in oxidative stress biomarker levels in brook trout in this extreme ecosystem.

Response to Reviewers: Response to Reviewer 5

Reviewer #5: In general, the authors improve the manuscript taking into account the comments/suggestions of the reviewers.
I recommend the publication of this study after some issues be addressed.
The aim of the study is interesting and important/innovative data was obtained.

Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. We hope our revision will meet with your approval.

Reviewer #5: In general, the English must be improved along the manuscript. Some parts of the manuscript are very confusing.

Authors' response: Thank you for your comment. The MS was revised by a professional English native speaker.

Reviewer #5: Specific comments: You refer along the manuscript that "...Liver, gills, kidney, spleen and muscle were analyzed for superoxide dismutase, catalase, glutathione peroxidases, glutathione reductase and S-transferase, and metallothionein for muscle..." You need to specify when you measure the levels and when you measure the activities.

Authors' response: Thank you for your comment. We measured the activity of SOD, CAT, GPx, Se-GPx, GR, GST and the levels of MTs. We pointed out it in the text.

Reviewer #5: An explanation about the evaluation of MT only in the muscle must also be included.

Authors' response: Thank you for your comment. In the present biomonitoring study, trace elements load and MTs were measured in muscle since the tissue represents the stable pool of trace elements for fish (Barwick and Maher, 2003). This last sentence was also added in the introduction.

Reviewer #5: Abstract. You need to be more specific and concise when you present your results.

Authors' response: Thank you for your comment. We revised the abstract accordingly.

Reviewer #5: Introduction. Improve the English: line 50-57. Improve the English along this section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Material and methods. In the Lines 175/176 substitute the complete names by the abbreviations: SOD, CAT...

Authors' response: Thank you for your comment. Done. Please, see lines 171-172.

Reviewer #5: Results. The presentation of the results is a little bit confusing. The authors must improve this section to be easier to understand the main results of the study.

Authors' response: Thank you for your comment. We revised the results section accordingly.

Reviewer #5: Line 247: Substitute "The mean concentration of trace elements in summer was..." by "The mean concentration of trace elements may be ranked as follow:..." The same for the Line 249.

Authors' response: Thank you for your comment. We rephrased the sentence.

Reviewer #5: The first time that you refer SeGPx is on the Results section. You need to refer this enzyme previously, at least on the Materials and Methods section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Sometimes along the Results section, it is difficult to understand if the results that you compare are significantly different or not. For example: Line: 260-262: "In spleen lower activity of SeGPx (70%) in females (Fig. 6), and GR and GST (up to 70%) in males (Fig. 7, 8) was measured in fish sampled 262 in autumn when compared to those of summer." It is significantly different? We need to see the figure to understand if it is or not. You need to clarify this along the text of the Results section when you compare the data.

Authors' response: Thank you for your comment. We revised this section accordingly.

Reviewer #5: I did not understand Line 290-291: "The separation between the organs from both seasons (summer on the right, 291 autumn on the left) suggested a difference in biomarkers concentration values." Can you clarify this?

Authors' response: Thank you for your comment. The biplot of loadings (variables) and score (observations) showed which organs (liver, gills, kidney, spleen and muscle) of brook trout is closest to them, and which variables (trace elements, biomarkers and physicochemical parameters) contributed to this grouping in the coordinate of Dim1 and Dim2. The separation between the organs from both seasons (summer on the right, autumn on the left) suggested a difference in biomarkers values.

Reviewer #5: Discussion. Improve the English along this section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Line 323: "The few studies on trace elements accumulation in biota from high-altitude lakes in general, and in Alpine lakes in particular, have mostly focused on fish." On Muscle of fish? Why you only measured on muscle?

Authors' response: Thank you for your comment. Yes, in fish muscle. We added this information in the text. We measured the trace element accumulation in muscle because it constitutes a stable pool of trace elements for fish (Barwick and Maher, 2003) (line 85).

Reviewer #5: Line 367/397: Did you measure the levels or the activity of CAT/SOD? I think that you measure activity but sometimes along the manuscript you refer levels.

Authors' response: Thank you for your comment. Yes, we measured the activity. We pointed out it throughout the text.

Reviewer #5: The Conclusion section does not show the main results of the study. The information of this section is too vague. You need to improve the section including the most relevant findings of the study, a take home message and future perspectives.

Authors' response: Thank you for your comment. Done. Please, see Conclusion section.

Reviewer #5: Figures. In Figure 3, is Superoxide dismutase concentration? It was activity, right? Put SOD on the caption of the figure to correspond to the complete name.

Authors' response: Thank you for your comment. Yes, activity. Done.

Reviewer #5: Figure 4: Put CAT on the caption of the figure to correspond to the complete name. The same for the figures 5-9.
Authors' response: Thank you for your comment. Done.

Response to Reviewer #6

Reviewer #6 1. Recommendation: major revision

2. Comments to Author:

Ms. Ref. No.: STOTEN-D-19-17682

Title: Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps). Overview and general recommendation:

Biomonitoring studies are constantly being developed field of research. More and more diagnostic tools are being proposed and monitoring of oxidative stress markers is part of this trend. The imbalance between production and elimination of Reactive Oxygen Species (ROS) leading to oxidative stress is valuable measure for organism metabolic and general health status. The methodology is accessible and well described, however, it has some limitations e.g. is species-specific; the results might be influenced by many factors. Including biochemical analysis into ecological research provides more comprehensive insight into the general problem aimed to be solved. The current study is quite well written. It describes several factors, which can affect the ecology of brook trout inhabiting extreme ecosystems, such as high-mountain lakes, and makes an attempt to verify this influence. In this case, the paper fits the journal scope. The authors put a big effort to carry out the studies in such remote site and conduct all the laboratory analysis, what should be highlighted and recognised. They distinguish between males and females, which is not common in oxidative stress research. The important support of their work is also quite detailed description of the study site as well as considering the diet composition. I find the research interesting and valuable, however, the authors did not avoid some major flaws. Although the introduction to the topic, study design and its performance generally meet the reviewer's expectations, the discussion of the obtained results is the weakest part of the manuscript. The authors indicate differences in measured parameters between tested tissues and sex, but they do not discuss it enough. There is almost no attempt to explain the cause of diverse results obtained for females compared to males, they are just mentioned. I am aware of the difficulty and effort this task requires; however, I think it would be more meaningful. Similar situation applies to the statement that food intake mainly affected the oxidative stress biomarkers. There is almost no discussion of the results as well as this issue and suddenly such conclusion appears. I encourage authors to rewrite the discussion part and consider the interpretation of obtained results. Therefore, I recommend the major revision of the paper.
Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. Furthermore, the MS was revised by a professional English native speaker. We hope our revision will meet with your approval.

Reviewer #6: Detailed comments:

Title. As I mentioned in the general comments, if authors decide to regard oxidative stress ecology in their manuscript, the obtained results should be discussed in line with ecological explanation. If not, it would be better to change 'ecology' to 'parameters'.

Authors' response: Thank you for your comment. We decided to maintain the "ecology" in the title, since we consider both biotic and abiotic parameters.

Reviewer #6: Keywords: brook trout and oxidative stress are words already mentioned in the title, thus I recommend exchanging them.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Highlights. Bullet points should capture the novel results of the research and/or new methods that were used. Now, the first three present what has been done in the study only (commonly used methods, nothing new or unique). The last one shows the results of the study, however, with the mistake (it should be food intake instead of feed intake).

Authors' response: Thank you for your comment. We revised the highlights following your suggestion.

Reviewer #6: Graphical abstract. Maybe it would be better to use another symbol for ice-free period than the leaf, especially when in the site description it is mentioned that the lake is located above the tree line. It is just a suggestion.

Authors' response: Thank you for your suggestion. We preferred to maintain the leaf since the emblem of the autumn.

Reviewer #6: Abstract. I recommend to change (finish after naming all the tested enzymes) the sentence starting in line 28 (Liver, gills, kidney, spleen and muscle were analyzed for superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase and S-transferase.) and start the new sentence: Bioaccumulation of Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se and Zn and metallothionein level were investigated for both sexes in muscles. Without change, it can be confusing as oxidative stress parameters were measured in different tissues and metallothionein in muscle only. Combining measures of heavy metals and metallothionein together seems more appropriate as all were analyzed in muscle.

Authors' response: Thank you for your suggestion. Done.

Reviewer #6: Introduction.

Line 53: I suggest to change like to such as.

Line 62: I suggest to change like for e.g..

Line 66: Once in the organism,... Once what? Maybe Once introduced to the organism,...

Line 67: I suggest to change ensuing to leading.

Line 83: add activity after antioxidant enzymes.

Authors' response: Thank you for your suggestions. Done.

Reviewer #6: Line 86: what about other factors such as temperature?

Authors' response: Thank you for your comment. We added few sentences following your suggestion. Please, see line 96-98.

Reviewer #6: Materials and Methods.

Line 99: remove is.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 100: remove in which.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 112: captured instead of capture.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 117: The homogeneity of fish sizes are uncertain as the range for each term and sex is more than 10 cm (extremally it reaches 18 cm in autumn for females).

Authors' response: Thank you for your comment. We collected fish entangled by gillnets. Thus, it was not possible to select fish for this study.

Reviewer #6: Line 123: change at level order for at order level and remove a before closer.

Line 128: at instead of in 5 sites.

Line 129: remove in before other.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 151: It would be good to mention what tissues were analyzed.

Authors' response: Thank you for your comment. We analysed Liver, gills, kidney, spleen and muscle. Please, see line 175.

Reviewer #6: Line 153: what about temperature during the centrifuging homogenates?

Line 168: The assay was measured in ... buffer.

Line 174: Please add the time and temperature of centrifuging.

Authors' response: Thank you for your comment. Temperature of centrifugation was 4°C.

Homogenates were centrifuged at 50,000 x g for 30 minutes. It was already inserted in the original MS. The sentence was completed as follows:

"Homogenates were centrifuged at 50,000 x g for 30 minutes at 4°C". For enzymatic analysis samples were homogenized with an UltraTurrax homogenizer in 100 mM potassium-phosphate buffer, pH 7.5, added with 2.5% sodium chloride (NaCl), 0.008 TIU ml⁻¹ aprotinin and 0.1 mg ml⁻¹ bacitracin. It was already inserted in the original MS.

Reviewer #6: Results. Line 207: Physico-chemical (with a dash to stay consequent).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 209 and 228: p value should be consequently presented either $p > 0.05$ or with particular result.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 212-216: This description is unnecessary as all these informations are already presented in the table and are not so relevant for the studies.

Authors' response: Thank you for your comment, but we think that the information are important for readers.

Reviewer #6: Lines 216-218: The diet analysis is poorly described considering the activities showed in the Materials and methods section. Especially, that the main conclusion is based on the food intake of the brook trout. This part of the results section need to be extended and presented with more details (e.g. table of food categories with, at least, frequency of occurrence).

Authors' response: Thank you for your comment. Done. Please see section 3.2.

Reviewer #6: Line 226: Add '>' between Ni (0.06) and Hg (0.02).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 235-237: I suggest to rewrite the sentence: In spleen, lower activity of SeGPx (70%) in females (Fig. 6), and GR and GST (up to 70%) in males (Fig. 7, 8) was recorded in fish from autumn samples compared to summer one.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 238: add comma after In muscle and rewrite the beginning of the sentence: instead of a marked lowering was measured for SOD... I suggest to write In muscle, significant decrease in SOD activity (up to 50%) in autumn was measured for females.

Line 240: add comma after tissue.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Discussion.

Lines 290 and 295: repetition of studied the, this could be easily solved.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 297: exchange how to that.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Reviewer #6: Lines 297-300: I do not understand the purpose for presenting the results of this particular studies with details. Why this one only (authors mention other papers on trace elements accumulation in fish)?

Authors' response: Thank you for your comment. We presented the results of other studies following the suggestion of other reviewers.

Reviewer #6: Line 303: change these for the one presented by other ...

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 305: I do not understand why the authors mention pedogenic sources as a source of contamination, especially when the sentence in line 313 partially contradict this statement.

Authors' response: Thank you for your comment. Since our results are in line with other studies on trace elements accumulation in high-mountain lakes, and no studies have been performed previously in Balma Lake, we supposed that trace elements amount detected in brook trout from Balma Lake have an origin both from anthropogenic and pedogenic sources. This sentence was added following other reviewer's suggestions.

Reviewer #6: Lines 307-309: This statement better suits materials and method section.

Authors' response: Thank you for your comment, but we think that this statement is essential to introduce the discussion about metallothionines.

Reviewer #6: Line 321: put in accordance with other studies in commas.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 330: in fish from autumn samples or sampled in autumn.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 349: strenghten is not appropriate and should be changed for strongly. I also recommend to move strongly after how (PCA analysis showed how strong samples from liver in the summer were related to oxidative....).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 356: remove for.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 359-360: Mentioned species is just an example, I would not say it is well known for the broad audience. This sentence should be changed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 375: remove a.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 377-383: Why? What are the basis and reasons of such assumption? Authors present it as a main result of their studies without any analysis and discussion.

Authors' response: Thank you for your comment. The results and discussion about food intake were implemented.

Reviewer #6: Line 382: food instead of feed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Conclusions.

Line 386: was instead of is.

Line 388: tissues instead of organs.

Line 392: food instead of feed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: References. All the papers from the reference list were cited, however, lack of consequence in their format is striking. According to the journal requirements, the journal titles should be abbreviated with dots. In some references, this rule is obeyed, in most not (lines to be corrected: 404, 407, 409, 415, 417, 423, 467, 474, 490, 497, 508, 511, 520, 531, 533, 537, 545, 548, 551, 553, 562, 564).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Table 2. Add \pm SD to Total Length - TL mean (cm).

Authors' response: Thank you for your comment. Done.

Reviewer #6: All the tables and figures were cited in the text.

Authors' response: Thanks.

Response to Reviewer #7

Reviewer #7: The manuscript titled "Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps)" (MS Number STOTEN-D-19-17682R1) by Pastorino P and colleagues investigate the environmental stressors pressure on brook trout (*Salvelinus fontinalis*) from Balma Lake, located in Piedmont Region (Cottian Alps, Northwest Italy). Main Physical-chemical parameters and nutrients were analyzed in water samples. Beside fish growth measurement, in fish muscle, liver, kidney, gills and spleen tissue, biomarkers of oxidative stress (superoxide dismutase, catalase, glutathione peroxidases, glutathione reductase and S-transferase, and metallothionein in fish muscle tissue) were analyzed. Furthermore, bioaccumulation level of trace elements (Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se and Zn) was analyzed in fish muscle tissue. Authors concluded that increased feed intake during the ice-free season mainly influenced the oxidative stress biomarkers.

In general, I recommend accepting the paper in STOTEN after minor corrections and additions.

Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. We hope our revision will meet with your approval.

Reviewer #7: General suggestions:

1. Exchange the term "Water nutrients" with "nutrients". Authors measured nutrients in the water!

Authors' response: Thank you for your comment. Done.

Reviewer #7: 2. Instead of "Physical-chemical", I would suggest the use of "Physico-chemical"

Authors' response: Thank you for your comment. Done.

Reviewer #7: HIGHLIGHTS Replace the order of the second and third line.

Authors' response: Thank you for your comment. The highlights were rewritten following the other reviewers' suggestions.

Reviewer #7: ABSTRACT. Line 23: additional stress instead of "additive stress"

Authors' response: Thank you for your comment. Done.

Reviewer #7: INTRODUCTION. Appropriate and acceptable, only a few small corrections:

Line 101 -102: see general suggestions, especially the term "nutrients characteristics of water" should be corrected

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 102 and 104: in both lines you have "c)". In line 104 should be "d)"

Authors' response: Thank you for your comment. Done.

Reviewer #7: MATERIALS AND METHODS. Appropriate and acceptable, only a few small corrections and one more explanation:

LINES 114-115. Correct the sentence (English is not correct; so the sentence is not clear enough)

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 126: "in 3 deep zone ones"??? What is this? Is it "at three depths"?

Authors' response: Thank you for your comment. We rephrased it.

Reviewer #7: Lines 134-139: three methods are cited, but measurement was made "by an adoption" of these three methods. If something is changed in methodology - it should be described. So, these adoptions of the methods should be explained! Otherwise, these methods could not be performed by others.

Authors' response: Thank you for your comment. We have taken this methodology from the instruction manual of spectrophotometer (Hanna Instrument). We didn't know other information.

Reviewer #7: Line 141: Delete "campaigns", "fish sampling" is just enough

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 142-144: The sentence "The choice of the two seasons was prescribed by the necessity to reach on foot the site during the ice free period." (Write the sentence in better English!)
Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 145: captured (add "d")
Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 174: instead of "in single" use "individually"
Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 177: use "enzymatic analysis" instead of "enzymes analysis"
Authors' response: Thank you for your comment. Done.

Reviewer #7: RESULTS. The results are presented in a clear manner. The figures used to show them are adequate. However, I would recommend:
Line 225: exchange "Physical-chemical and nutrients characteristics of water" with "Physico-chemical parameter and nutrient levels in water"
Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 259- 260: add "same" in front of seasons
Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 270: delete water in front of nutrients
Authors' response: Thank you for your comment. Done.

Reviewer #7: DISCUSSION Line 329: better use trace elements instead of heavy metals, because Se is not Heavy metal (listed in line 331)
Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 359-361: Comment: Alkalinity is not same as pH, so this statement ("explanation of this outcome") is not proper!
Authors' response: Thank you for your comment. We corrected it.

Reviewer #7: 411: same comment as for the line 270
Authors' response: Thank you for your comment. Done.

Reviewer #7: CONCLUSIONS. Clear. Hope my comments will help authors to improve their manuscript!!!
Authors' response: Thank you for your suggestions.



Ref: STOTEN-D-19-17682R1

To the Editor-in-Chief

Damià Barceló

Science of the Total Environment

Dear Editor-in-Chief,

on behalf of my co-authors, I would like to thank the reviewers for the positive and constructive suggestions and for the opportunity to revise our manuscript entitled “Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps)” for publication in *Science of the Total Environment*. We have studied the comments carefully and revised the manuscript accordingly, which we hope will meet with your final approval. In the section “Responses to Reviewers Comments”, we have provided detailed responses to the reviewers’ comments and illustrated the corrections made in the paper. All new changes are underlined in red in the manuscript. Furthermore, the manuscript was carefully read and revised by an English native speaker.

We are confident that the manuscript in the present form is ready for publication.

We remain at complete disposal for any further information you might require.

Kind regards

Paolo Pastorino

**Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake
(Cottian Alps)**

Paolo Pastorino^{1,2*}, Antonia Concetta Elia^{3**}, Barbara Caldaroni³, Vasco Menconi², Maria Cesarina
Abete², Paola Brizio², Marco Bertoli¹, Annalisa Zaccaroni⁴, Magara Gabriele³, Ambrosius Josef
Martin Dörr³, Elisabetta Pizzul¹, Marino Prearo¹

¹ Department of Life Sciences, University of Trieste, via Giorgieri 10, 34127 Trieste, Italy

² The Veterinary Medical Research Institute for Piemonte, Liguria and Valle d'Aosta, via Bologna
148, 10154 Torino, Italy

³ Department of Chemistry Biology and Biotechnology, University of Perugia, via Elce di Sotto 8,
06123 Perugia, Italy

⁴ Department of Veterinary Medical Science, University of Bologna, viale Vespucci 2, 47042
Cesenatico (FC), Italy

*Corresponding author: Paolo Pastorino, e-mail address: paolo.pastorino@izsto.it (P. Pastorino);

** Corresponding author: Antonia Concetta Elia, e-mail address: antonia.elia@unipg.it (A.C. Elia)

Response to Reviewer 5

Reviewer #5: In general, the authors improve the manuscript taking into account the comments/suggestions of the reviewers.

I recommend the publication of this study after some issues be addressed. The aim of the study is interesting and important/innovative data was obtained.

Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. We hope our revision will meet with your approval.

Reviewer #5: In general, the English must be improved along the manuscript. Some parts of the manuscript are very confusing.

Authors' response: Thank you for your comment. The MS was revised by a professional English native speaker.

Reviewer #5: Specific comments: You refer along the manuscript that "...Liver, gills, kidney, spleen and muscle were analyzed for superoxide dismutase, catalase, glutathione peroxidases, glutathione reductase and S-transferase, and metallothionein for muscle..." You need to specify when you measure the levels and when you measure the activities.

Authors' response: Thank you for your comment. We measured the activity of SOD, CAT, GPx, Se-GPx, GR, GST and the levels of MTs. We pointed out it in the text.

Reviewer #5: An explanation about the evaluation of MT only in the muscle must also be included.

Authors' response: Thank you for your comment. In the present biomonitoring study, trace elements load and MTs were measured in muscle since the tissue represents the stable pool of trace elements for fish (Barwick and Maher, 2003). This last sentence was also added in the introduction.

Reviewer #5: Abstract. You need to be more specific and concise when you present your results.

Authors' response: Thank you for your comment. We revised the abstract accordingly.

Reviewer #5: Introduction. Improve the English: line 50-57. Improve the English along this section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Material and methods. In the Lines 175/176 substitute the complete names by the abbreviations: SOD, CAT...

Authors' response: Thank you for your comment. Done. Please, see lines 171-172.

Reviewer #5: Results. The presentation of the results is a little bit confusing. The authors must improve this section to be easier to understand the main results of the study.

Authors' response: Thank you for your comment. We revised the results section accordingly.

Reviewer #5: Line 247: Substitute "The mean concentration of trace elements in summer was..." by "The mean concentration of trace elements may be ranked as follow:..." The same for the Line 249.

Authors' response: Thank you for your comment. We rephrased the sentence.

Reviewer #5: The first time that you refer SeGPx is on the Results section. You need to refer this enzyme previously, at least on the Materials and Methods section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Sometimes along the Results section, it is difficult to understand if the results that you compare are significantly different or not. For example: Line: 260-262: "In spleen lower activity of SeGPx (70%) in females (Fig. 6), and GR and GST (up to 70%) in males (Fig. 7, 8) was measured in fish sampled 262 in autumn when compared to those of summer." It is significantly different? We need to see the figure to understand if it is or not. You need to clarify this along the text of the Results section when you compare the data.

Authors' response: Thank you for your comment. We revised this section accordingly.

Reviewer #5: I did not understand Line 290-291: "The separation between the organs from both seasons (summer on the right, 291 autumn on the left) suggested a difference in biomarkers concentration values." Can you clarify this?

Authors' response: Thank you for your comment. The biplot of loadings (variables) and score (observations) showed which organs (liver, gills, kidney, spleen and muscle) of brook trout is closest to them, and which variables (trace elements, biomarkers and physicochemical parameters) contributed to this grouping in the coordinate of Dim1 and Dim2. The separation between the organs from both seasons (summer on the right, autumn on the left) suggested a difference in biomarkers values.

Reviewer #5: Discussion. Improve the English along this section.

Authors' response: Thank you for your comment. Done.

Reviewer #5: Line 323: "The few studies on trace elements accumulation in biota from high-altitude lakes in general, and in Alpine lakes in particular, have mostly focused on fish." On Muscle of fish? Why you only measured on muscle?

Authors' response: Thank you for your comment. Yes, in fish muscle. We added this information in the text. We measured the trace element accumulation in muscle because it constitutes a stable pool of trace elements for fish (Barwick and Maher, 2003) (line 85).

Reviewer #5: Line 367/397: Did you measure the levels or the activity of CAT/SOD? I think that you measure activity but sometimes along the manuscript you refer levels.

Authors' response: Thank you for your comment. Yes, we measured the activity. We pointed out it throughout the text.

Reviewer #5: The Conclusion section does not show the main results of the study. The information of this section is too vague. You need to improve the section including the most relevant findings of the study, a take home message and future perspectives.

Authors' response: Thank you for your comment. Done. Please, see Conclusion section.

Reviewer #5: Figures. In Figure 3, is Superoxide dismutase concentration? It was activity, right? Put SOD on the caption of the figure to correspond to the complete name.

Authors' response: Thank you for your comment. Yes, activity. Done.

Reviewer #5: Figure 4: Put CAT on the caption of the figure to correspond to the complete name. The same for the figures 5-9.

Authors' response: Thank you for your comment. Done.

Response to Reviewer #6

Reviewer #6 1. Recommendation: major revision

2. Comments to Author:

Ms. Ref. No.: STOTEN-D-19-17682

Title: Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps). Overview and general recommendation:

Biomonitoring studies are constantly being developed field of research. More and more diagnostic tools are being proposed and monitoring of oxidative stress markers is part of this trend. The imbalance between production and elimination of Reactive Oxygen Species (ROS) leading to oxidative stress is valuable measure for organism metabolic and general health status. The methodology is accessible and well described, however, it has some limitations e.g. is species-specific; the results might be influenced by many factors. Including biochemical analysis into ecological research provides more comprehensive insight into the general problem aimed to be solved. The current study is quite well written. It describes several factors, which can affect the ecology of brook trout inhabiting extreme ecosystems, such as high-mountain lakes, and makes an attempt to verify this influence. In this case, the paper fits the journal scope. The authors put a big effort to carry out the studies in such remote site and conduct all the laboratory analysis, what should be highlighted and recognised. They distinguish between males and females, which is not common in oxidative stress research. The important support of their work is also quite detailed description of the study site as well as considering the diet composition. I find the research interesting and valuable, however, the authors did not avoid some major flaws. Although the introduction to the topic, study design and its performance generally meet the reviewer's expectations, the discussion of the obtained results is the weakest part of the manuscript. The authors indicate differences in measured parameters between tested tissues and sex, but they do not discuss it enough. There is almost no attempt to explain the cause of diverse results obtained for females compared to males, they are just mentioned. I am aware of the difficulty and effort this task requires; however, I think it would be more meaningful. Similar situation applies to the statement that food intake mainly affected the oxidative stress biomarkers. There is almost no discussion of the results as well as this issue and suddenly such conclusion appears. I encourage authors to rewrite the discussion part and consider the interpretation of obtained results. Therefore, I recommend the major revision of the paper.

Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. Furthermore, the MS was revised by a professional English native speaker. We hope our revision will meet with your approval.

Reviewer #6: Detailed comments:

Title. As I mentioned in the general comments, if authors decide to regard oxidative stress ecology in their manuscript, the obtained results should be discussed in line with ecological explanation. If not, it would be better to change 'ecology' to 'parameters'.

Authors' response: Thank you for your comment. We decided to maintain the “ecology” in the title, since we consider both biotic and abiotic parameters.

Reviewer #6: Keywords: brook trout and oxidative stress are words already mentioned in the title, thus I recommend exchanging them.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Highlights. Bullet points should capture the novel results of the research and/or new methods that were used. Now, the first three present what has been done in the study only (commonly used methods, nothing new or unique). The last one shows the results of the study, however, with the mistake (it should be food intake instead of feed intake).

Authors' response: Thank you for your comment. We revised the highlights following your suggestion.

Reviewer #6: Graphical abstract. Maybe it would be better to use another symbol for ice-free period than the leaf, especially when in the site description it is mentioned that the lake is located above the tree line. It is just a suggestion.

Authors' response: Thank you for your suggestion. We preferred to maintain the leaf since the emblem of the autumn.

Reviewer #6: Abstract. I recommend to change (finish after naming all the tested enzymes) the sentence starting in line 28 (Liver, gills, kidney, spleen and muscle were analyzed for superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase and S-transferase.) and start the new sentence: Bioaccumulation of Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se and Zn and metallothionein level were investigated for both sexes in muscles. Without change, it can be confusing as oxidative stress parameters were measured in different tissues and metallothionein in muscle only. Combining measures of heavy metals and metallothionein together seems more appropriate as all were analyzed in muscle.

Authors' response: Thank you for your suggestion. Done.

Reviewer #6: Introduction.

Line 53: I suggest to change like to such as.

Line 62: I suggest to change like for e.g..

Line 66: Once in the organism,... Once what? Maybe Once introduced to the organism,...

Line 67: I suggest to change ensuing to leading.

Line 83: add activity after antioxidant enzymes.

Authors' response: Thank you for your suggestions. Done.

Reviewer #6: Line 86: what about other factors such as temperature?

Authors' response: Thank you for your comment. We added few sentences following your suggestion. Please, see line 96-98.

Reviewer #6: Materials and Methods.

Line 99: remove is.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 100: remove in which.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 112: captured instead of capture.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 117: The homogeneity of fish sizes are uncertain as the range for each term and sex is more than 10 cm (extremally it reaches 18 cm in autumn for females).

Authors' response: Thank you for your comment. We collected fish entangled by gillnets. Thus, it was not possible to select fish for this study.

Reviewer #6: Line 123: change at level order for at order level and remove a before closer.

Line 128: at instead of in 5 sites.

Line 129: remove in before other.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 151: It would be good to mention what tissues were analyzed.

Authors' response: Thank you for your comment. We analysed Liver, gills, kidney, spleen and muscle. Please, see line 175.

Reviewer #6: Line 153: what about temperature during the centrifuging homogenates?

Line 168: The assay was measured in ... buffer.

Line 174: Please add the time and temperature of centrifuging.

Authors' response: Thank you for your comment. Temperature of centrifugation was 4°C.

Homogenates were centrifuged at 50,000 x g for 30 minutes. It was already inserted in the original MS. The sentence was completed as follows: "Homogenates were centrifuged at 50,000 x g for 30 minutes at 4°C". For enzymatic analysis samples were homogenized with an UltraTurrax homogenizer in 100 mM potassium-phosphate buffer, pH 7.5, added with 2.5% sodium chloride (NaCl), 0.008 TIU ml⁻¹ aprotinin and 0.1 mg ml⁻¹ bacitracin. It was already inserted in the original MS.

Reviewer #6: Results. Line 207: Physico-chemical (with a dash to stay consequent).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 209 and 228: p value should be consequently presented either $p > 0.05$ or with particular result.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 212-216: This description is unnecessary as all these informations are already presented in the table and are not so relevant for the studies.

Authors' response: Thank you for your comment, but we think that the information are important for readers.

Reviewer #6: Lines 216-218: The diet analysis is poorly described considering the activities showed in the Materials and methods section. Especially, that the main conclusion is based on the food intake of the brook trout. This part of the results section need to be extended and presented with more details (e.g. table of food categories with, at least, frequency of occurrence).

Authors' response: Thank you for your comment. Done. Please see section 3.2.

Reviewer #6: Line 226: Add '>' between Ni (0.06) and Hg (0.02).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 235-237: I suggest to rewrite the sentence: In spleen, lower activity of SeGPx (70%) in females (Fig. 6), and GR and GST (up to 70%) in males (Fig. 7, 8) was recorded in fish from autumn samples compared to summer one.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 238: add comma after In muscle and rewrite the beginning of the sentence: instead of a marked lowering was measured for SOD... I suggest to write In muscle, significant decrease in SOD activity (up to 50%) in autumn was measured for females.

Line 240: add comma after tissue.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Discussion.

Lines 290 and 295: repetition of studied the, this could be easily solved.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 297: exchange how to that.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Reviewer #6: Lines 297-300: I do not understand the purpose for presenting the results of this particular studies with details. Why this one only (authors mention other papers on trace elements accumulation in fish)?

Authors' response: Thank you for your comment. We presented the results of other studies following the suggestion of other reviewers.

Reviewer #6: Line 303: change these for the one presented by other ...

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 305: I do not understand why the authors mention pedogenic sources as a source of contamination, especially when the sentence in line 313 partially contradict this statement.

Authors' response: Thank you for your comment. Since our results are in line with other studies on trace elements accumulation in high-mountain lakes, and no studies have been performed previously in Balma Lake, we supposed that trace elements amount detected in brook trout from Balma Lake have an origin both from anthropogenic and pedogenic sources. This sentence was added following other reviewer's suggestions.

Reviewer #6: Lines 307-309: This statement better suits materials and method section.

Authors' response: Thank you for your comment, but we think that this statement is essential to introduce the discussion about metallothionines.

Reviewer #6: Line 321: put in accordance with other studies in commas.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 330: in fish from autumn samples or sampled in autumn.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 349: strenghten is not appropriate and should be changed for strongly. I also recommend to move strongly after how (PCA analysis showed how strong samples from liver in the summer were related to oxidative....).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 356: remove for.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 359-360: Mentioned species is just an example, I would not say it is well known for the broad audience. This sentence should be changed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Line 375: remove a.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Lines 377-383: Why? What are the basis and reasons of such assumption? Authors present it as a main result of their studies without any analysis and discussion.

Authors' response: Thank you for your comment. The results and discussion about food intake were implemented.

Reviewer #6: Line 382: food instead of feed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: Conclusions.

Line 386: was instead of is.

Line 388: tissues instead of organs.

Line 392: food instead of feed.

Authors' response: Thank you for your comment. Done.

Reviewer #6: References. All the papers from the reference list were cited, however, lack of consequence in their format is striking. According to the journal requirements, the journal titles should be abbreviated with dots. In some references, this rule is obeyed, in most not (lines to be corrected: 404, 407, 409, 415, 417, 423, 467, 474, 490, 497, 508, 511, 520, 531, 533, 537, 545, 548, 551, 553, 562, 564).

Authors' response: Thank you for your comment. Done.

Reviewer #6: Table 2. Add \pm SD to Total Length - TL mean (cm).

Authors' response: Thank you for your comment. Done.

Reviewer #6: All the tables and figures were cited in the text.

Authors' response: Thanks.

Response to Reviewer #7

Reviewer #7: The manuscript titled "Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps)" (MS Number STOTEN-D-19-17682R1) by Pastorino P and colleagues investigate the environmental stressors pressure on brook trout (*Salvelinus fontinalis*) from Balma Lake, located in Piedmont Region (Cottian Alps, Northwest Italy). Main Physical-chemical parameters and nutrients were analyzed in water samples. Beside fish growth measurement, in fish muscle, liver, kidney, gills and spleen tissue, biomarkers of oxidative stress (superoxide dismutase, catalase, glutathione peroxidases, glutathione reductase and S-transferase, and metallothionein in fish muscle tissue) were analyzed. Furthermore, bioaccumulation level of trace elements (Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se and Zn) was analyzed in fish muscle tissue. Authors concluded that increased feed intake during the ice-free season mainly influenced the oxidative stress biomarkers. In general, I recommend accepting the paper in STOTEN after minor corrections and additions.

Authors' response: We are very pleased for your comments, thanks. We have studied your comments carefully and revised the manuscript accordingly. We hope our revision will meet with your approval.

Reviewer #7: General suggestions:

1. Exchange the term "Water nutrients" with "nutrients". Authors measured nutrients in the water!

Authors' response: Thank you for your comment. Done.

Reviewer #7: 2. Instead of "Physical-chemical", I would suggest the use of "Physico-chemical"

Authors' response: Thank you for your comment. Done.

Reviewer #7: HIGHLIGHTS Replace the order of the second and third line.

Authors' response: Thank you for your comment. The highlights were rewritten following the other reviewers' suggestions.

Reviewer #7: ABSTRACT. Line 23: additional stress instead of "additive stress"

Authors' response: Thank you for your comment. Done.

Reviewer #7: INTRODUCTION. Appropriate and acceptable, only a few small corrections:
Line 101 -102: see general suggestions, especially the term "nutrients characteristics of water" should be corrected

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 102 and 104: in both lines you have "c)". In line 104 should be "d)"

Authors' response: Thank you for your comment. Done.

Reviewer #7: MATERIALS AND METHODS. Appropriate and acceptable, only a few small corrections and one more explanation:

LINES 114-115. Correct the sentence (English is not correct; so the sentence is not clear enough)

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 126: "in 3 deep zone ones"??? What is this? Is it "at three depths"?

Authors' response: Thank you for your comment. We rephrased it.

Reviewer #7: Lines 134-139: three methods are cited, but measurement was made "by an adoption" of these three methods. If something is changed in methodology - it should be described. So, these adoptions of the methods should be explained! Otherwise, these methods could not be performed by others.

Authors' response: Thank you for your comment. We have taken this methodology from the instruction manual of spectrophotometer (Hanna Instrument). We didn't know other information.

Reviewer #7: Line 141: Delete "campaigns", "fish sampling" is just enough

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 142-144: The sentence "The choice of the two seasons was prescribed by the necessity to reach on foot the site during the ice free period." (Write the sentence in better English!)

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 145: captured (add "d")

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 174: instead of "in single" use "individually"

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 177: use "enzymatic analysis" instead of "enzymes analysis"

Authors' response: Thank you for your comment. Done.

Reviewer #7: RESULTS. The results are presented in a clear manner. The figures used to show them are adequate. However, I would recommend:

Line 225: exchange "Physical-chemical and nutrients characteristics of water" with "Physico-chemical parameter and nutrient levels in water"

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 259- 260: add "same" in front of seasons

Authors' response: Thank you for your comment. Done.

Reviewer #7: Line 270: delete water in front of nutrients

Authors' response: Thank you for your comment. Done.

Reviewer #7: DISCUSSION Line 329: better use trace elements instead of heavy metals, because Se is not Heavy metal (listed in line 331)

Authors' response: Thank you for your comment. Done.

Reviewer #7: Lines 359-361: Comment: Alkalinity is not same as pH, so this statement ("explanation of this outcome") is not proper!

Authors' response: Thank you for your comment. We corrected it.

Reviewer #7: 411: same comment as for the line 270

Authors' response: Thank you for your comment. Done.

Reviewer #7: CONCLUSIONS. Clear. Hope my comments will help authors to improve their manuscript!!!

Authors' response: Thank you for your suggestions.

1 **Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake**
2 **(Cottian Alps)**

3
4 Paolo Pastorino^{1,2*}, Antonia Concetta Elia^{3**}, Barbara Caldaroni³, Vasco Menconi², Maria Cesarina
5 Abete², Paola Brizio², Marco Bertoli¹, Annalisa Zaccaroni⁴, Magara Gabriele³, Ambrosius Josef
6 Martin Dörr³, Elisabetta Pizzul¹, Marino Prearo¹

7
8 ¹ Department of Life Sciences, University of Trieste, via Giorgieri 10, 34127 Trieste, Italy
9 ² The Veterinary Medical Research Institute for Piemonte, Liguria and Valle d'Aosta, via Bologna
10 148, 10154 Torino, Italy
11 ³ Department of Chemistry Biology and Biotechnology, University of Perugia, via Elce di Sotto 8,
12 06123 Perugia, Italy
13 ⁴ Department of Veterinary Medical Science, University of Bologna, viale Vespucci 2, 47042
14 Cesenatico (FC), Italy

15
16 *Corresponding author: Paolo Pastorino, e-mail address: paolo.pastorino@izsto.it (P. Pastorino);

17
18 ** Corresponding author: Antonia Concetta Elia, e-mail address: antonia.elia@unipg.it (A.C. Elia)

19
20 **Abstract**

21 High-mountain lakes are pristine ecosystems characterized by extreme environmental conditions.
22 The atmospheric transport of pollutants from lowlands may add further stress to organisms
23 inhabiting these environments. We investigated the environmental stress pressure on brook trout
24 (*Salvelinus fontinalis*) from a high-mountain lake in the Cottian Alps (Piedmont, northwest Italy).
25 To do this, males and females of brook trout were sampled from Balma Lake in summer (August)
26 and autumn (October) 2017 in order to assess the influence of trace elements accumulation and
27 environmental parameters (physicochemical parameters and nutrient characteristics of water) on
28 oxidative stress biomarkers. Bioaccumulation of Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se, and Zn and
29 metallothionein levels were measured in muscle tissue of males and females. Liver, gills, kidney,
30 and spleen tissue samples were analyzed for superoxide dismutase, catalase, total glutathione
31 peroxidase, selenium-dependent glutathione peroxidase, glutathione reductase, and glutathione S-
32 transferase activity. Analysis of environmental parameters showed changes in biomarker levels with
33 seasonal variations. Water temperature was significantly higher in summer than autumn (Wilcoxon
34 test; $p = 0.0078$), while pH was significantly higher in autumn than in summer (Wilcoxon test; $p =$

0.0078). Sex-related differences in oxidative stress biomarkers in tissues were unremarkable, whereas seasonal variability of oxidative stress biomarkers was observed, with major differences occurred for liver in summer and for gills, kidney, spleen and muscle in autumn. Positive correlations between environmental parameters and biomarkers were noted. Major fluctuations in water temperature, pH, Cu, Pb and Hg produced changes in **biomarker levels; however**, increased food intake during the ice-free season was probably the main factor that influenced changes in oxidative stress biomarker levels in brook trout in this extreme ecosystem.

Keywords: **Alpine lakes; extreme ecosystems; oxidative stress biomarkers; trace elements**

1. Introduction

Alpine lakes are remote, extreme ecosystems under harsh climatic conditions (Catalán et al., 2006). **The ice-free season lasts for few months, generally from mid-June to late October. During this brief period of ideal conditions, some aquatic organisms can complete their life cycle before the snow covers the lakes again.** Oligotrophic water conditions, UV radiation intensity, together with extreme temperatures allow for the development of a few dominant but well-adapted species (Sommaruga, 2001; Füreder et al., 2006; Pastorino et al., 2019a). These characteristics underlie the negative correlation between altitude and biodiversity (Rott, 1988; Starkweather, 1990). Due to their remote location, Alpine lakes are often considered pristine, unpolluted ecosystems (Pastorino et al., 2019b). Since the 1980s, however, they have been affected by the global anthropogenic impact **of** pollutants transported from lowland emission sources and the introduction of alien species (Tiberti et al., 2014; Pastorino et al., 2020). These ecosystems have low resilience to disturbances and can be particularly sensitive to the release of fish species for recreational angling, with important consequences along the entire trophic chain (Tiberti et al., 2014; Milardi et al., 2016; Perrine, 2017; Raposeiro et al., 2017). For example, the release of salmonids, especially brook trout (*Salvelinus fontinalis*), has had a dramatic outcome for zooplanktonic, macrobenthic, and amphibian communities in several Alpine lakes (Tiberti et al., 2014).

Alpine lakes are also a receptor for medium atmospheric transported (MRAT) contaminants (Ferrario et al., 2017), as observed in the Arctic (Hung et al., 2016), which is subject to the long-range transport potential (LRTP) of many chemicals. Altitudinal transport in the European Alps can occur over relatively short distances from sources of pollution in the industrialized areas of Germany, Switzerland, Austria, and northern Italy (Poma et al., 2017). The input of contaminant compounds into Alpine lakes is controlled by atmospheric deposition and condensation (Schmid et

69 al., 2007). The contaminants are bioaccumulated by the organisms inhabiting these ecosystems.
70 Because fish occupy the uppermost trophic level, they provide an excellent bioindicator for the
71 atmospheric medium-long range input of persistent organic compounds such as pesticides,
72 brominated flame retardants (Schmid et al., 2007), and trace elements (Rognerud et al., 2002; Yang
73 et al., 2007).

74 Moreover, fish are used as sentinel organisms to detect environmental contamination (Squadrone et
75 al., 2013, 2014, 2016). They provide a useful model for assessing the impact of pollutants on
76 biological functions such as detoxification (Elia et al., 2010). Assessment of contaminants in
77 aquatic organisms can estimate and quantify the bioavailable fraction that might have the potential
78 to induce an effect. However, because determination of body concentrations alone does not provide
79 valuable data about the effects, quantification of some biological responses is necessary to evaluate
80 the health state of contaminant-exposed organisms (Elia et al., 2010). Metals, for instance, are pro-
81 oxidants that exert oxidative stress via reactive oxygen species (ROS) production and **cause** critical
82 changes in cellular biotransformation/detoxification pathways (Lushchak, 2016). **Biomarker levels**
83 **can be also influenced by abiotic factors such as pH, dissolved oxygen content, and water**
84 **temperature (Sroda and Cossu-Leguille, 2011). Water temperature is a major factor in physiological**
85 **processes in fish and can induce the production of ROS (Lushchak, 2011).**

86 Oxidative stress results from an imbalance between pro-oxidants such as ROS and the protective
87 antioxidant system. Mechanisms involve the activity of numerous antioxidant enzymes, including
88 superoxide dismutase (SOD, EC 1.15.1.1), catalase (CAT, EC 1.11.1.6), **selenium-dependent**
89 **glutathione peroxidase (Se-GPx, EC 1.11.1.9)**, glutathione reductase (GR, EC 1.6.4.2), glutathione
90 S-transferases (GST, EC 2.5.1.18), as well as the rates of metal-trapping molecules such as reduced
91 glutathione and metallothioneins (MTs). They are important protective metabolic pathways that are
92 used as biomarkers of pollutant-induced oxidative stress. Selected oxidative stress biomarkers have
93 proven useful to assess the impact of a range of metals in aquatic organisms (Al Kaddissi et al.,
94 2014; Cozzari et al., 2015; Elia et al., 2006, 2007a, 2007b, 2010). Furthermore, contaminant levels
95 and antioxidant enzyme **activity** in aquatic organisms may change with the season and in response
96 to biological and environmental pressures (Monserrat et al., 2007). This poses a limitation on field
97 studies, because biochemical response may be linked to either fish physiology or exposure to
98 contaminants.

99 To our best knowledge, no studies are available about oxidative stress in fish from Balma Lake, an
100 Alpine lake in Piedmont (Cottian Alps, northwest Italy). Originally, the lake was fishless and brook
101 trout was released for recreational fishing. With the present study we investigated: a) the
102 **physicochemical parameters** and nutrient characteristics of lake water; b) the biometric

103 characteristics and stomach contents of brook trout (*S. fontinalis*); c) the trace element accumulation
104 in muscle because it constitutes a stable pool of trace elements for fish (Barwick and Maher, 2003);
105 d) the biomarkers of oxidative stress in muscle, liver, kidney, gills, and spleen in male and female
106 individuals captured in Balma Lake in summer and autumn 2017.

107

108 2. Materials and Methods

109 2.1 Study site

110 Balma Lake (45° 02' 14'' N; 07° 10' 52'' E) is located at 2100 m above sea level in the
111 municipality of Coazze, a small town 40 kilometers from Turin (Piedmont, northwest Italy). It is a
112 typical glacial-origin lake in the Cottian Alps, within the SCI/ZSC IT1110006 - Orsiera Rocciavère.
113 The lake is located above the tree line and is covered by ice from November to early June.
114 Originally, the lake was fishless, and *S. fontinalis* was released for recreational fishing during the
115 1970s (Pastorino et al., 2020). The main substrate of the area is composed of ophiolite metamorphic
116 bedrock. The main source of impact is the medium-long distance transport of pollutants from the
117 plain, grazing activities, and angling during the ice-free period. No previous studies or data about
118 the lake's hydrochemistry, taxa composition, and trace element accumulation are available. During
119 summer 2017 morphometric and bathymetric survey of the lake was carried out by GeoStudio RC
120 (Giaveno, Italy) using flying and floating drones. The lake measures 414 m in perimeter, 1.21 ha in
121 surface area, and 6.42 m maximum depth in the central zone (Fig. 1).

122

123 2.2 Physicochemical parameters and nutrients of lake water

124 During both sampling periods the main physicochemical parameters were monitored at 5 sites in the
125 littoral zone (in the upper centimeters of water) and 3 sites in the deep zone (in the water column)
126 (Fig. 1). Water temperature (°C), dissolved oxygen (% saturation; mg L⁻¹), conductivity (µS cm⁻¹),
127 and pH were recorded using field meters (HI 9033 conductivity meter, HI 9125 pH/ORP meter, HI
128 9147 oximeter, Hanna Instruments Inc. Woonsocket, RI, USA). Three replicates were carried out
129 for each parameter. Water samples were collected in sterile containers (three 1-L bottles for each
130 site), taking care not to include sediment particles, and then brought to the laboratory in a
131 refrigerated container within a few hours. Concentrations of NH₄⁺ (mg L⁻¹), NO₃⁻ (mg L⁻¹), and
132 PO₄³⁻ (mg L⁻¹) were measured using a multi-parameter benchtop photometer (HI 83200-02, Hanna
133 Instruments Inc.). NO₃⁻ (mg L⁻¹) concentration was obtained by measuring absorbance at 525 nm
134 via an adaptation of the cadmium reduction method (APHA et al., 1998); NH₄⁺ (mg L⁻¹)
135 concentration was obtained by measuring the absorbance at 420 nm (ASTM, 2015) via adaptation

136 of the Nessler method; finally, PO_4^{3-} (mg L^{-1}) concentration was obtained by measuring absorbance
137 at 610 nm via adaptation of the ascorbic acid method (APHA et al., 1998).

138

139 *2.3 Fish sampling*

140 Fish sampling campaigns were carried out during summer (August) and autumn (October) 2017.
141 These months were chosen so that we could reach the sampling site on foot during the ice-free
142 period. Permission for sampling was granted by the competent authority (Città Metropolitana di
143 Torino; authorization n. 176-19040/2017). Fish were **captured** using 4 multimesh gillnets (36 x 1.8
144 m) divided into 6 panels of different mesh size (10 to 38 mm) to capture all size classes
145 indiscriminately, except for offspring. The gillnets were randomly placed in the lake (Fig. 1) for 3
146 hours and then recovered. For each sampling period, 20 individuals (Table 2) were sacrificed after
147 deep anesthesia with a lethal concentration (200 mg kg^{-1}) of tricaine methanesulfonate (MS-222)
148 dissolved in water. The fish were necropsied, sexed, weighed, and measured for total length in the
149 field. Immediately thereafter, samples of gill, liver, spleen, kidney, and muscle of each specimen
150 were collected, packed in dry ice, and transported to the laboratory.

151 Stomach contents analysis was performed to obtain information about fish diet and to characterize
152 seasonal variations. Stomachs were preserved in 70% alcohol, and the contents were inspected by
153 stereomicroscopy (Zeiss Stemis V8, Jena, Germany). The ingested prey was identified **to the order**
154 **or family level**, since closer identification was precluded by the digestion status of the organisms.
155 To describe the diet, prey frequency of occurrence (Fi) (Tiberti et al., 2016) was calculated for both
156 seasons.

157

158 *2.4 Trace elements in fish muscle*

159 Trace elements in fish muscle from males and females were analyzed by inductively coupled
160 plasma-mass spectrometry (ICP-MS Xseries II, Thermo Scientific, Bremen, Germany).
161 Determination of Al, As, Cd, Cr, Cu, Fe, Ni, Pb, Se, and Zn was performed following protocols
162 reported by Squadrone et al. (2016). Hg concentration was determined on a direct mercury analyzer
163 (DMA-80 Analyzer, Milestone, Shelton, CT, USA). Analytical performance was verified by
164 processing certified reference materials (Oyster Tissue - SRM 1566b from the National Institute of
165 Standard and Technology), along with blank reagents in each analytical session. Table S1 presents
166 the reference material values and the percentages of recovery. The analytical method was validated
167 according to ISO/IEC 17025 (general requirements for the competence of testing and calibration
168 laboratories).

169

170 2.5 Biochemical analyses

171 A total of 40 specimens of *S. fontinalis*, 26 females (12 in August and 14 in October) and 14 males
172 (8 in summer and 6 in autumn) were examined **individually** for oxidative stress biomarkers. **Liver,**
173 **gills, kidney, spleen, and muscle were analyzed for SOD, CAT, GPx, Se-GPx, GR, and GST**
174 **activity, and MT levels only for muscle.** For **enzymatic analysis, the** samples were homogenized
175 with an UltraTurrax homogenizer in 100 mM potassium-phosphate buffer, pH 7.5, added with 2.5%
176 sodium chloride (NaCl), 0.008 TIU ml⁻¹ aprotinin and 0.1 mg ml⁻¹ bacitracin. The homogenates
177 were centrifuged at 50,000 x g for 30 minutes **at 4°C**. Cytosolic fractions were used to determine
178 antioxidant biomarker **activity**. Biochemical analyzes were performed according to the methods
179 reported in Elia et al. (2017). Briefly, SOD **activity** was assessed at 550 nm in 50 mM Na₂CO₃
180 buffer, pH 10, 0.1 mM EDTA, 500 mM cytochrome C and 1 mM hypoxanthine and xanthine
181 oxidase. Cytochrome C reduction by the xanthine/hypoxanthine complex was evaluated by
182 comparison with a standard SOD unit curve. CAT **activity** was measured at 240 nm after the
183 decrease in absorbance following the consumption of H₂O₂. The assay was carried out in NaH₂PO₄
184 buffer + Na₂HPO₄ 100 mM pH 7 and H₂O₂ 24 mM. Total **glutathione peroxidase (GPx) and**
185 **selenium-dependent glutathione peroxidase (Se-GPx) activity** was determined at 340 nm in
186 NaH₂PO₄ + Na₂HPO₄ 100 mM buffer, pH 7.5, 1 mM EDTA, 0.12 mM NADPH (b-nicotinamide
187 adenine dinucleotide), 2 mM GSH, 1 U of GR (glutathione reductase), 1 mM NaN₃ and H₂O₂ 0.6
188 mM for Se-GPx or 1 mM DTT and 0.8 mM cumene hydroperoxide for GPx. GR **activity** was
189 measured at 340 nm in NaH₂PO₄ + Na₂HPO₄ 100 mM buffer, pH 7, 1 mM GSSG (oxidized
190 glutathione), and 0.06 mM NADPH. GST **activity** was measured at 340 nm using CDNB (1-chloro-
191 2,4-dinitrobenzene) as substrate. The assay was carried out in 100 mM NaH₂PO₄ + Na₂HPO₄ 100
192 mM buffer, pH 6.5, 2 mM GSH and 2 mM CDNB. Concentration of cytosolic proteins was
193 determined according to the method of Lowry et al. (1951) and used to normalize biomarker
194 **activity**.

195 Metallothionein (MT) **levels** were measured in the muscle tissue of individuals of both sexes.
196 Samples were homogenized (1:4) in a buffer containing 0.02 M TRIS/HCl, 0.5 M sucrose, 0.1 mg
197 ml⁻¹ bacitracin, 0.008 tiu ml⁻¹ aprotinin, 87 µg ml⁻¹ phenylmethylsulfonyl fluoride (PMSF), and 0.1
198 µl ml⁻¹ α-mercaptoethanol. The homogenates were centrifuged at 14,500 x g **at 4°C** to obtain the
199 cytosolic fraction. The supernatants were purified using a chloroform/ethanol solution and then
200 HCl/ethanol to obtain the partially purified MT fraction. The pellets were washed with
201 ethanol/chloroform/TRIS/HCl (87/1/12) solution and suspended in 0.25 M NaCl. A destabilizing
202 solution (HCl 1N + EDTA 4 mM) and Ellman's reagent (DTNB: 5,5 dithiobis-2-nitrobenzoic acid)
203 were added to each sample. Sulphydril residue contents (-SH) were spectrophotometrically

quantified. Absorbance was measured at 412 nm and compared to that obtained from a standard curve with reduced glutathione (1 mg ml⁻¹ GSH). All biochemical analyses were performed in triplicate for each sample on a Varian spectrophotometer (Cary 50 Thermostat Cell Holder) at a constant temperature of 25°C.

208

2.6 Statistical analyses

Spearman's rank correlation coefficient (ρ S) was used to test for correlations between trace element concentration in fish muscle, water physicochemical parameters, nutrients, and oxidative stress biomarkers. Differences in the concentration of physicochemical parameters, nutrients, and trace elements between seasons were tested using the Wilcoxon test. Data from the biochemical analysis are reported as the mean and standard deviation (SD). Significant differences between sexes and months were tested using one-way ANOVA followed by Tukey's multiple comparison test. Homogeneity of variance was checked using Bartlett's test. The criterion for significance was set at $p < 0.05$. Principal component analysis (PCA) was performed to check for trends in trace elements, biomarkers, and physicochemical values between the sampling seasons (summer and autumn). Statistical analyses were performed using open source data analysis software RStudio® version 1.1.463 (RStudio, Inc.).

221

3. Results

3.1 Physicochemical and nutrient characteristics of lake water

Lake water temperature was significantly lower in autumn (range 7.20-8.80°C) than summer (range 14.60-16.10°C) (Wilcoxon test; $p = 0.0078$) in agreement with seasonal trends; no thermal stratification in the water column was observed, since the temperatures recorded at the deep sites were similar to those of the littoral sites in both seasons. pH values were significantly higher in autumn (range 7.53-7.90) than summer (range 6.52-7.31) (Wilcoxon test; $p = 0.0078$). No differences in oxygen saturation were recorded between summer (range 77-103%) and autumn (range 101-107%) (Wilcoxon test; $p > 0.05$). Water conductivity remained constant between summer and autumn (range 17-21 $\mu\text{S cm}^{-1}$) (Wilcoxon test; $p > 0.05$). PO_4^{3-} concentration was $\leq 0.02 \text{ mg L}^{-1}$ in both seasons. NH_4^+ level was $< 0.14 \text{ mg L}^{-1}$ at all sites, except for station 4 (0.20 mg L^{-1}) in summer. NO_3^- level was $< 9.20 \text{ mg L}^{-1}$ at all sites, except for site 4 (12 mg L^{-1}) in summer. There were no significant differences in PO_4^{3-} , NH_4^+ , and NO_3^- levels between seasons (Wilcoxon test; $p > 0.05$). Table 1 presents the changes in physicochemical and nutrient data (mean \pm SD).

236

3.2 Fish and stomach contents

237

Table 2 presents the average total length and weight of fish captured during summer and autumn 2017. Stomach contents analysis revealed the almost exclusive presence of terrestrial insect preys in both summer (94.19%) and autumn (81.04%). Preys belonged to the order Hymenoptera (77.74% and 66.39% in summer and autumn, respectively) and Coleoptera (16.45% and 14.65% in summer and autumn, respectively). Other taxa were present in extremely low proportions (Diptera Chironomidae larvae: 5.18% in summer and 2.79% in autumn).

3.3 Trace elements in fish muscle

Figure 2 presents the mean concentration and the relative standard deviation of each trace element detected in muscle tissue in summer and autumn. The mean concentration of trace elements was in the order: Zn (16.25) > Fe (8.78) > Al (1.49) > Se (0.67) > Cu (0.60) > Cr (0.14) > As (0.06) > Ni (0.06) > Pb (0.05) > Hg (0.01) mg Kg⁻¹. In autumn the mean concentration of trace elements was in the order: Zn (16.13) > Fe (10.63) > Al (2.03) > Se (0.49) > Cu (0.36) > Cr (0.16) > As (0.10) > Pb (0.10) > Ni (0.06) > Hg (0.02) mg Kg⁻¹. Cd was < LOQ (0.02 mg Kg⁻¹) in both seasons. There were no significant differences in trace element concentration between seasons (Wilcoxon test; $p > 0.05$ for all elements).

3.4 Biochemical analyses

In the liver, the activity of SOD, CAT, and GR was significantly lower in autumn (up to 70%) than summer in males and females (Tukey's test; $p < 0.05$) (Figs. 3-4, 7). In the gills, GPx activity was significantly higher in autumn (90%) mainly in the females (Tukey's test; $p < 0.05$); Se-GPx and GST activity was significantly higher (up to 2-fold) in males and females in autumn (Tukey's test; $p < 0.05$) (Figs. 5, 6, 8). In the kidney, only Se-GPx activity was significantly different between the seasons, being higher (up to 40%) in autumn (Tukey's test; $p < 0.05$) (Fig. 6). In the spleen, Se-GPx activity was significantly lower (70%) in the females (Tukey's test; $p < 0.05$) (Fig. 6). GR and GST activity was significantly higher (up to 70%) in males in autumn than in summer (Tukey's test; $p < 0.05$) (Fig. 7, 8). In muscle, SOD activity was significantly lower (up to 50%) in females in autumn (Tukey's test; $p < 0.05$), whereas CAT activity showed an opposite trend and was significantly higher in autumn (up to one-fold) (Tukey's test; $p < 0.05$) (Figs. 3-4). MT level was higher in autumn (up to one-fold) than in summer (Tukey's test; $p < 0.05$) (Fig. 9).

3.5 Spearman correlation matrix

Spearman correlation matrix revealed correlations between environmental parameters (trace elements, physicochemical parameters, nutrients) and oxidative stress biomarkers in muscle, gills,

liver, spleen, and kidney tissue for both seasons. Due to the multiplicity of positive correlations, Table S2 presents the correlation matrices (one for each organ). Only the most informative outcomes are presented and discussed for interpretation of biological response of *S. fontinalis* to environmental parameters. In detail, a significant positive correlation was found between MTs and Hg (ρ_S 0.787), MTs and Pb (ρ_S 0.787), MTs and Cu (ρ_S 0.683), MTs and pH (ρ_S 0.650), and CAT and pH (ρ_S 0.737) in muscle tissues of females captured in autumn; SOD and Cr (ρ_S 0.837) in the liver tissue of females captured in autumn; SOD and NO_3^- (ρ_S 0.750), Se-GPx and NO_3^- (ρ_S 0.750) in muscle and kidney tissue, respectively, of females captured in summer.

280

3.6 Principal Component Analysis (PCA)

The first two principal components (Dim1; Dim2) accounted for meaningful amounts of the total variance (58.2%), while the other components accounted for a relatively smaller fraction (Fig. 10). Dim1 accounted for 35.2% of the total variance and was positively correlated with the variables Ni, Se, temperature, oxygen, conductivity, and NO_3^- and negatively correlated with Al, Cr, Fe, As, Pb, Hg, and pH. Dim2 accounted for 23% of the total variance and was positively correlated with the variables GST, CAT, GPx, Se-GPx, SOD, and GR. The biplot of loadings (variables) and score (observations) shows which organs (liver, gills, kidney, spleen, and muscle) of brook trout are closest to them, and which variables (trace elements, biomarkers and physicochemical parameters) contributed to this grouping in the coordinate of Dim1 and Dim2. Separation of organs by season (summer on the right, autumn on the left) suggested a difference in biomarker values. In detail, the autumn samples of spleen, gill, muscle, kidney, and liver tissue are on the left in order of increasing value of Al, Cr, Fe, As, Pb, Hg, and pH. The summer samples of spleen, gill, muscle, kidney tissue are on the right in order of increasing value of NH_4^+ , NO_3^- , conductivity, oxygen, temperature, Zn, Cu, Ni, and Se. Remarkably, summer samples of liver tissue are well separated from other organs, following the trend in biomarker values.

297

4. Discussion

The environmental parameters of water oxygenation, pH, conductivity, and temperature are in line with those reported by Tiberti et al. (2010) for 12 alpine lakes in the Gran Paradiso National Park (Western Alps, Italy). Temperatures were lower in autumn than in summer according to seasonality. The temperature data recorded at the deep sites revealed no vertical layering in the lake because the shallow maximum depth (6.42 m) does not allow for the formation of a thermocline layer. The pH values are related to rock composition. Since Lake Bama lies over a granite bedrock, its pH values are lower than lakes on limestone or sandstone bedrock (Camarero et al., 2009). The pH values

were in line with the literature reported for high-altitude environments (Boggero et al., 2006; Füreder et al., 2006; Fjellheim et al., 2009). As expected for mountain lakes, the oxygenation levels were high. The water conductivity values were in line with published literature, as conductivity of silty-like lakes tends to be $< 50 \mu\text{S cm}^{-1}$ (Boggero et al., 2006; Füreder et al., 2006). No decrease in oxygen values correlated with depth was observed owing to the absence of temperature stratification. Nutrient levels (PO_4^{3-} , NO_3^- and NH_4^+) were in line with previous studies of Alpine lakes (Füreder et al., 2006; Camarero et al., 2009) and revealed an oligotrophic condition.

We observed a positive correlation between environmental parameters and biological response in the *S. fontinalis* from Balma Lake. The antioxidant response was tissue-specific; indeed, aerobic tissues such as the kidney, gills, spleen, and the liver in particular have a high potential for ROS production which is offset by protective mechanisms. Differently, muscle, which has a low content of mitochondria and a low-intensive oxidative metabolism, showed mild response to oxidative/reductive conditions. Moreover, the level of several biomarkers was related to the seasonal changes in the trace element concentration in some tissues.

The few studies on trace element accumulation in the biota from high-altitude lakes in general, and in Alpine lakes in particular, have focused largely on fish. Köck et al. (1996) studied the concentrations of Cd, Pb, Zn, and Cu in the muscle of arctic char (*Salvelinus alpinus*) from five oligotrophic Alpine lakes in northern Tyrol (Austria). Yang et al. (2007) studied the accumulation of trace elements in muscle of fish of the genus *Gymnocypris* (Cyprinidae) from high-mountain lakes in the Tibetan Plateau. Ni ranged from 0.094 to 0.12 mg kg^{-1} , Cu from 1.1 to 2.0 mg kg^{-1} , Zn from 4.4. to 6.9 mg kg^{-1} , As from 0.24 to 0.27 mg kg^{-1} , Se from 0.36 to 1.0 mg kg^{-1} , Cd from 0.024 to 0.025 mg kg^{-1} , and Pb from 0.047 to 0.079 mg kg^{-1} . Rognerud et al. (2002) found that Hg concentrations in fish muscle from high-mountain lakes in Europe ranged from 0.021 to 0.179 mg kg^{-1} . These data from previous studies demonstrate that high-mountain lakes function as a regional contaminant convergence zone for the medium and long-range atmospheric transport of contaminants. Since our results are in line with these findings, and because no previous studies have been performed to date, we assume that the trace element concentrations we detected in the brook trout from Balma Lake have both anthropogenic and pedogenic sources as their origin.

Metallothioneins have been widely considered as valuable biomarkers that reflect the level of trace elements in aquatic environments, where they act as metal trappers mainly of Cd, Hg, Pb, Cu, and Zn in fish (Bourdineaud et al., 2006; Morcillo et al., 2016). The higher levels of MTs we measured in autumn were related to the increased trace element concentration in this tissue, and we noted a strong correlation with Cu, Hg, and Pb levels in the females. This may suggest an alarming scenario, as it would signal an increased concentration of these elements in the environment.

340 Although no chemical analyses of the lake water were carried out, because of the peculiar
341 geomorphological characteristics of Balma Lake we can exclude an increase in such contaminants
342 during autumn.

343 Seasonal variation in metal concentrations in fish can be related to environmental factors such as
344 food availability, temperature, and biological cycle (Hermesz et al., 2001; Amiard et al., 2006;
345 Dragun et al., 2009). Furthermore, fish size may also play a key role in metal uptake (Wright and
346 Mason, 1999). In the present study, an increase in weight and length was measured in the fish from
347 both sampling seasons, and the frequency of occurrence of terrestrial invertebrates in the diet during
348 the ice-free season was in accordance with other studies performed on brook trout in other Alpine
349 lakes (Sotiropoulos et al., 2006; Tiberti et al., 2016). It is possible that fish growth due to a higher
350 intake of food in summer could have favored the uptake of heavy metals such as Pb and Hg.

351 This hypothesis is corroborated by a previous study on black scabbardfish *Aphanopus carbo*
352 (Trichiuridae), in which an exponential increase in total Hg load was found in all fish tissues in a
353 length-dependent manner (Bebianno et al., 2007). Furthermore, previous studies showed that
354 fluctuations toward high pH values can also play a key role in modulating metals uptake, affecting
355 their speciation and bioavailability (Playle, 1998). This outcome may explain the strong correlation
356 between pH and the two hydroxyl scavengers MTs and CAT in the female muscle tissue in autumn.

357 Trace elements also affected the activity of several enzymes in different tissues in the autumn fish
358 samples. We noted a strong correlation between Cr and SOD activity in the liver, as reported in
359 previous study in rock fish *Sebastes schlegelii* (Kim and Kang, 2016). Furthermore, a recent study
360 showed that Cr can induce conformational changes of CAT enzyme and reduce its activity
361 depending on its valence states and concentration (Chen et al., 2018). These findings may explain
362 the seasonal difference in CAT activity in the female muscle tissue.

363 In general, numerous elements can influence the activity of this biomarker, and As and Fe were the
364 elements most involved in modulating CAT enzyme activity in the tissues. Arsenic is a global
365 contaminant derived from natural or anthropogenic sources and a cause of great concern for
366 terrestrial and aquatic ecosystems (Elia et al., 2018). At high concentrations, arsenic may induce
367 oxidative stress by interacting with antioxidants and result in the accumulation of free radicals in
368 cells. Arsenite species can interact with sulfhydryl groups of biomolecules such as enzymes or
369 reduced glutathione (Elia et al., 2018). Furthermore, redox active metals such as iron generate ROS
370 or are involved in the Fenton route, leading to lipid peroxidation (Mahboob, 2013). Despite the
371 increase in trace element amount in autumn, the concentration of Pb, Hg, and Cd in fish muscle
372 mentioned in Regulation 1881/2006 (European Commission, 2006) was far below the established
373 threshold limit. This fact should be taken into account and may suggest the adaption of *S. fontinalis*

374 to seasonal changes rather than to oxidative stress. PCA analysis showed that summer samples of
375 liver tissue were **strongly** related to oxidative biomarker **level**, since liver tissue is the site of
376 multiple oxidative reactions and maximal free radical generation (Gul et al., 2004; Avci et al.,
377 2005).

378 For ectothermic organisms, temperature is a crucial abiotic factor (Hassan et al., 2017). Daily
379 temperature fluctuations and seasonal variations differ in their influence on fish (Buckley et al.,
380 2006; Place and Hofmann, 2004; Podrabsky and Somero, 2004). Wellness and growth are optimal
381 within a well-defined temperature range depending on the species (Godowsky and Caddell, 1991).
382 Temperatures deviating excessively from the optimum can exert harmful effects and induce
383 mortality (Rijnsdorp et al., 2009). In the present study, the marked drop in water temperature in
384 autumn was related to changes mainly in GPx activity in the male kidney and in the female spleen.
385 Moreover, higher temperature results in increased GPx **activity in other fish such** as the Antarctic
386 *Pagothenia borchgrevinki* (Almroth et al., 2015). Thermal changes have also been associated with
387 the up regulation of the NRF2 transcription factor, which is involved in the expression of
388 antioxidants via binding to the antioxidant responsive element (ARE) (Almroth et al., 2015).
389 Elevated GPx activity during summer may indicate strengthening of this fundamental defense line
390 against ROS. However, the lack of change in SOD and CAT activity, as well as the lower
391 concentration of trace elements in tissue, and the constant levels of the main **physicochemical**
392 parameters of water, except for pH and temperature, preclude an oxidative pressure scenario and
393 suggest an adaptive ability of *S. fontinalis* to higher temperature instead. On the other hand,
394 increased SOD activity in the muscle tissue of males and females is linked to an abiotic factor, such
395 as NO_3^- , in summer. Furthermore, nutrients also influenced SOD concentration in female muscle
396 and Se-GPx activity in female kidney in summer. In the aquatic environment, it is not unusual that
397 fish are simultaneously challenged by different abiotic factors. Conversely, at very high
398 concentrations, ammonia can induce a range of toxicological effects in fish, such as altered
399 metabolism, lack of growth, and mortality (Dosdat et al., 2003; Sinha et al., 2012, 2015). Ammonia
400 exposure can also lead to oxidative stress in fish (Sun et al., 2012; Sinha et al., 2014). A previous
401 study showed that conductivity plays a crucial role in maintaining the ammonia ionization
402 equilibrium (NH_3 and the non-toxic form NH_4^+) in aquatic environments (Sinha et al., 2015). In the
403 present study, the lack of change in conductivity and nutrients (e.g., phosphorus) between the
404 seasons suggests that the changes in antioxidant parameters may be related to food intake (**which**
405 **indirectly promotes metals uptake**) rather than to nutrient concentration. In their habitats, high-
406 mountain lakes included, fish are often exposed to periods of food insufficiency in response to
407 factors such as temperature, conductivity, and biological cycle (Pérez-Jiménez et al., 2007; Furné et

408 al., 2009). The increase in food intake during summer may also explain the fluctuation in
409 biomarkers of oxidative stress.

410

411 5. Conclusion

412 With this study we investigated the influence of trace element accumulation and environmental
413 parameters on oxidative stress biomarkers in male and female individuals of *S. fontinalis* from a
414 high-mountain lake during the ice-free period. Generally, positive correlations were found between
415 several environmental parameters and biomarkers. While oxidative stress biomarker levels were
416 similar for males and females, significant fluctuation between seasons due to biological and
417 environmental pressures was noted for several biomarkers. Although there was greater fluctuation
418 in temperature, pH, and trace elements (e.g., Cu, Pb and Hg) between seasons, which certainly
419 contributed to changes in biomarker levels, our findings indicate that increased food intake during
420 the ice-free season was probably the main factor that affected the oxidative stress response. Future
421 studies are needed to investigate other factors responsible for the changes in oxidative stress
422 biomarkers.

423

424 Acknowledgements

425 The authors would like to thank GeoStudio RC (Marco Rosa Clot, Luca Rosa Clot), Club Alpino
426 Italiano (Alfio Usseglio) and Ente di Gestione delle Aree Protette delle Alpi Cozie (Michele Ottino)
427 for their technical support.

428

429 References

430 Al Kaddissi, S., Legeay, A., Elia, A.C., Gonzalez, P., Floriani, M., Cavalieri, I., Massabuaum, J.C.,
431 Gilbin, R., Simon, O., 2014. Mitochondrial gene expression, antioxidant responses, and
432 histopathology after cadmium exposure. Environ. Toxicol. 29, 893-907.

433 Almroth, B.C., Asker, N., Wassmur, B., Rosengren, M., Jutfelt, F, Gräns, A., Sundell, K., Axelsson,
434 M., Sturve, J., 2015. Warmer water temperature results in oxidative damage in an Antarctic fish, the
435 bald notothen. J. Exp. Mar. Biol. Ecol. 468, 130-137.

436 Amiard, J.C., Amiard-Triquet, C., Barka, S., Pellerin, J., Rainbow, P.S., 2006. Metallothioneins in
437 aquatic invertebrates: their role in metal detoxification and their use as biomarkers. Aquat. Toxicol.
438 76, 160-202.

439 APHA, AWWA, WEF, 1998. Standard Methods for the Examination of Water and Wastewater,
 440 20th edition. American Public Health Association, Washington DC.

441 ASTM, 2015. Manual of Water and Environmental Technology, D1426-92, Nessler method.

442 Avci, A., Kačmaz, M., Durak, I., 2005. Peroxidation in muscle and liver tissues from fish in a
 443 contaminated river due to a petroleum refinery industry. *Ecotoxicol. Environ. Safe.* 60, 101-105.

444 Barwick, M., Maher, W., 2003. Biotransference and biomagnification of selenium copper,
 445 cadmium, zinc, arsenic and lead in a temperate seagrass ecosystem from Lake Macquarie Estuary,
 446 NSW, Australia. *Mar. Environ. Res.* 56, 471-502.

447 Bebianno, M.J., Santosa, C., Canário, J., Gouveia, N., Sena-Carvalho, D., Vale, C., 2007. Hg and
 448 metallothionein-like proteins in the black scabbardfish *Aphanopus carbo*. *Food. Chem. Toxicol.* 45,
 449 1443-1452.

450 Boggero, A., Füreder, L., Lencioni, V., Simcic, T., Thaler, B., Ferrarese, U., Lotter, A.F., Ettinger,
 451 R., 2006. Littoral chironomid communities of Alpine lakes in relation to environmental factors.
 452 *Hydrobiologia* 562, 145-165.

453 Bourdineaud J.P., Baudrimont M., González P., Moreau J.L., 2006. Challenging the model for
 454 induction of metallothionein gene expression. *Biochimie* 88, 1787-1792.

455 Buckley, B.A., Gracey, A.Y., Somero, G.N., 2006. The cellular response to heat stress in the goby
 456 *Gillichthys mirabilis*: a cDNA microarray and protein level analysis. *J. Exp. Biol.* 209, 2660-2677.

457 Camarero, L., Rogora, M., Mosello, R., Anderson, N.J., Barbieri, A., Botev, I., Kernan, M.,
 458 Kopáček, J., Korhola, A., Lotter, A.F., Muri, G., Postolache, C., Stuchik, E., Thies, H., Wright,
 459 R.F., 2009. Regionalization of chemical variability in European mountain lakes. *Freshw. Biol.* 54,
 460 2452-2469.

461 Catalán, J., Camarero, L., Felip, M., Pla S., Ventura, M., Buchaca, T., Bartumeus, F., De Mendoza,
 462 G., Miró, A., Casamayor, E.O., Medina-Sánchez, J.M., Bacardit, M., Altuna, M., Bartrons, M.,
 463 Díaz, de Quijano, D., 2006. High mountain lakes: extreme habitats and witnesses of environmental
 464 changes. *Limnetica* 25, 551-584.

465 Chen, L., Zhang, J., Zhu, Y., Zhang, Y., 2018. Interaction of chromium (III) or chromium (VI) with
 466 catalase and its effect on the structure and function of catalase: An in vitro study. Food. Chem. 244,
 467 378-385.

468 Cozzari, M., Elia, A.C., Pacini, N., Smith, B.D., Boyle, D., Rainbow, P.S., Khan, F.R., 2015.
 469 Bioaccumulation and oxidative stress responses measured in the estuarine ragworm (*Nereis*
 470 *diversicolor*) exposed to dissolved, nano- and bulk-sized silver. Environ. Pollut. 198, 32-40.

471 Dosdat, A., Ruyet, J.P., Coves, D., Dutto, G., Gasset, E., Le Roux, A., Lemarié, G., 2003. Effect of
 472 chronic exposure to ammonia on growth, food utilization and metabolism of the European sea bass
 473 (*Dicentrarchus labrax*). Aquat. Living Resour. 16, 509-520.

474 Dragun, Z., Podrug, M., Raspor, B., 2009. The assessment of natural causes of metallothionein
 475 variability in the gills of European chub (*Squalius cephalus* L.). Comp. Biochem. Physiol. C:
 476 Toxicol. Pharmacol. 150, 209e217.

477 Elia A.C., Dörr A.J.M, Abete M.C., Prearo M., 2010. Seasonal variability of detoxificant response
 478 and heavy metal accumulation in tissues of both sexes in *Tinca tinca* (L.) from Lake Trasimeno.
 479 Rev. Fish. Biol. Fisher. 20(3), 425-434.

480 Elia A.C., Magara G., Caruso C., Masoero L., Prearo M., Arsieni P., Caldaroni B., Scoparo M.,
 481 Dörr A.J.M., Salvati S., Brizio P., Squadrone S., Abete M.C., 2018. A comparative study on
 482 subacute toxicity of arsenic trioxide and dimethylarsinic acid on antioxidants and antioxidant-
 483 related enzymes in crandell rees feline kidney (CRFK), human hepatocellular carcinoma
 484 (PCL/PRF/5) and epithelioma papulosum cyprini (EPC) cell lines. J. Toxicol. Environ. Health part
 485 A 81(10), 333-348.

486 Elia, A.C., Dörr, A.J.M., Galarini, R., 2007a. Comparison of organochlorine pesticides, PCBs and
 487 heavy metal contamination and of detoxifying response in tissues of *Amerius melas* from Corbara,
 488 Alviano and Trasimeno Lakes, Italy. Bull. Environ. Contam. Toxicol. 78, 463-468.

489 Elia, A.C., Galarini, R., Dörr, A.J.M., Taticchi MI., 2007b. Heavy metal contamination and
 490 antioxidant response of a freshwater bryozoan (*Lophopus crystallinus* Pall., Phylactolaemata).
 491 Ecotox. Environ. Safe. 66, 188-194.

492 Elia, A.C., Galarini, R., Dörr, A.J.M., Taticchi MI. 2006. Bioaccumulation of heavy metals,
 493 organochlorine pesticides, and detoxification biochemical indexes in tissues of *Ictalurus melas* of
 494 Lake Trasimeno. Bull. Environ. Contam. Toxicol. 76, 132-139.

495 Elia, A.C., Magara, G., Righetti, M., Dörr, A.J.M., Scanzio, T., Pacini, N., Abete, M.C., Prearo, M.,
 496 2017. Oxidative stress and related biomarkers in cupric and cuprous chloride-treated rainbow trout.
 497 Environ. Sci. Pollut. R. 24, 10205-10219.

498 European Commission, 2006. Commission Regulation (EC) No 1881/2006 of 19 December 2006
 499 setting maximum levels for certain contaminants in foodstuffs. OJEC 364, 365-324.

500 Ferrario, C., Finizio, A., Villa S., 2017. Legacy and emerging contaminants in meltwater of three
 501 alpine glaciers. Sci. Total. Environ. 574, 350-357.

502 Fjellheim, A., Raddum, G.G., Vandvik, V., Cogălniceanu, D., Boggero, A., Brancelj, A., Galas, J.,
 503 Sporka, F., Vidinova, Y., Bitusik, P., Dumnicka, E., Gâldean, N., Kownacki, A., Krno, I., Preda, E.,
 504 Rîșnoveanu, G., Stuchlik, E., 2009. Diversity and distribution patterns of benthic invertebrates
 505 along alpine gradients. A study of remote European freshwater lakes. Adv. Limnol. 62, 167-190.

506 Füreder, L., Ettinger, R., Boggero, A., Thaler, B., Thies, H., 2006. Macroinvertebrate diversity in
 507 Alpine lakes: effects of altitude and catchment properties. Hydrobiologia 652, 123-144.

508 Furné, M., García-Gallego, M., Hidalgo, M.C., Morales, A.E., Domezain, A., Domezain, J., Sanz,
 509 A., 2009. Oxidative stress parameters during starvation and refeeding periods in Adriatic sturgeon
 510 (*Acipenser naccarii*) and rainbow trout (*Oncorhynchus mykiss*). Aquaculture Nutr. 15, 587-595.

511 Godowsky, D.M., Caddell, S.M., 1991. Effects of temperature on early-life-history stages of
 512 California halibut *Paralichthys californicus*. Fish Bulletin 89, 567-576.

513 Gul, S., Belge-Kurutas, E., Yildiz, E., Sahan, A. Doran, F., 2004. Pollution correlated modifications
514 of liver antioxidant systems and histopathology of fish (Cyprinidae) living in Seyhan Dam Lake,
515 Turkey. Environ. Int. 30, 605-609.

516 Hassan, M.K., Nahid, N.M., Bahar, A.N., Bhuiyan, M.M.R., Abdullah-Al-Shafi, M., Ahmed, K.,
517 2017. Dataset demonstrating the temperature effect on average output polarization for QCA based
518 reversible logic gates. Data in Brief, Elsevier 13, 713–716.

519 Hermes, E., Abrahm, M., Nemcsok, J., 2001. Tissue-specific expression of two metallothionein
520 genes in common carp during cadmium exposure and temperature shock. Biochem. Physiol. Part C
521 128, 457–465.

522 Hung, H., Katsoyianni, A.A., Brorström-Lundén, E., Olafsdottir, K., Aas, W., Breivik, K., Bohlin-
523 Nizzetto, P., Sigurdsson, A., Hakola, H., Bossi, R., Skov, H., Sverko, E., Barresi, E., Fellin, P.,
524 Wilson, S., 2016. Temporal trends of Persistent Organic Pollutants (POPs) in arctic air: 20 years of
525 monitoring under the Arctic Monitoring and Assessment Programme (AMAP). Environ. Pollut.
526 217, 52-61.

527 Kim, J.H., Kang, J.C., 2016. Oxidative stress, neurotoxicity, and metallothionein (MT) gene
528 expression in juvenile rock fish *Sebastes schlegelii* under the different levels of dietary chromium
529 (Cr^{6+}) exposure. Ecotox. Environ. Safe. 125, 78-84.

530 Köck, G., Triendl, M., Hofer, R., 1996. Seasonal patterns of metal accumulation in Arctic char
531 (*Salvelinus alpinus*) from an oligotrophic Alpine lake related to temperature. Can. J. Fish. Aquat.
532 Sci. 53, 780-786.

533 Lowry, O.H., Rosebrough, N.J., Farr A.L., Randall R.J., 1951. Protein measurement with the Folin
534 phenol reagent. J. Biol. Chem. 193, 265–75.

535 Lushchak, V.I., 2011. Environmentally induced oxidative stress in aquatic animals. Aquat. toxicol.
536 101(1), 13-30.

537 Lushchak, V.I., 2016. Contaminant-induced oxidative stress in fish: a mechanistic approach. Fish
538 Physiol. Biochem. 42, 711-747.

539 Mahboob, S., 2013. Environmental pollution of heavy metals as a cause of oxidative stress in fish: a
 540 review. *Life Sci. J.* 10(10s), 336-345.

541 Milardi, M.S. Siitonen, J., Lappalainen, A, Liljendahl, Weckström, J., 2016. The impact of trout
 542 introductions on macro- and micro-invertebrate communities of fishless boreal lakes. *J.*
 543 *Paleolimnol.* 55, 273-287.

544 Monserrat, J., Martinez, P., Geracitano, L., Amado, L., Martins, C., Pinho, G., Chaves, I., Cravo,
 545 M., Lima, J., Bianchini, A., 2007. Pollution biomarkers in estuarine animals: critical review and
 546 new perspectives. *Comp. Biochem. Physiol.* 146C, 221-234.

547 Morcillo, P., Esteban, M.A., Cuesta, A., 2016. Heavy metals produce toxicity, oxidative stress and
 548 apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere* 144, 225-233.

549 Pastorino P., Pizzul E., Bertoli M., Perilli S., Brizio P., Salvi G., Esposito G., Abete M.C., Prearo
 550 M., Squadrone S. (2019b). Macrobenthic invertebrates as bioindicators of trace elements in high-
 551 mountain lakes. *Environ. Sci. Pollut. Res.* <https://doi.org/10.1007/s11356-019-07325-x>

552 Pastorino, P., Polazzo, F., Bertoli, M., Santi, M., Righetti, M., Pizzu, E., Prearo, M., 2020.
 553 Consequences of fish introduction in fishless Alpine lakes: preliminary notes from a sanitary point
 554 of view. *Turk. J. Fish Aquat. Sc.* 20(1), 1-8.

555 Pastorino, P., Prearo, M., Pizzul, E., Bertoli, M., Francese, D.R., Menconi, V., Mugetti D., Bozzetta
 556 E., Varello, K. (2019a). Hepatic Steatosis in a Bullhead (*Cottus gobio*) Population from a High-
 557 Mountain Lake (Carnic Alps): Adaptation to an Extreme Ecosystem? *Water* 11(12), 2570.

558 Pérez-Jiménez, A., Guedes, M.J., Morales, A.E., Oliva-Teles, A., 2007. Metabolic responses to
 559 short starvation and refeeding in *Dicentrarchus labrax*. Effect of dietary composition. *Aquaculture*
 560 265, 325-335.

561 Perrine, S., 2017. Inferring fish abundance in lakes from subfossil midge (Diptera: Chironomidae)
 562 remains. University of Jyväskylä, Department of Biological and Environmental Science, Master
 563 Thesis, Jyväskylä, Finland.

564 Place, S.P., Hofmann, G.E., 2004. Temperature interactions of the molecular chaperone Hsc70 from
565 the eurythermal marine goby *Gillichthys mirabilis*. J. Exp. Biol. 204, 2675–2682.

566 Playle, R.C., 1998. Modelling metal interactions at fish gills. Sci. Total. Environ. 219, 147-163.

567 Podrabsky, J.E., Somero, G.N., 2004. Changes in gene expression associated with acclimation to
568 constant temperatures and fluctuating daily temperatures in an annual killifish, *Austrofundulus*
569 *limnaeus*. J. Exp. Biol. 207, 2237-2254.

570 Poma, G., Salerno, F., Roscioli, C., Novati, S., Guzzella, L., 2017. Persistent organic pollutants in
571 sediments of high-altitude Alpine ponds within Stelvio National Park, Italian Alps. Inland Waters 7,
572 34-44.

573 Raposeiro, P.M., Rubio M.J, González A., Hernández A., Sánchez-López G., Vázquez-Loureiro D.,
574 Rull V., Bao R., Costa A. C., Gonçalves V., Saéz A., Giralte S., 2017. Impact of the historical
575 introduction of exotic fishes on the chironomid community of Lake Azul (Azores Islands).
576 Palaeogeogr Palaeocl. 466, 77–88.

577 Rijnsdorp, A.D., Peck, M.A., Engelhard, G.H., Mollmann, C., Pinnegar, J.K., 2009. Resolving the
578 effect of climate change on fish populations. ICES J. Mar. Sci. 66, 1570-1583.

579 Rognerud, S., Grimalt, J.O., Rosseland, B.O., Fernandez, P., Hofer, R., Lackner, R., et al. 2002.
580 Mercury and organochlorine contamination in Brown trout (*Salmo trutta*) and Arctic charr
581 (*Salvelinus alpinus*) from high mountain lakes in Europe and the Svalbard archipelago. Water Air
582 Soil Pollut. 2, 209-32.

583 Rott, E., 1988. Some aspects of the seasonal distribution of flagellates in mountain lakes.
584 Hydrobiologia 161, 159-170.

585 Sinha, A.K., Abdelgawad, H., Giblen, T., Zinta, G., De Rop, M., Asard, H., Blust, R., De Boeck G.,
586 2014. Anti-oxidative defences are modulated differentially in three freshwater teleosts in response
587 to ammonia-induced oxidative stress. PLoS One, 9(4), e95319.

588 Sinha, A.K., Liew, H.J., Diricx, M., Blust, R., De Boeck, G., 2012. The interactive effects of
 589 ammonia exposure, nutritional status and exercise on metabolic and physiological responses in
 590 goldfish (*Carassius auratus* L.). *Aquat. Toxicol.* 109, 33–46.

591 Sinha, A.K., Rasoloniriana, R., Dasan, A.F., Pipralia, N., Blust, R., De Boeck, G., 2015. Interactive
 592 effect of high environmental ammonia and nutritional status on ecophysiological performance of
 593 European sea bass (*Dicentrarchus labrax*) acclimated to reduced seawater salinities. *Aquat.*
 594 *Toxicol.* 160, 39–56.

595 Schmid, P., Kohler, M., Gujer, E., Zennegg, M., Lanfranchi, M., 2007. Persistent organic
 596 pollutants, brominated flame retardants and synthetic musks in fish from remote alpine lakes in
 597 Switzerland. *Chemosphere* 67(9), S16-S21.

598 Sommaruga, R., 2001. The role of solar UV radiation in the ecology of alpine lakes. *J. Photochem.*
 599 *Photobiol.* 62, 35-42.

600 Sotiropoulos, J.C., Nislow, K.H., Ross M.R., 2006. Brook trout, *Salvelinus fontinalis*, microhabitat
 601 selection and diet under low summer stream flows. *Fisheries Manag. Ecol.* 13, 149-155.

602 Squadrone, S., Brizio, P., Stella, C., Prearo, M., Pastorino, P., Serracca, L., Ercolini, C., Abete,
 603 M.C., 2016. Presence of trace metals in aquaculture marine ecosystems of the northwestern
 604 Mediterranean Sea (Italy). *Environ. Poll.* 215, 77-83.

605 Squadrone, S., Ciccotelli, V., Favaro, L., Scanzio, T., Prearo, M., Abete, M.C., 2014. Fish
 606 consumption as a source of human exposure to perfluorinated alkyl substances in Italy: Analysis of
 607 two edible fish from Lake Maggiore. *Chemosphere* 114, 181-186.

608 Squadrone, S., Prearo, M., Brizio, P., Gavinelli, S., Pellegrino, M., Scanzio, T., Guarise, S.,
 609 Benedetto, A., Abete, M.C., 2013. Heavy metals distribution in muscle, liver, kidney and gill of
 610 European catfish (*Silurus glanis*) from Italian Rivers. *Chemosphere* 90, 358-365.

611 Sroda, S., Cossu-Leguille, C., 2011. Seasonal variability of antioxidant biomarkers and energy
 612 reserves in the freshwater gammarid *Gammarus roeseli*. *Chemosphere* 83(4), 538-544.

613 Starkweather, P.L., 1990. Zooplankton community structure of high elevation lakes: biogeographic
 614 and predator-prey interactions. Internationale Vereinigung für Theoretische und Angewandte
 615 Limnologie: Verhandlungen 24, 513-517.

616 Sun, H., Lü, K., Minter, E.J., Chen, Y., Yang, Z., Montagnes, D.J., 2012. Combined effects of
 617 ammonia and microcystin on survival, growth, antioxidant responses, and lipid peroxidation of
 618 bighead carp *Hypophthalmichthys nobilis* larvae. J. Hazard. Mater. 221-222, 213-219.

619 Tiberti, R., Tartari, G., Marchetto, A., 2010. Geomorphology and hydrochemistry of 12 Alpine
 620 lakes in the Gran Paradiso National Park, Italy. J. Limnol. 69, 242-256.

621 Tiberti, R., von Hardenberg, A., Bogliani, G., 2014. Ecological impact of introduced fish in high
 622 altitude lakes: a case of study from the European Alps. Hydrobiologia 724, 1-19.

623 Tiberti, R., Brighenti, S., Canedoli, C., Iacobuzio, R., Rolla, M., 2016. The diet of introduced brook
 624 trout (*Salvelinus fontinalis*; Mitchill, 1814) in an alpine area and a literature review on its feeding
 625 ecology. J. Limnol. 75, 488-507.

626 Wright, P., Mason, C.F., 1999. Spatial and seasonal variation in heavy metals in the sediments and
 627 biota of two adjacent estuaries, the Orwell and the Stour, in eastern England', Sci. Total. Environ.
 628 226, 139–156.

629 Yang, R., Yao, T., Xu, B., Jiang, G., Xin, X., 2007. Accumulation features of organochlorine
 630 pesticides and heavy metals in fish from high mountain lakes and Lhasa River in the Tibetan
 631 Plateau. Environ. Int. 33, 151–156.

632
 633
 634
 635
 636
 637
 638

639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664

Table 1. Mean and standard deviation (three replicates) of physicochemical parameters and nutrients measured in Balma Lake in summer (August) and autumn (Autumn) 2017.

665
666
667

	Summer (August)	Autumn (October)
Temperature (°C)	15.62 ± 0.77	8.24 ± 0.53
Dissolved oxygen concentration (mg L ⁻¹)	7.40 ± 0.89	7.29 ± 1.92
Oxygen saturation (%)	92.88 ± 8.52	104.75 ± 2.43
pH	6.64 ± 0.28	7.69 ± 0.67
Conductivity (µS cm ⁻¹)	18.29 ± 1.12	18.56 ± 1.05
NH ₄ ⁺ (mg L ⁻¹)	0.09 ± 0.07	0.09 ± 0.05
NO ₃ ⁻ (mg L ⁻¹)	8.55 ± 2.73	6.82 ± 2.14
PO ₄ ³⁻ (mg L ⁻¹)	0.01 ± 0.01	0.02 ± 0.01

677
678
679
680
681

Table 2. Biometric values of females and males of brook trout (*Salvelinus fontinalis*) from Balma Lake in summer and autumn 2017.

	Summer (August)		Autumn (October)	
Sex	Males	Females	Males	Females
Number of individuals	8	12	6	14
Weight - W mean ± SD (g)	69.9 ± 30.34	71.20 ± 28.34	88.47 ± 56.23	90.05 ± 60.03
W min. (g)	15.00	15.36	10.23	11.00
W max. (g)	146.00	150.00	187.00	193.00
Total Length - TL mean ± SD (cm)	17.77 ± 3.90	18.56 ± 4.20	19.23 ± 4.66	19.35 ± 5.72
TL min. (cm)	11.50	11.00	7.50	8.00
TL max. (cm)	23.00	24.00	25.50	26.00

690
691
692
693

694

695

696

697

698

699

700

701

702

703


704

705

706

707 **Figure 1.** Balma Lake: bathymetry and sampling sites for physicochemical parameters (L = littoral
708 sites; B = deep sites). Red rectangles indicate the placement of gillnets to sample fish in summer
709 (August) and autumn (October) 2017.

710



711

712

713

714

715

716

717 **Figure 2.** Trace elements (Log₁₀ concentration; mg kg⁻¹) detected in the muscle tissue of brook
718 trout in summer (August) and autumn (October) 2017.

719




Figure 3. Superoxide dismutase (SOD) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.






Figure 4. Catalase (CAT) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.




760 **Figure 5. Total glutathione peroxidase (GPx) activity in the liver, gills, kidney, spleen, and muscle**
 761 **tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S);**
 762 **Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences**
 763 **(Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.**

764



765

766 **Figure 6.** Selenium-dependent glutathione peroxidase (Se-GPx) activity in the liver, gills, kidney,
 767 spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard
 768 deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate
 769 statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling
 770 months for each tissue type.




785
786
787
788
789


790

791
792
793
794
795
796
797
798
799
800
801
802
803


Figure 7. Glutathione reductase (GR) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.




804 **Figure 8. Glutathione S-transferase (GST) activity in the liver, gills, kidney, spleen, and muscle**
 805 **tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S);**
 806 **Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences**
 807 **(Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.**




823 **Figure 9. Metallothionein (MTs) concentration in the muscle tissue of female and male brook trout.**
824 Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M).
825 Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the
826 sexes and the sampling months for each tissue type.



844 **Figure 10.** Biplot of score and loadings from principal component analysis. The scores of each
845 organ (gill, kidney, liver, muscle, and spleen) are denoted by a color and a symbol (largest symbol =
846 average value). Confidence ellipses plot convex hull values of each organ.





Highlights

- The influence of environmental factors on oxidative stress biomarkers was assessed
- Temperature, pH, Cu, Pb and Hg showed the major fluctuation during seasons
- A seasonal-linked variability of oxidative stress biomarkers level was recorded
- Both sexes showed similar level of oxidative stress biomarkers
- Food intake in summer mainly affected oxidative stress biomarkers

**Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake
(Cottian Alps)**

Paolo Pastorino^{1,2*}, Antonia Concetta Elia^{3**}, Barbara Caldaroni³, Vasco Menconi², Maria Cesarina Abete², Paola Brizio², Marco Bertoli¹, Annalisa Zaccaroni⁴, Magara Gabriele³, Ambrosius Josef Martin Dörr³, Elisabetta Pizzul¹, Marino Prearo¹

¹ Department of Life Sciences, University of Trieste, via Giorgieri 10, 34127 Trieste, Italy

² The Veterinary Medical Research Institute for Piemonte, Liguria and Valle d’Aosta, via Bologna 148, 10154 Torino, Italy

³ Department of Chemistry Biology and Biotechnology, University of Perugia, via Elce di Sotto 8, 06123 Perugia, Italy

⁴ Department of Veterinary Medical Science, University of Bologna, viale Vespucci 2, 47042 Cesenatico (FC), Italy

*Corresponding author: Paolo Pastorino, e-mail address: paolo.pastorino@izsto.it (P. Pastorino);

** Corresponding author: Antonia Concetta Elia, e-mail address: antonia.elia@unipg.it (A.C. Elia)

Abstract

High-mountain lakes are pristine ecosystems characterized by extreme environmental conditions. The atmospheric transport of pollutants from lowlands may add further stress to organisms inhabiting these environments. We investigated the environmental stress pressure on brook trout (*Salvelinus fontinalis*) from a high-mountain lake in the Cottian Alps (Piedmont, northwest Italy). To do this, males and females of brook trout were sampled from Balma Lake in summer (August) and autumn (October) 2017 in order to assess the influence of trace elements accumulation and

environmental parameters (physicochemical parameters and nutrient characteristics of water) on oxidative stress biomarkers. Bioaccumulation of Al, As, Cd, Cr, Cu, Fe, Hg, Pb, Ni, Se, and Zn and metallothionein levels were measured in muscle tissue of males and females. Liver, gills, kidney, and spleen tissue samples were analyzed for superoxide dismutase, catalase, total glutathione peroxidase, selenium-dependent glutathione peroxidase, glutathione reductase, and glutathione S-transferase activity. Analysis of environmental parameters showed changes in biomarker levels with seasonal variations. Water temperature was significantly higher in summer than autumn (Wilcoxon test; $p = 0.0078$), while pH was significantly higher in autumn than in summer (Wilcoxon test; $p = 0.0078$). Sex-related differences in oxidative stress biomarkers in tissues were unremarkable, whereas seasonal variability of oxidative stress biomarkers was observed, with major differences occurred for liver in summer and for gills, kidney, spleen and muscle in autumn. Positive correlations between environmental parameters and biomarkers were noted. Major fluctuations in water temperature, pH, Cu, Pb and Hg produced changes in biomarker levels; however, increased food intake during the ice-free season was probably the main factor that influenced changes in oxidative stress biomarker levels in brook trout in this extreme ecosystem.

Keywords: Alpine lakes; extreme ecosystems; oxidative stress biomarkers; trace elements

1. Introduction

Alpine lakes are remote, extreme ecosystems under harsh climatic conditions (Catalán et al., 2006). The ice-free season lasts for few months, generally from mid-June to late October. During this brief period of ideal conditions, some aquatic organisms can complete their life cycle before the snow covers the lakes again. Oligotrophic water conditions, UV radiation intensity, together with extreme temperatures allow for the development of a few dominant but well-adapted species (Sommaruga, 2001; Füreder et al., 2006; Pastorino et al., 2019a). These characteristics underlie the negative

53 correlation between altitude and biodiversity (Rott, 1988; Starkweather, 1990). Due to their remote
54 location, Alpine lakes are often considered pristine, unpolluted ecosystems (Pastorino et al., 2019b).
55 Since the 1980s, however, they have been affected by the global anthropogenic impact of pollutants
56 transported from lowland emission sources and the introduction of alien species (Tiberti et al.,
57 2014; Pastorino et al., 2020). These ecosystems have low resilience to disturbances and can be
58 particularly sensitive to the release of fish species for recreational angling, with important
59 consequences along the entire trophic chain (Tiberti et al., 2014; Milardi et al., 2016; Perrine, 2017;
60 Raposeiro et al., 2017). For example, the release of salmonids, especially brook trout (*Salvelinus*
61 *fontinalis*), has had a dramatic outcome for zooplanktonic, macrobenthic, and amphibian
62 communities in several Alpine lakes (Tiberti et al., 2014).

63 Alpine lakes are also a receptor for medium atmospheric transported (MRAT) contaminants
64 (Ferrario et al., 2017), as observed in the Arctic (Hung et al., 2016), which is subject to the long-
65 range transport potential (LRTP) of many chemicals. Altitudinal transport in the European Alps can
66 occur over relatively short distances from sources of pollution in the industrialized areas of
67 Germany, Switzerland, Austria, and northern Italy (Poma et al., 2017). The input of contaminant
68 compounds into Alpine lakes is controlled by atmospheric deposition and condensation (Schmid et
69 al., 2007). The contaminants are bioaccumulated by the organisms inhabiting these ecosystems.
70 Because fish occupy the uppermost trophic level, they provide an excellent bioindicator for the
71 atmospheric medium-long range input of persistent organic compounds such as pesticides,
72 brominated flame retardants (Schmid et al., 2007), and trace elements (Rognerud et al., 2002; Yang
73 et al., 2007).

74 Moreover, fish are used as sentinel organisms to detect environmental contamination (Squadrone et
75 al., 2013, 2014, 2016). They provide a useful model for assessing the impact of pollutants on
76 biological functions such as detoxification (Elia et al., 2010). Assessment of contaminants in
77 aquatic organisms can estimate and quantify the bioavailable fraction that might have the potential
78 to induce an effect. However, because determination of body concentrations alone does not provide

79 valuable data about the effects, quantification of some biological responses is necessary to evaluate
80 the health state of contaminant-exposed organisms (Elia et al., 2010). Metals, for instance, are pro-
81 oxidants that exert oxidative stress via reactive oxygen species (ROS) production and cause critical
82 changes in cellular biotransformation/detoxification pathways (Lushchak, 2016). Biomarker levels
83 can be also influenced by abiotic factors such as pH, dissolved oxygen content, and water
84 temperature (Sroda and Cossu-Leguille, 2011). Water temperature is a major factor in physiological
85 processes in fish and can induce the production of ROS (Lushchak, 2011).

86 Oxidative stress results from an imbalance between pro-oxidants such as ROS and the protective
87 antioxidant system. Mechanisms involve the activity of numerous antioxidant enzymes, including
88 superoxide dismutase (SOD, EC 1.15.1.1), catalase (CAT, EC 1.11.1.6), selenium-dependent
89 glutathione peroxidase (Se-GPx, EC 1.11.1.9), glutathione reductase (GR, EC 1.6.4.2), glutathione
90 S-transferases (GST, EC 2.5.1.18), as well as the rates of metal-trapping molecules such as reduced
91 glutathione and metallothioneins (MTs). They are important protective metabolic pathways that are
92 used as biomarkers of pollutant-induced oxidative stress. Selected oxidative stress biomarkers have
93 proven useful to assess the impact of a range of metals in aquatic organisms (Al Kaddissi et al.,
94 2014; Cozzari et al., 2015; Elia et al., 2006, 2007a, 2007b, 2010). Furthermore, contaminant levels
95 and antioxidant enzyme activity in aquatic organisms may change with the season and in response
96 to biological and environmental pressures (Monserrat et al., 2007). This poses a limitation on field
97 studies, because biochemical response may be linked to either fish physiology or exposure to
98 contaminants.

99 To our best knowledge, no studies are available about oxidative stress in fish from Balma Lake, an
100 Alpine lake in Piedmont (Cottian Alps, northwest Italy). Originally, the lake was fishless and brook
101 trout was released for recreational fishing. With the present study we investigated: a) the
102 physicochemical parameters and nutrient characteristics of lake water; b) the biometric
103 characteristics and stomach contents of brook trout (*S. fontinalis*); c) the trace element accumulation
104 in muscle because it constitutes a stable pool of trace elements for fish (Barwick and Maher, 2003);

105 d) the biomarkers of oxidative stress in muscle, liver, kidney, gills, and spleen in male and female
106 individuals captured in Balma Lake in summer and autumn 2017.

107

108 **2. Materials and Methods**

109 *2.1 Study site*

110 Balma Lake (45° 02' 14'' N; 07° 10' 52'' E) is located at 2100 m above sea level in the
111 municipality of Coazze, a small town 40 kilometers from Turin (Piedmont, northwest Italy). It is a
112 typical glacial-origin lake in the Cottian Alps, within the SCI/ZSC IT1110006 - Orsiera Rocciavère.
113 The lake is located above the tree line and is covered by ice from November to early June.
114 Originally, the lake was fishless, and *S. fontinalis* was released for recreational fishing during the
115 1970s (Pastorino et al., 2020). The main substrate of the area is composed of ophiolite metamorphic
116 bedrock. The main source of impact is the medium-long distance transport of pollutants from the
117 plain, grazing activities, and angling during the ice-free period. No previous studies or data about
118 the lake's hydrochemistry, taxa composition, and trace element accumulation are available. During
119 summer 2017 morphometric and bathymetric survey of the lake was carried out by GeoStudio RC
120 (Giaveno, Italy) using flying and floating drones. The lake measures 414 m in perimeter, 1.21 ha in
121 surface area, and 6.42 m maximum depth in the central zone (Fig. 1).

122

123 *2.2 Physicochemical parameters and nutrients of lake water*

124 During both sampling periods the main physicochemical parameters were monitored at 5 sites in the
125 littoral zone (in the upper centimeters of water) and 3 sites in the deep zone (in the water column)
126 (Fig. 1). Water temperature (°C), dissolved oxygen (% saturation; mg L⁻¹), conductivity (µS cm⁻¹),
127 and pH were recorded using field meters (HI 9033 conductivity meter, HI 9125 pH/ORP meter, HI
128 9147 oximeter, Hanna Instruments Inc. Woonsocket, RI, USA). Three replicates were carried out
129 for each parameter. Water samples were collected in sterile containers (three 1-L bottles for each
130 site), taking care not to include sediment particles, and then brought to the laboratory in a

131 refrigerated container within a few hours. Concentrations of NH_4^+ (mg L^{-1}), NO_3^- (mg L^{-1}), and
132 PO_4^{3-} (mg L^{-1}) were measured using a multi-parameter benchtop photometer (HI 83200-02, Hanna
133 Instruments Inc.). NO_3^- (mg L^{-1}) concentration was obtained by measuring absorbance at 525 nm
134 via an adaptation of the cadmium reduction method (APHA et al., 1998); NH_4^+ (mg L^{-1})
135 concentration was obtained by measuring the absorbance at 420 nm (ASTM, 2015) via adaptation
136 of the Nessler method; finally, PO_4^{3-} (mg L^{-1}) concentration was obtained by measuring absorbance
137 at 610 nm via adaptation of the ascorbic acid method (APHA et al., 1998).

138

139 *2.3 Fish sampling*

140 Fish sampling campaigns were carried out during summer (August) and autumn (October) 2017.
141 These months were chosen so that we could reach the sampling site on foot during the ice-free
142 period. Permission for sampling was granted by the competent authority (Città Metropolitana di
143 Torino; authorization n. 176-19040/2017). Fish were captured using 4 multimesh gillnets (36 x 1.8
144 m) divided into 6 panels of different mesh size (10 to 38 mm) to capture all size classes
145 indiscriminately, except for offspring. The gillnets were randomly placed in the lake (Fig. 1) for 3
146 hours and then recovered. For each sampling period, 20 individuals (Table 2) were sacrificed after
147 deep anesthesia with a lethal concentration (200 mg kg^{-1}) of tricaine methanesulfonate (MS-222)
148 dissolved in water. The fish were necropsied, sexed, weighed, and measured for total length in the
149 field. Immediately thereafter, samples of gill, liver, spleen, kidney, and muscle of each specimen
150 were collected, packed in dry ice, and transported to the laboratory.

151 Stomach contents analysis was performed to obtain information about fish diet and to characterize
152 seasonal variations. Stomachs were preserved in 70% alcohol, and the contents were inspected by
153 stereomicroscopy (Zeiss Stemis V8, Jena, Germany). The ingested prey was identified to the order
154 or family level, since closer identification was precluded by the digestion status of the organisms.
155 To describe the diet, prey frequency of occurrence (Fi) (Tiberti et al., 2016) was calculated for both
156 seasons.

157

158 2.4 Trace elements in fish muscle

159 Trace elements in fish muscle from males and females were analyzed by inductively coupled
160 plasma-mass spectrometry (ICP-MS Xseries II, Thermo Scientific, Bremen, Germany).
161 Determination of Al, As, Cd, Cr, Cu, Fe, Ni, Pb, Se, and Zn was performed following protocols
162 reported by Squadrone et al. (2016). Hg concentration was determined on a direct mercury analyzer
163 (DMA-80 Analyzer, Milestone, Shelton, CT, USA). Analytical performance was verified by
164 processing certified reference materials (Oyster Tissue - SRM 1566b from the National Institute of
165 Standard and Technology), along with blank reagents in each analytical session. Table S1 presents
166 the reference material values and the percentages of recovery. The analytical method was validated
167 according to ISO/IEC 17025 (general requirements for the competence of testing and calibration
168 laboratories).

169

170 2.5 Biochemical analyses

171 A total of 40 specimens of *S. fontinalis*, 26 females (12 in August and 14 in October) and 14 males
172 (8 in summer and 6 in autumn) were examined individually for oxidative stress biomarkers. Liver,
173 gills, kidney, spleen, and muscle were analyzed for SOD, CAT, GPx, Se-GPx, GR, and GST
174 activity, and MT levels only for muscle. For enzymatic analysis, the samples were homogenized
175 with an UltraTurrax homogenizer in 100 mM potassium-phosphate buffer, pH 7.5, added with 2.5%
176 sodium chloride (NaCl), 0.008 TIU ml⁻¹ aprotinin and 0.1 mg ml⁻¹ bacitracin. The homogenates
177 were centrifuged at 50,000 x g for 30 minutes at 4°C. Cytosolic fractions were used to determine
178 antioxidant biomarker activity. Biochemical analyzes were performed according to the methods
179 reported in Elia et al. (2017). Briefly, SOD activity was assessed at 550 nm in 50 mM Na₂CO₃
180 buffer, pH 10, 0.1 mM EDTA, 500 mM cytochrome C and 1 mM hypoxanthine and xanthine
181 oxidase. Cytochrome C reduction by the xanthine/hypoxanthine complex was evaluated by
182 comparison with a standard SOD unit curve. CAT activity was measured at 240 nm after the

183 decrease in absorbance following the consumption of H_2O_2 . The assay was carried out in NaH_2PO_4
184 buffer + Na_2HPO_4 100 mM pH 7 and H_2O_2 24 mM. Total glutathione peroxidase (GPx) and
185 selenium-dependent glutathione peroxidase (Se-GPx) activity was determined at 340 nm in
186 NaH_2PO_4 + Na_2HPO_4 100 mM buffer, pH 7.5, 1 mM EDTA, 0.12 mM NADPH (b-nicotinamide
187 adenine dinucleotide), 2 mM GSH, 1 U of GR (glutathione reductase), 1 mM NaN_3 and H_2O_2 0.6
188 mM for Se-GPx or 1 mM DTT and 0.8 mM cumene hydroperoxide for GPx. GR activity was
189 measured at 340 nm in NaH_2PO_4 + Na_2HPO_4 100 mM buffer, pH 7, 1 mM GSSG (oxidized
190 glutathione), and 0.06 mM NADPH. GST activity was measured at 340 nm using CDNB (1-chloro-
191 2,4-dinitrobenzene) as substrate. The assay was carried out in 100 mM NaH_2PO_4 + Na_2HPO_4 100
192 mM buffer, pH 6.5, 2 mM GSH and 2 mM CDNB. Concentration of cytosolic proteins was
193 determined according to the method of Lowry et al. (1951) and used to normalize biomarker
194 activity.

195 Metallothionein (MT) levels were measured in the muscle tissue of individuals of both sexes.
196 Samples were homogenized (1:4) in a buffer containing 0.02 M TRIS/HCl, 0.5 M sucrose, 0.1 mg
197 ml^{-1} bacitracin, 0.008 tiu ml^{-1} aprotinin, 87 $\mu\text{g ml}^{-1}$ phenylmethylsulfonyl fluoride (PMSF), and 0.1
198 $\mu\text{l ml}^{-1}$ α -mercaptoethanol. The homogenates were centrifuged at 14,500 x g at 4°C to obtain the
199 cytosolic fraction. The supernatants were purified using a chloroform/ethanol solution and then
200 HCl/ethanol to obtain the partially purified MT fraction. The pellets were washed with
201 ethanol/chloroform/TRIS/HCl (87/1/12) solution and suspended in 0.25 M NaCl. A destabilizing
202 solution (HCl 1N + EDTA 4 mM) and Ellman's reagent (DTNB: 5,5 dithiobis-2-nitrobenzoic acid)
203 were added to each sample. Sulphydril residue contents (-SH) were spectrophotometrically
204 quantified. Absorbance was measured at 412 nm and compared to that obtained from a standard
205 curve with reduced glutathione (1 mg ml^{-1} GSH). All biochemical analyses were performed in
206 triplicate for each sample on a Varian spectrophotometer (Cary 50 Thermostat Cell Holder) at a
207 constant temperature of 25°C.

208

209 2.6 Statistical analyses

210 Spearman's rank correlation coefficient (ρ_S) was used to test for correlations between trace element
211 concentration in fish muscle, water physicochemical parameters, nutrients, and oxidative stress
212 biomarkers. Differences in the concentration of physicochemical parameters, nutrients, and trace
213 elements between seasons were tested using the Wilcoxon test. Data from the biochemical analysis
214 are reported as the mean and standard deviation (SD). Significant differences between sexes and
215 months were tested using one-way ANOVA followed by Tukey's multiple comparison test.
216 Homogeneity of variance was checked using Bartlett's test. The criterion for significance was set at
217 $p < 0.05$. Principal component analysis (PCA) was performed to check for trends in trace elements,
218 biomarkers, and physicochemical values between the sampling seasons (summer and autumn).
219 Statistical analyses were performed using open source data analysis software RStudio® version
220 1.1.463 (RStudio, Inc.).

222 3. Results

223 3.1 Physicochemical and nutrient characteristics of lake water

224 Lake water temperature was significantly lower in autumn (range 7.20-8.80°C) than summer (range
225 14.60-16.10°C) (Wilcoxon test; $p = 0.0078$) in agreement with seasonal trends; no thermal
226 stratification in the water column was observed, since the temperatures recorded at the deep sites
227 were similar to those of the littoral sites in both seasons. pH values were significantly higher in
228 autumn (range 7.53-7.90) than summer (range 6.52-7.31) (Wilcoxon test; $p = 0.0078$). No
229 differences in oxygen saturation were recorded between summer (range 77-103%) and autumn
230 (range 101-107%) (Wilcoxon test; $p > 0.05$). Water conductivity remained constant between
231 summer and autumn (range 17-21 $\mu\text{S cm}^{-1}$) (Wilcoxon test; $p > 0.05$). PO_4^{3-} concentration was \leq
232 0.02 mg L^{-1} in both seasons. NH_4^+ level was $< 0.14 \text{ mg L}^{-1}$ at all sites, except for station 4 (0.20 mg
233 L^{-1}) in summer. NO_3^- level was $< 9.20 \text{ mg L}^{-1}$ at all sites, except for site 4 (12 mg L^{-1}) in summer.

234 There were no significant differences in PO_4^{3-} , NH_4^+ , and NO_3^- levels between seasons (Wilcoxon
235 test; $p > 0.05$). Table 1 presents the changes in physicochemical and nutrient data (mean \pm SD) .
236

237 3.2 Fish and stomach contents

238 Table 2 presents the average total length and weight of fish captured during summer and autumn
239 2017. Stomach contents analysis revealed the almost exclusive presence of terrestrial insect preys in
240 both summer (94.19%) and autumn (81.04%). Preys belonged to the order Hymenoptera (77.74%
241 and 66.39% in summer and autumn, respectively) and Coleoptera (16.45% and 14.65% in summer
242 and autumn, respectively). Other taxa were present in extremely low proportions (Diptera
243 Chironomidae larvae: 5.18% in summer and 2.79% in autumn).
244

245 3.3 Trace elements in fish muscle

246 Figure 2 presents the mean concentration and the relative standard deviation of each trace element
247 detected in muscle tissue in summer and autumn. The mean concentration of trace elements was in
248 the order: Zn (16.25) > Fe (8.78) > Al (1.49) > Se (0.67) > Cu (0.60) > Cr (0.14) > As (0.06) > Ni
249 (0.06) > Pb (0.05) > Hg (0.01) mg Kg^{-1} . In autumn the mean concentration of trace elements was in
250 the order: Zn (16.13) > Fe (10.63) > Al (2.03) > Se (0.49) > Cu (0.36) > Cr (0.16) > As (0.10) > Pb
251 (0.10) > Ni (0.06) > Hg (0.02) mg Kg^{-1} . Cd was < LOQ (0.02 mg Kg^{-1}) in both seasons. There were
252 no significant differences in trace element concentration between seasons (Wilcoxon test; $p > 0.05$
253 for all elements).
254

255 3.4 Biochemical analyses

256 In the liver, the activity of SOD, CAT, and GR was significantly lower in autumn (up to 70%) than
257 summer in males and females (Tukey's test; $p < 0.05$) (Figs. 3-4, 7). In the gills, GPx activity was
258 significantly higher in autumn (90%) mainly in the females (Tukey's test; $p < 0.05$); Se-GPx and
259 GST activity was significantly higher (up to 2-fold) in males and females in autumn (Tukey's test; p

260 < 0.05) (Figs. 5, 6, 8). In the kidney, only Se-GPx activity was significantly different between the
261 seasons, being higher (up to 40%) in autumn (Tukey's test; $p < 0.05$) (Fig. 6). In the spleen, Se-GPx
262 activity was significantly lower (70%) in the females (Tukey's test; $p < 0.05$) (Fig. 6). GR and GST
263 activity was significantly higher (up to 70%) in males in autumn than in summer (Tukey's test; $p <$
264 0.05) (Fig. 7, 8). In muscle, SOD activity was significantly lower (up to 50%) in females in autumn
265 (Tukey's test; $p < 0.05$), whereas CAT activity showed an opposite trend and was significantly
266 higher in autumn (up to one-fold) (Tukey's test; $p < 0.05$) (Figs. 3-4). MT level was higher in
267 autumn (up to one-fold) than in summer (Tukey's test; $p < 0.05$) (Fig. 9).

268

269 3.5 Spearman correlation matrix

270 Spearman correlation matrix revealed correlations between environmental parameters (trace
271 elements, physicochemical parameters, nutrients) and oxidative stress biomarkers in muscle, gills,
272 liver, spleen, and kidney tissue for both seasons. Due to the multiplicity of positive correlations,
273 Table S2 presents the correlation matrices (one for each organ). Only the most informative
274 outcomes are presented and discussed for interpretation of biological response of *S. fontinalis* to
275 environmental parameters. In detail, a significant positive correlation was found between MTs and
276 Hg (ρ_S 0.787), MTs and Pb (ρ_S 0.787), MTs and Cu (ρ_S 0.683), MTs and pH (ρ_S 0.650), and CAT
277 and pH (ρ_S 0.737) in muscle tissues of females captured in autumn; SOD and Cr (ρ_S 0.837) in the
278 liver tissue of females captured in autumn; SOD and NO_3^- (ρ_S 0.750), Se-GPx and NO_3^- (ρ_S 0.750)
279 in muscle and kidney tissue, respectively, of females captured in summer.

280

281 3.6 Principal Component Analysis (PCA)

282 The first two principal components (Dim1; Dim2) accounted for meaningful amounts of the total
283 variance (58.2%), while the other components accounted for a relatively smaller fraction (Fig. 10).
284 Dim1 accounted for 35.2% of the total variance and was positively correlated with the variables Ni,
285 Se, temperature, oxygen, conductivity, and NO_3^- and negatively correlated with Al, Cr, Fe, As, Pb,

Hg, and pH. Dim2 accounted for 23% of the total variance and was positively correlated with the variables GST, CAT, GPx, Se-GPx, SOD, and GR. The biplot of loadings (variables) and score (observations) shows which organs (liver, gills, kidney, spleen, and muscle) of brook trout are closest to them, and which variables (trace elements, biomarkers and physicochemical parameters) contributed to this grouping in the coordinate of Dim1 and Dim2. Separation of organs by season (summer on the right, autumn on the left) suggested a difference in biomarker values. In detail, the autumn samples of spleen, gill, muscle, kidney, and liver tissue are on the left in order of increasing value of Al, Cr, Fe, As, Pb, Hg, and pH. The summer samples of spleen, gill, muscle, kidney tissue are on the right in order of increasing value of NH_4^+ , NO_3^- , conductivity, oxygen, temperature, Zn, Cu, Ni, and Se. Remarkably, summer samples of liver tissue are well separated from other organs, following the trend in biomarker values.

297

298 **4. Discussion**

The environmental parameters of water oxygenation, pH, conductivity, and temperature are in line with those reported by Tiberti et al. (2010) for 12 alpine lakes in the Gran Paradiso National Park (Western Alps, Italy). Temperatures were lower in autumn than in summer according to seasonality. The temperature data recorded at the deep sites revealed no vertical layering in the lake because the shallow maximum depth (6.42 m) does not allow for the formation of a thermocline layer. The pH values are related to rock composition. Since Lake Bama lies over a granite bedrock, its pH values are lower than lakes on limestone or sandstone bedrock (Camarero et al., 2009). The pH values were in line with the literature reported for high-altitude environments (Boggero et al., 2006; Füreder et al., 2006; Fjellheim et al., 2009). As expected for mountain lakes, the oxygenation levels were high. The water conductivity values were in line with published literature, as conductivity of silty-like lakes tends to be $< 50 \mu\text{S cm}^{-1}$ (Boggero et al., 2006; Füreder et al., 2006). No decrease in oxygen values correlated with depth was observed owing to the absence of temperature

311 stratification. Nutrient levels (PO_4^{3-} , NO_3^- and NH_4^+) were in line with previous studies of Alpine
312 lakes (Füreder et al., 2006; Camarero et al., 2009) and revealed an oligotrophic condition.

313 We observed a positive correlation between environmental parameters and biological response in
314 the *S. fontinalis* from Balma Lake. The antioxidant response was tissue-specific; indeed, aerobic
315 tissues such as the kidney, gills, spleen, and the liver in particular have a high potential for ROS
316 production which is offset by protective mechanisms. Differently, muscle, which has a low content
317 of mitochondria and a low-intensive oxidative metabolism, showed mild response to
318 oxidative/reductive conditions. Moreover, the level of several biomarkers was related to the
319 seasonal changes in the trace element concentration in some tissues.

320 The few studies on trace element accumulation in the biota from high-altitude lakes in general, and
321 in Alpine lakes in particular, have focused largely on fish. Köck et al. (1996) studied the
322 concentrations of Cd, Pb, Zn, and Cu in the muscle of arctic char (*Salvelinus alpinus*) from five
323 oligotrophic Alpine lakes in northern Tyrol (Austria). Yang et al. (2007) studied the accumulation
324 of trace elements in muscle of fish of the genus *Gymnocypris* (Cyprinidae) from high-mountain
325 lakes in the Tibetan Plateau. Ni ranged from 0.094 to 0.12 mg kg⁻¹, Cu from 1.1 to 2.0 mg kg⁻¹, Zn
326 from 4.4. to 6.9 mg kg⁻¹, As from 0.24 to 0.27 mg kg⁻¹, Se from 0.36 to 1.0 mg kg⁻¹, Cd from 0.024
327 to 0.025 mg kg⁻¹, and Pb from 0.047 to 0.079 mg kg⁻¹. Rognerud et al. (2002) found that Hg
328 concentrations in fish muscle from high-mountain lakes in Europe ranged from 0.021 to 0.179 mg
329 kg⁻¹. These data from previous studies demonstrate that high-mountain lakes function as a regional
330 contaminant convergence zone for the medium and long-range atmospheric transport of
331 contaminants. Since our results are in line with these findings, and because no previous studies have
332 been performed to date, we assume that the trace element concentrations we detected in the brook
333 trout from Balma Lake have both anthropogenic and pedogenic sources as their origin.

334 Metallothioneins have been widely considered as valuable biomarkers that reflect the level of trace
335 elements in aquatic environments, where they act as metal trappers mainly of Cd, Hg, Pb, Cu, and
336 Zn in fish (Bourdineaud et al., 2006; Morcillo et al., 2016). The higher levels of MTs we measured

337 in autumn were related to the increased trace element concentration in this tissue, and we noted a
338 strong correlation with Cu, Hg, and Pb levels in the females. This may suggest an alarming
339 scenario, as it would signal an increased concentration of these elements in the environment.
340 Although no chemical analyses of the lake water were carried out, because of the peculiar
341 geomorphological characteristics of Balma Lake we can exclude an increase in such contaminants
342 during autumn.

343 Seasonal variation in metal concentrations in fish can be related to environmental factors such as
344 food availability, temperature, and biological cycle (Hermesz et al., 2001; Amiard et al., 2006;
345 Dragun et al., 2009). Furthermore, fish size may also play a key role in metal uptake (Wright and
346 Mason, 1999). In the present study, an increase in weight and length was measured in the fish from
347 both sampling seasons, and the frequency of occurrence of terrestrial invertebrates in the diet during
348 the ice-free season was in accordance with other studies performed on brook trout in other Alpine
349 lakes (Sotiropoulos et al., 2006; Tiberti et al., 2016). It is possible that fish growth due to a higher
350 intake of food in summer could have favored the uptake of heavy metals such as Pb and Hg.

351 This hypothesis is corroborated by a previous study on black scabbardfish *Aphanopus carbo*
352 (Trichiuridae), in which an exponential increase in total Hg load was found in all fish tissues in a
353 length-dependent manner (Bebianno et al., 2007). Furthermore, previous studies showed that
354 fluctuations toward high pH values can also play a key role in modulating metals uptake, affecting
355 their speciation and bioavailability (Playle, 1998). This outcome may explain the strong correlation
356 between pH and the two hydroxyl scavengers MTs and CAT in the female muscle tissue in autumn.

357 Trace elements also affected the activity of several enzymes in different tissues in the autumn fish
358 samples. We noted a strong correlation between Cr and SOD activity in the liver, as reported in
359 previous study in rock fish *Sebastes schlegelii* (Kim and Kang, 2016). Furthermore, a recent study
360 showed that Cr can induce conformational changes of CAT enzyme and reduce its activity
361 depending on its valence states and concentration (Chen et al., 2018). These findings may explain
362 the seasonal difference in CAT activity in the female muscle tissue.

363 In general, numerous elements can influence the activity of this biomarker, and As and Fe were the
364 elements most involved in modulating CAT enzyme activity in the tissues. Arsenic is a global
365 contaminant derived from natural or anthropogenic sources and a cause of great concern for
366 terrestrial and aquatic ecosystems (Elia et al., 2018). At high concentrations, arsenic may induce
367 oxidative stress by interacting with antioxidants and result in the accumulation of free radicals in
368 cells. Arsenite species can interact with sulfhydryl groups of biomolecules such as enzymes or
369 reduced glutathione (Elia et al., 2018). Furthermore, redox active metals such as iron generate ROS
370 or are involved in the Fenton route, leading to lipid peroxidation (Mahboob, 2013). Despite the
371 increase in trace element amount in autumn, the concentration of Pb, Hg, and Cd in fish muscle
372 mentioned in Regulation 1881/2006 (European Commission, 2006) was far below the established
373 threshold limit. This fact should be taken into account and may suggest the adaption of *S. fontinalis*
374 to seasonal changes rather than to oxidative stress. PCA analysis showed that summer samples of
375 liver tissue were strongly related to oxidative biomarker level, since liver tissue is the site of
376 multiple oxidative reactions and maximal free radical generation (Gul et al., 2004; Avci et al.,
377 2005).

378 For ectothermic organisms, temperature is a crucial abiotic factor (Hassan et al., 2017). Daily
379 temperature fluctuations and seasonal variations differ in their influence on fish (Buckley et al.,
380 2006; Place and Hofmann, 2004; Podrabsky and Somero, 2004). Wellness and growth are optimal
381 within a well-defined temperature range depending on the species (Godowsky and Caddell, 1991).
382 Temperatures deviating excessively from the optimum can exert harmful effects and induce
383 mortality (Rijnsdorp et al., 2009). In the present study, the marked drop in water temperature in
384 autumn was related to changes mainly in GPx activity in the male kidney and in the female spleen.
385 Moreover, higher temperature results in increased GPx activity in other fish such as the Antarctic
386 *Pagothenia borchgrevinki* (Almroth et al., 2015). Thermal changes have also been associated with
387 the up regulation of the NRF2 transcription factor, which is involved in the expression of
388 antioxidants via binding to the antioxidant responsive element (ARE) (Almroth et al., 2015).

Elevated GPx activity during summer may indicate strengthening of this fundamental defense line against ROS. However, the lack of change in SOD and CAT activity, as well as the lower concentration of trace elements in tissue, and the constant levels of the main physicochemical parameters of water, except for pH and temperature, preclude an oxidative pressure scenario and suggest an adaptive ability of *S. fontinalis* to higher temperature instead. On the other hand, increased SOD activity in the muscle tissue of males and females is linked to an abiotic factor, such as NO_3^- , in summer. Furthermore, nutrients also influenced SOD concentration in female muscle and Se-GPx activity in female kidney in summer. In the aquatic environment, it is not unusual that fish are simultaneously challenged by different abiotic factors. Conversely, at very high concentrations, ammonia can induce a range of toxicological effects in fish, such as altered metabolism, lack of growth, and mortality (Dosdat et al., 2003; Sinha et al., 2012, 2015). Ammonia exposure can also lead to oxidative stress in fish (Sun et al., 2012; Sinha et al., 2014). A previous study showed that conductivity plays a crucial role in maintaining the ammonia ionization equilibrium (NH_3 and the non-toxic form NH_4^+) in aquatic environments (Sinha et al., 2015). In the present study, the lack of change in conductivity and nutrients (e.g., phosphorus) between the seasons suggests that the changes in antioxidant parameters may be related to food intake (which indirectly promotes metals uptake) rather than to nutrient concentration. In their habitats, high-mountain lakes included, fish are often exposed to periods of food insufficiency in response to factors such as temperature, conductivity, and biological cycle (Pérez-Jiménez et al., 2007; Furné et al., 2009). The increase in food intake during summer may also explain the fluctuation in biomarkers of oxidative stress.

410

411 5. Conclusion

412 With this study we investigated the influence of trace element accumulation and environmental
413 parameters on oxidative stress biomarkers in male and female individuals of *S. fontinalis* from a
414 high-mountain lake during the ice-free period. Generally, positive correlations were found between

several environmental parameters and biomarkers. While oxidative stress biomarker levels were similar for males and females, significant fluctuation between seasons due to biological and environmental pressures was noted for several biomarkers. Although there was greater fluctuation in temperature, pH, and trace elements (e.g., Cu, Pb and Hg) between seasons, which certainly contributed to changes in biomarker levels, our findings indicate that increased food intake during the ice-free season was probably the main factor that affected the oxidative stress response. Future studies are needed to investigate other factors responsible for the changes in oxidative stress biomarkers.

Acknowledgements

The authors would like to thank GeoStudio RC (Marco Rosa Clot, Luca Rosa Clot), Club Alpino Italiano (Alfio Usseglio) and Ente di Gestione delle Aree Protette delle Alpi Cozie (Michele Ottino) for their technical support.

References

- Al Kaddissi, S., Legeay, A., Elia, A.C., Gonzalez, P., Floriani, M., Cavalieri, I., Massabuaum, J.C., Gilbin, R., Simon, O., 2014. Mitochondrial gene expression, antioxidant responses, and histopathology after cadmium exposure. *Environ. Toxicol.* 29, 893-907.
- Almroth, B.C., Asker, N., Wassmur, B., Rosengren, M., Jutfelt, F., Gräns, A., Sundell, K., Axelsson, M., Sturve, J., 2015. Warmer water temperature results in oxidative damage in an Antarctic fish, the bald notothen. *J. Exp. Mar. Biol. Ecol.* 468, 130-137.
- Amiard, J.C., Amiard-Triquet, C., Barka, S., Pellerin, J., Rainbow, P.S., 2006. Metallothioneins in aquatic invertebrates: their role in metal detoxification and their use as biomarkers. *Aquat. Toxicol.* 76, 160-202.
- APHA, AWWA, WEF, 1998. Standard Methods for the Examination of Water and Wastewater, 20th edition. American Public Health Association, Washington DC.

441 ASTM, 2015. Manual of Water and Environmental Technology, D1426-92, Nessler method.

442 Avci, A., Kačmaz, M., Durak, I., 2005. Peroxidation in muscle and liver tissues from fish in a
 443 contaminated river due to a petroleum refinery industry. *Ecotoxicol. Environ. Safe.* 60, 101-105.

444 Barwick, M., Maher, W., 2003. Biotransference and biomagnification of selenium copper,
 445 cadmium, zinc, arsenic and lead in a temperate seagrass ecosystem from Lake Macquarie Estuary,
 446 NSW, Australia. *Mar. Environ. Res.* 56, 471-502.

447 Bebianno, M.J., Santosa, C., Canário, J., Gouveia, N., Sena-Carvalho, D., Vale, C., 2007. Hg and
 448 metallothionein-like proteins in the black scabbardfish *Aphanopus carbo*. *Food. Chem. Toxicol.* 45,
 449 1443-1452.

450 Boggero, A., Füreder, L., Lencioni, V., Simcic, T., Thaler, B., Ferrarese, U., Lotter, A.F., Ettinger,
 451 R., 2006. Littoral chironomid communities of Alpine lakes in relation to environmental factors.
 452 *Hydrobiologia* 562, 145-165.

453 Bourdineaud J.P., Baudrimont M., González P., Moreau J.L., 2006. Challenging the model for
 454 induction of metallothionein gene expression. *Biochimie* 88, 1787-1792.

455 Buckley, B.A., Gracey, A.Y., Somero, G.N., 2006. The cellular response to heat stress in the goby
 456 *Gillichthys mirabilis*: a cDNA microarray and protein level analysis. *J. Exp. Biol.* 209, 2660-2677.

457 Camarero, L., Rogora, M., Mosello, R., Anderson, N.J., Barbieri, A., Botev, I., Kernan, M.,
 458 Kopáček, J., Korhola, A., Lotter, A.F., Muri, G., Postolache, C., Stuchik, E., Thies, H., Wright,
 459 R.F., 2009. Regionalization of chemical variability in European mountain lakes. *Freshw. Biol.* 54,
 460 2452-2469.

461 Catalán, J., Camarero, L., Felip, M., Pla S., Ventura, M., Buchaca, T., Bartumeus, F., De Mendoza,
 462 G., Miró, A., Casamayor, E.O., Medina-Sánchez, J.M., Bacardit, M., Altuna, M., Bartrons, M.,
 463 Díaz, de Quijano, D., 2006. High mountain lakes: extreme habitats and witnesses of environmental
 464 changes. *Limnetica* 25, 551-584.

465 Chen, L., Zhang, J., Zhu, Y., Zhang, Y., 2018. Interaction of chromium (III) or chromium (VI) with
 466 catalase and its effect on the structure and function of catalase: An in vitro study. Food. Chem. 244,
 467 378-385.

468 Cozzari, M., Elia, A.C., Pacini, N., Smith, B.D., Boyle, D., Rainbow, P.S., Khan, F.R., 2015.
 469 Bioaccumulation and oxidative stress responses measured in the estuarine ragworm (*Nereis*
 470 *diversicolor*) exposed to dissolved, nano- and bulk-sized silver. Environ. Pollut. 198, 32-40.

471 Dosdat, A., Ruyet, J.P., Coves, D., Dutto, G., Gasset, E., Le Roux, A., Lemarié, G., 2003. Effect of
 472 chronic exposure to ammonia on growth, food utilization and metabolism of the European sea bass
 473 (*Dicentrarchus labrax*). Aquat. Living Resour. 16, 509-520.

474 Dragun, Z., Podrug, M., Raspor, B., 2009. The assessment of natural causes of metallothionein
 475 variability in the gills of European chub (*Squalius cephalus* L.). Comp. Biochem. Physiol. C:
 476 Toxicol. Pharmacol. 150, 209e217.

477 Elia A.C., Dörr A.J.M, Abete M.C., Prearo M., 2010. Seasonal variability of detoxificant response
 478 and heavy metal accumulation in tissues of both sexes in *Tinca tinca* (L.) from Lake Trasimeno.
 479 Rev. Fish. Biol. Fisher. 20(3), 425-434.

480 Elia A.C., Magara G., Caruso C., Masoero L., Prearo M., Arsieni P., Caldaroni B., Scoparo M.,
 481 Dörr A.J.M., Salvati S., Brizio P., Squadrone S., Abete M.C., 2018. A comparative study on
 482 subacute toxicity of arsenic trioxide and dimethylarsinic acid on antioxidants and antioxidant-
 483 related enzymes in crandell rees feline kidney (CRFK), human hepatocellular carcinoma
 484 (PCL/PRF/5) and epithelioma papulosum cyprini (EPC) cell lines. J. Toxicol. Environ. Health part
 485 A 81(10), 333-348.

486 Elia, A.C., Dörr, A.J.M., Galarini, R., 2007a. Comparison of organochlorine pesticides, PCBs and
 487 heavy metal contamination and of detoxifying response in tissues of *Amerius melas* from Corbara,
 488 Alviano and Trasimeno Lakes, Italy. Bull. Environ. Contam. Toxicol. 78, 463-468.

489 Elia, A.C., Galarini, R., Dörr, A.J.M., Taticchi MI., 2007b. Heavy metal contamination and
 490 antioxidant response of a freshwater bryozoan (*Lophopus crystallinus* Pall., Phylactolaemata).
 491 Ecotox. Environ. Safe. 66, 188-194.

492 Elia, A.C., Galarini, R., Dörr, A.J.M., Taticchi MI. 2006. Bioaccumulation of heavy metals,
 493 organochlorine pesticides, and detoxification biochemical indexes in tissues of *Ictalurus melas* of
 494 Lake Trasimeno. Bull. Environ. Contam. Toxicol. 76, 132-139.

495 Elia, A.C., Magara, G., Righetti, M., Dörr, A.J.M., Scanzio, T., Pacini, N., Abete, M.C., Prearo, M.,
 496 2017. Oxidative stress and related biomarkers in cupric and cuprous chloride-treated rainbow trout.
 497 Environ. Sci. Pollut. R. 24, 10205-10219.

498 European Commission, 2006. Commission Regulation (EC) No 1881/2006 of 19 December 2006
 499 setting maximum levels for certain contaminants in foodstuffs. OJEC 364, 365-324.

500 Ferrario, C., Finizio, A., Villa S., 2017. Legacy and emerging contaminants in meltwater of three
 501 alpine glaciers. Sci. Total. Environ. 574, 350-357.

502 Fjellheim, A., Raddum, G.G., Vandvik, V., Cogălniceanu, D., Boggero, A., Brancelj, A., Galas, J.,
 503 Sporka, F., Vidinova, Y., Bitusik, P., Dumnicka, E., Gâldean, N., Kownacki, A., Krno, I., Preda, E.,
 504 Rîșnoveanu, G., Stuchlik, E., 2009. Diversity and distribution patterns of benthic invertebrates
 505 along alpine gradients. A study of remote European freshwater lakes. Adv. Limnol. 62, 167-190.

506 Füreder, L., Ettinger, R., Boggero, A., Thaler, B., Thies, H., 2006. Macroinvertebrate diversity in
 507 Alpine lakes: effects of altitude and catchment properties. Hydrobiologia 652, 123-144.

508 Furné, M., García-Gallego, M., Hidalgo, M.C., Morales, A.E., Domezain, A., Domezain, J., Sanz,
 509 A., 2009. Oxidative stress parameters during starvation and refeeding periods in Adriatic sturgeon
 510 (*Acipenser naccarii*) and rainbow trout (*Oncorhynchus mykiss*). Aquaculture Nutr. 15, 587-595.

511 Godowsky, D.M., Caddell, S.M., 1991. Effects of temperature on early-life-history stages of
 512 California halibut *Paralichthys californicus*. Fish Bulletin 89, 567-576.

513 Gul, S., Belge-Kurutas, E., Yildiz, E., Sahan, A. Doran, F., 2004. Pollution correlated modifications
514 of liver antioxidant systems and histopathology of fish (Cyprinidae) living in Seyhan Dam Lake,
515 Turkey. Environ. Int. 30, 605-609.

516 Hassan, M.K., Nahid, N.M., Bahar, A.N., Bhuiyan, M.M.R., Abdullah-Al-Shafi, M., Ahmed, K.,
517 2017. Dataset demonstrating the temperature effect on average output polarization for QCA based
518 reversible logic gates. Data in Brief, Elsevier 13, 713–716.

519 Hermes, E., Abrahm, M., Nemcsok, J., 2001. Tissue-specific expression of two metallothionein
520 genes in common carp during cadmium exposure and temperature shock. Biochem. Physiol. Part C
521 128, 457–465.

522 Hung, H., Katsoyianni, A.A., Brorström-Lundén, E., Olafsdottir, K., Aas, W., Breivik, K., Bohlin-
523 Nizzetto, P., Sigurdsson, A., Hakola, H., Bossi, R., Skov, H., Sverko, E., Barresi, E., Fellin, P.,
524 Wilson, S., 2016. Temporal trends of Persistent Organic Pollutants (POPs) in arctic air: 20 years of
525 monitoring under the Arctic Monitoring and Assessment Programme (AMAP). Environ. Pollut.
526 217, 52-61.

527 Kim, J.H., Kang, J.C., 2016. Oxidative stress, neurotoxicity, and metallothionein (MT) gene
528 expression in juvenile rock fish *Sebastes schlegelii* under the different levels of dietary chromium
529 (Cr^{6+}) exposure. Ecotox. Environ. Safe. 125, 78-84.

530 Köck, G., Triendl, M., Hofer, R., 1996. Seasonal patterns of metal accumulation in Arctic char
531 (*Salvelinus alpinus*) from an oligotrophic Alpine lake related to temperature. Can. J. Fish. Aquat.
532 Sci. 53, 780-786.

533 Lowry, O.H., Rosebrough, N.J., Farr A.L., Randall R.J., 1951. Protein measurement with the Folin
534 phenol reagent. J. Biol. Chem. 193, 265–75.

535 Lushchak, V.I., 2011. Environmentally induced oxidative stress in aquatic animals. Aquat. toxicol.
536 101(1), 13-30.

537 Lushchak, V.I., 2016. Contaminant-induced oxidative stress in fish: a mechanistic approach. Fish
538 Physiol. Biochem. 42, 711-747.

539 Mahboob, S., 2013. Environmental pollution of heavy metals as a cause of oxidative stress in fish: a
 540 review. *Life Sci. J.* 10(10s), 336-345.

541 Milardi, M.S. Siitonen, J., Lappalainen, A., Liljendahl, Weckström, J., 2016. The impact of trout
 542 introductions on macro- and micro-invertebrate communities of fishless boreal lakes. *J.*
 543 *Paleolimnol.* 55, 273-287.

544 Monserrat, J., Martinez, P., Geracitano, L., Amado, L., Martins, C., Pinho, G., Chaves, I., Cravo,
 545 M., Lima, J., Bianchini, A., 2007. Pollution biomarkers in estuarine animals: critical review and
 546 new perspectives. *Comp. Biochem. Physiol.* 146C, 221-234.

547 Morcillo, P., Esteban, M.A., Cuesta, A., 2016. Heavy metals produce toxicity, oxidative stress and
 548 apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere* 144, 225-233.

549 Pastorino P., Pizzul E., Bertoli M., Perilli S., Brizio P., Salvi G., Esposito G., Abete M.C., Prearo
 550 M., Squadrone S. (2019b). Macrobenthic invertebrates as bioindicators of trace elements in high-
 551 mountain lakes. *Environ. Sci. Pollut. Res.* <https://doi.org/10.1007/s11356-019-07325-x>

552 Pastorino, P., Polazzo, F., Bertoli, M., Santi, M., Righetti, M., Pizzu, E., Prearo, M., 2020.
 553 Consequences of fish introduction in fishless Alpine lakes: preliminary notes from a sanitary point
 554 of view. *Turk. J. Fish Aquat. Sc.* 20(1), 1-8.

555 Pastorino, P., Prearo, M., Pizzul, E., Bertoli, M., Francese, D.R., Menconi, V., Mugetti D., Bozzetta
 556 E., Varello, K. (2019a). Hepatic Steatosis in a Bullhead (*Cottus gobio*) Population from a High-
 557 Mountain Lake (Carnic Alps): Adaptation to an Extreme Ecosystem? *Water* 11(12), 2570.

558 Pérez-Jiménez, A., Guedes, M.J., Morales, A.E., Oliva-Teles, A., 2007. Metabolic responses to
 559 short starvation and refeeding in *Dicentrarchus labrax*. Effect of dietary composition. *Aquaculture*
 560 265, 325-335.

561 Perrine, S., 2017. Inferring fish abundance in lakes from subfossil midge (Diptera: Chironomidae)
 562 remains. University of Jyväskylä, Department of Biological and Environmental Science, Master
 563 Thesis, Jyväskylä, Finland.

564 Place, S.P., Hofmann, G.E., 2004. Temperature interactions of the molecular chaperone Hsc70 from
 565 the eurythermal marine goby *Gillichthys mirabilis*. J. Exp. Biol. 204, 2675–2682.

566 Playle, R.C., 1998. Modelling metal interactions at fish gills. Sci. Total. Environ. 219, 147-163.

567 Podrabsky, J.E., Somero, G.N., 2004. Changes in gene expression associated with acclimation to
 568 constant temperatures and fluctuating daily temperatures in an annual killifish, *Austrofundulus*
 569 *limnaeus*. J. Exp. Biol. 207, 2237-2254.

570 Poma, G., Salerno, F., Roscioli, C., Novati, S., Guzzella, L., 2017. Persistent organic pollutants in
 571 sediments of high-altitude Alpine ponds within Stelvio National Park, Italian Alps. Inland Waters 7,
 572 34-44.

573 Raposeiro, P.M., Rubio M.J, González A., Hernández A., Sánchez-López G., Vázquez-Loureiro D.,
 574 Rull V., Bao R., Costa A. C., Gonçalves V., Saéz A., Giralte S., 2017. Impact of the historical
 575 introduction of exotic fishes on the chironomid community of Lake Azul (Azores Islands).
 576 Palaeogeogr Palaeocl. 466, 77–88.

577 Rijnsdorp, A.D., Peck, M.A., Engelhard, G.H., Mollmann, C., Pinnegar, J.K., 2009. Resolving the
 578 effect of climate change on fish populations. ICES J. Mar. Sci. 66, 1570-1583.

579 Rognerud, S., Grimalt, J.O., Rosseland, B.O., Fernandez, P., Hofer, R., Lackner, R., et al. 2002.
 580 Mercury and organochlorine contamination in Brown trout (*Salmo trutta*) and Arctic charr
 581 (*Salvelinus alpinus*) from high mountain lakes in Europe and the Svalbard archipelago. Water Air
 582 Soil Pollut. 2, 209-32.

583 Rott, E., 1988. Some aspects of the seasonal distribution of flagellates in mountain lakes.
 584 Hydrobiologia 161, 159-170.

585 Sinha, A.K., Abdelgawad, H., Giblen, T., Zinta, G., De Rop, M., Asard, H., Blust, R., De Boeck G.,
 586 2014. Anti-oxidative defences are modulated differentially in three freshwater teleosts in response
 587 to ammonia-induced oxidative stress. PLoS One, 9(4), e95319.

588 Sinha, A.K., Liew, H.J., Diricx, M., Blust, R., De Boeck, G., 2012. The interactive effects of
 589 ammonia exposure, nutritional status and exercise on metabolic and physiological responses in
 590 goldfish (*Carassius auratus* L.). *Aquat. Toxicol.* 109, 33–46.

591 Sinha, A.K., Rasoloniriana, R., Dasan, A.F., Pipralia, N., Blust, R., De Boeck, G., 2015. Interactive
 592 effect of high environmental ammonia and nutritional status on ecophysiological performance of
 593 European sea bass (*Dicentrarchus labrax*) acclimated to reduced seawater salinities. *Aquat.*
 594 *Toxicol.* 160, 39–56.

595 Schmid, P., Kohler, M., Gujer, E., Zennegg, M., Lanfranchi, M., 2007. Persistent organic
 596 pollutants, brominated flame retardants and synthetic musks in fish from remote alpine lakes in
 597 Switzerland. *Chemosphere* 67(9), S16-S21.

598 Sommaruga, R., 2001. The role of solar UV radiation in the ecology of alpine lakes. *J. Photochem.*
 599 *Photobiol.* 62, 35-42.

600 Sotiropoulos, J.C., Nislow, K.H., Ross M.R., 2006. Brook trout, *Salvelinus fontinalis*, microhabitat
 601 selection and diet under low summer stream flows. *Fisheries Manag. Ecol.* 13, 149-155.

602 Squadrone, S., Brizio, P., Stella, C., Prearo, M., Pastorino, P., Serracca, L., Ercolini, C., Abete,
 603 M.C., 2016. Presence of trace metals in aquaculture marine ecosystems of the northwestern
 604 Mediterranean Sea (Italy). *Environ. Poll.* 215, 77-83.

605 Squadrone, S., Ciccotelli, V., Favaro, L., Scanzio, T., Prearo, M., Abete, M.C., 2014. Fish
 606 consumption as a source of human exposure to perfluorinated alkyl substances in Italy: Analysis of
 607 two edible fish from Lake Maggiore. *Chemosphere* 114, 181-186.

608 Squadrone, S., Prearo, M., Brizio, P., Gavinelli, S., Pellegrino, M., Scanzio, T., Guarise, S.,
 609 Benedetto, A., Abete, M.C., 2013. Heavy metals distribution in muscle, liver, kidney and gill of
 610 European catfish (*Silurus glanis*) from Italian Rivers. *Chemosphere* 90, 358-365.

611 Sroda, S., Cossu-Leguille, C., 2011. Seasonal variability of antioxidant biomarkers and energy
 612 reserves in the freshwater gammarid *Gammarus roeseli*. *Chemosphere* 83(4), 538-544.

613 Starkweather, P.L., 1990. Zooplankton community structure of high elevation lakes: biogeographic
614 and predator-prey interactions. Internationale Vereinigung für Theoretische und Angewandte
615 Limnologie: Verhandlungen 24, 513-517.

616 Sun, H., Lü, K., Minter, E.J., Chen, Y., Yang, Z., Montagnes, D.J., 2012. Combined effects of
617 ammonia and microcystin on survival, growth, antioxidant responses, and lipid peroxidation of
618 bighead carp *Hypophthalmichthys nobilis* larvae. J. Hazard. Mater. 221-222, 213-219.

619 Tiberti, R., Tartari, G., Marchetto, A., 2010. Geomorphology and hydrochemistry of 12 Alpine
620 lakes in the Gran Paradiso National Park, Italy. J. Limnol. 69, 242-256.

621 Tiberti, R., von Hardenberg, A., Bogliani, G., 2014. Ecological impact of introduced fish in high
622 altitude lakes: a case of study from the European Alps. Hydrobiologia 724, 1-19.

623 Tiberti, R., Brighenti, S., Canedoli, C., Iacobuzio, R., Rolla, M., 2016. The diet of introduced brook
624 trout (*Salvelinus fontinalis*; Mitchill, 1814) in an alpine area and a literature review on its feeding
625 ecology. J. Limnol. 75, 488-507.

626 Wright, P., Mason, C.F., 1999. Spatial and seasonal variation in heavy metals in the sediments and
627 biota of two adjacent estuaries, the Orwell and the Stour, in eastern England', Sci. Total. Environ.
628 226, 139–156.

629 Yang, R., Yao, T., Xu, B., Jiang, G., Xin, X., 2007. Accumulation features of organochlorine
630 pesticides and heavy metals in fish from high mountain lakes and Lhasa River in the Tibetan
631 Plateau. Environ. Int. 33, 151–156.

632

633

634

635

636

637

638

639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654

Table 1. Mean and standard deviation (three replicates) of physicochemical parameters and nutrients measured in Balma Lake in summer (August) and autumn (Autumn) 2017.

657
658
659
660
661
662
663
664

665

666

667

676

677

678

679

680

681

682

683

684

685

	Summer (August)	Autumn (October)
Temperature (°C)	15.62 ± 0.77	8.24 ± 0.53
Dissolved oxygen concentration (mg L ⁻¹)	7.40 ± 0.89	7.29 ± 1.82
Oxygen saturation (%)	92.88 ± 8.52	104.75 ± 2.43
pH	6.64 ± 0.28	7.69 ± 0.12
Conductivity (µS cm ⁻¹)	18.29 ± 1.12	18.56 ± 1.07
NH ₄ ⁺ (mg L ⁻¹)	0.09 ± 0.07	0.09 ± 0.07
NO ₃ ⁻ (mg L ⁻¹)	8.55 ± 2.73	6.82 ± 2.14
PO ₄ ³⁻ (mg L ⁻¹)	0.01 ± 0.01	0.02 ± 0.01


Table 2. Biometric values of females and males of brook trout (*Salvelinus fontinalis*) from Balma Lake in summer and autumn 2017.

	Summer (August)		Autumn (October)	
Sex	Males	Females	Males	Females
Number of individuals	8	12	6	14
Weight - W mean ± SD (g)	69.9 ± 30.34	71.20 ± 28.34	88.47 ± 56.23	90.05 ± 60.03
W min. (g)	15.00	15.36	10.23	11.00
W max. (g)	146.00	150.00	187.00	193.00
Total Length - TL mean ± SD (cm)	17.77 ± 3.90	18.56 ± 4.20	19.23 ± 4.66	19.35 ± 5.72
TL min. (cm)	11.50	11.00	7.50	8.00
TL max. (cm)	23.00	24.00	25.50	26.00

694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710

711 **Figure 1.** Balma Lake: bathymetry and sampling sites for physicochemical parameters (L = littoral
712 sites; B = deep sites). Red rectangles indicate the placement of gillnets to sample fish in summer
713 (August) and autumn (October) 2017.

714



715

716

717

718

719

720

721 **Figure 2.** Trace elements (Log₁₀ concentration; mg kg⁻¹) detected in the muscle tissue of brook
722 trout in summer (August) and autumn (October) 2017.

723




Figure 3. Superoxide dismutase (SOD) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.






Figure 4. Catalase (CAT) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.



764 **Figure 5.** Total glutathione peroxidase (GPx) activity in the liver, gills, kidney, spleen, and muscle
 765 tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S);
 766 Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences
 767 (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.

768



769

770 **Figure 6.** Selenium-dependent glutathione peroxidase (Se-GPx) activity in the liver, gills, kidney,
771 spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard
772 deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate
773 statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling
774 months for each tissue type.




Figure 7. Glutathione reductase (GR) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.






Figure 8. Glutathione S-transferase (GST) activity in the liver, gills, kidney, spleen, and muscle tissue of female and male brook trout. Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M). Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the sexes and the sampling months for each tissue type.



827 **Figure 9.** Metallothionein (MTs) concentration in the muscle tissue of female and male brook trout.
828 Data are reported as mean \pm standard deviation. Summer (S); Autumn (A); females (F); males (M).
829 Lower case letters indicate statistically significant differences (Tukey test, $p < 0.05$) between the
830 sexes and the sampling months for each tissue type.

831



832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848 **Figure 10.** Biplot of score and loadings from principal component analysis. The scores of each
849 organ (gill, kidney, liver, muscle, and spleen) are denoted by a color and a symbol (largest symbol =
850 average value). Confidence ellipses plot convex hull values of each organ.




Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)




Figure 6
[Click here to download high resolution image](#)




Figure 7
[Click here to download high resolution image](#)




Figure 8
[Click here to download high resolution image](#)





Figure 9
[Click here to download high resolution image](#)



[Click here to download high resolution image](#)




Table S2

[Click here to download Supplementary material for on-line publication only: correlation matrices.xlsx](#)

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

--