



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Treatment with the flavonoid 7,8-Dihydroxyflavone: a promising strategy for a constellation of body and brain disorders

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Emili M., Guidi S., Uguagliati B., Giacomini A., Bartesaghi R., Stagni F. (2022). Treatment with the flavonoid 7,8-Dihydroxyflavone: a promising strategy for a constellation of body and brain disorders. *CRITICAL REVIEWS IN FOOD SCIENCE AND NUTRITION*, 62(1), 13-50 [10.1080/10408398.2020.1810625].

Availability:

This version is available at: <https://hdl.handle.net/11585/775197> since: 2024-04-09

Published:

DOI: <http://doi.org/10.1080/10408398.2020.1810625>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4
5 **Title: Treatment with the Flavonoid 7,8-Dihydroxyflavone: a promising Strategy for a**
6
7
8 **Constellation of Body and Brain Disorders**
9

10
11
12 **Authors:** Marco Emili ^a, Sandra Guidi ^a, Beatrice Uguagliati ^a, Andrea Giacomini ^a, Renata
13
14 Bartesaghi ^a, Fiorenza Stagni ^b
15

16
17 **Address:**

18
19 a) Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy.

20
21 b) Department for Life Quality Studies, University of Bologna, Rimini, Italy.
22
23
24
25
26

27 **Corresponding author:**

28 Dr. FIORENZA STAGNI

29 Neurophysiology Research Unit, Department for Life Quality Studies (QUVI)

30 University of Bologna

31 Corso D'Augusto 237, 47921, Rimini, Italy

32 Via San Donato 19/2, 40127, Bologna, Italy

33 Tel +39 051 2095635

34 Tel +39 051 2091721

35 E-mail fiorenza.stagni2@unibo.it
36
37

38 **Co-Corresponding author:**

39 Prof. RENATA BARTESAGHI

40 Department of Biomedical and Neuromotor Sciences, Physiology Building

41 University of Bologna

42 Piazza di Porta San Donato 2

43 I-40126 BOLOGNA BO, Italy

44 Tel +39 051 2091727; Fax +39 051 2091737

45 E-mail renata.bartesaghi@unibo.it
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **ABSTRACT**
4
5
6
7

8 Flavonoids have long been known to exert benefits in various health problems. Among them, the BDNF
9 mimetic 7,8-Dihydroxyflavone (7,8-DHF) is emerging as a potential treatment for a constellation of brain
10 and body pathologies. During the past 10 years, more than 180 preclinical studies have explored the efficacy
11 of 7,8-DHF in animal models of different pathologies. The current review intends to be an exhaustive survey
12 of these studies. By providing detailed information on the rationale of the experimental design and outcome
13 of treatment, we will give the reader tools to critically interpret the achievement obtained so far. If we put
14 together each individual piece of this complex mosaic, a picture emerges that is full of promise regarding the
15 potential usefulness of 7,8-DHF for human treatment. Much has been done so far and we believe that the
16 time is now ripe to move from the bench to the bedside, in order to establish whether supplementation with
17 7,8-DHF may serve as therapy or, at least, as adjuvant for the treatment of pathologies affecting brain and
18 body functioning.
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 **Key words:** Nutraceuticals; 7,8-Dihydroxyflavone; BDNF-TrkB system; Antioxidants; Neuroprotection;
34 Body pathophysiology
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LIST OF ABBREVIATIONS

1
2
3
4
5 7M8H-flavone: 7-methoxy-8-hydroxyflavone
6 7,8-DHF: 7,8-Dihydroxyflavone
7 A β 40 and A β 42: Beta amyloid 40, Beta amyloid 42
8 AD: Alzheimer's disease
9 ALS: Amyotrophic lateral sclerosis
10 AMPK: Adenosine Monophosphate-Activated Protein Kinase
11 APP: Amyloid precursor protein
12 ASD: Autism Spectrum Disorder
13 BACE1: β -secretase 1
14 BBB: Blood-brain barrier
15 Bcl-2: B-cell lymphoma-2
16 β -CTF: β -secretase-cleaved C-terminal fragment
17 BDNF: Brain-derived Neurotrophic Factor
18 CA1, CA3: Cornu Ammonis 1, Cornu Ammonis 3
19 CaMKII: Ca²⁺/calmodulin-dependent protein kinase II
20 CAT: Catalase
21 CCI: Controlled Cortical Impact Model
22 CFC: Contextual Fear Conditioning
23 COII: Mitochondrial cytochrome oxidase II
24 CREB: c-AMP response element-binding protein
25 CTX-B: Cyclothaxin-B
26 DAG: Diacylglycerol
27 DG: dentate gyrus
28 DOX: Doxorubicin
29 DS: Down syndrome
30 ER- α : Estrogen receptor α
31 FMRP: Fragile X mental retardation protein
32 FPI: Fluid Percussion Injury
33 FST: Forced swim test
34 FXS: Fraxile X syndrome
35 GAP-43: Growth Associated Protein 43
36 GSH: Glutathione
37 HC: Hair cells
38 HD: Huntington's disease
39 HI: Hypoxia-ischemia
40 HO-1: Heme oxygenase 1
41 iNOS: Inducible nitric oxide synthase
42 IP3: inositol-1,4,5-triphosphate
43 I/R: Ischemia/Reperfusion
44 L&M: Learning and memory
45 LPS: Lipopolysaccharide
46 LTP: Long-term potentiation
47 MAPK: Mitogen-activated protein kinase
48 MCAO: Middle cerebral artery occlusion
49 MECP2: methyl-CpG binding protein 2
50 METH: Methamphetamine
51 MDA: Malonildialdeide
52 MDD: Major depressive disorder
53 MMPs: Matrix metalloproteinases
54 Mn-SOD: Mn-superoxide dismutase
55 mTOR: mammalian target of rapamycin
56 MWM: Morris Water Maze
57 NO: nitric oxide
58
59
60

1
2
3 NGF: Nerve growth factor
4 NOR: Novel Object Recognition
5 Nrf2: Nuclear factor erythroid 2-related factor 2
6 OA: Osteoarthritis
7 OGD: Oxygen-glucose deprivation
8 OPA1: Optic atrophy 1
9 PD: Parkinson's disease
10 PFC: Prefrontal cortex
11 PGC-1 α : Peroxisome proliferator-activated receptor- γ coactivator 1-alpha
12 PI3K: Phosphatidylinositol 3-kinase
13 PLC: Phospholipase C
14 PPI: Prepulse Inhibition
15 Rb: Retinoblastoma protein
16 RGC: Retinal ganglion cells
17 ROS: Reactive oxygen species
18 RTT: Rett syndrome
19 SGN: Spiral ganglion neurons
20 SOD: Superoxide dismutase
21 Shc: Src-homology 2-domain
22 Sp1: Specificity protein 1
23 SPT: Sucrose Preference Test
24 TBI: Traumatic brain injury
25 Trk: Tropomyosin receptor kinase
26 TFAM: Mitochondrial transcription factor 1
27 TST: Tail suspension test
28 UCP1: Uncoupling protein 1
29 VrK3: Vaccinia-related kinase 3
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. INTRODUCTION

Flavonoids are low molecular weight compound that belong to a class of plant secondary metabolites having a polyphenolic structure. They are not synthesized by animals but can be assumed with the diet through fruits, vegetables and certain beverages (Panche, Diwan, and Chandra 2016). Flavonoids are categorized into 6 subgroups (Flavones, Flavonols, Isoflavonoids, Flavanones, Chalcones, Anthocyanins), based on the carbon of the C ring on which the B ring is attached and the degree of unsaturation and oxidation of the C ring (Panche, Diwan, and Chandra 2016). In addition to their role in plants, flavonoids appear to play a number of protective functions in the human body due to their antioxidant, cytotoxic, anticancer, anti-inflammatory, antiviral, antiallergic, cardioprotective, and hepatoprotective properties (Nijveldt et al. 2001; Panche, Diwan, and Chandra 2016; Tapas, Sakarkar, and Kakde 2008). For this reason, flavonoids are now considered as an indispensable component in a variety of nutraceutical, and pharmaceutical applications.

7,8-Dihydroxyflavone (7,8-DHF) is a flavonoid belonging to the subclass of flavones that, similarly to other flavonoids, was known to exert anticancer action (Le Bail et al. 1998) and prevention of oxidative stress (Zhang et al. 2009). The discovery ten years ago that 7,8-DHF is a bioactive high-affinity agonist of the tropomyosin-related kinase B (TrkB) receptor for the brain-derived neurotrophic factor (BDNF) that crosses the blood brain barrier (BBB) and provokes TrkB autophosphorylation and activation of downstream signaling pathways represented a milestone in the history of this flavonoid (Jang et al. 2010). Various brain disorders that are related to deficits in BDNF-TrkB signaling cannot be “cured” with exogenous BDNF due to its unfavorable pharmacokinetic properties. 7,8-DHF, however, is a BDNF mimetic that crosses the BBB and, therefore, can be pharmacologically exploited to treat BDNF-related brain disorders. Indeed, the pioneering study by Jang et al. provided preliminary evidence for a positive effect of treatment with 7,8-DHF *in vivo* for stroke and Parkinson disease (Jang et al. 2010). This evidence heavily drew the attention of the scientific community and prompted a number of studies aimed at establishing the therapeutic potential of 7,8-DHF in different brain diseases. The growing interest for 7,8-DHF as potential medicine is documented by results of a search in Pub Med using the keyword “7,8-Dihydroxyflavone”. This search yielded a total 203 publications, 16 of which were in the period 1992-2009 and 187 in the period 2010-2020. Moreover, t

1
2
3
4 publication rate in the last decade has grown exponentially, as shown in Fig. 1. This is not surprising
5
6 considering the therapeutic potential of BDNF and, hence, 7,8-DHF in a variety of neurological disorders. In
7
8 addition, 7,8-DHF being a flavonoid has the added value of being an antioxidant, which may be of relevance
9
10 by itself and because BDNF-related pathologies may also be associated with metabolic dysregulation and
11
12 oxidative stress.

13
14 The available reviews regarding the therapeutic potential of 7,8-DHF were mainly focused on brain
15
16 disorders (Du, and Hill 2015; Liu, Chan, and Ye 2016). It must be noted, however, that 7,8-DHF exerts
17
18 important effect on numerous body organs and functions, which anticipates a wide scope of action of this
19
20 molecule in the framework of human health. For this reason, we thought it may be important to include in
21
22 this review studies that have examined the effects of 7,8-DHF both on the brain and body. This review was
23
24 additionally prompted by the idea to give the reader detailed information of most of the preclinical studies on
25
26 7,8-DHF published so far. Our description will be analytical in terms of i) categorization of the effects of
27
28 7,8-DHF, disease by diseases and ii) abundance of experimental details (e.g., doses, duration of treatment,
29
30 utilized strain, age, etc.). Each section is organized in a brief premise, that provides propaedeutic
31
32 information, the description of the actual studies on a given topic, and a brief comment on the potential
33
34 significance of the experimental evidence. In addition, we have summarized in two tables (Table 1 and Table
35
36 2) the animal models, doses, route of administration used in all the reviewed studies and in a third table
37
38 (Table 3) the studies that have investigated a given brain parameter (e.g., neurogenesis, spine density etc.) to
39
40 help the reader quickly gather available evidence on the topic/s of interest. We hope that this effort may
41
42 provide a clear and exhaustive view of the state of the art of preclinical research regarding the benefits of
43
44 7,8-DHF. This, in turn, may help to design new targeted studies founded on the existing body of knowledge
45
46 and to individuate possible gaps to be addressed in future research aimed at expanding the characterization of
47
48 this very promising molecule.
49
50
51
52
53

54 **2. BDNF AND TRKB SIGNALING**

55 **2.1 Brief overview on neurotrophins**

56
57 Neurotrophins are a family of protein regulators of neural survival, development, function, and plasticity in
58
59 the central and peripheral nervous system (see (Huang, and Reichardt 2001)). Nerve growth factor (NGF)
60

1
2
3 was the first identified and best characterized member of this family (Levi-Montalcini 1987). There are four
4 neurotrophins characterized in mammals: NGF, BDNF, neurotrophin-3 (NT-3), and neurotrophin-4/5 (NT-
5 4/5). Neurotrophins bind two types of receptors: the tropomyosin receptor kinase (Trk) and the so called low-
6 affinity NGF receptor p75 (p75NTR). The Trk receptors mediate specificity of the neurotrophic functions of
7 neurotrophins. There are three main types of Trk receptors: TrkA is a receptor for NGF, NT-6 and NT-7;
8 TrkB is a receptor for BDNF and NT-4/5; and TrkC is a receptor for NT-3. Since 7,8-DHF specifically binds
9 to the TrkB receptor for BDNF, we will focus here on the BDNF-TrkB system.

10
11 **2.2 Synthesis of BDNF** The human *BDNF* gene (~70 kb) consists of 11 exons, located on chromosome 11,
12 and its different splicing enables formation of transcripts specific to various tissues and responsive to
13 different stimuli (see(Cattaneo et al. 2016)). These transcripts, however, are translated and processed into
14 identical pro-BDNF protein. The BDNF protein is synthesized and folded in the endoplasmic reticulum as a
15 precursor form, pre-pro-BDNF (~ 27 kDa). The pre-region is then cleaved, resulting in formation of pro-
16 BDNF. The pro-BDNF is then further processed *via* the Golgi apparatus where the pro-domain is
17 proteolytically cleaved off to form the pro-domain (of 129 amino acids) and the mature domain m-BDNF (of
18 118 amino acids) also simply named BDNF. The ratio of pro-BDNF to m-BDNF varies between particular
19 stages of brain development and regions. While in the early postnatal period higher concentration of pro-
20 BDNF is reported, m-BDNF prevails in adulthood (Yang et al. 2014a). While pro-BDNF interacts
21 preferentially with the p75NTR, m-BDNF binds the TrkB receptor.

23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 **2.3 The TrkB receptor**

44
45 The TrkB receptor is encoded by a single *TrkB* gene, the *NTRK2* gene encoding 24 exons (350 kbp) located
46 on humans chromosome 9q22 (see (Pradhan, Noakes, and Bellingham 2019)). The TrkB protein consists of
47 three different domains: an extracellular ligand binding domain (encoded by exons 5-14), a transmembrane
48 domain (encoded by exon 15) and an intracellular tyrosine kinase domain (encoded by exons 20-24). Exon
49 12 encodes an IG-like domain that has been postulated to be the region responsible for binding specificity to
50 its ligand, BDNF (Middlemas, Lindberg, and Hunter 1991). The extracellular transmembrane domain and the
51 intracellular cytoplasmic domain of the TrkB protein full length (TrkB F.L.) act as phosphorylation-
52 dependent docking sites and rapidly transmits the effects of neurotrophin binding to downstream network.

1
2
3 Alternative splicing at exon 18 of the primary transcript produces truncated TrkB isoforms (TrkB-T1 and
4 TrkB-T-shc in humans; T1 and T2 in rat) that lack the intracellular tyrosine kinase domain (Fig. 2) and
5 cannot elicit rapid intracellular signaling (Stoilov, Castren, and Stamm 2002). BDNF activation of TrkB-T1
6 and TrkB-T2 has been shown to increase the rate of acidic metabolites release from the cell (Baxter et al.
7 1997). TrkB-T1 can behave as a dominant negative isoform by competing with neurotrophins and forming
8 heterodimers with the TrkB F.L. In human, the isoform TrkB-T1 is predominantly expressed in the brain but
9 also detected in heart, kidney and pancreas (see (Gupta et al. 2013a)).

17 **2.4 BDNF-TrkB signaling**

19 The binding of BDNF to the TrkB receptor, causes receptor dimerization and autophosphorylation of
20 intracellular tyrosine residues (Y705/6), which results in the formation of the phosphorylated-TrkB receptor
21 (see (Rantamaki et al. 2007)). This, in turn, launches autophosphorylation of other tyrosine residues situated
22 outside the activation loop (Y515 and Y816) that serve as docking sites for adaptor proteins. Phosphorylated
23 TrkB sets in motion three interconnected signaling transduction pathway with specific cellular functions
24 (Fig. 3): a) Phosphatidylinositol 3-kinase (PI3K), b) mitogen-activated protein kinase (MAPK) and c)
25 phospholipase C γ (PLC γ) (see (Chao 2003; Rantamaki et al. 2007)).

26 a) Activation of the PI3K pathway incorporates combined actions of Ras at the Tyr515 residue, which
27 activates the PI3K/Akt and MEK/MAPK pathways. The **PI3K/Akt-related pathway** results in the activation
28 of B-cell lymphoma 2 (Bcl-2)-associated agonist of cell death (BAD) and inhibition of the kinase glycogen
29 synthase kinase 3 β (GSK-3 β), which in turn leads to a reduction in the phosphorylation of collapsing
30 response mediator protein 2 (CRMP-2), a protein that is highly expressed in the developing of nervous
31 system and that regulates axonal outgrowth and dendritic branching. Activation of mammalian target of
32 rapamycin (mTOR) by PI3K/Akt pathway enhances local BDNF translation to dendrites at active synapses,
33 enhances dendritic growth and branching.

34 b) Activation of the TrkB receptor at Tyr490 and Tyr515 residue causes the docking of Src-homology 2-
35 domain (Shc) adaptor protein at these tyrosine sites and recruits growth factor receptor bound protein 2
36 (grb2) which binds with GTPase Ras to form a complex, and initiates activation of the **MAPK/ERK**
37 **pathway** that is an important regulator of cell survival and neuronal differentiation. The MAPK/ERK
38 signaling cascade is able to phosphorylate and activate the transcription factor cAMP response element-

1
2
3 binding protein (CREB). Phosphorylated CREB is then translocated into the nucleus, where it induces BDNF
4 transcription by binding to BDNF promoters. Binding to BDNF promoters drives BDNF expression to
5 regulate neuronal survival, differentiation and synaptic plasticity. Moreover, ERK/CREB regulates the
6 transcription of genes required for dendritic growth including proteins that regulate cytoskeleton dynamics
7 (e.g., Arc and cypin).
8
9

10
11
12
13 c) The phosphorylation of the TrkB receptor at its Tyr816 residue activates the **PLC γ pathway** with
14 consequent intracellular generation of inositol-1,4,5-triphosphate (IP₃) and 1,2-diacylglycerol (DAG). The
15 PLC γ /IP₃ pathway induces the release of calcium from intracellular stores. Calcium activates
16 Ca²⁺/calmodulin-dependent protein kinase (CAMKII) which in turn induces CREB phosphorylation. The
17 generation of DAG activates PKC that is translocated to the membrane for further activation and
18 phosphorylation of ERK leading to survival, neuritic outgrowth, and synaptic plasticity. Activation of PLC γ
19 appears to be dispensable for increasing dendritic branching but is required to induce hippocampal long-term
20 potentiation (LTP).
21
22
23
24
25
26
27
28
29

30 **2.5 TrkB activation without neurotrophins**

31
32 The TrkB receptor can execute autophosphorylation activity and downstream signaling without stimulation
33 by its ligand. BDNF-independent TrkB activation occurs *via* a mechanism known as trans-activation (see
34 (Chao 2003; Sasi et al. 2017)). For instance, activation of TrkB receptors in the absence of neurotrophins can
35 be mediated by the G-protein-coupled adenosine 2A receptor (A2A-R) or the dopamine D1 receptor. This
36 effect is mediated by Src family of protein tyrosine kinases. Transactivating agents of TrkB include pituitary
37 adenylate cyclase-activating polypeptide (PACAP), epidermal growth factor (EGF), zinc, and H₂O₂ (Huang,
38 and McNamara 2012).
39
40
41
42
43
44
45
46

47 **2.6 TrkB inhibitors**

48
49 K252a is an alkaloid (467.48 Da) that is able to penetrate the cell and act as a potent Trk receptor inhibitor
50 (Boulle et al. 2012). However, the lack of specificity for TrkB, as compared to TrkA and TrkC, constitutes a
51 major limitation for the use of K252a. More recently, two specific and potent TrkB antagonists have been
52 developed, ANA-12 and cyclotraxin-B. ANA-12 (407.49 Da), which has been identified by an *in silico*
53 screening approach, crosses the BBB, binds the extracellular domain of TrkB, prevents BDNF-induced TrkB
54 activation, and abolishes the biological effects of BDNF but not those of NGF or NT-3 on TrkA- and TrkC-
55
56
57
58
59
60

1
2
3 expressing cells (Cazorla et al. 2011). Cyclotraxin-B (CTX-B), a cyclic peptide (1200 Da), is a TrkB antagonist
4 that crosses the BBB and allosterically alters the conformation of the receptor, leading to long-lasting inhibition
5 of BDNF activity. The action of this molecule is mediated by the link with the TrkB binding extracellular
6 domains, that are not critical for the interaction with BDNF but rather involved in its activation capacity (Cazorla
7 et al. 2010). CTX-B inhibits the two main signaling pathways downstream of TrkB, the MAPK- and PLC γ -
8 pathway, with similar amplitude of inhibitory effects (Cazorla et al. 2010). Very recently, a novel inhibitor has
9 been discovered, GZD2202, that has a moderate selectivity between Trk B/C and TrkA (Zou et al. 2019).
10 GZD2202 dose-dependently inhibits the phosphorylation of TrkB and its downstream signaling proteins Akt,
11 ERK, PLC γ .
12
13
14
15
16
17
18
19
20
21
22

23 **3. PHARMACOLOGY AND SIGNALING OF 7,8-DHF**

24 **3.1 Pharmacology of 7,8-DHF and its prodrugs**

25 **3.1.1 Sources and Chemistry**

26
27 7,8-DHF is a flavone substituted by hydroxy groups at positions 7 and 8 (Fig. 4). It is naturally present in
28 *Godmania aesculifolia*, *Tridax procumbens* and various species of *Primula*. *Godmania aesculifolia* is a
29 species of flowering plant belonging to the Bignoniaceae family. Bignonia plants are widely used in
30 traditional medicine as remedy for skin ailments, postpartum hemorrhage, malaria, diabetes and pneumonia
31 (Mahmoud et al. 2019). Flavonoids are the most abundant class in different *Bignonia* plants and are
32 contained in the roots and leaves. *Tridax procumbens* L. (*Tridax*) is a flowering plant belonging to the
33 Asteraceae family that has been traditionally in use from ancient times (Beck et al. 2018). *Tridax* has various
34 pharmacological properties including immunomodulatory, antioxidant, anti-hepatotoxic, analgesic,
35 antidiabetic, anti-inflammatory, antifungal, and antimicrobial activities. Flavonoids are found in the leaves and
36 other organs. Different substances such as powdered leaves, decoctions, oils, teas, leaf juice, skin
37 cataplasms have been produced using this species (Beck et al. 2018). Flowers, leaves and roots of various
38 *Primula* species contain different flavonoids (Colombo et al. 2014; Jäger et al. 2006). *Primula* flower and
39 root preparations, which are usually available as herbal tea to be drunk and in liquid forms to be taken by
40 mouth are approved by EMA and can be used for coughs associated with colds (Web reference 1; Web
41 reference 2). Flowers, leaves and roots are used in Danish folk medicine against convulsions (Jäger et al.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2006). 7,8-DHF was identified from 2,000 biologically active compounds through the screening of the
4
5 Spectrum Collection Library as a high affinity TrkB agonist, that provokes dimerization and phosphorylation
6
7 of TrkB, and activation of downstream signaling cascades (Jang et al. 2010). A structure-activity relationship
8
9 study suggests that the catechol group (7,8-dihydroxy on the A ring) is indispensable for the agonistic
10
11 activity, in particular the 8-hydroxy group in A ring is essential for the TrkB stimulatory effect (Liu et al.
12
13 2010).

14 15 **3.1.2 Pharmacokinetics**

16
17 The *in vivo* investigation of 7,8-DHF pharmacokinetics profile determined that the plasma half-life of 7,8-
18
19 DHF in mice, after oral gavage of 50 mg/kg, was about 134 min. The plasma concentration peaked at 10 min
20
21 with 70 ng/ml and was still detectable after 8 h (5 ng/ml). 7,8-DHF was able to cross the BBB and peaked at
22
23 10 min with concentration of 50 ng/g of brain, at 4 h its concentration was 7 ng/g and at 6 h was below the
24
25 quantification limits (Liu et al. 2013). Catechol-containing compounds are usually cleared in the circulatory
26
27 system and undergo oxidation and conjugation (glucuronidation, sulfation and methylation). From a
28
29 chromatographic analysis of plasma samples, metabolites were detected in addition to the parent drug,
30
31 including 7-methoxy-8-hydroxyflavone (7M8H-flavone), 7-hydroxy-8-methoxyflavone (7H8M-flavone) and
32
33 the glucuronidated product of 7,8-DHF (Liu et al. 2013). In plasma, 7M8H-flavone was detected after 3 min,
34
35 its concentration peaked at 10 min (1.0 ng/ml), decreased to 0.2 ng/ml at 4 h and was below the
36
37 quantification limits at 6 h (Liu et al. 2013). In brain samples, 7M8H-flavone was below the limit of
38
39 quantification, whereas the 8M7H-flavone was observable 3 min after oral administration and still detectable
40
41 at 240 min (Liu et al. 2013). The *in vivo* agonistic effects exerted by 7,8-DHF are thought to mainly result
42
43 from the activity of 7,8-DHF and its metabolite 8M7H-flavone (Liu et al. 2013). Indeed, Catechol-O-
44
45 methyltransferase (COMT) inhibition reduced the TrkB activation exerted by 7,8-DHF, indicating that the
46
47 methylated metabolite contributes to TrkB receptor activation after oral administration (Liu et al. 2013). A
48
49 study in monkey plasma showed that 7,8-DHF has longer half-life in monkeys compared to mice. The half-
50
51 life of 7,8-DHF was approximately 4-8 h during chronic oral treatment with 30 mg/kg/day (He et al., 2016).
52
53 The study of 7,8-DHF metabolism in monkeys confirmed the methylation, glucuronidation and sulfation
54
55 clearance mechanisms observed in mice (He et al. 2016; Sun et al. 2017).

56 57 **3.1.3 Bioavailability**

1
2
3 Oral administration of 7,8-DHF in mice elicits robust TrkB activation in the brain, suggesting that this
4 compound is orally-bioavailable (Liu et al. 2010). Both the methylated metabolites of 7,8-DHF are able to
5 activate TrkB in mouse brain and in primary neurons (Liu et al. 2013). However, the catechol group makes 7,8-
6 DHF labile for fast metabolism, including sulfation, glucuronidation, methylation. Growing evidence suggests
7 that intestinal microbiota increase the bioavailability and physiological function of dietary polyphenols.
8 These metabolites may have larger bioavailability and exert larger biological effect than the parent compound
9 (Murota, Nakamura, and Uehara 2018). No specific evidence exists regarding the role of gut microbiota in the
10 metabolism of 7,8-DHF. However, it is conceivable that bioactive phenolic acids derived from the
11 catabolism of 7,8-DHF elicited by gut microbiota may increase the biological activity of 7,8-DHF and that
12 interpersonal variability in the microbiota composition may underlie different biological responses following
13 oral administration of 7,8-DHF. Moreover, an O-methyltransferase SpOMT2884, originating from
14 *Streptomyces peucetius* ATCC 27952 has been shown to convert 7,8-DHF into 7-hydroxy-8-methoxyflavone in
15 an *in vitro* system (Koirala et al. 2014). This metabolite is more stable than 7,8-DHF and exhibits potent
16 antioxidant effects without inducing cytotoxicity. Interestingly, *Streptomyces* has been shown to be present in
17 gut microbiome (Bolourian, and Mojtahedi 2018), suggesting that these and possibly other gut bacteria may
18 impact on the bioavailability and activity of 7,8-DHF. Several prodrugs of 7,8-DHF have been synthesized and
19 tested in order to improve the bioavailability of 7,8-DHF. A recent study identified the compounds R13
20 and R7 as the prodrugs with the best pharmacokinetic profile (Chen et al. 2018). R13 improved the oral
21 bioavailability from 4.6% to 10.5% (Chen et al. 2018) and R7 from 4.6% to 82.2% (Liu et al. 2013). R13
22 appeared to be the prodrug with the best pharmacokinetics profile. Accordingly, it was found to be readily
23 hydrolyzed into 7,8-DHF in liver microsomes and following oral administration of 78 mg/kg of R13 (equal to
24 50 mg/kg of 7,8-DHF) the concentration of 7,8-DHF in the mouse brain was significantly greater (>8 ng/gr
25 even after 4 h) than the parent compound DHF (below the quantification limit at 6 h) (Chen et al. 2018).

3.1.4 Toxicity

26 Exposure to 7,8-DHF did not impair viability of the human kidney cell line HK-2 up to a concentration of 200
27 (Ma et al. 2016). Accordingly, an $IC_{50} > 100 \mu M$ has been established for cytotoxicity in HepG2 cells exposed for
28 24 h to 7,8-DHF (Kozics, Valovicova, and Slamenova 2011). Neuronal cell death assay revealed
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that 7,8-DHF is non-toxic for primary cortical neurons up to 5 μ M (Liu et al. 2010). In mice treated with 5
4 mg/kg/day of 7,8-DHF for 3 weeks no pathological changes were found in the kidney, liver, lung, muscle,
5 spleen, cortex, hippocampus, heart, intestine and testis in comparison with the vehicle treated counterparts
6 (Liu et al. 2013). Moreover, from complete blood count analysis no significant difference appeared between
7 7,8-DHF-treated and control mice. This is consistent with another long-term experiment where female mice
8 receiving 7,8-DHF (~2.4 mg/kg/day) for 20 weeks displayed normal complete blood count values (Chan et
9 al. 2015). In addition, a study in monkeys showed that chronic treatment with 7,8-DHF for 7 months with a
10 dose of 30 mg/kg/day, a dose higher than the one commonly used in rodents (5 mg/kg), did not cause toxicity
11 (assessed by evaluating hematology and liver, heart, muscle, kidney and pancreas histology) and health
12 problems (He et al. 2016). Taken together, these data suggest that chronic treatment with 7,8-DHF does not
13 exert toxic effects.

24 **3.2 7,8-DHF signaling**

25 **3.2.1 TrkB-dependent signaling**

26 7,8-DHF specifically binds the TrkB (but not TrkA or TrkC) receptor extracellular domain with a binding
27 constant $K_d = 320$ nM, acting as a selective TrkB agonist and mimicking the physiological actions of BDNF
28 (Jang et al. 2010; Liu et al. 2014). 7,8-DHF appears, however, to bind to a different site of TrkB extracellular
29 domain in comparison with BDNF (Liu et al. 2013). Similarly to BDNF, 7,8-DHF induces phosphorylation
30 of TrkB at Tyr515, Tyr706, and Tyr816 although 7,8-DHF-induced phosphorylation of Tyr515 and Tyr816
31 is less robust in comparison with that induced by BDNF (Liu et al. 2014). 7,8-DHF-mediated TrkB
32 phosphorylation leads to activation of the MAPK, PI3K/Akt, and ERK1/2 pathways in a time frame that is
33 comparable to BDNF and in a dose-dependent manner (Jang et al. 2010; Liu, Chan, and Ye 2016; Liu et al.
34 2010; Liu et al. 2014; Liu et al. 2013). Internalization of the neurotrophin-Trk complex, leading to formation
35 of signaling endosomes and ubiquitination/degradation, plays an important role in signal transduction. While
36 BDNF is more potent than 7,8-DHF in stimulating TrkB internalization and early endosomes delivery in the
37 first 10 min, 7,8-DHF produces a more robust endocytic response than BDNF at 60 min after stimulation.
38 BDNF induces TrkB ubiquitination 10 min after stimulation and its ubiquitination signals correlates with its
39 phosphorylation pattern. While BDNF-triggered TrkB phosphorylation peaks at 10 min, decreases at 60 min
40 and fades away at 180 min, 7,8-DHF-triggered TrkB phosphorylation lasts for more than 3 h, without

1
2
3 inducing TrkB ubiquitination or degradation (Liu et al. 2014). These findings support the idea that 7,8-DHF
4 and BDNF activate TrkB with different mechanisms and with a different time-course.
5
6

7 **3.2.2 TrkB-independent signaling**

8
9 A recent study in cell assays failed to confirm that 7,8-DHF activates the TrkB receptor (Boltaev et al. 2017).
10 Further studies are needed to explain this discrepancy which may be due to the used cell model, culture
11 condition and duration of treatment. In any case, the possibility must be taken into account that 7,8-DHF
12 may induce transactivation of the TrkB receptor without the need to binding to it. Indeed, some studies show
13 that some effects (in particular, antioxidant effects) of 7,8-DHF do not require the TrkB receptor as they are
14 generated in conditions in which the receptor is pharmacologically inhibited or in cells that do not express
15 the TrkB receptor (Garcia-Diaz Barriga et al. 2017; He et al. 2018; Huai et al. 2014; Ryu et al. 2014a). In
16 addition, there is evidence for uptake of various flavonoids and their metabolites into the cytosol of several
17 cell types, including cells of the central nervous system (CNS) (Ferrara, and Thompson 2019; Mukai et al.
18 2011; Spencer, Abd-el-Mohsen, and Rice-Evans 2004). If this also holds for 7,8-DHF, its cell uptake will
19 provide the possibility of a TrkB-independent direct modulation of signaling pathways such as the MAPK or
20 PI3K/Akt pathway (Williams, Spencer, and Rice-Evans 2004). In addition, being an antioxidant, 7,8-DHF
21 might exert a direct antioxidant action within the cell which may contribute to its powerful antioxidant
22 effects (Han et al. 2014).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **4. ACTIONS OF 7,8-DHF IN THE CENTRAL NERVOUS SYSTEM**

42 BDNF is present in almost all brain regions and is involved in a number of biological functions that include
43 regulation of neurogenesis, gliogenesis, synaptogenesis, neuroprotection, synaptic plasticity, memory and
44 cognition (see (Kowianski et al. 2018)). In view of its multifaceted roles, it is not surprising that BDNF
45 dysregulation is involved in the pathogenesis of a plethora of brain disorders. Although BDNF may represent
46 a beneficial therapeutic agent against these disorders, its poor brain delivery and short half-life represent a
47 serious issue. In view of its chemical and functional properties (Liu, Chan, and Ye 2016), however, 7,8-DHF
48 may represent a key therapeutic alternative to the use of BDNF. In this section we will review preclinical
49 studies showing that 7,8-DHF exerts a myriad of benefits in the CNS.
50
51
52
53
54
55
56
57
58
59

60 **4.1 Genetic disorders**

4.1.1 Down syndrome

Brief premise. Down syndrome (DS) is a genetic pathology due to triplication of chromosome 21. The brain of individuals with DS is hypotrophic starting from fetal life stages due to neurodevelopmental defects that include neurogenesis reduction, dendritic pathology, and altered brain wiring (Stagni et al. 2018). These defects underlie the typical cognitive impairment in DS. In spite of numerous preclinical studies (Stagni et al. 2015), no effective treatment is currently available for DS. Reduced BDNF levels have been found in the DS brain (see (Stagni et al. 2017)), suggesting that this defect may concur to alter brain development in DS. Based on this premise, three studies have explored the effects of 7,8-DHF in the Ts65Dn mouse model of DS.

The evidence. Stagni et al. (Stagni et al. 2017) treated Ts65Dn mice with 7,8-DHF in the early postnatal period, a critical time window for hippocampal development. Pups treated with 7,8-DHF from P3 to P15 underwent an improvement in hippocampal neurogenesis, rescue of total granule cell number, granule neuron dendritic spine density, and hippocampal synaptophysin levels. These effects were accompanied by an increase in hippocampal p-TrkB (Y816) and p-ERK1/2 levels. Mice treated with 7,8-DHF from infancy to adolescence (P3-P45) exhibited full rescue of learning and memory (L&M) (Morris Water Maze test, MWM). A second study (Giacomini et al. 2019) showed that the positive effects of treatment during infancy on hippocampal development did not translate into a behavioral benefit in adulthood if treatment was discontinued, indicating the necessity of chronic administration of 7,8-DHF. This study additionally shows that in 4-month-old mice treatment with 7,8-DHF for 40 days with the same dose that was effective in pups had no effect on neurogenesis, L&M (MWM) and hippocampal p-TrkB (Y816) levels. At variance with this latter study, Parrini et al. (Parrini et al. 2017) showed that adult Ts65Dn mice treated with 7,8-DHF for 4 weeks underwent restoration of hippocampal neurogenesis, LTP in field CA1, memory (contextual fear conditioning, CFC; Novel Object recognition, NOR) and an increase in hippocampal p-TrkB levels. This discrepancy may be explained by the higher dose of 7,8-DHF (~ 22 mg/kg) used by Parrini et al., while Giacomini et al. used a dose of 5 mg/kg.

Significance. These studies show that 7,8-DHF fully restores hippocampal development and cognitive performance if administered during early life stages. Its effects, however, are ephemeral suggesting that continuous administration is needed. This may be feasible in view of the lack of toxicity of 7,8-DHF. Thus,

1
2
3 in view of its safe profile, treatment with 7,8-DHF may represent a promising therapeutic opportunity for
4 children with DS.
5

6 7 **4.1.2 Rett syndrome** 8

9 **Brief premise.** Rett syndrome (RTT), is a rare disease characterized by intellectual disability and a host of
10 neurological hallmarks that include stereotypic hand movements, impaired motor coordination, autonomic
11 dysfunction, seizure and loss of language skills. RTT is due to loss of function of a gene encoding the
12 transcription factor methyl-CpG binding protein 2 (MECP2). MECP2 is a transcription regulator of several
13 genes, including BDNF (Li, and Pozzo-Miller 2014). Accordingly, strategies for improving BDNF signaling
14 appear to exert some benefit in RTT mouse models (Li, and Pozzo-Miller 2014).
15
16
17
18
19
20
21

22 **The evidence.** A single study examined the effect of treatment with 7,8-DHF in a model of RTT (Johnson et
23 al. 2012). *Mecp2* mutant mice that were treated with 7,8-DHF for one month lived significantly longer in
24 comparison with untreated mice, displayed a delay in weight loss, an improvement in the size of
25 hippocampal CA1 neuronal nuclei, in voluntary locomotor distance (wheel-running activity) and in the
26 breathing irregularity that is typical of RTT.
27
28
29
30
31

32 **Significance.** The effects of 7,8-DHF in the *Mecp2* mutant mouse mimic some of the effects obtained by
33 manipulating BDNF levels in RTT models (Li, and Pozzo-Miller 2014). This predicts the potential
34 usefulness of treatment with the BDNF mimetic 7,8-DHF in RTT patients to compensate for the loss of
35 BDNF expression. Further evidence, however, is necessary to confirm and better characterize the efficacy of
36 7,8-DHF in RTT.
37
38
39
40
41
42

43 **4.1.3 Fragile X syndrome** 44

45 Fragile X syndrome (FXS) is characterized by moderate-to-severe intellectual disability. It is caused by
46 silencing of the fragile X mental retardation 1 (*FMR1*) gene with consequent absence/reduction of the fragile
47 X mental retardation protein (FMRP), an RNA-binding protein that regulates the translation of neuronal
48 mRNAs at synapses (Kumari, and Gazy 2019). The absence of FMRP causes abnormal dendritic spine
49 formation and altered synaptic plasticity. Absence of FMRP reduces BDNF expression and hampers TrkB-
50 mediated protein synthesis (see (Tian et al. 2015)) suggesting that BDNF-TrkB signaling may play a role in
51 the pathophysiology of FXS.
52
53
54
55
56
57
58
59
60

1
2
3 **The evidence.** Tian et al. (Tian et al. 2015) tested the effects of 7,8-DHF in the *Fmr1* KO mouse model of
4 FXS. Weaning mice that received 7,8-DHF for 4 weeks exhibited restoration of L&M (MWM, CFC), and of
5 the aberrantly high spine density in neurons of field CA1, and reinstatement of p-TrkB (Y816) levels and
6 downstream targets (p-CaMKII and p-PKC α / β II) in synaptosomes of the hippocampus and amygdala.
7 Moreover, treatment enhanced the phosphorylation of GluA1- and GluA2-containing AMPARs. In a recent
8 study, Seese et al. (Seese et al. 2020) examined the effect of 7,8-DHF in adult *Fmr1* KO mice. In
9 hippocampal slices from *Fmr1* KO mice exposure to 7,8-DHF normalized the reduced magnitude of LTP. In
10 *in vivo* experiments, mice were acutely (one injection), semi-chronically (4 days) or chronically (one month)
11 treated with 7,8-DHF. Irrespective of its duration, treatment rescued spatial memory (object location
12 memory). In addition, a single injection of 7,8-DHF prior to training was sufficient to rescue visual
13 recognition memory (NOR). Finally, acute treatment with 7,8-DHF increased activation of TrkB (Y515) in
14 stratum radiatum of field CA1.

15
16
17
18
19
20
21
22
23
24
25
26
27
28 **Significance.** These two studies demonstrate that 7,8-DHF improves synaptic structure, hippocampal
29 synaptic plasticity and memory in a model of FXS. Interestingly, a single administration of 7,8-DHF is
30 sufficient to restore LTP and spatial/object memory, which suggests a fast restoring effect of the synaptic
31 plasticity alterations that characterize FXS. While these results are encouraging in the framework of potential
32 therapies for individuals with FXS, the duration of the beneficial effects of treatment on synaptic functioning
33 and cognitive performance remains a key unanswered issue.

4.2 Psychiatric disorders

4.2.1 Autism

44
45 **Brief premise.** Autism Spectrum Disorder (ASD) is a developmental disorder characterized by deficits in
46 social behavior, such as inability to initiate social interactions or develop relationships, lack of social or
47 emotional reciprocity, lack of interest in the emotions of others and communication deficits (Lai, Lombardo,
48 and Baron-Cohen 2014). Various animal models incorporating ASD-relevant behavioral phenotypes have
49 been developed, to identify the alterations underlying this disorder and to test possible pharmacotherapies.

50
51
52
53
54
55
56 **The evidence.** Rhine et al. (Rhine et al. 2019) studied the effects of different drugs, including 7,8-DHF, on
57 the behavioral phenotype of two autism mouse models, BTBR and Shank3B. Mice received 7,8-DHF 5 or 7
58 days before a battery of behavioral tests and on each testing days (60 min before the start). BTBR mice
59
60

1
2
3 showed restoration of sociability (three-chamber test and social sniffing) and Shank3B mice showed an
4 improvement in learning but not memory (MWM). The effect of 7,8-DHF was tested in the vaccinia related
5 kinase 3 (*Vrk3*) KO mouse model generated by Kang et al. (Kang et al. 2017). The *Vrk3*-KO mouse shows
6 unusual social interactions and repetitive behaviors similarly to autism subjects. Acute treatment with 7,8-
7 DHF restored hippocampal levels of p-TrkB (Y706) and social interaction (three-chamber test) but the latter
8 effect was not retained at 7 days after treatment cessation. Chronic treatment restored spine density in CA1
9 pyramidal neurons and suppressed elevated grooming activity, improved abnormal social activity and L&M
10 (passive avoidance and NOR). These behavioral effects were retained for several weeks after treatment
11 cessation. Lee (Lee, and Han 2019) induced an autistic-like behavior in a heterozygous mouse for the D2
12 receptor by administering an early-life stress (through maternal separation) to the pups. A three-day
13 treatment with 7,8-DHF increased social interactions (three-chamber test), suppressed increased grooming
14 and increased the expression of *TrkB*, *Bdnf*, *Mecp2* and *Hdac2* genes in the dorsal striatum. Inhibition of
15 TrkB in the dorsal striatum during treatment with 7,8-DHF by injections of TrkB-siRNA prevented the
16 treatment-induced behavioral rescue.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33 **Significance.** ASD is heterogeneous in etiology (interaction between environmental factors and genetic
34 variants) and symptoms. Thus, ASD models can only approximate the complexity of the human condition.
35 Yet, the studies reviewed above suggest that 7,8-DHF could be a pharmacological strategy to ameliorate the
36 defects in social behavior, a common central feature of ASD.

4.2.2 Schizophrenia

42
43 **Brief premise.** Schizophrenia, a complex psychiatric disorder that arises from genetic susceptibility and
44 environmental factors, is characterized by a combination of psychotic symptoms and cognitive deficits such
45 as impairments in L&M, attention, and executive functions. Reduced BDNF levels in serum (Toyooka et al.
46 2002) and reduced expression of BDNF in multiple cortical areas (Ray, Shannon Weickert, and Webster
47 2014) have been found in subjects with schizophrenia suggesting that impairment of BDNF-TrkB signaling
48 may contribute to the cognitive deficits that characterize this pathology.

49
50
51
52
53
54
55
56 **The evidence.** Yang et al. examined the effects of 7,8-DHF in a rat model of schizophrenia obtained by
57 blocking NMDA receptors through administration of MK-801 during the neonatal period (P7-P10) (Yang et
58 al. 2014b). At 2 months of age these rats were treated with 7,8-DHF for 14 days. Treatment restored LTP in
59

1
2
3 field CA1, learning performance (MWM), hippocampal levels of p-TrkB (Y515 and 816), p-ERK1/2, p-
4
5 CaMKII, p-CREB, and p-GluR1. Since gestational exposure to infection appears to be involved in the
6
7 etiology of schizophrenia, maternal immune activation can be used to obtain animal models of schizophrenia.
8
9 In the first of three studies, Han et al. treated pregnant mice with polyriboinosinic-polyribocytidylic acid
10
11 [poly(I:C)], a potent inducer of immune response (Han et al. 2016). At 4 weeks of age, the offspring was
12
13 treated with 7,8-DHF for 4 weeks and then behaviorally tested. Mice exposed to poly(I:C) exhibited cognitive
14
15 deficits (NOR), PPI deficits of the acoustic startle response, reduced levels of BDNF and p-TrkB (Y706) in
16
17 the prefrontal cortex (PFC) and field CA1, and reduced parvalbumin and peroxisome proliferator-activated
18
19 receptor- γ coactivator 1- α (PGC- α) immunoreactivity in the PFC and CA1. All these defects were
20
21 completely restored in mice treated with 7,8-DHF. In view of the association between increased circulating
22
23 levels of complement protein C1q and schizophrenia, a subsequent study investigated the effect of the same
24
25 treatment schedule on C1q levels in adult offspring (Han, Zhang, and Hashimoto 2017). Treatment with 7,8-
26
27 DHF restored the excessive C1q levels in the PFC, suggesting a positive impact of 7,8-DHF on the immune
28
29 system. A further study was aimed at investigating whether supplementation of 7,8-DHF from pregnancy to
30
31 weaning can prevent the onset of cognitive deficit (Han et al. 2017). Pregnant mice, treated with poly(I:C)
32
33 from E12 to E17, received 7,8-DHF in drinking water from embryonic day E12 to 3-weeks after delivery. An
34
35 evaluation of the effects of 7,8-DHF in offspring aged 12 weeks showed that 7,8-DHF prevented the onset of
36
37 cognitive defects (NOR) and restored the reduced levels of BDNF and p-TrkB (Y706) in the PFC.
38
39

40
41 **Significance.** These studies show that 7,8-DHF improves the cognitive defects associated with schizophrenia
42
43 through activation of TrkB signaling. One explanation for the development of schizophrenia is that the
44
45 disorders begins in utero following infections, obstetric complications or excess stress levels. These
46
47 preclinical studies suggest that supplementation with 7,8-DHF during pregnancy may prevent the subsequent
48
49 transition to schizophrenia in adulthood. Further studies are needed to confirm the preventive effect of 7,8-
50
51 DHF in other models of fetal disturbances and to better characterize the molecular mechanisms whereby 7,8-
52
53 DHF exerts its positive effects.
54

55 56 **4.2.3 Major depressive disorder** 57 58 59 60

1
2
3 **Brief premise.** Major depressive disorder (MDD) represents the most common and debilitating of psychiatric
4 disorder, affecting 17% of the population at some point in their lifetime. Despite the current plethora of
5 antidepressant therapies, approximately two third of patient with depression fail to respond to first-line
6 antidepressant treatments. Depressed patients exhibit low circulating levels of BDNF (Sen, Duman, and
7 Sanacora 2008) and a postmortem study found that patients with a history of major depression who had
8 committed suicide had reduced levels of BDNF and TrkB in the hippocampus and PFC, two brain regions
9 essential for mood regulation and cognitive function (Dwivedi et al. 2003). Accumulating evidence suggests
10 that enhancement of BDNF-TrkB signaling mediates the therapeutic effects of antidepressant drugs and that
11 a BDNF-dependent increase in hippocampal neurogenesis plays a key role in the beneficial effects of
12 antidepressants. Accordingly, ablation of TrkB in hippocampal neural progenitor cells prevented
13 antidepressant-induced neurogenesis and rendered mice behaviorally insensitive to chronic antidepressant
14 treatment (Li et al. 2008). Moreover, in accordance with this hypothesis, intracerebral administration of
15 BDNF has been shown to induce long-lasting antidepressant-like effects in models of depression (Hoshaw,
16 Malberg, and Lucki 2005).

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32 **The evidence.** Based on these lines of evidence, Liu et al. sought to establish whether 7,8-DHF elicits
33 potential antidepressant effects, similarly to BDNF (Liu et al. 2010). In mice treated with 7,8-DHF for 21
34 days, the swimming immobility time during the forced swim test (FST) was significantly decreased,
35 suggesting an antidepressant effect. This effect was associated with an increase in neurogenesis in the
36 dentate gyrus (DG). This study also used the *TrkB*^{F616A} knockin mouse because this mouse harbors a *TrkB*
37 F616A mutation which allows blocking of TrkB receptor kinase activity by the inhibitor 1NMPP1. *TrkB*^{F616A}
38 mice treated with 1NMPP1 one day before administration of 7,8-DHF for 5 days, did not show any reduction
39 in the swimming immobility in the FST, suggesting that the antidepressant effects of 7,8-DHF are mediated
40 by TrkB-dependent signaling. A subsequent study showed that the 7,8-DHF methylated metabolite 7M8H-
41 flavone promoted hippocampal neurogenesis and displayed antidepressant effect (FST and tail suspension
42 test, TST) by activating the TrkB receptor (Liu et al. 2013). Based on evidence that inflammatory processes
43 have a role in the pathophysiology of MDD, a study examined potential antidepressant effects of 7,8-DHF in
44 an inflammation-related mouse model of depression (Zhang et al. 2014a). In this model, systemic
45 administration of the bacterial endotoxin lipopolysaccharide (LPS) induced depression-like behavior
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 resulting from activation of pro-inflammatory cytokine signaling and decreased BDNF and p-TrkB (Y706)
4 levels in CA3, DG and PFC. These mice were injected with 7,8-DHF 23 h after LPS administration. 7,8-
5 DHF restored the LPS-induced reduction in spine density of pyramidal neurons in CA3, DG, and PFC,
6 normalized p-TrkB levels in all these regions and improved depression-like behavior in the FST and TST.
7 The effects of 7,8-DHF on LPS-induced depressive behavior were blocked by pretreatment with the TrkB
8 antagonist ANA-12. The antidepressant effect of 7,8-DHF was also examined in the social defeat stress
9 model of depression (Zhang et al. 2015). Mice that were identified as susceptible to social defeat stress,
10 received a single injection of 7,8-DHF. Treatment elicited antidepressant effects immediately after 7,8-DHF
11 administration (FST and TST). In addition, 7,8-DHF restored the reduced levels of the synaptic markers
12 PSD-95 and GluA1 in CA3, DG, and PFC at 4 but not 8 days after the injection, suggesting that a single dose
13 of 7,8-DHF elicits a rapid but not sustained antidepressant response. The effects of a single bilateral infusion
14 of 7,8-DHF into specific brain regions (medial PFC, CA3 or DG) were investigated in a learned helplessness
15 (LH) rat model of depression (Shirayama et al. 2015). LH rats that received a microinjection of 1.0 pmol 7,8-
16 DHF into the PFC, CA3 or DG underwent an improvement in the conditioned avoidance test, whereas
17 injections of 0.1 pmol did not exert any antidepressant effects, suggesting that the dose of 7,8-DHF is critical
18 for inducing antidepressant effects. Zhang et al. investigated the effects of 7,8-DHF in a chronic mild stress
19 model of depression (Zhang et al. 2016). In this model, depressive-like state develops gradually over time in
20 response to stressful conditions, recapitulating the human condition. Mice were exposed to various mild
21 stresses (e.g. food or water deprivation) before and during treatment with 7,8-DHF for 28 days. 7,8-DHF
22 reverted depressive-like behavior assessed with sucrose preference test (SPT) and novelty suppressed
23 feeding test, restored levels of synaptophysin and PSD95 in the PFC and normalized the reduced levels of
24 BDNF and p-TrkB (Y515). These effects were abolished by pretreatment with the TrkB antagonist K252a.
25 Consistently with the study in mice (Zhang et al. 2016), 7,8-DHF reverted depressive-like behaviors in rats
26 exposed to chronic mild stress for 8 weeks and treated with 7,8-DHF during the last 4 weeks (Chang et al.
27 2016). The therapeutic impact of 7,8-DHF alone or in combination with the antidepressant fluoxetine was
28 investigated in a model of chronic unpredictable mild stress during the perimenopausal period, when the
29 possibility to develop depression increases due to a reduction in levels of the estrogen hormone (Amin et al.
30 2020). Adult C57BL/6J female mice were first subjected to bilateral ovariectomy and then were exposed to a

1
2
3 battery of stressors in combination with drugs administration (7,8-DHF or 7,8-DHF+fluoxetine
4
5 18/mg/kg/day) for 28 days. 7,8-DHF alone or in combination with fluoxetine improved in a similar manner
6
7 depressive-like behavior (FST and SPT). The combined treatment, however, was more effective in reducing
8
9 NF- κ B and inducible nitric oxide synthase (iNOS) levels (markers of brain inflammation), increasing the
10
11 reduced number of neurons and microglia and reverting the downregulation of BDNF/TrkB/ERK1/2 and
12
13 PI3K/Akt/mTOR signaling in comparison with treatment with 7,8-DHF alone.

14
15 **Significance.** These studies show that 7,8-DHF has beneficial effects in different models of depression
16
17 suggesting that it may be considered as a new therapeutic agent for MDD of different etiology and may
18
19 represent an effective treatment for patients resistant to first-line antidepressant treatments. These findings
20
21 additionally indicate that activation of TrkB and its downstream signaling cascade in the hippocampus and
22
23 PFC play a crucial role in mediating the antidepressant effects of 7,8-DHF.

24 25 26 **4.2.4 Anxiety and Stress**

27
28 **Brief premise.** Anxiety and stress disorders are among the most prevalent neuropsychiatric disorders.
29
30 Anxiety disorders may fall along a continuum that ranges from specific fear-based reactivity to more diffuse
31
32 and prolonged stress (Duval, Javanbakht, and Liberzon 2015). Based on the notion that BDNF expression is
33
34 affected by stress, a few studies have explored the effect of 7,8-DHF in anxiety/stress-models.

35
36 **The evidence.** In a first study, Andero et al. (Andero et al. 2011) examined the effects of 7,8-DHF on the
37
38 acquisition and extinction of cue-dependent conditioned fear. Mice were subjected to cued fear conditioning
39
40 (a tone followed by a foot shock) and the expression of fear (mean percentage of time spent freezing) was
41
42 assessed after 24 h. Mice injected with 7,8-DHF 1 h prior to fear conditioning showed increased freezing,
43
44 indicating an enhancement of fear learning. On the next day, mice were given 7,8-DHF 1 h prior to
45
46 extinction training. These mice froze less than did the vehicle group indicating enhancement of extinction. A
47
48 similar treatment-induced enhancement of extinction was also obtained in mice subjected to immobilization
49
50 stress (a putative model of post-traumatic stress disorder) before the cued-fear conditioning protocol. In a
51
52 second study, Andero et al. (Andero et al. 2012) analyzed the effects of 7,8-DHF on spatial memory in rats
53
54 exposed to a traumatic stressor event consisting in immobilization for 2 h. Rats received 7,8-DHF 2 h prior
55
56 or 8 h after termination immobilization and L&M were assessed after three days with the MWM test.
57
58 Treatment with 7,8-DHF either before or after the stressor event prevented the spatial memory deficit caused
59
60

1
2
3 by stress. Choi et al. (Choi, Gourley, and Ressler 2012) used BDNF-floxed mice to knockdown *Bdnf*
4 bilaterally in the prelimbic cortex and examined the outcome in a cue-dependent fear conditioning protocol.
5 While untreated *Bdnf* knockdown mice exhibited a reduced expression of learned fear, mice injected with
6 7,8-DHF 24 h after fear conditioning training expressed fear similarly to the control mice. TrkB^{F616A} mice
7 that received an infusion of the TrkB inhibitor 1NMPP1 in the prelimbic cortex before fear conditioning
8 training failed to show fear learning, confirming the role of TrkB signaling in this region in this behavior.
9 Sanz-Garcia et al. (Sanz-Garcia et al. 2016) examined the effect of 7,8-DHF in rats exposed to an
10 immobilization stress for 2 h. At 2 days after the stressor, rats that had received a single dose of 7,8-DHF 8 h
11 after the stress exhibited an improvement in the stress-induced impairment of spatial memory (MWM) and
12 normalization of the stress-induced increase in LTP in field CA1. Barfield et al. (Barfield et al. 2017)
13 exposed adolescent mice to the stress hormone cortisol from P31 to P47. As adults, these mice exhibited
14 reduced spine density in the medial PFC, increased habit behavior (i.e. lack of goal-directed behavior) in a
15 test of instrumental contingency degradation, and increased immobility in the FST, a typical stress-related
16 depressive behavior. In cortisol-treated mice, co-treatment with 7,8-DHF (from P39 to P47) blocked cortisol-
17 induced habit biases and reduced the immobility time in the FST. In addition, 7,8-DHF blocked the p-ERK42
18 deficit due to cortisol exposure and increased the levels of PSD95 in the ventral hippocampus. The same
19 group studied the effects of 7,8-DHF on behavioral flexibility after adolescence exposure to cortisol
20 (Barfield, and Gourley 2017). Male and female mice treated with cortisol from P31 to P42 were tested at P56
21 with an instrumental reversal learning task, a task commonly used to model cognitive stress related deficits.
22 Cortisol-exposed females, but not males, exhibited perseverative errors. Co-treatment with 7,8-DHF in the
23 period P39-47 eliminated cortisol-induced errors in females and improved the performance in males.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 **Significance.** Although the issue as to whether animal species can really be used as “models” of anxiety and
48 stress, given the highly subjective nature of these emotions, they may help predict the efficacy of
49 pharmacological treatments. These studies suggest that 7,8-DHF may represent a potential treatment for the
50 prevention/improvement of stress-related memory alterations.
51
52
53
54

55 **4.3 Addiction**

56 **4.3.1 Drug abuse**

57
58
59
60

1
2
3 **Brief premise.** Abuse of substances such as cannabis, cocaine, and amphetamine-type stimulants has severe
4
5 effects on brain and behavior. Both reward seeking and inhibitory control (that requires behavioral
6
7 flexibility), contribute to drug abuse vulnerability (Perry et al. 2011). All addictive drugs have in common
8
9 that they enhance dopaminergic reward synaptic function in the nucleus accumbens (Gardner 2011).
10
11 Preclinical and clinical studies have attempted to cure drug addiction by pharmacologically targeting
12
13 addiction-related systems in the brain (Liu, and Li 2018). BDNF plays roles in many types of plasticity
14
15 including drug addiction. Physical exercise, a condition that is known to increase BDNF levels, appears to
16
17 exert some benefit (Lynch et al. 2013), However, it must be underlined that mixed results are reported
18
19 regarding the role of BDNF in drug-seeking behavior in animals, as BDNF can either enhance reinstatement
20
21 or have protective properties, being stage-dependent (Li, and Wolf 2015).
22
23

24 **The evidence.**

25
26 **4.3.1.1 Cocaine.** Choi et al. (Choi, Gourley, and Ressler 2012) trained prelimbic cortex-specific *Bdnf*
27
28 knockdown mice in a cocaine-conditioned place preference paradigm. These mice had attenuated preference
29
30 for the cocaine-associated chamber. In contrast, mice receiving 7,8-DHF immediately post-training
31
32 underwent restoration of cocaine-associated place preference suggesting a role of TrkB signaling in drug
33
34 seeking behavior. A study in rats examined the effect of 7,8-DHF on extinction of drug seeking in a cocaine-
35
36 induced conditioned place preference task (Otis, Fitzgerald, and Mueller 2014). Systemic injections of 7,8-
37
38 DHF or infralimbic infusions of BDNF strengthened extinction of drug seeking whereas blockade of
39
40 infralimbic TrkB receptors with the inhibitor ANA-12 disrupted consolidation of extinction. A recent study
41
42 (DePoy, Allen, and Gourley 2016) sought to establish whether adolescent cocaine exposure has an effect on
43
44 decision-making strategies in adulthood. Since males but not females with a history of escalating self-
45
46 administration of cocaine developed a bias towards habit-based behaviors the rest of the study focused on
47
48 females. In females, low, ‘stable’ cocaine-reinforced response rates during adolescence were associated with
49
50 cocaine-conditioned context preference in adulthood. Administration of 7,8-DHF 60 min before the test
51
52 mitigated context-elicited reward-seeking responses, suggesting a role of TrkB signaling in this behavior.
53
54

55
56 **4.3.1.2 3,4-Methylenedioxypyrovalerone.** A very recent study has examined the effects of 7,8-DHF on
57
58 behavioral sensitization (progressive and enduring increase in the motor stimulant response to drugs) induced
59
60 by MDPV, a popular and potent psychostimulant, or cocaine (Duart-Castells et al. 2019). Mice received

1
2
3 either MDPV or cocaine for 5 days and their locomotor activity was examined during and 10 days after the
4 last drug administration. Both drugs enhanced locomotor activity and induced a higher locomotor response
5 after withdrawal (behavioral sensitization). Pretreatment with 7,8-DHF 30 min prior to MDPV injections
6 blocked MDPV behavioral sensitization and this effect was prevented by co-administration of the TrkB
7 receptor inhibitor ANA-12. In contrast, 7,8-DHF had no effect on cocaine-dependent behavioral
8 sensitization, suggesting involvement of a different signaling pathway.

9
10
11
12
13
14
15 *4.3.1.2 Methamphetamine.* Ren et al. examined the effect of 7,8-DHF on the prepulse inhibition (PPI) deficit
16 of the acoustic startle response caused by methamphetamine (METH) (Ren et al. 2013). While a single dose
17 of METH impaired PPI, administration of 7,8-DHF 30 min before METH improved PPI. This improvement
18 was prevented by co-administration of the TrkB inhibitor ANA-12. In addition, 7,8-DHF attenuated the
19 METH-induced increase in dopamine levels in the striatum, suggesting that 7,8-DHF may improve PPI
20 through inhibition of dopamine release. In a subsequent study, the same group examined the effects of 7,8-
21 DHF on the hyperlocomotion induced by METH (Ren et al. 2014). Pretreatment with 7,8-DHF, reduced the
22 hyperlocomotion induced by a single administration of METH, in a dose-dependent manner. Repeated
23 administration of METH (once daily for 5 days) caused an increase in hyperlocomotion after 7 days
24 (behavioral sensitization) that was attenuated by injection of 7,8-DHF before METH. In addition, 7,8-DHF
25 mitigated the METH-induced reduction in the expression of the dopamine transporter in the striatum and this
26 effect was prevented by the TrkB inhibitor ANA-12. In a third study the same group examined the role of
27 TrkB signaling on METH withdrawal symptoms (Ren et al. 2015). Mice were subjected to repeated
28 administration of METH (once daily for 5 days). During withdrawal (from day 12 to 25) mice received either
29 ANA-12 or 7,8-DHF. Depressive-like behavior (TST and FST) due to METH withdrawal on day 28, was
30 improved by ANA-12 but not by 7,8-DHF suggesting that increased BDNF–TrkB signaling may play a role
31 in the behavioral abnormalities during withdrawal and, consequently, that a BDNF mimetic such as 7,8-DHF
32 is not a suitable treatment for METH-induced depressive-like-behavior.

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53 **Significance.** These studies show that 7,8-DHF may have a positive impact on some (e.g. extinction) but not
54 all addiction-related behaviors, which emphasizes the complexity of the BDNF-dependent mechanisms
55 underlying drug addiction and poses some caveats for the use of 7,8-DHF as pharmacological adjunct for
56 therapies against drug abuse.
57
58
59
60

4.3.2 Alcohol abuse

Brief premise. Alcoholism is a brain disorder associated with compulsive and escalating consumption of alcohol. BDNF, in addition to being implicated in the neurobiology of drug addiction, appears to play a role in alcohol use disorders (Olsen et al. 2019; Pandey 2016). Decreased levels of BDNF in some brain areas, such as the medial PFC, striatum, hippocampus, and amygdala, are associated with excessive alcohol drinking behaviors and correction of the deficits in BDNF levels attenuates these behaviors, suggesting that BDNF is a negative regulator of alcohol consumption (Pandey 2016).

The evidence. Pandey et al. have examined the possible neuroprotective effect of 7,8-DHF on behavioral and neurochemical changes induced by alcohol and high fat diet (Pandey et al. 2020). Results of alcohol-treated rats only will be described here. Rats received alcohol in the drinking water for 12 weeks. Another group received in addition to alcohol 7,8-DHF during the last 4 weeks. The alcohol-induced impairment in hippocampus-dependent L&M (MWM) was improved by co-treatment with 7,8-DHF. In addition, 7,8-DHF restored the alcohol-induced increase in malonildialdeide (MDa) and nitrite hippocampal levels and the reduction in glutathione (GSH) levels, indicating amelioration of oxido-nitrosative stress. Finally, the alcohol-upregulated mRNA levels of NF- κ B, iNOS, and caspase-3 (indicative of apoptosis) and downregulated levels of nuclear factor erythroid 2-related factor 2 (Nrf2), heme oxygenase 1 (HO-1; indicative of cellular resistance to oxidants) were normalized by 7,8-DHF. Hogarth et al. have explored the possibility that activation of the TrkB receptor *via* 7,8-DHF may modify voluntary alcohol intake reinstatement (Hogarth, Djouma, and van den Buuse 2020). Sprague Dawley rats were trained for about 7 weeks to self-administer through lever presses a 10% ethanol solution (acquisition). During extinction (removal of ethanol reward) animals were treated for two weeks with 7,8-DHF. Following extinction, both male and female rats exhibited alcohol-primed reinstatement (alcohol consumption). While this effect was not altered in male rats by pretreatment with 7,8-DHF, in female rats reinstatement was significantly greater following 7,8-DHF treatment compared to vehicle. Thus, in contrast with the hypothesis of this work, i.e. that 7,8-DHF-mediated TrkB activation would reduce alcohol reinstatement, 7,8-DHF had an opposite effect, at least in females.

Significance. These studies show that while 7,8-DHF may be of benefit for the protection of the brain from the anatomical and functional damages caused by alcohol, it appears to have a negative effect, at least

1
2
3 regarding reinstatement of alcohol abuse in females. Further studies are necessary in order to delineate the
4 potential benefits of 7,8-DHF as treatment for alcohol use disorders.
5
6

7 **4.4 Brain injury**

8 **4.4.1 Traumatic brain injury**

9
10
11 **Brief premise.** Traumatic brain injury (TBI) results from a violent blow to the head or when an object pierces
12 the skull and enters the brain. The pathogenesis of TBI results from primary and secondary injuries that
13 induce temporary or permanent neurological deficits (Galgano et al. 2017; Prins et al. 2013). The
14 primary injury is due to the mechanical impact of the brain. The secondary injury consists in
15 neurochemical alterations that include changes in glucose cerebral metabolism, energy crisis
16 with mitochondrial impairments, accumulation of free radicals, and inflammation (Galgano et al.
17 2017; Prins et al. 2013). Numerous studies conducted in animal models of TBI sought to discover
18 treatments that prevent further brain damage due to the secondary injury and promote functional recovery
19 (Galgano et al. 2017). Commonly used model of TBI are the Fluid Percussion Injury (FPI) model based
20 on hydraulically induced pressure transients on the brain and the Controlled Cortical Impact Model (CCI)
21 that uses a rigid impactor to deliver mechanical energy. Various studies, reviewed below, have
22 examined the potential benefit of 7,8-DHF in these models.
23
24
25
26
27
28
29
30
31
32
33
34
35

36
37 **The evidence.** Agrawal et al. examined the effects of 7,8-DHF on memory retention and brain energy
38 homeostasis in a rat model of FPI (Agrawal et al. 2015). Rats were first trained (learning) on the Barnes
39 maze test for 5 days and then subjected to FPI. In the following 7 days rats received 7,8-DHF or vehicle and
40 then were tested by the Barnes maze for memory retention, their brains were then removed and used for
41 various analyses. In rats treated with 7,8-DHF the memory impairment caused by TBI was fully rescued.
42 Administration of the TrkB receptor inhibitor K252a prior to TBI prevented the memory rescue. Treatment
43 with 7,8-DHF improved the TBI-induced reduction in the hippocampal expression of p-TrkB (Y816),
44 counteracted the reduction of molecules associated with brain plasticity as p-CREB, growth associated-
45 protein 43 (GAP-43), and syntaxin-3. In addition, 7,8-DHF normalized the levels of molecules involved in
46 mitochondrial biogenesis as PGC-1 α , mitochondrial transcription factor 1 (TFAM), mitochondrial
47 cytochrome oxidase II (COII) and brain energy metabolism as adenosine monophosphate-activated protein
48 kinase (AMPK) and sirtuin 1 (SIRT1). A second study from the same group (Krishna et al. 2017) used a
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 substantially similar protocol but examined the effect of exercise and treatment with 7,8-DHF alone or in
4 combination. Sprague-Dawley rats were subjected to FPI after a training of 5 days on the Barnes maze
5 (learning). Thereafter rats received 7,8-DHF with or without voluntary wheel running for 7 days. The TBI-
6 induced memory impairment and reduction in hippocampal p-TrkB levels were increased by treatment with
7 either 7,8-DHF or exercise but were further increased by the combined application of 7,8-DHF and exercise.
8
9

10
11 Likewise, the reduction in the levels of p-CREB, GAP-43, syntaxin-1, p-CaMKII, and AMPK was
12 restored by treatment with either 7,8-DHF or exercise and this effect was enhanced by their combination.
13
14

15
16 Wu et. al. (Wu et al. 2014) examined the effects of 7,8-DHF in a mouse CCI model. Mice received 7,8-DHF
17 10 min after TBI injury and in the following 3 days. Treatment caused an improvement in various
18 neurological tests, reduced the ipsilateral contusion volume, the number of degenerating neurons and
19 apoptotic cells, the expression levels of apoptotic markers and counteracted the TBI-induced reduction in p-
20 TrkB (Y705) and p-Akt, with no effect on p-ERK1/2 (suggesting activation of TrkB and downstream PI3K/
21 Akt signaling). Intracerebroventricular administration of the TrkB receptor inhibitor K252a and the PI3K
22 inhibitor LY294002 before TBI abolished the reduction in contusion volume induced by 7,8-DHF. Chen et
23 al. (Chen et al. 2015) examined the effect of 7,8-DHF on the survival of hippocampal immature neurons in a
24 mouse CCI model. Mice received 7,8-DHF 1 h prior to TBI injury and their brain was collected after 24 h.
25
26

27
28 Treatment with 7,8-DHF reduced the number of dying neurons, increased the number of new granule
29 neurons in the DG, and increased hippocampal levels of p-TrkB (Y816). Treatment with the TrkB inhibitor
30 ANA12 before 7,8-DHF administration prevented the positive effects of 7,8-DHF on the number of dying
31 neurons and of new granule neurons. Zaho et al. (Zhao et al. 2016b) examined the effect of 7,8-DHF on
32 neurogenesis and dendritogenesis in the DG of a CCI mouse model. Mice received 7,8-DHF 1 h after TBI
33 and in the following two weeks. Treated mice showed an increase in the number of new granule neurons,
34 and an improvement in length and number of dendritic branches of the granule neurons. In a subsequent
35 study (Zhao et al. 2016a), the same group analyzed the effect of 7,8-DHF on neocortical neurons of layer II/
36 III in a CCI mouse model. Mice received 7,8-DHF 1 h after TBI and in the following 3 days. Treatment
37 with 7,8-DHF did not ameliorate the TBI-induced cortical lesion. Treatment however, reduced dendritic
38 swelling (indicative of degeneration) of layer II/III neurons, restored the number and length of apical and
39 basal dendritic branches and the density of dendritic spines. Treated mice showed no improvement in L&M
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 (MWM) and a partial improvement in the rotarod test, indicating that the protective effects of 7,8-DHF on
4
5 cortical neurons did not translate into a behavioral improvement.
6

7 **Significance.** Experimental studies in animal models of TBI have not been successfully translated into
8
9 clinical therapies so far. The results obtained following treatment with 7,8-DHF in rat and mouse models of
10
11 TBI show that treatment may be beneficial in terms of brain plasticity, energy homeostasis, protection from
12
13 apoptosis and dendritic degeneration although its effects on behavior are very limited. It must be underlined
14
15 that these studies used a relatively short period of treatment with 7,8-DHF (maximum 14 days). It remains to
16
17 be established whether optimization of the treatment schedule (duration and doses) may have a larger impact
18
19 on the damages caused by TBI.
20

21 22 **4.4.2 Stroke/Ischemia**

23
24 **Brief premise.** Cerebral ischemia consists in a reduction in the flow to the brain causing reduction in the
25
26 supply of glucose and oxygen. Ischemic stroke is due to the block of brain blood vessels by a clot formation,
27
28 and hemorrhagic stroke is caused by the rupture of blood vessels and blood leakage. Stroke causes oxidative
29
30 stress, excitotoxicity, inflammation and apoptosis. Various models of perinatal and adult ischemic stroke
31
32 have been developed (Northington 2006; Sommer 2017). A widely used animal model of stroke consists in
33
34 the permanent occlusion of the middle cerebral artery (MCAO). Another model is obtained with transient
35
36 occlusion (ischemia) followed by reperfusion (I/R). In this model, the absence of oxygen during ischemia
37
38 favors inflammation and oxidative stress when circulation is restored (reoxygenation injury). Another model
39
40 that mimics neonatal hypoxia-ischemia (HI) is obtained by exposing pups to hypoxia after ligation of the
41
42 common carotid artery which results in reproducible brain injury ipsilateral to the carotid ligation. Numerous
43
44 preclinical studies in animal models of stroke have shown protective effects of various therapies, although
45
46 most of the clinical trials do not have shown any success so far (Kaur, Prakash, and Medhi 2013).
47

48
49 **The evidence.** The first evidence regarding the effect of 7,8-DHF in stroke was obtained by Jang et al. (Jang
50
51 et al. 2010) in the TrkB^{F616A} knockin mouse. Mice were injected with 7,8-DHF 2 h before occlusion of the
52
53 middle cerebral artery and their brains were removed after 2 days. Pretreatment with 7,8-DHF largely
54
55 reduced the infarct volume and this effect was prevented in mice that had received INMPP1 1 day before
56
57 MCAO. Based on the notion that perinatal ischemia in preterm infants mainly targets the oligodendrocyte
58
59 lineage, Hung et al. (Hung et al. 2013) explored in a model of HI the effects of thyroxin (T4), a key hormone
60

1
2
3 for oligodendrocyte maturation and that is known to upregulate BDNF levels, and of 7,8-DHF. Rat pups
4 underwent HI at P7 and received on P7, P9, and P11 either T4 or 7,8-DHF. T4 attenuated axonal injury,
5 astrocytosis, and microgliosis, and increased preoligodendrocyte survival. 7,8-DHF, similarly to T4,
6 improved the hypomyelination caused by ischemic injury. In the study by Uluc et al. (Uluc et al. 2013) mice
7 pups subjected to HI received 7,8-DHF at 10 min after HI and for the following 1 or 6 days. At 72 h after HI
8 mice exhibited an increase in the number of degenerating neurons in hippocampal field CA1. While
9 treatment with 7,8-DHF had no effect in males, it reduced the number of degenerating cells in females. In
10 both sexes, 7,8-DHF enhanced hippocampal p-TrkB levels but this increase was more pronounced in
11 females. At 21 days after HI, motor learning was improved by 7,8-DHF in males and females in a similar
12 manner. At 51 days after HI, spatial learning (MWM) was improved by treatment with 7,8-DHF in females
13 but not in males. The study by Cikla et al. (Cikla et al. 2016) was prompted by evidence that the brains of
14 male pups are more susceptible to perinatal hypoxia and ischemia. Pups underwent permanent occlusion of
15 the left carotid, subjected to hypoxia and were treated with 7,8-DHF at 10 min after HI and in the following
16 2 days. While no sex differences were found in p-TrkB (Y705) levels in the hippocampus of sham mice, at 3
17 days post HI there was a higher hippocampal TrkB phosphorylation in 7,8-DHF-treated females compared
18 with males, suggesting that this difference may account for the larger neuroprotection observed in females.
19 Interestingly, following HI there was an increase in hippocampal ER α expression that was more pronounced
20 in females. Estrogen receptor α (ER α) is coupled to Src family kinase (SFK) activation which, in turn,
21 enhances the activation of the TrkB receptor, in part by direct phosphorylation of TrkB (Y705/Y706) and in
22 response to ligands such as 7,8-DHF. Thus, the sexually differentiated phosphorylation of TrkB following HI
23 and treatment with the TrkB agonist 7,8-DHF appears to be due to sex-related differences in the expression
24 of ER α . In a study by Wang et al. (Wang et al. 2014) rats were subjected to MCAO for 90 min followed by
25 reperfusion (ischemia/reperfusion, I/R) for 24 h and received 7,8-DHF immediately after MCAO. Treatment
26 with 7,8-DHF improved the I/R-induced neurological deficits and the infarct volume, counteracted the
27 increase in the cortical expression levels of cleaved caspase-3 and Bcl-2 homologous antagonist/killer (Bax)
28 and the reduction in Bcl-2 levels, reduced oxidative stress, evaluated based on the content of MDA and the
29 activity of antioxidant enzymes as GSH, glutathione peroxidase (GSH-PX), superoxide dismutase (SOD).
30 Finally, 7,8-DHF inhibited nitric oxide (NO) production and iNOS expression, and suppressed the NF-kB
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 pathway, indicating reduction of inflammation. These effects were accompanied by an increase in p-
4 TrkB levels. In an *in vitro* study, Tecuatl et al. (Tecuatl et al. 2018) investigated the effect of 7,8-DHF
5 on the ischemic damage in hippocampal fields CA1 and CA3. Rat hippocampal slices were exposed to an
6 Oxygen-Glucose Deprivation (OGD) solution for 10 min and then re-perfused with normal artificial
7 cerebrospinal fluid. OGD caused a permanent suppression of CA1 but not CA3 field potentials. The basal
8 levels of p-TrkB (Y816) were higher in CA3 than in CA1 and OGD caused a larger increase in p-TrkB in
9 CA3 in comparison with CA1. Pre-incubation with 7,8-DHF (or BDNF) enhanced the recovery of CA3
10 but not CA1 response after OGD, suggesting that the TrkB receptor plays a central role in the different
11 vulnerability of CA3 and CA1.

12
13 **Significance.** These studies show that 7,8-DHF can protect against cerebral I/R injury by reducing
14 apoptotic cell death, neuronal degeneration, oxidative stress and neuroinflammation. These effects
15 translate into amelioration of hippocampal evoked potentials and behavior. Similarly to models of
16 addiction, treatment with 7,8-DHF appears to exert sexually dimorphic effects in models of ischemia.

17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Brief premise. Brain radiation, used to treat brain tumors, may be associated with brain injury and
cognitive impairment that can occur in up to 50-90% of patients (Greene-Schloesser et al. 2012).
Preclinical studies in rodents have shown that irradiation damages hippocampal neurogenesis and causes
neuroinflammation and cognitive impairment (Greene-Schloesser et al. 2012), suggesting that
treatments that counteracts these effects may ameliorate radiation-induced brain injury.

The evidence. In the study by Yang et al. (Yang et al. 2016), mice received 5 Gy of cranial irradiation
and after one week were treated with 7,8-DHF for 3 or 5 weeks. At 4 weeks after completion of
7,8-DHF administration mice showed restoration of spatial (Barnes Maze), contextual (CFC), and working
memory (Y maze), of hippocampal neurogenesis and granule cell dendritic spine loss. Treatment, in
addition, increased the levels of p-TrkB (Y816) and its downstream targets ERK and Akt.

Significance. This study shows that 7,8-DHF exerts remarkably positive effects on the brain subjected
radiation. These effects span from restoration of hippocampal neurogenesis to prevention of dendritic and
dendritic spine regression. Importantly, thanks to these neuroanatomical effects, 7,8-DHF restores cognitive
performance.

4.5 Neurodegeneration

4.5.1 Alzheimer's disease

Brief premise. Alzheimer's disease (AD) is the leading cause of dementia in individuals over 65. This progressive neurodegenerative disease is characterized by brain extracellular neuritic plaques of amyloid beta ($A\beta$) and intracellular neurofibrillary tangles consisting of hyperphosphorylated microtubule-associated Tau protein, resulting in synaptic deficit and neuronal death at advanced stage of the disease. In addition to these main neuropathological hallmarks, the levels of BDNF and TrkB receptor are reduced in the hippocampus and some cortical areas of AD patients, even at the early AD stages (Ferrer et al. 1999; Zhang et al. 2012), suggesting that impairment of BDNF-TrkB signaling may contribute to the pathogenesis of AD.

Accordingly, central BDNF administration or gene delivery to AD animal models exert beneficial effects, including restoration of L&M (Nagahara et al. 2009). Based on evidence of the efficacy of BDNF against AD pathology, 7,8-DHF was tested in the preclinical field of AD in order to overcome the problems related to the delivery and scarce half-life of BDNF *in vivo*.

The evidence. A first study investigated the effects of 7,8-DHF in the 5XFAD mouse model of AD (Devi, and Ohno 2012). This model develops cerebral amyloid plaques at 2 months of age and shows synaptic dysfunction and memory impairment by 4-5 months of age. Mice aged 12-15 months received 7,8-DHF for 14 days. Treatment completely restored the reduced hippocampal levels both of the phosphorylated (Y705) and total form of TrkB receptor, without affecting the reduced BDNF levels. The activation of TrkB receptor was accompanied by rescue of spatial working memory (Y maze) and by reduction of the excessive levels of (β -secretase) BACE1, β -CTF (β -secretase-cleaved C-terminal fragment), and of $A\beta$ 40 and $A\beta$ 42. A subsequent study in the 5XFAD model tested the effects of 7,8-DHF at an earlier stage of disease progression (Zhang et al. 2014b). Mice received 7,8-DHF for 4 months starting from 2 months of age, at the beginning of plaque deposition. Administration of 7,8-DHF reversed the loss of dendritic spines and synapses in the hippocampal field CA1 and restored of hippocampal LTP and hippocampus-dependent L&M (MWM). At the molecular levels, 7,8-DHF elicited robust brain TrkB phosphorylation (Y816) and also increased p-Akt and p-ERK1/2 levels. Treatment also reduced the density of hippocampal $A\beta$ plaques but, in contrast with the previous study, did not affect total $A\beta$ 42 levels. Exploration of the effects of 7,8-DHF *in vitro* showed that 7,8-DHF (500 nM) promoted dendritic arborization of primary cortical neurons and

1
2
3 reduced the apoptotic rate induced by pre-aggregated A β , both in cortical and locus coeruleus neurons. The
4 anti-apoptotic effect was abolished by the TrkB receptor inhibitor K252. In a further study, one-month-old
5 5XFAD mice were treated with 7,8-DHF for 2 months in order to establish whether 7,8-DHF can prevent the
6 AD-related phenotypes (Aytan et al. 2018). 5XFAD mice showed, similarly to individuals with AD, a
7 reduction in hippocampal glutamate levels and an increase in choline and phosphocholine levels. These
8 alterations were restored by treatment. In addition, treatment prevented the pruning of the dendritic tree of
9 cortical neurons of layer V although it had a very modest effect on spine density. Finally, 7,8-DHF reduced the
10 A β plaque burden and A β 42 levels, although this latter effect was not statistically significant. Chen et al.
11 investigated the effects of R13, a prodrug of 7,8-DHF, in 2-month-old 5XFAD mice (Chen et al. 2018). All
12 tested doses of R13 prevented the loss of dendritic spines and synapses in field CA1, improved LTP, and
13 restored L&M (MWM). These positive effects were associated with increased hippocampal levels of p-TrkB
14 (Y816), p-Akt, and p-ERK. In addition, treatment reduced amyloid plaques in the hippocampus and frontal
15 cortex and the levels of A β 40, but not A β 42. Furthermore, R13 reduced the enzymatic activity of delta-
16 secretase (AEP; asparagine endopeptidase) with a consequently reduction in the levels of APP/Tau
17 pathological fragments (APP N373, APP N585, and Tau N368) and pro-inflammatory cytokines (IL-1 β , IL-6,
18 and TNF- α). Gao et al. (Gao et al. 2016) investigated the effects of 7,8-DHF in the Tg2576 mouse, one of the
19 most widely used AD models that overexpresses the human APP with the Swedish mutation, resulting in
20 elevated A β levels and amyloid plaques. In 10-month-old Tg2576 mice a single injection of 7,8-DHF was
21 ineffective on field CA1 spine loss and L&M impairment (MWM). Conversely, chronic administration of
22 7,8-DHF for 4 weeks fully rescued spine density of CA1 pyramidal neurons and L&M. In addition, in
23 hippocampal synaptoneurosomal fractions, chronic treatment increased the reduced protein levels of both
24 GluA1 and GluA2, two subunits of the glutamate AMPA receptor. This effect was mediated by activation of
25 TrkB (Y816) and its downstream PI3K/Akt, Ras/ERK, and PLC γ /CaMKII signaling pathways because it was
26 prevented by the TrkB inhibitor cyclotraxin B, U0126 (a Ras-ERK pathway inhibitor), Wortmannin (an Akt
27 phosphorylation inhibitor), and KN-93 (a CaMKII inhibitor). No treatment effects, however, were found in
28 the excessive hippocampal levels of APP and A β 42. The effect of a single injection of 7,8-DHF on spatial
29 memory (object location memory test) was investigated in the APP^{swe}/PS1^{dE9} mouse model of AD at 7
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 months of age, when hippocampal functional deficits start to be present (Bollen et al. 2013). Mice that
4 received 7,8-DHF 3 h after the first exploration trial of the test underwent an improvement in their
5 discrimination performance following a retention period of 24 h, suggesting an acute effect of 7,8-DHF on
6 spatial memory. An improvement of memory (CFC test) was found in APP^{swe}/PS1^{dE9} male mice aged 9
7 months, following chronic administration of 7,8-DHF for 10 days (Hsiao et al. 2014). The effects of 7,8-
8 DHF were also examined in the CaM/Tet-DT_A mouse, an inducible model of neuronal loss in field CA1 (but not
9 CA3), DG and entorhinal cortex that replicates the neurodegenerative feature of the AD brain (Castello et al.
10 2014). Administration of 7,8-DHF to 3.5-5-month-old mice for 2 weeks increased the density of thin
11 immature dendritic spines of CA1 pyramidal neurons, with no effect on the mushroom and stubby spines, and
12 caused an improvement in L&M performance (MWM). In contrast with all above studies, 7,8-DHF failed
13 to affect AD-related phenotypes in the APP23/PS45 mouse model of AD (Zhou et al. 2015). Treatment
14 with 7,8-DHF for 4 weeks starting from 6 weeks of age had no effect on the levels of APP, CTFs, BACE1, A β
15 42 and A β 40, and A β plaque formation. Furthermore, 7,8-DHF did not improve L&M deficits (MWM).
16 Positive effects of treatment with 7,8-DHF on AD-like pathological dysfunction were observed in rats treated
17 with scopolamine, a non-selective muscarinic receptor antagonist that, by blocking cholinergic signaling,
18 induces A β deposition, oxidative stress, synaptic dysfunction and L&M impairment (Chen et al. 2014).
19 Administration of 7,8-DHF restored LTP in the DG and decreased the scopolamine-induced excessive levels of
20 hippocampal A β 42 and A β 40. In addition, 7,8-DHF reduced hippocampal lipid peroxidation, caused an
21 increase in MDA levels, a reduction in GSH levels and in the activity of SOD, revealing an antioxidant
22 effect. These effects were accompanied by restoration of L&M (MWM) and hippocampal levels of p-TrkB
23 (Y516) and its downstream molecules p-Akt and p-ERK1.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 **Significance.** This evidence in different AD models suggests that 7,8-DHF may represent a promising
49 therapeutic strategy not only for the prevention/delay of AD but also for the amelioration of
50 pathophysiological processes associated with the late stage of disease progression. Further studies are needed in
51 order to establish the optimum effective dose of 7,8-DHF and the duration of its effects. While the positive effects
52 of 7,8-DHF on synaptic dysfunction and memory impairments seem to be directly mediated by the activation
53 of TrkB and its downstream signaling pathways, the molecular mechanisms underlying the effects of 7,8-DHF
54 on APP processing and A β aggregation remain to be elucidated.
55
56
57
58
59
60

4.5.2 Parkinson's disease

Brief premise. Parkinson's disease (PD), which is the second-most common neurodegenerative disorder, is characterized by loss of dopaminergic neurons in the substantia nigra of the midbrain - possibly due to intracellular overexpression of α -synuclein and/or oxidative injury - and diminished content of dopamine in the striatum. These alterations cause impairment of dopamine-regulated motor behavior (bradykinesia, postural instability, rigidity and tremor) (Poewe et al. 2017). Although various preclinical studies have shown that the delivery of neurotrophic factors, including BDNF, to the brain protects dopaminergic neurons and alleviates PD symptoms, the problem of the delivery of trophic factors in humans remains a challenge (Tome et al. 2017).

The evidence. The first evidence that 7,8-DHF may be of benefit for PD derives from a model of PD obtained by administering the dopaminergic toxicant MPTP (Jang et al. 2010). Eight-week-old mice were treated with 7,8-DHF for 14 days. On day 7, mice were treated with MPTP and were killed on day 14. 7,8-DHF reduced the neurotoxic effects of MPTP, as revealed by preservation of the expression of tyrosine hydroxylase (a marker of dopaminergic neurons) in the striatum and reduced the levels of activated caspase-3 in the substantia nigra. In a study by Sconce et al. (Sconce et al. 2015), mice received 7,8-DHF starting at midpoint (week 3) of a 4-week-MTPT administration and for an additional 2 weeks. Mice treated with 7,8-DHF maintained about 50% of tyrosine hydroxylase levels in the dorsolateral striatum, with no effect, however, in the substantia nigra underwent restoration of motor deficits, as assessed through gait dynamics. In addition, treated mice underwent an increase in the levels of p-TrkB (Y817), and p-ERK1/2 (but not of p-Akt) in the dorsolateral striatum and substantia nigra compared to untreated mice. In a third study in the MPTP model (Li et al. 2016), mice received 7,8-DHF from the first day of a 5-day-long MPTP treatment until 9 days after discontinuation of MPTP treatment. Mice that were treated MTPT alone were impaired in motor performance (rotarod test, pole test and wire suspensions test), exhibited loss of dopaminergic neurons in the substantia nigra and of dopaminergic terminals in the striatum, increased levels of α -synuclein, and reduced levels of p-TrkB in the substantia nigra and striatum. All these defects were improved/restored in mice that had received 7,8-DHF. In addition, 7,8-DHF restored the levels of the endogenous antioxidants GSH and SOD in the substantia nigra and striatum. The protective effects of 7,8-DHF have been also examined in a rotenone-induced animal model of PD (Nie et al. 2019). Rats were injected with rotenone

1
2
3 daily for 5 weeks in order to induce a PD phenotype. Administration of 7,8-DHF during treatment with
4 rotenone improved behavioral performance (distance travelled and speed in an open field test), reduced the
5 loss of dopaminergic neurons in the substantia nigra and of dopaminergic terminals in the striatum, and
6 reduced the rotenone-induced increase in p-MAPK, p-Tau and p- α -synuclein. The effects of 7,8-DHF were
7 associated with increased levels of p-TrkB (Y816) and p-Akt. The study by He et al. (He et al. 2016)
8 examined the effect of 7,8-DHF in a primate model (Macaca Fascicularis; 8-year-old) of PD obtained with
9 intracerebroventricular injection of MPP+, the toxic metabolite of MPTP, delivered over a period of seven
10 months. 7,8-DHF was administered 2 weeks earlier than MPP+ injections and during the following 7
11 months. An evaluation of dopaminergic neurons in the substantia nigra showed loss of neurons in the
12 monkey that had not received 7,8-DHF. In contrast, 7,8-DHF-treated monkeys had a similar number of
13 neurons as wild-type monkey indicating a powerful neuroprotective effect of treatment. No effect of either
14 MPP+ or 7,8-DHF were found on astrogliosis, microglia activation and neurogenesis in the substantia nigra.

15
16
17
18
19
20
21
22
23
24
25
26
27
28 **Significance.** Taken together, these studies show that 7,8-DHF has a positive impact in different models of
29 PD through TrkB-dependent mechanisms. The differences in the activated pathways downstream of the
30 TrkB receptor may be due to differences in the employed model of PD and/or duration of treatment.
31 Whatever the mechanism, all these studies show that 7,8-DHF has beneficial effects in PD models,
32 suggesting that it may represent a possible treatment for alleviating the symptoms of PD.
33
34
35
36
37
38

39 **4.5.3 Huntington's disease**

40
41 **Brief premise.** Huntington's disease (HD) is a neurodegenerative disorder caused by an expanded CAG
42 repeat in the huntingtin gene. This mutation causes widespread neurodegeneration, with massive loss of the
43 medium spiny neurons in the striatum, motor impairment and cognitive decline (Vonsattel, and DiFiglia
44 1998). It has been suggested that a reduction in brain BDNF levels due to mutant huntingtin underlies
45 neurodegeneration in HD (Zuccato, and Cattaneo 2007). The reduced BDNF levels are paralleled by a
46 reduction in TrkB receptor. In the R6/1 mouse model of HD, administration of exogenous BDNF can
47 improve the HD-related neurological phenotype, indicating that the remaining TrkB receptors are sufficient
48 to mediate BDNF-dependent signaling (Zuccato, and Cattaneo 2007). Accordingly, various studies have
49 suggested the potential benefits of BDNF in HD which, in turn, has prompted studies that have exploited the
50 BDNF mimetic 7,8-DHF.
51
52
53
54
55
56
57
58
59
60

1
2
3 **The evidence.** A study in the HD mouse model N171-82Q has examined the effect of 7,8-DHF on HD-
4 related phenotypes (Jiang et al. 2013). Mice received 7,8-DHF from 6 weeks of age until they were 20-week-
5 old. Treated mice displayed an improvement in motor function (balance beam), an increase in their lifespan,
6 and attenuation of brain atrophy (MRI scans *in vivo*), and an increase in p-TrkB (Y706) and p-MAPK levels. A
7 study in the mouse R6/1 model of HD examined the effects of 7,8-DHF *in vitro* and *in vivo* (Garcia-Diaz
8 Barriga et al. 2017). In cultures of striatal neurons from R6/1 mice, 7,8-DHF was found to phosphorylate the
9 TrkB receptor at Y816 (but not Y515), leading to activation of PLC γ 1 downstream signaling. These effects
10 were associated with an increase in the number and length of neurites and improvement of neuronal
11 excitability. R6/1 mice chronically treated with 7,8-DHF, starting at pre-symptomatic stages (8 weeks) and
12 continuing up to 20 weeks of age, partially improved motor performance in the rotarod test, completely
13 reversed the deficit in the NOR test, partially recovered the soma size of the medium spiny neurons, and
14 underwent a reduction in mutant huntingtin aggregation in the striatum. At the molecular level, *in vivo*
15 treatment with 7,8-DHF reverted the reduction in p-TrkB (Y816) and increased the levels of p-PLC γ 1. In
16 addition, treatment restored the reduced levels of neuronal nitric oxide synthase (nNOS) and reduced the
17 excessive levels of iNOS, indicating an effect on NO metabolism and, thus, on neuroinflammation. Finally,
18 in the Hdh(Q7) striatal immortalized cell line exposed to H₂O₂, 7,8-DHF was able to prevent cell death in the
19 presence of the TrkB inhibitor K252a, indicating that 7,8-DHF protects against oxidative stress with a
20 mechanism independent from TrkB activation.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40
41 **Significance.** These studies show that 7,8-DHF has a positive impact in models of HD through TrkB-
42 dependent and independent mechanisms. Although 7,8-DHF does not restore the phenotype of HD, it may
43 have a therapeutic potential for the amelioration of the motor and cognitive deficits in this pathology.
44
45
46

47 4.5.4 Amyotrophic lateral sclerosis

48
49 **Brief premise.** Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disorder characterized by
50 degeneration of motor neurons, progressive motor weakness and, ultimately, death due to respiratory failure.
51 Various neurotrophic factors have been investigated pre-clinically but with disappointing results in clinical
52 trials (Henriques, Pitzer, and Schneider 2010).
53
54
55
56

57
58 **The evidence.** The possible neuroprotective effect of 7,8-DHF has been investigated in the SOD1^{G93A} mouse, a
59 transgenic ALS mouse model (Korkmaz et al. 2014). Mice aged 30 days received 7,8-DHF until 105 days
60

1
2
3 of age. Treatment prevented the motor decline (rotarod test) that started to appear at 60 days of age.
4
5 An evaluation of the number of motor neurons in the ventral horn of the spinal cord and of dendritic
6
7 spine density at the end of treatment showed that untreated SOD1^{G93A} mice had a reduced number of
8
9 motor neurons and a reduced dendritic spine density. Treatment prevented neuron loss and fully restored
10
11 dendritic spine density.
12

13
14 **Significance.** Since the only available study in a model of ALS did not examine the TrkB receptor and
15
16 its downstream signaling, it remains to be established whether the beneficial effects of 7,8-DHF were
17
18 mediated by the TrkB receptor or rather by antioxidant actions. Nevertheless, this study shows that 7,8-
19
20 DHF may be a possible tool to mitigate the outcome of ALS.
21

22 **4.6 Actions of 7,8-DHF in the healthy brain**

23
24 **Brief premise.** The studies mentioned above were aimed at establishing the therapeutic potential of 7,8-
25
26 DHF in a variety of brain diseases. A few studies show that 7,8-DHF also improves the anatomy and
27
28 function in the healthy brain.
29

30
31 **The evidence.** Zimmerman et al. (Zimmermann et al. 2017) examined the effect of 7,8-DHF on goal
32
33 directed action selection in normal mice using a test of instrumental contingency degradation. Mice are
34
35 trained to generate two food-reinforced responses, then the likelihood that one response will be reinforced
36
37 is reduced (action-outcome contingency degradation). During a subsequent probe test, preferential
38
39 engagement of the response that is likely to be reinforced is considered “goal-directed”. While
40
41 control mice failed to differentiate between the responses that were likely to be reinforced, mice
42
43 injected with 7,8-DHF immediately following the training session exhibited an increase in the preference
44
45 for the response likely to be reinforced, indicating that 7,8-DHF enhances action-selection. In addition,
46
47 treatment increased dendritic spine density on neurons in the orbitofrontal cortex. Bollen et al.
48
49 examined the effects of 7,8-DHF on memory consolidation in normal healthy rats (Bollen et al. 2013).
50
51 Rats were subjected to a NOR test and the effects of different doses of 7,8-DHF administered either
52
53 immediately after or 3 h after the learning trial carried out after 24 h were tested. Animals that had
54
55 received 7,8-DHF exhibited a better discrimination index (time of exploration of the new object in
56
57 comparison with the old object) in comparison with vehicle treated rats. Perez-Rando et al. (Perez-Rando et
58
59 al. 2018) examined the effects of 7,8-DHF in Thy1-YFP transgenic mice, a strain in which layer V
60
cortical pyramidal neurons are labeled with the fluorescent protein YFP,

1
2
3 which allows observation of dendritic spine dynamics *in vivo* with 2-photon microscopy. Mice received 7,8-
4 DHF for 14 days and were behaviorally examined before and at the end of treatment. 7,8-DHF did not
5 modify locomotion or anxiety related behavior (Open Field Test) but improved object recognition (modified
6 version of the NOR) in comparison with the control group. Regarding spine dynamics, treatment improved the
7 gain rate, loss rate, and stability rate of spines of neocortical pyramidal neurons, providing the first *in vivo*
8 evidence of how 7,8-DHF modifies spine dynamics in healthy animals. Even in the absence of specific
9 pathologies, memory undergoes a “physiological” decline with aging. The study by Zeng et al. (Zeng et al.
10 2012b) was aimed to establish whether treatment with 7,8-DHF has a protective effect on age-related
11 memory decline. This study used aged rats (20- or 30-month-old) that revealed memory impairment in
12 comparison with young rats (assessed by pretesting them with the MWM test). These rats were treated with
13 7,8-DHF for 34 days and tested with the MWM 10 days after beginning of treatment and in the last week of
14 treatment. Treated rats of either age showed full restoration of L&M, although the older group showed a
15 restoration in the second retest only. An evaluation of LTP in hippocampal slices from some of the
16 behaviorally tested rats showed that LTP in field CA1 was restored by 7,8-DHF in the 20-month group and
17 improved in the 30-month group. Likewise, dendritic spine density of CA1 pyramidal neurons was restored by
18 7,8-DHF in the 20-month group and improved in the 30-month group. In both age groups, treatment with 7,8-
19 DHF increased the age-related reduction in the hippocampal levels of p-TrkB (Y816), p-ERK1/2, p-CREB,
20 p-CaMKII, and p-GluR1. In a subsequent study (Zeng et al. 2012a) the same group demonstrated that 25-month-
21 old rats exhibited impairment for contextual and cued fear conditioning. Treatment with 7,8-DHF for 4 weeks
22 prevented the decline in fear conditioning, restored LTP in the amygdala, spine density, p-TrkB (Y515 and
23 Y816), p-CaMKII, p-ERK1/2, p-CREB, and p-GluR1 levels in the amygdala, hippocampus, and PFC.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47 **Significance.** These studies show that 7,8-DHF may be used in normal subjects to favor development of
48 dendritic spines, which are the substrate of synaptic plasticity associated with L&M and, consequently, to
49 boost memory performance. Importantly, treatment with 7,8-DHF may represent a suitable treatment to
50 combat and prevent the age-related memory decline.
51
52
53
54
55
56
57

58 **5. ACTIONS OF 7,8-DHF IN THE PERIPHERAL NERVOUS SYSTEM**

5.1 Retina

Brief premise. In the retina BDNF-TrkB signaling has a key role in the development of inner retinal neuronal circuits but not in the rod function and synaptic transmission to bipolar cells (Grishanin et al. 2008). Retinal ganglion cells (RGCs) express high levels of both full-length and truncated forms of TrkB and while TrkB signaling is not required for survival of RGCs during the period of target-dependent survival, it reduces degeneration of RGCs in adult animals (Rohrer et al. 2001). BDNF and TrkB are also important for the differentiation and survival of retinal pigment epithelium (Liu, Zhu, and Eide 1997).

The evidence. In cultures of primary rat RGCs and the retinal neuronal precursor RGC-5 cell line, treatment with 7,8-DHF activated the TrkB receptor, Akt and ERK1/2 (Gupta et al. 2013b). Pretreatment with 7,8 DHF protected cells against the injury caused by glutamate-induced excitotoxicity and H₂O₂-induced oxidative stress and prevented the activation of the apoptotic protein caspase-3. While exposure to glutamate or H₂O₂ decreased by 50% TrkB kinase activity, pretreatment with 7,8 DHF caused a substantial protection of TrkB activity. Recent evidence shows that 7,8-DHF also has a positive impact on retinal pigment epithelial cells (Yu et al. 2018). ARPE-19 cells (a human retinal pigment epithelial cell line) were treated with D-glucose in order to induce apoptosis. Pretreatment with 7,8-DHF prior to glucose, increased the percentage of non-apoptotic cells. Treatment with the TrkB antagonist K252a abrogated this effect and caused downregulation of p-TrkB. A study *in vivo* sought to establish whether 7,8-DHF can protect the immature retina from early HI injury (Huang et al. 2018). Rat pups were treated with 7,8-DHF 2 h and 18 h after induction of HI injury and their retinas were examined at different time points. Treatment with 7,8-DHF improved the functionality of the retina (assessed with electroretinography), counteracted gliosis and reduction in the number of RGCs and favored the survival of proliferating inner retinal cells, including Müller glia, enhancing their trans-differentiation to bipolar cells. The positive effect of 7,8-DHF was prevented by inhibition of the ERK pathway with the ERK inhibitor PD98059, suggesting that these effects were mediated by TrkB-ERK signaling. A recent study (Daly et al. 2017) used a zebrafish model of inherited blindness (*dye^{ucd6}*) in order to establish whether it is possible to pharmacologically improve visual function. Treatment of the mutant larvae with HDAC inhibitors restored retinal morphology and visual function (assessed with functional tests and electroretinography). Treatment with 7,8-DHF alone was sufficient to rescue retinal morphology, reduce the number of dying cells and rescue visual function. The latter effect was greater than that elicited by the most

1
2
3 effective among the HDAC inhibitors. The 7,8-DHF-induced rescue of the visual function was prevented by
4
5 co-treatment with the TrkB inhibitor ANA-12.
6

7 **Significance.** This evidence envisages the possibility that systemic or local administration of 7,8-DHF may be
8
9 a useful therapy for disorders, such as glaucoma, associated with degeneration of RGCs or with
10
11 degeneration of retinal pigment epithelial cells, as well for inherited blindness. In addition, systemic
12
13 administration of 7,8-DHF may be a potential strategy to improve retinal development and function
14
15 following perinatal ischemic injury.
16

17 18 **5.2 Cochlea**

19
20 **Brief premise.** Spiral ganglion neurons (SGN) and cochlear hair cells (HC) express the TrkB receptor
21
22 (Knipper et al. 1996; Liu et al. 2011) and development and maintenance of SGN is critically modulated by
23
24 BDNF (and NT3) (Leake et al. 2011). Degeneration of the HC and SGN underlies age-related hearing loss.
25
26 Following damage of the HC, the peripheral neurites of the SGN gradually degenerate but their cell bodies
27
28 persist for months after degeneration of the HC. Although the delivery of BDNF to the inner ear has been
29
30 shown to exert beneficial effect, its use is hampered by a poor pharmacokinetics.
31

32
33 **The evidence.** An approach *in vitro* and *in vivo* explored the effect of both 7,8-DHF and another agonist of
34
35 the TrkB receptor, 7,8,3'-trihydroxyflavone (7,8,3'-THF), in organotypic cultures of cochlea from C57BL/6
36
37 and TrkB^{F616A} mice and an *in vivo* model of degeneration of the SGN, the conditional connexin26 (cCx26)
38
39 null mouse (Yu et al. 2013). We will mention here the effects of 7,8-DHF only. In the *in vitro* experiments,
40
41 exposure to 7,8-DHF increased the density of SGN and neurite density and length in comparison with the no
42
43 treatment condition, with a potency like that of BDNF. The surviving SGN were functional because they
44
45 could generate inward Na⁺ currents and action potentials. In *in vivo* experiments in cCx26 null mice, 7,8-
46
47 DHF was delivered locally to the cochlea at P2 and P30 and its effects were examined at P60. Mice exhibited a
48
49 greater density of SGN in the treated in comparison with the untreated cochlea and a reduced threshold of the
50
51 electrically evoked auditory brain stem response, which is expression of the function of SGN. A more recent
52
53 study in mice (Kempfle et al. 2018) sought to improve the local delivery of 7,8-DHF to the inner ear through
54
55 the round window membrane by conjugating it with a bisphosphonate (Ris-DHF), based on the rationale
56
57 that bisphosphonates have strong affinity to bone mineral and may prolong the permanence of 7,8-DHF in the
58
59 inner ear. Treatment with either Ris-DHF or 7,8-DHF caused an increase in neurite length of
60

1
2
3 SGN. This effect was abrogated by co-treatment with the TrkB inhibitor ANA-12. In explants of the organ of
4 Corti with attached synapses and exposed to kainate, in order to cause excitotoxic damage of the synapses
5 between the HC and SGN (ribbon synapses), treatment with either Ris-DHF or 7,8-DHF caused an increase
6 in the number of synapses, indicating a regenerating effect.
7
8
9

10
11 **Significance.** These two studies provide proof-of-principle evidence that 7,8-DHF locally delivered to the
12 cochlea may be a suitable therapy for improving damage to the HC and SGN and hearing loss. It remains to
13 be established whether systemic administration of 7,8-DHF elicits comparable effects.
14
15

16 17 **5.3 Peripheral nerves**

18
19 **Brief premise.** Although axons in injured peripheral nerves have the capacity to regenerate and reinnervate
20 their targets, functional recovery is very poor. In the context of nerve regeneration, BDNF appears to play a
21 key role by binding to the TrkB receptor in the regenerating axons. Local treatment with BDNF promotes
22 enhanced early axon regeneration in mice (Wilhelm et al. 2012).
23
24

25
26 **The evidence.** English et al. (English et al. 2013) examined the effects of 7,8-DHF on axon regeneration in
27 wild type and TrkB knockout (*SLICK::trkB^{fl/fl}*) mice. In experiments *in vitro* they found that axon
28 regeneration of the cut sciatic nerve was enhanced by topical treatment with 7,8-DHF. This effect did not
29 take place in mice in which the neurotrophin receptor TrkB was selectively knocked out in neurons,
30 indicating requirement of TrkB signaling. In *in vivo* experiments, mice received 7,8-DHF for 2 weeks after
31 sciatic nerve transection. Systemic treatment was as effective as topical application in enhancing the
32 elongation of regenerating axons. Moreover, treatment increased muscle reinnervation (assessed as
33 amplitude of the response evoked in the gastrocnemius by proximal stimulation of the sciatic nerve) for at
34 least 6 weeks after the end of treatment.
35
36

37
38 **Significance.** Despite many advances in the field, the problem of axon regeneration in the peripheral (as well
39 as central) nervous system remains a challenge and long-lasting denervation may lead to progressive atrophy
40 of the target organs (Fenrich, and Gordon 2004). Systemic or local administration of BDNF as treatment for
41 axon regeneration poses several practical problems. The study by English et al. provides new hopes for the
42 development of effective strategy to treat peripheral nerve injury with systemic administration of 7,8-DHF.
43
44
45
46

47 48 49 50 51 52 53 54 55 56 57 58 59 60 **6. ACTIONS OF 7,8-DHF ON PERIPHERAL TISSUES**

1
2
3 Although BDNF is widely expressed in the CNS and is important for neuronal survival and
4 differentiation it also plays important roles in non-neuronal peripheral tissue.
5
6

7 **6.1 Cardiovascular system**

8 **6.1.1 Heart**

9
10
11 **Brief premise.** BDNF is expressed in the heart (Nakagomi et al. 2015) and plays a fundamental role in
12 heart development (Donovan et al. 2000) and angiogenesis of the heart vascular beds (Kermani, and
13 Hempstead 2007). TrkB receptor have been found in the myocardium (Okada et al. 2012) and
14 BDNF-TrkB signaling represents an important pathway in the heart mechanics, being required for the
15 heart both to fully contract and relax (Feng et al. 2015). Two recent studies provide novel evidence for
16 a pharmacological effect of 7,8-DHF on cardiac damage.
17
18
19
20
21
22
23

24 **The evidence.** Wang et al. showed that treatment with 7,8-DHF improved cardiac dysfunction
25 (ejection fraction, fractional shortening, diastolic and systolic ventricular volume) and attenuated
26 cardiac tissue injury in a myocardial ischemic mouse model (Wang et al. 2019). In the h9c2 cell line of
27 cardiomyocytes exposed to H₂O₂, treatment with 7,8-DHF increased cell viability and inhibited
28 excessive mitochondrial fission through Akt activation. This activation inhibited the proteolytic
29 cleavage of optic atrophy 1 (OPA1), a mitochondrial inner membrane protein that regulates
30 fusion and cristae structure. These effects were mediated by the TrkB receptor, as they were
31 prevented by the TrkB inhibitor ANA-12. A second study investigated the effect of 7,8-DHF on
32 cardiotoxicity induced by Doxorubicin (DOX), an anthracycline used as chemotherapy drug in cancer
33 patients (Zhao et al. 2019). In a DOX mouse model, treatment with 7,8-DHF improved cardiac function
34 (ejection fraction, fractional shortening) and attenuated cardiac tissue injury. In h9c2 cells exposed to
35 DOX, 7,8-DHF increased cell viability, mitochondrial respiration and membrane potential, and the
36 expression of OPA1. The beneficial effects of 7,8-DHF on cardiomyocytes were prevented by the
37 antagonist of the TrkB receptor ANA-12.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **Significance.** These two studies provide novel evidence for a pharmacological effect of 7,8-DHF on
54 cardiac mitochondrial dynamics and ischemia/toxicity-induced heart damage, suggesting a potential
55 therapeutic role of 7,8-DHF in ischemic and metabolic heart diseases.
56
57
58
59

60 **6.1.2 Vessels and blood pressure**

1
2
3 **Brief premise.** TrkB receptors exist in human and rat vascular endothelia and smooth muscles (Donovan et
4 al. 2000; Donovan et al. 1995) and neurotrophins appear to play an important role in regulating the response
5 of vascular smooth muscle cells to injury. Very few studies have examined the effects of 7,8-DHF on
6 vascular dynamics.
7
8
9

10
11 **The evidence.** One of these studies shows that 7,8-DHF induced relaxation of pre-constricted rat aortic rings
12 and that this effect was mediated by an action of 7,8-DHF on both endothelia and smooth muscle cells (Huai et
13 al. 2014). The dilating effect was not mediated by the TrkB receptor because it was not prevented by its
14 inhibitor ANA-12. Rather it appeared to be mediated by the NO/cGMP pathway and through a blocking (in a
15 manner not identified) of both intracellular Ca^{2+} release and extracellular Ca^{2+} influx in the smooth muscle
16 cells. Consistently with the dilating effects observed *in vitro*, intravenous injection of 7,8-DHF significantly
17 reduced the blood pressure of spontaneously hypertensive rats, with no effect on heart rate. This effect
18 occurred within 20 min and disappeared at 60 min after the injection. In addition of being involved in
19 vasodilation by acting on vascular endothelia, 7,8-DHF appears to exert a protective role on vascular
20 endothelial cells. A study in a human umbilical vein endothelial cell line EA.hy926 (Wang et al. 2015) showed
21 that 7, 8-DHF protected the cells from the damage caused by exposure to H_2O_2 by counteracting apoptosis,
22 mitigating release of inflammatory factors (IL-1 β , ICAM-1, and TNF), and inhibiting the generation of
23 reactive oxygen species (ROS). These effects were abrogated by the TrkB antagonist ANA-12. **Significance.**
24 Dysfunction of vascular endothelia favors development of vascular diseases (such as hypertension or
25 atherosclerosis). These two studies suggest that intake of 7,8-DHF may represent a possible prophylaxis
26 against vascular diseases.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **6.2 Pulmonary system**

46
47 **Brief premise.** BDNF and its receptors, including the TrkB receptor, are expressed in different lung
48 structures such as nerves, immune cells, bronchial and alveolar epithelium, smooth muscle, fibroblasts and
49 vascular endothelium (Nakagomi et al. 2015; Prakash, and Martin 2014). BDNF is emerging as a particularly
50 important player in the lung and airways physiology and pathophysiology (Prakash, and Martin 2014).
51
52
53

54 **The evidence.** Cultures of airway smooth muscle exposed to BDNF undergo an increase in Ca^{2+} response to
55 acetylcholine (ACh) and histamine and this effect is mimicked by 7,8-DHF, suggesting that 7,8-DHF,
56 similarly to BDNF, may enhance airway muscle cell contractility (Abcejo et al. 2012). A study by Helan et
57
58
59
60

1
2
3 al. (Helan et al. 2014) showed that human pulmonary artery endothelial cells secrete BDNF in response to
4 hypoxia which, in turn, induces an increase in TrkB-FL/TrkB-T1 ratio, upregulates the expression of hypoxia-
5 inducible factor 1 α (HIF-1 α), that is known to regulate expression of nitric oxide synthase genes and increase
6 iNOS levels and the production of NO. The same effects were elicited by exposure of 7,8-DHF, suggesting a
7 role of BDNF-TrkB (and, hence, 7,8-DHF) signaling in vascular diseases involving hypoxia, such as
8 pulmonary hypertension. A study in Chinese hamster lung fibroblast cells examined the effects of 7,8-DHF on
9 H₂O₂-induced cell damage (Zhang et al. 2009). 7,8-DHF protected lung fibroblasts by scavenging intracellular
10 ROS, inhibiting apoptosis and activating the Akt signaling pathway. A study in the same cell model examined
11 the mechanisms whereby 7,8-DHF exerts cytoprotective effects against oxidative stress, focusing in particular
12 on the antioxidant enzyme HO-1 (Ryu et al. 2014b). The results show that 7,8-DHF protected the cells by
13 increasing the expression of HO-1 and Nrf2, a transcription factor that plays a critical role in the
14 transcriptional regulation of HO-1. Activation of Nrf2 was mediated *via* ERK phosphorylation. Thus, the
15 ERK/Nrf2/HO-1 pathway appears to play an important role in the effects of 7,8-DHF against oxidative stress
16 in lung fibroblasts.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33 **Significance.** These studies indicate that 7,8-DHF affects the function of pulmonary airway smooth muscle
34 cells, vascular endothelial cells and fibroblasts. 7,8-DHF appears to enhance airway muscle cells
35 contractility, thereby increasing airway resistance. On the other hand, 7,8-DHF may have a positive impact
36 on pulmonary hypertension by enhancing the production of NO by endothelial cells. These two opposite
37 effects must be taken into account when administering 7,8-DHF to patients with airway or vascular lung
38 diseases. 7,8-DHF, in addition, appears to ameliorate pulmonary fibroblast function by reducing oxidative
39 stress. Since oxidative stress is implicated in the pathogenesis of pulmonary fibrosis, treatment with 7,8-DHF
40 may positively impact lung function in pathologies involving the lung parenchyma.

41 42 43 44 45 46 47 48 49 **6.3 Gastrointestinal system**

50 51 52 **6.3.1 Intestine**

53
54 **Brief premise.** BDNF is present in the intestine where it is released from mucosal enteroendocrine cells and
55 from intrinsic primary afferent neurons innervating the mucosa. In the intestine, TrkB receptors have been
56 identified in enteric neurons and glia, epithelial cells of the mucosa and smooth muscle cells (see (Al-Qudah
57 et al. 2014)).
58
59
60

1
2
3 **The evidence.** Al-Qudah et al (Al-Qudah et al. 2014) by using strips of intestinal longitudinal smooth muscle
4 demonstrated that BDNF alone does not enhance smooth muscle contraction but it enhances the contraction
5 induced by muscarinic receptor stimulation with charbacol (a commonly used cholinomimetic activator of
6 muscle strip contractions) and that this effect was mediated by activation of the TrkB receptors and the
7 phosphorylation of PLC, most likely PLC γ . Exposure to 7,8-DHF enhanced the contraction induced by
8 muscarinic receptor stimulation, similarly to BDNF. A more recent study has examined the effects of 7,8-
9 DHF on gastric motility *in vitro* and *in vivo* (He et al. 2018). 7,8-DHF alone, did not induce contraction of
10 gastric strips but enhanced charbacol-induced motility. The inhibitor of the TrkB receptor ANA-12 partially
11 blocked this effect. Adult rats orally fed with 7,8-DHF for 7 days exhibited an increase in the velocity of
12 gastric emptying, indicating that *in vivo* 7,8-DHF alone can affect gastric motility.

13
14
15
16
17
18
19
20
21
22
23
24 **Significance.** The increased contractility of smooth muscle associated with gut inflammation has
25 been attributed to enhancement of BDNF-TrkB signaling. Therefore, treatment with 7,8-DHF may have
26 adverse effects on intestinal motility in pathologies associated with gut inflammation. On the other hand,
27 7,8-DHF may be a useful agent for enhancing gastric motility in diseases causing alterations in
28 stomach emptying dynamics.

34 6.3.2 Liver

35
36
37 **Brief premise.** The TrkB receptor is present in the liver of newborn and adult mice (Garcia-Suarez et
38 al. 2006b). In neonate mice (0 to 15 days) the TrkB receptor is present in monocyte-macrophage-dendritic
39 cells scattered throughout the organ. In adult mice (3- and 6-months-old) it is restricted to nerve fibers.

40
41
42 Detectable levels of the two main TrkB ligands, the BDNF and NT-4/5 (Lommatzsch et al. 2005) are also
43 present in the liver at different ages.

44
45
46
47 **The evidence.** High-fat diet and alcohol intake induce liver inflammation and injury. A study in rats
48 subjected to high fat diet and/or alcohol intake for twelve weeks examined the possibility that concomitant
49 treatment with 7,8-DHF during the last 4 weeks may attenuate liver injury (Kumar et al. 2019). This study
50 shows that treatment with 7,8-DHF had a beneficial effect on liver histopathology, restored serum levels of
51 biomarkers of hepatic damage (aspartate aminotransferase and alanine aminotransferase), oxido-nitrosative
52 markers, and the inflammatory cytokine IL-1 β . The diet-induced down-regulation of Nrf-2 and HO-1 and up-
53 regulation of NF-kB and iNOS mRNA expression resulted ameliorated by treatment with 7,8-DHF.
54
55
56
57
58
59
60

1
2
3 **Significance.** This study shows that 7,8-DHF has an hepatoprotective action against liver toxicity induced
4
5 by alcohol and high-fat diet *via* a reduction of oxido-nitrosative stress and NF-kB activation,
6
7 suggesting a potential therapeutic role of 7,8-DHF supplementation in diseases causing liver hepatotoxicity.
8

9 **6.4 Genitourinary system**

10 **6.4.1 Oocytes**

11 **Brief premise.** In the ovary, BDNF secreted by granulosa and cumulus cells acts on TrkB
12
13 receptors expressed in oocytes and enhances first polar body extrusion and promotes the *in vitro*
14
15 development of zygotes (Kawamura et al. 2005). Oxidative stress is heavily involved in the
16
17 physiology of the female reproductive tract and influences the outcomes of assisted reproductive
18
19 technology in human and animal reproduction (Agarwal et al. 2012; Choi et al. 2013). Since 7,8-DHF in
20
21 addition to be a BDNF mimetic has antioxidant properties, it is conceivable that exposure to 7,8-DHF favors
22
23 oocyte maturation in a dual manner. **The evidence.** Choi et al. (Choi et al. 2013) cultured porcine oocytes
24
25 in media supplemented with 7,8-DHF during maturation and culture *in vitro*, after parthenogenic
26
27 activation. Treatment with 7,8-DHF (1 μ M) caused increased oocyte cytoplasmic maturation and
28
29 accelerated attainment of the blastocysts stage. This effect was accompanied by an increase in GSH
30
31 levels, reduction of ROS generation, and reduction of Bcl-2 homologous antagonist/killer (BAK1, pro-
32
33 apoptotic) and increase of Bcl-2-like 1 (BCL2L1, antiapoptotic) protein expression. The contribution of
34
35 the TrkB receptor in these effects remains to be established. **Significance.** In human reproduction,
36
37 increased implantation and clinical pregnancy rates are reported when antioxidant-supplemented medium
38
39 is used in assisted reproductive technology (Agarwal et al. 2012). However, although low
40
41 concentration of 7,8-DHF improves oocyte maturation, higher concentrations have negative effects (Choi
42
43 et al. 2013). Oxidative stress is known to affects female reproductive abilities (Agarwal et al. 2012).
44
45 It remains to be established whether supplementation with 7,8-DHF may have a positive effect on
46
47 reproduction *in vivo*.
48
49
50
51
52
53

54 **6.4.2 Kidney tubular cells**

55 **Brief premise.** BDNF is expressed in the kidney (Nakagomi et al. 2015) and the kidney possesses
56
57 TrkB receptors identical to those in the brain (Garcia-Suarez et al. 2006a), suggesting a role of TrkB
58
59 and its ligands in the control of renal function.
60

1
2
3 **The evidence.** The possible protective function of 7,8-DHF against hypoxia has been examined in the human
4 proximal tubular cell line HK-2 (Ma et al. 2016). This study shows that HK-2 cells express the TrkB receptor
5 and that 7,8-DHF increased their proliferation rate at concentrations of 50-200 μ M but had adverse effects at
6 a higher concentration (250 μ M). In cells exposed to hypoxia 7,8-DHF improved cell viability,
7 possibly through an increase in the expression of cysteine-rich protein 61(CYR61), an extracellular matrix-
8 associated signaling protein that regulates inflammation (Lau 2011). In addition, 7,8-DHF reduced the
9 hypoxia-induced increase in the expression of CCAAT/enhancer-binding protein homologous protein
10 (CHOP), a key regulator of endoplasmatic reticulum (ER) stress. This study did not investigate the pathways
11 involved in the effects of 7,8-DHF on HK-2 cells.

12
13
14
15
16
17
18
19
20
21
22 **Significance.** Acute kidney injury, which is generated in most cases by hypoxia, is characterized by severe
23 renal damage, in particular of the cells of the proximal tubule. This study suggests that treatment with of 7,8-
24 DHF may protect the tubular cells from the hypoxic damage thereby preserving their key role in the renal
25 function.

30 **6.5 Musculoskeletal system**

32 **6.5.1 Muscle**

33
34
35 **Brief premise.** Muscle cells produce BDNF that by acting in an autocrine/paracrine manner regulates
36 muscle physiology. TrkB-mediated signaling is relevant to the maintenance of function and
37 structure at the neuromuscular junction, but not muscle fibers, even though muscle fibers also may
38 express TrkB receptors (Mantilla et al. 2014). Recent data show that BDNF is essential for specification of
39 glycolytic muscle fiber type (Delezie et al. 2019). In addition, BDNF produced locally by muscle cells in
40 response to contraction enhances fat oxidation in skeletal muscle (Matthews et al. 2015).

41
42
43
44
45
46
47 **The evidence.** A study using a diaphragm-phrenic nerve preparation from TrkB^{F616A} mice (sensitive to
48 inhibition by 1NMPP1) examined the effect of 7,8-DHF on muscle fiber properties and neuromuscular
49 transmission (Mantilla, and Ermilov 2012). 7,8-DHF did not directly affect either the twitch force or the
50 fatigability. Treatment with 7,8-DHF, however, reduced the contribution of neuromuscular transmission
51 failure to muscle fatigue which is in agreement with the role of neurotrophins in neuromuscular transmission
52 (Mantilla et al. 2014). This effect was prevented by inhibition of the TrkB receptor with 1NMPP1. A
53 subsequent study explored the possibility that chronic treatment with 7,8-DHF may mitigate the age-related
54
55
56
57
58
59
60

1
2
3 diaphragm neuromuscular transmission failure and sarcopenia (Greising et al. 2017). TrkB^{F616A} mice aged 18
4 months received 7,8-DHF for 6 months and a diaphragm-phrenic nerve preparation from these mice was
5 used to evaluate the effects of treatment. An evaluation of neuromuscular transmission failure, muscle force,
6 and fiber cross-sectional areas at the end of treatment showed no beneficial effect of 7,8-DHF, suggesting
7 that treatment with 7,8-DHF is not useful in order to mitigate age-related neuromuscular functional
8 alterations.

9
10
11
12
13
14
15 **Significance.** Although 7,8-DHF appears to enhance neuromuscular transmission *via* activation of the TrkB
16 receptor in a diaphragm-phrenic nerve preparation, it fails to exert any benefit in age related neuromuscular
17 transmission failure. Further studies should address the potential benefit of 7,8-DHF in neuromuscular
18 junction diseases, such as myasthenia gravis, that are characterized by failure of the neuromuscular synapse
19 due to autoimmune or toxic events.

20 21 22 23 24 25 26 **6.5.2 Articular Chondrocytes**

27
28 **Brief premise.** Articular chondrocytes express neurotrophins and various neurotrophin receptors, including the
29 TrkB receptor and neurotrophins are considered to be of relevance for the articular cartilage and the
30 inflammation process in arthritis (Grimsholm et al. 2008).

31
32
33
34 **The evidence.** Cai et al. (Cai et al. 2019) investigated whether 7,8-DHF can protect cartilage from oxidative
35 stress and delay the progression of articular cartilage destruction in a mouse model of osteoarthritis (OA),
36 induced by sectioning the medial meniscotibial ligament. In cultures of primary chondrocytes, 7,8-DHF
37 activated the Nrf2 signaling pathway and attenuated H₂O₂-induced oxidative stress. OA mice received
38 7,8-DHF once a week postoperatively and their joints were harvested 4 and 8 weeks postoperatively. While
39 vehicle-treated mice exhibited a loss of knee joint cartilage, mice treated with 7,8-DHF exhibited a reduced
40 loss both at 4 and 8 weeks. In the cartilage of the joints, 7,8-DHF suppressed the increase in mRNA levels of
41 TNF- α , IL-1 β , IL-6, and MMP-1, MMP-3 and MMP-13 that occurred in OA mice in comparison with the
42 sham group and increased the expression of HO-1 and Nrf2. This suggests that 7,8-DHF protects the damage
43 cartilage through an antioxidant action.

44
45
46
47
48
49
50
51
52
53
54
55
56 **Significance.** No cure exists for osteoarthritis, a well-known cause of disability around the globe that
57 consumes a substantial amount of healthcare resources. Treatment for OA includes pain relievers and anti-
58 inflammatory drugs and, ultimately, surgery. The study reported above suggests that treatment with 7,8-DHF
59

1
2
3 may represent a suitable therapy that by preserving the articular cartilage may delay/reduce the necessity of
4 symptomatic and/or surgical interventions.
5
6

7 **6.6 Skin**

8
9 **Brief premise.** Neurotrophins and their receptors play a role in the control of skin homeostasis (Botchkarev et
10 al. 1998; Botchkarev et al. 2006), which is not surprising considering the common ectodermal origin of skin
11 and nervous system. In addition, development and maintenance of cutaneous mechanoreceptors has also been
12 shown to depend on BDNF and its TrkB receptor (Garcia-Piqueras et al. 2019). Most of the cells of the
13 epidermis are represented by keratinocytes and most of the cells of the dermis are represent by fibroblasts.
14 Human keratinocytes secrete all neurotrophins and express TrkA and TrkC but do not express the TrkB F.L.,
15 but express the truncated isoform of TrkB (Marconi et al. 2003). In contrast, mouse keratinocytes express the
16 TrkB receptor (Botchkarev et al. 1998). Human dermal fibroblasts secrete all neurotrophins and express all
17 neurotrophin receptors, including the TrkB receptor (Palazzo et al. 2019).
18
19

20
21
22 **The evidence.** Skin aging is characterized by destruction of collagen fibrils due to increased activity of matrix
23 metalloproteinases (MMPs; enzymes that degrade the extracellular matrix) secreted by epidermal
24 keratinocytes and dermal fibroblasts. A study by Choi et al. (Choi, Lee, and Park 2017) examined the effects of
25 7,8-DHF on Hs68 cells (human dermal fibroblasts) treated with TNF- α , an inflammatory cytokine that mediates
26 skin aging and activates MMPs. 7,8-DHF restored the reduction in procollagen caused by exposure to TNF- α
27 and lowered the levels of MMP1, indicating that 7,8-DHF counteracts the reduction in collagen *via* an induction
28 in the synthesis of collagen and a reduction of MMP levels. In addition, 7,8-DHF reduced the increase in ROS
29 production due to exposure to TNF- α . This effect was associated with an increase in the expression levels of the
30 antioxidant enzymes catalase (CAT), Mn-superoxide dismutase (Mn-SOD), and HO-1. These effects were
31 accompanied by suppression of TNF- α -induced activation of the MAPKs and Akt pathways. The effect of 7,8-
32 DHF against oxidative stress has been investigated in human keratinocytes (HaCaT cells), with a focus on
33 HO-1, in view of its powerful cytoprotective properties against oxidative damage (Ryu et al. 2014a). HaCaT
34 cells treated with 7,8-DHF exhibited an increased expression of HO-1 and Nrf2. In cells exposed to H₂O₂ or
35 ultraviolet B (UVB) radiation to cause oxidative stress, treatment with 7,8-DHF improved cell viability and
36 increased the levels of p-ERK and p-PI3K/Akt. ERK and PI3K inhibitors reduced but not abrogated the effects
37 of 7,8-DHF. Since the HaCaT cells do not express functional
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 TrkB receptors (Rossler, and Thiel 2004), it ensues that 7,8-DHF must exert its effects *via* a
4
5 TrkB-independent mechanism.
6

7 **Significance.** Skin aging, which is due to intrinsic and environmental factors, ultimately promotes
8
9 skin inflammation, impaired wound repair, and increased risk of skin cancer. Slowing down skin aging
10
11 may be important in order to preserve the role of the skin, i.e. to protect the underlying tissues, and to
12
13 prevent the occurrence of skin cancer. The results of the two studies reported above, that were obtained *in*
14
15 *vitro*, show that skin cells exposed to 7,8-DHF result protected from oxidative and radiation
16
17 damage. It may be interesting to establish whether a similar effect is elicited *in vivo* by systemic or topic
18
19 administration of 7,8-DHF.
20
21

22 **6.7 Body mass and energy metabolism**

23
24 **Brief premise.** BDNF and the TrkB receptor are expressed in brain centers involved in energy
25
26 homeostasis and BDNF plays a key role in the regulation of body weight and energy balance (see
27
28 (Nakagomi et al. 2015)). BDNF is expressed in the adipose tissue and the TrkB receptor is expressed in
29
30 the adipose tissue at higher levels in comparison with other peripheral tissues (Nakagomi et al. 2015). In
31
32 mice, a high-caloric diet increases BDNF expression and reduces TrkB receptor expression and
33
34 adipocyte-specific BDNF or TrkB deficiency causes obesity and hyperphagia, suggesting an important
35
36 role of BDNF in the peripheral control of energy homeostasis (Nakagomi et al. 2015).
37
38

39 **The evidence.** Based on evidence that oxidative stress induces lipid accumulation and the suppressive effects
40
41 of antioxidants on adipogenesis, a study *in vitro* has examined the anti-obesity effects of 7,8-DHF in 3T3-L1
42
43 preadipocyte cells (Choi et al. 2016). Low concentrations of 7,8-DHF did not affect cell viability, whereas
44
45 high concentrations ($>20\mu\text{M}$) induced apoptotic cell death. During differentiation of 3T3-L1 preadipocytes
46
47 to mature adipocytes, concentrations of 7,8-DHF $<20\mu\text{M}$ reduced the accumulation of lipid droplets and the
48
49 expression of CCAAT/enhancer-binding protein- α (C/EBP- α), C/EBP- β , and peroxisome proliferator
50
51 activated receptor- γ (PPAR- γ), factors that regulate fat storage. In addition, 7,8-DHF reduced the ROS
52
53 accumulation that takes place in adipocytes during adipogenesis and increased the expression of antioxidant
54
55 enzymes such as Mn-SOD, CAT, heme HO-1. MAPK proteins such as ERK, and p38 are activated by ROS
56
57 and this accelerates the differentiation of preadipocyte cells. Treatment with 7,8-DHF reduced the increase in
58
59 p-ERK and p-p38 levels in differentiated cells, suggesting involvement of these pathways in its antioxidant
60

1
2
3 actions. The anti-obesity action of 7,8-DHF has been examined *in vivo* in adult female mice (Chan et al.
4
5 2015). This study was conducted in females because in males 7,8-DHF did not reduce the body weight.
6
7 Chronic administration of 7,8-DHF for 22 weeks attenuated the obesity induced by high fat diet by activating
8
9 the TrkB receptor in muscle. Specifically, treatment reduced the increase in white adipose tissue and the
10
11 hepatic concentration of cholesterol in the liver and triglyceride and free fatty acids in the liver and muscle,
12
13 improved insulin sensitivity and glucose tolerance, reduced the percentage of body fat mass, and increased
14
15 the metabolic rate. At the molecular level, these effects were accompanied by enhanced expression of
16
17 uncoupling protein 1 (UCP1) and AMPK activity in skeletal muscle, p-ERK (liver and muscle) and p-CREB
18
19 (liver, muscle, and white adipose tissue). In muscle-specific TrkB knockout mice, treatment with 7,8-DHF
20
21 failed to mitigate the diet-induced obesity, indicating that the anti-obesity activity of 7,8-DHF was muscular
22
23 TrkB-dependent. A subsequent study (Wood et al. 2018) sought to establish whether treatment with 7,8-DHF
24
25 reverses the metabolic impairment in mice that are already obese. Mouse females aged 8 weeks were fed
26
27 with high fat diet for 13 weeks. Thereafter mice received 7,8-DHF for 9 weeks. Treated mice exhibited a
28
29 reduction in body weight in comparison with control animal, due to a reduction in white adipose tissue, and a
30
31 reduction in hyperlipidemia. In the muscle of treated mice there was an increase in the levels of p-AMPK, p-
32
33 ERK, p-CREB and PGC- α . All these factors play a key role in mitochondrial biogenesis. Accordingly, in
34
35 the muscle of treated mice there was an increase in various mitochondrial markers, including UCP1, and in
36
37 the expressions of TFAM and nuclear respiratory factor 1(Nrf1), two master transcriptional factors of genes
38
39 underlying mitochondrial biogenesis. In line with an increase in mitochondrial biogenesis, treated mice
40
41 exhibited a higher O₂ consumption and an increase in total energy expenditure. Interestingly, treatment also
42
43 improved insulin sensitivity in obese mice.
44
45
46

47 **Significance.** These studies suggest that chronic activation of the muscular TrkB receptor by 7,8-DHF may
48
49 represent a potential strategy against development of obesity. Obesity is a major risk factor for numerous
50
51 diseases including diabetes, and cardiovascular diseases, Therefore, treatment with 7,8-DHF may have an
52
53 impact that is not limited to obesity but may involve serious health issues of the modern society.
54
55 Interestingly, the study by Chan et al. shows that 7,8-DHF has an anti-obesity effect in females but not in
56
57 males (Chan et al. 2015). This issue opens the way to further studies aimed at identifying the causes of the
58
59 sexually dimorphic response to the anti-obesity (and possibly other) effects of 7,8-DHF.
60

7. ANTICANCER ACTION OF 7,8-DHF

Brief premise. Flavonoids have been shown to be potent therapeutic candidates for the treatment of various types of cancer due to their promotion of apoptotic cell death (Abotaleb et al. 2018). Their effects have been documented so far in a variety of carcinomas, which are tumors of epithelial origin. A few studies have specifically examined the potential of 7,8-DHF as anticancer molecule in cell lines of different types of carcinoma.

The evidence.

Breast cancer cell line. The initial evidence for an anticancer action of 7,8-DHF was obtained in a human breast cancer cell line (MCF-7), which is a cell line estrogen hormone-dependent (Le Bail et al. 1998). The goal of the study was to evaluate the anti-proliferative activity of several flavonoids (tested at doses of 10-50,000 nM) in the presence or absence of estradiol. In the absence of estradiol, 7,8-DHF induced a small increment in cell proliferation and had no anti-proliferative activity even at the highest concentration. However, in the presence of estradiol it inhibited cell proliferation at the concentration of 50,000 nM suggesting that it could serve as anti-breast cancer molecule.

Hepatocellular carcinoma cell line. A study in the human hepatocellular carcinoma cell line (HepG2) (Kozics, Valovicova, and Slamenova 2011) examined the protective effects of ten flavonoids, including 7,8-DHF, against the genotoxic effects induced by B(a)P, a potent mutagen and carcinogenic agent that results from incomplete combustion of organic matter at high temperature (e.g. tobacco smoke, grilled meats). A cytotoxicity assay of different concentration of flavonoid (1-100 μ M) showed that 7,8-DHF had a cytotoxicity lower in comparison with most of the other tested flavonoids ($IC_{50} > 100 \mu$ M). Cells exposed to B(a)P exhibited micronuclei, which is indicative of carcinogenicity, and DNA lesions. 7,8-DHF inhibited the increase in cells with micronuclei at all tested concentrations (2.5-25 μ M) and DNA lesions at concentrations of 5-25 μ M. This study thus shows that 7,8-DHF can exert protective effects against genotoxicity at concentration well below those inducing cytotoxicity.

Monocytic leukemia cell line. Park et al. (Park et al. 2012) tested the anti-proliferative effects of 7,8-DHF in U937 human monocytic leukemia cells. Cultures exposed to 7,8-DHF (70 μ M) exhibited a reduced viability due to increased apoptosis. FACS analysis showed that most of the cells exposed to 7,8-DHF were in the G1

1
2
3 phase of the cell cycle. This effect was associated with reduction in the protein levels of cyclin E,
4 upregulation of p27, and reduced phosphorylation of retinoblastoma protein (Rb). The protein p27 prevents
5 the activation of the cyclin E/Cdk2 complex which by phosphorylating Rb promotes G1 progression. These
6 findings provide a mechanistic link between exposure to 7,8-DHF and arrest of the cells in G1 and
7 consequent apoptosis. The cellular pathways that underlie the effects of 7,8-DHF on the cell cycle, however,
8 remain to be established. In a second study (Park et al. 2013), the same group investigated the mechanisms
9 underlying the apoptotic effect of 7,8-DHF in U937 human monocytic leukemia cells. Cultures exposed to
10 7,8-DHF (5-100 μM) exhibited a reduced viability at concentrations $\geq 20 \mu\text{M}$ and in cultures exposed to 70
11 μM concentration of 7,8-DHF there was an increase in apoptotic cell death. These cultures exhibited an
12 increase in various death membrane receptor-related protein and a reduction in inhibitor of apoptosis proteins
13 (IAP; a group of negative regulators of both caspases and cell death). In addition, 7,8-DHF modulated the
14 expression of members of the Bcl-2 family (that acts on the mitochondrion) and released cytochrome c from
15 mitochondria, indicating an involvement of the mitochondrial pathway in its apoptotic effects. The effects of
16 7,8-DHF were mediated by activation of ERK and JNK because inhibitors of ERK and JNK counteracted the
17 7,8-DHF-induced apoptosis.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 *Oral squamous cell carcinoma cell lines.* Lee et al. (Lee et al. 2015) examined the anticancer effect of 7,8-
36 DHF in the oral squamous cell carcinoma cell lines HN22 and HSC4. Cultures exposed to 7,8-DHF (5-40
37 μM) exhibited a reduced viability at concentrations $\geq 5 \mu\text{M}$ and an increase in apoptotic cell death at
38 concentrations of 10-40 μM . Treatment with 7,8-DHF (10-40 μM) reduced the expression of specificity
39 protein 1 (Sp1), a transcription factor that is involved in many cellular processes and plays pivotal roles in
40 cell proliferation and metastasis of various tumors. The reduced expression of Sp1 was accompanied by
41 increased expression of cleaved caspase-3 levels, suggesting, that the reduction in Sp1-levels may take part
42 in the treatment-induced apoptosis increase.

43
44
45
46
47
48
49
50
51
52 *Malignant melanoma cell line.* Sim et al. (Sim, Sohng, and Jung 2016) examined the chemotherapeutic
53 potential and underlying mechanisms of action of 7,8-DHF on malignant melanoma cells using B16F10
54 melanoma cells. Cells were treated with α -MSH, a hormone that stimulates skin and hair melanogenesis and
55 favors melanoma development, and co-exposed to 7,8-DHF. Treatment with 7,8-DHF (0.4-100 μM)
56 inhibited cell proliferation ($\text{IC}_{50} = 9.04 \mu\text{M}$) and colony-forming ability and reduced cell migration capacity.
57
58
59
60

1
2
3 Treatment, in addition, reduced the formation of melanin and tyrosinase activity (the enzyme controlling the
4 production of melanin), suggesting inhibition of differentiation. These effects were associated with
5 downregulation of microphthalmia-associated transcription factor (MITF), a transcription factor associated
6 with melanoma development, and its main downstream transcription target, *HLF1 α* and tyrosine-protein
7 kinase MET (c-MET), through a reduction in cAMP level.
8
9

10
11 **Significance.** These studies clearly show that 7,8-DHF has an anticancer activity *via* inhibition of cell
12 proliferation and enhancement of apoptosis at concentrations higher than those that enhance proliferation.
13 While they provide a hope that a natural compound such as 7,8-DHF may represent a weapon against cancer,
14 the issues of efficacy, toxicity, and safety of large doses of 7,8-DHF *in vivo* still remains to be addressed. In
15 this connection, a recent study must be mentioned that has created 7,8-DHF conjugated to nanoparticles as a
16 tool for targeted antitumoral nanomedicine (Choi et al. 2018) which may provide a tool to overcome the
17 problem of administration of large doses of 7,8-DHF.
18
19
20
21
22
23
24
25
26
27
28
29

30 8. CONCLUSIONS

31
32 The over 100 preclinical studies reviewed here show that treatment with 7,8-DHF has beneficial effects in a
33 number of brain (Fig. 5) and body (Fig. 6) diseases. In most cases the effects consist in a full functional
34 rescue or at least in an improvement. In very few cases (six studies) treatment exerted no benefit (Tables 1
35 and 2) or had a negative effect (one study, Table 1). It must be noted that the majority of the *in vivo* studies
36 used a dose of 5.0 mg/kg/day of 7,8-DHF administered either i.v. or i.p. and showed positive outcome of
37 such a treatment in most cases. Very few studies administered 7,8-DHF *per os* (see Tables 1 and 2) using
38 doses equal to or higher than 5.0 mg/kg/day. It remains to be established whether or not the lack of effects in
39 some of the latter studies is attributable to the scarce absorption rate of 7,8-DHF through the gut epithelium
40 (Chen et al. 2019). Thus, the issue of the appropriate dose of 7,8-DHF when administered *per os* needs still
41 to be better elucidated. Most of the effects of 7,8-DHF appear to be mediated through the TrkB receptor, i.e.
42 *via* a BDNF mimetic action, although some of its effects may exploit additional pathways. The doses
43 exerting positive effects did not cause toxicity, suggesting that 7,8-DHF may represent a safe treatment.
44 Since most of the preclinical studies were carried out in males (see Tables 1 and 2), the question as to
45 whether comparable effects are elicited in females remains open. In addition, a sexually dimorphic response
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to treatment with 7,8-DHF has been found in 6 studies (see Table1 and 2), emphasizing the need to further
4 explore this issue. All *in vivo* studies reviewed here did not report any side effect of even high doses of 7,8-
5 DHF. Yet, despite its enormous benefits, possible adverse effects of 7,8-DHF on brain activity cannot be
6 ignored. For instance, it has been shown that in rat hippocampal-entorhinal cortex slices although 7,8-DHF
7 did not initiate any epileptiform activity it increased the frequency of ictal and interictal events induced by
8 the convulsant 4 aminopyridine (Aydin-Abidin, and Abidin 2019). This effect needs to be better elucidated
9 in the framework of chronic treatments with 7,8-DHF.
10
11

12
13 We must be aware that the experimental evidence showing extremely positive effects of 7,8-DHF in a
14 constellation of disorders has been obtained in animal disease models (in most cases mouse models) and/or in
15 the simplified conditions of *in vitro* experiments. Yet, the large body of coherent evidence accumulated so far
16 in a diversity of models of the same pathology and in different experimental settings encourages us to
17 believe that 7,8-DHF may be proposed as a natural medicine to be tested in future clinical trials for a number
18 of brain and body diseases.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 **Acknowledgements**

35 This study was supported by a grant from Fondazione e Assicurazioni Generali, Italy, to RB.
36
37
38
39
40

41 **Disclosure statement**

42 The authors declare no potential conflict of interest.
43
44
45
46

47 **Author contributions**

48 R. Bartesaghi and F. Stagni conceived, wrote, revised the manuscript and created the tables. M. Emili
49 contributed to write the manuscript and created the figures. S. Guidi, A. Giacomini and B. Uguagliati
50 contributed to write the manuscript.
51
52
53
54
55
56
57

58 **Bibliography**

59
60

- 1
2
3 Abcejo, A. J., V. Sathish, D. F. Smelter, B. Aravamudan, M. A. Thompson, W. R. Hartman, C. M.
4 Pabelick, and Y. S. Prakash. 2012. Brain-derived neurotrophic factor enhances calcium regulatory
5 mechanisms in human airway smooth muscle. *PLoS One*. 7: e44343.
- 6 Abotaleb, M., S. M. Samuel, E. Varghese, S. Varghese, P. Kubatka, A. Liskova, and D. Busselberg.
7 2018. Flavonoids in Cancer and Apoptosis. *Cancers (Basel)*. 11.
- 8 Agarwal, A., A. Aponte-Mellado, B. J. Premkumar, A. Shaman, and S. Gupta. 2012. The effects of
9 oxidative stress on female reproduction: a review. *Reprod Biol Endocrinol*. 10: 49.
- 10 Agrawal, R., E. Noble, E. Tyagi, Y. Zhuang, Z. Ying, and F. Gomez-Pinilla. 2015. Flavonoid derivative
11 7,8-DHF attenuates TBI pathology via TrkB activation. *Biochimica et biophysica acta*. 1852: 862-
12 872.
- 13 Al-Qudah, M., C. D. Anderson, S. Mahavadi, Z. L. Bradley, H. I. Akbarali, K. S. Murthy, and J. R.
14 Grider. 2014. Brain-derived neurotrophic factor enhances cholinergic contraction of longitudinal
15 muscle of rabbit intestine via activation of phospholipase C. *Am J Physiol Gastrointest Liver*
16 *Physiol*. 306: G328-337.
- 17 Amin, N., S. Xie, X. Tan, Y. Chen, Q. Ren, B. O. A. Botchway, S. Hu, Y. Ma, Z. Hu, and M. Fang.
18 2020. Optimized integration of fluoxetine and 7, 8-dihydroxyflavone as an efficient therapy for
19 reversing depressive-like behavior in mice during the perimenopausal period. *Prog*
20 *Neuropsychopharmacol Biol Psychiatry*. 101: 109939.
- 21 Andero, R., N. Daviu, R. M. Escorihuela, R. Nadal, and A. Armario. 2012. 7,8-dihydroxyflavone, a
22 TrkB receptor agonist, blocks long-term spatial memory impairment caused by immobilization stress
23 in rats. *Hippocampus*. 22: 399-408.
- 24 Andero, R., S. A. Heldt, K. Ye, X. Liu, A. Armario, and K. J. Ressler. 2011. Effect of 7,8-
25 dihydroxyflavone, a small-molecule TrkB agonist, on emotional learning. *Am J Psychiatry*. 168:
26 163-172.
- 27 Aydin-Abidin, S., and I. Abidin. 2019. 7,8-Dihydroxyflavone potentiates ongoing epileptiform activity
28 in mice brain slices. *Neurosci Lett*. 703: 25-31.
- 29 Aytan, N., J. K. Choi, I. Carreras, L. Crabtree, B. Nguyen, M. Lehar, J. K. Blusztajn, B. G. Jenkins, and
30 A. Dedeoglu. 2018. Protective effects of 7,8-dihydroxyflavone on neuropathological and
31 neurochemical changes in a mouse model of Alzheimer's disease. *Eur J Pharmacol*. 828: 9-17.
- 32 Barfield, E. T., K. J. Gerber, K. S. Zimmermann, K. J. Ressler, R. G. Parsons, and S. L. Gourley. 2017.
33 Regulation of actions and habits by ventral hippocampal trkB and adolescent corticosteroid
34 exposure. *PLoS Biol*. 15: e2003000.
- 35 Barfield, E. T., and S. L. Gourley. 2017. Adolescent Corticosterone and TrkB Pharmac-Manipulations
36 Sex-Dependently Impact Instrumental Reversal Learning Later in Life. *Front Behav Neurosci*. 11:
37 237.
- 38 Baxter, G. T., M. J. Radeke, R. C. Kuo, V. Makrides, B. Hinkle, R. Hoang, A. Medina-Selby, D. Coit, P.
39 Valenzuela, and S. C. Feinstein. 1997. Signal transduction mediated by the truncated trkB receptor
40 isoforms, trkB.T1 and trkB.T2. *The Journal of neuroscience : the official journal of the Society for*
41 *Neuroscience*. 17: 2683-2690.
- 42 Beck, S., H. Mathison, T. Todorov, C.-J. E, and O. R. Kopp. 2018. A Review of Medicinal Uses and
43 Pharmacological Activities of Tridax Procumbens (L). *Journal of Plant Studies*. 7: 19-35.
- 44 Bollen, E., T. Vanmierlo, S. Akkerman, C. Wouters, H. M. Steinbusch, and J. Prickaerts. 2013. 7,8-
45 Dihydroxyflavone improves memory consolidation processes in rats and mice. *Behav Brain Res*.
46 257: 8-12.
- 47 Bolourian, A., and Z. Mojtahedi. 2018. Streptomyces, shared microbiome member of soil and gut, as
48 'old friends' against colon cancer. *FEMS Microbiology Ecology*. 94: fiy120.
- 49 Boltaev, U., Y. Meyer, F. Tolibzoda, T. Jacques, M. Gassaway, Q. Xu, F. Wagner, Y. L. Zhang, M.
50 Palmer, E. Holson, et al. 2017. Multiplex quantitative assays indicate a need for reevaluating
51 reported small-molecule TrkB agonists. *Sci Signal*. 10.
- 52 Botchkarev, V. A., P. Welker, K. M. Albers, N. V. Botchkareva, M. Metz, G. R. Lewin, S. Bulfone-
53 Paus, E. M. Peters, G. Lindner, and R. Paus. 1998. A new role for neurotrophin-3: involvement in
54 the regulation of hair follicle regression (catagen). *Am J Pathol*. 153: 785-799.
- 55 Botchkarev, V. A., M. Yaar, E. M. Peters, S. P. Raychaudhuri, N. V. Botchkareva, A. Marconi, S. K.
56 Raychaudhuri, R. Paus, and C. Pincelli. 2006. Neurotrophins in skin biology and pathology. *J Invest*
57 *Dermatol*. 126: 1719-1727.

- 1
2
3 Boulle, F., G. Kenis, M. Cazorla, M. Hamon, H. W. Steinbusch, L. Lanfumey, and D. L. van den Hove.
4 2012. TrkB inhibition as a therapeutic target for CNS-related disorders. *Prog Neurobiol.* 98: 197-
5 206.
- 6 Cai, D., W. Feng, J. Liu, L. Jiang, S. Chen, T. Yuan, C. Yu, H. Xie, D. Geng, and J. Qin. 2019. 7,8-
7 Dihydroxyflavone activates Nrf2/HO-1 signaling pathways and protects against osteoarthritis. *Exp*
8 *Ther Med.* 18: 1677-1684.
- 9 Castello, N. A., M. H. Nguyen, J. D. Tran, D. Cheng, K. N. Green, and F. M. LaFerla. 2014. 7,8-
10 Dihydroxyflavone, a small molecule TrkB agonist, improves spatial memory and increases thin
11 spine density in a mouse model of Alzheimer disease-like neuronal loss. *PLoS One.* 9: e91453.
- 12 Cattaneo, A., N. Cattane, V. Begni, C. M. Pariante, and M. A. Riva. 2016. The human BDNF gene:
13 peripheral gene expression and protein levels as biomarkers for psychiatric disorders. *Transl*
14 *Psychiatry.* 6: e958.
- 15 Cazorla, M., A. Jouveneau, C. Rose, J. P. Guilloux, C. Pilon, A. Dranovsky, and J. Premont. 2010.
16 Cyclotraxin-B, the first highly potent and selective TrkB inhibitor, has anxiolytic properties in mice.
17 *PLoS One.* 5: e9777.
- 18 Cazorla, M., J. Premont, A. Mann, N. Girard, C. Kellendonk, and D. Rognan. 2011. Identification of a
19 low-molecular weight TrkB antagonist with anxiolytic and antidepressant activity in mice. *J Clin*
20 *Invest.* 121: 1846-1857.
- 21 Chan, C. B., M. C. Tse, X. Liu, S. Zhang, R. Schmidt, R. Otten, L. Liu, and K. Ye. 2015. Activation of
22 muscular TrkB by its small molecular agonist 7,8-dihydroxyflavone sex-dependently regulates
23 energy metabolism in diet-induced obese mice. *Chem Biol.* 22: 355-368.
- 24 Chang, H. A., Y. H. Wang, C. S. Tung, C. B. Yeh, and Y. P. Liu. 2016. 7,8-Dihydroxyflavone, a
25 Tropomyosin-Kinase Related Receptor B Agonist, Produces Fast-Onset Antidepressant-Like Effects
26 in Rats Exposed to Chronic Mild Stress. *Psychiatry Investig.* 13: 531-540.
- 27 Chao, M. V. 2003. Neurotrophins and their receptors: a convergence point for many signalling
28 pathways. *Nat Rev Neurosci.* 4: 299-309.
- 29 Chen, C., X. H. Li, S. Zhang, Y. Tu, Y. M. Wang, and H. T. Sun. 2014. 7,8-dihydroxyflavone
30 ameliorates scopolamine-induced Alzheimer-like pathologic dysfunction. *Rejuvenation Res.* 17: 249-
31 254.
- 32 Chen, C., Z. Wang, Z. Zhang, X. Liu, S. S. Kang, Y. Zhang, and K. Ye. 2018. The prodrug of 7,8-
33 dihydroxyflavone development and therapeutic efficacy for treating Alzheimer's disease.
34 *Proceedings of the National Academy of Sciences of the United States of America.* 115: 578-583.
- 35 Chen, L., X. Gao, S. Zhao, W. Hu, and J. Chen. 2015. The Small-Molecule TrkB Agonist 7, 8-
36 Dihydroxyflavone Decreases Hippocampal Newborn Neuron Death After Traumatic Brain Injury. *J*
37 *Neuropathol Exp Neurol.* 74: 557-567.
- 38 Chen, Y., F. Xue, G. Xia, Z. Zhao, C. Chen, Y. Li, and Y. Zhang. 2019. Transepithelial transport
39 mechanisms of 7,8-dihydroxyflavone, a small molecular TrkB receptor agonist, in human intestinal
40 Caco-2 cells. *Food Funct.* 10: 5215-5227.
- 41 Choi, D. C., S. L. Gourley, and K. J. Ressler. 2012. Prelimbic BDNF and TrkB signaling regulates
42 consolidation of both appetitive and aversive emotional learning. *Transl Psychiatry.* 2: e205.
- 43 Choi, J. W., C. W. Lee, J. Lee, D. J. Choi, J. K. Sohng, and Y. I. Park. 2016. 7,8-Dihydroxyflavone
44 inhibits adipocyte differentiation via antioxidant activity and induces apoptosis in 3T3-L1
45 preadipocyte cells. *Life Sci.* 144: 103-112.
- 46 Choi, J. W., J. Lee, and Y. I. Park. 2017. 7,8-Dihydroxyflavone attenuates TNF-alpha-induced skin
47 aging in Hs68 human dermal fibroblast cells via down-regulation of the MAPKs/Akt signaling
48 pathways. *Biomed Pharmacother.* 95: 1580-1587.
- 49 Choi, J. Y., J. T. Kang, S. J. Park, S. J. Kim, J. H. Moon, I. M. Saadeldin, G. Jang, and B. C. Lee. 2013.
50 Effect of 7,8-dihydroxyflavone as an antioxidant on in vitro maturation of oocytes and development
51 of parthenogenetic embryos in pigs. *J Reprod Dev.* 59: 450-456.
- 52 Choi, Y. R., H. J. Kim, G. Y. Ahn, M. J. Lee, J. R. Park, D. R. Jun, T. K. Ryu, J. W. Park, E. Shin, and
53 S. W. Choi. 2018. Fabrication of dihydroxyflavone-conjugated hyaluronic acid nanogels for targeted
54 antitumoral effect. *Colloids Surf B Biointerfaces.* 171: 690-697.
- 55 Cikla, U., V. Chanana, D. B. Kintner, E. Udho, J. Eickhoff, W. Sun, S. Marquez, L. Covert, A. Otles, R.
56 A. Shapiro, et al. 2016. ERalpha Signaling Is Required for TrkB-Mediated Hippocampal
57 Neuroprotection in Female Neonatal Mice after Hypoxic Ischemic Encephalopathy(1,2,3). *eNeuro.*
58 3.

- 1
2
3 Colombo, P. S., G. Flamini, M. S. Christodoulou, G. Rodondi, S. Vitalini, D. Passarella, and G. Fico.
4 2014. Farinose alpine Primula species: phytochemical and morphological investigations.
5 *Phytochemistry*. 98: 151-159.
- 6 Daly, C., L. Shine, T. Heffernan, S. Deeti, A. L. Reynolds, J. J. O'Connor, E. T. Dillon, D. J. Duffy, W.
7 Kolch, G. Cagney, et al. 2017. A Brain-Derived Neurotrophic Factor Mimetic Is Sufficient to
8 Restore Cone Photoreceptor Visual Function in an Inherited Blindness Model. *Sci Rep*. 7: 11320.
- 9 Delezie, J., M. Weihrauch, G. Maier, R. Tejero, D. J. Ham, J. F. Gill, B. Karrer-Cardel, M. A. Ruegg, L.
10 Tabares, and C. Handschin. 2019. BDNF is a mediator of glycolytic fiber-type specification in
11 mouse skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of*
12 *America*. 116: 16111-16120.
- 13 DePoy, L. M., A. G. Allen, and S. L. Gourley. 2016. Adolescent cocaine self-administration induces
14 habit behavior in adulthood: sex differences and structural consequences. *Transl Psychiatry*. 6: e875.
- 15 Devi, L., and M. Ohno. 2012. 7,8-dihydroxyflavone, a small-molecule TrkB agonist, reverses memory
16 deficits and BACE1 elevation in a mouse model of Alzheimer's disease. *Neuropsychopharmacology*
17 *: official publication of the American College of Neuropsychopharmacology*. 37: 434-444.
- 18 Donovan, M. J., M. I. Lin, P. Wiegand, T. Ringstedt, R. Kraemer, R. Hahn, S. Wang, C. F. Ibanez, S.
19 Rafii, and B. L. Hempstead. 2000. Brain derived neurotrophic factor is an endothelial cell survival
20 factor required for intramyocardial vessel stabilization. *Development*. 127: 4531-4540.
- 21 Donovan, M. J., R. C. Miranda, R. Kraemer, T. A. McCaffrey, L. Tessarollo, D. Mahadeo, S. Sharif, D.
22 R. Kaplan, P. Tsoulfas, L. Parada, et al. 1995. Neurotrophin and neurotrophin receptors in vascular
23 smooth muscle cells. Regulation of expression in response to injury. *Am J Pathol*. 147: 309-324.
- 24 Du, X., and R. A. Hill. 2015. 7,8-Dihydroxyflavone as a pro-neurotrophic treatment for
25 neurodevelopmental disorders. *Neurochem Int*. 89: 170-180.
- 26 Duart-Castells, L., R. Lopez-Arnau, S. Vizcaino, J. Camarasa, D. Pubill, and E. Escubedo. 2019. 7,8-
27 Dihydroxyflavone blocks the development of behavioral sensitization to MDPV, but not to cocaine:
28 Differential role of the BDNF-TrkB pathway. *Biochem Pharmacol*. 163: 84-93.
- 29 Duval, E. R., A. Javanbakht, and I. Liberzon. 2015. Neural circuits in anxiety and stress disorders: a
30 focused review. *Ther Clin Risk Manag*. 11: 115-126.
- 31 Dwivedi, Y., H. S. Rizavi, R. R. Conley, R. C. Roberts, C. A. Tamminga, and G. N. Pandey. 2003.
32 Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in
33 postmortem brain of suicide subjects. *Arch Gen Psychiatry*. 60: 804-815.
- 34 English, A. W., K. Liu, J. M. Nicolini, A. M. Mulligan, and K. Ye. 2013. Small-molecule trkB agonists
35 promote axon regeneration in cut peripheral nerves. *Proceedings of the National Academy of*
36 *Sciences of the United States of America*. 110: 16217-16222.
- 37 Feng, N., S. Huke, G. Zhu, C. G. Tocchetti, S. Shi, T. Aiba, N. Kaludercic, D. B. Hoover, S. E. Beck, J.
38 L. Mankowski, et al. 2015. Constitutive BDNF/TrkB signaling is required for normal cardiac
39 contraction and relaxation. *Proceedings of the National Academy of Sciences of the United States of*
40 *America*. 112: 1880-1885.
- 41 Fenrich, K., and T. Gordon. 2004. Canadian Association of Neuroscience review: axonal regeneration in
42 the peripheral and central nervous systems--current issues and advances. *Can J Neurol Sci*. 31: 142-
43 156.
- 44 Ferrara, B. T., and E. P. Thompson. 2019. A method for visualizing fluorescence of flavonoid
45 therapeutics in vivo in the model eukaryote Dictyostelium discoideum. *Biotechniques*. 66: 65-71.
- 46 Ferrer, I., C. Marin, M. J. Rey, T. Ribalta, E. Goutan, R. Blanco, E. Tolosa, and E. Marti. 1999. BDNF
47 and full-length and truncated TrkB expression in Alzheimer disease. Implications in therapeutic
48 strategies. *J Neuropathol Exp Neurol*. 58: 729-739.
- 49 Galgano, M., G. Toshkezi, X. Qiu, T. Russell, L. Chin, and L. R. Zhao. 2017. Traumatic Brain Injury:
50 Current Treatment Strategies and Future Endeavors. *Cell Transplant*. 26: 1118-1130.
- 51 Gao, L., M. Tian, H. Y. Zhao, Q. Q. Xu, Y. M. Huang, Q. C. Si, Q. Tian, Q. M. Wu, X. M. Hu, L. B.
52 Sun, et al. 2016. TrkB activation by 7, 8-dihydroxyflavone increases synapse AMPA subunits and
53 ameliorates spatial memory deficits in a mouse model of Alzheimer's disease. *J Neurochem*. 136:
54 620-636.
- 55 Garcia-Diaz Barriga, G., A. Giralt, M. Anglada-Huguet, N. Gaja-Capdevila, J. G. Orlandi, J. Soriano, J.
56 M. Canals, and J. Alberch. 2017. 7,8-dihydroxyflavone ameliorates cognitive and motor deficits in a
57 Huntington's disease mouse model through specific activation of the PLCgamma1 pathway. *Human*
58 *molecular genetics*. 26: 3144-3160.
- 59
60

- 1
2
3 Garcia-Piqueras, J., Y. Garcia-Mesa, L. Carcaba, J. Feito, I. Torres-Parejo, B. Martin-Biedma, J. Cobo,
4 O. Garcia-Suarez, and J. A. Vega. 2019. Ageing of the somatosensory system at the periphery: age-
5 related changes in cutaneous mechanoreceptors. *J Anat.* 234: 839-852.
- 6 Garcia-Suarez, O., T. Gonzalez-Martinez, A. Germana, D. F. Monjil, J. R. Torrecilla, R. Laura, I. Silos-
7 Santiago, J. L. Guate, and J. A. Vega. 2006a. Expression of TrkB in the murine kidney. *Microsc Res*
8 *Tech.* 69: 1014-1020.
- 9 Garcia-Suarez, O., T. Gonzalez-Martinez, M. Perez-Perez, A. Germana, M. A. Blanco-Gelaz, D. F.
10 Monjil, E. Ciriaco, I. Silos-Santiago, and J. A. Vega. 2006b. Expression of the neurotrophin receptor
11 TrkB in the mouse liver. *Anat Embryol (Berl).* 211: 465-473.
- 12 Gardner, E. L. 2011. Addiction and brain reward and antireward pathways. *Adv Psychosom Med.* 30: 22-
13 60.
- 14 Giacomini, A., F. Stagni, M. Emili, B. Uguagliati, R. Rimondini, R. Bartesaghi, and S. Guidi. 2019.
15 Timing of Treatment with the Flavonoid 7,8-DHF Critically Impacts on Its Effects on Learning and
16 Memory in the Ts65Dn Mouse. *Antioxidants (Basel).* 8.
- 17 Grishanin, R. N., H. Yang, X. Liu, K. Donohue-Rolfe, G. C. Nune, K. Zang, B. Xu, J. L. Duncan, M. M.
18 neurotrophin receptors in articular chondrocytes and inflammatory infiltrates in knee joint arthritis.
19 *Cells Tissues Organs.* 188: 299-309.
- 20
21
22 Greene-Schloesser, D., M. E. Robbins, A. M. Peiffer, E. G. Shaw, K. T. Wheeler, and M. D. Chan. 2012.
23 Radiation-induced brain injury: A review. *Front Oncol.* 2: 73.
- 24 Greising, S. M., A. K. Vasdev, W. Z. Zhan, G. C. Sieck, and C. B. Mantilla. 2017. Chronic TrkB agonist
25 treatment in old age does not mitigate diaphragm neuromuscular dysfunction. *Physiol Rep.* 5.
26 Lavan, D. R., Copenhagen, et al. 2008. Retinal TrkB receptors regulate neural development in the
27 inner, but not outer, retina. *Mol Cell Neurosci.* 38: 431-443.
- 28 Grimsholm, O., Y. Guo, I. Ny, and S. Forsgren. 2008. Expression patterns of neurotrophins and
29
30 Gupta, V. K., Y. You, V. B. Gupta, A. Klistorner, and S. L. Graham. 2013a. TrkB receptor signalling:
31 implications in neurodegenerative, psychiatric and proliferative disorders. *Int J Mol Sci.* 14: 10122-
32 10142.
- 33 Gupta, V. K., Y. You, J. C. Li, A. Klistorner, and S. L. Graham. 2013b. Protective effects of 7,8-
34 dihydroxyflavone on retinal ganglion and RGC-5 cells against excitotoxic and oxidative stress. *J*
35 *Mol Neurosci.* 49: 96-104.
- 36 Han, M., J. C. Zhang, and K. Hashimoto. 2017. Increased Levels of C1q in the Prefrontal Cortex of
37 Adult Offspring after Maternal Immune Activation: Prevention by 7,8-Dihydroxyflavone. *Clin*
38 *Psychopharmacol Neurosci.* 15: 64-67.
- 39 Han, M., J. C. Zhang, X. F. Huang, and K. Hashimoto. 2017. Intake of 7,8-dihydroxyflavone from
40 pregnancy to weaning prevents cognitive deficits in adult offspring after maternal immune
41 activation. *Eur Arch Psychiatry Clin Neurosci.* 267: 479-483.
- 42 Han, M., J. C. Zhang, W. Yao, C. Yang, T. Ishima, Q. Ren, M. Ma, C. Dong, X. F. Huang, and K.
43 Hashimoto. 2016. Intake of 7,8-Dihydroxyflavone During Juvenile and Adolescent Stages Prevents
44 Onset of Psychosis in Adult Offspring After Maternal Immune Activation. *Sci Rep.* 6: 36087.
- 45 Han, X., S. Zhu, B. Wang, L. Chen, R. Li, W. Yao, and Z. Qu. 2014. Antioxidant action of 7,8-
46 dihydroxyflavone protects PC12 cells against 6-hydroxydopamine-induced cytotoxicity. *Neurochem*
47 *Int.* 64: 18-23.
- 48 He, B., Z. Qu, Z. Tian, K. Zhao, L. Wei, and L. Ma. 2018. 7,8-dihydroxyflavone enhanced cholinergic
49 contraction of rat gastric smooth muscle via augmenting muscarinic M3 receptor expression. *Clin*
50 *Exp Pharmacol Physiol.* 45: 1170-1180.
- 51 He, J., Z. Xiang, X. Zhu, Z. Ai, J. Shen, T. Huang, L. Liu, W. Ji, and T. Li. 2016. Neuroprotective
52 Effects of 7, 8-dihydroxyflavone on Midbrain Dopaminergic Neurons in MPP(+)-treated Monkeys.
53 *Sci Rep.* 6: 34339.
- 54 Helan, M., B. Aravamudan, W. R. Hartman, M. A. Thompson, B. D. Johnson, C. M. Pabelick, and Y. S.
55 Prakash. 2014. BDNF secretion by human pulmonary artery endothelial cells in response to hypoxia.
56 *J Mol Cell Cardiol.* 68: 89-97.
- 57 Henriques, A., C. Pitzer, and A. Schneider. 2010. Neurotrophic growth factors for the treatment of
58 amyotrophic lateral sclerosis: where do we stand? *Frontiers in neuroscience.* 4: 32.
- 59 Hogarth, S. J., E. Djouma, and M. van den Buuse. 2020. 7,8-Dihydroxyflavone Enhances Cue-
60 Conditioned Alcohol Reinstatement in Rats. *Brain Sci.* 10.

- 1
2
3 Hoshaw, B. A., J. E. Malberg, and I. Lucki. 2005. Central administration of IGF-I and BDNF leads to
4 long-lasting antidepressant-like effects. *Brain Res.* 1037: 204-208.
- 5 Hsiao, Y. H., H. C. Hung, S. H. Chen, and P. W. Gean. 2014. Social interaction rescues memory deficit
6 in an animal model of Alzheimer's disease by increasing BDNF-dependent hippocampal
7 neurogenesis. *The Journal of neuroscience : the official journal of the Society for Neuroscience.* 34:
8 16207-16219.
- 9 Huai, R., X. Han, B. Wang, C. Li, Y. Niu, R. Li, and Z. Qu. 2014. Vasorelaxing and antihypertensive
10 effects of 7,8-dihydroxyflavone. *Am J Hypertens.* 27: 750-760.
- 11 Huang, E. J., and L. F. Reichardt. 2001. Neurotrophins: roles in neuronal development and function.
12 *Annu Rev Neurosci.* 24: 677-736.
- 13 Huang, H. M., C. C. Huang, M. H. Tsai, Y. C. Poon, and Y. C. Chang. 2018. Systemic 7,8-
14 Dihydroxyflavone Treatment Protects Immature Retinas Against Hypoxic-Ischemic Injury via
15 Muller Glia Regeneration and MAPK/ERK Activation. *Invest Ophthalmol Vis Sci.* 59: 3124-3135.
- 16 Huang, Y. Z., and J. O. McNamara. 2012. Neuroprotective effects of reactive oxygen species mediated
17 by BDNF-independent activation of TrkB. *The Journal of neuroscience : the official journal of the*
18 *Society for Neuroscience.* 32: 15521-15532.
- 19 Hung, P. L., C. C. Huang, H. M. Huang, D. G. Tu, and Y. C. Chang. 2013. Thyroxin treatment protects
20 against white matter injury in the immature brain via brain-derived neurotrophic factor. *Stroke.* 44:
21 2275-2283.
- 22 Jäger, A. K., B. Gauguin, A. Adersen, and L. Gudiksen. 2006. Screening of plants used in Danish folk
23 medicine to treat epilepsy and convulsions. *J Ethnopharmacol.* 105: 294-300.
- 24 Jang, S. W., X. Liu, M. Yepes, K. R. Shepherd, G. W. Miller, Y. Liu, W. D. Wilson, G. Xiao, B.
25 Bianchi, Y. E. Sun, et al. 2010. A selective TrkB agonist with potent neurotrophic activities by 7,8-
26 dihydroxyflavone. *Proceedings of the National Academy of Sciences of the United States of America.*
27 107: 2687-2692.
- 28 Jiang, M., Q. Peng, X. Liu, J. Jin, Z. Hou, J. Zhang, S. Mori, C. A. Ross, K. Ye, and W. Duan. 2013.
29 Small-molecule TrkB receptor agonists improve motor function and extend survival in a mouse
30 model of Huntington's disease. *Human molecular genetics.* 22: 2462-2470.
- 31 Johnson, R. A., M. Lam, A. M. Punzo, H. Li, B. R. Lin, K. Ye, G. S. Mitchell, and Q. Chang. 2012. 7,8-
32 dihydroxyflavone exhibits therapeutic efficacy in a mouse model of Rett syndrome. *J Appl Physiol*
33 *(1985).* 112: 704-710.
- 34 brain-derived neurotrophic factor (BDNF) promotes the development of oocytes into
35 preimplantation embryos. *Proceedings of the National Academy of Sciences of the United States of*
36 *America.* 102: 9206-9211.
- 37 Kang, M. S., T. Y. Choi, H. G. Ryu, D. Lee, S. H. Lee, S. Y. Choi, and K. T. Kim. 2017. Autism-like
38 behavior caused by deletion of vaccinia-related kinase 3 is improved by TrkB stimulation. *J Exp*
39 *Med.* 214: 2947-2966.
- 40 Kaur, H., A. Prakash, and B. Medhi. 2013. Drug therapy in stroke: from preclinical to clinical studies.
41 *Pharmacology.* 92: 324-334.
- 42 Kawamura, K., N. Kawamura, S. M. Mulders, M. D. Sollewijn Gelpke, and A. J. Hsueh. 2005. Ovarian
43 Kempfle, J. S., K. Nguyen, C. Hamadani, N. Koen, A. S. Edge, B. A. Kashemirov, D. H. Jung, and C. E.
44 McKenna. 2018. Bisphosphonate-Linked TrkB Agonist: Cochlea-Targeted Delivery of a
45 Neurotrophic Agent as a Strategy for the Treatment of Hearing Loss. *Bioconjug Chem.* 29: 1240-
46 1250.
- 47 Kermani, P., and B. Hempstead. 2007. Brain-derived neurotrophic factor: a newly described mediator of
48 angiogenesis. *Trends Cardiovasc Med.* 17: 140-143.
- 49 Knipper, M., U. Zimmermann, K. Rohbock, I. Kopschall, and H. P. Zenner. 1996. Expression of
50 neurotrophin receptor trkB in rat cochlear hair cells at time of rearrangement of innervation. *Cell*
51 *Tissue Res.* 283: 339-353.
- 52 Koirala, N., R. P. Pandey, P. Parajuli, H. J. Jung, and J. K. Sohng. 2014. Methylation and subsequent
53 glycosylation of 7,8-dihydroxyflavone. *J Biotechnol.* 184: 128-137.
- 54 Korkmaz, O. T., N. Aytan, I. Carreras, J. K. Choi, N. W. Kowall, B. G. Jenkins, and A. Dedeoglu. 2014.
55 7,8-Dihydroxyflavone improves motor performance and enhances lower motor neuronal survival in
56 a mouse model of amyotrophic lateral sclerosis. *Neurosci Lett.* 566: 286-291.
- 57
58
59
60

- 1
2
3 Kowianski, P., G. Lietzau, E. Czuba, M. Waskow, A. Steliga, and J. Morys. 2018. BDNF: A Key Factor
4 with Multipotent Impact on Brain Signaling and Synaptic Plasticity. *Cell Mol Neurobiol.* 38: 579-
5 593.
- 6 Kozics, K., Z. Valovicova, and D. Slamenova. 2011. Structure of flavonoids influences the degree
7 inhibition of Benzo(a)pyrene - induced DNA damage and micronuclei in HepG2 cells. *Neoplasma.*
8 58: 516-524.
- 9 Krishna, G., R. Agrawal, Y. Zhuang, Z. Ying, A. Paydar, N. G. Harris, L. F. F. Royes, and F. Gomez-
10 Pinilla. 2017. 7,8-Dihydroxyflavone facilitates the action exercise to restore plasticity and
11 functionality: Implications for early brain trauma recovery. *Biochim Biophys Acta Mol Basis Dis.*
12 1863: 1204-1213.
- 13 Kumar, D., D. K. Dwivedi, M. Lahkar, and A. Jangra. 2019. Hepatoprotective potential of 7,8-
14 Dihydroxyflavone against alcohol and high-fat diet induced liver toxicity via attenuation of oxido-
15 nitrosative stress and NF-kappaB activation. *Pharmacol Rep.* 71: 1235-1243.
- 16 Kumari, D., and I. Gazy. 2019. Towards Mechanism-Based Treatments for Fragile X Syndrome. *Brain*
17 *Sci.* 9.
- 18 Lai, M. C., M. V. Lombardo, and S. Baron-Cohen. 2014. Autism. *Lancet.* 383: 896-910.
- 19 Lau, L. F. 2011. CCN1/CYR61: the very model of a modern matricellular protein. *Cell Mol Life Sci.* 68:
20 3149-3163.
- 21 Le Bail, J. C., F. Varnat, J. C. Nicolas, and G. Habrioux. 1998. Estrogenic and antiproliferative activities
22 on MCF-7 human breast cancer cells by flavonoids. *Cancer Lett.* 130: 209-216.
- 23 Leake, P. A., G. T. Hradek, A. M. Hetherington, and O. Stakhovskaya. 2011. Brain-derived neurotrophic
24 factor promotes cochlear spiral ganglion cell survival and function in deafened, developing cats. *J*
25 *Comp Neurol.* 519: 1526-1545.
- 26 Lee, R. H., J. C. Shin, K. H. Kim, Y. H. Choi, J. I. Chae, and J. H. Shim. 2015. Apoptotic effects of 7,8-
27 dihydroxyflavone in human oral squamous cancer cells through suppression of Sp1. *Oncol Rep.* 33:
28 631-638.
- 29 Lee, Y., and P. L. Han. 2019. Early-Life Stress in D2 Heterozygous Mice Promotes Autistic-like
30 Behaviors through the Downregulation of the BDNF-TrkB Pathway in the Dorsal Striatum. *Exp*
31 *Neurobiol.* 28: 337-351.
- 32 Levi-Montalcini, R. 1987. The nerve growth factor 35 years later. *Science.* 237: 1154-1162.
- 33 Li, W., and L. Pozzo-Miller. 2014. BDNF deregulation in Rett syndrome. *Neuropharmacology.* 76 Pt C:
34 737-746.
- 35 Li, X., and M. E. Wolf. 2015. Multiple faces of BDNF in cocaine addiction. *Behav Brain Res.* 279: 240-
36 254.
- 37 Li, X. H., C. F. Dai, L. Chen, W. T. Zhou, H. L. Han, and Z. F. Dong. 2016. 7,8-dihydroxyflavone
38 Ameliorates Motor Deficits Via Suppressing alpha-synuclein Expression and Oxidative Stress in the
39 MPTP-induced Mouse Model of Parkinson's Disease. *CNS Neurosci Ther.* 22: 617-624.
- 40 Li, Y., B. W. Luikart, S. Birnbaum, J. Chen, C.-H. Kwon, S. G. Kernie, R. Bassel-Duby, and L. F.
41 Parada. 2008. TrkB regulates hippocampal neurogenesis and governs sensitivity to antidepressive
42 treatment. *Neuron.* 59: 399-412.
- 43 Liu, C., C. B. Chan, and K. Ye. 2016. 7,8-dihydroxyflavone, a small molecular TrkB agonist, is useful
44 for treating various BDNF-implicated human disorders. *Translational neurodegeneration.* 5: 2.
- 45 Liu, J. F., and J. X. Li. 2018. Drug addiction: a curable mental disorder? *Acta Pharmacol Sin.* 39: 1823-
46 1829.
- 47 Liu, W., A. Kinnefors, M. Bostrom, and H. Rask-Andersen. 2011. Expression of TrkB and BDNF in
48 human cochlea-an immunohistochemical study. *Cell Tissue Res.* 345: 213-221.
- 49 Liu, X., C.-B. Chan, S.-W. Jang, S. Pradoldej, J. Huang, K. He, L. H. Phun, S. France, G. Xiao, Y. Jia, et
50 al. 2010. A synthetic 7,8-dihydroxyflavone derivative promotes neurogenesis and exhibits potent
51 antidepressant effect. *J Med Chem.* 53: 8274-8286.
- 52 Liu, X., O. Obianyo, C. B. Chan, J. Huang, S. Xue, J. J. Yang, F. Zeng, M. Goodman, and K. Ye. 2014.
53 Biochemical and biophysical investigation of the brain-derived neurotrophic factor mimetic 7,8-
54 dihydroxyflavone in the binding and activation of the TrkB receptor. *J Biol Chem.* 289: 27571-
55 27584.
- 56 Liu, X., Q. Qi, G. Xiao, J. Li, H. R. Luo, and K. Ye. 2013. O-methylated metabolite of 7,8-
57 dihydroxyflavone activates TrkB receptor and displays antidepressant activity. *Pharmacology.* 91:
58 185-200.

- 1
2
3 Liu, Z. Z., L. Q. Zhu, and F. F. Eide. 1997. Critical role of TrkB and brain-derived neurotrophic factor in
4 the differentiation and survival of retinal pigment epithelium. *The Journal of neuroscience : the*
5 *official journal of the Society for Neuroscience*. 17: 8749-8755.
- 6 Lommatzsch, M., D. Quarcoo, O. Schulte-Herbruggen, H. Weber, J. C. Virchow, H. Renz, and A.
7 Braun. 2005. Neurotrophins in murine viscera: a dynamic pattern from birth to adulthood. *Int J Dev*
8 *Neurosci*. 23: 495-500.
- 9 Lynch, W. J., A. B. Peterson, V. Sanchez, J. Abel, and M. A. Smith. 2013. Exercise as a novel treatment
10 for drug addiction: a neurobiological and stage-dependent hypothesis. *Neurosci Biobehav Rev*. 37:
11 1622-1644.
- 12 Ma, R., J. Zhang, X. Liu, S. Yue, Q. Zhao, and Y. Xu. 2016. 7,8-DHF Treatment Induces Cyr61
13 Expression to Suppress Hypoxia Induced ER Stress in HK-2 Cells. *Biomed Res Int*. 2016: 5029797.
- 14 Mahmoud, B., A. Hamed, M. Samy, and M. Kamel. 2019. Phytochemical and biological overview of
15 genus "Bignonia" (1969-2018) *J. Adv. Biomed. & Pharm. Sci*. 2: 83-97.
- 16 Mantilla, C. B., and L. G. Ermilov. 2012. The novel TrkB receptor agonist 7,8-dihydroxyflavone
17 enhances neuromuscular transmission. *Muscle Nerve*. 45: 274-276.
- 18 Mantilla, C. B., J. M. Stowe, D. C. Sieck, L. G. Ermilov, S. M. Greising, C. Zhang, K. M. Shokat, and
19 G. C. Sieck. 2014. TrkB kinase activity maintains synaptic function and structural integrity at adult
20 neuromuscular junctions. *J Appl Physiol (1985)*. 117: 910-920.
- 21 Marconi, A., M. Terracina, C. Fila, J. Franchi, F. Bonte, G. Romagnoli, R. Maurelli, C. M. Failla, M.
22 Dumas, and C. Pincelli. 2003. Expression and function of neurotrophins and their receptors in
23 cultured human keratinocytes. *J Invest Dermatol*. 121: 1515-1521.
- 24 Matthews, V. B., M. B. Astrom, M. H. Chan, C. R. Bruce, K. S. Krabbe, O. Prelovsek, T. Akerstrom, C.
25 Yfanti, C. Broholm, O. H. Mortensen, et al. 2015. Erratum to: Brain-derived neurotrophic factor is
26 produced by skeletal muscle cells in response to contraction and enhances fat oxidation via
27 activation of AMP-activated protein kinase. *Diabetologia*. 58: 854-855.
- 28 Middlemas, D. S., R. A. Lindberg, and T. Hunter. 1991. trkB, a neural receptor protein-tyrosine kinase:
29 evidence for a full-length and two truncated receptors. *Mol Cell Biol*. 11: 143-153.
- 30 Mukai, R., J. Terao, Y. Shirai, N. Saito, and Ashida, H. 2011. Determination of Subcellular Localization
31 of Flavonol in Cultured Cells by Laser Scanning, Laser Scanning, Theory and Applications, Chau-
32 Chang Wang, IntechOpen. Chapter 12: 215-232.
- 33 Murota, K., Y. Nakamura, and M. Uehara. 2018. Flavonoid metabolism: the interaction of metabolites
34 and gut microbiota. *Biosci Biotechnol Biochem*. 82: 600-610.
- 35 Nagahara, A. H., D. A. Merrill, G. Coppola, S. Tsukada, B. E. Schroeder, G. M. Shaked, L. Wang, A.
36 Minamino. 2015. Role of the central nervous system and adipose tissue BDNF/TrkB axes in
37 metabolic regulation. *NPJ Aging Mech Dis*. 1: 15009.
- 38 Nakagomi, A., S. Okada, M. Yokoyama, Y. Yoshida, I. Shimizu, T. Miki, Y. Kobayashi, and T.
39 Bresch, A. Kim, J. M. Conner, et al. 2009. Neuroprotective effects of brain-derived neurotrophic
40 factor in rodent and primate models of Alzheimer's disease. *Nat Med*. 15: 331-337.
- 41
42
43 Nie, S., K. Ma, M. Sun, M. Lee, Y. Tan, G. Chen, Z. Zhang, Z. Zhang, and X. Cao. 2019. 7,8-
44 Dihydroxyflavone Protects Nigrostriatal Dopaminergic Neurons from Rotenone-Induced
45 Neurotoxicity in Rodents. *Parkinsons Dis*. 2019: 9193534.
- 46 Nijveldt, R. J., E. van Nood, D. E. van Hoorn, P. G. Boelens, K. van Norren, and P. A. van Leeuwen.
47 2001. Flavonoids: a review of probable mechanisms of action and potential applications. *Am J Clin*
48 *Nutr*. 74: 418-425.
- 49 Northington, F. J. 2006. Brief update on animal models of hypoxic-ischemic encephalopathy and
50 neonatal stroke. *ILAR J*. 47: 32-38.
- 51 Okada, S., M. Yokoyama, H. Toko, K. Tateno, J. Moriya, I. Shimizu, A. Nojima, T. Ito, Y. Yoshida, Y.
52 Kobayashi, et al. 2012. Brain-derived neurotrophic factor protects against cardiac dysfunction after
53 myocardial infarction via a central nervous system-mediated pathway. *Arterioscler Thromb Vasc*
54 *Biol*. 32: 1902-1909.
- 55 Olsen, D., M. Kaas, J. Lundhede, S. Molgaard, A. Nykjaer, M. Kjolby, S. D. Ostergaard, and S. Glerup.
56 2019. Reduced Alcohol Seeking and Withdrawal Symptoms in Mice Lacking the BDNF Receptor
57 SorCS2. *Frontiers in pharmacology*. 10: 499.
- 58 Otis, J. M., M. K. Fitzgerald, and D. Mueller. 2014. Infralimbic BDNF/TrkB enhancement of GluN2B
59 currents facilitates extinction of a cocaine-conditioned place preference. *The Journal of neuroscience*
60 *: the official journal of the Society for Neuroscience*. 34: 6057-6064.

- 1
2
3 Palazzo, E., A. Marconi, C. Pincelli, and M. I. Morasso. 2019. Do DLX3 and CD271 Protect Human
4 Keratinocytes from Squamous Tumor Development? *Int J Mol Sci.* 20.
5 Panche, A. N., A. D. Diwan, and S. R. Chandra. 2016. Flavonoids: an overview. *J Nutr Sci.* 5: e47.
6 Pandey, S. C. 2016. A Critical Role of Brain-Derived Neurotrophic Factor in Alcohol Consumption. *Biol*
7 *Psychiatry.* 79: 427-429.
8 Pandey, S. N., M. Kwatra, D. K. Dwivedi, P. Choubey, M. Lahkar, and A. Jangra. 2020. 7,8-
9 Dihydroxyflavone alleviated the high-fat diet and alcohol-induced memory impairment: behavioral,
10 biochemical and molecular evidence. *Psychopharmacology (Berl).*
11 Park, H. Y., G. Y. Kim, J. W. Hyun, N. D. Kim, C. G. Kim, W. J. Kim, Y. H. Yoo, and Y. H. Choi.
12 2012. 7,8-dihydroxyflavone induces G1 arrest of the cell cycle in U937 human monocytic leukemia
13 cells via induction of the Cdk inhibitor p27 and downregulation of pRB phosphorylation. *Oncol Rep.*
14 28: 353-357.
15 Park, H. Y., G. Y. Kim, T. K. Kwon, H. J. Hwang, N. D. Kim, Y. H. Yoo, and Y. H. Choi. 2013.
16 Apoptosis induction of human leukemia U937 cells by 7,8-dihydroxyflavone hydrate through
17 modulation of the Bcl-2 family of proteins and the MAPKs signaling pathway. *Mutat Res.* 751: 101-
18 108.
19 Parrini, M., D. Ghezzi, G. Deidda, L. Medrihan, E. Castroflorio, M. Alberti, P. Baldelli, L. Cancedda,
20 and A. Contestabile. 2017. Aerobic exercise and a BDNF-mimetic therapy rescue learning and
21 memory in a mouse model of Down syndrome. *Sci Rep.* 7: 16825.
22 Perez-Rando, M., E. Castillo-Gomez, C. Bueno-Fernandez, and J. Nacher. 2018. The TrkB agonist 7,8-
23 dihydroxyflavone changes the structural dynamics of neocortical pyramidal neurons and improves
24 object recognition in mice. *Brain Struct Funct.* 223: 2393-2408.
25 Perry, J. L., J. E. Joseph, Y. Jiang, R. S. Zimmerman, T. H. Kelly, M. Darna, P. Huettl, L. P. Dwoskin,
26 and M. T. Bardo. 2011. Prefrontal cortex and drug abuse vulnerability: translation to prevention and
27 treatment interventions. *Brain Res Rev.* 65: 124-149.
28 Poewe, W., K. Seppi, C. M. Tanner, G. M. Halliday, P. Brundin, J. Volkmann, A. E. Schrag, and A. E.
29 Lang. 2017. Parkinson disease. *Nat Rev Dis Primers.* 3: 17013.
30 Pradhan, J., P. G. Noakes, and M. C. Bellingham. 2019. The Role of Altered BDNF/TrkB Signaling in
31 Amyotrophic Lateral Sclerosis. *Front Cell Neurosci.* 13: 368.
32 Prakash, Y. S., and R. J. Martin. 2014. Brain-derived neurotrophic factor in the airways. *Pharmacol*
33 *Ther.* 143: 74-86.
34 Prins, M., T. Greco, D. Alexander, and C. C. Giza. 2013. The pathophysiology of traumatic brain injury
35 at a glance. *Dis Model Mech.* 6: 1307-1315.
36 Rantamaki, T., P. Hendolin, A. Kankaanpaa, J. Mijatovic, P. Piepponen, E. Domenici, M. V. Chao, P. T.
37 Mannisto, and E. Castren. 2007. Pharmacologically diverse antidepressants rapidly activate brain-
38 derived neurotrophic factor receptor TrkB and induce phospholipase-Cgamma signaling pathways in
39 mouse brain. *Neuropsychopharmacology.* 32: 2152-2162.
40 Ray, M. T., C. Shannon Weickert, and M. J. Webster. 2014. Decreased BDNF and TrkB mRNA
41 expression in multiple cortical areas of patients with schizophrenia and mood disorders. *Transl*
42 *Psychiatry.* 4: e389.
43 Ren, Q., M. Ma, C. Yang, J. C. Zhang, W. Yao, and K. Hashimoto. 2015. BDNF-TrkB signaling in the
44 nucleus accumbens shell of mice has key role in methamphetamine withdrawal symptoms. *Transl*
45 *Psychiatry.* 5: e666.
46 Ren, Q., J. C. Zhang, Y. Fujita, M. Ma, J. Wu, and K. Hashimoto. 2013. Effects of TrkB agonist 7,8-
47 dihydroxyflavone on sensory gating deficits in mice after administration of methamphetamine.
48 *Pharmacol Biochem Behav.* 106: 124-127.
49 Ren, Q., J. C. Zhang, M. Ma, Y. Fujita, J. Wu, and K. Hashimoto. 2014. 7,8-Dihydroxyflavone, a TrkB
50 agonist, attenuates behavioral abnormalities and neurotoxicity in mice after administration of
51 methamphetamine. *Psychopharmacology (Berl).* 231: 159-166.
52 Rhine, M. A., J. M. Parrott, M. N. Schultz, T. M. Kazdoba, and J. N. Crawley. 2019. Hypothesis-driven
53 investigations of diverse pharmacological targets in two mouse models of autism. *Autism Res.* 12:
54 401-421.
55 Rohrer, B., M. M. LaVail, K. R. Jones, and L. F. Reichardt. 2001. Neurotrophin receptor TrkB activation
56 is not required for the postnatal survival of retinal ganglion cells in vivo. *Exp Neurol.* 172: 81-91.
57
58
59
60

- 1
2
3 Rossler, O. G., and G. Thiel. 2004. Brain-derived neurotrophic factor-, epidermal growth factor-, or A-
4 Raf-induced growth of HaCaT keratinocytes requires extracellular signal-regulated kinase. *Am J*
5 *Physiol Cell Physiol.* 286: C1118-1129.
- 6 Ryu, M. J., K. A. Kang, M. J. Piao, K. C. Kim, J. Zheng, C. W. Yao, J. W. Cha, H. S. Chung, S. C. Kim,
7 E. Jung, et al. 2014a. 7,8-Dihydroxyflavone protects human keratinocytes against oxidative stress-
8 induced cell damage via the ERK and PI3K/Akt-mediated Nrf2/HO-1 signaling pathways. *Int J Mol*
9 *Med.* 33: 964-970.
- 10 Ryu, M. J., K. A. Kang, M. J. Piao, K. C. Kim, J. Zheng, C. W. Yao, J. W. Cha, C. L. Hyun, H. S.
11 Chung, J. C. Park, et al. 2014b. Effect of 7, 8-dihydroxyflavone on the up-regulation of Nrf2-
12 mediated heme oxygenase-1 expression in hamster lung fibroblasts. *In Vitro Cell Dev Biol Anim.* 50:
13 549-554.
- 14 Sanz-Garcia, A., S. Knafo, I. Pereda-Perez, J. A. Esteban, C. Venero, and A. Armario. 2016.
15 Administration of the TrkB receptor agonist 7,8-dihydroxyflavone prevents traumatic stress-induced
16 spatial memory deficits and changes in synaptic plasticity. *Hippocampus.* 26: 1179-1188.
- 17 Sasi, M., B. Vignoli, M. Canossa, and R. Blum. 2017. Neurobiology of local and intercellular BDNF
18 signaling. *Pflugers Arch.* 469: 593-610.
- 19 Sconce, M. D., M. J. Churchill, C. Moore, and C. K. Meshul. 2015. Intervention with 7,8-
20 dihydroxyflavone blocks further striatal terminal loss and restores motor deficits in a progressive
21 mouse model of Parkinson's disease. *Neuroscience.* 290: 454-471.
- 22 Seese, R. R., A. A. Le, K. Wang, C. D. Cox, G. Lynch, and C. M. Gall. 2020. A TrkB agonist and
23 ampakine rescue synaptic plasticity and multiple forms of memory in a mouse model of intellectual
24 disability. *Neurobiology of disease.* 134: 104604.
- 25 Sen, S., R. Duman, and G. Sanacora. 2008. Serum brain-derived neurotrophic factor, depression, and
26 antidepressant medications: meta-analyses and implications. *Biol Psychiatry.* 64: 527-532.
- 27 Shirayama, Y., C. Yang, J. C. Zhang, Q. Ren, W. Yao, and K. Hashimoto. 2015. Alterations in brain-
28 derived neurotrophic factor (BDNF) and its precursor proBDNF in the brain regions of a learned
29 helplessness rat model and the antidepressant effects of a TrkB agonist and antagonist. *Eur*
30 *Neuropsychopharmacol.* 25: 2449-2458.
- 31 Sim, D. Y., J. K. Sohng, and H. J. Jung. 2016. Anticancer activity of 7,8-dihydroxyflavone in melanoma
32 cells via downregulation of alpha-MSH/cAMP/MITF pathway. *Oncol Rep.* 36: 528-534.
- 33 Sommer, C. J. 2017. Ischemic stroke: experimental models and reality. *Acta Neuropathol.* 133: 245-261.
- 34 Spencer, J. P., M. M. Abd-el-Mohsen, and C. Rice-Evans. 2004. Cellular uptake and metabolism of
35 flavonoids and their metabolites: implications for their bioactivity. *Arch Biochem Biophys.* 423: 148-
36 161.
- 37 Stagni, F., A. Giacomini, M. Emili, S. Guidi, and R. Bartsaghi. 2018. Neurogenesis impairment: An
38 early developmental defect in Down syndrome. *Free radical biology & medicine.* 114: 15-32.
- 39 Stagni, F., A. Giacomini, S. Guidi, E. Ciani, and R. Bartsaghi. 2015. Timing of therapies for Down
40 syndrome: the sooner, the better. *Frontiers in behavioral neuroscience.* 9: 265.
- 41 Stagni, F., A. Giacomini, S. Guidi, M. Emili, B. Uguagliati, M. E. Salvalai, V. Bortolotto, M. Grilli, R.
42 Rimondini, and R. Bartsaghi. 2017. A flavonoid agonist of the TrkB receptor for BDNF improves
43 hippocampal neurogenesis and hippocampus-dependent memory in the Ts65Dn mouse model of DS.
44 *Exp Neurol.* 298: 79-96.
- 45 Stoilov, P., E. Castren, and S. Stamm. 2002. Analysis of the human TrkB gene genomic organization
46 reveals novel TrkB isoforms, unusual gene length, and splicing mechanism. *Biochem Biophys Res*
47 *Commun.* 290: 1054-1065.
- 48 Sun, T., S. Chen, H. Huang, T. Li, W. Yang, and L. Liu. 2017. Metabolic profile study of 7, 8-
49 dihydroxyflavone in monkey plasma using high performance liquid chromatography-tandem mass
50 spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.* 1061-1062: 97-102.
- 51 Tapas, A., D. Sakarkar, and R. Kakde. 2008. Flavonoids as Nutraceuticals: A Review. . *Trop J Pharm*
52 *Res.* 7.
- 53 Tecuatl, C., G. Herrera-Lopez, A. Martin-Avila, B. Yin, S. Weber, G. Barrionuevo, and E. J. Galvan.
54 2018. TrkB-mediated activation of the phosphatidylinositol-3-kinase/Akt cascade reduces the
55 damage inflicted by oxygen-glucose deprivation in area CA3 of the rat hippocampus. *Eur J*
56 *Neurosci.* 47: 1096-1109.
- 57
58
59
60

- 1
2
3 Tian, M., Y. Zeng, Y. Hu, X. Yuan, S. Liu, J. Li, P. Lu, Y. Sun, L. Gao, D. Fu, et al. 2015. 7, 8-
4 Dihydroxyflavone induces synapse expression of AMPA GluA1 and ameliorates cognitive and spine
5 abnormalities in a mouse model of fragile X syndrome. *Neuropharmacology*. 89: 43-53.
- 6 Tome, D., C. P. Fonseca, F. L. Campos, and G. Baltazar. 2017. Role of Neurotrophic Factors in
7 Parkinson's Disease. *Curr Pharm Des*. 23: 809-838.
- 8 Toyooka, K., K. Asama, Y. Watanabe, T. Muratake, M. Takahashi, T. Someya, and H. Nawa. 2002.
9 Decreased levels of brain-derived neurotrophic factor in serum of chronic schizophrenic patients.
10 *Psychiatry Res*. 110: 249-257.
- 11 Uluc, K., P. Kendigelen, E. Fidan, L. Zhang, V. Chanana, D. Kintner, E. Akture, C. Song, K. Ye, D.
12 Sun, et al. 2013. TrkB receptor agonist 7, 8 dihydroxyflavone triggers profound gender- dependent
13 neuroprotection in mice after perinatal hypoxia and ischemia. *CNS Neurol Disord Drug Targets*. 12:
14 360-370.
- 15 Vonsattel, J. P., and M. DiFiglia. 1998. Huntington disease. *J Neuropathol Exp Neurol*. 57: 369-384.
- 16 Wang, B., N. Wu, F. Liang, S. Zhang, W. Ni, Y. Cao, D. Xia, and H. Xi. 2014. 7,8-dihydroxyflavone, a
17 small-molecule tropomyosin-related kinase B (TrkB) agonist, attenuates cerebral ischemia and
18 reperfusion injury in rats. *J Mol Histol*. 45: 129-140.
- 19 Wang, B., Q. Zhang, R. Yao, X. Liu, and Z. Qu. 2015. 7, 8-Dihydroxyflavone Protects an Endothelial
20 Cell Line from H2O2 Damage. *PLoS One*. 10: e0135345.
- 21 Wang, Z., S. P. Wang, Q. Shao, P. F. Li, Y. Sun, L. Z. Luo, X. Q. Yan, Z. Y. Fan, J. Hu, J. Zhao, et al.
22 2019. Brain-derived neurotrophic factor mimetic, 7,8-dihydroxyflavone, protects against myocardial
23 ischemia by rebalancing optic atrophy 1 processing. *Free radical biology & medicine*. 145: 187-197.
- 24 Wilhelm, J. C., M. Xu, D. Cucoranu, S. Chmielewski, T. Holmes, K. S. Lau, G. J. Bassell, and A. W.
25 English. 2012. Cooperative roles of BDNF expression in neurons and Schwann cells are modulated
26 by exercise to facilitate nerve regeneration. *The Journal of neuroscience : the official journal of the*
27 *Society for Neuroscience*. 32: 5002-5009.
- 28 Williams, R. J., J. P. Spencer, and C. Rice-Evans. 2004. Flavonoids: antioxidants or signalling
29 molecules? *Free radical biology & medicine*. 36: 838-849.
- 30 Wood, J., M. C. L. Tse, X. Yang, D. Brobst, Z. Liu, B. P. S. Pang, W. S. Chan, A. M. Zaw, B. K. C.
31 Chow, K. Ye, et al. 2018. BDNF mimetic alleviates body weight gain in obese mice by enhancing
32 mitochondrial biogenesis in skeletal muscle. *Metabolism*. 87: 113-122.
- 33 Wu, C. H., T. H. Hung, C. C. Chen, C. H. Ke, C. Y. Lee, P. Y. Wang, and S. F. Chen. 2014. Post-injury
34 treatment with 7,8-dihydroxyflavone, a TrkB receptor agonist, protects against experimental
35 traumatic brain injury via PI3K/Akt signaling. *PLoS One*. 9: e113397.
- 36 Yang, J., L. C. Harte-Hargrove, C. J. Siao, T. Marinic, R. Clarke, Q. Ma, D. Jing, J. J. Lafrancois, K. G.
37 Bath, W. Mark, et al. 2014a. proBDNF negatively regulates neuronal remodeling, synaptic
38 transmission, and synaptic plasticity in hippocampus. *Cell Rep*. 7: 796-806.
- 39 Yang, P., D. Leu, K. Ye, C. Srinivasan, J. R. Fike, and T. T. Huang. 2016. Cognitive impairments
40 following cranial irradiation can be mitigated by treatment with a tropomyosin receptor kinase B
41 agonist. *Exp Neurol*. 279: 178-186.
- 42 Yang, Y. J., Y. K. Li, W. Wang, J. G. Wan, B. Yu, M. Z. Wang, and B. Hu. 2014b. Small-molecule
43 TrkB agonist 7,8-dihydroxyflavone reverses cognitive and synaptic plasticity deficits in a rat model
44 of schizophrenia. *Pharmacol Biochem Behav*. 122: 30-36.
- 45 Yu, Q., Q. Chang, X. Liu, Y. Wang, H. Li, S. Gong, K. Ye, and X. Lin. 2013. Protection of spiral
46 ganglion neurons from degeneration using small-molecule TrkB receptor agonists. *The Journal of*
47 *neuroscience : the official journal of the Society for Neuroscience*. 33: 13042-13052.
- 48 Yu, X., Q. Liu, X. Wang, H. Liu, and Y. Wang. 2018. 7,8-Dihydroxyflavone ameliorates high-glucose
49 induced diabetic apoptosis in human retinal pigment epithelial cells by activating TrkB. *Biochem*
50 *Biophys Res Commun*. 495: 922-927.
- 51 Zeng, Y., Y. Liu, M. Wu, J. Liu, and Q. Hu. 2012a. Activation of TrkB by 7,8-dihydroxyflavone
52 prevents fear memory defects and facilitates amygdalar synaptic plasticity in aging. *J Alzheimers*
53 *Dis*. 31: 765-778.
- 54 Zeng, Y., F. Lv, L. Li, H. Yu, M. Dong, and Q. Fu. 2012b. 7,8-dihydroxyflavone rescues spatial memory
55 and synaptic plasticity in cognitively impaired aged rats. *J Neurochem*. 122: 800-811.
- 56 Zhang, F., Z. Kang, W. Li, Z. Xiao, and X. Zhou. 2012. Roles of brain-derived neurotrophic
57 factor/tropomyosin-related kinase B (BDNF/TrkB) signalling in Alzheimer's disease. *J Clin*
58 *Neurosci*. 19: 946-949.
- 59
60

- 1
2
3 Zhang, J. C., J. Wu, Y. Fujita, W. Yao, Q. Ren, C. Yang, S. X. Li, Y. Shirayama, and K. Hashimoto.
4 2014a. Antidepressant effects of TrkB ligands on depression-like behavior and dendritic changes in
5 mice after inflammation. *Int J Neuropsychopharmacol.* 18.
6 Zhang, J. C., W. Yao, C. Dong, C. Yang, Q. Ren, M. Ma, M. Han, and K. Hashimoto. 2015. Comparison
7 of ketamine, 7,8-dihydroxyflavone, and ANA-12 antidepressant effects in the social defeat stress
8 model of depression. *Psychopharmacology (Berl).* 232: 4325-4335.
9 Zhang, M. W., S. F. Zhang, Z. H. Li, and F. Han. 2016. 7,8-Dihydroxyflavone reverses the depressive
10 symptoms in mouse chronic mild stress. *Neurosci Lett.* 635: 33-38.
11 Zhang, R., K. A. Kang, M. J. Piao, D. O. Ko, Z. H. Wang, W. Y. Chang, H. J. You, I. K. Lee, B. J. Kim,
12 S. S. Kang, et al. 2009. Preventive effect of 7,8-dihydroxyflavone against oxidative stress induced
13 genotoxicity. *Biol Pharm Bull.* 32: 166-171.
14 Zhang, Z., X. Liu, J. P. Schroeder, C. B. Chan, M. Song, S. P. Yu, D. Weinschenker, and K. Ye. 2014b.
15 7,8-dihydroxyflavone prevents synaptic loss and memory deficits in a mouse model of Alzheimer's
16 disease. *Neuropsychopharmacology.* 39: 638-650.
17 Zhao, J., J. Du, Y. Pan, T. Chen, L. Zhao, Y. Zhu, Y. Chen, Y. Zheng, Y. Liu, L. Sun, et al. 2019.
18 Activation of cardiac TrkB receptor by its small molecular agonist 7,8-dihydroxyflavone inhibits
19 doxorubicin-induced cardiotoxicity via enhancing mitochondrial oxidative phosphorylation. *Free*
20 *radical biology & medicine.* 130: 557-567.
21 Zhao, S., X. Gao, W. Dong, and J. Chen. 2016a. The Role of 7,8-Dihydroxyflavone in Preventing
22 Dendrite Degeneration in Cortex After Moderate Traumatic Brain Injury. *Mol Neurobiol.* 53: 1884-
23 1895.
24 Zhao, S., A. Yu, X. Wang, X. Gao, and J. Chen. 2016b. Post-Injury Treatment of 7,8-Dihydroxyflavone
25 Promotes Neurogenesis in the Hippocampus of the Adult Mouse. *J Neurotrauma.* 33: 2055-2064.
26 Zhou, W., X. Li, D. Huang, W. Zhou, T. Li, and W. Song. 2015. No significant effect of 7,8-
27 dihydroxyflavone on APP processing and Alzheimer-associated phenotypes. *Curr Alzheimer Res.*
28 12: 47-52.
29 Zimmermann, K. S., J. A. Yamin, D. G. Rainnie, K. J. Ressler, and S. L. Gourley. 2017. Connections of
30 the Mouse Orbitofrontal Cortex and Regulation of Goal-Directed Action Selection by Brain-Derived
31 Neurotrophic Factor. *Biol Psychiatry.* 81: 366-377.
32 Zou, J., Z. Zhang, F. Xu, S. Cui, C. Qi, J. Luo, Z. Wang, X. Lu, Z. Tu, X. Ren, et al. 2019. GZD2202, a
33 novel TrkB inhibitor, suppresses BDNF-mediated proliferation and metastasis in neuroblastoma
34 models. *J Drug Target.* 27: 442-450.
35 Zuccato, C., and E. Cattaneo. 2007. Role of brain-derived neurotrophic factor in Huntington's disease.
36 *Prog Neurobiol.* 81: 294-330.
37
38
39

40 Web References

- 41 Web reference 1 [https://www.ema.europa.eu/en/documents/herbal-summary/primula-flower-](https://www.ema.europa.eu/en/documents/herbal-summary/primula-flower-summary-public_it.pdf)
42 [summary-public_it.pdf](https://www.ema.europa.eu/en/documents/herbal-summary/primula-flower-summary-public_it.pdf)
43 Web reference 2 [https://www.ema.europa.eu/en/documents/herbal-report/final-assessment-report-](https://www.ema.europa.eu/en/documents/herbal-report/final-assessment-report-primula-veris-l/primula-elatior-l-hill-radix_en.pdf)
44 [primula-veris-l/primula-elatior-l-hill-radix_en.pdf](https://www.ema.europa.eu/en/documents/herbal-report/final-assessment-report-primula-veris-l/primula-elatior-l-hill-radix_en.pdf)
45
46
47
48

49 LEGEND TO FIGURES

50
51
52
53 Fig. 1. Number of articles by year of publication that have examined the effects of 7,8-DHF in animal or
54 cellular models in the period January 1990-June 2020. Source: PubMed using keywords “7,8-
55 Dihydroxyflavone” or “7,8-DHF”.
56
57
58
59
60

1
2
3 Fig. 2. TrkB receptor isoforms. TrkB full length (TrkB F.L.) bears a tyrosine kinase domain that
4 autophosphorylates upon BDNF-triggered receptor dimerization. TrkB autophosphorylation, in turn, elicits
5 activation of the cellular pathways summarized in Fig. 3. The truncated form of TrkB (TrkB-T1) lacks the
6 tyrosine kinase domain and does not activate these pathways. Dimerization of TrkB-T1 with TrkB F.L. has a
7 dominant negative effect.
8
9
10
11
12
13
14

15
16 Fig. 3. TrkB signaling pathways. Dimerization of the TrkB receptor leads to transphosphorylation of the
17 autophosphorylation loops (Tyr705/6; not shown here) and phosphorylation of tyrosine residues 490, 515
18 and 816. Phosphorylation and recruitment of adaptors to Tyr515 and 490 launches the MAPK signaling
19 cascade (Ras/Raf/MEK/ERK) and Akt signaling cascade. Phosphorylation of Tyr 816 launches the PLC γ
20 signaling cascade. Abbreviations: Akt, protein kinase B; CaM, Calmodulin; CaMKII, Ca²⁺/calmodulin-
21 dependent protein kinases; CREB, cAMP response element binding protein; DAG, diacylglycerol; ERK,
22 extracellular signal-regulated kinase; IP3, inositol triphosphate; PI3K, phosphoinositide 3-kinases; PKC,
23 protein kinase C; PLC γ , phospholipase C γ ; Shc, Src-homology 2- domain.
24
25
26
27
28
29
30
31
32
33
34

35 Fig 4. Chemical structure of 7,8-dihydroxyflavone (7,8-Dihydroxy-2-phenyl-4*H*-1-benzopyran-4-one).
36
37
38

39 Fig. 5. Brain disorders positively affected by treatment with 7,8-DHF. 7,8-DHF also positively impacts
40 memory in the healthy brain. Abbreviation: MDPV, 3,4-Methylenedioxypropylvalerone.
41
42
43
44

45 Fig. 6. Body organs positively affected by treatment 7,8-DHF.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Effect of treatment with 7,8-dihydroxyflavone on the central and peripheral nervous system.

Condition/Tissue/Function	Model	Gender	Age	Treatment Duration	Dose and route of administration	Notes	Reference
GENETIC DISORDERS							
Down syndrome	Ts65Dn mice	M & F	P3	13 D or 45 D	5 mg/kg/day s.c.		(Stagni et al. 2017)
	Ts65Dn mice	M	16-20 W	4 W	~ 22 mg/kg/day p.o.		(Parrini et al. 2017)
	Ts65Dn mice	M & F	P3	15 D	5 mg/kg/day s.c.	NB	(Giacomini et al. 2019)
Rett		M	4 Mo	40 D	5 mg/kg/day i.p.	NB	(Giacomini et al. 2019)
	Mecp2 mutant mice	N.A.	4 W	4 W	80 mg/l p.o. dose N.A		(Johnson et al. 2012)
Fragile X	<i>Fmr1</i> KO mice	M	21 D	4 W	~ 345 mg/kg/day p.o.		(Tian et al. 2015)
	<i>Fmr1</i> KO mice	M	3-5 Mo	1 D or 4 D	5 mg/kg/twice day i.p.		(Seese et al. 2020)
	<i>Fmr1</i> KO mice	M	3-5 Mo	1 Mo	0.08 mg/ml p.o. dose N.A.		(Seese et al. 2020)
	<i>Fmr1</i> KO mice (slices)	M	3-4 Mo	30 min	1 µM		(Seese et al. 2020)
PSYCHIATRIC DISORDERS							
Autism	Shank3B mouse	M & F	6-16 W	5 D *	5 mg/kg/day i.p.		(Rhine et al. 2019)
	BTBR mice	M & F	6-16 W	7 D *	2.5, 5.0, 7.5, 15.0 mg/kg/day i.p.		(Rhine et al. 2019)
Schizophrenia	VRK3-KO mice	M & F	12 W	3 D	10 mg/kg/day i.p.		(Kang et al. 2017)
	VRK3-KO mice	M & F	4 W	8 W	10 mg/kg/week i.p.		(Kang et al. 2017)
	D2 ^{+/-} C57BL/6 mice	M & F	8-10 W	3 D	10 mg/kg/day i.p.		(Lee, and Han 2019)
	Sprague-Dawley rats	M	2 Mo	14 D	5 mg/kg/day i.p		(Yang et al. 2014)
	ddY mice	M	4 W	4 W	1 mg/ml p.o. dose N.A.		(Han et al. 2016)
	ddY mice	M	4 W	4 W	1 mg/ml p.o. dose N.A.		(Han, Zhang, and Hashimoto 2017)
	ddY mice	M	E10	E10 to 3 W	1 mg/ml p.o. dose N.A.		(Han et al. 2017)
Major depressive disorder	C57BL/6J mice	M	2-3 Mo	21 D	5 mg/kg/day p.o.		(Liu et al. 2010)
	C57BL/6 mice	M	8 W	1 injection	1, 3, 10 mg/kg, i.p.		(Zhang et al. 2014a)
	C57Bl/6 mice	M	8 W	1 injection	10 mg/kg, i.p		(Zhang et al. 2015)
	Sprague-Dawley rats	M	7 W	1 infusion	0.1, 1.0 pmol b.i.		(Shirayama et al. 2015)
	C57BL/6 mice	M	8 W	28 D	10, 20 mg/kg/day i.p.		(Zhang et al. 2016)

		Sprague-Dawley rats	M	8 W	2 W	5, 20 mg/kg i.p.		(Chang et al. 2016)
		C57BL/6J mice	F	Adult	28 D	5 mg/kg/day i.p.		(Amin et al. 2020)
	Anxiety and stress disorder	Sprague-Dawley rats	M	2 Mo	1 injection	5 mg/kg s.c.		(Andero et al. 2012)
		C57BL/6J mice	M	2-4 Mo	1 injection	5 mg/kg i.p.		(Andero et al. 2011)
		Sprague-Dawley rats	M	60 D	1 injection	5 mg/kg s.c.		(Sanz-Garcia et al. 2016)
		<i>bdnf</i> -floxed and TrkB ^{F616A} mice	M	2-4 Mo	1 injection	5 mg/kg i.p.		(Choi, Gourley, and Ressler 2012)
		C57BL/6 mice	M	31 D	8 D	3, 10 mg/kg/day i.p.		(Barfield et al. 2017)
		C57BL/6 mice	M & F	31 D	8 D	3 mg/kg/day i.p.	SD	(Barfield, and Gourley 2017)
	ADDICTION							
	Cocaine	<i>bdnf</i> -floxed mice	M	2-4 Mo	1 injection	5 mg/kg i.p.		(Choi, Gourley, and Ressler 2012)
		Rats	M	3-6 Mo	1 injection	5 mg/kg i.p.		(Otis, Fitzgerald, and Mueller 2014)
		C57BL/6 mice	M & F	Adult	1 injection	5 mg/kg i.p.		(DePoy, Allen, and Gourley 2016)
	MDPV	Swiss CD-1 mice	F	41-44 D	5 D	10 mg/kg/day i.p.		(Duart-Castells et al. 2019)
	Methamphetamine	ddy mice	M	8 W	1 injection	3.0, 10, 30 mg/kg, i.p.		(Ren et al. 2013)
		BALB/cAnCrlCrlj	M	10 W	1 injection	3.0, 10, 30 mg/kg i.p.		(Ren et al. 2014)
		C57BL/6 mice	M	8 W	14 D	10 mg/kg/day i.p.	NB	(Ren et al. 2015)
	Alcohol	Wistar rats	M	8-10 W	4 W	5 mg/kg/day i.p.		(Pandey et al. 2020)
		Sprague Dawley rats	M & F	6 W	2 W	5 mg/kg/day i.p.	NB, AE, SD	(Hogarth, Djouma, and van den Buuse 2020)
	BRAIN INJURY							
	Traumatic Brain Injury	Sprague-Dawley rats	M	2 Mo	7 D	5 mg/kg/day i.p.		(Agrawal et al. 2015)
		Sprague-Dawley rats	M	8-10 W	7 D	5 mg/kg/day i.p.		(Krishna et al. 2017)
		C57BL/6J mice	M	8-10 W	4 D	20, 50 mg/kg/day i.p.		(Wu et al. 2014)
		C57BL/6J mice	M	8-10 W	1 injection	5 mg/kg i.p.		(Chen et al. 2015)
		C57BL/6J mice	M	8-10 W	15 D	5 mg/kg/day i.p.		(Zhao et al. 2016b)
		C57BL/6J mice	M	8-10 W	4 D	5 mg/kg/day i.p.		(Zhao et al. 2016a)
	Stroke/Ischemia	TrkB ^{F616A} mice	M	N.A.	1 injection	5 mg/kg i.p.		(Jang et al. 2010)
		Rats	M	7 D	5 D	5 mg/kg/every other day i.p.		(Hung et al. 2013)
		C57BL/6J mice	M & F	9 D	1 or 7 D	5 mg/kg/day i.p.	SD	(Uluc et al. 2013)

	C57BL/6J mice	M & F	9 D	3 D	5 mg/kg/day i.p.	SD	(Cikla et al. 2016)
	Sprague–Dawley rats	M	Adult	1 injection	5 mg/kg i.p.		(Wang et al. 2014)
	Sprague–Dawley rats (slices)	N.A.	35-45 D	–	25 µM		(Tecuatl et al. 2018)
Irradiation	C57BL/6J mice	M	2 Mo	3 or 5 W	5 mg/kg/day s.c. or o.p.		(Yang et al. 2016)
NEURODEGENERATION							
Alzheimer Disease	5XFAD mice	M & F	12-15 Mo	10 D	5 mg/kg/day i.p.		(Devi, and Ohno 2012)
	5XFAD mice	N.A.	2 Mo	2 Mo	5 mg/kg/day p.o.		(Zhang et al. 2014b)
	5XFAD mice	F	1 Mo	2 Mo	5 mg/kg/day i.p.		(Aytan et al. 2018)
	5XFAD mice	N.A.	2 Mo	3 Mo	§ 7.2, 21.8, 43.6 mg/kg/day p.o.		(Chen et al. 2018)
	Tg2576 mice	M	10 Mo	4 W	~ 200 mg/kg/day p.o.		(Gao et al. 2016)
	APPswe/PS1dE9 mice	M	7 Mo	1 injection	0.1 mg/kg i.p.		(Bollen et al. 2013)
	APPswe/PS1dE9 mice	M	9 Mo	10 D	5 mg/kg/day i.p.		(Hsiao et al. 2014)
	CaM/Tet-DT _A mice	M & F	3.5-5 Mo	10 D	5 mg/kg/day i.p.		(Castello et al. 2014)
	APP23/PS45 mice	N.A.	6 W	4 W	5 mg/kg/day i.p.	NB	(Zhou et al. 2015)
Parkinson Disease	Sprague-Dawley rats	M	Adult	4 W	1 mg/kg/day i.p.		(Chen et al. 2014)
	C57BL/6 mice+MPTP	N.A.	8 W	14 D	5 mg/kg/day p.o.		(Jang et al. 2010)
	C57BL/6 mice+MPTP	M	8 W	4 W	5 mg/kg/day i.p.		(Sconce et al. 2015)
	C57BL/6 mice+MPTP	N.A.	Adult	14 D	5 mg/kg/day i.p.		(Li et al. 2016)
	Sprague–Dawley rats+Rot	M	Adult	5 W	5 mg/kg/day i.p.		(Nie et al. 2019)
	Macaca Fascicularis+ MPP+	M	8 Y	7 Mo	30 mg/kg/day po.		(He et al. 2016)
Hungtington’s Disease	N171-82Q mice	M	6 W	14 W	5 mg/kg/day p.o.		(Jiang et al. 2013)
	R6/1 mice	M	8 W	12 W	5 mg/kg/day p.o.		(Garcia-Diaz Barriga et al. 2017)
Amyotrophic Lateral Sclerosis	SOD1 ^{G93A} mice	M	30 D	75 D	5 mg/kg/day i.p.		(Korkmaz et al. 2014)
HEALTHY BRAIN							
	Wistar rats	M	3 Mo	1 injection	0.1, 0.3, 1, 3 mg/kg p.o.		(Bollen et al. 2013)
	Thy1-YFP mice	N.A.	2 Mo	14 D	5 mg/kg/day p.o.		(Perez-Rando et al. 2018)
	Sprague-Dawley rats	M	22&30 Mo	34 D	5 mg/kg/day i.p.		(Zeng et al. 2012b)
	Sprague-Dawley rats	M	24 Mo	4 W	5 mg/kg/day i.p.		(Zeng et al. 2012a)
	C57BL/6 mice	M	8 W	1 injection	5 mg/kg		(Zimmermann et al. 2017)

1
2 **PERIPHERAL**
3 **NERVOUS SYSTEM**

4 Retina	Rat RGCs, RGC-5 cells	–	–	10 min-16 h	100 nM	(Gupta et al. 2013)
	ARPE-19 cells	–	–	48 h	0.01-20 µM	(Yu et al. 2018)
	Sprague-Dawley rats	N.A.	7 D	2 injections	5 mg/kg i.p	(Huang et al. 2018)
	Zebrafish dye ^{ucd6}	N.A.	Embryos	2 D	10 µM	(Daly et al. 2017)
9 Cochlea	Cochlear cultures from C57BL/6 and TrkB ^{F616A} mice	M & F		72 h	300 nM	(Yu et al. 2013)
	cCx26 null mice	M & F	2 D	at 2 D and 30 D	200 µM local application	(Yu et al. 2013)
	SGN cultures CBA/CaJ mice	N.A.	4 D	48 h	400 nM	(Kempfle et al. 2018)
15 Nerves	C57B6 and <i>SLICK::trkB^{fl/fl}</i> mice	M & F	N.A.	2 W	500 nM local application	(English et al. 2013)
	C57B6 and <i>SLICK::trkB^{fl/fl}</i> mice	M & F	N.A.	2 W	5 mg/kg/day, i.p	(English et al. 2013)

21
22 Summary of the treatment methodology of the studies that have examined the effects of 7,8-DHF on the central and peripheral nervous system. Abbreviations: AE,
23 adverse effects; b.i., brain infusion; D, day; F, female; h, hour; i.p., intraperitoneal; M, male; min, minutes; Mo, month; MDPV, 3,4-Methylenedioxypropylvalerone;
24 MPP+, 1-methyl-4-phenylpyridinium; MTPP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; N.A., not available; NB, no benefit; o.p., osmotic pump; P, postnatal day;
25 p.o., per os; s.c., subcutaneous; Rot, Rotenone; SD, sex differences; W, week; Y, year. * plus one injection before behavioral testing; § prodrug R13.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 2. Effect of treatment with 7,8-Dihydroxyflavone on peripheral tissues.

Organ	Model	Gender	Age	Treatment Duration	Dose and route of administration	Notes	Reference
Heart	Kunming mice	N.A.	Adult	4 W	5 mg/kg/day i.p.		(Wang et al. 2019)
	H9c2 cells	–	–	24 h	100 μ M		(Wang et al. 2019)
	Kunming mice	N.A.	Adult	3 W	5 mg/kg/day i.p.		(Zhao et al. 2019)
	H9c2 cells	–	–	24 h	1-600 μ M		(Zhao et al. 2019)
Blood Vessels	Rat aortic rings from Wistar rats	M	16 W	acute (minutes)	1-500 μ M		(Huai et al. 2014)
	Wistar rats	M	16 W	1 injection	2.5 mg/kg i.v.		(Huai et al. 2014)
	EA.hy926 cells	–	–	24 h	1-100 μ M		(Wang et al. 2015)
Lung	Human ASM	–	–	24 h	500 nM		(Abcejo et al. 2012)
	Human PAEC	–	–	24 h	N.A.		(Helan et al. 2014)
	Chinese hamster V79-4 lung fibroblasts	–	–	24 h	10 μ g/ml		(Zhang et al. 2009)
	Chinese hamster V79-4 lung fibroblasts	–	–	24 h	0.1, 1, 10 μ g/ml		(Ryu et al. 2014b)
	Muscle strips from the intestine of New Zealand rabbits	N.A.	N.A.	60 min	1 μ M		(Al-Qudah et al. 2014)
Intestine	Muscle strip from the stomach of rats	M	N.A.	5-60 min	1- 100 μ M		(He et al. 2018)
	Rats	M	N.A.	7 D	0.2, 0.8, 3.2 mM in 1 ml/day p.o.		(He et al. 2018)
Liver	Wistar rats	M	Adult	4 W	5 mg/kg/day i.p.		(Kumar et al. 2019)
Oocytes	Porcine oocyte cultures	–	–	7 D	0, 1, 5, 10 μ M		(Choi et al. 2013)
Kidney	HK-2 cells	–	–	1-12 h	50-250 μ M		(Ma et al. 2016)
Muscle	Diaphragm muscle preparations from TrkB ^{F616A} mice	M	Adult	30 min	10 μ M		(Mantilla, and Ermilov 2012)
	TrkB ^{F616A} mice	M	18 Mo	6 Mo	5 mg/kg/day p.o.	NB	(Greising et al. 2017)
Articular Chondrocytes	Primary chondrocytes from B6 mice	N.A.	5 D	8 h	0-10 μ M		(Cai et al. 2019)
	C57BL/6 mice	N.A.	8-10 W	4 or 8 W	5 mg/kg/week i.p.		(Cai et al. 2019)

Table 3. Studies grouped by examined brain parameter.

PARAMETER	REFERENCE
Neurogenesis/Differentiation	(Chen et al. 2015; Giacomini et al. 2019; He et al. 2016; Huang et al. 2018; Hung et al. 2013; Liu et al. 2010; Parrini et al. 2017; Stagni et al. 2017; Yang et al. 2016; Zhao et al. 2016b)
Apoptosis/Neurodegeneration	(Amin et al. 2020; Chen et al. 2015; Daly et al. 2017; Gupta et al. 2013; He et al. 2016; Huang et al. 2018; Jang et al. 2010; Korkmaz et al. 2014; Li et al. 2016; Nie et al. 2019; Pandey et al. 2020; Sconce et al. 2015; Uluc et al. 2013; Wang et al. 2014; Wu et al. 2014; Yu et al. 2013; Yu et al. 2018; Zhang et al. 2014b; Zhao et al. 2016a)
Denditic arbor/Neurites	(Aytan et al. 2018; Garcia-Diaz Barriga et al. 2017; Kempfle et al. 2018; Yu et al. 2013; Zhang et al. 2014b; Zhao et al. 2016a; Zhao et al. 2016b)
Spine density	(Aytan et al. 2018; Barfield et al. 2017; Castello et al. 2014; Gao et al. 2016; Kang et al. 2017; Korkmaz et al. 2014; Perez-Rando et al. 2018; Stagni et al. 2017; Tian et al. 2015; Yang et al. 2016; Zeng et al. 2012a; Zeng et al. 2012b; Zhang et al. 2014a; Zhang et al. 2014b; Zhao et al. 2016a; Zimmermann et al. 2017)
Synaptic proteins	(Agrawal et al. 2015; Barfield et al. 2017; Gao et al. 2016; Krishna et al. 2017; Stagni et al. 2017; Tian et al. 2015; Zeng et al. 2012a; Zhang et al. 2015; Zhang et al. 2016)
Inflammation	(Amin et al. 2020; Garcia-Diaz Barriga et al. 2017; Wang et al. 2014).
Oxidative stress	(Chen et al. 2014; Garcia-Diaz Barriga et al. 2017; Gupta et al. 2013; Li et al. 2016; Pandey et al. 2020; Wang et al. 2014; Yang et al. 2014)
Cell bioenergetics	(Agrawal et al. 2015; Han et al. 2016; Krishna et al. 2017)
LTP	(Chen et al. 2014; Parrini et al. 2017; Sanz-Garcia et al. 2016; Seese et al. 2020; Yang et al. 2014; Zeng et al. 2012a; Zeng et al. 2012b; Zhang et al. 2014b)
Hippocampus-dependent memory (MWM, NOR, CFC)	(Agrawal et al. 2015; Andero et al. 2012; Andero et al. 2011; Bollen et al. 2013; Castello et al. 2014; Chen et al. 2014; Devi, and Ohno 2012; Gao et al. 2016; Garcia-Diaz Barriga et al. 2017; Giacomini et al. 2019; Han et al. 2016; Hsiao et al. 2014; Kang et al. 2017; Krishna et al. 2017; Pandey et al. 2020; Parrini et al. 2017; Perez-Rando et al. 2018; Sanz-Garcia et al. 2016; Seese et al. 2020; Stagni et al. 2017; Tian et al. 2015; Uluc et al. 2013; Yang et al. 2016; Yang et al. 2014; Zeng et al. 2012b; Zhang et al. 2014b; Zhao et al. 2016a; Zhou et al. 2015)
TrkB signaling	(Agrawal et al. 2015; Amin et al. 2020; Andero et al. 2011; Chen et al. 2014; Chen et al. 2015; Cikla et al. 2016; Devi, and Ohno 2012; Garcia-Diaz Barriga et al. 2017; Giacomini et al. 2019; Gupta et al. 2013; Han et al. 2017; Han et al. 2016; Jiang et al. 2013; Krishna et al. 2017; Li et al. 2016; Nie et al. 2019; Parrini et al. 2017; Sconce et al. 2015; Seese et al. 2020; Stagni et al. 2017; Uluc et al. 2013; Wang et al. 2014; Wu et al. 2014; Yang et al. 2016; Yang et al. 2014; Yu et al. 2018; Zeng et al. 2012a; Zeng et al. 2012b; Zhang et al. 2014a; Zhang et al. 2016; Zhang et al. 2014b)
TrkB inhibition	(Agrawal et al. 2015; Chen et al. 2015; Choi, Gourley, and Ressler 2012; Daly et al. 2017; Duarte-Castells et al. 2019; Gao et al. 2016; Garcia-Diaz Barriga et al. 2017; Jang et al. 2010; Kang et al. 2017; Kempfle et al. 2018; Liu et al. 2010; Otis, Fitzgerald, and Mueller 2014; Ren et al. 2015; Ren et al. 2013; Ren et al. 2014; Wu et al. 2014; Yu et al. 2018; Zhang et al. 2014a; Zhang et al. 2016; Zhang et al. 2014b; Zimmermann et al. 2017)

The reported studies have examined the effects of 7,8-DHF on the indicated brain parameters. The parameter “TrkB inhibition” refers to studies that have used TrkB inhibitors in order to demonstrate whether the effects of 7,8-DHF were mediated by the TrkB receptor. Abbreviations; CFC, contextual fear conditioning; LTP, long-term potentiation; MWM, Morris water maze; NOR, novel object recognition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 2




Fig. 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 6