



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

alessandra di francesco, elena baraldi (2021). How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches. *BIOLOGICAL CONTROL*, 152(January 2021), 1-6 [10.1016/j.biocontrol.2020.104456].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/775215> since: 2020-10-20

*Published:*

DOI: <http://doi.org/10.1016/j.biocontrol.2020.104456>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

## Journal Pre-proofs

How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches

Alessandra Di Francesco, Elena Baraldi

PII: S1049-9644(20)30683-6  
DOI: <https://doi.org/10.1016/j.biocontrol.2020.104456>  
Reference: YBCON 104456

To appear in: *Biological Control*

Received Date: 16 October 2019  
Revised Date: 30 June 2020  
Accepted Date: 8 October 2020

Please cite this article as: Di Francesco, A., Baraldi, E., How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches, *Biological Control* (2020), doi: <https://doi.org/10.1016/j.biocontrol.2020.104456>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0>)

1 **How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans***  
2 **against *Monilinia laxa* on peaches**

3  
4 Alessandra Di Francesco\*, Elena Baraldi

5  
6 *Department of Agricultural Sciences, University of Bologna, Viale G. Fanin, 46, 40127 Bologna, Italy*

7 *\*corresponding author: alessand.difrancesc3@unibo.it*

8  
9 **Abstract**

10 In the present study *Aureobasidium pullulans* strains L1 and L8 were shown both *in vitro* and *in*  
11 *vivo* to compete for iron with *Monilinia laxa* through the secretion of siderophores (1.2 and 1.4 mg  
12 ml<sup>-1</sup>, respectively) and to prevent postharvest fruit decay of peaches. The two strains reduced  
13 mycelial growth and conidial germination of *M. laxa* specially in presence of lower iron  
14 concentrations (5 and 10 µg l<sup>-1</sup> FeCl<sub>3</sub>), confirming a better efficacy when nutrients are scarce. In *in*  
15 *vivo* assay, *A. pullulans* L1 and L8 strains inhibited pathogen virulence, reducing by 83.5% and  
16 84.4% on average respectively the peach lesion diameter for each tested iron solutions (5, 10, 20 µg  
17 l<sup>-1</sup> FeCl<sub>3</sub>). The highest iron solution slowed down the antagonists' action and conversely increase  
18 the pathogen aggressiveness. Results shows that *A. pullulans* L1 and L8 strains compete with *M.*  
19 *laxa* for iron, so revealing new biocontrol aspects. Both strains showed the capability to decrease  
20 the accumulation of iron competing with some fungal pathogens and reducing their virulence. These  
21 results provide new perspectives for the use of biocontrol agents in agriculture.

22  
23 **Keywords:** Yeast - Iron – Stone fruit – *Monilinia laxa* - Siderophore

24  
25 **1. Introduction**

26 *Monilinia laxa* is the most common species in European stone fruit orchards, and able to cause  
27 postharvest losses that reach high values (59%) (Larena et al., 2005). *Monilinia* control depends on  
28 an integrated strategy based on fungicide use and cultural practices. However, to overcome the  
29 issues related to the use of fungicides, alternative pathogen control strategies have been  
30 investigated, such as the use of biocontrol agents (BCAs) (Di Francesco et al., 2016).  
31 *Aureobasidium pullulans* L1 and L8 strains showed a high capability to control *Monilinia* spp. on  
32 stone fruit as previously reported by Mari et al. (2012).

33 Among the various modes of action involved in the control of postharvest diseases by antagonistic  
34 yeasts, competition for nutrients and space is considered one of the most important, particularly to  
35 control wound pathogens for which the availability of exogenous nutrients is crucial during the  
36 early stage of infection (Elad and Chet 1987, Mekbib et al., 2011, Bautista-Rosales et al., 2013, Di  
37 Francesco et al., 2016, 2017, 2018a).

38 In fact, previous studies reported how the antifungal ability of *A. pullulans* (Janisiewicz et al., 2000;  
39 Bencheqroun et al., 2006), *Pichia caribbica* (Xu et al., 2013), and *Pichia guilliermondii*  
40 (Chanchaichaovivat et al., 2008) against *Penicillium expansum*, *Rhizopus stolonifer*, and  
41 *Colletotrichum capsici* respectively, can be influenced by the presence or the absence of exogenous  
42 nutrients (Druvefors et al., 2005, Bencheqroun et al., 2006, Liu et al., 2010).

43 Besides the role of competing for nutrients, iron may also play a role in the Biocontrol Agents  
44 (BCAs) modes of action (Raaska and Mattila-Sandholm 1995) through the production of chelating  
45 molecules (siderophores).

46 These are molecules able to chelate low weight ferric compounds enhancing the effectiveness of  
47 BCAs by subtracting iron to pathogen inhibiting its growth and metabolic activity (Riquelme,  
48 1996). Microorganisms can produce a variety of chelating agents, which solubilize ferric iron and  
49 transport it into the cell (Liu et al., 2013, Calvente et al., 2001a, Sanz Ferramola et al., 2013). Under  
50 competitive conditions when nutrient availability is a limiting factor, the production of siderophores  
51 can represent a distinctive feature for a microorganism. It is known that yeasts produce only a type

52 of siderophore: hydroxamate (Riquelme, 1996). The siderophores production could play an  
53 important role in the biocontrol of pathogens because by iron sequestering their growth and  
54 metabolic activity were inhibited or slowed down (Riquelme 1996). Calvente et al. (2001a)  
55 demonstrated that rhodotorulic acid, a siderophore produced by *Rhodotorula glutinis* improved the  
56 biological control of blue rot of apples caused by *P. expansum*. Also, *Bacillus subtilis* produced  
57 siderophores playing an important role in the control of *Fusarium oxysporum* (Yu et al., 2011).  
58 Other studies reported the siderophores ability to inhibit the growth of certain pathogenic fungi, like  
59 *Pythium ultimum* and *Sclerotinia sclerotiorum* (Hamdan et al., 1991, McLoughlin et al., 1992),  
60 establishing that these molecules could be considered as a potential mechanism of action against  
61 several fungal pathogens. Parafati et al. (2015) hypothesized that competition for iron was as one of  
62 the main mechanisms of action for *Metschnikowia pulcherrima* against *Botrytis cinerea*.  
63 In a previous study, *A. pullulans* L1 and L8 strains yet showed competition for nutrients like amino  
64 acids and space (Di Francesco et al., 2017, 2018a) against *Monilinia laxa* and *B. cinerea*.  
65 In the present work, we investigated for the first time the involvement of iron in the biocontrol  
66 activity of both strains against the brown rot agent of peaches. Specifically, the aims of this study  
67 were to determine the ability of L1 and L8 to produce siderophores through i) the chrome azurol S  
68 (CAS) assay, ii) the spectrophotometer quantification, and iii) the *in vitro* and *in vivo* competition  
69 for iron with *M. laxa* by testing three different concentrations of  $\text{FeCl}_3$  (5, 10 and 20  $\mu\text{g ml}^{-1}$ ).

70

## 71 2. Materials and methods

72

### 73 2.1 Fruit

74 Peaches (*Prunus persica* (L.) Batsch) cv 'Redhaven' were harvested at commercial maturity in  
75 orchards located in Cadriano (Bologna, Italy). Fruit-were stored at 0 °C and used within 5 days from  
76 harvest and wounded by a sterile nail (3x3x3 mm) on opposite sides of the equatorial area.

77

## 78 2.2 Pathogen and antagonists

79 *Monilinia laxa* (ML4 strain from CRIOF-DipSA collection) (Di Francesco et al., 2017) was chosen  
80 for the present study for its aggressiveness and grown as by Martini et al. (2016). Conidia from  
81 pathogen colonies 10 days-old, grown on tomato agar (20 g of Agar Technical, Oxoid Basingstoke,  
82 Hampshire, UK; 750 ml distilled water to which 250 ml tomato sauce was added after sterilization)  
83 at 25 °C, were collected and suspended in sterile distilled water containing 0.05 %-(v/v) Tween 80.  
84 The concentration of each conidia suspension was quantified with haematocytometer and adjusted  
85 to a concentration of  $10^5$  conidia  $\text{ml}^{-1}$  with sterile distilled water. *Aureobasidium pullulans* L1 and  
86 L8 strains (Di Francesco et al., 2018b), were maintained on nutrient yeast dextrose agar (NYDA: 8  
87 g of nutrient broth, 5 g of yeast extract, 10 g of dextrose and 15 g of agar in 1 L of distilled water) at  
88 4 °C until use. Two days before trials, each antagonist was inoculated on NYDA and incubated at  
89 25 °C for 2 days. The yeast cells were collected in sterile distilled water and adjusted with  
90 haematocytometer to a concentration of  $10^8$  cells  $\text{ml}^{-1}$ .

## 92 2.3 CAS assay

93 CAS-blue agar was prepared according to Schwyn and Neilands (1987). Dishes were prepared with  
94 30 ml of culturing medium for each antagonist and fungus: NYDA for L1 and L8 and PDA for *M.*  
95 *laxa*. Then the solidified growth media was cut in two halves, one of which was replaced by CAS-  
96 blue agar. The dishes were inoculated **placing in the centre of the borderline**, between the two  
97 media, a 6-mm-plug of L1, L8 or *M. laxa* derived from a colony in active growth. The plates were  
98 incubated in the dark at 25 °C until the agar colour-change. The CAS reaction was determined by  
99 measuring the advance of the colour-change in the CAS-blue agar from blue to different shades of  
100 red. The control plates were not inoculated but incubated under the same conditions described  
101 above. The sample unit was represented by 10 dishes and the experiment was repeated twice.

102

## 103 2.4 Spectrophotometer siderophore assay

104 For the siderophore production by yeasts and *M. laxa*, one loop of L1 and L8 cells and one of *M.*  
105 *laxa* conidia were transferred in a siderophore solution as described in Wang et al. (2009) and  
106 incubated at 25 °C at 200 rpm for 24 h. In addition, L1 or L8 cells and *M. laxa* were co-cultured in  
107 the siderophore culture solution and incubated in the same conditions above mentioned. Aliquots of  
108 5 ml of each sample's suspension were inoculated into 45 ml of new siderophore solution and  
109 suddenly incubated for 120 h at 25 °C at 200 rpm. The supernatant of the centrifuged (5.000 rpm  
110 for 20 min) cultures (500 µl) was added to 2.5 ml of a solution containing 5 mM FeCl<sub>3</sub> and used for  
111 quantitative determination of siderophore by a spectrophotometer at 440 nm (Infinite® 200 PRO-  
112 Tecan) after 30 min of incubation at 25 °C. A standard curve (0, 0.05, 0.1, 0.25, 0.5, 0.75 and 1.0  
113 mg ml<sup>-1</sup>) was prepared with deferoxamine mesylate (Sigma-Aldrich) (one type of hydroxamates) as  
114 chelating agent standard (Calvente et al., 2001). The siderophore amount was extrapolated from the  
115 deferoxamine mesylate standard curve. The sample unit was represented by three flasks (replicates)  
116 for each condition. The experiment was repeated twice.

117

### 118 2.5 *In vitro* competition for iron

119 The competition for nutrient exerted by L1 and L8 *A. pullulans* strains on *M. laxa* mycelial growth  
120 was investigated through the dual interaction between the pathogen and the yeast strains. The  
121 experiment conducted with the two strains and the pathogen was assessed on Minimal Medium  
122 Agar (MMA: Na<sub>2</sub>HPO<sub>4</sub> 6 g, KH<sub>2</sub>PO<sub>4</sub> 3 g, NaCl 0.5 g, NH<sub>4</sub>Cl 1 g, MgSO<sub>4</sub> 1M, CaCl<sub>2</sub> 0.1M, Sucrose  
123 30 g, Agar technical 20 g) amended with 0, 5, 10 and 20 µg ml<sup>-1</sup> of FeCl<sub>3</sub> (Saravanakumar et al.,  
124 2008). *Monilinia laxa* mycelial plugs (6 mm of diameter) from 7-day-old colony were inoculated at  
125 30 mm of distance from the plate edge. *Aureobasidium pullulans* L1 and L8 cells were taken from  
126 48 h culture on NYDA by sterile loop and were streaked at 30 mm from the edge of the other side  
127 of the plate. Plates were incubated at 25 °C for 5 days and the radial growth of pathogen mycelium  
128 was measured. Plates inoculated only with *M. laxa* plug represented the control. The sample unit  
129 was represented by five plates (replicates) for each condition. The experiment was performed twice.

130

131 2.6 Influence of iron concentrations on antagonists' biocontrol activity against *Monilinia laxa*  
132 germination and germ tube elongation

133 The competition activity for iron of L1 and L8 was assayed on the germination and germ tube  
134 elongation of *M. laxa* conidia. Tissue culture plates (Costar, Corning Inc., Corning, NY) and culture  
135 plate inserts Millicell-CM (Millipore Corp., Bedford, MA) were used as reported by Janisiewicz et  
136 al., (2000) with some modifications (Di Francesco et al., 2017). Aliquots (120  $\mu\text{l}$ ) of minimal  
137 medium broth (without agar) amended with 0, 5, 10 and 20  $\mu\text{g ml}^{-1}$  of  $\text{FeCl}_3$  were dispensed in the  
138 wells of culture plates with 40  $\mu\text{l}$  of the pathogen conidia suspension ( $10^5$  conidia  $\text{ml}^{-1}$ ), while the  
139 same aliquots of L1 or L8 cell suspensions ( $10^8$  cells  $\text{ml}^{-1}$ ) were dispensed inside the cylinder  
140 inserts, without physical contact between antagonist and pathogen. The plates were placed at 25 °C  
141 on a rotary shaker at 50 rpm; after 6 h of incubation, cylinders were removed from the wells and 20  
142  $\mu\text{l}$  of the medium were transferred to a glass slide for microscope (Nikon Eclipse TE2000–E)  
143 observations. The percentage of conidia germination and the germ tube elongation ( $\mu\text{m}$ ) were  
144 determined (90 conidia per treatment, 3 microscopic fields with 30 conidia each). Cells without iron  
145 solutions **addition** were considered as a control. The experiment was performed twice.

146

147 2.7 *In vivo* competition for iron

148 Peach fruits (20) cv 'Redhaven' were wounded and inoculated with 20  $\mu\text{l}$  of each yeast suspension  
149  $10^8$  cell  $\text{ml}^{-1}$ ; after 1 h at room temperature fruit were inoculated with 20  $\mu\text{l}$  of the pathogen conidia  
150 suspension ( $10^5$  cells  $\text{ml}^{-1}$ ) and right after its drying, 20  $\mu\text{l}$  of iron solutions, at concentrations above  
151 mentioned, were added. Fruit treated with sterile distilled water instead of yeast cell suspensions  
152 represented the control. Fruit were stored at 20 °C **and 90% of relative humidity (RH)** for 7 days  
153 and the lesion diameters determined by the pathogen were recorded. **The sample unit was**  
154 **represented by 45 peaches (15 for replicate) for each antagonist, iron concentration, and control.**  
155 The experiment was repeated twice.



156

## 157 2.8 Statistical analysis

158 Data were statistically handled by one-way analysis of variance (ANOVA). Statistical comparison  
159 of means was carried out to reveal the differences between treatments using Tukey's HSD Test  
160 ( $\alpha = 0.05$ ). Data were reported as mean values  $\pm$  standard error (SE) of two experiments. Before  
161 analysis of data, homogeneity of variance was tested by the Kruskal–Wallis test. All analyses were  
162 performed with Statgraphics software (version centurion 15.0).

163

## 164 3. Results

165

### 166 3.1 *Aureobasidium pullulans* L1, L8 strains siderophore production

167 In order to verify the siderophore production by the antagonists and *M. laxa*, CAS-blue agar was  
168 used as screening assay for the target microorganisms. Our results showed that both L1 and L8  
169 induced a colour change of CAS-blue agar from blue to dark orange indicating that both *A.*  
170 *pullulans* strains are siderophore producers. However, despite the two strains required the same  
171 time (twelve days) to change the CAS-blue-agar from blue to dark orange, the colour change halo  
172 produced by both yeasts was different in size, being 20 mm for L1, and 27 mm for L8 (Table 1)  
173 (Fig. 1). In fact, L1 strain after 12 days of incubation did not determined a total colour change of  
174 CAS-blue agar half, as instead L8. Conversely, *M. laxa* produced a barrage between the two media  
175 without any colour change of the indicative agar. Control plates, without plug inoculation, not  
176 changed colour after incubation conditions.

177

### 178 3.2 Spectrophotometer quantification of siderophores

179 To quantify the siderophore production by *A. pullulans* L1 and L8 strains and the influence of *M.*  
180 *laxa* on this activity, a spectrophotometer assay with deferoxamine mesylate standard curve was  
181 conducted. Both strains showed a considerable activity, producing 1.2 and 1.4 mg ml<sup>-1</sup> of

182 siderophore by L1 and L8 respectively, while *M. laxa* produced only 0.64 mg ml<sup>-1</sup> of siderophore, a  
183 quantity not detectable by CAS-blue-agar.

184 Siderophore quantity was not influenced by *M. laxa* showing the same amount with respect to the  
185 yeasts grown alone (Fig. 2).

186

### 187 3.3 *In vitro* competition for iron

188 Competition for iron was tested co-culturing in MM plates amended with FeCl<sub>3</sub> (0, 5, 10, or 20 µg  
189 ml<sup>-1</sup>) L1 and L8 strains with *M. laxa* isolate. As shown by Table 2, both *A. pullulans* L1 and L8  
190 strains reduced *M. laxa* mycelium diameter by 41.3% and 43.3%, respectively when iron  
191 concentration was not present in MM.

192 L1 and L8 competition was mainly stimulated by the presence of 10 µg ml<sup>-1</sup> and 20 µg ml<sup>-1</sup> of FeCl<sub>3</sub>,  
193 both leading to an inhibition of fungal pathogen mycelial growth corresponding to 45% and 47.6%  
194 and 43% and 45%, respectively.

195 *Aureobasidium pullulans* strain L8 was slightly more effective with respect to L1 in competing for  
196 iron, determining a higher inhibition of *M. laxa* mycelial growth on average by 44.7% and 42.9%,  
197 respectively.

198

### 199 3.4 Influence of iron concentrations on *M. laxa* germination and germ tube elongation

200 After 6 h of incubation at 25 °C in MM broth without iron and in MM broth with 5 µg ml<sup>-1</sup> of FeCl<sub>3</sub>,  
201 *M. laxa* conidial germination was reduced by both L1 and L8 strain of 50%, as showed in Table 3.

202 The presence of 10 µg ml<sup>-1</sup> of FeCl<sub>3</sub> in the culture medium induced a higher antagonistic activity by  
203 both strains against *M. laxa* conidial germination with respect to the other iron concentrations (5  
204 and 20 µg ml<sup>-1</sup> of FeCl<sub>3</sub>). In effect, conidial germination of *M. laxa* was reduced by 62.5% and  
205 68.7%, respectively by L1 and L8. Conversely, 20 µg ml<sup>-1</sup> of FeCl<sub>3</sub> inhibited the strains antagonistic  
206 activity, that showed a reduction of *M. laxa* conidia germination of ~35%.

207 As regards to *M. laxa* germ tube length, the highest inhibition values exerted by L1 and L8 were  
208 mainly showed with 5 and 10  $\mu\text{g ml}^{-1}$  of  $\text{FeCl}_3$  solution by 59.5%, 63.1% and by 65.9%, 67.6%,  
209 respectively. Nevertheless, 10  $\mu\text{g ml}^{-1}$  of  $\text{FeCl}_3$  stimulated mostly pathogen germ tube elongation  
210 (10.4%) with respect to the other concentrations (Table 4). Anyway, L8 strain was always slightly  
211 more effective with respect to L1 in determining a higher inhibition of *M. laxa* conidial germination  
212 and germ tube elongation on average by 52.1%, 50.2%, and by 57.7%, 57.1%, respectively.

213

### 214 3.5 *In vivo* competition for iron

215 The influence of the addition of aliquots of iron solutions at different concentrations on fungal  
216 virulence and yeasts antagonistic activity on peach fruit was measured. *Monilinia laxa* growth on  
217 peach fruit in absence of antagonists was slightly enhanced by 6.8% only at  $\text{FeCl}_3$  concentrations of  
218 10 or 20  $\mu\text{g ml}^{-1}$  whereas no influence was observed on the pathogen aggressiveness at 5  $\mu\text{g ml}^{-1}$ ,  
219 the lowest  $\text{FeCl}_3$  concentrations (Fig. 3).

220 However, L1 and L8 *A. pullulans* strains showed a great capability to inhibit *M. laxa*  
221 aggressiveness, reducing the peach lesion diameters by 83.5% and 84.4% on average respectively,  
222 at each iron concentration. Furthermore, the highest concentration of iron (20  $\mu\text{g ml}^{-1}$ ) seemed to  
223 slow down the antagonists action and conversely increase the pathogen aggressiveness. In fact, both  
224 L1 and L8 controlled *M. laxa* incidence in presence of 20  $\mu\text{g ml}^{-1}$   $\text{FeCl}_3$  on average by 88% with  
225 respect to 92%, in presence of the other iron concentrations (data not reported).

226

## 227 4. Discussion

228 *Aureobasidium pullulans* (L1 and L8) are considered promising alternatives to fungicides in the  
229 reduction of the stone fruits postharvest diseases' incidence caused by *Monilinia* spp. (Mari et al.,  
230 2012; Di Francesco et al., 2017). In our work, competition for nutrients was studied and in  
231 particular the competition for iron, a nutrient that played a significant role in biocontrol interactions  
232 (Raaska and Mattila-Sandholm 1995).

233 As known, iron is an essential nutrient for all organisms (Wang et al., 2009) acting as a cofactor for  
234 enzymes and regulatory proteins involved in many cellular processes (Miethke and Marahiel 2007;  
235 Li et al., 2018) but not promptly available due to the low solubility in alkaline environments. One of  
236 the main strategy used by microorganisms (such as yeasts) and by plants to obtain iron is the  
237 secretion of siderophores (Li et al., 2018), which are low-molecular-weight compounds that chelate  
238 iron in the extracellular phase re-entering it in the cells by specific membrane transporters (Hider  
239 and Kong 2010). More than 500 siderophores have been discovered from microorganisms and  
240 plants and their chemical diversity often renders a variety of biological functions beyond capture  
241 iron (Ho et al., 2019). Yeasts produce only hydroxamate-type siderophore (Riquelme 1996) derived  
242 from the amino acid ornithine and classified into four structural families: fusarines, coprogens,  
243 ferrichromes and rhodotorulic acid (Johnson 2008). Wang et al., (2009) showed that *A. pullulans*  
244 could produce 1.1 mg ml<sup>-1</sup> hydroxamate type siderophore. In our experiments, both L1 and L8 were  
245 evaluated for their capability to produce siderophore by CAS screening assay (Schwyn and  
246 Neilands 1987). This assay was used for the screening of siderophore producing microorganisms by  
247 the induction of a colour change of the growth medium: from blue to orange, red, or purple.  
248 Our results showed that L1 and L8 strains produced siderophore, displaying after twelve days of  
249 incubation a colour change of the CAS agar, from blue to red. Moreover, L8 strain induced a more  
250 intense agar colour change and extended halo with respect to L1. Conversely, *M. laxa* was not able  
251 to produce siderophores in solid medium but produced a clear barrage between the common fungal  
252 culture medium (PDA) and the siderophore screening agar (CAS). In fact, CAS agar assay is mainly  
253 useful for the identification of microorganisms capable to produce siderophores in high  
254 concentrations and characterized by a high affinity for iron (III). Most likely, the assay was not so  
255 sensitive to detect the low siderophores concentration produced by *M. laxa*.  
256 In effect, these results were supported by the quantitative analysis conducted with the  
257 spectrophotometer. Both yeasts were found to produce high levels of chelating molecules (1.2 and  
258 1.4 mg ml<sup>-1</sup>, respectively) whit respect to the pathogen that seemed instead to be a poor siderophores

259 producer (0.64 mg ml<sup>-1</sup>). Probably siderophores role vary among different fungal pathosystems  
260 (Chen et al., 2013) and according to the strain intrinsic characteristics. In fact, the importance of  
261 siderophores for fungal pathogenesis was firstly demonstrated for the pathogens such as  
262 *Cochliobulus heterostrophus* (Lee et al., 2005; Oide et al., 2006), *Alternaria* spp. (Chen et al.,  
263 2013), *Ustilago maydis* (Mei et al., 1993) and for the human pathogen *Aspergillus fumigatus*  
264 (Hissen et al., 2005; Schrettl et al., 2007; Chen et al., 2013).

265 In our study, the siderophore production by the yeasts was not affected by the presence of *M. laxa*,  
266 probably for the nutritional and biochemical preferences of the fungus. According to these results,  
267 we can suppose that *M. laxa* probably mainly compete for other nutrients rather than for iron; on the  
268 other side, *A. pullulans* strains L1 and L8 showed a high effective antagonistic action against *M.*  
269 *laxa* in presence of iron (10 µg l<sup>-1</sup> FeCl<sub>3</sub>), displaying a great capability to create critical conditions  
270 for the pathogen growth.

271 As other microorganisms (Droby et al., 1989, Bencheqroun et al., 2006) both L1 and L8, reduced  
272 their antagonistic efficacy with the addition of high concentrations of exogenous nutrients, e.g. with  
273 20 µg l<sup>-1</sup> of FeCl<sub>3</sub> both in *in vitro* and in *in vivo* assays.

274 In fact, the highest reduction of conidial germination, brown rot severity and incidence (data not  
275 shown) in fruit was revealed in presence of lower iron concentrations, confirming that both strains  
276 displayed their antagonistic efficacy when exogenous nutrients were scarce (Calvente et al., 1999,  
277 Calvo et al., 2008, Sanz Ferramola et al., 2013). Plante and Labbè (2019) showed how in  
278 *Schizosaccaromyces pombe* the siderophore synthetase expression remains upregulated under low  
279 iron conditions.

280 According with previous observations on nutritional requirements of *Monilinia* spp. germination  
281 (Byrde and Willetts 1977), in the present research, the germination rate of the pathogen conidia  
282 resulted low when grown in minimal medium. This confirmed that the target pathogen is nutrient-  
283 dependent and needs an adequate amount of supplements for its growth: both conidial germination  
284 and hyphal development (Bencheqroun et al., 2006, Di Francesco et al., 2017). Moreover, the two

285 BCAs notably reduced *M. laxa* germ tube elongation: a fungal characteristic directly related to the  
286 pathogen aggressiveness (Fujii 2004). In fact, one critical step of germination is the germ tube  
287 elongation that can be critically blocked by iron deficiency (Plante and Labbè 2019), in this case  
288 caused by the antagonistic action of both L1 and L8 to compete for nutrients.

289 In conclusion, *A. pullulans* L1 and L8 strains can compete with *M. laxa* for nutrients such as amino  
290 acids as previously reported by Di Francesco et al. (2017, 2018a) and in the present study for iron, a  
291 vital element for cell proliferation and survival for almost all organisms. Therefore, the ability of *A.*  
292 *pullulans* strains L1 and L8 to acquire iron can counteract the expression of fungal pathogen genes  
293 required for the production of siderophores and virulence in fruits (Chen et al., 2013).

294 This information can be very important to exploit L1 and L8 activity in a commercial formulation  
295 characterized by a right and efficient composition. More investigations are necessary to isolate and  
296 characterize siderophores produced by both yeast strains and possibly verify their environmental  
297 involvement in absorption and metabolization of metals. Nevertheless, in our study the strong  
298 efficacy exerted by L1 and L8 strains can be attributed most likely to a synergic activity of different  
299 mechanisms of action.

300

### 301 **Conflict of Interest**

302 The authors declare that they have no conflict of interests.

303

### 304 **Authors contribution**

305 Dr. Alessandra Di Francesco conducted all the experiments, interpreted data, and drafted the  
306 manuscript. Prof. Elena Baraldi coordinated the experimental plan, has corrected, and approved the  
307 final draft of the manuscript.

308

### 309 **References**

310

- 311 Bautista-Rosales, P. U., Calderon-Santoya, M., Servin-Villegas, R., Ochoa-Alvarez, N. A.,  
312 Ragazzo-Sanchez, J. A., 2013. Action mechanisms of the yeast *Meyerozyma caribbica* for the  
313 control of the phytopathogen *Colletotrichum gloesporioides* in mangoes. *Biol. Control* 65, 293-301  
314
- 315 Bencheqroun, S. K., Bajji, M., Massart, S., EI-Jaafari, S., Jijakli, M. H., 2006. *In vitro* and *in situ*  
316 study of postharvest apple blue mould biocontrol by *Aureobasidium pullulans*: evidence for the  
317 involvement of competition for nutrients. *Postharvest Biol. Technol.* 46, 28–135  
318
- 319 Byrde, R. J. W., Willetts, H. J., 1977. *The Brown Rot Fungi of Fruit: Their Biology and Control*.  
320 Pergamon Press, Oxford, **United Kingdom**  
321
- 322 Calvente, V., Benuzzi, D., Sanz de Tosetti, M. I., 1999. Antagonistic action of siderophores from  
323 *Rhodotorula glutinis* upon the postharvest pathogen *Penicillium expansum*. *Int. Biodet. Biodeg. J.*  
324 43,167-172  
325
- 326 Calvente, V., de Orellano, M. E., Sansone, G., Benuzzi, D., Sanz de Tosetti, M. I., **2001a**. Effect of  
327 nitrogen source and pH on siderophore production by *Rhodotorula* strains and their application to  
328 biocontrol of phytopathogenic moulds. *J. Ind. Microbiol. Biotechnol.* 26, 226-229  
329
- 330 Calvente, V., Orellano, M. E., Sansone, G., Sanz de Tosetti, M. I., **2001b**. A simple agar plate assay  
331 for screening siderophore producer yeasts. *J. Microbiol. Meth.* 47, 273-279  
332
- 333 Calvo, J., Calvente, V., Orellano, M. E., Benuzzi, D., Sanz de Tosetti, M. I., 2008. Control of  
334 *Penicillium expansum* and *Botrytis cinerea* on apple fruit by mixture of bacteria and yeast. *Food*  
335 *Bioprocess Tech.* 3, 644-650  
336

- 337 Chanchaichaovivat, A., Panijpan, B., Ruenwongsa, P., 2008. Putative modes of action of *Pichia*  
338 *guilliermondii* strain R13 in controlling chilli anthracnose after harvest. *Biol. Control* 47, 207-215  
339
- 340 Chen, L. H., Lin, C. H., Chung, K. R., 2013. A nonribosomal peptide synthetase mediates  
341 siderophore production and virulence in the citrus fungal pathogen *Alternaria alternata*. *Mol. Plant*  
342 *Pathol.* 14, 497–505  
343
- 344 Di Francesco A., Martini C., Mari M., 2016. Biological control of postharvest diseases by microbial  
345 antagonists: how many mechanisms of action? *Eur. J. Plant Pathol.* 145, 711-717  
346
- 347 Di Francesco, A., Ugolini, L., D'aquino, S., Pagnotta, E., Mari, M., 2017. Biocontrol of *Monilinia*  
348 *laxa* by *Aureobasidium pullulans* strains: Insights on competition for nutrients and space. *Int. J.*  
349 *Food Microbiol.* 248, 32-38  
350
- 351 Di Francesco, A., Mari, M., Ugolini, L., Baraldi, E., 2018a. Effect of *Aureobasidium pullulans*  
352 strains against *Botrytis cinerea* on kiwifruit during storage and on fruit nutritional composition.  
353 *Food Microbiol.* 72, 67-72  
354
- 355 Di Francesco, A., Calassanzio, M., Ratti, C., Folchi, A., Baraldi, E., 2018b. Molecular  
356 characterization of the two postharvest biological control agents *Aureobasidium pullulans* L1 and  
357 L8. *Biol. Control* 123, 53-59  
358
- 359 Droby, S., Chalutz, E., Wilson, C. L., Wisniewski, M., 1989. Characterization of the biocontrol  
360 activity of *Debaryomyces hansenii* in the control of *Penicillium digitatum* on grape fruit. *Can. J.*  
361 *Microbiol.* 35, 794  
362



- 363 Druvefors, U. A., Passoth, V., Schnurer, J., 2005. Nutrient effects on biocontrol of *Penicillium*  
364 *roqueforti* by *Pichia anomala* J121 during airtight storage of wheat. J. Appl. Environ. Microbiol.  
365 71, 1865-1869  
366
- 367 Elad, Y., Chet, I., 1987. The role of competition for nutrients in biocontrol of *Pythium* damping off  
368 by bacteria. Phytopathology 77, 190-195  
369
- 370 Fujii, T. 2004. Germination of spore and decomposition of apple fruit tissue by hypha in  
371 *Penicillium expansum*. Mycotoxins 54, 2  
372
- 373 Hamdan, H., Weller, D. M., Thomashow, L. S., 1991. Relative importance of fluorescent  
374 siderophores and other factors in biological control of *Gaeumannomyces graminis* var. *Tritici* by  
375 *Pseudomonas fluorescens* 2-79 and M4-80R. J. Appl. Environ. Microbiol. 57, 3270-3277  
376
- 377 Hider RC, Kong X, 2010. Chemistry and biology of siderophores. Natural product reports 27, 637-  
378 657  
379
- 380 Hissen, A. H. T., Wan, A. N. C., Warwas, M. L., Pinto, L. J., Moore, M. M., 2005. The *Aspergillus*  
381 *fumigatus* siderophore biosynthetic gene *sidA*, encoding L-ornithine N5-oxygenase, is required for  
382 virulence. Infect. Immun. 73, 5493-5503  
383
- 384 Ho, Y. N., Lee, H. J., Hsieh, C. T., Peng, C. C., Yang, Y. L., 2019. Chemistry and biology of  
385 salicyl-capped siderophores. Nat. Prod. Chem. 59, 431-49  
386

- 387 Janisiewicz, W. J., Tworkoski, T. J., Sharer, C., 2000. Characterizing the mechanism of biological  
388 control of postharvest diseases on fruits with a simple method to study competition for nutrients.  
389 *Phytopathology* 90, 1196–1200  
390
- 391 Johnson, L., 2008. Iron and siderophores in fungal-host interactions. *Mycol. Res.* 112, 170–183  
392
- 393 Larena, I., Torres, R., De Cal, A., Linan, M., Melgarejo, P., Domenichini, P., Bellini, A., Mandrin,  
394 J.F., Lichou, J., Ochoa de Eribe, X., Usall J., 2005. Biological control of postharvest brown rot  
395 (*Monilinia* spp.) of peaches by field applications of *Epicoccum nigrum*. *Biol. Control* 32, 305-310  
396
- 397 Lee, B. N., Kroken, S., Chou, D. Y. T., Robbertse, B., Yoder, O. C., Turgeon, B. G., 2005.  
398 Functional analysis of all nonribosomal peptide synthetases in *Cochliobolus heterostrophus* reveals  
399 a factor, NPS6, involved in virulence and resistance to oxidative stress. *Eukaryotic Cell* 4, 545–555  
400
- 401 Li, C., Zhu, L., Pan, D., Li, S., Xiao, H., Zhang, Z., Shen, X., Wang, Y., Long, M., 2018.  
402 Siderophore-mediated iron acquisition enhances resistance to oxidative and aromatic compound  
403 stress in JMP134. *Appl. Environ. Microbiol.* 85  
404
- 405 Liu, X., Fang, W., Liu, L., Yu, T., Lou, B., Zheng, X., 2010. Biological control of postharvest sour  
406 rot of citrus by two antagonistic yeasts. *Lett. Appl. Microbiol.* 51, 30–35  
407
- 408 Liu, P., Luo, L., Long, C., 2013. Characterization of competition for nutrients in the biocontrol of  
409 *Penicillium italicum* by *Kloeckera apiculata*. *Biol. Control* 67, 157-162  
410

- 411 Mari, M., Martini, C., Guidarelli, M., Neri, F., 2012. Postharvest biocontrol of *Monilinia laxa*,  
412 *Monilinia fructicola* and *Monilinia fructigena* on stone fruit by two *Aureobasidium pullulans*  
413 strains. *Biol. Control* 60, 132-140
- 414
- 415 Martini, C., Guidarelli, M., Di Francesco, A., Ceredi, G., Mari, M., 2016. Characterization of  
416 thiophanate methyl resistance in Italian *Monilinia fructicola* isolates. *J. Plant Pathol.* 98 (3)
- 417
- 418 McLoughlin, T. J., Quinn, J. P., Bettermann, A., Bookland, R., 1992. *Pseudomonas cepacia*  
419 suppression of sunflower wilt fungus and role of antifungal compounds in controlling the disease.  
420 *Appl. Environ. Microbiol.* 58, 1760–1763
- 421
- 422 Mei, B. G., Budde, A. D., Leong, S. A., 1993. Sid1, a gene initiating siderophore biosynthesis in  
423 *Ustilago maydis*: molecular characterization, regulation by iron, and role in phytopathogenicity.  
424 *Proc. Natl. Acad. Sci. USA* 90, 903–90
- 425
- 426 Mekbib, S. B., Regnier, T. J. C., Korsten, L., 2011. Efficacy and mode of action of yeast  
427 antagonists for control of *Penicillium digitatum* in oranges. *Trop. Plant Pathol.* 36, 233–240
- 428
- 429 Miethke, M., Marahiel, M. A., 2007. Siderophore-based iron acquisition and pathogen control.  
430 *Microb. Mol. Biology Rev.* 71, 413-51
- 431
- 432 Oide, S., Moeder, W., Krasnoff, S., Gibson, D., Haas, H., Yoshioka, K., Turgeon, B. G., 2006.  
433 NPS6, encoding a nonribosomal peptide synthetase involved in siderophore-mediated iron  
434 metabolism, is a conserved virulence determinant of plant pathogenic ascomycetes. *Plant Cell* 18,  
435 2836–2853

436

437 Parafati, L., Vitale, A., Restuccia, C., Cirvilleri, G., 2015. Biocontrol ability and action mechanism  
438 of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape.  
439 Food Microbiol. 47, 85-92

440

441 Plante, S., Labbè, S., 2019. Spore germination requires ferrichrome biosynthesis and the  
442 siderophore transporter Str1 in *Schizosaccharomyces pombe*. Genetics 211, 893-911

443

444 Raaska, L., Mattila-Sandholm, T., 1995. Effects of iron level on the antagonistic action of  
445 siderophores from non-pathogenic *Staphylococcus* spp. J. Microbiol. Biotechnol. 15, 480-485

446

447 Riquelme, M., 1996. Fungal siderophores in plant-microbe interactions. Microbiology 12, 537–546

448

449 Sanz Ferramola, M. I., Benuzzi, D., Calvente, V., Calvo, J., Sansone, G., Cerutti, S., Raba, J., 2013.  
450 The use of siderophores for improving the control of postharvest diseases in stored fruits and  
451 vegetables. Microbial pathogens and strategies for combating them: science, technology and  
452 education (A. Méndez-Vilas, Ed.), pp. 1385-1394

453

454 Saravanakumar, D., Ciavorella, A., Spadaro, D., Garibaldi, A., Gullino, M. L., 2008.  
455 *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and  
456 *Penicillium expansum* in apple through iron depletion. Postharvest Biol. Technol. 49, 121-128

457

458 Schrettl, M., Bignell, E., Kragl, C., Sabiha, Y., Loss, O., Eisendle, M., Wallner, A., Arst, H. N.,  
459 Haynes, K., Haas, H., 2007. Distinct roles for intra- and extra-cellular siderophores during  
460 *Aspergillus fumigatus* infection. PLoS Pathogen 28, 1195-207

461

- 462 Schwyn, B., Neilands, J. B., 1987. Universal chemical assay for the detection and determination of  
463 siderophores. *Anal. Biochem.* 160, 47-56  
464
- 465 Wang, W., Chi, Z., Liu, G., Buzdar, M. A., Chi, Z., Gu, Q., 2009. Chemical and biological  
466 characterization of siderophore produced by the marine-derived *Aureobasidium pullulans* HN6.2  
467 and its antibacterial activity. *Biometals* 22, 965–972  
468
- 469 Xu, B., Zhang, H., Chen, K., Xu, Q., Yao, Y., Gao, H., 2013. Biocontrol of postharvest *Rhizopus*  
470 decay of peaches with *Pichia caribbica*. *Curr. Microbiol.* 67, 255-261  
471
- 472 Yu, X., Ai, C., Xin, L., Zhou, G., 2011. The siderophore producing bacterium, *Bacillus subtilis*  
473 CAS15, has a biocontrol effect on *Fusarium* wilt and promotes the growth of pepper. *Eur. J. Soil*  
474 *Biol.* 47, 138


475

476

477 Figure 1

478 CAS assay performed with *Aureobasidium pullulans* L1, L8 strains and *Monilinia laxa* (isolate

479 ML4).



480

481

482


483

484

485

486 Figure 2

487 Quantitative ( $\text{mg ml}^{-1}$ ) determination of siderophore at 440 nm by spectrophotometer, on488 deferoxamine mesylate standard curve, of *Aureobasidium pullulans* L1 and L8 strains and489 *Monilinia laxa* (M), alone and co-cultured. Data are the means of two independent490 experiments  $\pm$  standard error, each consisting of three flasks per treatment. Different letters indicate491 significant differences at  $\alpha=0.05$  according to Tukey's HSD Test.



492


493

494

495 Figure 3

496 *In vivo* antagonistic effect of L1 and L8 *Aureobasidium pullulans* strains on *Monilinia laxa* in peach  
 497 fruits. Fruits were artificially inoculated with 20  $\mu$ l of yeasts cell suspensions ( $10^8$  cells ml<sup>-1</sup>); after  
 498 air drying they were inoculated with 20  $\mu$ l of *M. laxa* conidia suspension ( $10^5$  conidia ml<sup>-1</sup>) and  
 499 finally with an iron suspension (0, 5, 10 and 20  $\mu$ g ml<sup>-1</sup> FeCl<sub>3</sub>) (0, 5, 10, 20). Control consisted of  
 500 peaches inoculated with water instead of yeasts suspensions. Data are the means of two independent  
 501 experiments  $\pm$  standard error, each consisting of 45 fruit per treatment. For each treatment group  
 502 (Control, L1, and L8) different letters indicate significant differences at  $\alpha = 0.05$  according to  
 503 *Tukey's* HSD Test.

504



505

506

507

508

509

510

511 Table 1

512 CAS assay for analysis of siderophore production by *Aureobasidium pullulans* (L1 and L8) and  
 513 *Monilinia laxa* (isolate ML4). Data are the means of two independent experiments  $\pm$  standard error,  
 514 each consisting of ten replicates per treatment. Different letters within the same column indicate  
 515 significant differences at  $\alpha = 0.05$  according to Tukey's HSD Test.

516

517

<i>Microorganism</i>	<i>Growth time</i> (days)*	<i>Color change</i>	<i>CAS reaction</i> **
<i>Monilinia laxa</i>	10 $\pm$ 0.4b	-	0 $\pm$ 0.0a
L1	12 $\pm$ 0.0a	dark orange	20 $\pm$ 1.9b
L8	12 $\pm$ 0.0a	dark orange	27 $\pm$ 1.5c



518

519 \*Days required for the fungal mycelium to cover the non-CAS half of the plate.

520 \*\* mm of advance of colour change front in the CAS blue agar after three weeks of incubation.

521

522

523 Table 2

524 Mycelium diameter (mm) of *Monilinia laxa* growth in dual culture with *Aureobasidium pullulans*525 L1 and L8 strains in minimal medium agar plates amended with 0, 5, 10 and 20  $\mu\text{g ml}^{-1}$   $\text{FeCl}_3$  and

526 incubated at 25 °C for 5 days. Control plates were inoculated only with pathogen plug (6 mm).

527 Data are the means of two independent experiments  $\pm$  standard error, each consisting of five528 replicates per treatment. Within the same  $\text{FeCl}_3$  treatment (column), the same lower-case letters529 represent no significant differences at  $\alpha = 0.05$  according to Tukey's HSD Test. Within the same

530 yeast strains or pathogen isolate (control) (row), the same upper-case letters represent no significant

531 differences at  $\alpha = 0.05$  according to Tukey's HSD Test.

532

533

	Treatment ( $\text{FeCl}_3$ )			
	0 $\mu\text{g ml}^{-1}$	5 $\mu\text{g ml}^{-1}$	10 $\mu\text{g ml}^{-1}$	20 $\mu\text{g ml}^{-1}$
Control	60.0 $\pm$ 0.6cA	60.0 $\pm$ b1.2bA	63.0c $\pm$ 1.0cB	61.0 $\pm$ 1.5cAB
L1	35.2 $\pm$ 1.9bB	35.0a $\pm$ 0.7aB	34.7b $\pm$ 1.3bA	34.9 $\pm$ 1.1bA
L8	34.0 $\pm$ 1.1aB	34.4a $\pm$ 1.0aB	33.0a $\pm$ 0.8aA	33.6 $\pm$ 0.8aAB

534

535

536

537

538 Table 3

539 Effect of *Aureobasidium pullulans* (L1 and L8) on conidia germination (%) of *Monilinia laxa*540 grown in culture medium (minimal medium) amended with 5, 10 or 20  $\mu\text{g ml}^{-1}$   $\text{FeCl}_3$  and incubated

541 at 25°C for 6 hours. Data are the means of two independent experiments  $\pm$  standard error, each  
 542 consisting of 90 conidia per treatment. Within the same FeCl<sub>3</sub> treatment (column), the same lower-  
 543 case letters represent no significant differences at  $\alpha = 0.05$  according to *Tukey's HSD Test*.

544 Within the pathogen isolate (control) and yeast strains (row), the same upper-case letters represent  
 545 no significant differences at  $\alpha = 0.05$  according to *Tukey's HSD Test*.

546

547

548

	Treatment (FeCl <sub>3</sub> )			
	0 $\mu\text{g ml}^{-1}$	5 $\mu\text{g ml}^{-1}$	10 $\mu\text{g ml}^{-1}$	20 $\mu\text{g ml}^{-1}$
Control	40 $\pm$ 1.3cA	41 $\pm$ 1.8bA	48 $\pm$ 0.8cC	43 $\pm$ 1.5bB
L1	20 $\pm$ 2.1bB	19 $\pm$ 1.7aB	18 $\pm$ 1.4bA	28 $\pm$ 2.1aC
L8	19 $\pm$ 0.9aB	19 $\pm$ 1.5aB	15 $\pm$ 1.2aA	29 $\pm$ 1.8aC

549

550

551

552 Table 4

553 Effect of *Aureobasidium pullulans* (L1 and L8) on *Monilinia laxa* conidia germ tube elongation  
 554 ( $\mu\text{m}$ ) grown in liquid culture medium (minimal medium) amended with 0, 5, 10 and 20  $\mu\text{g ml}^{-1}$   
 555 FeCl<sub>3</sub> at 25 °C for 6 hours. Data are the means of two independent experiments  $\pm$  standard error,  
 556 each consisting of 90 conidia per treatment. Within the same FeCl<sub>3</sub> treatment (column), the same  
 557 lower-case letters represent no significant differences significant differences at  $\alpha = 0.05$  according  
 558 to *Tukey's HSD Test*. Within the pathogen isolate (control) and yeast strains (row), the same upper-  
 559 case letters represent no significant differences significant differences at  $\alpha = 0.05$  according to  
 560 *Tukey's HSD Test*.

561

	Treatment (FeCl <sub>3</sub> )			
	0µg ml <sup>-1</sup>	5 µg ml <sup>-1</sup>	10 µg ml <sup>-1</sup>	20 µg ml <sup>-1</sup>
Control	16.2±1.3cA	16.3±1.6cA	18.2±2.1cB	16.5±1.8bA
L1	7.7±2.1aB	6.6±1.8bA	6.2±2.3bA	8.1±2.1aC
L8	8.3±1.5bC	6.0±1.6aB	5.7±3.5aA	8.2±1.8aC

562

563

564 **Conflict of Interest**

565 The authors declare that they have no conflict of interests.

566

567 **Authors contribution**

568 Dr. Alessandra Di Francesco conducted all the experiments, interpreted data, and drafted the  
569 manuscript. Prof. Elena Baraldi coordinated the experimental plan, has corrected, and approved the  
570 final draft of the manuscript.

571

572

573 **Highlights**

574

575 - Competition for nutrients is one of the most important mechanism of action for BCAs

576 - Iron plays a role in BCAs modes of action through the production of siderophores

577 - Siderophores can inhibit the growth of certain pathogenic fungi

578 - *Aureobasidium pullulans* L1 and L8 strains compete for iron against *Monilinia laxa*

579

580