

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Effects of 17 performance, carcass and raw ham quality parameters on ham weight loss at first salting in heavy pigs, a meat quality indicator for the production of high quality dry-cured hams

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Dall'Olio, S., Aboagye, G., Nanni Costa, L., Gallo, M., Fontanesi, L. (2020). Effects of 17 performance, carcass and raw ham quality parameters on ham weight loss at first salting in heavy pigs, a meat quality indicator for the production of high quality dry-cured hams. MEAT SCIENCE, 162(April 2020), 1-7 [10.1016/j.meatsci.2019.108012].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/777175> since: 2020-11-02

*Published:*

DOI: <http://doi.org/10.1016/j.meatsci.2019.108012>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

**Effects of 17 performance, carcass and raw ham quality parameters on ham weight loss at first salting in heavy pigs, a meat quality indicator for the production of high quality dry-cured hams**

Stefania Dal 'Olio<sup>a</sup>, Gizella Aboagye<sup>a</sup>, Leonardo Nanni Costa<sup>a</sup>, Maurizio Gallo<sup>b</sup>, Luca Fontanesi<sup>a</sup>

<sup>a</sup> Department of Agricultural and Food Sciences (DISTAL), Division of Animal Sciences,

University of Bologna, Viale Fanin 46, 40127 Bologna, Italy

<sup>b</sup> Associazione Nazionale Allevatori Suini, Via Nizza 53, 00198 Roma, Italy

Corresponding author: tel. +39 051 2096531

E-mail: [stefania.dallolio@unibo.it](mailto:stefania.dallolio@unibo.it) (S. Dal 'Olio)

**Running title:** Ham weight loss at first salting and carcass and ham quality traits

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

### **Highlights**

- Ham weight loss at first salting (HWLFS) was evaluated in Italian Large White pigs.
- HWLFS was affected by slaughter day and visible intermuscular fat of trimmed hams.
- Residual correlations between HWLFS and 17 traits were determined.
- HWLFS could be predicted by several parameters.
- Backfat thickness, lean cuts and pH<sub>24h</sub> were the most important predictors.

## Abstract

Ham weight loss at first salting (HWLFS) is a meat quality parameter used to assess the suitability of the hams for salting and seasoning. The relationships between HWLFS and 17 performance, carcass and raw ham quality parameters were investigated in 260 performance tested Italian Large White heavy pigs. HWLFS was affected by slaughter day and visible intermuscular fat of trimmed hams whereas sex did not affect its variability. Residual correlations of HWLFS with backfat thickness (BFT;  $r = -0.51$ ) and lean cuts (LC;  $r = +0.51$ ) were stronger than with ham weight at trimming ( $r = +0.40$ ) and after first salting ( $r = +0.37$ ). Significant correlations of HWLFS with fresh ham quality traits ranged from  $+0.16$  to  $-0.25$ . BFT, LC and  $\text{pH}_{24\text{h}}$  were the main predictors of HWLFS in the regression model. Results from this study indicated that higher ham fat coverage and  $\text{pH}_{24\text{h}}$  and lower LC could reduce HWLFS of green hams for Protected Designation of Origin products.

**Keyword:** Heavy pig; Italian Large White; Ham quality; Ham weight loss at first salting; Phenotypic correlations.

## 1. Introduction

The Italian pig industry is mainly oriented towards the production of heavy carcasses to provide suitable raw meat material for typical processed products, including dry-cured hams of the Protected Designation of Origin (PDO) system. Italian heavy pigs are slaughtered at an average live weight of 160 kg ( $\pm 10\%$ ) and when they are at least nine months old (EEC, 1992). This long period is essential to obtain raw hams of the right weight, muscle maturity and subcutaneous fat (15-30 mm of thickness) that minimize seasoning losses of the hams over the processing period (Bosi & Russo, 2004). PDO dry-cured hams processing technology is standardized and involves only the addition of sea salt (which acts as a bacteriostatic agent, it reduces water activity and plays a role on

69 taste of the final product), the control of ambient conditions (i.e. temperature, relative humidity and  
70 air flow) and the duration of the ripening (Pagliarini et al., 2016; Parolari, 1996). Briefly, the  
71 processing of PDO dry-cured ham, such as Parma ham, includes the following steps: i) cooling, by  
72 keeping legs in cold rooms for 24 hours after slaughter to attain uniform temperature; ii) trimming,  
73 the procedure that removes external fat and rind needed to obtain the typical ham shape and to help  
74 the salting phase; iii) two salting steps (the first lasting seven days in a salting room; after which  
75 salt residuals are removed, legs are re-salted and stored for 15-18 days in another salting rooms); iv)  
76 resting for 60-90 days, to obtain a homogeneous salt distribution inside the leg muscles; v) washing  
77 from the excess of salt; and then drying; vi) pre-maturation period; vii) smearing or greasing; and  
78 then vii) final maturation (Pagliarini et al., 2016).

79 PDO dry cured ham production requires a processing time of at least 12 months (for PDO  
80 Parma ham) or 13 months (for PDO San Daniele ham). The key changes that occur during the  
81 processing period are related to water loss of the legs, salt intake, lipolysis and proteolysis (Čandek-  
82 Potokar & Škrlep, 2012). When the curing process is standardized, the final quality of dry cured  
83 hams (i.e. typical texture and flavour sensorial characteristics and processing yield) is primarily  
84 determined by the intrinsic characteristics of the fresh hams (Bosi & Russo, 2004). Sensory  
85 properties developed by the hams are mainly due to physical, and biochemical changes caused by  
86 endogenous proteolytic and lipolytic muscle enzymes that work during the drying and  
87 ripening/maturation phases (Toldrá, Aristoy, & Flores, 2000; Toldrá & Flores, 1998; Zhou, et al.,  
88 2019c,b).

90 To produce suitable legs to maximize dry-cured ham processing yield, breeding programmes  
91 for Italian heavy pigs, beside performance traits (average daily gain and feed:gain ratio), carcass  
92 traits (backfat thickness, weight of neck and loin and weight of hams), and meat quality (visible  
93 intermuscular fat of trimmed hams), includes ham weight loss after the first salting step (HWLFS).  
94 This latter characteristic, that is determined just 8-9 days after slaughter, is a parameter used in the  
95 pig selection programmes with the aim to minimize the total weight loss of the hams up to the end  
96

of the seasoning period. The weight losses at these two periods (at first salting and at the end of the processing period) have high phenotypic and genetic correlations (0.57-0.73 and 0.49-0.73, respectively (Bosi & Russo, 2004). Phenotypic correlations between HWLFS and traits related to performances, carcass and ham quality traits have been poorly investigated in heavy pig breeds (Bosi & Russo, 2004; Russo, Nanni Costa, Lo Fiego, & De Grossi, 1993).

The objectives of this study are 1) to investigate the phenotypic relationships between HWLFS and 17 parameters related to production performance, carcass and ham muscles in Italian Large White heavy pigs, 2) to determine the predictive parameters of HWLFS using a stepwise multiple regression analysis.

## **2. Material and methods**

Animal care and slaughter of the animals used in this study were performed in compliance with the European rules [Council Regulation (EC) No. 1/2005 and Council Regulation (EC) No. 1099/2009] on the protection of animals during transport and related operations and at the time of the killing. All slaughter procedures were monitored by the veterinary team appointed by the Italian Ministry of Health and sampling occurred with the permission of the National Pig Breeder Association (ANAS; <http://www.anas.it>).

### **2.1. Animals**

A total of 260 Italian Large White pigs (172 entire gilts and 88 castrated males obtained from 79 sires) were involved in this study. These pigs were from triplets of siblings of the same litter (two females and one castrated male) individually performance tested at the central station of ANAS for the genetic evaluation of a candidate boar from the same litter (sib-testing). The test period began when piglets were approximately 30 kg live weight and it ended when the pigs were slaughtered at about 8 months following the rules of the test station.

All pigs were genotyped for the ryanodine receptor 1 (*RYR1*) g.1843C>T mutation (Fujii et al., 1991), causing the Pale Soft and Exudative meat defect and for the p.R200Q mutation of the protein kinase AMP-activated non-catalytic subunit gamma 3 (*PRKAG3*) gene (Milan et al., 2000), causing the acid meat defect. Genotyping was carried out as previously described by Fontanesi et al. (2008).

## **2.2. In vivo performances of the animals**

Nutritive level of the animals in the performance testing period till slaughter was *quasi ad libitum*, meaning that about 60% of the pigs were able to ingest the entire supplied ration. The feeding regime and the diets are already described by Fontanesi et al. (2010). Feed intake was individually recorded daily and live weight was measured every two weeks. The data obtained were then used to calculate average daily gain (ADG, kg/d) and feed/gain ratio (FGR) for each pig.

## **2.3. Slaughtering and carcass traits**

The days before slaughter, the pigs were weighed in the performance station (thereafter referred as slaughter weight; SW, kg). Animals at the end of test were transported to a commercial slaughterhouse (six slaughtering days) located 24.5 km from the test station (the full sibs were slaughtered in two different slaughter days). After unloading, pigs were immediately stunned and bled in a lying position. Carcasses were then divided in commercial cuts. Within 3 h after slaughter, backfat thickness (BFT, mm) at the level of *Gluteus medius* muscle, weight of necks and loins (referred as lean cuts, LC; kg), and weight of raw hams (HW, kg) were measured in the commercial abattoir.

## **2.4. Ham technological traits**

At a single processing plant, all raw hams were chilled for 24 h and then trimmed. The two hams of each pig were re-weighed at the end of the trimming line (mean trimmed ham weight;

158 HW\_TR, kg), and after six days from the first salting step (HW\_FS, kg). Mean ham weight loss  
159 (HWL) after trimming [HWLTR = (HW - HW\_TR), in kg] and after first salting [HWLFS =  
160 (HW\_TR - HW\_FS), in kg] were then calculated.

161  
162  
163 **2.5. Raw ham quality traits measurements**  
164

165 Visible intermuscular fat (VIF) is the level of fat deposition (infiltration) subjectively  
166 evaluated on muscles of trimmed hams (which includes mainly *semitendinosus*, *semimembranosus*,  
167 *biceps femoris*, *gastrocnemius* and *gracilis* muscles) by a trained personnel with the scoring system  
168 described by Fontanesi et al. (2017). VIF in this trial was coded as follows: 0, when exposed muscle  
169 area of both legs of the same pig did not show VIF; 1, when muscles from only one leg showed  
170 VIF; 2, when muscles from both legs showed VIF.

171 Subsequent measures on the hams were assessed on the *Semimembranosus* muscle (SM). On  
172 the slaughter line, pH was measured at 2 h *post mortem* (pH<sub>2h</sub>), and at 24 h *post mortem* (pH<sub>24h</sub>)  
173 using a portable Crison pH-meter equipped with an Ingold Xerolite glass electrode (Model 5232,  
174 Crison, Modena, Italy) and an automatic temperature compensation probe.

175 At 2 h *post mortem*, samples taken from SM were snap frozen in liquid nitrogen and freeze-  
176 dried for the determination of muscle carbohydrates as described by Fontanesi et al. (2008). Briefly,  
177 content of lactate was determined by UV method using the L-Lactic acid BioAnalysis kit  
178 (Boehringer Mannheim/R-Biopharma, R-Biopharma GmbH, Darmstadt, Germany). Glycogen (first  
179 degraded to glucose with amyloglucosidase from *Aspergillus niger*), glucose and glucose-6-  
180 phosphate were determined using the D-Glucose Enzymatic BioAnalysis kit (Boehringer Mannheim/R-  
181 Biopharma, Milan, Italy) and expressed as  $\mu\text{mol}$  of lactic acid equivalent per gram of fresh muscle  
182 ( $\mu\text{mol/g}$ ). Glycolytic potential (GP), a measure of all components present in the muscle that can  
183 potentially be converted into lactate during post-mortem glycolysis, was calculated as  $2[\text{glycogen} +$   
184  $\text{glucose} + \text{glucose-6-phosphate}] + [\text{lactate}]$  and expressed as  $\mu\text{mol/g}$  (Monin &  
185 Sellier, 1985).



Activity of the endopeptidase cathepsin B activity (Catb activity) was analyzed on SM samples collected at 24 h *post mortem*. Activity was measured using N-CBZ-L arginyl-L-arginine as the fluorescent substrate (expressed as nmol of 7-amino-4-methylcoumarin released/min/g muscle; nmol AMC/min/g. Parolari, Virgili, & Schivazappa, 1994; Russo et al., 2000; Schivazappa et al., 2002).

## 2.6. Statistical analyses

Descriptive statistics (mean, standard deviation, minimum and maximum values, and coefficient of variation) were calculated using the PROC MEANS procedure of Statistical Analysis System (SAS Inst. INC., Cary, NC, USA, version 9.4.).

A multivariate analysis of variance was performed using the PROC GLM procedure of SAS. The model included sex (castrated males and gilts) and VIF (three levels: 0, 1 and 2) as fixed effects and slaughter day (six levels) as random effect. The least square means (LSM) and standard errors (SE) were compared using the PDIF option. Residual correlation coefficients were used to determine the relationships between traits and were graphically displayed as heatmap using the *gplots* package in R environment (R Core Team, 2018).

The observations of HWLFS were clustered using the FASTCLUS procedure in SAS. Two classes of HWLFS referred as low ( $190 \pm 21$  g, range from 135 g to 221 g) and high ( $252 \pm 24$  g, range from 224 g to 317 g) were defined. In this procedure, observations that were very close to each other were assigned to the same cluster by an algorithm for minimizing the sum of squared distances from the clustered means. The distribution of HWLFS clusters by VIF (absence and presence of VIF, i.e.  $VIF = 0$  and  $VIF > 0$  including 1 and 2 classes of VIF) were analyzed using the PROC FREQ procedure in SAS with the chi square test option. In this analysis two classes of VIF were used (instead of three) since difference between LSM of HWLFS when  $VIF = 1$  and  $VIF = 2$  was not significant.

Furthermore, the stepwise regression analysis of SAS was performed with the aim of selecting input variables from a set of explanatory variables of performance, carcass, technological and meat characteristics that most contributed to HWLFS. On the basis of the results of variance and cluster analyses, the stepwise regression analysis was carried out for all pigs and separately for pigs grouped based on visible intermuscular fat ( $VIF = 0$  and  $VIF > 0$ ). To assess multicollinearity, the variance inflation factor option was used (values were all  $< 10$ ). The  $P$  value,  $R^2$  (partial and cumulative) and root mean square error (RMSE) were used as parameters to determine the accuracy of the prediction.

### 3. Results and discussion

#### 3.1. Description of the *in vivo* performance, carcass and technological traits

All pigs were reared under the same environmental conditions and feeding regime and subjected to the same pre- and post-slaughter handling, which are all important factors of variations for the quality of hams (Čandek-Potokar & Škrlep, 2012). The pigs did not carry the g.1843T allele of the *RYR1* gene (Fujii et al., 1991) and the p.200Q allele of the *PRKAG3* gene (Milan et al., 2000), the two major negative mutations for meat quality traits in pigs (Čandek-Potokar & Škrlep, 2012; Cherel et al., 2010).

Descriptive statistics of the measured traits are reported in Table 1. The variability in HWLFS as measured by the coefficient of variation ( $CV = 17.4\%$ ) was high. This ham quality parameter was on average equal to  $220 \pm 38$  g, corresponding to a loss percentage in this phase of  $1.75 \pm 0.28\%$ . Similar mean values ( $1.93 \pm 0.59\%$ ), but with higher variability ( $CV = 30.6\%$ ), were previously reported by Schivazappa et al. (2002) in three Italian heavy pig breeds, Italian Duroc, Italian Landrace and Italian Large White, with higher losses in this latter breed (2.16%).

Among the *in vivo* performance and carcass traits, BFT of *gluteus medius* showed the highest variability (mean and SD =  $26.5 \pm 5.4$  mm;  $CV = 20.5\%$ ), although the specific aim of selection of heavy pigs is focused on maintaining a constant value of BFT because an insufficient fat covering

of the ham causes increased seasoning losses and negatively impacts organoleptic characteristics of dry cured hams (Bosi & Russo, 2004; Čandek-Potokar & Škrlep, 2012). Raw ham weight (HW) was on average  $15.4 \pm 1.3$  kg (CV = 8.5%). After trimming the hams (with mean of  $12.6 \pm 1.1$  kg) lost on average  $2.8 \pm 0.4$  kg of their weight in fat and muscle, corresponding to a loss percentage at trimming phase of  $18.3 \pm 1.7\%$ . Similar values of HW and HW\_TR were previously reported in Italian Large White animals (Russo et al., 1993). In heavy pigs for Parma ham production, the trimmed legs should preferably weigh between 12 and 14 kg, but in no case be less than 10 kg (EEC, 1992).

### 3.2 Description of the ham quality traits

For visible intermuscular fat (VIF), defined as the subjective estimation of exposed fat depots between the ham muscle groups, 40.8% of the hams had absence of VIF in both legs while 25.0% showed the presence of VIF in one ham with the remaining 34.2% showing the presence of VIF in both hams.

The pH values were much less variable (pH<sub>2h</sub>: CV = 4.1% and pH<sub>24h</sub>: CV = 3.7%), in agreement with what was previously reported by other authors in both light and heavy pigs (Ramos, Serenius, Stalder, & Rothschild, 2007; Sturaro, Gallo, Noventa, & Carnier, 2008; Ventura et al., 2011). Several samples were not in the range of 5.6 - 6.2 recommended for pH<sub>24h</sub> in fresh hams intended for curing (Arnau, 2004). Although the pigs did not carry the *PRKAG3* p.200Q allele, known to decrease the ultimate pH and pork quality characteristics (Cherel et al., 2010), eight samples exhibited low pH<sub>24h</sub> values (<5.4) compatible with the acid meat defect. This indicated that low muscle pH<sub>24h</sub> can occur in pigs that do not carry the negative allele of the *PRKAG3* gene. Conversely, nine samples had an insufficient decrease in the final pH (pH<sub>24h</sub> above 6.2).

Among the investigated traits, the highest variability of the fresh hams was observed for glycogen (CV = 47.7 %) lactate (CV = 28.2 %) and GP (CV = 22.1%). In pigs that do not carry the negative allele at the *RN* locus (i.e. animals with the recessive genotype at the *PRKAG3* p.R200Q

polymorphic site), the muscle glycogen level is mainly determined by husbandry and feeding practices of the animals, transport conditions, climatic and environmental factors, handling procedures at slaughter and muscle type (England et al., 2016; Larzul, Monin, Sellier & Le Roy, 1998; Scheffler & Gerrard, 2007). Lactate was the main component ( $56.4 \pm 15.9$  %) of GP, as already reported in other studies (Maribo, Støier, & Jørgensen, 1999; Przybylski, Sionek, Jaworska, & Santé-Lhoutellier, 2016). The contribution of lactate in GP reported in this study was lower than those in other investigations. This difference could probably be due to the difference in the sampling time (Maribo et al., 1999; Przybylski et al., 2016). However, according to Maribo et al. (1999) the determination of GP is not affected by the different *post mortem* sampling time.

In our samples, Catb activity ( $1.16 \pm 0.23$  nmol AMC min<sup>-1</sup> g<sup>-1</sup> meat; CV = 19.6%) was lower than what was on average obtained in other Italian heavy pigs (i.e. 1.36 nmol AMC min<sup>-1</sup> g<sup>-1</sup> and CV = 22.8%, Sturaro et al., 2008; 1.55 nmol AMC min<sup>-1</sup> g<sup>-1</sup> and CV = 18%, Virgili, Schivazappa, Parolari, Bordini, & Degni, 1998). Catb activity has been found to exhibit variability depending on genetic type, breeding, age at slaughter, type of muscle, season and, during the processing period of the legs, by several other factors such as salt, temperature, humidity and time of ripening (Arnau, Guerrero, & Sárraga, 1998; Mora, Escudero, & Toldrá, 2016; Sárraga, Gil, & García-Regueiro, 1993; Sturaro, Gallo, Noventa, & Carnier, 2008; Virgili, Parolari, Schivazappa, Bordini, & Borri, 1995; Zhou, et al., 2019a). Sturaro et al. (2008) reported a slight decrease in Catb activity when the carcass weight increased in crossbred of Large White boar line x crossbred sows (1.38 nmol AMC/min/g for lighter carcasses and 1.33 nmol AMC/min/g for heavier carcasses;  $P < 0.05$ ). In addition, Sárraga et al. (1993) identified lower Catb activity in green hams from heavy pigs than light pigs, and they suggested that lower levels in heavy pigs could be attributed to their slower protein turnover.

### **3.2 Effects of sex, slaughter days and visible intermuscular fat**

Investigated traits and parameters that were used in the correlation analyses were adjusted for the effects of three different variation factors (Čandek-Potokar & Škrlep, 2011; Ramos et al., 2007) such as sex, slaughter days and VIF. Probability values of factors included in the model and coefficient of determination (R-square in percentage) are reported in Table 2. Coefficients of determination was higher for HWLFS (32.0%).

Sex did not affect HWLFS as well as ADG, FGR and HW at different stages ( $P > 0.05$ ; Tables 2 and S1). Castrated males had thicker BFT and higher losses at trimming phase compared to gilts. LC from gilts were heavier than males (Tab. S1). The thicker BFT of barrows and the greater yields of lean cuts from gilts are in agreement with previous studies carried out on heavy pigs (Latorre, García-Belenguer, & Ariño, 2008; Schiavon et al., 2015).

Sex did not show any effect on GP, its components, pH measurements and Catb activity of fresh hams ( $P > 0.05$ ), except for glycogen that was higher in castrated males. Other studies reported no effect of sex on several meat quality traits, including some analyzed in this study such as pH<sub>24h</sub>, Catb activity and glycolytic components (Armero et al., 1999; Monin & Sellier, 1985; Peloso, Lopes, Gomide, Guimarães, & Carneiro, 2010; Schivazappa, Virgili, & Parolari, 1992). Nevertheless, Sturaro et al. (2008) identified slightly higher Catb activity in gilts than castrated males.

The visible intermuscular fat of trimmed hams was a significant source of variation for HWLFS ( $P < 0.001$ ; Tables 2 and S2). In particular, the pigs with absence of VIF in both hams (VIF = 0) had significantly higher least square means of HWLFS compared to those of the pigs of the other two classes of VIF (i.e. 1 = presence of VIF in one and 2 = presence of VIF in both hams;  $P < 0.001$ ). Furthermore, VIF affected the variability of other traits such as BFT, LC, HW\_TR, HW\_FS, pH<sub>24h</sub>, glycogen and GP (Table 2 and S2). The pigs with absence of VIF in both hams (VIF = 0) had lower BFT and pH<sub>24h</sub> and higher LC, HW\_TR, HW\_FS and glycogen than the pigs with presence of VIF in at least one ham. The comparison between the least square means of the extreme (0 and 2) classes of VIF resulted significant for GP ( $P = 0.044$ ).

The day of slaughter (which may incorporate differences in pens, batch, and month of slaughtering) was a significant source of variation ( $P < 0.05$ ) for all traits under study except for HW at trimming and at post first salting stages ( $P = > 0.05$ ).

### 3.3 Correlations between HWLFS and *in vivo* performance, carcass and ham technological traits

HWLFS is a quality trait used to assess the aptitude of the meat for salting to obtain legs with optimal yield. Almost all correlations of HWLFS with *in vivo* performance, carcass and technological ham traits were significant (Figure 1. Table S3). As expected, HWLFS was negatively correlated with BFT ( $r = -0.51$ ,  $P < 0.0001$ ), confirming that an adequate level of BFT is beneficial to minimizing HW loss (Bosi & Russo, 2004). This is due to lower water content of adipose tissue than muscular tissue (5-15% vs 70-75%) and to the positive barrier effect of fat on water which drips out from the hams through diffusion and evaporation during the processing period (Bosi & Russo, 2004; Ramos et al., 2007). Conversely, HWLFS was positively correlated with LC ( $r = +0.51$ ;  $P < 0.0001$ ), indicating that the selection for carcass conformation (muscularity) tends to worsen this quality characteristic of the ham. The relationships between HWLFS and HW at different stages were positives and significant ( $r = +0.34/+0.40$ ;  $P < 0.0001$ ). Čandek-Potokar & Škrlep (2011) observed higher daily moisture losses in hams with heavier weight due to their larger contact area for exchange. Other studies showed no or very low correlations between HW and seasoning losses or yield (Čandek-Potokar & Škrlep, 2012; Ramos et al., 2007).

### 3.4 Correlations between HWLFS and ham quality traits

The correlations between HWLFS and fresh meat quality traits were generally low (Figure 1. Table S4). HWLFS was correlated negatively with  $pH_{24h}$  ( $r = -0.25$ ,  $P < 0.0001$ ). This result is in agreement with previous studies reporting higher weight losses during the initial phase of curing in hams exhibiting low  $pH_{24h}$  values (Čandek-Potokar & Škrlep, 2011; Schivazappa et al., 2002). The

correlations of HWLFS with glycogen ( $r = +0.22$ ) and GP ( $r = +0.16$ ) were significant ( $P < 0.05$ ), indicating that higher glycogen and GP resulted in increased weight loss during the first salting process (Nanni Costa, Lo Fiego, Pantano, & Russo, 1998). The coefficient of correlation between HWLFS and GP observed in the current study was similar to those observed for commercial hybrid heavy pigs and Duroc x (Landrace x Large White) crossbreeds (Nanni Costa et al., 1998). These results confirmed the detrimental effect of high GP on ham production and pork quality attributes (Hamilton, Miller, Ellis, McKeith, & Wilson, 2003). HWLFS did not appear to be correlated with Catb activity.

### 3.5 Correlations between raw ham quality traits

Initial pH measured at 2 h was moderately correlated with  $pH_{24h}$  ( $r = +0.58$ , Table S4). The correlation between  $pH_{2h}$  and lactate was significant and negative ( $r = -0.48$ ), no correlation was found between  $pH_{24h}$  and lactate. The correlation between glycogen and  $pH_{24h}$  ( $r = -0.50$ ) suggested that glycogen, responsible for *post mortem* glycolysis, lead to hams with low  $pH_{24h}$ . The correlation between GP and  $pH_{24h}$  resulted negative ( $r = -0.54$ ;  $P < 0.0001$ ), confirming that an increased GP indicates a high level of final lactate that can be accumulated in the post mortem muscles (Hamilton et al., 2003; Maribo et al., 1999; Monin & Sellier, 1985; Przybylski et al., 2016). However, other authors found that GP was only weakly associated with  $pH_{24h}$  in a synthetic Duroc line (England et al., 2016).

In agreement with the acidic properties of cathepsin B,  $pH_{24h}$  was negatively correlated with Catb activity. However, the magnitude of the correlation was limited ( $r = -0.19$ ;  $P = 0.0024$ ). Low  $pH_{24h}$  may favour the release of acid enzymes from the lysosomes, enhancing proteolytic activity in muscular tissue. Proteolysis and Catb activity is beneficial to the sensory quality of dry-cured hams, but high residual Catb at the end of processing period has been associated with textural (excessive softness, pastiness), flavor (i.e. bitter flavor), and appearance related (formation of white films on the ham surface) problems of dry-cured hams (Russo et al., 2000; Schivazappa et al., 2002; Virgili

et al., 1998). GP was more correlated with glycogen concentration ( $r = +0.75$ ;  $P < 0.0001$ ), than lactate accumulation ( $r = +0.40$ ;  $P < 0.0001$ ). This is not surprising given that GP was determined in the early *post mortem* period.

382  
383  
384  
385  
386

### 3.6 Cluster analysis and relationships between HWLFS and VIF

The distribution (count and frequency) of HWLFS clusters (low and high) by fatness classes of visible intermuscular fat ( $VIF = 0$  and  $VIF > 0$ ) was significant ( $P = 0.0012$ ; Table S5). About 51% of the hams clustered in the group with high HWLFS were scored to have no VIF on both legs (Fig. 2). In the cluster based on low HWLFS, only 31.4% of samples were scored to have no VIF.

These results confirmed that, beside subcutaneous level (BFT), HWLFS is partially influenced by intermuscular fat depot in the hams. However, an excess in VIF is a defect that depreciates the final product, as sliced hams with high intermuscular fat content are not accepted by consumers (Kouba & Sellier, 2011). Thus, HWLFS and VIF appear to be antagonistic traits in ham quality.

395  
396  
397  
398

### 3.7 Stepwise multiple regression analyses predicting HWLFS

Stepwise multiple regression analyses were performed to identify explanatory variables of performance, carcass, technological and raw ham quality traits that were most effective in predicting HWLFS (Table 3). For the entire sample, the predictors that satisfied the  $P < 0.05$  to enter into the regression model were BFT, LC,  $pH_{24h}$ , SW and HW, which explained 55.9% of the variation in HWLFS. BFT and LC accounted for 33.0% and 13.8% of the variation, respectively, confirming the importance of these variables in the prediction of HWLFS. As expected, BFT,  $pH_{24h}$  and SW had negative regression coefficients, confirming that higher values of these traits are beneficial for lower losses during the curing process.



In the sample with absence of visible intermuscular fat ( $VIF = 0$ ), 50.5% of the variation was explained by BFT, HW\_TR and  $pH_{24h}$ . BFT (27.4%), and HW\_TR (20.2%) explained a higher proportion of the variation. The predictors when VIF was present in one or both hams ( $VIF > 0$ ) were BFT, LC and  $pH_{24h}$  explaining 50.6% of the variation in HWLFS. BFT and LC explained 34.2 and 11.8% of the variation, respectively.

According to our data the aptitude of the ham for seasoning is mainly affected by an increase in BFT and  $pH_{24h}$  and a decrease in LC supporting the notion that the right proportion of BFT, LC, VIF, and SW is required to prevent inappropriate losses during ripening.

The phenotypic correlations and stepwise regression analyses reported in this study provided information that can be used to minimize the inappropriate loss of weight during the curing of raw hams.

#### 4. Conclusions

This study identified significant phenotypic residual correlations between HWLFS and 17 parameters, including performance, carcass and ham quality traits.

From the results a higher SM muscle  $pH_{24h}$  is desirable and associated with improved quality of meat as well as ham yield in the early stage of salting. Furthermore  $pH_{24h}$  was among the main explanatory variables influencing HWLFS, aside BTF and LC.

BFT alone explained a higher proportion of the variability in HWLFS, which confirms the importance of this variable in predicting HWLFS. In addition to BFT, VIF of trimmed hams appears to influence HWLFS and could be used as quality criteria in the evaluation of weight loss during the curing process. These results further pointed out the importance to use mature animals and carcasses of the right weight to obtain raw hams that could be suitable for the salting and seasoning processes.

#### Conflict of interest

The authors declare that they do not have competing interests.

## Acknowledgments

The authors would like to thank ANAS for their collaboration during sampling. This work was funded by University of Bologna RFO (2017-2018).

## References

- Armero, E., Flores, M., Toldrá, F., Barbosa, J. A., Olivet, J., Pla, M., & Baselga, M. (1999). Effects of pig sire type and sex on carcass traits, meat quality and sensory quality of dry-cured ham. *Journal of the Science of Food and Agriculture*, 79, 1147-1154.
- Arnau, J. (2004). Ham production. In W. K. Jensen, C. Devine, & M. Dikeman (Eds.), *Encyclopedia of meat sciences* (pp. 557–567). Oxford: Elsevier Academic Press.
- Arnau, J., Guerrero, L., & Sárraga, C. (1998). The effect of green ham pH and NaCl concentration on cathepsin activities and the sensory characteristics of dry-cured hams. *Journal of the Science of Food and Agriculture*, 77, 387-392.
- Bosi, P., & Russo, V. (2004). The production of the heavy pig for high quality processed products. *Italian Journal of Animal Science*, 3, 309-321.
- Čandek-Potokar, M., & Škrlep, M. (2011). Dry ham (“Kraški pršut”) processing losses as affected by raw material properties and manufacturing practice. *Journal of Food Processing and Preservation*, 35, 96–111.
- Čandek-Potokar, M., & Škrlep, M. (2012). Factors in pig production that impact the quality of dry-cured ham: a review. *Animal*, 6, 327-338.
- Cherel, P., Glénisson, J., Figwer, P., Pires, J., Damon, M., Franck, M., & Le Roy, P. (2010). Updated estimates of HAL n and RN– effects on pork quality: Fresh and processed loin and ham. *Meat Science*, 86, 949-954.

EEC, 1992. Parma ham. Protected Designation of Origin. General rules, dossier pursuant to article 4 of council regulation (EEC) no 2081/92 of July, 14<sup>th</sup> 1992. *Official Journal European Communities*, L208, 9-14.

England, E. M., Matarneh, S. K., Oliver, E. M., Apaoblaza, A., Scheffler, T. L., Shi, H., & Gerrard, D. E. (2016). Excess glycogen does not resolve high ultimate pH of oxidative muscle. *Meat Science*, 114, 95-102.

Fontanesi, L., Davoli, R., Costa, Nanni Costa, L., Beretti, F., Scotti, E., Tazzoli, M., ... & Russo, V. (2008). Investigation of candidate genes for glycolytic potential of porcine skeletal muscle: Association with meat quality and production traits in Italian Large White pigs. *Meat Science*, 80, 780-787.

Fontanesi, L., Schiavo, G., Galimberti, G., Bovo, S., Russo, V., Gallo, M., & Buttazzoni, L. (2017). A genome-wide association study for a proxy of intermuscular fat level in the Italian Large White breed identifies genomic regions affecting an important quality parameter for dry-cured hams. *Animal Genetics* 48, 459-465.

Fontanesi, L., Speroni, C., Buttazzoni, L., Scotti, E., Dall'Olio, S., Nanni Costa, L., ... & Russo, V. (2010). The insulin-like growth factor 2 (IGF2) gene intron3-g. 3072G> A polymorphism is not the only *Sus scrofa* chromosome 2p mutation affecting meat production and carcass traits in pigs: evidence from the effects of a cathepsin D (CTSD) gene polymorphism. *Journal of Animal Science*, 88, 2235-2245.

Fujii, J., Otsu, K., Zorzato, F., De Leon, S., Khanna, V. K., Weiler, J. E., ... & MacLennan, D. H. (1991). Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science*, 253, 448-451.

Hamilton, D. N., Miller, K. D., Ellis, M., McKeith, F. K., & Wilson, E. R. (2003). Relationships between longissimus glycolytic potential and swine growth performance, carcass traits, and pork quality. *Journal of Animal Science*, 81, 2206-2212.

495 Kouba, M., & Sellier, P. (2011). A review of the factors influencing the development of  
 496 intermuscular adipose tissue in the growing pig. *Meat Science*, 88, 213-220.

497 Larzul, C., Monin, G., Sellier, P., & Le Roy, P. (1998). Genetic variability of muscle glycolytic  
 498 potential in pigs. *Productions Animales (France)*, 11, 183-197.

499 Latorre, M. A., García-Belenguer, E., & Ariño, L. (2008). The effects of sex and slaughter weight  
 500 on growth performance and carcass traits of pigs intended for dry-cured ham from Teruel  
 501 (Spain). *Journal of Animal Science*, 86, 1933-1942.

502 Maribo, H., Støier, S., & Jørgensen, P. F. (1999). Procedure for determination of glycolytic  
 503 potential in porcine m. longissimus dorsi. *Meat Science*, 51, 191-193.

504 Milan, D., Jeon, J. T., Looft, C., Amarger, V., Robic, A., Thelander, M., ... & Ronne, H. (2000). A  
 505 mutation in PRKAG3 associated with excess glycogen content in pig skeletal muscle.  
 506 *Science*, 288, 1248-1251.

507 Monin, G., & Sellier, P. (1985). Pork of low technological quality with a normal rate of muscle pH  
 508 fall in the immediate post-mortem period: The case of the Hampshire breed. *Meat Science*, 13, 49-  
 509 63.

510 Mora, L., Escudero, E., & Toldrá, F. (2016). Characterization of the peptide profile in Spanish  
 511 Teruel, Italian Parma and Belgian dry-cured hams and its potential bioactivity. *Food Research*  
 512 *International*, 89, 638-646.

513 Nanni Costa, L., Lo Fiego, D. P., Pantano, A., & Russo, V. (1998). Relationship between glycolytic  
 514 potential and technological quality of meat and dry-cured Parma ham in the Italian heavy pig.  
 515 *CIHEAM-Options Mediterranees*, 999, 227-231.

516 Pagliarini, E., Laureati, M., Dinnella, C., Monteleone, E., Proserpio, C., & Piasentier, E. (2016).  
 517 Influence of pig genetic type on sensory properties and consumer acceptance of Parma, San  
 518 Daniele and Toscano dry-cured hams. *Journal of the Science of Food and Agriculture*, 96, 798-  
 519 806.

520 Parolari, G. (1996). Achievements, needs and perspectives in dry-cured ham technology: the  
 521 example of Parma ham. *Food Science and Technology International*, 2, 69-78.

522 Parolari, G., Virgili, R., & Schivazappa, C. (1994). Relationship between cathepsin B activity and  
 523 compositional parameters in dry-cured hams of normal and defective texture. *Meat Science*,  
 524 38, 117-122.

525 Peloso, J.V., Lopes, P.S., Gomide, L.A.M., Guimarães, S.E.F., & Carneiro, P.L.S. (2010). Carcass  
 526 and ham quality characteristics of heavy pigs from different genetic groups intended for the  
 527 production of dry-cured hams. *Meat Science*, 86, 371-376.

528 Przybylski, W., Sionek, B., Jaworska, D., & Santé-Lhoutellier, V. (2016). The application of  
 529 biosensors for drip loss analysis and glycolytic potential evaluation. *Meat Science*, 117, 7-11.

530 R Core Team (2018). R: A Language and Environment for Statistical Computing; R Foundation for  
 531 Statistical Computing: Vienna, Austria.

532 Ramos, A. M., Serenius, T. V., Stalder, K. J., & Rothschild, M. F. (2007). Phenotypic correlations  
 533 among quality traits of fresh and dry-cured hams. *Meat Science*, 77, 182-189.

534 Russo, V., Buttazzoni, L., Baiocco, C., Davoli, M. R., Nanni Costa, N. L., Schivazappa, O. C., &  
 535 Virgili, P. C. (2000). Heritability of muscular cathepsin B activity in Italian Large White pigs.  
 536 *Journal of Animal Breeding and Genetics*, 117, 37-42.

537 Russo, V., Nanni Costa, L., Lo Fiego, D. P., & De Grossi, A. (1993). Early estimation of seasoning  
 538 loss in Parma ham production. In *International Congress of Meat Science and Technology*  
 539 (Vol. 37, pp. 926-926).

540 Sárraga, C., Gil, M., & García-Regueiro, J. A. (1993). Comparison of calpain and cathepsin (B, L  
 541 and D) activities during dry-cured ham processing from heavy and light Large White pigs.  
 542 *Journal of the Science of Food and Agriculture*, 62, 71-75.

543 Scheffler, T. L., & Gerrard, D. E. (2007). Mechanisms controlling pork quality development: The  
 544 biochemistry controlling postmortem energy metabolism. *Meat Science*, 77, 7-16.

545 Schiavon, S., Carraro, L., Dalla Bona, M., Cesaro, G., Carnier, P., Tagliapietra, F., ... & Crovetto,  
 546  
 547 G. M. (2015). Growth performance, and carcass and raw ham quality of crossbred heavy pigs  
 548 from four genetic groups fed low protein diets for dry-cured ham production. *Animal Feed*  
 549 *Science and Technology*, 208, 170-181.

550 Schivazappa, C., Degni, M., Nanni Costa, L., Russo, V., Buttazzoni, L., & Virgili, R. (2002).  
 551  
 552 Analysis of raw meat to predict proteolysis in Parma ham. *Meat Science*, 60, 77-83.  
 553

554 Schivazappa, C., Virgili, R., & Parolari, G. (1992). Enzimi proteolitici nel prosciutto stagionato. II:  
 555  
 556 Correlazione con i parametri chimici del muscolo fresco. *Industria Conserve*, 67, 413-416.

556 Sturaro, E., Gallo, L., Noventa, M., & Carnier, P. (2008). The genetic relationship between  
 557  
 558 enzymatic activity of cathepsin B and firmness of dry-cured hams. *Meat Science*, 79, 375-381.

558 Toldrá, F., & Flores, M. (1998). The role of muscle proteases and lipases in flavor development  
 559  
 560 during the processing of dry-cured ham. *Critical Reviews in Food Science*, 38, 331-352.

560 Toldrá, F., Aristoy, M. C., & Flores, M. (2000). Contribution of muscle aminopeptidases to flavor  
 561  
 562 development in dry-cured ham. *Food Research International*, 33, 181-185.

562 Ventura, H. T., Lopes, P. S., Peloso, J. V., Guimarães, S. E., Carneiro, A. P. S., & Carneiro, P. L.  
 563  
 564 (2011). A canonical correlation analysis of the association between carcass and ham traits in  
 565  
 566 pigs used to produce dry-cured ham. *Genetics and Molecular Biology*, 34, 415-455.

565 Virgili, R., Parolari, G., Schivazappa, C., Bordini, C. S., & Borri, M. (1995). Sensory and texture  
 566  
 567 quality of dry-cured ham as affected by endogenous cathepsin B activity and muscle  
 568  
 569 composition. *Journal of Food Science*, 60, 1183-1186.

568 Virgili, R., Schivazappa, C., Parolari, G., Bordini, C. S., & Degni, M. (1998). Proteases in fresh  
 569  
 570 pork muscle and their influence on bitter taste formation in dry-cured ham. *Journal of Food*  
 571  
 572 *Biochemistry*, 22, 53-63.

571 Zhou, C. Y., Pan, D. D., Bai, Y., Li, C. B., Xu, X. L., Zhou, G. H., & Cao, J. X. (2019). Evaluating  
 572  
 573 endogenous protease of salting exudates during the salting process of Jinhua ham. *LWT*, 101,  
 76-82.

574      Zhou, C. Y., Wang, C., Tang, C. B., Dai, C., Bai, Y., Yu, X. B., ... & Cao, J. X. (2019). Label-free  
575      proteomics reveals the mechanism of bitterness and adhesiveness in Jinhua ham. *Food*  
576      *Chemistry*, 125012.

577      Zhou, C. Y., Wu, J. Q., Tang, C. B., Li, G., Dai, C., Bai, Y., ... & Cao, J. X. (2019). Comparing the  
578      proteomic profile of proteins and the sensory characteristics in Jinhua ham with different  
579      processing procedures. *Food Control*, 106694.

**Table 1**

Descriptive statistics of traits measured in 260 Italian Large White pigs.

Traits	Mean	Standard deviation (SD)	Min	Max	CV (%)
<i>Production (in vivo performances) and carcass traits</i>					
Average daily gain, ADG; kg/d	0.777	0.083	0.494	0.960	10.7
Feed:gain ratio, FGR	3.95	0.38	3.25	5.77	9.6
Slaughter weight, kg; SW	148.8	11.5	108.3	173.6	7.7
Slaughter age, d; SA	239.7	8.8	220	305	3.7
Backfat thickness of <i>gluteus medius</i> m., BFT; mm	26.51	5.43	13.00	42.00	20.5
Necks and loins weight (referred as lean cuts), LC; kg	28.59	2.78	19.56	36.80	9.7
Raw ham weight, HW; kg	15.41	1.31	11.03	19.24	8.5
<i>Ham technological traits<sup>1</sup></i>					
Ham weight after trimming, HW_TR; kg	12.59	1.10	9.00	16.05	8.8
Ham weight post first salting, HW_FS; kg	12.37	1.09	8.84	15.74	8.8
Ham weight loss, HWL					
Trimming, HWLTR; kg	2.82	0.36	1.94	3.99	12.8
First salting, HWLFS; g	220	38	135	317	17.4
<i>Raw ham quality traits<sup>2</sup></i>					
pH at 2 h <i>post mortem</i> , pH <sub>2h</sub>	5.94	0.24	5.41	6.80	4.1
pH at 24 h <i>post mortem</i> , pH <sub>24h</sub>	5.67	0.21	5.30	6.54	3.7
Glycogen <sup>3</sup>	46.92	22.40	7.33	135.74	47.7
Lactate <sup>3</sup>	56.40	15.92	23.28	116.93	28.2
Glycolytic potential, GP <sup>3</sup>	103.32	22.85	49.23	182.73	22.1
Cathepsin B activity, Catb <sup>4</sup>	1.16	0.23	0.6	2.0	19.6

<sup>1</sup> Averaged from left and right hams.<sup>2</sup> Measured on the *Semimembranosus* muscle.<sup>3</sup> Expressed as  $\mu\text{mol}$  of lactic acid equivalent per g of fresh muscle ( $\mu\text{mol/g}$ ).<sup>4</sup> Expressed as nmol of 7-amino-4-methylcoumarin release/min/g muscle (nmol AMC/min/g).



**Table 2**

Effects of sex, slaughter day and visible intermuscular fat of trimmed hams on traits measured in Italian Large White pigs. Probability of the effects and coefficient of determination ( $R^2$ ) of the model.

Traits <sup>1</sup>	Sex (df = 1)	Slaughter day (df = 5)	VIF (df = 2) <sup>2</sup>	R <sup>2</sup> , %
	P values <sup>3</sup>			
Average daily gain, ADG; kg/d	0.393	<b>0.002</b>	0.700	8.2
Feed:gain ratio, FGR	0.262	<b>&lt;0.001</b>	0.466	17.8
Slaughter weight, kg; SW	0.335	<b>0.024</b>	0.312	5.8
Slaughter age, d; SA	0.191	0.400	0.571	3.6
Backfat thickness of <i>Gluteus medius</i> m., BFT; mm	<b>0.010</b>	<b>&lt;0.001</b>	<b>0.001</b>	16.9
Lean cuts, LC; kg	<b>0.041</b>	<b>0.035</b>	<b>0.049</b>	8.3
Ham weight,				
Raw, HW; kg	0.724	<b>0.031</b>	0.054	5.6
Trimming, HW_TR; kg	0.358	0.122	<b>0.017</b>	5.4
Post first salting, HW_FS; kg	0.363	0.162	<b>0.026</b>	4.9
Ham weight loss				
Trimming, HWLTR; kg	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.553	18.2
First salting, HWLFS; g	0.472	<b>&lt;0.001</b>	<b>&lt;0.001</b>	32.0
pH <sub>2h</sub>	0.768	<b>0.015</b>	0.446	6.2
pH <sub>24h</sub>	0.726	<b>&lt;0.001</b>	<b>0.031</b>	12.0
Glycogen; μmol/g	<b>0.049</b>	<b>0.002</b>	<b>0.001</b>	11.5
Lactate; μmol/g	0.518	<b>0.001</b>	0.238	8.9
Glycolytic potential, GP; μmol/g	0.147	<b>0.018</b>	<b>0.044</b>	7.7
Cathepsin B activity, Catb; nmol AMC/min/g	0.569	<b>0.031</b>	0.803	5.0


<sup>1</sup> Traits are defined in Table 1.

<sup>2</sup> VIF= subjective assessment of visual intermuscular fat of trimmed hams (0= absence in both hams, 1= presence in one ham, 2= presence in both hams).

<sup>3</sup> Significant effects ( $P < 0.05$ ) are in bold.

**Figure 1**

Heatmap showing the residual correlations of ham weight loss at first salting (HWLFS) with production, carcass and ham technological traits of Italian Large white pigs. Traits are defined in Table 1. Data of residual correlations are shown in Tables S3-S5.



**Table 3**

The influence of carcass and ham quality traits on HWLFS using stepwise regression.

Group, number	Explanatory variables <sup>1</sup>	Parameter estimate $\pm$ SE <sup>2</sup>	Partial R <sup>2</sup> , %	P	Cumulative R <sup>2</sup> , %	RMSE <sup>3</sup>
All pigs, N. = 260	Intercept	0.4034 $\pm$ 0.0503		<0.001		
	BFT	-0.0030 $\pm$ 0.0004	33.0	<0.001		
	LC	0.0042 $\pm$ 0.0012	13.8	<0.001		
	pH <sub>24h</sub>	-0.0417 $\pm$ 0.0078	3.6	<0.001		
	SW	-0.0015 $\pm$ 0.0004	3.0	<0.001		
	HW	0.0152 $\pm$ 0.0027	2.5	0.003		
					55.9	0.0256
Absence of visible intermuscular fat (VIF = 0), N. = 106	Intercept	0.3759 $\pm$ 0.0989		<0.001		
	BFT	-0.0040 $\pm$ 0.0005	27.4	<0.001		
	HW_TR	0.0118 $\pm$ 0.0029	20.2	<0.001		
	pH <sub>24h</sub>	-0.0405 $\pm$ 0.0165	2.9	0.003		
					50.5	0.0281
Presence of visible intermuscular fat >0 (VIF> 0), N. = 154	Intercept	0.3693 $\pm$ 0.0607		<0.001		
	BFT	-0.0033 $\pm$ 0.0004	34.2	<0.001		
	LC	0.0044 $\pm$ 0.0008	11.8	<0.001		
	pH <sub>24h</sub>	-0.0333 $\pm$ 0.0089	4.6	0.003		
					50.6	0.0256


<sup>1</sup>Traits are defined in Table 1.

<sup>2</sup>Standard error.

<sup>3</sup>Root mean square error.

**Figure 2**

Distribution (%) of hams into weight loss at first salting (HWLFS) groups according to visible intermuscular fat (VIF) classes. Hams were grouped into two classes of HWLFS (low =  $190 \pm 21$  g, ranging from 135 g to 221 g; high =  $252 \pm 24$  g, ranging from 224 g to 317 g) based on the FASTCLUS procedure of SAS. VIF: subjective assessment of absence (0) or presence (> 0) of visual intermuscular fat of trimmed hams.



**Table S1**

Least square means (LSM)  $\pm$  standard errors (SE) of investigated traits in castrated males and gilts of Italian Large White pigs.

Traits <sup>1</sup>	Castrated males	Gilts	Gender p-values <sup>2</sup>
	LSM $\pm$ SE	LSM $\pm$ SE	
Average daily gain, ADG; kg/d	0.783 $\pm$ 0.009	0.774 $\pm$ 0.006	0.393
Feed:gain ratio, FGR	4.00 $\pm$ 0.04	3.95 $\pm$ 0.03	0.262
Slaughter weight, SW; kg	149.83 $\pm$ 1.26	148.35 $\pm$ 0.89	0.335
Slaughter age, SA; g	238.71 $\pm$ 0.86	240.09 $\pm$ 0.61	0.191
Backfat thickness of <i>gluteus medius</i> m., BFT; mm	27.95 $\pm$ 0.56	26.17 $\pm$ 0.40	<b>0.001</b>
Lean cuts, LC; kg	28.00 $\pm$ 0.30	28.76 $\pm$ 0.21	<b>0.041</b>
Ham weight			
Raw, HW; kg	15.43 $\pm$ 0.14	15.37 $\pm$ 0.10	0.724
Trimming, HW_TR; kg	12.47 $\pm$ 0.12	12.61 $\pm$ 0.09	0.358
Post first salting, HW_FS, kg	12.26 $\pm$ 0.12	12.39 $\pm$ 0.08	0.363
Ham weight loss			
Trimming, HWLTR; kg	2.96 $\pm$ 0.04	2.76 $\pm$ 0.03	<b>&lt;0.001</b>
First salting, HWLFS; g	217 $\pm$ 4	220 $\pm$ 2	0.472
pH <sub>2h</sub>	5.93 $\pm$ 0.03	5.94 $\pm$ 0.02	0.769
pH <sub>24h</sub>	5.68 $\pm$ 0.02	5.67 $\pm$ 0.02	0.726
Glycogen	49.82 $\pm$ 2.38	44.08 $\pm$ 1.68	<b>0.049</b>
Lactate	56.25 $\pm$ 1.71	57.61 $\pm$ 1.21	0.518
Glycolytic potential, GP	106.08 $\pm$ 2.47	101.69 $\pm$ 1.75	0.147
Cathepsin B activity, Catb	1.14 $\pm$ 0.02	1.16 $\pm$ 0.02	0.569

<sup>1</sup> Traits are defined in Table 1.

<sup>2</sup> The significant P values are indicated in bold.

**Table S2**

Least square means (LSM)  $\pm$  standard errors (SE) of traits investigated in different classes of visible intermuscular fat (VIF) of Italian Large White pigs.

Traits <sup>1</sup>	VIF = 0 <sup>2</sup>	VIF = 1 <sup>2</sup>	VIF = 2 <sup>2</sup>	VIF p-values <sup>4</sup>
	LSM $\pm$ SE <sup>3</sup>	LSM $\pm$ SE <sup>3</sup>	LSM $\pm$ SE <sup>3</sup>	
Average daily gain, ADG; kg/d	0.778 $\pm$ 0.009	0.785 $\pm$ 0.011	0.773 $\pm$ 0.009	0.691
Feed:gain ratio, FGR	3.93 $\pm$ 0.04	4.01 $\pm$ 0.05	3.98 $\pm$ 0.04	0.466
Slaughter weight, SW; kg	149.83 $\pm$ 1.26	148.35 $\pm$ 0.89	148.35 $\pm$ 0.89	0.312
Slaughter age, SA; g	238.71 $\pm$ 0.86	240.09 $\pm$ 0.61	240.09 $\pm$ 0.61	0.571
Backfat thickness of <i>gluteus medius</i> m., BFT; mm	24.97 $\pm$ 0.55 <sup>B, D</sup>	28.37 $\pm$ 0.66 <sup>C</sup>	27.84 $\pm$ 0.57 <sup>A</sup>	<b>0.001</b>
Lean cuts, LC; kg	29.03 $\pm$ 0.30 <sup>A</sup>	28.10 $\pm$ 0.36 <sup>B</sup>	28.01 $\pm$ 0.31 <sup>B</sup>	<b>0.049</b>
Ham weight				
Raw, HW; kg	15.69 $\pm$ 0.14 <sup>a</sup>	15.34 $\pm$ 0.17	15.17 $\pm$ 0.15 <sup>b</sup>	0.054
Trimming, HW_TR; kg	12.84 $\pm$ 0.12 <sup>a, c</sup>	12.45 $\pm$ 0.14 <sup>b</sup>	12.34 $\pm$ 0.12 <sup>d</sup>	<b>0.017</b>
Post first salting, HW_FS, kg	12.60 $\pm$ 0.12 <sup>a</sup>	12.24 $\pm$ 0.14 <sup>b, c</sup>	12.13 $\pm$ 0.12 <sup>d</sup>	<b>0.026</b>
Ham weight loss				
Trimming, HWLTR; kg	2.85 $\pm$ 0.04	2.89 $\pm$ 0.04	2.84 $\pm$ 0.04	0.553
First salting, HWLFS; g	238 $\pm$ 03 <sup>C</sup>	210 $\pm$ 4 <sup>D</sup>	207 $\pm$ 4 <sup>D</sup>	<b>&lt;0.001</b>
pH <sub>2h</sub>	5.94 $\pm$ 0.03	5.96 $\pm$ 0.03	5.91 $\pm$ 0.03	0.446
pH <sub>24h</sub>	5.63 $\pm$ 0.02 <sup>b</sup>	5.70 $\pm$ 0.03 <sup>a</sup>	5.70 $\pm$ 0.02 <sup>a</sup>	<b>0.031</b>
Glycogen	54.00 $\pm$ 2.35 <sup>a, A</sup>	46.33 $\pm$ 2.82 <sup>b</sup>	40.52 $\pm$ 2.43 <sup>B</sup>	<b>0.001</b>
Lactate	55.01 $\pm$ 1.69	56.48 $\pm$ 2.03	59.30 $\pm$ 1.75	0.238
Glycolytic potential, GP	109.01 $\pm$ 2.44 <sup>a</sup>	102.81 $\pm$ 2.93	99.83 $\pm$ 2.52 <sup>b</sup>	<b>0.044</b>
Cathepsin B activity, Catb	1.16 $\pm$ 0.02	1.14 $\pm$ 0.03	1.15 $\pm$ 0.03	0.803

<sup>1</sup> Traits are defined in Table 1.

<sup>2</sup> VIF: subjective assessment of visual intermuscular fat of trimmed hams (0 = absence in both hams, 1 = presence in one ham, 2 = presence in both hams).

<sup>3</sup> Different lower case superscript letters in the same line indicate statistically significant differences of classes (a,b: P<0.05; c,d: P<0.01; A,B: P< 0.001; C,D: P<0.0001).

<sup>4</sup> The significant P values are indicated in bold.

**Table S3**

Residual correlations<sup>1</sup> of ham weight loss at first salting (HWLFS) with production, carcass and ham technological traits of Italian Large white pigs.

	HWLF S g	ADG kg/d	FGR	SW Kg	AS d	BFT mm	LC kg	HW kg	HW_TR kg	HW_FS kg	HWLTR kg
HWLFS g	1	<b>+0.19</b> <b>0.0020</b>	<b>-0.27</b> <b>0.0001</b>	<b>+0.19</b> <b>0.0021</b>	0.00 0.9931	<b>-0.51</b> <b>&lt;0.0001</b>	<b>+0.51</b> <b>&lt;0.0001</b>	<b>+0.34</b> <b>&lt;0.0001</b>	<b>+0.40</b> <b>&lt;0.0001</b>	<b>+0.37</b> <b>&lt;0.0001</b>	0.03 0.5870
ADG kg/d		1	<b>-0.79</b> <b>&lt;0.0001</b>	<b>0.72</b> <b>&lt;0.0001</b>	<b>-0.40</b> <b>&lt;0.0001</b>	0.06 0.3741	<b>+0.59</b> <b>&lt;0.0001</b>	<b>+0.66</b> <b>&lt;0.0001</b>	<b>+0.63</b> <b>0.0001</b>	<b>+0.63</b> <b>&lt;0.0001</b>	<b>+0.51</b> <b>&lt;0.0001</b>
FGR			1	<b>-0.38</b> <b>&lt;0.0001</b>	<b>+0.44</b> <b>&lt;0.0001</b>	+0.12 0.0516	<b>-0.41</b> <b>&lt;0.0001</b>	<b>-0.39</b> <b>&lt;0.0001</b>	<b>-0.39</b> <b>&lt;0.0001</b>	<b>-0.38</b> <b>&lt;0.0001</b>	<b>-0.23</b> <b>0.0002</b>
SW kg				1	+0.03 0.6551	<b>+0.19</b> <b>0.0031</b>	<b>+0.79</b> <b>&lt;0.0001</b>	<b>+0.89</b> <b>&lt;0.0001</b>	<b>+0.85</b> <b>&lt;0.0001</b>	<b>+0.85</b> <b>&lt;0.0001</b>	<b>+0.69</b> <b>&lt;0.0001</b>
AS d					1	+0.10 0.1226	+0.03 0.6098	+0.02 0.7767	+0.02 0.6950	+0.03 0.6915	-0.01 0.8572
BFT mm						1	<b>-0.20</b> <b>0.0011</b>	+0.07 0.2669	-0.03 0.6219	-0.02 0.7956	<b>+0.38</b> <b>&lt;0.0001</b>
LC kg							1	<b>+0.77</b> <b>&lt;0.0001</b>	<b>+0.80</b> <b>&lt;0.0001</b>	<b>+0.79</b> <b>&lt;0.0001</b>	<b>+0.41</b> <b>&lt;0.0001</b>
HW kg								1	<b>+0.98</b> <b>&lt;0.0001</b>	<b>+0.98</b> <b>&lt;0.0001</b>	<b>+0.70</b> <b>&lt;0.0001</b>
HW_TR kg									1	<b>+0.99</b> <b>&lt;0.0001</b>	<b>+0.53</b> <b>&lt;0.0001</b>
HW_FS kg										1	<b>+0.54</b> <b>&lt;0.0001</b>
HWLTR kg											1

<sup>1</sup> The values for the phenotypic correlations are shown in the upper rows, the lower rows report the P values, significant correlations (at least  $P < 0.05$ ) are indicated in bold. Traits are defined in Table 1.



**Table S4**

Residual correlations<sup>1</sup> of ham weight loss at first salting (HWLFS) with raw ham quality traits of Italian Large White pigs.

	HWLFS g	pH <sub>2h</sub>	pH <sub>24h</sub>	Glycogen μmol/g	Lactate μmol/g	GP μmol/g	Catb nmol AMC/min/g
HWLFS g	1	-0.10 0.1303	<b>-0.25</b> <b>&lt;0.0001</b>	<b>+0.22</b> <b>0.0004</b>	-0.08 0.2117	<b>+0.16</b> <b>0.0113</b>	+0.08 0.1815
pH <sub>2h</sub>		1	<b>+0.58</b> <b>&lt;0.0001</b>	-0.06 0.3383	<b>-0.48</b> <b>&lt;0.0001</b>	<b>-0.39</b> <b>&lt;0.0001</b>	-0.10 0.1048
pH <sub>24h</sub>			1	<b>-0.50</b> <b>&lt;0.0001</b>	-0.07 0.2534	<b>-0.54</b> <b>&lt;0.0001</b>	<b>-0.19</b> <b>0.0024</b>
Glycogen μmol/g				1	<b>-0.31</b> <b>&lt;0.0001</b>	<b>+0.75</b> <b>&lt;0.0001</b>	+0.11 0.0800
Lactate μmol/g					1	<b>+0.40</b> <b>&lt;0.0001</b>	+0.02 0.7001
GP μmol/g						1	+0.12 0.0504
Catb nmol AMC/min/							1

<sup>1</sup> The values for the phenotypic correlations are shown in the upper rows, the lower rows report the P values, significant correlations (at least  $P <$

0.05) are indicated in bold. Traits are defined in Table 1.

**Table S5**

Distribution of hams into weight loss at first salting (HWLFS)<sup>1</sup> groups according to visible intermuscular fat (VIF)<sup>2</sup>

Traits	HWLFS clusters <sup>1</sup>		P-value
	Low Number (%)	High Number (%)	
Visible intermuscular fat, VIF <sup>2</sup>	137	123	0.0012
0 = absence in both hams	43 (31.4)	63 (51.2)	
> 0 = presence in one or both hams	94 (69.6)	60 (48.8)	

<sup>1</sup> Hams were grouped into two classes of HWLFS (low =  $190 \pm 21$  g, ranging from 135 g to 221 g; high =  $252 \pm 24$  g, ranging from 224 g to 317 g) based on FASTCLUS procedure of SAS.

<sup>2</sup> VIF: subjective assessment of visual intermuscular fat of trimmed hams.




Figure 1




Figure 2