

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Meta-analysis of the effects of venous super-drainage in deep inferior epigastric artery perforator flaps for breast reconstruction

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Pignatti, M., Pinto, V., Giorgini Federico A, Lozano Miralles, M.E., Cannamela, G., D'Arpa, S., et al. (2021). Meta-analysis of the effects of venous super-drainage in deep inferior epigastric artery perforator flaps for breast reconstruction. *MICROSURGERY*, 41(2), 186-195 [10.1002/micr.30682].

Availability:

This version is available at: <https://hdl.handle.net/11585/779595> since: 2020-12-30

Published:

DOI: <http://doi.org/10.1002/micr.30682>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the accepted manuscript of:

Pignatti, M, Pinto, V, Giorgini, FA, et al. *Meta-analysis of the effects of venous super-drainage in deep inferior epigastric artery perforator flaps for breast reconstruction*. Microsurgery. 2020; 1– 10.

Final version available: <https://doi.org/10.1002/micr.30682>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

**A meta-analysis of the effects of Venous super-drainage in Deep Inferior Epigastric
Perforator flaps for breast reconstruction.**

Marco Pignatti ^a, Valentina Pinto ^b, Federico A. Giorgini ^{b,e}, Maria Elisa Lozano Miralles ^{b,e}, Giacomo Cannamela ^c, Salvatore D'Arpa ^d, Riccardo Cipriani ^b, Giorgio De Santis ^e

M. Pignatti, MD, FEBOPRAS

Plastic Surgery, Policlinico di Sant'Orsola - DIMES, University of Bologna, Italy

V. Pinto, MD, PhD

Plastic Surgery, Policlinico di Sant'Orsola - Bologna, Italy

F. A. Giorgini, MD

Plastic Surgery, Policlinico di Sant'Orsola - Bologna, Italy

&

Plastic Surgery, Policlinico di Modena, University of Modena and Reggio Emilia, Italy

M. E. Lozano Miralles, MD

Plastic Surgery, Policlinico di Sant'Orsola - Bologna, Italy

&

Plastic Surgery, Policlinico di Modena, University of Modena and Reggio Emilia, Italy

G. Cannamela, Data scientist.

Milan. Italy

S. D'Arpa, MD, PhD

Plastic and Reconstructive Surgery, Department of Surgical, Oncological and Oral Sciences (DICHIORONS). University of Palermo, Palermo, Italy

R. Cipriani, MD

Plastic Surgery, Policlinico di Sant'Orsola - Bologna, Italy

G. De Santis, MD

Plastic Surgery, Policlinico di Modena, University of Modena and Reggio Emilia, Italy

Corresponding author

Marco Pignatti, MD, FEBOPRAS

Via Massarenti, 9

1
2
3 40138 Bologna, Italy
4 mrpignatti@gmail.com
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A meta-analysis of the effects of Venous super-drainage in Deep Inferior Epigastric artery Perforator flaps for breast reconstruction.

Abstract

Introduction

Venous congestion is the most common vascular complication of the Deep Inferior Epigastric artery Perforator (DIEP) flaps. Adding a second venous drainage by anastomosing a flap vein and a recipient vein (super-drainage) is considered the solution of choice. Evidence to support this procedure, had not yet been confirmed by an analysis of the literature. We aimed to provide this evidence.

Materials and Methods

We searched the literature (MedLine, Scopus, EMBASE, Cochrane Library, and Google Scholar), for studies discussing venous congestion and venous super-drainage in DIEP flap for breast reconstruction. 13/35 articles compared results between one or two venous anastomoses. Meta-analysis was performed following PRISMA guidelines . Pooled risk ratio (RRs) for congestion, fat necrosis, partial necrosis, and total necrosis with corresponding 95% confidence intervals (CI) were calculated using a fixed-effect model with the Mantel–Haenszel method. The need to return to surgery (95% CI) was estimated with a random effect model using the DerSimonian & Liard method.

Results

We showed a statistically significant advantage of super-drainage to reduce the venous congestion of the flap (RR: 0.12, 95% CI: 0.04 – 0.34, p -value < 0,001), partial flap necrosis (RR: 0.50, 95% CI: 0.30 - 0.84, p -value 0,008), total flap necrosis (RR: 0.31, 95% CI: 0.11 - 0.85, p -value 0.023), and the need to take the patient back to surgery for perfusion-related complications (RR: 0,45, 95% CI: 0.21 - 0.99, p value 0.048).

Conclusions

Performing a second venous anastomosis between the SIEV and a recipient vein (venous superdrainage) reduces venous congestion and related complications in DIEP flaps for breast reconstruction.

Keywords: DIEP, venous congestion, meta-analysis, super-charging, super-drainage

INTRODUCTION

The Deep Inferior Epigastric artery Perforator (DIEP) flap, is considered the gold standard for autologous breast reconstruction.¹ Flap failure is reported to occur in 1 to 5 % of cases.² However, the frequency of partial failures and their consequences is higher.³ The unsatisfactory results are frequently due to perfusion-related complications, developing in up to 17% of DIEP flaps.⁴ Venous congestion is the most common of the above-mentioned perfusion-related complications, with a reported incidence ranging from 2% to 20%^{5,6} representing one of the main causes of fat necrosis, partial flap necrosis and, less frequently, complete necrosis of the flap.³ The etiology of venous congestion is multifactorial⁷⁻¹² with reversible and non-reversible causes.

When the causes of venous congestion are not reversible intraoperatively, treatment of the congestion itself is necessary and usually consists in increasing venous drainage of the flap by adding a second venous anastomosis between the flap and a recipient vein. The term flap supercharge or, semantically more correctly, venous super-drainage is usually employed.

Several papers have been published highlighting the problem of DIEP's venous congestion, and describing the surgical solutions available. **Although there is no controversy regarding the use of flap super-drainage, which is the standard solution in case of venous congestion, evidence of the efficacy of this procedure on flap necrosis and vascular complications had not yet been reported.**

To try to clarify the matter, we conducted a literature search statistically evaluating with a meta-analysis the relevant studies looking for evidence on the role of DIEP flap's venous super-drainage in decreasing flap complications.

MATERIALS AND METHODS

We performed a meta-analysis of the literature evaluating the role of a second venous anastomosis to prevent the complications of DIEP flap's congestion. A literature search was conducted using MedLine, Scopus, EMBASE, Cochrane Library, and Google Scholar to identify and include all citations from 1994 to June 2019. To reduce inclusion bias, two of the Authors (VP and FG) performed, separately, the initial article search and selection with pertinent keywords ("Deep inferior epigastric perforator" OR "DIEP" AND "flap" [All Fields] AND venous congestion [All Fields] AND ("additional anastomosis"[Subheading] AND "superficial inferior epigastric vein"[All Fields] OR "SIEV"[All Fields] AND "supercharging" OR "superdrain" OR "super-drainage") (Figure 1)

Duplicated articles, isolated abstracts, case reports, correspondence and letters were excluded and only full-text articles in English regarding DIEP flaps and venous congestion were considered.

We followed the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA statement)¹³.

We considered eligible the retrieved publications based on PICOS criteria (Patients, Interventions, Control/comparison, Outcomes, and Study Design), and analyzed by meta-analysis the incidence of perfusion-related complications: venous congestion, flap necrosis (total and partial), fat necrosis and need to take the patient back to surgery for perfusion-related complications. The most relevant data collected from the publications found were: first author, year of publication, number of patients and events, type of the second anastomosis and recipient vein, and operative time.

Statistical Analysis

The Mantel-Haenszel test in R was used to calculate Pooled RRs and corresponding 95% confidence intervals (CI) for perfusion-related complications. The heterogeneity among studies

was analyzed by using the I^2 test. We used a fixed-effects model for I^2 ranging from 0 to 30%; in the case of $I^2 > 30\%$, we applied a random effect model using the DerSimonian & Liard method.

For each test p value $\leq 0,05$ was set for significance

RESULTS

After exclusion of non-relevant papers, we identified 35 eligible articles and performed a qualitative evaluation following the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA statement).¹³ Of these articles, 13 compared results between one or two venous anastomoses in DIEP flaps for breast reconstruction^{3,14-25} and were included in our study. All the studies that we reviewed were retrospective, except the one by Ayestaray,¹⁶ which is prospective and randomized.

Overall, our study analyzed 3094 DIEPs, of which, 1279 received a second venous anastomosis ("case group"); and 1815 received only one venous anastomosis ("control group"). Venous congestion occurred in 38 out of 1121 flaps, with an overall mean rate of 3.4%. The venous congestion rate in each study ranged from 0.9% to 36.5%. Regarding other perfusion-related complications, our literature review revealed 1.5 % total flap loss (34/2278) (ranging from 0% to 9.6%), 6% partial flap loss (78/1294) (0% to 28.8%), 11.9% fat necrosis (217/1824) (7.2% to 34.6%), 11.1% take back to surgery for perfusion-related complications (187/1678) (0.9% to 21.2%).

Of the 13 articles^{3,14-25} included in our meta-analysis, 12^{3,14-18,20-25} reported the second anastomosis to have been performed between the SIEV and a recipient vein of the thorax, while in one study¹⁹ the second anastomosis was done using as flap vein the SIEV or a second DIEV, without additional information being provided. In all the articles included in our systematic review, performing an adjunctive anastomosis variably increased the operation time from no difference to 105 minutes (TABLE 1).

1
2
3 In more detail we found that anastomosing the SIEV of the flap with a recipient vein provided a
4 statistically significant advantage in terms of venous congestion (RR: 0.12, 95% CI: 0.04 – 0.34) *p*-
5 value < 0,001) (Figure 2) and to prevent flap necrosis both partial (RR: 0.50, 95% CI: 0.30 - 0.84, *p*-
6 value 0,008) (Figure 3) and total (RR: 0.31, 95% CI: 0.11 - 0.85, *p*-value 0.023) (Figure 4). A second
7 venous drainage showed a statistically significant advantage also regarding the need to take the
8 patient back to surgery for perfusion-related complications (RR: 0,45, 95% CI: 0.21 - 0.99, *p* value
9 0.048) (Figure 5). Conversely, the advantage of a second venous anastomosis was not significantly
10 beneficial to prevent fat necrosis RR: 0.83, 95% CI: 0.63 - 1.10 reporting a *p*-value 0.19). (Figure 6)
11
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **DISCUSSION**

26
27
28
29 Although the DIEP flap is considered the gold standard for breast reconstruction, venous
30 congestion can cause complete or partial flap failure and several other complications.
31
32

33
34 Our work, by reviewing the updated pertinent literature, aimed to provide evidence on
35 the effect of super-drainage in terms of venous congestion and related complications. In case of
36 intraoperative venous congestion, the reversible causes can be identified and corrected. In
37 particular, anastomotic failure requires resection of the anastomotic site, removal of any
38 thrombus present, rinsing of the donor and recipient vessels with a heparinized solution (and,
39 depending on the clinical situation, time since the beginning of occlusion and kind of flap, flap
40 perfusion with thrombolytic agents), and vessels' re-anastomosis. Vessel kinking can be solved by
41 repositioning the entire pedicle of the flap by supporting it with a piece of fat or muscle, by
42 stabilizing it with fibrin glue or by applying a stitch between the kinked vessel and the surrounding
43 tissue to straighten its course. A mild to moderate torsion of a long pedicle can be solved simply
44 by repositioning of the pedicle and therefore redistributing the torsion on its entire length. In
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

more severe cases the anastomosis will need to be redone. If, in a large flap, the venous congestion is stable and only involves the distal portion of the flap, because of insufficient drainage despite effective outflow through the pedicle, resection of that area may be sufficient.

The second venous anastomosis, aimed to obtain a super-drainage of the flap, can be performed

a) between the deep venous system of the flap and a comitant vein of the recipient vessels (comitant DIEV-IMV)^{19,26}, b) between the deep venous system of the flap and a vein of the thorax (comitant DIEV-vein of thorax)²⁷, c) between the SIEV and the DIEV or to one of its side-branches (SIEV-DIEV)^{1,5,17,28,29}

d) between SIEV and a vein of the thorax^{3,14-22,24}.

In physiologic conditions, the dominant venous drainage to the lower abdominal skin and fat is provided by the superficial venous system through the inferior epigastric vein (SIEV).^{5, 15} All the papers included in our meta-analysis used the SIEV as a second vein of the flap to increase venous output. Table 1. The SIEV can be anastomosed, as said, to a vein of the deep venous system of the flap (intra-flap anastomosis) 1,5 or a vein of the patient's chest (extra-flap anastomosis).¹⁴

Unfavorable anatomy of the venous system is considered among the most important, not-reversible, causes of venous congestion. Several studies^{5,6,10,30} attribute the clinical manifestations of venous congestion and the so-called perfusion-related complications to the lack of sufficient venous drainage from all areas of the flap (superficial, deep, ipsilateral, contralateral). Not all the factors related to the flap's venous anatomy can be modified by the surgeon. Therefore accurate planning based on vascular anatomy is of utmost importance to prevent venous congestion of the flap. Computed tomographic angiography and magnetic resonance angiography permit to visualize the arterial system and, even more importantly, the venous connections between the deep and superficial systems, allowing to choose the most favorable perforators for optimal perfusion and

1
2
3 drainage.^{7,10} The ideal perforator will have a large caliber, a central position in the flap and
4
5 numerous connections between the deep and superficial venous systems.³¹
6
7

8 Once venous congestion occurs, and intraoperatively the reversible causes have been excluded, it
9
10 is necessary to treat the venous congestion itself, by venous super-drainage.
11
12

13 One previously published meta-analysis, from Lee et al³² showed that the use of super-drainage
14
15 has a statistical advantage over venous congestion but only a trend toward a decreased risk of
16
17 congestion-related complications. Further studies were, therefore, suggested. A detailed
18
19 comparison with that previous meta-analysis³², is presented in TABLE 2. Our analysis includes the
20
21 six articles examined by Lee and seven additional ones, reporting on large groups of patients^{22,23}
22
23 and on a prospective randomized study¹⁶. (TABLE 1). Statistical methods were comparable but the
24
25 larger amount of data, allowed us to achieve statistical evidence that the use of super-drainage,
26
27 employing a second venous anastomosis between the SIEV and a recipient vein, reduces venous
28
29 congestion, prevents partial and total flap necrosis, and the need to take the patient back to
30
31 surgery. Unfortunately, the available data did not show a statistically significant advantage of a
32
33 second venous anastomosis to prevent fat necrosis, a common complication (6% to 17.4%) in DIEP
34
35 flaps.
36
37
38
39
40

41
42 However, from the results of our meta-analysis, there seems to be an advantage also in
43
44 performing a second venous anastomosis in flaps without venous congestion. Therefore, the
45
46 choice of performing routinely a second venous anastomosis should be considered.
47
48
49
50

51
52 A comparison between the super-drainage technique consisting in anastomosis of the SIEV to a
53
54 recipient vein (the only one for which we have demonstrated a statistical advantage), and other
55
56 techniques, such as DIEV to recipient vein, intermittent SIEV catheter drainage, or the application
57
58
59
60

of leeches, was not feasible, based on the data available in the literature. The same problem exists regarding which vein the SIEV was anastomosed (e.g., retrograde IMV vs cephalic turndown).

A limitation of our study consists in the difficulty to achieve solid conclusions when working on mainly retrospective articles , none of which with an evidence level I or II (TABLE 1). However, our study follows the PRISMA guidelines ¹³, which warrant the quality of a meta-analysis.

Our study is the first to report statistically significant results on the subject of super-drainage. Our meta-analysis of the literature demonstrated the usefulness of venous superdrainage when performing DIEP flaps in postmastectomy breast reconstruction, an elective surgery burdened with relevant emotional implications for the patient. The use of an adjunctive venous anastomosis as a preventive measure could be considered.

CONCLUSIONS

In conclusion, our meta-analysis shows that venous super-drainage, i.e. performing a second venous anastomosis between the superficial venous system and a recipient vein, provides a statistically significant advantage in terms of venous congestion and related complications in DIEP flaps for breast reconstruction.

REFERENCES

1) Bartlett EL, Zavlin D, Menn ZK, Spiegel AJ. Algorithmic Approach for Intraoperative Salvage of Venous Congestion in DIEP Flaps. J Reconstr Microsurg. 2018; 34(06): 404–412.

- 1
2
3 2) Lie KH, Barker AS, Ashton MW. A classification system for partial and complete DIEP flap
4
5 necrosis based on a review of 17,096 DIEP flaps in 693 articles including analysis of 152 total flap
6
7 failures. *Plast Reconstr Surg*. 2013;132:1401–1408
8
9
10
11
12
13 3) Santanelli F, Longo B, Cagli B, Pugliese P, Sorotos M, Paolini G. Predictive and protective factors
14
15 for partial necrosis in DIEP flap breast reconstruction: does nulliparity bias flap viability? *Ann Plast*
16
17 *Surg*. 2015; 74(1): 47-51.
18
19
20
21
22
23 4) Peeters WJ, Nanhekhan L, Van Ongeval C, Fabrè G, Vandevoort M. Fat necrosis in deep inferior
24
25 epigastric perforator flaps: an ultrasound-based review of 202 cases. *Plast Reconstr Surg*. 2009;
26
27 124(6):1754-1758.
28
29
30
31
32
33 5) Sbitany H, Mirzabeigi MN, Kovach SJ, Wu LC, Serletti JM. Strategies for recognizing and
34
35 managing intraoperative venous congestion in abdominally based autologous breast
36
37 reconstruction. *Plast Reconstr Surg*. 2012; 129(04):809–815.
38
39
40
41
42
43 6) Kim DY, Lee TJ, Kim EK, Yun J, Eom JS. Intraoperative venous congestion in free transverse
44
45 rectus abdominis musculocutaneous and deep inferior epigastric artery perforator flaps during
46
47 breast reconstruction: a systematic review. *Plast Surg (Oakv)*. 2015; 23(04):255–259.
48
49
50
51
52
53
54 7) Schaverien MV, Ludman CN, Neil-Dwyer J et al. Relationship between venous congestion and
55
56 intraflap venous anatomy in DIEP flaps using contrast-enhanced magnetic resonance angiography.
57
58 *Plast Reconstr Surg*. 2010; 126(2):385-392.
59
60

1
2
3
4
5
6 8) Blondeel PN, Arnstein M, Verstraete K, et al. Venous congestion and blood flow in free
7
8 transverse rectus abdominis myocutaneous and deep inferior epigastric perforator flaps. *Plast*
9
10 *Reconstr Surg*. 2000; 106(6):1295–1299.
11
12

13
14
15 9) Tran NV, Buchel EW, Convery PA. Microvascular complications of DIEP flaps. *Plast Reconstr*
16
17 *Surg*. 2007; 119(5):1397–1405; discussion 1406-1408.
18
19
20

21
22
23 10) Rozen WM, Pan WR, Le Roux CM, Taylor GI, Ashton MW. The venous anatomy of the anterior
24
25 abdominal wall: An anatomical and clinical study. *Plast Reconstr Surg*. 2009;124(3): 848–853.
26
27
28

29
30 11) Lundberg J, Mark H. Avoidance of complications after the use of deep inferior epigastric
31
32 perforator flaps for reconstruction of the breast. *Scand J Plast Reconstr Surg Hand Surg*. 2006;
33
34 40(2):79–81.
35
36
37

38
39
40 12) Nahabedian MY, Tsangaris T, Momen B. Breast reconstruction with the DIEP flap or the
41
42 muscle-sparing (MS-2) free TRAM flap: is there a difference? *Plast Reconstr Surg*. 2005;
43
44 115(2):436–444-discussion 445-446.
45
46
47

48
49
50 13) Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for
51
52 systematic reviews and metanalyses: the PRISMA statement. *PLoS Med*. 2009; 21; 6(7): e1000097.
53
54
55

56
57 14) Eom JS, Sun SH, Lee TJ. Selection of the recipient veins for additional anastomosis of the
58
59 superficial inferior epigastric vein in breast reconstruction with free transverse rectus abdominis
60

1
2
3 musculocutaneous or deep inferior epigastric artery perforator flaps. Ann Plast Surg. 2011; 67(5):
4
5 505-509.
6
7
8
9

10
11 15) Enajat M, Rozen WM, Whitaker IS, Smith JM, Acosta R. A single center comparison of one
12
13 versus two venous anastomoses in 564 consecutive DIEP flaps: investigating the effect on venous
14
15 congestion and flap survival. Microsurgery. 2010; 30(3):185-191.
16
17
18
19

20
21 16) Ayestaray B, Yonekura K, Motomura H, Ziade M. A Comparative Study Between Deep Inferior
22
23 Epigastric Artery Perforator and Thoracoacromial Venous Supercharged Deep Inferior Epigastric
24
25 Artery Perforator Flaps. Ann Plast Surg. 2016; 76(1):78-82.
26
27
28
29

30
31 17) Xin Q, Luan J, Mu H, Mu L. Augmentation of venous drainage in deep inferior epigastric
32
33 perforator flap breast reconstruction: efficacy and advancement. J Reconstr Microsurg. 2012
34
35 Jun;28(5):313-8
36
37
38
39
40
41

42
43 18) Ali R, Bernier C, Lin YT, Ching WC, Rodriguez EP, Cardenas-Mejia A et al. Surgical strategies to
44
45 salvage the venous compromised deep inferior epigastric perforator flap. Ann Plast Surg. 2010
46
47 Oct;65(4):398-406.
48
49
50
51
52
53

54
55 19) La Padula S, Hersant B, Noel W, Niddam J, Hermeziu O, Bouhassira J et al. Use of the
56
57 retrograde limb of the internal mammary vein to avoid venous congestion un DIEP flap breast
58
59
60

1
2
3 reconstruction. Further evidences of a reliable and time-sparing procedure. Microsurgery. 2016;
4
5 36(6):4476-452.
6
7
8
9

10 20) Boutros SG. Double venous system drainage in deep inferior epigastric perforator flap breast
11
12 reconstruction: a single-surgeon experience. Plast Reconstr Surg. 2013;131(4): 671-676.
13
14
15
16

17 21) Al Hindi A, Ozil C, Rem K, Rausky J, Moris V, Guillier D, Binder JP, Revol M, Cristofari S.
18
19 Intraoperative superficial inferior epigastric vein preservation for venous compromise prevention
20
21 in breast reconstruction by deep inferior epigastric perforator flap. Ann Chir Plast Esthet. 2019
22
23 Jun;64(3):245-250.
24
25
26
27
28
29

30 22) Ochoa O, Pisano S, Chrysopoulou M, Ledoux P, Arishita G, Nastala C. Salvage of intraoperative
31
32 deep inferior epigastric perforator flap venous congestion with augmentation of venous outflow:
33
34 flap morbidity and review of the literature. Plast Reconstr Surg Glob Open. 2013 Nov 7;1(7):e52.
35
36
37
38
39

40 23) Unukovych D, Gallego CH, Aineskog H, Rodriguez-Lorenzo A1, Mani M. Predictors of
41
42 Reoperations in Deep Inferior Epigastric Perforator Flap Breast Reconstruction. Plast Reconstr Surg
43
44 Glob Open. 2016 Aug 29;4(8):e1016. doi: 10.1097/GOX.0000000000001016. eCollection 2016 Aug.
45
46
47
48
49

50 24) Al-Dhamin A, Bissell MB, Prasad V, Morris SF. The use of retrograde limb of internal mammary
51
52 vein in autologous breast reconstruction with DIEAP flap: anatomical and clinical study. Ann Plast
53
54 Surg. 2014 Mar;72(3):281-4.
55
56
57
58
59
60

- 1
2
3 25) Lee KT, Lee JE, Nam SJ, Mun GH. Ischaemic time and fat necrosis in breast reconstruction with
4 a free deep inferior epigastric perforator flap. J Plast Reconstr Aesthet Surg. 2013 Feb;66(2):174-
5 81.
6
7
8
9
10
11
12
13 26) Marck KW, van der Biezen JJ, Dol JA. Internal mammary artery and vein supercharge in TRAM
14 flap breast reconstruction. Microsurgery. 1996;17(7):371-4.
15
16
17
18
19
20 27) Fansa H. Internal Mammary Artery and Vein Perforator Vessels as Troubleshooter Recipient
21 Vessels. Plast Reconstr Surg Glob Open. 2019; 20;7(3):e2148.
22
23
24
25
26
27 28) Rohde C, Keller A. Novel technique for venous augmentation in a free deep inferior epigastric
28 perforator flap. Ann Plast Surg. 2005 Nov;55(5):528-30.
29
30
31
32
33
34 29) Liu TS, Ashjian P, Festekjian J. Salvage of congested deep inferior epigastric perforator flap
35 with a reverse flow venous anastomosis. Ann Plast Surg. 2007 Aug;59(2):214-7.
36
37
38
39
40
41
42 30) Schaverien M, Saint-Cyr M, Arbique G, Brown SA. Arterial and venous anatomies of the deep
43 inferior epigastric perforator and superficial inferior epigastric artery flaps. Plast Reconstr Surg.
44 2008; 121(6):1909–1919.
45
46
47
48
49
50
51 31) Davis CR, Jones L, Tillett RL, Richards H, Wilson SM. Predicting venous congestion before DIEP
52 breast reconstruction by identifying atypical venous connections on preoperative CTA imaging.
53 Microsurgery. 2019 Jan;39(1):24-31.
54
55
56
57
58
59
60

32) Lee KT, Mun GH. Benefits of super-drainage using SIEV in DIEP flap breast reconstruction: A systematic review and meta-analysis. Microsurgery. 2017; 37(1):75-83.

CRediT authorship contribution statement

Marco Pignatti: Conceptualization, Data curation, Formal analysis, Writing - original draft, Writing - review & editing. Valentina Pinto: Conceptualization, Data curation, Formal analysis, Writing - original draft.

Federico A. Giorgini: Conceptualization, Data curation.

Maria Elisa Lozano Miralles: Conceptualization, Data curation, Formal analysis. Salvatore D'Arpa: Conceptualization, Data curation Writing - original draft, review and editing.

Riccardo Cipriani: Conceptualization, Formal analysis, Writing - review & editing.

Giorgio De Santis: Conceptualization, Formal analysis, Writing - original draft, Writing - review & editing.

Disclosure statement:

The authors do not have any conflict of interest to disclose

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LEGENDS to Figures and Tables

Figure 1

Flow diagram that shows selection of articles.

Figure 2

Venous congestion in DIEP flaps.

Articles and related forest plots dealing with the complication

VC: number of flaps presenting Venous Congestion

The column "Cases" represents the total number of flaps where an adjunctive second vein anastomosis (SIEV-recipient vein) was performed. The column "Controls" represents the total number of flaps where a single vein anastomosis (DIEV-main vein) was performed

Figure 3

Partial necrosis in DIEP flaps.

Articles and related forest plots dealing with the complication

PN: number of flaps presenting Partial Necrosis

Figure 4

Total necrosis in DIEP flaps.

Articles and related forest plots dealing with the complication

TN: number of flaps presenting Total Necrosis

Figure 5

Need to take back to surgery for perfusion-related complications in DIEP flaps.

Articles and related forest plots dealing with each complication

TB: number of flaps needing Take Back to surgery

Figure 6

Fat necrosis

Articles and related forest plots dealing with the complication

FN: number of flaps presenting Fat Necrosis

TABLE 1.

Studies included in our meta-analysis

OT: operating time, NA: not available, TAV: thoracoacromial vein, TDV: thoracodorsal vein, ICV: intercostal vein, IMV: internal mammary vein, LTV: lateral thoracic vein, IMVr: internal mammary vein retrograde branch, IMVp: internal mammary vein perforator, SCV: scapular circumflex vein, SUV: subscapular vein, AV: axillary vein

“Cases”: flaps with a second venous anastomosis. “Controls”: flaps with a single venous anastomosis

In bold are the 7 studies that were not analyzed in the previously published meta-analysis by Lee³²,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 2.

Comparison between our meta-analysis and the one from Lee and Mun

RR: Risk Ratio. Cases: flaps with a second venous anastomosis. Controls: only one venous anastomosis

In bold are the 7 studies that were not analyzed in the previously published meta-analysis by Lee³², “Cases”: flaps with a second venous anastomosis. “Controls”: flaps with a single venous anastomosis




Figure 1

355x266mm (96 x 96 DPI)




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

TABLE 1. Studies included in our meta-analysis

Study	Study type	Flap 2° vein	Recipient vein	N. cases	N. controls	Mean OT cases (minutes)	Mean OT controls (minutes)	Difference in OT (minutes)
Ali, 2010	Retrospective cohort	SIEV	IMV	21	11	Na	Na	Na
Enajat, 2010	Retrospective cohort	SIEV	Cephalic vein	291	273	383	385	- 2
Eom, 2011	Retrospective cohort	SIEV	LTV, branch of the TAV, IMVp	45	108	Na	Na	Na
Lee, 2012	Retrospective cohort	SIEV	Na	18	68	Na	Na	Na
Xin, 2012	Retrospective cohort	SIEV	TDV, LTV, ICV, SIEV, DIEV	32	47	396	366	+ 30
Boutros, 2013	Retrospective cohort	SIEV	IMVp, IMV	311	41	Na	Na	Na
Ochoa, 2013	Retrospective cohort	SIEV	IMV	87	639	314	253	+ 61
Al-Dhamin, 2014	Retrospective cohort	SIEV	IMVr	31	17	Na	Na	Na
Santanelli, 2015	Retrospective cohort	SIEV	SCV, TDV, IMV, SUV, LTV	173	74	Na	Na	Na
Ayestaray, 2016	Prospective cohort, randomized	SIEV	TAV	23	29	510	405	+ 105
La Padula, 2016	Retrospective cohort	SIEV or 2 nd DIEV	IMVr	36	38	Na	Na	Na
Unukovych, 2016	Retrospective cohort	SIEV	Na	211	272	Na	Na	Na
Al Hindi, 2019	Retrospective cohort	SIEV	Cephalic vein	14	184	461	356	+ 105

1 OT: operating time, Na: not available, TAV: thoracoacromial vein, TDV: thoracodorsal vein, ICV: intercostal vein, IMV: internal mammary vein, LTV: lateral
2 thoracic vein, IMVr: internal mammary vein retrograde branch, IMVp: internal mammary vein perforator, SCV: scapular circumflex vein, SUV: subscapular
3 vein, AV: axillary vein
4

5 “Cases”: flaps with a second venous anastomosis. “Controls”: flaps with a single venous anastomosis
6

7 In bold are the 7 studies that were not analyzed in the previous published metanalysis by Lee, 2017
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

	Lee and Mun, 2017	Pignatti et al, 2020
<i>Articles included</i>	Enajat, 2010 Lee, 2012 Xin, 2012 Santanelli, 2015 Boutros, 2013 Al-Dhamin, 2014	Ali, 2010 Enajat, 2010 Eom, 2011 Lee, 2012 Xin, 2012 Boutros, 2013 Ochoa, 2013 Al-Dhamin, 2014 Santanelli, 2015 Ayestaray, 2016 La Padula, 2016 Unukovich, 2016 Al Hindi, 2019
<i>Statistical Analysis</i>	Pooled RRs and corresponding 95% confidence intervals (CI) for perfusion-related complications were calculated by the Mantel–Haenszel test in RevMan 5.3. The heterogeneity among studies was evaluated with the I2 test. When I2 ranged from 0 to 30% a fixed- effects model was used. When the value of I2 exceeded 30%, a random effect model was used.	The Mantel-Haenszel test in R was used to calculate Pooled RRs and corresponding 95% confidence intervals (CI) for perfusion-related complications. The heterogeneity among studies was analyzed by using the I2 test. We used a fixed-effects model for I2 ranging from 0 to 30%; in the case of I2> 30%, we applied a random effect model using the DerSimonian & Liard method.
<i>N. of flaps. TOTAL</i> <i>- N. of cases</i> <i>- N. of controls</i>	1376 - 842 - 534	3094 - 1279 - 1815
<i>Total flap loss</i>	RR: 0.97, 95% CI: 0.36-2.57; p = 0.94	RR: 0.31, 95% CI: 0.11-0.85; p = 0.023
<i>Partial flap lost</i>	RR: 0.59, 95% CI: 0.18-1.94; p = 0.39	RR: 0.50, 95% CI: 0.30-0.84; p = 0.008
<i>Fat necrosis</i>	RR: 0.87, 95% CI: 0.58-1.30; p = 0.49	RR: 0.83, 95% CI: 0.63-1.10; p = 0.19
<i>Venous congestion</i>	RR: 0.06, 95% CI: 0.01-0.51; p = 0.01	RR: 0.12, 95% CI: 0.04-0.34; p < 0,001
<i>Conclusion</i>	“The present review demonstrated that superdrainage using SIEV reduces the risk of flap congestion notably, while having little influence on flap survival. With regard to partial flap necrosis including partial flap loss and fat necrosis, general trends toward decreased risks were observed.	“In conclusion, our meta-analysis shows, that venous super-drainage, i.e. performing a second venous anastomosis between the superficial venous system and a recipient vein, provides a statistical advantage in terms of venous congestion and related complications in DIEP flaps

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	However, statistical significance was not achieved and further studies would be needed."	for breast reconstruction."
--	--	-----------------------------

TABLE 2. Comparison between our meta-analysis and the one from Lee and Mun

RR: Risk Ratio. Cases: flaps with a second venous anastomosis. Controls: only one venous anastomosis

The 7 studies that were not analyzed in the previous published metanalysis by Lee, 2017 appear in **bold**.

In the fourth row, the results with statistical evidence appear in **bold**.