

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Role of prenatal magnetic resonance imaging in fetuses with isolated mild or moderate ventriculomegaly in the era of neurosonography: international multicenter study

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Di Mascio D., Khalil A., Thilaganathan B., Rizzo G., Buca D., Liberati M., et al. (2020). Role of prenatal magnetic resonance imaging in fetuses with isolated mild or moderate ventriculomegaly in the era of neurosonography: international multicenter study. *ULTRASOUND IN OBSTETRICS & GYNECOLOGY*, 56(3), 340-347 [10.1002/uog.21974].

Availability:

This version is available at: <https://hdl.handle.net/11585/781248> since: 2020-11-18

Published:

DOI: <http://doi.org/10.1002/uog.21974>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Role of prenatal magnetic resonance imaging in fetuses with isolated mild or moderate ventriculomegaly in the era of neurosonography: a multicenter study

The ENSO working group

Corresponding Author:

Francesco D'Antonio, MD, PhD
Fetal Medicine Unit
Department of Medical and Surgical Sciences,
Department of Obstetrics and Gynecology
University of Foggia
Viale L. Pinto, 71100 Foggia, Italy
E-mail address: francesco.dantonio@unifg.it

Short title: MRI in fetal ventriculomegaly

Keywords: ventriculomegaly, central nervous system, fetal magnetic resonance imaging, MRI, fetal ultrasound, neurosonography, prenatal diagnosis.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/uog.21974

CONTRIBUTION

What are the novel findings of this work?

The rate of fetal anomalies missed at US and detected only at fetal MRI is lower than what was previously reported in the published literature. The large majority of anomalies detected exclusively on MRI involve mainly migration disorders and hemorrhage which can be difficult to detect on US and tend to have a later presentation during pregnancy

What are the clinical implications of this work?

This is the largest study exploring the role of fetal brain MRI in detecting associated anomalies not diagnosed on US in fetuses with mild and moderate VM. This study support the practice of MRI assessment in every fetus with a prenatal diagnosis of VM, although parents can be reassured of the low risk of associated anomalies, when VM is isolated.

ABSTRACT

Objectives: To assess the role of fetal magnetic resonance imaging (MRI) in detecting associated anomalies in fetuses presenting with mild and moderate isolated ventriculomegaly (VM) undergoing multiplanar ultrasound (US) evaluation of fetal brain.

Methods: Multicenter, retrospective, cohort study involving 15 referral fetal medicine centers in Italy, United Kingdom, and Spain. Inclusion criteria were fetuses affected by isolated VM on US, defined as VM with normal karyotype and no other additional central nervous system (CNS) and extra-CNS anomalies on US, undergoing detailed assessment of fetal brain via a multiplanar approach as suggested by ISUOG guidelines on fetal neurosonogram. The primary outcome of the study was to report the rate of additional CNS anomalies detected exclusively at prenatal MRI and missed at US, while the secondary aim was to estimate the incidence of additional anomalies detected exclusively after birth and missed at prenatal imaging (US and MRI). Sub-group analyses according to the gestational age at MRI (< vs \geq 24 weeks of gestation), laterality of VM (unilateral vs bilateral), and severity of dilatation (mild vs moderate VM) were also performed.

Results: Five-hundreds and fifty-six fetuses with a prenatal diagnosis of isolated fetal VM on US were included in the analysis. Additional structural anomalies were detected at prenatal MRI and missed at US in 5.4% (95% CI 3.8-7.6) of cases. When considering the type of anomalies, supratentorial intra-cranial hemorrhage was detected on MRI in 26.7% of fetuses, while polymicrogyria and lissencephaly in 20.0% and 13.3% of cases, respectively. Hypoplasia of the corpus callosum was detected at MRI in 6.7% of cases, while dysgenesis in 3.3%. Fetuses with an associated anomaly detected only at MRI were more likely to have moderate compared to mild VM (60.0% vs 17.7%, $p < 0.001$), while there was no significant difference between the proportion of cases with bilateral VM between the two groups ($p = 0.2$). The results of the logistic regression analysis showed that maternal body mass index (OR: 0.85, 95% CI 0.7-0.99, $p = 0.030$), the presence of moderate VM (OR: 5.8, 95% CI 2.6-13.4, $p < 0.001$) and gestational age at MRI ≥ 24 weeks of gestation (OR: 4.1, 95% CI 1.1-15.3, $p = 0.038$) were independently associated with the probability of detecting associated anomalies at MRI. Associated anomalies were detected exclusively at birth and missed at prenatal imaging in 3.8% of cases.

Conclusions: The rate of associated fetal anomalies missed at US and detected only at fetal MRI in fetuses with isolated mild and moderate VM undergoing neurosonography is lower than that

previously reported. The large majority of these anomalies are difficult to detect on ultrasound. The findings from this study support the practice of MRI assessment in every fetus with a prenatal diagnosis of VM, although parents can be reassured of the low risk of associated anomalies, when VM is isolated on US.

Accepted Article

INTRODUCTION

Ventriculomegaly (VM) is the most common brain anomaly diagnosed during fetal life and encompasses a large spectrum of conditions characterized by a dilatation of the lateral ventricles of the brain, typically defined as greater than 10 mm at the level of the atria.¹⁻⁷

VM is frequently classified according to the degree of ventricular dilatation as mild (10-12 mm), moderate (13-15 mm), or severe (>15 mm), with higher degrees of dilation being associated with an increased risk of neurodevelopmental delay.⁸

The cause, severity and presence of associated anomalies are the major determinants in predicting the outcome of fetuses affected by VM. Thus, the main issue when approaching a fetus with VM is to rule out central nervous system (CNS) and extra-CNS anomalies.⁶ The International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) recommends that every fetus presenting with isolated VM on ultrasound should undergo a thorough multiplanar examination through axial, coronal and sagittal views of the fetal brain, possibly performed with a high-resolution vaginal probe. Moreover, a detailed assessment of the entire fetal anatomy, including fetal echocardiography, should also be undertaken.⁹

Ultrasound (US) is the primary imaging tool for the assessment of fetal brain, while fetal magnetic resonance imaging (MRI) has been shown to detect additional anomalies in 20% to 50% of cases.¹⁰⁻¹¹

In a recent systematic review, we found that in fetuses diagnosed with isolated VM, the rate of CNS anomalies detected exclusively on MRI was lower than that previously reported when a multiplanar assessment of the fetal brain is undertaken.^{9,12} However, the heterogeneity among the included studies make the results of this systematic review difficult to generalize.

The primary aim of this study was to evaluate the role of fetal MRI in detecting associated anomalies in fetuses presenting with isolated mild and moderate VM undergoing multiplanar neurosonography and to ascertain whether the incidence of such anomalies is dependent upon the degree and laterality of ventricular dilatation and gestational age at MRI. The secondary aim was to estimate the incidence of additional anomalies detected exclusively after birth and missed at prenatal imaging (US and MRI).

METHODS

Study design and participants

This was a multicenter, retrospective, cohort study involving 15 referral centers in Italy, United Kingdom, and Spain. The study included pregnant women who had fetal brain MRI following the diagnosis of isolated VM performed by neurosonography from January 2010 to March 2019. Only cases with at post-natal imaging or post-mortem examination, in case of termination of pregnancy or fetal demise, were included. The clinical records were examined and data were collected in a dedicated merged database.

Inclusion criteria

- Fetuses affected by isolated mild (10-12 mm), moderate (13-15 mm) VM at US, defined as VM with no other additional CNS and extra-CNS on the scan
- Detailed assessment of fetal brain via a multiplanar approach as suggested by 2007 ISUOG guidelines on fetal neurosonogram⁹
- Detailed fetal assessment including echocardiography
- Normal karyotype (including chromosomal microarray, when available)
- Negative infection screening (including CMV and Toxoplasmosis)
- Maternal Age \geq 18 years
- Gestational age \geq 18 weeks

Exclusion criteria

- Fetuses affected by severe ($>$ 15 mm) VM at US cases of severe VM
- Cases affected by chromosomal anomalies
- Cases affected by additional CNS and extra-CNS anomalies at the time of initial diagnosis
- Cases affected by congenital infections
- US protocol unclear or unavailable

Outcomes

The primary outcome of the study was to establish the rate of additional CNS anomalies detected exclusively on MRI and confirmed at birth in fetuses with a prenatal diagnosis of isolated VM

following dedicated neurosonography. The secondary aim was to estimate the incidence of additional anomalies detected exclusively after birth and missed at prenatal imaging (US and MRI). Sub-group analyses according to the gestational age at MRI (< vs \geq 24 weeks of gestation), laterality of VM (unilateral vs bilateral VM), and the severity of dilatation (mild vs moderate VM) were also performed.

For the purpose of the analysis, additional CNS anomalies were classified into:

- Callosal anomalies, including complete and partial agenesis (ACC), hypoplasia (HCC) and dysgenesis of the corpus callosum
- Septal anomalies, including all the anomalies characterized by a primary defect involving the septum pellucidum with a normally present corpus callosum
- Posterior fossa anomalies, including all defects involving the cerebellar vermis and/or hemispheres
- Intra-ventricular hemorrhage
- Cortical anomalies, including all abnormalities associated with a primary defect in neuronal migration towards the cortical surface of the brain
- Peri-ventricular heterotopia
- Other white matter anomalies
- Peri-ventricular cysts
- Complex brain anomalies, including all defects characterized by the presence of multiple intra-cranial anomalies
- Other cerebral anomalies

We did not consider biometric variation in brain structures, such as mega cisterna magna, increased or reduced degree of ventricular dilatation or of cranial size, as associated anomalies.

Statistical analysis

We investigated the relationship between the presence of ventriculomegaly-associated structural anomalies, assessed through fetal MRI (primary outcome) and a number of maternal and fetal

characteristics, including mother's age and body mass index (BMI), ventriculomegaly severity and laterality, gestational age at US and MRI assessment. As a secondary outcome, we investigated the relationship between the same maternal/fetal parameters and a postnatal diagnosis of ventriculomegaly-associated anomalies in fetuses with isolated VM (following negative prenatal imaging).

The potential association between all recorded maternal and fetal parameters and the two outcomes were first evaluated with standard univariate analyses (chi-squared test for categorical variables; Kruskal-Wallis test for continuous variables). For the secondary outcome, no multivariate analysis could be performed, because the number of newborns with a postnatal diagnosis of structural anomalies was limited to 11, and the likelihood of overfitting was too high.

As regards the primary outcome, we investigated the potential independent predictors of a fetal MRI diagnosis of ventriculomegaly-associated anomalies with a twofold approach. First, we performed a random-effect logistic regression, with hospital region as the cluster unit. A stepwise forward process was used for model building, and the following criteria were adopted for covariates selection, which were limited to four in every step of the analysis to reduce the risk of overfitting: (1) $p < 0.05$ at univariate analyses; (2) clinical significance; (3) the interval, expressed in weeks, between US and MRI examinations included a priori as a continuous variable. To avoid multicollinearity between the mean dilatation of cerebral ventricle (in mm) and the severity of ventriculomegaly (classified into "mild" and "moderate" according to the mm of dilatation), only the latter covariate was included in the model as a categorical variable. Standard post-estimation tests were used to check the validity of the final model, performing multicollinearity and influential observation analyses (using standardized residuals, change in Pearson and deviance chi-square).¹³⁻¹⁴ Second, all analyses were repeated after multiple imputation with bootstrap option for missing values ($m=5$, *mi* Stata command),¹³⁻¹⁴ as 37% of the mother's BMI values were not reported. The results of the complete model were very similar to those of the random-effect logistic regression, and only the results of the model without missing imputation have been shown to avoid redundancy. Statistical significance was defined as a two-sided p -value < 0.05 for all analyses,¹⁵ which were carried out using Stata, version 13.1 (Stata Corp., College Station, Texas, USA, 2013). This study was reported following the STROBE guidelines.¹⁶

RESULTS

Characteristics of the women

Five-hundred and fifty-six fetuses with a prenatal diagnosis of isolated fetal VM were included in the analysis. The general characteristics of the study population are shown in Table 1. The mean maternal age was 32.0 ± 5.9 years, while the mean BMI was 24.6 ± 4.1 . The mean gestational age at MRI was 26.7 ± 4.4 , with 30.9% of the scans performed before, while 69.1% performed at or after 24 weeks of gestation. Of the included cases, 36.5% (95% CI 32.6-40.4; 203/556) were affected by bilateral, while 63.5% (95% CI 59.4-67.4; 353/556) by unilateral VM. Finally, VM was mild (10-12 mm) in 80.0% (95% CI 76.5-83.2; 445/556) and moderate (13-15 mm) in 20.0% (95% CI 16.9-23.5; 111/556) of the cases.

Synthesis of the results

Table 2 shows the results of the primary and secondary outcomes of study. Additional structural anomalies were detected exclusively by MRI in 5.4% (95% CI 3.8-7.6; 30/556) of cases. When considering the type of anomalies, supra-tentorial intra-cranial hemorrhage was detected exclusively at MRI in 26.7% (8/30) of cases, while polymicrogyria and lissencephaly in 20.0% (6/30) and 13.3% (4/30) of cases, respectively. HCC was detected only at MRI in 6.7% (2/30) of cases, while dysgenesis in 3.3% (1/30).

The rate of associated anomaly detected exclusively at MRI was significantly higher in fetuses affected by moderate VM (60%), compared with fetuses affected by mild VM (17.7%) ($p < 0.001$), while there was no significant difference between the proportion of cases with bilateral VM between the two groups ($p = 0.2$). The majority (90.0%) of fetuses with associated anomalies had MRI performed at ≥ 24 weeks of gestation (Supplementary Table 1).

The results of the logistic regression analysis showed that maternal BMI (OR: 0.85, 95% CI 0.7-0.99, $p = 0.030$), the presence of moderate VM (OR: 5.8, 95% CI 2.6-13.4, $p < 0.001$) and gestational age at MRI ≥ 24 weeks of gestation (OR: 4.1, 95% CI 1.1-15.3, $p = 0.038$) were independently associated with the probability of detecting associated anomalies at MRI (Table 3).

After birth, 3.8% (11/278) of cases with an isolated VM confirmed at prenatal MRI had associated anomalies detected exclusively at post-natal imaging (cranial ultrasound and/or post-natal MRI). Mean ventricular dilatation was significantly higher in fetuses with a post-natal diagnosis of associated anomalies compared to those who had isolated VM confirmed at post-natal imaging (13.0 ± 1.5 vs 11.5 ± 1.3 , $p = 0.002$). Furthermore, fetuses with a post-natal diagnosis of associated anomalies were more likely to be affected by bilateral 3 decimal points (81.8% vs 50.0%, $p = 0.04$) and moderate (63.6% vs 18.7%, $p < 0.001$) VM compared to those who were confirmed to be isolated after birth (Supplementary Table 2).

DISCUSSION

Summary of the main findings

The findings of this study show that in fetuses with prenatal diagnosis of isolated VM examined using multiplanar neurosonography, the rate of additional structural anomalies detected exclusively by fetal brain MRI was 5.4%. The most common type of anomalies included supra-tentorial intracranial hemorrhage, and neuronal migration disorders. Factors independently associated with a higher incidence of additional findings detected exclusively at fetal MRI were gestational age at MRI ≥ 24 weeks, moderate ventricular dilatation and maternal BMI. Finally, the rate of associated anomalies detected exclusively after birth and missed at prenatal imaging was 3.8%.

Strengths and limitations

Large sample size, inclusion of cases examined using a multiplanar approach as proposed by ISUOG guidelines on fetal neurosonography and stratification of the analyses according to the degree and laterality of ventricular dilatation represent the main strengths of this study. The retrospective non-randomized design represents the main limitation of the study and led to challenges in obtaining all the details on the imaging for all the fetuses in the participating centers, with some incomplete follow-up and some missing data, mostly related to the postnatal MRI or US and karyotyping. Furthermore, although 75.5% of fetuses underwent MRI two weeks or less after neurosonogram, 24.5% of fetuses underwent MRI after three to more than six weeks after US; therefore, it is likely that a prolonged interval between US and MRI may have somehow influenced our results. Finally, the present study included only cases having neurosonography performed in each participating center without exploring its feasibility according to different maternal and fetal characteristics.

Implications for clinical practice and research

Isolated VM represents a considerable diagnostic dilemma as it can be an apparently benign finding, but can also be associated with chromosomal abnormalities, congenital infections, cerebral vascular accidents or hemorrhage, and other fetal cerebral and extracerebral abnormalities.⁵⁻⁷ Even when isolated, VM has been reported to increase the risk of neurodevelopmental delay in childhood and adolescence. The prevalence of neurodevelopmental disorders in fetuses with both unilateral

and bilateral mild VM has been reported to be only slightly higher than that of the general population.¹⁷⁻¹⁹ Conversely, neurodevelopmental delay is higher in case of VM associated with additional CNS anomalies, and in case of severe ventricular dilatation.²⁰⁻²²

The ISUOG guidelines on sonographic examination of the fetal CNS suggest that a multiplanar assessment of fetal head should be always performed when a suspicion of brain anomaly is raised at the basic US examination.⁹ Furthermore, in a recent consult series on mild VM, the Society for Maternal-Fetal Medicine (SMFM) has stated that MRI may be considered in cases of mild or moderate fetal VM, although it may be of less value if the women undergone detailed US assessment by an individual with specific experience and expertise in sonographic imaging of the fetal brain.⁸

MRI is the imaging technique of choice in analyzing brain anomalies postnatally. Compared to US, MRI has a better regional resolution, thus being theoretically superior to US in detecting abnormalities of the cortical development. Based on these findings, the current practice today suggests a prenatal MRI examination in the late second, early third trimester of pregnancy, although there is no complete agreement among different authors regarding the need and time of MRI examination. However, the large majority of studies on MRI does not specify which type of ultrasound imaging protocol was adopted to assess fetal brain, and this is fundamental as most of CNS anomalies cannot be easily detected on the standard axial plane of fetal brain.^{12,23-29}

In 2017, a large prospective, multicenter study involving 16 centers across the United Kingdom (the MERIDIAN Study) was designed to evaluate the diagnostic and clinical role of fetal MRI in fetuses with a previous US with a suspicion or diagnosis of CNS anomaly.¹¹ This study showed a much greater diagnostic accuracy of MRI compared to US (93% versus 68%), with additional findings detected exclusively on MRI in 49% of brain abnormalities and in up to 19.4% when focusing on the subgroup analysis of fetuses with mild and moderate VM.²³ Of note, the rate of associated callosal anomalies detected exclusively on MRI was lower in our cohort compared to the MERIDIAN study (10% versus 55% of cases of failed commissuration).²³

In the present study, the large majority of anomalies detected exclusively on prenatal MRI included neuronal migration (lissencephaly, heterotopia) or acquired anomalies (i.e. hemorrhage) which could be difficult to diagnose on ultrasound.³⁰⁻³⁴ In this scenario, fetal MRI should always be performed in order to rule out associated anomalies which can be potentially missed on ultrasound. Nevertheless, parents should be reassured regarding the low risk of such additional anomalies.

In the present study, MRI performed at or after 24 weeks was an independent factor which was associated with the risk of additional anomalies detected exclusively at MRI. This finding might appear quite intuitive, as some abnormalities, such as malformations of cortical development, migration disorders and hemorrhage can become evident only later during pregnancy. Not surprisingly, the most frequent type of anomalies detected only at MRI in this study were hemorrhage and migration disorders, anomalies presenting mostly after the second trimester of pregnancy.³⁵ However, some authors have suggested that early MRI (before 24 weeks of gestations) may be reliable and has similar accuracy as MRI performed later in gestation,¹¹⁻¹² with the advantage of an early diagnosis which can allow parents the option of earlier diagnosis, especially in countries where termination of pregnancy is not legal beyond the second trimester.

Conclusions

The rate of fetal anomalies missed at US and detected only at fetal MRI is lower than what was previously reported in the published literature when a multiplanar examination of fetal brain performed. The large majority of anomalies detected exclusively on MRI involve mainly migration disorders and hemorrhage which cannot be easily detected on ultrasound, thus highlight the need for an MRI examination in fetuses with a prenatal diagnosis of VM undergoing neurosonography. Future research investigating prenatal diagnosis of isolated VM should aim at a multicenter prospective approach mostly to establish the optimal timing and the frequency of MRI examinations.

Disclosure:

Authors report no conflict of interest.

The European NeuroSONography (the ENSO) working group

Daniele Di Mascio*,^{1,2} Asma Khalil,^{3,4} Basky Thilaganathan,³ Giuseppe Rizzo,^{5,6} Danilo Buca,⁷ Marco Liberati,⁷ Claudio Celentano,⁷ Karen Melchiorre,⁸ Massimo Caulo,⁹ Gianluigi Pilu,¹⁰ Ginevra Salsi,¹⁰ Francesco Toni,¹¹ Tamara Stampalija,^{12,13} Iliaria Fantasia,¹² Giulia Luise,¹² Massimo Gregori,¹⁴ Paolo Volpe,¹⁵ Claudiana Olivieri,¹⁵ Antonella Giacotti,¹ Valentina D'Ambrosio,¹ Roberto Brunelli,¹ Pierluigi Benedetti Panici,¹ Lucia Manganaro,¹⁶ Amanda Antonelli,¹⁶ Giada Ercolani,¹⁶ Lucia Pasquini,¹⁷ Giulia Masini,¹⁷ Marco Di Maurizio,¹⁸ Christoph Lees,^{19,20} Gabriella Bracalente,²¹ José Morales-Roselló,^{22,23} Gabriela Loscalzo,²² Gabriele Saccone,²⁴ Luigi Carbone,²⁴ Laura Sarno,²⁴ Giuseppe Maria Maruotti,²⁴ Fulvio Zullo,²⁴ Tullio Ghi,²⁵ Tiziana Frusca,²⁵ Andrea Dall'Asta,^{19,20,25} Nicola Volpe,²⁵ Francesca Ormitti,²⁶ Silvia Buongiorno,^{22,23,27} Marco De Santis,²⁷ Luisa D'Oria,²⁷ Antonio Lanzone,²⁷ Federico Prefumo,^{29,29} Lorenzo Pinelli,³⁰ Emma Bertucci,³¹ Filomena Giulia Sileo,³¹ Maria Elena Flacco,³² Lamberto Manzoli,³² Iliaria Giangiordano,³³ Luciana Masticci,³⁴ Gabriella Meccariello,³⁴ Lorenzo Vasciaveo,³⁴ Luigi Nappi,³⁴ Alessandra Familiari,³⁵ Giovanni Scambia,²⁷ Vincenzo Berghella,² Francesco D'Antonio**,³⁴

*first author; **last author.

- 1: Department of Maternal and Child Health and Urological Sciences, Sapienza University of Rome, Italy
- 2: Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Sidney Kimmel Medical College of Thomas Jefferson University, Philadelphia, USA
- 3: Fetal Medicine Unit, Saint George's Hospital, London, United Kingdom
- 4: Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute, St George's University of London, United Kingdom
- 5: Division of Maternal and Fetal Medicine, Ospedale Cristo Re, University of Rome Tor Vergata, Rome, Italy
- 6: Department of Obstetrics and Gynecology The First I.M. Sechenov Moscow State Medical University; Moscow Russia
- 7: Department of Obstetrics and Gynecology, University of Chieti, Italy
- 8: Department of Obstetrics and Gynecology, Santo Spirito Hospital, Pescara, Italy
- 9: Department of Neuroscience, Imaging and Clinical Sciences, Institute for Advanced Biomedical Technologies, "G. D'Annunzio" University of Chieti, Italy
- 10: Unit of Obstetrics, Department of Medical and Surgical Sciences, University of Bologna, Italy
- 11: IRCCS Istituto delle Scienze Neurologiche UOC Neuroradiologia, Bologna, Italy
- 12: Unit of Fetal Medicine and Prenatal Diagnosis, Institute for Mother and Child Health IRCCS Burlo Garofolo, Trieste, Italy
- 13: Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste, Italy.
- 14: Unit of Paediatric Radiology, Institute for maternal and child health IRCCS Burlo Garofolo
- 15: Fetal Medicine Unit, Di Venere and Sarcone Hospitals, Bari, Italy
- 16: Department of Radiology, Anatomic-Pathology and Oncology, Sapienza University of Rome, Italy
- 17: Fetal Medicine Unit, Department for Women and Children Health, AOU Careggi, Florence, Italy
- 18: UOC Radiodiagnostica, AOU Meyer, Florence, Italy
- 19: Department of Surgery and Cancer, Institute of Reproductive and Developmental Biology, Imperial College London, United Kingdom
- 20: Centre for Fetal Care, Queen Charlotte's and Chelsea Hospital, Imperial College Healthcare NHS Trust, London, United Kingdom
- 21: Fetal Medicine Unit, Department of Obstetrics and Gynecology, Osp. Cà Foncello Treviso, Italy
- 22: Servicio de Obstetricia, Hospital Universitario y Politécnico La Fe, Valencia, Spain
- 23: Departamento de Pediatría Obstetricia y Ginecología, Universidad de Valencia, Spain
- 24: Department of Neuroscience, Reproductive Sciences and Dentistry, School of Medicine, Federico II University of Naples, Italy
- 25: Department of Medicine and Surgery, Unit of Surgical Sciences, Obstetrics and Gynecology, University of Parma, Italy
- 26: Department of Radiology - Azienda Ospedaliera Universitaria di Parma, Italy
- 27: Department of "Scienze della Salute della Donna, del Bambino e di Sanità Pubblica" - Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy
- 28: Division of Obstetrics and Gynecology, Department of Clinical and Experimental Sciences, University of Brescia, Italy

- 29: Division of Obstetrics and Gynecology, ASST Spedali Civili, Brescia, Italy
- 30: Neuroradiology Unit, Paediatric Neuroradiology Section, ASST Spedali Civili, Brescia, Italy
- 31: Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of Modena and Reggio Emilia, AOU Policlinico, Modena, Italy
- 32: Department of Medical Sciences, University of Ferrara, Italy
- 33: Department of Child and Adolescent Neuropsychiatry, Local Health Unit of Avezzano-Sulmona-L'Aquila, Sulmona, Italy
- 34: Fetal Medicine Unit, Department of Medical and Surgical Sciences, University of Foggia, Italy
- 35: Department of Obstetrics and Gynecology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy.

REFERENCES

1. Achiron R, Schimmel M, Achiron A, Mashiach S. Fetal mild idiopathic lateral ventriculomegaly: is there a correlation with fetal trisomy? *Ultrasound Obstet Gynecol* 1993; **3**:89-92.
2. Alagappan R, Browing PD, Laorr A, McGahan JP. Distal lateral ventricular atrium: reevaluation of normal range. *Radiology* 1994; **193**:405-408.
3. Nomura ML, Barini R, De Andrade C, Milanez H, Simoni RZ, Peralta CF, Machado IN, Zambelli H, Maio KT. Congenital hydrocephalus: gestational age and neonatal outcomes. *Arch Gynecol Obstet* 2010; **282**:607-611.
4. Garel C, Luton D, Oury JF, Gressens P. Ventricular dilatations. *Childs Nerv Syst* 2003; **19**:517-523.
5. Signorelli M, Tiberti A, Valsariati D, Molin E, Cerri V, Groli C, Bianchi UA. Width of fetal lateral ventricular atrium between 10 and 12 mm: a simple variation of the norm? *Ultrasound Obstet Gynecol* 2004; **23**:14-18.
6. Melchiorre K, Bhide A, Gika AD, Pilu G, Papageorghiou AT. Counseling in isolated mild fetal ventriculomegaly. *Ultrasound Obstet Gynecol* 2009; **34**:212-224.
7. Shizuo OI. Controversies in definition and classification of hydrocephalus. *Neurol Med Chir* 2010; **50**:859-869.
8. Society for Maternal-Fetal Medicine (SMFM); Fox NS, Monteagudo A, Kuller JA, Craigo S, Norton ME. Mild fetal ventriculomegaly: diagnosis, evaluation, and management. *Am J Obstet Gynecol* 2018; **219**:B2-B9.
9. ISUOG Guidelines. Sonographic examination of the fetal central nervous system: guidelines for performing the 'basic examination' and the 'fetal neurosonogram' *Ultrasound Obstet Gynecol* 2007; **29**:109–116.
10. Rossi AC, Prefumo F. Additional value of fetal magnetic resonance imaging in the prenatal diagnosis of central nervous system anomalies: a systematic review of the literature. *Ultrasound Obstet Gynecol* 2014; **44**:388–393.
11. Griffiths PD, Bradburn M, Campbell MJ, Cooper CL, Graham R, Jarvis D, Kilby MD, Mason G, Mooney C, Robson SC, Wailoo A; MERIDIAN collaborative group. Use of

- MRI in the diagnosis of fetal brain abnormalities in utero (MERIDIAN): a multicentre, prospective cohort study. *Lancet* 2017; **389**:538–546.
12. Di Mascio D, Sileo FG, Khalil A, Rizzo G, Persico N, Brunelli R, Giancotti A, Benedetti Panici P, Acharya G, D'Antonio F. Systematic review and meta-analysis on the role of prenatal magnetic resonance imaging in the era of fetal neurosonography: mild and moderate ventriculomegaly. *Ultrasound Obstet Gynecol* 2019; **54**:164-171.
13. Pizzi C, Costa GM, Santarella L, Flacco ME, Capasso L, Bert F, Manzoli L. Depression symptoms and the progression of carotid intima-media thickness: A 5-year follow-up study. *Atherosclerosis* 2014; **233**:530-536.
14. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011; **30**:377-399.
15. McNamee R. Regression modelling and other methods to control confounding. *Occup Environ Med* 2005; **62**:500-506.
16. Von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP for the STROBE Initiative. The strengthening the reporting of the observational studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007; **370**:1453-1457.
17. Scala C, Familiari A, Pinas A, Papageorgiou AT, Bhide A, Thilaganathan B, Khalil A. Perinatal and long-term outcome in fetuses diagnosed with isolated unilateral ventriculomegaly: systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2017; **49**:450-459.
18. Pagani G, Thilaganathan B, Prefumo F. Neurodevelopmental outcome in isolated mild fetal ventriculomegaly: Systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2014; **44**:254-260.
19. Thorup E, Jensen LN, Bak GS, Ekelund CK, Greisen G, Jorgensen DS, Hellmuth SG, Wulff C, Petersen OB, Pedersen LH, Tabor A. Neurodevelopmental disorder in children believed to have isolated mild ventriculomegaly prenatally. *Ultrasound Obstet Gynecol* 2019; **54**:182-189.
20. Weichert J, Hartge D, Krapp M, Germer U, Gembruch U, Axt-Flidner R. Prevalence, characteristics and perinatal outcome of fetal ventriculomegaly in 29.000 pregnancies

followed at a single institution. *Fetal Diagn Ther* 2010; **27**:142–148.

21. Carta S, Kaelin Agten A, Belcaro C, Bhide A. Outcome of fetuses with prenatal diagnosis of isolated severe bilateral ventriculomegaly: systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2018; **52**:165-173.
22. Dall'Asta A, van Oostrum NHM, Basheer SN, Paramasivam G, Ghi T, Galli L, Groenenberg IA, Tangi A, Accorsi P, Echevarria M, Angeles Rodríguez Perez M, Albaiges Baiget G, Prefumo F, Frusca T, Go AT, Lees CC. Etiology and Prognosis of Severe Ventriculomegaly Diagnosed at Late Gestation. *Ultraschall Med* 2018; **39**:675-689.
23. Griffiths PD, Brackley K, Bradburn M, Connolly DJ, Gawne-Cain ML, Griffiths DI, Kilby MD, Mandefield L, Mooney C, Robson SC, Vollmer B, Mason G. Anatomical subgroup analysis of the MERIDIAN cohort: ventriculomegaly. *Ultrasound Obstet Gynecol* 2017; **50**:736-744.
24. Griffiths P, Reeves M, Morris J, Mason G, Russell S, Paley M, Whitby E. A prospective study of fetuses with isolated ventriculomegaly investigated by antenatal sonography and in utero MR imaging. *Am J Neuroradiol* 2010; **31**:106–111.
25. Lavongtheung A, Jedraszak G, Naepels P, Tourneux P, Gondry-Jouet C, Le MoingAG, Gondry J, Chevreau J. Should isolated fetal ventriculomegaly measured below 12 mm be viewed as a variant of the norm? Results of a 5-year experience in a prenatal referral center. *J Matern Fetal Neonatal Med* 2018; **31**:2325–2331.
26. Mehlhorn, AJ, Morin CE, Wong-You-Cheong JJ, Contag SA. Mild fetal cerebral ventriculomegaly: prevalence, characteristics, and utility of ancillary testing in cases presenting to a tertiary referral center. *Prenat Diagn* 2017; **37**:647–657.
27. Kandula T, Fahey M, Chalmers R, Edwards A, Shekleton P, Teoh M, Clark J, Goergen SK. Isolated ventriculomegaly on prenatal ultrasound: what does fetal MRI add? *J Med Imaging Radiat Oncol* 2015; **59**:154–162.
28. Morris JE, Rickard S, Paley MNJ, Griffiths PD, Rigby A, Whitby EH. The value of in-utero magnetic resonance imaging in ultrasound diagnosed foetal isolated cerebral ventriculomegaly. *Clin Radiol* 2007; **62**:140–144.

29. Gezer NS, Gezer C, Ekin A, Yesilirmak DC, Solmaz U, Dogan A, Guleryuz H. Obstetric and neurodevelopmental outcome in fetal cerebral ventriculomegaly. *Clin Exp Obstet Gynecol* 2016; **43**:490–494.
30. Dhouib A, Blondiaux E, Moutard ML, Billette de Villemeur T, Chalard F, Jouannic JM, Ducou le Pointe H, Garel C. Correlation between pre- and postnatal cerebral magnetic resonance imaging. *Ultrasound Obstet Gynecol* 2011; **38**:170-178.
31. Pugash D, Hendson G, Dunham CP, Dewar K, Money DM, Prayer D. Sonographic assessment of normal and abnormal patterns of fetal cerebral lamination. *Ultrasound Obstet Gynecol* 2012; **40**:642-651.
32. Greco P, Resta M, Vimercati A, Dicuonzo F, Loverro G, Vicino M, Selvaggi L. Antenatal diagnosis of isolated lissencephaly by ultrasound and magnetic resonance imaging. *Ultrasound Obstet Gynecol* 1998; **12**:276-279.
33. Blondiaux E, Sileo C, Nahama-Allouche C, Moutard ML, Gelot A, Jouannic JM, Ducou le Pointe H, Garel C. Periventricular nodular heterotopia on prenatal ultrasound and magnetic resonance imaging. *Ultrasound Obstet Gynecol* 2013; **42**:149–155.
34. Ghi T, Simonazzi G, Perolo A, Savelli L, Sandri F, Bernardi B, Santini D, Bovicelli L, Pilu G. Outcome of antenatally diagnosed intracranial hemorrhage: case series and review of the literature. *Ultrasound Obstet Gynecol* 2003; **22**:121-130.
35. Malinger G, Lerman-Sagie T, Watemberg N, Rotmensch S, Lev D, Glezerman M. A normal second-trimester ultrasound does not exclude intracranial structural pathology. *Ultrasound Obstet Gynecol* 2002; **20**:51-56.

Table 1. Selected gestational and fetal characteristics in a sample of singleton pregnancies with a prenatal sonographic diagnosis of ventriculomegaly

<i>variables</i>	N = 556
<i>General characteristics:</i>	
Mean maternal age in years (SD)	32.0 (5.9)
Mean maternal BMI in kg/m ² (SD)	24.6 (4.1)
Gestational age at prenatal MRI diagnosis of additional anomalies in weeks:	
- Mean gestational age at diagnosis (SD)	26.7 (4.4)
- Diagnosis <24 weeks, %	30.9
- Diagnosis ≥24 weeks, %	69.1

Interval between prenatal US and MRI examinations in weeks:

- Mean interval (SD)	2.2 (2.5)
- ≤2 weeks, %	75.5
- 3-5 weeks, %	11.3
- ≥6 weeks, %	13.2

Characteristics of fetal ventriculomegaly:

Bilateral ventriculomegaly, %	36.5
Severity, %	
- Mild	80.0
- Moderate	20.0
Mean ventricular dilatation in mm (SD):	
- Overall	11.6 (1.3)
- Mild ventriculomegaly	11.1 (0.7)
- Moderate ventriculomegaly	13.8 (0.8)
Mean dilatation of the controlateral ventricle in mm (SD)**	11.8 (1.4)

SD: Standard deviation; US: ultrasound; MRI, magnetic resonance imaging

*Data available only for 425 and 148 fetuses, respectively. **Based on 203 fetuses with bilateral ventriculomegaly

Table 2. Primary and secondary outcomes

Outcomes

Fetuses with additional structural anomalies detected through prenatal MRI, %	5.4
---	-----

Type of prenatal-diagnosed anomaly, % *	
- Hemorrhage	26.7
- Polymicrogyria	20.0
- Lissencephaly	13.3
- Corpus Callosum hypoplasia	6.7
- Periventricular heterotopia	6.7
- Others	26.6
Newborn with additional structural anomalies detected through postnatal MRI, % **	3.8

* N=30 fetuses with additional anomalies detected through prenatal MRI

** Analyses restricted to 289 newborn (both the fetuses with a prenatal diagnosis of structural anomaly and the newborn without a postnatal MRI exam were excluded).

Table 3. Logistic regression models evaluating the potential independent predictors of a prenatal MRI diagnosis of ventriculomegaly-associated anomalies

<i>Covariates</i>	Adjusted OR (95% CI)	P value
<i>Model 1 *</i>		
Maternal BMI, 1-unit increase	0.85 (0.74-0.99)	0.030
Severity of ventriculomegaly, moderate vs mild	5.84 (2.55-13.4)	<0.001
Interval between US and MRI assessment, 1-week increase	1.00 (0.84-1.21)	0.919
Gestational age at MRI diagnosis, \geq versus $<$ 24 weeks	4.07 (1.09-15.3)	0.038

* Random-effect logistic regression with Hospital region as the cluster level.