ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Contrasting patterns of tree features, lichen, and plant diversity in managed and abandoned old-growth chestnut orchards of the northern Apennines (Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*
Pezzi, G., Gambini, S., Buldrini, F., Ferretti, F., Muzzi, E., Maresi, G., et al. (2020). Contrasting patterns of tree features, lichen, and plant diversity in managed and abandoned old-growth chestnut orchards of the northern Apennines (Italy). FOREST ECOLOGY AND MANAGEMENT, 470-471, 1-10 [10.1016/j.foreco.2020.118207].

*Availability:*
This version is available at: https://hdl.handle.net/11585/783156 since: 2021-03-01

*Published:*

DOI: http://doi.org/10.1016/j.foreco.2020.118207

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/).
When citing, please refer to the published version.

(Article begins on next page)

07 October 2026

# Contrasting patterns of tree features, lichen, and plant diversity in managed and abandoned old-growth chestnut orchards of the northern Apennines (Italy)

Giovanna Pezzi[a], Simone Gambini[a], Fabrizio Buldrini[a]*, Fabrizio Ferretti[b], Enrico Muzzi[c], Giorgio Maresi[d], Juri Nascimbene[a]

[a] Department of Biological, Geological, and Environmental Sciences, Università di Bologna. Via Irnerio 42, 40126 Bologna, Italy (giovanna.pezzi@unibo.it, fabrizio.buldrini@unibo.it, simone.gambini2@studio.unibo.it, juri.nascimbene@unibo.it)

[b] CREA Research Centre for Forestry and Wood, v.le Santa Margherita 80, 52100 Arezzo, Italy (fabrizio.ferretti@crea.gov.it)

[c] Department of Agricultural and Food Sciences, Università di Bologna. Viale G. Fanin 44, 40127 Bologna, Italy (enrico.muzzi@unibo.it)

[d] Centre for Technology Transfer, Fondazione Edmund Mach. Via E. Mach 1, I-38010 San Michele all'Adige, TN, Italy (giorgio.maresi@fmach.it)

* Corresponding author; fabrizio.buldrini@unibo.it

**Abstract**

In mountain regions of southern Europe, old-growth chestnut orchards maintained by traditional management were a key component of the economic, cultural, and ecological heritage. Currently, many stands are abandoned due to decreased economic sustainability even though, according to European policies, the loss of traditionally managed old-growth chestnut orchards should be contrasted to prevent biodiversity loss. In this study, we preliminarily mapped the remnants of old-growth chestnut orchards across a region of the northern Apennines (Italy) with a strong tradition of chestnut orchard cultivation. Then, we assessed the effects of management/abandonment in terms of

tree features (e.g. size, crown structure, health conditions), occurrence and abundance of target epiphytic lichens, and richness and composition of understory vegetation. Our results revealed contrasting patterns of tree features, lichen, and plant diversity in managed and abandoned old-growth chestnut orchards of the northern Apennines, supporting the view that traditional management is fundamental for the long-term maintenance of healthy veteran trees, the enhancement of epiphytic lichens related to old-growth conditions, and plant diversity. This indicates that 1000 years of 'chestnut civilization' represent a cultural heritage that benefits nature conservation, promoting a virtuous interplay between human activities and biodiversity. For this reason, policies aimed at sustaining traditional management in old-growth chestnut orchards are indispensable to avoid the degradation and loss of this habitat and its centuries-old cultural and ecological legacy.

**Keywords**

*Castanea sativa*; Chestnut diseases; Failure risk assessment; Intermediate disturbance hypothesis; *Caliciales*; *Lobaria pulmonaria*

## Introduction

In mountain regions of southern Europe, including Italy, *Castanea sativa* Mill. orchards were a key component of the economic, cultural and ecological heritage. Since the early Middle Ages, chestnut orchards provided one of the main staple foods (Pitte, 1986; Conedera et al., 2004; Squatriti, 2013), and influenced the lifestyle of mountain people, leading to a phenomenon called 'chestnut civilization' (Gabrielli, 1994; Arnaud et al., 1997; Conedera et al., 2004). These anthropogenic woods modelled the landscape and were maintained by traditional management, including mowing, grazing, and tree pruning.

Although a first decline coincided with the eighteenth-century Little Ice Age, on Italian mountains the massive abandonment of many chestnut orchards occurred after the mid-20$^{th}$ century. After World War II, strong social-economic changes caused an exodus towards cities, provoking the loss of traditional cultivation and lifestyles (Arnaud et al., 1997; Conedera and Krebs, 2008; Bounous, 2014). Besides these social dynamics, pests and pathogens acted as disturbance factors with both direct and indirect effects, i.e. influencing perceptions and attitudes of chestnut growers towards the cultivation (Turchetti et al., 2012). In the middle of the 20$^{th}$ century, the spread of *Cryphonectria parasitica* (Murr.) Barr, the agent of chestnut blight, forced the abandonment of many stands, which then started to evolve as mixed woods (Mondino, 1991; Romane et al., 1995; Arnaud et al., 1997; Paci et al., 2000; Conedera et al., 2001). In Italy, since the beginning of the 20$^{th}$ century, chestnut orchards have decreased by about 90%, from more than 608.000 ha (Vigiani, 1908) to 60.000 ha that are currently cultivated (Bounous, 2014).

According to the conservation polices of the European Union, the loss of traditionally managed old-growth chestnut orchards should be contrasted to prevent biodiversity loss (Arnaud et al., 1997; Piussi and Pettenella, 2000). Old-growth chestnut orchards with semi-natural undergrowth are included in the list of habitats worthy of conservation according to the Council Directive 92/43/EEC. In general, their conservation is likely dependent on moderate, but relatively continuous, traditional management (e.g., Gondard et al., 2001, 2007). In this perspective, they

represent a tangible expression of an intangible cultural heritage, and a tight bond between man and forest (see UNESCO Convention) that should be preserved.

Traditionally managed old-growth chestnut orchards may host species-rich plant communities (Gondard et al., 2001; Barbati and Marchetti, 2005), including species of conservation concern, such as several orchids or species related to dry grasslands, which may be threatened by abandonment. While management intensification (i.e. frequent mowing) is unlikely to occur due to the scarce economical value of this cultivation, abandonment is expected to drive major changes in plant community composition and richness (Nascimbene et al., 2014, 2016).

In old-growth chestnut orchards, the occurrence of veteran trees provides refugia to organisms that depend on this virtually missing habitat across European forests (Krebs et al., 2008). This is the case for epiphytic organisms that require long ecological continuity, in particular the forest lichen *Lobaria pulmonaria* (L.) Hoffm. In Italy, chestnut orchards are among the main habitats for this species, often hosting luxuriant populations (e.g. Matteucci et al., 2012) that were likely originated by propagules from the surrounding beech or oak forests. However, contrary to surrounding forests that may have been intensively exploited, chestnut orchards provide more suitable ecological conditions for this lichen. At the tree level, tree features (e.g. size, age, health conditions) are the main drivers for these species, while at the stand level forest management is the main driver with contrasting effects. Intensive management is usually highly detrimental, while abandonment of management leading to closed canopy stands may also negatively affect this species (Nascimbene et al., 2013a, 2013b, 2016; Brunialti et al., 2015). Besides local factors, also landscape features may influence the occurrence and abundance of this lichen, whose dispersal is mainly related to vegetative propagules over relatively short distances (Löbel et al., 2009). Due to its peculiar ecology and its easy detectability it is considered a valuable indicator of forest sites that host high lichen diversity and species of conservation concern (Nascimbene et al., 2010). Among these, species belonging to the order *Caliciales* are associated with old-growth stands characterized by long ecological continuity (Selva, 1994; Brunialti et al., 2015).

In this study, we preliminarily mapped the remnants of old-growth chestnut orchards across a region of the northern Apennines with a strong tradition of chestnut orchard cultivation. Then, we assessed the effects of management/abandonment in terms of tree features (e.g. size, crown structure, health conditions), occurrence and abundance of target epiphytic lichens (*L. pulmonaria*, *Caliciales*), and richness and composition of understory vegetation. Due to the low intensity practices applied in managed orchards, we expect trees to have different and healthier conditions compared to abandoned orchards. Moreover, we expect to find more suitable conditions for *L. pulmonaria* in managed compared to abandoned orchards, and more diverse and richer plant communities that also include species of conservation concern related to semi-open habitats. Finally, we expect that the abundance of *L. pulmonaria* could be a reliable indicator of species related to old-growth stands, as in the case of the lichens belonging to the *Caliciales*.

## Materials and Methods

### *Study area*

The study was carried out in a 1935 $km^2$ area that includes the chestnut belt of the Apennines between Bologna and Modena (centroid 44.2561645 N, 10.9453453 E; Fig. 1a). Its altitudinal range is about from 300 up to 1000 m a.s.l., from the deciduous *Quercus*-dominated forests to the *Fagus sylvatica* belt (Ubaldi et al., 1993). Chestnut orchards cover only 2% of this area and are closely linked to the local tradition and cultural heritage (Pezzi et al., 2019). Traditional management was carried out by regular mowing (twice or thrice a year) and pruning of trees, sometimes topped. Contrary to other Mediterranean areas, fire and *slupatura* (burn inside the trunk cavities and mechanical removal of damaged wood) were rarely used to sanitize decayed trunks (Fenaroli, 1946; Seijo et al., 2017, 2018). Burrs and pruned branches (and even leaves, when they were not used for other purposes) were generally removed, burned in single burning places with localized and controlled fires.

Thanks also to specific quality trade-marks and promotion initiatives, only *marroni* (i.e. the top quality fruits) have benefitted from a renewed interest in chestnut cultivation since the 1980s (Pezzi et al., 2017). In these forests, ink disease caused by *Phytophthora cambivora* (Petri) Buism. has been recorded since the last century and initially caused great concern about the survival of chestnut orchards due to the rapid spread of its attacks (Quattrocchi, 1938). Chestnut blight caused by *Cryphonectria parasitica* occurred for the last seventy years but has been controlled by the natural spread of hypovirulence (Turchetti et al., 2008). Asian chestnut gall wasp (*Dryocosmus kuriphilus* Yasumatsu) was first reported in 2007 and its parasitoid *Torymus sinensis* Kamijo was first released in 2010 (Vai et al., 2014).




**Figure 1.** a) Location of the study area (blue triangle) within Italy; b) thalli of *Lobaria pulmonaria* (L.) Hoffm. (grey-green to grey-brown, on the right) and *Lobarina scrobiculata* (Scop.) Nyl. (blue-

grey, on the left) on the trunk of a veteran chestnut tree (the thalli are wet); view of an old-growth

managed (c) and abandoned (d) chestnut stand. Photographs taken by G. Pezzi, July 2018 (a, b, c), and G. Maresi, July 2018 (d). [2-column fitting image, colour online only]

*Sample selection*

As the pattern of old-growth chestnut orchards was largely misunderstood and spatial information was almost lacking, we preliminarily conducted informal interviews to map old-growth stands within the study area (January 2017 to April 2018). We first turned to people in charge of chestnut associations, then to owners/managers and local experts. New contacts were added via snowball sampling (Hislop et al., 2004; Tongco, 2007). We stopped when no additional suitable stands were identified, or when the same stand had been previously mentioned. In total, 48 people were interviewed. These were asked to inform us of chestnut stands at least 0.2 ha wide and with a majority (> 70%) of trees with a diameter at breast height (DBH) at least of 1 m. Despite the lack of a strong correlation between diameter and age for these formations, according to the current literature a chestnut tree with a DBH of about 1 m is centuries old (Krebs et al., 2005; Temel et al., 2009). Preliminary surveys were carried out to check the congruence of the sites with the required standard. Then, each chestnut stand was mapped in QGIS 3.4 (www.qgis.org) by visual interpretation of current digital orthophotographs coupled to a forest map. Overall, 20 old-growth chestnut orchards were mapped. Their surface varied between 0.2 to 4.4 ha (Table 1), mainly with a northern aspect, at an altitude ranging between 600 and 1030 m. Slope was on average 18.3° (± 12.1°). Canopy cover was on average 82.5% (± 18.3%). 12 stands were still traditionally managed (11 through mowing, 1 through grazing) and 8 were abandoned (Fig. 1c,d). Furthermore, 17 were multi-varietal stands, while only 3 were mono-varietal stands (*marroni* stands).

The field survey was carried out in June-July 2018 to obtain information on the main tree features, on the occurrence and abundance of target epiphytic lichens (i.e. *L. pulmonaria* and *Caliciales*), and

undergrowth vegetation. Each stand was sampled through an 85 m transect, along which 10 chestnut trees were selected and five 5 × 5 m vegetation plots were regularly placed.

**Table 1**. General features of the 20 old-growth chestnut orchards. For each examined stand, centroid coordinates are given in WGS 84 as reference system.

| Stand | Lat. N (°) | Long. E (°) | Extent (ha) | Altitude (m a.s.l.) | Slope (°) | Exposure | Management |
|---|---|---|---|---|---|---|---|
| ALBE | 44.23431569 | 10.92732514 | 0.35 | 820 | 30 | NW | Managed |
| ALP2 | 44.13343623 | 10.88385259 | 0.40 | 930 | 10 | NW | |
| BAL1 | 44.26506962 | 11.34018064 | 0.53 | 650 | 20 | N | |
| GAGG | 44.24136697 | 10.99360049 | 0.56 | 670 | 10 | NE | |
| MALA | 44.22446988 | 10.92814309 | 0.51 | 900 | 5 | NE | |
| MART | 44.21683909 | 11.37867011 | 0.57 | 730 | 40 | S | |
| MONT | 44.23195568 | 10.92689686 | 0.28 | 800 | 30 | NW | |
| PIGH | 44.22164423 | 10.93021957 | 0.20 | 950 | 5 | W | |
| SPON | 44.13549313 | 10.89902546 | 0.35 | 890 | 10 | NW | |
| PRIA | 44.27244870 | 11.35271828 | 0.72 | 640 | 20 | N | |
| STAN | 44.22920382 | 10.92679194 | 0.46 | 840 | 10 | W | |
| TRES | 44.13208347 | 10.90276305 | 0.44 | 940 | 30 | NW | |
| ALP1 | 44.13327507 | 10.88298493 | 0.66 | 890 | 40 | NW | Abandoned |
| BAL2 | 44.26424548 | 11.33968890 | 0.40 | 670 | 30 | N | |
| BURC | 44.10592682 | 10.92572239 | 4.06 | 1030 | 20 | S | |
| POGG | 44.10400122 | 10.92358048 | 4.39 | 1015 | 25 | S | |
| POR1 | 44.13296076 | 11.09445196 | 0.55 | 860 | 0 | NW | |
| POR2 | 44.13339436 | 11.09565897 | 0.57 | 870 | 5 | NE | |
| SERR | 44.21515106 | 11.09893862 | 1.60 | 670 | 10 | NE | |
| TORR | 44.32923316 | 11.04493297 | 1.04 | 600 | 15 | NW | |

*Assessment of tree features*

For each selected tree, DBH, tree height, crown insertion and size were measured (Klamkin, 1971, 1976). A measuring tape was used for diameters, while a Vertex III Haglof hypsometer was adopted to measure height and crown insertion. The visible crown extension in each of the four cardinal directions was measured by vertically looking up. Then vitality, presence of diseases, structural condition and failure risk were evaluated. Vitality was expert-based assessed on crown condition (Turchetti et al., 2012), adopting the following scale: vigorous plant, slightly suffering, suffering, declining, dead. The three main diseases of chestnut were considered in the evaluation of healthy

status. Chestnut blight was assessed by counting the number of healing, healed, virulent, and intermediate cankers on the crown and trunk (see Turchetti et al., 2008). Each tree was assigned to one of the following two levels: predominance of hypovirulent (healing and healed cankers), or virulent infections. Ink disease was evaluated considering three classes: no symptoms, early symptoms (rarefied foliage and small and yellowing leaves), dead tree with completely dead crown and brown flames from collar. We also recorded old and recent attacks. Asian chestnut gall wasp (hereafter ACGW) was assessed by means of an inspection of the crown. Four levels of parasite presence were adopted: no galls observed, from 1 to 10 galls, from 11 to 100 galls, more than 100 galls (heavy infestation). When possible, up to 5 galls were collected and opened to assess the presence of the parasitoid *Torymus sinensis*, whose occurrence was recorded.

The state of preservation of the trees was assessed considering the presence of damages and structural defects (see Table 2) in different parts of the trees (crown, trunk and collar), following the indications reported by the International society of Arboriculture (Smiley et al., 2017). The level of damage and related failure risk was assessed by adapting the risk classes reported by Smiley et al. (2017): low risk (no damage); moderate risk (low level of damages, management needed); high risk (medium degree of damages, problems for tree stability and need of management); extreme risk (high degree of damages, tree stability strongly affected and few or no possibilities of management).

**Table 2**. Summary of structural defects considered in assigning score of damage to the single parts of the trees (crown, trunk and collar).

| Tree part | Defect type |
|---|---|
| Crown | Asymmetry, dead branches, old or recent pruning wounds, other wounds, decaying wood with rot and/or cavities, fruiting bodies |
| Trunk | Leaning, wounds, decaying wood with rot and/or cavities (% of diameter), fruiting bodies |
| Collar and root | Wounds, decaying wood with rot and/or cavities (% of diameter), fruiting bodies, excavation or soil movement in the root area, soil cracking |

*Lichen and plant survey*

196 On each selected tree, the surface covered by *L. pulmonaria* was assessed from the base up to the
197 crown insertion as a proxy for its abundance. The total extent occupied by the species (hereafter
198 abundance) was obtained as a sum of the areas of the thalli occurring on each trunk. The area of
199 each thallus was calculated as the product of the two major axes of the thallus measured by a ruler
200 positioned on a pole (Benesperi et al., 2018). The occurrence of all the lichen species belonging to
201 *Caliciales* was recorded in a north-exposed 30 × 180 cm plot placed with its shorter side at the base
202 of the trunk. In each of the five 5x5 m plots placed along the transect, the occurrence of all vascular
203 plant species was recorded. Nomenclature of lichens follows Nimis (2016), while for plants it
204 follows Pignatti et al. (2017-2019).
205
206 *Data analysis*
207 We used Canonical Discriminant Analysis (*candisc* package; Friendly and Fox, 2017) to test the
208 effect of management/abandonment on tree features: DBH, crown size, crown insertion, height,
209 vitality, structural defects and failure risk.
210 We tested the effect of management/abandonment, tree DBH (mean value at the stand level), and
211 the landscape composition (500 m and 3000 m spatial extent) on abundance of *L. pulmonaria* at
212 stand level. Multiple linear regression was performed with a stepwise AIC method for the variable
213 selection (*vegan* package; Oksanen et al., 2019). Landscape composition (i.e. amount of beech,
214 chestnut, and oak woods) was evaluated in 500 m and 3000 m radius circular buffers centred on the
215 centroid of each old-growth chestnut orchard. The Forest Map (scale 1: 10000) of the Emilia-
216 Romagna Region (available at http://ambiente.regione.emilia-romagna.it/it/parchi-
217 natura2000/foreste/le-foreste-dellemilia-romagna/le-foreste-in-emilia-romagna) was used as a
218 source of data for landscape analysis. Furthermore, the capability of *L. pulmonaria* abundance to
219 predict the species richness of *Caliciales* at the stand level was tested using the Spearman's
220 correlation coefficient.

Finally, we tested the effect of management/abandonment on plant species richness and composition. Differences in species richness between managed and abandoned stands was evaluated by Pearson's test. A Non-Metric Multidimensional Scaling (NMDS) based on Bray–Curtis dissimilarity matrix was used to display the pattern of plant species composition (*vegan* package; Oksanen et al., 2019). An Indicator Species Analysis (*indicspecies* package, ver. 1.7.8; De Cáceres 2019) was used to identify species that were overrepresented in managed and in abandoned stands (i.e. indicator species).

All the analyses were performed using R 3.6.1 for statistical computing and graphics (R Core Team, 2019).

## Results

*Tree features*

Two hundred chestnut trees were sampled. The average DBH was 1.3 m (range: 0.7-3.0 m). The most represented diameter classes ranged between 1.00 and 1.70 m (Fig. 2); classes from 1.90 to 3.00 m prevailed in abandoned stands. The average tree height was 9.3 m (range: 4.2-16.7 m), crown insertion 3.3 m (range 1.7-6.0 m) and crown size 221 $m^2$ (range: 5-570 $m^2$).

In general, vigorous trees prevailed (Fig. 3a); only 13 dead trees were found, 12 of which in abandoned stands. The good vitality conditions were also related to low levels of damage caused by pests and diseases. Chestnut blight was recorded in 98.5% of the trees (Appendix 1), prevalently with hypovirulent infections; only single and sporadic recent virulent infections were found on the crown. Ink disease was only recorded as old attacks in 8 trees (6 of them already dead) clustered in 5 stands (3 abandoned, 2 managed). ACGW was recorded on 86 trees, with less than 10 galls per tree. Furthermore, 86% of 120 examined galls contained *T. sinensis* larvae.

However, the failure risk ranged from moderate to extreme (Fig. 3b) since all the trees showed some structural problems, mainly related to the presence of wood decay and holes caused by old

246 pruning or other types of wound. Several trees had heavy branches with weak attachment to the
247 trunk.
248 On the whole (Fig. 4), a positive and significant correlation was found between management and
249 tree height ($P$ = 0.000) and crown size ($P$ = 0.014). In contrast, in abandoned stands trees showed
250 significantly higher values of DBH, defects of the crown, trunk, and collar and failure risk ($P$ =
251 0.000), while vitality was lower than in managed sites ($P$ = 0.001).
252




254 **Figure 2.** Frequency of the diameter classes of the 200 chestnut trees surveyed. Diameter classes
255 are expressed in cm. [2-column fitting image]
256
257




259 **Figure 3.** Pattern of vitality (a) and failure risk (b) of the 200 chestnut trees surveyed. [2-column
260 fitting image]

261

262 **Figure 4.** Results of Canonical Discriminant Analysis of features measured for the sampled trees
263 stratified under management conditions (M= Managed, A = Abandoned). Levels of significance: *
264 $P < 0.05$, ** $P < 0.01$. [2-column fitting image; colour online only]




*Target epiphytic lichens*

*L. pulmonaria* (Fig. 1b) occurred on 38 trees (DBH = 1.32 ± 0.38 m) clustered in 11 old-growth chestnut orchards (8 managed, see Appendix 2). In two stands the species was found together with its close relative *Lobarina scrobiculata* (Scop.) Nyl. *L. pulmonaria* abundance ranged from 50 to 3800 cm² at the tree level and from 50 to 14505 cm² at the stand level. The abundance of *L. pulmonaria* was only significantly predicted by increasing beech woods in the landscape (Table 3), both at 500 m ($P = 0.001$) and at 3000 m ($P = 0.009$) buffers. In the latter model also management was significant ($P = 0.029$), predicting higher abundance in managed stands.

11 *Caliciales* belonging to the genera *Acolium*, *Calicium*, *Chaenotheca*, and *Chaenothecopsis* were found in 13 stands (Appendix 2). The most common were *Calicium adspersum* (17 records in 10 stands), and *C. salicinum* (12 records in 6 stands). A positive relationship was found between the

number of species belonging to *Caliciales* and the abundance of *L. pulmonaria* ($\rho$ = 0.619, $P$ = 0.0056).

**Table 3.** Regression models of factors affecting abundance of *L. pulmonaria* at stand level. Model 1 considers habitat amounts at 500 m ring buffer, while Model 2 at 3000 m. M = managed; A = abandoned.

| Model | | Estimate | Std. Error | t value | Pr(>\|t\|) |
|---|---|---|---|---|---|
| 1 | (Intercept) | -1.046 | 11.402 | -0.092 | 0.928 |
| | M/A | 24.023 | 13.450 | 1.786 | 0.092 |
| | Beech woods (500 m) | 11.859 | 2.812 | 4.217 | 0.001 |
| 2 | (Intercept) | -97.138 | 54.789 | -1.773 | 0.095 |
| | M/A | 32.081 | 13.386 | 2.397 | 0.029 |
| | Beech woods (3000 m) | 3.422 | 1.162 | 2.946 | 0.009 |

*Pattern of plant diversity*

Overall, 313 plant species were found. Species richness at the stand level ranged from 32 to 106 species. Some species worthy of conservation were found among orchids: *Cephalanthera longifolia*, *C. rubra*, *Epipactis helleborine*, *Listera ovata*, *Neotinea maculata*, *Platanthera bifolia*, and *P. chlorantha*. Additional species of conservation concern, which are protected at regional level (Regional Law 24-1-1977, n. 2), were *Dianthus armeria*, *D. balbisii* and *Gentiana asclepiadea*. 40 occurrences of these species were recorded in managed stands and 20 in abandoned stands.

Managed stands had a significantly higher species richness than abandoned stands ($P$ = 0.0014). In managed stands, species richness ranged from 72 to 106 species (average: 87.42 ± 3.42), whereas in abandoned stands it ranged from 32 to 88 species (average: 61.25 ± 6.80). The effect of management on plant communities was also confirmed by the pattern of species composition, as indicated by the NMDS plot (stress 0.096, Fig. 5). The Indicator Species Analysis revealed 22 species significantly overrepresented in managed stands and 7 in abandoned stands (Table 4). In managed stands, grassland species prevailed, whereas in abandoned stands wood and pre-forestry species prevailed.




300 **Figure 5.** NMDS plot of the plant species composition pattern in managed and abandoned stands.
301 Red dots: managed stands; green dots: abandoned stands. Stress = 0.096. [2-column fitting image;
302 colour online only]

303

304 **Table 4**. List of indicator species for managed and abandoned stands. The *p*-value is also given.

| Stand type | List of species |
|---|---|
| Managed | *Anthoxanthum odoratum* and *Lotus corniculatus* ($P < 0.01$), *Lychnis flos-cuculi*, *Holcus lanatus*, *Genista tinctoria*, *Cynosurus cristatus*, *Campanula rapunculus*, *Plantago lanceolata*, *Achillea collina*, *Trisetum flavescens*, *Danthonia decumbens*, *Polygala vulgaris*, *Aira elegantissima, Leontodon hispidus*, *Schedonorus pratensis*, *Taraxacum* gr. *officinale*, *Calluna vulgaris*, *Leucanthemum* gr. *vulgare*, *Trifolium campestre*, *Helianthemum nummularium*, *Rhinanthus alectorolophus* ($P < 0.05$) |
| Abandoned | *Fraxinus ornus*, *Rosa canina*, *Lactuca muralis*, *Cytisus scoparius*, *Melica uniflora*, *Abies alba*, *Teucrium scorodonia* ($P < 0.05$) |

305

306 **Discussion**

307 Our results revealed contrasting patterns of tree features, lichen, and plant diversity in managed and
308 abandoned old-growth chestnut orchards of the northern Apennines, supporting the view that

309 traditional management is fundamental for the long-term maintenance of healthy veteran trees, the
310 enhancement of epiphytic lichens related to old-growth conditions, and plant diversity.
311 Our old-growth chestnut orchards survived the three main diseases that are currently quite
312 compatible with tree health. For example, blight damages are low, usually involving only single
313 desiccated branches, and hypovirulence predominated as in other Italian chestnut areas (Turchetti et
314 al., 2008). Also ink disease rarely occurs in the investigated stands, even if the disease remains as a
315 potential risk. ACGW is present only with few galls always parasitized by *Torymus sinensis*, thus
316 confirming the effectiveness of the biological control (Vai et al., 2014). However, in abandoned
317 orchards these reduced damages likely accumulate over the years, producing the spectral aspect of
318 these stands (Turchetti and Maresi, 2008). Moreover, in abandoned stands crown size, and height
319 are lower possibly due to competition with invading trees, while crown, trunk and collar defects due
320 to wood decay are higher. Light reduction, competition and decay are worsened by the age of the
321 plants, producing the failure or the death of the old trees, whose main trunk is often replaced by
322 clones of the same tree (when some roots survive) as if it were coppice. Consistently, our results
323 indicate that in abandoned stands failure risk is significantly higher than in managed stands and
324 overall tree vitality is lower.
325 In managed orchards the occurrence of veteran trees arranged in an open forest structure and the
326 absence of shoots that cover the trunks may explain the higher abundance of the forest lichen *L.*
327 *pulmonaria* (Nascimbene et al., 2013) as compared to abandoned stands, where increasing canopy
328 closure and the decay of the more ancient trees coupled with re-sprouting of new shoots around
329 remnant trunks may reduce habitat suitability for this large lichen. In these anthropogenic habitats,
330 the maintenance of the most valuable species is thus related to low intensity, traditional
331 management (Nascimbene et al., 2014). The exclusive occurrence of the declining and red-listed
332 lichen *Lobarina scrobiculata* (Nascimbene et al., 2013c, 2016) and the higher occurrence of
333 *Caliciales* (whose species richness is predicted by the abundance of *L. pulmonaria*) in managed
334 orchards clearly corroborate the importance of these anthropogenic habitats for epiphyte

conservation. However, our results also stress the role of landscape composition around the old-growth chestnut orchards, indicating that the abundance of *L. pulmonaria* is maximized in stands embedded in a beech forest-dominated landscape. The huge historical herbarium records, available since the first half of 19$^{th}$ century, suggest a long lasting persistence and a higher commonness of both *L. pulmonaria* and *Lobarina scrobiculata* in our study area (Bertoloni, 1867; Saccardo and Fiori, 1894; Zanfrognini, 1902; see also Appendix 3), where they established luxuriant populations in both beech and chestnut stands (Bertoloni, 1867). In this historical scenario, old-growth beech forests may be considered the primary habitat for these lichens, while anthropogenic chestnut stands likely provided a suitable secondary habitat that allowed their survival when most beech stands were converted to coppice. Frequent disturbance in coppice stands coupled with the absence of old trees and the muffling of the trunks by the excess of shoots may have caused the rarefaction of these lichens in beech stands, while the more stable ecological conditions and the occurrence of veteran trees in old-growth chestnut orchards provided a valuable refuge. In this perspective, the effect of the landscape found in our study may reflect a historical condition in which the old-growth beech forest originally supplied propagules for the establishment of the species in old-growth chestnut orchards that currently are likely the main propagule hot-spots in the mountain forest landscape of the Northern Apennines (Matteucci et al., 2012).

The role of managed old-growth chestnut orchards for biodiversity is also confirmed for plant communities (e.g. Gondard et al., 2001, 2007; Nascimbene et al., 2014), with significantly higher species richness than in abandoned stands. Moreover, our results indicate that the abandonment of management practices (i.e. mowing and/or grazing) triggers compositional shifts that may cause the loss of several species of conservation concern, as in the case of many orchids. In general, plant communities of the managed stands host species related to open habitats (*Molino-Arrhenatheretea* Tüxen 1937 or *Festuco valesiacae-Brometea erecti* Br.-Bl. & Tüxen ex Br.-Bl. 1949), while in abandoned stands pre-forestry and woody species tend to prevail (*Laburno-Ostryion* Ubaldi 1980 and/or *Fagetalia sylvaticae* Pawłowski in Pawłowski, Sokołowski & Wallisch 1928; Ubaldi et al.,

1993; Pezzi et al., 2011). Historical records (Bertoloni, 1867) corroborate the view that the floristic composition of managed stands remained almost unchanged for decades, further underlining the role of traditional management for maintaining stable ecological conditions (Tomaselli, 1989).

## Conclusions

Our study supports the view that old-growth chestnut orchards are a perturbation-dependent system (Vogl, 1980; Gondard et al., 2001) whose effectiveness for biodiversity conservation depends on the maintenance of low-intensity management. In this perspective, mowing and pruning are still the most effective practices for preserving the old-growth chestnut orchard habitat and its associated biodiversity. However, pruning needs to be light and not frequent, aiming to restore a suitable architectural structure, cutting only dangerous branches and saving biodiversity niches (Zapponi et al., 2017). Topping and heavier pruning used in the past are likely the main causes of the current precarious status of the surviving trees. Moreover, also sporadic fire as a sanitation method or to burn mowed and pruned material should be avoided, since it may negatively impact on long-term soil conservation. Our results indicate that 1000 years of *chestnut civilization* represent a cultural heritage that benefits nature conservation, promoting a virtuous interplay between human activities and biodiversity. For this reason, policies aimed at sustaining traditional and even more sustainable management in old-growth chestnut orchards are indispensable to avoid the degradation and loss of this habitat and its centuries-old cultural and ecological legacy.

## Acknowledgements

We thank Dr. Renzo Panzacchi (President of the Consorzio Castanicoltori Appennino Bolognese), Mr. Erminio Bernardi (chestnut grower, President of the Associazione Naturalmente, Montese - MO), Dr. Luigi Vezzalini (Technical coordinator of the Associazione nazionale Città del Castagno), Dr. Maurizio Musolesi (chestnut grower), Prof. Giuseppe Nanni (president of the Consorzio Castanicoltori di Granaglione), Umberto Biagi (chestnut grower) for helping us in contacting

chestnut growers and finding chestnut stands. We thank all the chestnut growers that allowed us to perform the relevés in their properties and gave us useful information and support for our work. In addition, we thank Dr. David Bianco (Ente di Gestione per i Parchi e la Biodiversità dell'Emilia orientale), for allowing us to perform the researches in the Natura 2000 sites. Prof. Davide Ubaldi (formerly Università di Bologna) and Dr. Alessandro Alessandrini helped us in the identification of some plant species and Domenico Puntillo (Università della Calabria) in the identification of *Caliciales* species. Michael Webb (Modern English Study Centre, Bologna) kindly revised the original English text.

**Declaration of interest**: none.

## References

Arnaud, M.T., Chassany, J.P., Dejean, R., Ribart, J., Queno, L., 1997. Economic and ecological consequences of the disappearance of traditional practices related to chestnut groves. J. Environ. Manage. 49, 373–391. https://doi.org/10.1006/jema.1995.0120

Barbati, A., Marchetti, M., 2005. Forest Types for Biodiversity Assessment (FTBAs) in Europe: the revised classification scheme. In: Marchetti, M. (ed.), Monitoring and indicators of forest biodiversity in Europe - from ideas to operationality. European Forest Institute Proceedings, 51, pp. 105-126.

Benesperi, R., Nascimbene, J., Lazzaro, L., Bianchi, E., Tepsich, A., Longinotti, S., Giordani, P. 2018. Successful conservation of the endangered forest lichen *Lobaria pulmonaria* requires knowledge of fine-scale population structure. Fungal Ecol. 33, 65-71. https://doi.org/10.1016/j.funeco.2018.01.006

Bertoloni, G., 1867. Vegetazione dei monti di Porretta e dei suoi prodotti vegetali. In Bertoloni, G., illustrazione delle terme di Porretta e del suo territorio pubblicata per deliberazione del consiglio provinciale di Bologna. Regia Tipografia, Bologna, pp. 193-332.

Bounous, G., 2014. Il castagno. Risorsa multifunzionale in Italia e nel mondo. Edagricole, Bologna.

Brunialti, G., Frati, L., Ravera, S., 2015. Structural variables drive the distribution of the sensitive lichen *Lobaria pulmonaria* in Mediterranean old-growth forests. Ecol. Ind. 53, 37-42. https://doi.org/10.1016/j.ecolind.2015.01.023

Conedera, M, Krebs, P., 2008. History, present situation and perspective of chestnut cultivation in Europe. In: Abreu, C.G., Peixoto, F.P., Gomes-Laranjo, J. (eds.), Proceedings of the second Iberian Chestnut Congress, Vila Real (P), June 20–22th 2007. Acta Hort. 784, 23–27. http://dx.doi.org/10.17660/ActaHortic.2008.784.1

Conedera, M., Krebs, P., Tinner, W., Pradella, M., Torriani, D., 2004. The cultivation of *Castanea sativa* (Mill.) in Europe. from its origin to its diffusion on a continental scale. Veget. Hist. Archaeobot. 13(3), 161-179. http://dx.doi.org/10.1007/s00334-004-0038-7

Conedera, M., Stanga, P., Oester, B., Bachmann, P., 2001. Different post-culture dynamics in abandoned chestnut orchards and coppices. For. Snow Landsc. Res. 76(3), 487-492.

Connel, J.H., 1978. Diversity in tropical rainforest and coral reefs. Science 199, 1302-1310.

De Cáceres, M, 2019. How to use the indicspecies package (ver. 1.7.8). Retrieved from https://cran.r-project.org/web/packages/indicspecies/

Fenaroli, L., 1946. Il castagno. Trattati di Agricoltura, vol. 1, 222 pp. Roma, Ramo Editoriale degli Agricoltori.

Friendly, M., Fox, J., 2017. candisc: visualizing generalized canonical discriminant and canonical correlation analysis. R package version 0.8–0. Retrieved from https://CRAN.R-project.org/package=candisc

Gabrielli, A., 1994. La civiltà del castagno. Monti e boschi 65, 3.

Gondard, H., Romane, F., Grandjanny, M., Li, J., Aronson, J., 2001. Plant species diversity changes in abandoned chestnut (*Castanea sativa*) groves in southern France. Biodivers. Conserv. 10, 189-207. https://doi.org/10.1023/A:1008997625523

438 Gondard, H., Santa Regina, I., Salazar S., Peix A., Romane, F., 2007. Effect of forest management
439 on plant species diversity in *Castanea sativa* stands in Salamanca (Spain) and the Cévennes
440 (France). Scientific Research and Essay 2 (2), 062-070. Available online at
441 http://www.academicjournals.org/SREHislop, M., Twery, M., Vihemäki, H., 2004. Involving people
442 in forestry: a toolbox for public involvement in forest and woodland planning. Forestry
443 Commission, Edinburgh.
444 Klamkin, M.S., 1971. Elementary approximations to the area of n-dimensional ellipsoids. Am.
445 Math. Monthly 78 (3), 280-283. https://dx.doi.org/10.2307/2317530
446 Klamkin, M.S., 1976. Corrections to "elementary approximations to the area of N-dimensional
447 ellipsoids" (This Monthly, 78 (1971) 280–283). Am. Math. Monthly 83 (6), 478-478.
448 https://doi.org/10.1080/00029890.1976.11994150
449 Krebs, P., Conedera, M., Fonti, P., 2005. The inventory of the giant chestnut trees in Southern
450 Switzerland. Acta Hort. 693, 171. https://doi.org/10.17660/ActaHortic.2005.693.20
451 Krebs, P., Moretti, M., Conedera, M., 2008. Castagni monumentali nella Svizzera sudalpina.
452 Importanza geostorica, valore ecologico e condizioni sanitarie. Sherwood 14 (1), 5-10.
453 Löbel, S., Snäll, T., Rydin, H., 2009. Mating system, reproduction mode and diaspore size affect
454 metacommunity diversity. J. Ecol. 97, 176–185. https://doi.org/10.1111/j.1365-2745.2008.01459.x
455 Macagno, A.L.M., Gobbi, M., Franceschini, A., Lencioni, V., 2012. New record of *Osmoderma*
456 *eremita* (Scopoli, 1763) (Coleoptera: Cetoniidae) in chestnut trees in Trentino (Eastern Italian Alps).
457 Studi Trent. Sci. Nat. 92, 37–41.
458 Matteucci, E., Benesperi, R., Giordani, P., Piervittori, R., Isocrono, D., 2012. Epiphytic lichen
459 communities in chestnut stands in Central-North Italy. Biologia 67 (1), 61-70.
460 https://doi.org/10.2478/s11756-011-0145-8
461 Mondino, G.P., 1991. Caratteristiche dei boschi di sostituzione e loro tendenze evolutive. In:
462 AA.VV., I boschi italiani. Valori naturalistici e problemi di gestione. Atti del Simposio. Castiglione
463 de' Pepoli (Bologna), 9 ottobre 1989. Soc. Emil. Pro Montibus et Silvis, Bologna, pp. 53-61.

464 Nascimbene, J., Brunialti, G., Ravera, S., Frati, L., Caniglia, G., 2010. Testing *Lobaria pulmonaria*
465 (L.) Hoffm. as an indicator of lichen conservation importance of Italian forests. Ecol. Ind. 10 (2),
466 353-360. https://doi.org/10.1016/j.ecolind.2009.06.013
467 Nascimbene, J., Dainese, M., Sitzia, T., 2013a. Contrasting responses of epiphytic and dead wood-
468 dwelling lichen diversity to forest management abandonment in silver fir mature woodlands. For.
469 Ecol. Manage. 289, 325-332 https://doi.org/10.1016/j.foreco.2012.10.052
470 Nascimbene, J., Benesperi, R., Brunialti, G., Catalano, I., Vedove, M.D., Grillo, M., Isocrono, D.,
471 Matteucci, E., Potenza, G., Puntillo, D., Puntillo, M., Ravera, S., Rizzi, G., Giordani, P., 2013b.
472 Patterns and drivers of β-diversity and similarity of *Lobaria pulmonaria* communities in Italian
473 forests. J. Ecol. 101, 493-505. https://doi.org/10.1111/1365-2745.12050
474 Nascimbene, J., Nimis, P.L., Ravera, S., 2013c. Evaluating the conservation status of epiphytic
475 lichens of Italy: A red list. Plant Biosyst. 147 (4), 898-904.
476 https://doi.org/10.1080/11263504.2012.748101
477 Nascimbene, J., Fontana, V., Spitale D., 2014. A multi-taxon approach reveals the effect of
478 management intensity on biodiversity in Alpine larch grasslands. Sci Tot Env 487, 110–116
479 https://doi.org/10.1016/j.scitotenv.2014.04.013
480 Nascimbene, J., Casazza, G., Benesperi, R., Catalano, I., Cataldo, D., Grillo, M., Isocrono, D.,
481 Matteucci, E., Ongaro, S., Potenza, G., Puntillo, D., Ravera, S., Zedda, L., Giordani, P., 2016.
482 Climate change fosters the decline of epiphytic *Lobaria* species in Italy. Biol. Conserv. 201, 377-
483 384. https://doi.org/10.1016/j.biocon.2016.08.003
484 Nimis, P.L., 2016. The Lichens of Italy. A Second Annotated Catalogue. EUT, Trieste.
485 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'
486 Hara, R., Simpson, G., Solymos, P., Henry, M., Stevens, M., Szoecs, E., Wagner, H., 2019. vegan:
487 Community Ecology Package. R package version 2.5-6. https://CRAN.R-
488 project.org/package=vegan

489 Paci, M., Maltoni, A., Tani, A., 2000. I castagneti abbandonati della Toscana: dinamismo e proposte
490 gestionali. In: Bucci, G., Minotta, G., Borghetti, M. (eds), Applicazioni e prospettive per la ricerca
491 forestale in Italia, Atti II congresso SISEF (Bologna, 20–23 October 1999), pp. 9–16.
492 Pezzi, G., Lucchi, E., Maresi, G., Ferretti, F., Viaggi, D., Frascaroli, F., 2017. Abandonment or
493 survival? Understanding the future of *Castanea sativa* stands in function of local attitude (Northern
494 Apennine, Italy). Land Use Policy. 61, 564-574. https://doi.org/10.1016/j.landusepol.2016.10.049
495 Pezzi, G., Donati, D., Muzzi, E., Conedera, M., Krebs, P., 2019: Using chorographic sources to
496 reconstruct past agro-forestry systems. A methodological approach based on the study case of the
497 northern Apennines. Landsc. Res. https://doi.org/10.1080/01426397.2019.1624700
498 Pezzi, G., Maresi, G., Conedera, M., Ferrari, C., 2011. Woody species composition of chestnut
499 stands in the Northern Apennines: the result of 200 years of changes in land use. Landsc. Ecol. 26
500 (10), 1463–1476. https://doi.org/10.1007/s10980-011-9661-8
501 Pignatti, S., Guarino, R., La Rosa, M., 2017-2019. Flora d'Italia, II ed. Edagricole, Bologna.
502 Pitte, J.R., 1986. Homme et paysage du châtaignier de l'Antiquité à nos jours. Librairie A. Fayard,
503 Paris.
504 Piussi, P., Pettenella, D., 2000. Spontaneous afforestation of fallows in Italy. In: Weber, N. (ed.),
505 NEWFOR new forest for Europe: afforestation at the turn of the century. Proceeding of the
506 scientific symposium, Freiburg, 16–17 February 2000, pp. 151–163.
507 Quattrocchi, G., 1938. Il miglioramento dei castagneti dell'Appennino Bolognese. Stabilimento
508 Grafico F. Lega, Faenza.
509 R Core Team, 2019. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
510 Foundation for Statistical Computing. Accessed 1 october 2019. https://www.R-project.org/.
511 Romane, F., Hauter, S., Valerino, L., 1995. Factors affecting biodiversity in chestnut (*Castanea*
512 *sativa* Mill.) ecosystems along a gradient from coppice to orchard in the Cevennes Mountains
513 (Southern France). In: Romane, F., (ed.), Sustainability of mediterranean ecosystems. Case study of
514 the chestnut forest. Ecosystem Research Report 19, EUR 15727 EN, pp. 103–109

515 Saccardo, F., Fiori, A., 1894. Contribuzione alla lichenologia del Modenese e del Reggiano. Atti
516 Soc. Nat. Mat. Modena 28, 198-225.
517 Seijo, F., Cespedes B., Zaval, G., 2018. Traditional fire use impact in the aboveground carbon stock
518 of the chestnut forests of Central Spain and its implications for prescribed burning. Sci. Total
519 Environ., 625, 1405-1414. https://doi.org/10.1016/j.scitotenv.2017.12.079
520 Seijo, F., Millington, J. D. A., Gray, R., Mateo, L. H., Sangüesa-Barreda, G., Camarero, J.J., 2017.
521 Divergent Fire Regimes in Two Contrasting Mediterranean Chestnut Forest Landscapes. Human
522 Ecol. (New York), 45(2), 205-219. https://doi.org/10.1007/s10745-016-9879-9
523 Selva, S.B., 1994. Lichen diversity and stand continuity in the northern hardwoods and spruce-fir
524 forests of northern New England and western New Brunswick. Bryologist 97, 424-429.
525 https://doi.org/10.2307/3243911
526 Smiley, E.T., Matheny, N., Lilly, S., 2017. Best management practices: tree risk assessment. Second
527 edition. ISA, International Society of Arboriculture.
528 Squatriti, P., 2013. Landscape and change in early Medieval Italy: chestnuts, economy, and culture.
529 Cambridge University Press, Cambridge and New York. https://doi.org/10.1017/S00387134150015
530 Temel, F., Ozalp, M., 2009. Monumental *Castanea sativa* Mill. individuals on the slopes of Genya
531 Mountain, Artvin, Turkey. Acta Hort. 815, 171-178.
532 http://dx.doi.org/10.17660/ActaHortic.2009.815.22
533 Tomaselli, M., 1989. Osservazioni sul dinamismo di alcuni boschi decidui in Alta Lunigiana
534 (Toscana NW). Monti e Boschi 2, 53.
535 Tongco, M., 2007. Purposive sampling as a tool for informant selection. Ethnobot. Res. Appl. 5,
536 147-158.
537 Turchetti, T., Maresi, G., 2008. Biological control and management of chestnut diseases. Integrated
538 management of diseases caused by fungi, phytoplasma and bacteria, In: Ciancio, A., Mukerji, K.G.
539 (Eds.), Integrated Management of Diseases Caused by Fungi, Phytoplasma and Bacteria. Springer,
540 the Netherlands, pp. 85-118. http://dx.doi.org/10.1007/978-1-4020-8571-0

Turchetti, T., Ferretti, F., Maresi, G., 2008. Natural spread of *Cryphonectria parasitica* and persistence of hypovirulence in three Italian coppiced chestnut stands. For. Pathol. 38 (4), 227–243. https://doi.org/10.1111/j.1439-0329.2008.00557.x

Turchetti, T., Pennacchio, F., D'Acqui, L.P., Maresi, G., Pedrazzoli, F., 2012. Interventi per la gestione dei castagneti invasi dal cinipide. Forest@ 9, 227–235. http://dx.doi.org/10.3832/efor0701-009

Ubaldi, D., Zanotti, A.L., Puppi, G., Maurizzi, S., 1993. I boschi di *Laburno-Ostryion* in Emilia-Romagna. Ann. Bot. (Roma) 51 (10), 157-170.

Vai, N., Colla, R., Mazzoli, L., Bariselli, M., 2014. The regional project for biological control of the Chinese gall wasp *Dryocosmus kuriphilus* in Emilia-Romagna. In: Giordano, L., Ferrini, F., Gonthier, P. (eds.), European Conference of Arboriculture − Planning the Green City: Relationships between trees and infrastructures: Conference and Abstracts Book, pp. 87-88.

Vigiani, D., 1908. Il castagno. Tipografia e Litografia Carlo Cassone, Casale.

Vogl, R.J., 1980. The ecological factors that produce perturbation-dependent ecosystems. In: Cairns, J. (Ed.), The recovery process in damaged ecosystems. Ann. Arbor. Science Publishers Inc., Ann. Arbor, Michigan, pp. 63-94.

Zanfrognini, C., 1902. Flora lichenologica dell'Emilia. Nuovo Giorn. Bot. Ital. 9, 190-211.

Zapponi, L., Mazza, G., Farina, A., Fedrigoli, L., Mazzocchi, F., Roversi, P.F., Sabbatini Peverieri, G., Mason, F., 2017. The role of monumental trees for the preservation of saproxylic biodiversity: re-thinking their management in cultural landscapes. In: Campanaro, A., Hardersen, S., Sabbatini Peverieri, G., Carpaneto, G.M. (Eds.), Monitoring of saproxylic beetles and other insects protected in the European Union. Nature Conservation 19, 231-243. https://doi.org/10.3897/natureconservation.19.12464

**CR CRediT author statement**

Giovanna Pezzi, Giorgio Maresi, Fabrizio Ferretti, Simone Gambini, Fabrizio Buldrini, Juri Nascimbene: *Conceptualization, Methodology*

Giovanna Pezzi, Giorgio Maresi, Fabrizio Ferretti, Simone Gambini, Fabrizio Buldrini, Juri Nascimbene: *Data curation*

Enrico Muzzi: *Statistical Analysis*

Giovanna Pezzi, Giorgio Maresi, Juri Nascimbene: *Supervision*

Giovanna Pezzi, Giorgio Maresi, Fabrizio Ferretti, Fabrizio Buldrini, Enrico Muzzi, Juri Nascimbene: *Writing-Original draft preparation*

Giovanna Pezzi, Giorgio Maresi, Fabrizio Ferretti, Fabrizio Buldrini, Enrico Muzzi, Juri Nascimbene: *Writing-Reviewing, and Editing*