

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

The cohomology rings of the unordered configuration spaces of elliptic curves

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

The cohomology rings of the unordered configuration spaces of elliptic curves / Pagaria, Roberto. - In: ALGEBRAIC AND GEOMETRIC TOPOLOGY. - ISSN 1472-2739. - STAMPA. - 20:6(2020), pp. 2995-3012. [10.2140/agt.2020.20.2995]

*Availability:*

This version is available at: <https://hdl.handle.net/11585/784664> since: 2020-12-17

*Published:*

DOI: <http://doi.org/10.2140/agt.2020.20.2995>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**Pagaria, R. (2020). The cohomology rings of the unordered configuration spaces of elliptic curves. Algebraic and Geometric Topology, 20(6), 2995-3012**

The final published version is available online at  
<https://dx.doi.org/10.2140/agt.2020.20.2995>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

## **The cohomology rings of the unordered configuration spaces of elliptic curves**

ROBERTO PAGARIA

We study the cohomology ring of the configuration space of unordered points in the two dimensional torus. In particular, we compute the mixed Hodge structure on the cohomology, the action of the mapping class group, the structure of the cohomology ring and we prove the formality over the rationals.

55R80

### **Introduction**

We fully describe the cohomology with rational coefficients of the configuration spaces of unordered points in an elliptic curve (frequently called torus).

Configuration spaces of points are related to physics (state spaces of non-colliding particles on a manifold), robotics (motion planning), knot theory, and topology. Configuration spaces give invariants of the homeomorphism type of the base space. In the algebraic setting, configuration spaces are open in the moduli spaces of points.

Since the literature is very extensive, we compare our work only with the main results on the (co-)homology of configuration spaces. The first computation of the cohomology algebra of configuration spaces is due to Arnol'd [1, 2] in the case of  $\mathbb{R}^2$ . This result has been generalized by Cohen, Lada, and May [9] to the configuration space of  $\mathbb{R}^n$  and later by Goresky and MacPherson [17]. Partially additive results have been obtained: by Bödighheimer and Cohen [6] for once-punctured oriented surfaces, by the same authors and Taylor [7] for odd dimensional manifolds, and by Drummond-Cole and Knudsen [11] for surfaces in general. However there is no description of the ring structure; we provide it in the case of elliptic curves. The Betti numbers  $C^n(X)$  are described in the following cases: for  $X = \mathbb{P}^2(\mathbb{R})$  by Wang [30], for  $X$  a sphere by Salvatore [26], for  $X = \mathbb{P}^2(\mathbb{C})$  by Felix and Tanré [14] and for elliptic curves by Maguire and Schiessl [23, 27].

In this paper we improve the previous results on configuration spaces in an elliptic curve in three ways. We describe:

- the mixed Hodge structure on the cohomology ([Theorem 3.3](#)),

- the action of the mapping class group ([Theorem 3.3](#)),
- the ring structure ([Theorem 4.1](#)).

The formality result over the rationals is proven in [Theorem 4.3](#).

We prove these results using the Križ model [[22](#), [28](#), [5](#), [12](#)] and the representation theory on it [[3](#), [4](#)].

In [Section 1](#) we recall the Križ model, then in [Section 2](#) we improve the result on the decomposition of the Križ model into irreducible representations, see [Theorem 2.9](#). Descriptions of the mixed Hodge structure and of the action of the mapping class group are obtained in [Section 3](#) by computing the cohomology of the model. Finally, the ring structure is presented in the last section.

## 1 The Križ model

Let  $E$  be an elliptic curve and consider the configuration space of  $n$  ordered distinct points

$$\mathcal{F}^n(E) \stackrel{\text{def}}{=} \{\underline{p} \in E^n \mid p_i \neq p_j\}.$$

The symmetric group  $\mathfrak{S}_n$  acts on  $\mathcal{F}^n(E)$  by permuting the coordinates and the quotient is the configuration space of  $n$  unordered points

$$\mathcal{C}^n(E) \stackrel{\text{def}}{=} \mathcal{F}^n(E)/\mathfrak{S}_n.$$

We also consider the space  $\mathcal{M}^n(E)$ , defined by

$$\mathcal{M}^n(E) \stackrel{\text{def}}{=} \{\underline{p} \in \mathcal{F}^n(E) \mid \sum p_i = 0\}.$$

Notice that there exists a non canonical isomorphism  $\mathcal{F}^n(E) \cong E \times \mathcal{M}^n(E)$ .

In this section we recall a rational model for the cohomology algebra of  $\mathcal{F}^n(E)$ . The model is a commutative differential bi-graded algebra (dga) that can be obtained in two different ways: as a specialization of the Križ model for the configuration spaces or as the second page of the Leray spectral sequence (also known as the Totaro spectral sequence) for elliptic arrangements. Our main references for the first approach are [[22](#), [3](#), [4](#)] and for the second one are [[28](#), [12](#), [5](#)]. In the following we define the models for the cohomology of  $\mathcal{F}^n(E)$  and of  $\mathcal{M}^n(E)$ .

Let  $\Lambda$  be the exterior algebra over  $\mathbb{Q}$  with generators

$$\{x_i, y_i, \omega_{i,j}\}_{1 \leq i < j \leq n}.$$

We set the degree of each  $x_i$  and  $y_i$  equal to  $(1, 0)$  and the degree of  $\omega_{i,j}$  equal to  $(0, 1)$ . Define the differential  $d: \Lambda \rightarrow \Lambda$  of bi-degree  $(2, -1)$  on generators as follows:  $dx_i = 0$  and  $dy_i = 0$  for  $i = 1, \dots, n$  and

$$d\omega_{i,j} \stackrel{\text{def}}{=} (x_i - x_j)(y_i - y_j).$$

For the sake of notation we set  $\omega_{i,j} := \omega_{j,i}$  for  $i > j$ .

We define the dga  $A^{\bullet,\bullet}$  as the quotient of  $\Lambda$  by the following relations:

$$(x_i - x_j)\omega_{i,j} = 0 \quad \text{and} \quad (y_i - y_j)\omega_{i,j} = 0, \\ \omega_{i,j}\omega_{j,k} - \omega_{i,j}\omega_{k,i} + \omega_{j,k}\omega_{k,i} = 0.$$

Notice that the ideal is preserved by the differential map, thus the differential  $d: A^{\bullet,\bullet} \rightarrow A^{\bullet,\bullet}$  is well defined.

**Remark 1.1** The model  $A^{\bullet,\bullet}$  coincides with the Križ model  $E^\bullet$  introduced in [22] up to shifting the degrees, ie

$$A^{p,q} \cong E_q^{p+q}.$$

The dga  $E^\bullet$  is a rational model for  $X$ , as shown in [22, Theorem 1.1].

In order to study the cohomology of  $A^{\bullet,\bullet}$  we need to introduce the elements  $u_{i,j} = x_i - x_j$ ,  $v_{i,j} = y_i - y_j$  and  $\gamma = \sum_{i=1}^n x_i$ ,  $\bar{\gamma} = \sum_{i=1}^n y_i \in A^{1,0}$ .

We define the dga  $B^{\bullet,\bullet}$  as the subalgebra of  $A^{\bullet,\bullet}$  generated by  $u_{i,j}, v_{i,j}$  and  $\omega_{i,j}$  for  $1 \leq i < j \leq n$ . Let  $D^{\bullet,0}$  be the subalgebra of  $A^{\bullet,\bullet}$  generated by  $\gamma$  and  $\bar{\gamma}$  endowed with the zero differential map. Notice that

$$(1) \quad A^{\bullet,\bullet} \cong B^{\bullet,\bullet} \otimes_{\mathbb{Q}} D^{\bullet,0}$$

as differential algebras and that  $D^{\bullet,0}$  is the cohomology ring of the elliptic curve  $E$ .

The mixed Hodge structure on the cohomology of algebraic varieties defines a bigrading compatible with the algebra structure (see [10, p.81] or [29, Theorem 8.35]). In our case the bigrading given by the mixed Hodge structure coincides with the one given by the Leray spectral sequence as shown by Totaro [28, Theorem 3] and by Gorinov [18]. Explicitly, the subspace  $A^{p,q}$  has weight  $p + 2q$  and degree  $p + q$ .

The following result is a particular case of [5, Theorem 3.3] and of [12, Theorem 1.2].

**Theorem 1.2** *The cohomology algebra of  $\mathcal{F}^n(E)$  (or of  $\mathcal{M}^n(E)$ ) with rational coefficients is isomorphic to the cohomology of the dga  $A^{\bullet,\bullet}$  (respectively of  $B^{\bullet,\bullet}$ ). Moreover, the  $n^2$ -sheeted covering*

$$E \times \mathcal{M}^n(E) \rightarrow \mathcal{F}^n(E) \\ (q, p) \mapsto (p_i + q)_{i=1, \dots, n}$$

*induces the isomorphism of eq. (1).*

## 2 Representation theory on the Križ model

Now we study the action of the symmetric group  $\mathfrak{S}_n$  and of  $SL_2(\mathbb{Q})$  on the algebras  $A^{\bullet,\bullet}$  and  $B^{\bullet,\bullet}$ . Those actions are given by a geometric action on  $\mathcal{F}^n(E)$ . For general reference about the representation theory of the Lie group and of the Lie algebra we refer to [19] and to [15], respectively. The cases of  $SL_2(\mathbb{C})$  and of  $\mathfrak{sl}_2(\mathbb{C})$  can be found in [16].

### 2.1 Definition of the actions

Consider the action of  $\mathfrak{S}_n$  on  $\mathcal{F}^n(E)$  defined by

$$\sigma^{-1} \cdot (p_1, \dots, p_n) = (p_{\sigma(1)}, \dots, p_{\sigma(n)})$$

for all  $\sigma \in \mathfrak{S}_n$ . This induces an action on  $A^{\bullet,\bullet}$  and on  $B^{\bullet,\bullet}$  defined by

$$\begin{aligned}\sigma^{-1}(x_i) &= x_{\sigma(i)}, \\ \sigma^{-1}(y_i) &= y_{\sigma(i)}, \\ \sigma^{-1}(\omega_{i,j}) &= \omega_{\sigma(i),\sigma(j)}\end{aligned}$$

for all  $1 \leq i < j \leq n$  and all  $\sigma \in \mathfrak{S}_n$ .

The mapping class group  $\text{MCG}(E)$  acts naturally on  $\mathcal{F}^n(E)$  and on  $\mathcal{C}^n(E)$ .

**Theorem 2.1** (Theorem 2.5 [13]) *The mapping class group  $\text{MCG}(E)$  of the torus is isomorphic to  $SL_2(\mathbb{Z})$  and the isomorphism is given by the natural action of  $\text{MCG}(E)$  on  $H^1(E; \mathbb{Z})$ .*

Let  $f$  be an automorphism of  $E$ , the map induces the following vertical morphisms

$$\begin{array}{ccc} \mathcal{F}^n(E) & \hookrightarrow & E^n \\ f^n_{|\mathcal{F}^n(E)} \downarrow & & \downarrow f^n \\ \mathcal{F}^n(E) & \hookrightarrow & E^n \end{array}$$

and by functoriality of the Leray spectral sequence it induces the action of  $SL_2(\mathbb{Z})$  on  $A^{\bullet,\bullet}$ . We explicitly describe this action on the generators  $\omega_{i,j}$ ,  $x_i$ , and  $y_i$ : since  $f^n: E^n \rightarrow E^n$  fixes the divisor  $\{p_i = p_j\}$ , then  $f \cdot \omega_{i,j} = \omega_{i,j}$ . The other generators belongs to  $A^{1,0} = H^1(E^n) \cong H^1(E)^{\otimes n}$ . Therefore the action of  $\text{MCG}(E) \cong SL_2(\mathbb{Z})$  on  $A^{1,0}$  is given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot x_i = ax_i + cy_i \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot y_i = bx_i + dy_i.$$

This action extends to  $SL_2(\mathbb{Q})$  and since the actions of  $\mathfrak{S}_n$  and of  $SL_2(\mathbb{Q})$  commute, then  $A^{\bullet,\bullet}$ ,  $B^{\bullet,\bullet}$  and  $D^{\bullet,0}$  become  $\mathfrak{S}_n \times SL_2(\mathbb{Q})$ -modules.

## 2.2 Decomposition into $\mathfrak{S}_n$ -representations

We recall a result of [3, Theorem 3.15] on the decomposition of  $A^{\bullet,\bullet}$  into  $\mathfrak{S}_n$ -modules. The notations used here follow the ones in [3].

Let  $L_* = (\lambda_1, \dots, \lambda_t)$  be a partition of the number  $n$ , ie  $\lambda_i \in \mathbb{N}_+$  and  $\sum_{i=1}^t \lambda_i = n$ . We mark all blocks with labels in  $\{1, x, y, xy\}$ , an ordered basis of  $H^*(E)$ . The order is  $1 \prec x \prec y \prec xy$ .

**Definition 2.2** A marked partition  $(L_*, H_*)$  is a partition  $L_* \vdash n$  together with marks  $H_* = (h_1, \dots, h_t)$ ,  $h_i \in \{1, x, y, xy\}$  such that: if  $\lambda_i = \lambda_{i+1}$  then  $h_i \succeq h_{i+1}$ .

Let  $C_k$  be the cyclic group of order  $k$ . For any partition  $L_* \vdash n$  define  $C_{L_*}$  as the product of the cyclic groups  $C_{\lambda_i}$  for  $i = 1, \dots, t$ . It acts on  $\{1, \dots, n\}$  in the natural way. Consider a marked partition  $(L_*, H_*)$  and define  $N_{L_*, H_*}$  as the group that permutes the blocks of  $L_*$  with the same labels. The group  $N_{L_*, H_*}$  is a product of symmetric groups. Call  $Z_{L_*, H_*}$  the semidirect product  $C_{L_*} \rtimes N_{L_*, H_*}$ .

**Example 2.3** Let  $(L_*, H_*)$  be the marked partition  $L_* = (5, 5, 5, 5, 1, 1, 1) \vdash 23$  and  $H_* = (xy, xy, xy, 1, x, x, x)$ . The group  $C_{L_*} \cong (\mathbb{Z}/5\mathbb{Z})^4 < \mathfrak{S}_{23}$  is generated by  $(1, 2, 3, 4, 5), (6, 7, 8, 9, 10), (11, 12, 13, 14, 15)$ , and  $(16, 17, 18, 19, 20)$ . The subgroup  $N_{L_*, H_*} \cong \mathfrak{S}_3 \times \mathfrak{S}_3$  is generated by the permutations  $(1, 6)(2, 7)(3, 8)(4, 9)(5, 10)$ ,  $(1, 11)(2, 12)(3, 13)(4, 14)(5, 15)$ ,  $(21, 22)$ , and  $(21, 23)$ . Finally,  $Z_{L_*, H_*}$  is a group isomorphic to  $(\mathbb{Z}/5\mathbb{Z} \wr \mathfrak{S}_3) \times \mathbb{Z}/5\mathbb{Z} \times \mathfrak{S}_3$ .

Given two representations  $V, W$  of two groups  $G$  and  $H$  respectively, define the tensor representation  $V \boxtimes W$  of  $G \times H$  by the vector space  $V \otimes W$  with the action  $(g, h)(v \otimes w) = g(v) \otimes h(w)$ .

We define the following one-dimensional representations. Let  $\varphi_n$  be a faithful character of the cyclic group and  $\varphi_{L_*}$  the character of  $C_{L_*} \cong \mathbb{Z}/\lambda_1\mathbb{Z} \times \dots \times \mathbb{Z}/\lambda_t\mathbb{Z}$  given by

$$\varphi_{L_*} \stackrel{\text{def}}{=} \text{sgn}_n|_{C_{L_*}} \cdot (\varphi_{\lambda_1} \boxtimes \dots \boxtimes \varphi_{\lambda_t}).$$

Recall that the degree  $\deg$  of  $1, x, y, xy$  are respectively  $0, 1, 1, 2$ . Let  $\alpha_{L_*, H_*}$  be the one dimensional representation of  $N_{L_*, H_*} \cong \mathfrak{S}_{\mu_1} \times \dots \times \mathfrak{S}_{\mu_t}$  defined on generators by

$$\alpha_{L_*, H_*}(\sigma) \stackrel{\text{def}}{=} (-1)^{\lambda + \deg(h) + 1},$$

where  $\sigma$  is the permutation that exchange two blocks of size  $\lambda$  and label  $h$ . Set  $\xi_{L_*, H_*}$  to be the one dimensional representation of  $Z_{L_*, H_*}$  such that  $\text{Res}_{C_{L_*}}^{Z_{L_*, H_*}} \xi_{L_*, H_*} = \varphi_{L_*}$  and  $\text{Res}_{N_{L_*, H_*}}^{Z_{L_*, H_*}} \alpha_{L_*, H_*}$ .

We define  $|L_*| = n - t$  for a partition  $L_* = (\lambda_1, \dots, \lambda_t)$  of  $n$  and for a mark  $H_*$  the numbers  $|H_*| = \sum_{i=1}^t \deg(h_i)$  and  $\|H_*\| = |\{i \mid h_i = x\}| - |\{i \mid h_i = y\}|$ .

**Definition 2.4** Let  $(L_*, H_*)$  be a marked partition and set  $p = |L_*|$  and  $q = |H_*|$ . Define  $A_{L_*, H_*} \subseteq A^{p,q}$  as the  $\mathfrak{S}_n$ -subrepresentation generated by the following element:

$$m_{L_*, H_*} \stackrel{\text{def}}{=} \prod_{i=1}^t (h_i)_{l_i+1} \prod_{j=1}^{\lambda_i-1} \omega_{l_i+j, l_i+j+1},$$

where  $l_i = \sum_{k < i} \lambda_k$ .

**Theorem 2.5** ([3, Theorem 3.15]) *The  $\mathfrak{S}_n$ -representation  $A^{p,q}$  decomposes as*

$$A^{p,q} = \bigoplus_{\substack{|L_*|=q \\ |H_*|=p}} A_{L_*, H_*}.$$

Moreover:

$$A_{L_*, H_*} \otimes_{\mathbb{Q}} \mathbb{C} \cong \text{Ind}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \xi_{L_*, H_*}.$$

**Remark 2.6** Observe that:

- The  $n$ -Lie operad over the complex numbers is described as the  $\mathfrak{S}_n$ -representation induced from  $C_n$  by a faithful character.
- The direct sum over the natural numbers of all homological Križ models is isomorphic to the Chevalley-Eilenberg complex  $\text{CE}(\mathfrak{g}_E[1])$ , where the Lie algebra  $\mathfrak{g}_E$  is  $H^*(E; \mathbb{C}) \otimes \text{Lie}_n$ .

These two facts gives a conceptual explanation to [Theorem 2.5](#). For references see [\[8, 21, 20\]](#).

**Example 2.7** Consider the marked partition  $(L_*, H_*)$  of [Theorem 2.3](#). We have  $|L_*| = 16$ ,  $|H_*| = 9$  and  $\|H_*\| = 3$ . The element generating  $A_{L_*, H_*}$  is the following:

$$m_{L_*, H_*} = x_1 y_1 x_6 y_6 x_{11} y_{11} x_{21} x_{22} x_{23} \omega_{1,2} \omega_{2,3} \omega_{3,4} \omega_{4,5} \omega_{6,7} \omega_{7,8} \omega_{8,9} \omega_{9,10} \cdot \\ \omega_{11,12} \omega_{12,13} \omega_{13,14} \omega_{14,15} \omega_{16,17} \omega_{17,18} \omega_{18,19} \omega_{19,20}$$

The characters associated with  $(L_*, H_*)$  are shown in the following table.

	(1, 2, 3, 4, 5)	(16, 17, 18, 19, 20)	(1, 6)(2, 7)(3, 8)(4, 9)(5, 10)	(21, 22)
$\varphi$	$\zeta_5$	$\zeta_5$		
$\alpha$			1	-1
$\xi$	$\zeta_5$	$\zeta_5$	1	-1



### 2.3 Decomposition into $\mathfrak{S}_n \times SL_2(\mathbb{Q})$ -representations

Let  $T = \{H_t\} \cong \mathbb{Q}^*$  be the maximal torus in  $SL_2(\mathbb{Q})$  generated by the diagonal matrices  $H_t = \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}$ . Let  $\mathbb{V}_1$  be the irreducible representation  $\mathbb{Q}^2$  with the standard action of matrix-vector multiplication and let  $\mathbb{V}_k = S^k \mathbb{V}_1$  be the irreducible representation given by the symmetric power of  $\mathbb{V}_1$ . The representation  $\mathbb{V}_k$  has dimension  $k + 1$  and can be view as  $\mathbb{Q}[x, y]_k$ , ie the vector space of homogeneous polynomials in two variables. The action of  $T$  on the monomials is given by  $H_t \cdot x^a y^{k-a} = t^{2a-k} x^a y^{k-a}$ , thus  $\mathbb{V}_k$  decomposes, as representations of  $T$

$$(2) \quad \mathbb{V}_k = \bigoplus_{a=0}^k V(2a - k),$$

where  $V(2a - k)$  is the subspace where  $H_t$  acts with character  $t^{2a-k}$ , ie the subspace generated by  $x^a y^{k-a}$ . The algebraic group  $SL_2(\mathbb{Q})$  is reductive, so by [24, Theorem 22.42] each representation of  $SL_2(\mathbb{Q})$  is semisimple. The Fundamental Theorem [24, Theorem 22.2] says that each irreducible representation of  $SL_2(\mathbb{Q})$  is isomorphic to the representation  $\mathbb{V}_k$  described above, for an unique  $k \in \mathbb{N}$ .

As a consequence we can decompose a representation  $V$  of  $SL_2(\mathbb{Q})$  using its decomposition  $V = \bigoplus_{a \in \mathbb{Z}} V(a)^{\oplus n_a}$  as representation of  $T$ : indeed  $V \cong \bigoplus_{k \in \mathbb{N}} \mathbb{V}_k^{\oplus m_k}$  as representation of  $SL_2(\mathbb{Q})$ , where  $m_k = n_k - n_{k+2}$ . By setting  $V = \mathbb{V}_m \otimes \mathbb{V}_n$ , we obtain the following formula for  $n \leq m$ :

$$\mathbb{V}_m \otimes \mathbb{V}_n \cong \mathbb{V}_{m+n} \oplus \mathbb{V}_{m+n-2} \oplus \cdots \oplus \mathbb{V}_{m-n}.$$

As observed in Section 2.1, the group  $SL_2(\mathbb{Q})$  acts trivially on  $\omega_{i,j}$  for all  $1 \leq i < j \leq n$  and the two dimensional subspace generated by  $x_i$  and  $y_i$  is isomorphic to  $\mathbb{V}_1$  as representation of  $SL_2(\mathbb{Q})$ .

We will use the decomposition of Theorem 2.5 to obtain a decomposition of  $A^{\bullet, \bullet}$  into  $\mathfrak{S}_n \times SL_2(\mathbb{Q})$ -modules. Define  $A_a^{p,q}$  as the following direct sum:

$$A_a^{p,q} \stackrel{\text{def}}{=} \bigoplus_{\substack{|L_*|=q \\ |H_*|=p, \|H_*\|=a}} A_{L_*, H_*}.$$

Let  $E = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in SL_2(\mathbb{Q})$  and consider the map  $\iota: A^{p,q} \rightarrow A^{p,q}$  defined by  $v \mapsto v - E \cdot v$ . Notice that  $\iota(x_i) = 0$ ,  $\iota(y_i) = x_i$  and  $\iota(\omega_{i,j}) = 0$ , for all  $i \neq j$ , hence it induces a restricted map  $\iota_a: A_a^{p,q} \rightarrow A_{a+2}^{p,q}$ .

**Lemma 2.8** *For all  $a \geq 0$ , the map  $\iota_a$  is surjective.*

**Proof** Let  $W$  be an irreducible  $SL_2(\mathbb{Q})$ -representation in  $A^{p,q}$  and notice that  $\iota(W) \subseteq W$ . By the arbitrariness of  $W$ , it is enough to prove that  $W \cap A_{a+2}^{p,q}$  is contained in  $\text{Im } \iota_a$ .

Since  $W \cong \mathbb{V}_k$  for some  $k$ , we can reduce to work in  $\mathbb{V}_k$ . If  $k < a+2$  or  $k \not\equiv a \pmod{2}$ , then the torus  $T$  acts with weight  $a+2$  on  $A_{a+2}^{p,q}$ . The weight  $a+2$  does not appear in  $W$ , therefore  $W \cap A_{a+2}^{p,q} = 0$ . Otherwise,  $k = a+2+2b$  for some  $b \in \mathbb{N}$  and  $W \cap A_{a+2}^{p,q}$  is the one dimensional vector space generated by the image of  $x^{a+2+b}y^b \in \mathbb{V}_k$ . The identity

$$(\text{Id} - E) \cdot \frac{x^{a+1+b}y^{b+1}}{b+1} = x^{a+2+b}y^b,$$

completes the proof.  $\square$

**Theorem 2.9** *The algebra  $A^{\bullet,\bullet}$  decomposes as  $\mathfrak{S}_n \times SL_2(\mathbb{Q})$ -representation in the following way:*

$$A^{p,q} \cong \bigoplus_{a=0}^p \ker \iota_a \boxtimes \mathbb{V}_a.$$

Moreover,  $\dim \ker \iota_a$  is the difference between  $\dim A_a^{p,q}$  and  $\dim A_{a+2}^{p,q}$ .

**Proof** The assertion on the dimension of  $\ker \iota_a$  follows from [Theorem 2.8](#). Let  $W_a$  be the  $\mathfrak{S}_n \times SL_2(\mathbb{Q})$ -subrepresentation generated by  $\ker \iota_a$ . It is isomorphic to  $\ker \iota_a \boxtimes \mathbb{V}_a$  because all elements in  $\ker \iota_a$  are of highest weight  $a$ . Thus  $A^{p,q} \supseteq \sum_{a=0}^p W_a$  and the sum is direct since the  $SL_2(\mathbb{Q})$ -representations  $\mathbb{V}_a$  are pairwise non-isomorphic. We have proven that  $A^{p,q} \supseteq \bigoplus_{a=0}^p W_a$  and  $W_a \cong \ker \iota_a \boxtimes \mathbb{V}_a$ . We complete the proof by a dimensional reasoning:

$$\begin{aligned} \dim A^{p,q} &= \sum_{a=-p}^p \dim A_a^{p,q} = -\dim A_0^{p,q} + 2 \sum_{a=0}^p \dim A_a^{p,q} \\ &= \sum_{a=0}^p (\dim A_a^{p,q} - \dim A_{a+2}^{p,q})(a+1) = \sum_{a=0}^p \dim \ker \iota_a \cdot \dim \mathbb{V}_a \\ &= \dim \bigoplus_{a=0}^p W_a. \end{aligned}$$

Since  $A^{p,q}$  and  $\bigoplus_{a=0}^p W_a$  have the same dimension, they are equal.  $\square$

Define the  $\mathfrak{S}_n$ -invariant subalgebra of  $A^{\bullet,\bullet}$  by  $UA^{\bullet,\bullet}$  and of  $B^{\bullet,\bullet}$  by  $UB^{\bullet,\bullet}$ . Obviously we have  $UA^{\bullet,\bullet} = UB^{\bullet,\bullet} \otimes_{\mathbb{Q}} D^{\bullet,\bullet}$ . We use the previous calculation to compute  $UA^{\bullet,\bullet}$ .

**Corollary 2.10** *For  $q > p+1$  we have  $UA^{p,q} = 0$ .*

**Proof** Let  $\mathbb{1}_n$  be the trivial representation of  $\mathfrak{S}_n$ . We use [Theorem 2.5](#) to show that

$$\langle \mathbb{1}_n, A^{p,q} \rangle_{\mathfrak{S}_n} = 0$$

for  $q > p + 1$ . Indeed, it is enough to prove that

$$\langle \mathbb{1}_n, \text{Ind}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \xi_{L_*, H_*} \rangle_{\mathfrak{S}_n} = 0$$

for all  $(L_*, H_*)$  with  $|L_*| = q$  and  $|H_*| = p$ . By Frobenius reciprocity we have

$$\langle \mathbb{1}_n, \text{Ind}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \xi_{L_*, H_*} \rangle_{\mathfrak{S}_n} = \langle \text{Res}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \mathbb{1}_n, \xi_{L_*, H_*} \rangle_{Z_{L_*, H_*}}$$

Since the representations in the right hand side are one-dimensional the value of  $\langle \mathbb{1}_n, \text{Ind}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \xi_{L_*, H_*} \rangle_{\mathfrak{S}_n}$  is non zero if and only if  $\xi_{L_*, H_*} = \mathbb{1}$ .

By definition  $\xi_{L_*, H_*} = \mathbb{1}$  is equivalent to  $\varphi_{L_*} = \mathbb{1}$  and  $\alpha_{L_*, H_*} = \mathbb{1}$ . From the fact that  $\varphi_k = \text{sgn}_k$  only for  $k = 1, 2$ ,  $\psi_{L_*} = \mathbb{1}$  if and only if  $\lambda_i = 1, 2$  for all  $i = 1, \dots, t$ . The condition  $\alpha_{L_*, H_*} = \mathbb{1}$  implies that the only marked blocks of  $(L_*, H_*)$  that appear more than once are the ones with  $\lambda_i = 2$  and  $\deg(h_i) = 1$  or the ones with  $\lambda_i = 1$  and  $\deg(h_i) \neq 1$ .

Consequently,  $\langle \mathbb{1}_n, \text{Ind}_{Z_{L_*, H_*}}^{\mathfrak{S}_n} \xi_{L_*, H_*} \rangle_{\mathfrak{S}_n} \neq 0$  only if  $L_* = (2^q, 1^{n-2q})$  and the degree of  $h_i$  is 1 for  $i < q$ , this implies  $p \geq q - 1$  contrary to our hypothesis.  $\square$

**Corollary 2.11** For  $q > p + 1$  we have  $UB^{p,q} = 0$ .  $\square$

### 3 The additive structure of the cohomology

We compute the cohomology with rational coefficients of the unordered configuration spaces of  $n$  points, taking care of the mixed Hodge structure and of the action of  $SL_2(\mathbb{Q})$ . The integral cohomology groups are known only for small  $n$  in [\[25, Table 2\]](#), where a cellular decomposition of ordered configuration spaces is given. In this section, we use the calculation of the Betti numbers of  $\mathcal{C}^n(E)$  to determine the Hodge polynomial in the Grothendieck ring of  $SL_2(\mathbb{Q})$ .

Observe that  $H^*(\mathcal{C}^n(E)) = H^*(\mathcal{F}^n(E))^{\mathfrak{S}_n}$  by the Transfer Theorem. Define the series

$$T(u, v) = \frac{1 + u^3 v^4}{(1 - u^2 v^3)^2} = 1 + 2u^2 v^3 + u^3 v^4 + 3u^4 v^6 + 2u^5 v^7 + \dots$$

and let  $T_n(u, v)$  be its truncation at degree  $n$  in the variable  $u$ .

The computation of the Betti numbers of unordered configuration space of  $n$  points in an elliptic curve was done simultaneously by [\[11\]](#), [\[23\]](#), and [\[27\]](#) in different generality. We point to the last reference because [\[27, Theorem\]](#) fits exactly our generality.

**Theorem 3.1** *The Poincaré polynomial of  $\mathcal{C}^n(E)$  is  $(1+t)^2 T_{n-1}(t, 1)$ .*

We use the notation  $Vu^k v^h$  to denote a vector space  $V$  in degree  $k$  with a Hodge structure of weight  $h$ . The Grothendieck ring of  $SL_2(\mathbb{Q})$  is the free  $\mathbb{Z}$ -module with basis given by  $[V]$  for all finite-dimensional irreducible representations  $V$  of  $SL_2(\mathbb{Q})$  and product defined by the tensor product of representations.

**Definition 3.2** The Hodge polynomial of  $\mathcal{C}^n(E)$  with coefficients in the Grothendieck ring of  $SL_2(\mathbb{Q})$  is

$$\sum_{i=0}^{2n} \sum_{k=i}^{2i} \left[ W_k H^i(\mathcal{C}^n(E); \mathbb{Q}) / W_{k-1} H^i(\mathcal{C}^n(E); \mathbb{Q}) \right] u^i v^k,$$

where  $W_k H^i(\mathcal{C}^n(E); \mathbb{Q})$  is the weight filtration on  $H^i(\mathcal{C}^n(E); \mathbb{Q})$ . The ordinary Hodge polynomial is

$$\sum_{i=0}^{2n} \sum_{k=i}^{2i} \dim_{\mathbb{Q}} \left( W_k H^i(\mathcal{C}^n(E); \mathbb{Q}) / W_{k-1} H^i(\mathcal{C}^n(E); \mathbb{Q}) \right) u^i v^k.$$

We prove a stronger version of [Theorem 3.1](#).

**Theorem 3.3** *The Hodge polynomial of  $\mathcal{C}^n(E)$  with coefficients in the Grothendieck ring of  $SL_2(\mathbb{Q})$  is*

$$(3) \quad ([\mathbb{V}_0] + [\mathbb{V}_1]uv + [\mathbb{V}_0]u^2v^2) \left( \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} [\mathbb{V}_i]u^{2i}v^{3i} + \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - 1} [\mathbb{V}_{i-1}]u^{2i+1}v^{3i+1} \right)$$

and the ordinary Hodge polynomial is  $(1+uv)^2 T_{n-1}(u, v)$ .


[Figure 1](#) represents the module  $H^{\bullet,*}(UB)$  that corresponds to the right factor of eq. (3).

### 3.1 Some elements in cohomology

**Definition 3.4** Let  $\alpha, \bar{\alpha} \in A^{1,1}, \beta \in A^{1,2}$  be the elements

$$\begin{aligned} \alpha &\stackrel{\text{def}}{=} \sum_{i,k < h} (x_i - x_k) \omega_{k,h} \\ \bar{\alpha} &\stackrel{\text{def}}{=} \sum_{i,k < h} (y_i - y_k) \omega_{k,h} \\ \beta &\stackrel{\text{def}}{=} \sum_{i,j,k < h} (3x_i - x_j - 2x_k)(y_j - y_k) \omega_{k,h} \end{aligned}$$

where the sum is taken over pairwise distinct indices  $i, j, k, h$  with  $k < h$ .

Figure 1: The algebra  $H^{\bullet,\bullet}(UB)$  as representation of  $SL_2(\mathbb{Q})$ .

Notice that the elements  $\alpha$  and  $\bar{\alpha}$  are defined only for  $n > 2$  and  $\beta$  for  $n > 3$ . Remember that  $\gamma, \bar{\gamma} \in D^1$  were already defined as  $\sum_i x_i$  and  $\sum_i y_i$ .

**Lemma 3.5** *The element  $\alpha$  belongs to  $UB^{1,1}$ , is non-zero, and  $d\alpha = 0$ .*

**Proof** First observe that  $\alpha = \sum_{i,k < h} u_{i,k} \omega_{k,h} \in B^{1,1}$ . For all  $\sigma \in \mathfrak{S}_n$  we have

$$\sigma\alpha = \sum_{i,k < h} u_{\sigma(i),\sigma(k)} \omega_{\sigma(k),\sigma(h)} = \alpha,$$

since  $u_{\sigma(i),\sigma(k)} \omega_{\sigma(k),\sigma(h)} = u_{\sigma(i),\sigma(h)} \omega_{\sigma(k),\sigma(h)}$  in  $A$ . The elements  $x_i \omega_{k,h}$  and  $x_k \omega_{k,h}$  are linearly independent, so it is enough to observe that the coefficient of  $x_3 \omega_{1,2}$  is 1. This proves that  $\alpha \neq 0$ . Finally, we compute  $d\alpha$ :

$$\begin{aligned} d\alpha &= \sum_{i,k < h} x_i d\omega_{k,h} - x_k d\omega_{k,h} \\ &= \sum_{i,k < h} x_i(x_k - x_h)(y_k - y_h) + x_k x_h(y_k - y_h) \\ &= \sum_{i,k,h} x_i x_k y_k - x_i x_h y_k + x_k x_h y_k \\ &= - \sum_{i,k,h} x_i x_h y_k = 0, \end{aligned}$$

where all sums are taken over pairwise distinct indices and the first two with the additional condition  $k < h$ .  $\square$

**Lemma 3.6** *The element  $\beta$  belongs to  $UB^{1,2}$ , is non-zero, and  $d\beta = 0$ .*

**Proof** Observe that

$$\beta = \sum_{i,j,k < h} (u_{i,j} + 2u_{i,k})v_{j,k}\omega_{k,h} \in B^{1,2}$$

and that  $\sigma\beta = \beta$  for all  $\sigma \in \mathfrak{S}_n$  by the relations  $u_{j,k}\omega_{k,h} = u_{j,h}\omega_{k,h}$  and  $v_{j,k}\omega_{k,h} = v_{j,h}\omega_{k,h}$ . Consider the map  $\varphi: A \rightarrow \mathbb{Q}$  defined on generators by  $\varphi(\omega_{1,2}) = 1$ ,  $\varphi(x_3) = 1$  and  $\varphi(y_4) = 1$  and zero on the other generators. The map  $\varphi$  is well defined and  $\varphi(\beta) = 3$ , thus  $\beta \neq 0$ . Using the computation in the proof of [Theorem 3.5](#), we can observe that  $d(\sum_{i,j,h < k} 3x_i(y_j - y_k)\omega_{k,h}) = 0$ . The claim  $d\beta = 0$  follows from:

$$\begin{aligned} d(\beta) &= d\left(\sum_{j,k,h} (x_j + 2x_k)(y_j - y_k)\omega_{k,h}\right) \\ &= \sum_{j,k < h} x_j y_j d\omega_{k,h} + x_j y_k (x_k - x_h) y_h - 2x_k y_j x_h (y_k - y_h) - 2x_k y_k x_h y_h \\ &= \sum_{j,k,h} x_j y_j x_k y_k - x_i y_j x_h y_k + x_j y_k x_k y_h - 2x_k y_j x_h y_k - x_k y_k x_h y_h \\ &= 0, \end{aligned}$$

where the indexes of the sums are pairwise distinct.  $\square$

**Lemma 3.7** *For  $n > 2q$  the element  $\alpha^q$  is non-zero.*

**Proof** We show that the coefficient of the monomial  $m = x_1\omega_{1,2}x_3\omega_{3,4} \dots x_{2q-1}\omega_{2q-1,2q}$  (defined for  $n \geq 2q$ ) in  $\alpha^q$  is non-zero for  $n > 2q$ . Initially, we will prove that the coefficient is

$$(4) \quad a_q = q! \sum_{\sigma \in \mathfrak{S}_q} \text{sgn}(\sigma) (2-n)^{|\text{Fix } \sigma|} 2^{q-|\text{Fix } \sigma|},$$

then that  $a_q \neq 0$  for  $n > 2q$ .

We start with the following identity:

$$\alpha^q = \left( \sum_{j < k} \sum_{i \neq j,k} (x_i - x_j) \omega_{j,k} \right)^q = \sum_{J,K} \prod_{l=1}^q \left( \sum_{i \neq j_l, k_l} (x_i - x_{j_l}) \right) \omega_{j_l, k_l},$$

where the sum is taken over all multi-indexes  $J = \{j_1, \dots, j_q\}$  and  $K = \{k_1, \dots, k_q\}$  with  $j_l < k_l$  for all  $l = 1, \dots, q$ . Since, for all  $l = 1, \dots, q$ , the pair of indexes  $(2l-1, 2l)$  appear in the monomial  $m$ , the monomial  $m$  arise only in the terms with the

following property  $\mathcal{P}$ : there exist  $\sigma \in \mathfrak{S}_q$  such that  $j_l = 2\sigma(l) - 1$  and  $k_l = 2\sigma(l)$  for all  $l = 1, \dots, q$ . We further manipulate it:

$$\begin{aligned} \sum_{(J,K) \in \mathcal{P}} \prod_{l=1}^q \left( \sum_{i \neq j_l, k_l} (x_i - x_{j_l}) \right) \omega_{j_l, k_l} &= \sum_{\sigma \in \mathfrak{S}_q} \prod_{l=1}^q \left( \sum_{i \neq 2\sigma(l)-1, 2\sigma(l)} (x_i - x_{2\sigma(l)-1}) \right) \omega_{2\sigma(l)-1, 2\sigma(l)} \\ &= q! \prod_{l=1}^q \left( \sum_{i \neq 2l-1, 2l} (x_i - x_{2l-1}) \right) \omega_{2l-1, 2l} \\ &= q! \prod_{l=1}^q \left( 2 \sum_{\substack{i=1, \dots, q \\ i \neq l}} x_{2i-1} + \sum_{i > 2q} x_i - (n-2)x_{2l-1} \right) \omega_{2l-1, 2l}, \end{aligned}$$

where the last equality is obtained using  $x_j \omega_{j,k} = x_k \omega_{j,k}$ . Since the variable  $x_i$  with  $i > 2q$  does not appear in  $m$ , the coefficient  $a_q$  is equal to the coefficient of  $x_1 x_3, \dots, x_{2q-1}$  in  $q! \prod_{l=1}^q (2 \sum_{i \neq l} x_{2i-1} - (n-2)x_{2l-1})$ . We have:

$$\begin{aligned} \prod_{l=1}^q (2 \sum_{i \neq l} x_{2i-1} - (n-2)x_{2l-1}) &= \sum_{\sigma \in \mathfrak{S}_q} \prod_{l=1}^q c_{\sigma(l), l} x_{2\sigma(l)-1} \\ &= \sum_{\sigma \in \mathfrak{S}_q} \text{sgn } \sigma \prod_{l=1}^q c_{\sigma(l), l} x_{2l-1} \\ &= \sum_{\sigma \in \mathfrak{S}_q} \text{sgn } \sigma (2-n)^{|\text{Fix } \sigma|} 2^{q-|\text{Fix } \sigma|} \prod_{l=1}^q x_{2l-1}, \end{aligned}$$

where  $c_{k,l} = 2-n$  if  $k = l$  and  $c_{k,l} = 2$  otherwise. Putting all together we obtain [Equation 4](#).

We claim that

$$(5) \quad \sum_{\sigma \in \mathfrak{S}_q} \text{sgn}(\sigma) x^{|\text{Fix } \sigma|} = (x-1)^{q-1} (x+q-1),$$

since both sides are the determinant of the matrix

$$\begin{pmatrix} x & 1 & \cdots & 1 \\ 1 & x & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & x \end{pmatrix}.$$

The left hand side of eq (5) is obtained by using the Laplace formula for the determinant and the right hand side by relating the that determinant to the characteristic polynomial

$(-t)^{q-1}(q-t)$  of the matrix  $\begin{pmatrix} 1 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 1 \end{pmatrix}$ . We use eq (5) with  $x = \frac{2-n}{2}$  to obtain:

$$a_q = q!2^q \sum_{\sigma \in \mathfrak{S}_q} \text{sgn}(\sigma) \left( \frac{2-n}{2} \right)^{|\text{Fix } \sigma|} = q!2^q \left( \frac{-n}{2} \right)^{q-1} \left( \frac{2q-n}{2} \right)$$

Thus  $a_q = (-1)^q q! n^{q-1} (n-2q)$  that is non-zero for  $n > 2q$ .  $\square$

**Lemma 3.8** For  $n > 2q + 1$  the element  $\alpha^{q-1}\beta$  is non-zero.

**Proof** Let us rewrite  $\beta$  as

$$\begin{aligned} \beta = \sum_{i,j,k < h} x_i y_j \omega_{k,h} - 2(n-3) \sum_{i,k < h} (x_i y_k + x_k y_i) \omega_{k,h} - (n-3) \sum_{i,k < h} x_i y_i \omega_{k,h} + \\ + 2(n-2)(n-3) \sum_{k < h} x_k y_k \omega_{k,h}. \end{aligned}$$

Let  $b_q$  be the coefficient in  $\alpha^{q-1}\beta$  of the monomial

$$x_1 \omega_{1,2} x_3 \omega_{3,4} \dots x_{2q-1} \omega_{2q-1,2q} y_{2q+1}.$$

This monomial is defined for  $n \geq 2q + 1$  and we will show that  $b_q \neq 0$  for  $n > 2q + 1$ . The number  $b_q$  coincides with the coefficient of the same monomial in the product

$$\alpha^{q-1} \left( \sum_{i,j,k < h} x_i y_j \omega_{k,h} - 2(n-3) \sum_{i,k < h} x_k y_i \omega_{k,h} \right).$$

With further manipulation, we obtain that  $b_q$  is the coefficient of the above monomial in the expression

$$3\alpha^q y_{2q+1} + n\alpha^{q-1} \sum_{k < h} x_k \omega_{k,h} y_{2q+1}.$$

Using the computation in the proof of Theorem 3.7 we obtain

$$\begin{aligned} b_q &= 3(-1)^q q! n^{q-1} (n-2q) + nq(-1)^{q-1} (q-1)! n^{q-2} (n-2q+2) \\ &= 2(-1)^q q! n^{q-1} (n-2q-1). \end{aligned}$$

The number  $b_q$  is non zero for  $n > 2q + 1$ .  $\square$

**Proof of Theorem 3.3** It is enough to prove that the Hodge polynomial of  $UB$  in the Grothendieck ring of  $SL_2(\mathbb{Q})$  is

$$\sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} [\mathbb{V}_i] u^{2i} v^{3i} + \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - 1} [\mathbb{V}_{i-1}] u^{2i+1} v^{3i+1}$$



Observe that  $\text{Im } d^{q,p} = 0$  for  $q > p + 1$  by [Theorem 2.11](#). From [Theorem 3.5](#) and [Theorem 3.7](#) we have that the elements  $\alpha^k$  for  $2k < n$  generate as  $SL_2(\mathbb{Q})$ -module a subspace of dimension at least  $k + 1$  in  $H^{k,k}(UB, \mathfrak{d})$ . Analogously, from [Theorem 3.6](#) and [Theorem 3.8](#) the elements  $\alpha^{k-1}\beta$  for  $2k + 1 < n$  generate as  $SL_2(\mathbb{Q})$ -module a subspace of dimension at least  $k$  in  $H^{k,k+1}(UB, \mathfrak{d})$ . Since the Betti numbers of  $UB^{\bullet,\bullet}$  ([Theorem 3.1](#)) coincides with the above dimensions, we have that  $H^{2k}(UB) \cong \mathbb{V}_k u^{2k} v^{3k}$  and  $H^{2k+1}(UB) \cong \mathbb{V}_{k-1} u^{2k+1} v^{3k+1}$ .  $\square$

## 4 The cohomology ring

In this section we determine the cup product structure in the cohomology of  $\mathcal{C}^n(E)$  and we prove the formality result.

In the following we consider graded algebras with an action of  $SL_2(\mathbb{Q})$ . We will write  $(x_i \mid i \in I)_{SL_2(\mathbb{Q})}$  for the ideal generated by the elements  $Mx_i$  for all  $M \in SL_2(\mathbb{Q})$  and all  $i \in I$ .

**Theorem 4.1** *The cohomology ring of  $\mathcal{C}^n(E)$  is isomorphic to*

$$\Lambda^\bullet \mathbb{V}_1 \otimes S^\bullet \mathbb{V}_1[b] / (a^{\lfloor \frac{n+1}{2} \rfloor}, a^{\lfloor \frac{n}{2} \rfloor} b, b^2)_{SL_2(\mathbb{Q})},$$

where  $a$  is a non-zero degree-one element in  $V(1) \subset \mathbb{V}_1$  and  $b$  an  $SL_2(\mathbb{Q})$ -invariant indeterminate of degree 3.

**Proof** It is enough to prove that  $H^\bullet(UB) \cong S^\bullet \mathbb{V}_1[b] / (a^{\lfloor \frac{n+1}{2} \rfloor}, a^{\lfloor \frac{n}{2} \rfloor} b, b^2)_{SL_2(\mathbb{Q})}$ . Define the morphism  $\varphi: S^\bullet \mathbb{V}_1[b] / (a^{\lfloor \frac{n+1}{2} \rfloor}, a^{\lfloor \frac{n}{2} \rfloor} b, b^2)_{SL_2(\mathbb{Q})} \rightarrow H^\bullet(UB)$  that sends  $a, b$  to  $\alpha, \beta$  respectively. It is well defined because  $H^k(UB) = 0$  for  $k \geq n$  and  $\beta^2 = 0$  since it has odd degree. The map  $\varphi$  is surjective since  $H^\bullet(UB)$  is generated by  $\alpha^i$  and  $\alpha^i \beta$  as  $SL_2(\mathbb{Q})$ -module by [Theorem 3.3](#). A dimensional reasoning shows the injectivity of the map  $\varphi$ .  $\square$

**Corollary 4.2** *The cohomology  $H^\bullet(\mathcal{C}^n(E))$  is generated as an algebra in degrees one, two and three.*

**Proof** A minimal set of generators is given by  $\alpha, \bar{\alpha}, \beta, \gamma, \bar{\gamma}$ .  $\square$

**Corollary 4.3** *The space  $\mathcal{C}^n(E)$  is formal over the rationals.*

**Proof** We prove that  $UB$  is formal. Consider the subalgebra  $K^{\bullet,\bullet}$  of  $UB^{\bullet,\bullet}$  generated by  $\alpha, \bar{\alpha}, \beta$  endowed with the zero differential. It is concentrated in degrees  $(i, i)$  and  $(i, i + 1)$  because  $\beta^2 = 0$ . Since  $K \cap \text{Im } d = 0$  (Theorem 2.11),  $K \hookrightarrow UB$  is a quasi-isomorphism. The fact that  $K \cong H(UB)$  implies that the algebra  $UB$  is formal. As a consequence  $UA$  is formal. The space  $C^n(E)$  is formal since our model  $UA$  is equivalent to the Sullivan model.  $\square$

## Acknowledgements

I would like to thank the referee for several useful suggestions.

## References

- [1] **V I Arnol'd**, *The cohomology ring of the colored braid group*, Mathematical notes of the Academy of Sciences of the USSR 5 (Feb 1969) 138–140
- [2] **V I Arnold**, *On some topological invariants of algebraic functions*, from “Vladimir I. Arnold - Collected Works: Hydrodynamics, Bifurcation Theory, and Algebraic Geometry 1965-1972” (A B Givental, B A Khesin, A N Varchenko, V A Vassiliev, O Y Viro, editors), Springer Berlin Heidelberg, Berlin, Heidelberg (2014) 199–221
- [3] **S Ashraf, H Azam, B Berceanu**, *Representation theory for the Križ model*, Algebr. Geom. Topol. 14 (2014) 57–90
- [4] **H Azam**, *Cohomology groups of configuration spaces of Riemann surfaces*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) 58(106) (2015) 33–47
- [5] **C Bibby**, *Cohomology of abelian arrangements*, Proc. Amer. Math. Soc. 144 (2016) 3093–3104
- [6] **C-F Bökigheimer, F R Cohen**, *Rational cohomology of configuration spaces of surfaces*, from “Algebraic topology and transformation groups (Göttingen, 1987)”, Lecture Notes in Math. 1361, Springer, Berlin (1988) 7–13
- [7] **C-F Bökigheimer, F Cohen, L Taylor**, *On the homology of configuration spaces*, Topology 28 (1989) 111–123
- [8] **C Chevalley, S Eilenberg**, *Cohomology theory of Lie groups and Lie algebras*, Trans. Amer. Math. Soc. 63 (1948) 85–124
- [9] **F R Cohen, T J Lada, J P May**, *The homology of iterated loop spaces*, Lecture Notes in Mathematics, Vol. 533, Springer-Verlag, Berlin-New York (1976)
- [10] **P Deligne**, *Poids dans la cohomologie des variétés algébriques*, from “Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974), Vol. 1”, Canad. Math. Congress, Montreal, Que. (1975) 79–85

- [11] **G C Drummond-Cole, B Knudsen**, *Betti numbers of configuration spaces of surfaces*, J. Lond. Math. Soc. (2) 96 (2017) 367–393
- [12] **C Dupont**, *The Orlik-Solomon model for hypersurface arrangements*, Ann. Inst. Fourier (Grenoble) 65 (2015) 2507–2545
- [13] **B Farb, D Margalit**, *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*, Princeton University Press, Princeton, NJ (2012)
- [14] **Y Félix, D Tanré**, *The cohomology algebra of unordered configuration spaces*, J. London Math. Soc. (2) 72 (2005) 525–544
- [15] **W Fulton, J Harris**, *Representation theory*, volume 129 of *Graduate Texts in Mathematics*, Springer-Verlag, New York (1991) A first course, Readings in Mathematics
- [16] **R Goodman, N R Wallach**, *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*, Springer, Dordrecht (2009)
- [17] **M Goresky, R MacPherson**, *Stratified Morse theory*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, Springer-Verlag, Berlin (1988)
- [18] **A G Gorinov**, *A purity theorem for configuration spaces of smooth compact algebraic varieties*, arXiv e-prints (Feb 2017) [arXiv:1702.08428](https://arxiv.org/abs/1702.08428)
- [19] **B Hall**, *Lie groups, Lie algebras, and representations*, volume 222 of *Graduate Texts in Mathematics*, second edition, Springer, Cham (2015) An elementary introduction
- [20] **N Idrissi**, *The Lambrechts-Stanley model of configuration spaces*, Invent. Math. 216 (2019) 1–68
- [21] **B Knudsen**, *Betti numbers and stability for configuration spaces via factorization homology*, Algebr. Geom. Topol. 17 (2017) 3137–3187
- [22] **I Križ**, *On the rational homotopy type of configuration spaces*, Ann. of Math. (2) 139 (1994) 227–237
- [23] **M Maguire**, *Computing cohomology of configuration spaces*, arXiv e-prints (December 2016) 55 [arXiv:1612.06314](https://arxiv.org/abs/1612.06314)
- [24] **J S Milne**, *Algebraic groups*, volume 170 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge (2017) The theory of group schemes of finite type over a field
- [25] **F Napolitano**, *On the cohomology of configuration spaces on surfaces*, J. London Math. Soc. (2) 68 (2003) 477–492
- [26] **P Salvatore**, *Configuration spaces on the sphere and higher loop spaces*, Math. Z. 248 (2004) 527–540
- [27] **C Schiessl**, *Betti numbers of unordered configuration spaces of the torus*, arXiv e-prints (February 2016) 7 [arXiv:1602.04748](https://arxiv.org/abs/1602.04748)
- [28] **B Totaro**, *Configuration spaces of algebraic varieties*, Topology 35 (1996) 1057–1067

- [29] **C Voisin**, *Hodge theory and complex algebraic geometry. I*, volume 76 of *Cambridge Studies in Advanced Mathematics*, english edition, Cambridge University Press, Cambridge (2007) Translated from the French by Leila Schneps
- [30] **J H Wang**, *On the braid groups for  $\mathbf{RP}^2$* , J. Pure Appl. Algebra 166 (2002) 203–227

Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italia

[roberto.pagaria@sns.it](mailto:roberto.pagaria@sns.it)