

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Phytoplasma presence in carrot seedlings

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Satta E., Carminati G., Bertaccini A. (2020). Phytoplasma presence in carrot seedlings. AUSTRALASIAN PLANT DISEASE NOTES, 15(1), 1-4 [10.1007/s13314-020-0377-y]. *Published Version:*

Availability:

[This version is available at: https://hdl.handle.net/11585/785591 since: 2020-12-26](https://hdl.handle.net/11585/785591)

Published:

[DOI: http://doi.org/10.1007/s13314-020-0377-y](http://doi.org/10.1007/s13314-020-0377-y)

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

> This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

> > (Article begins on next page)

Phytoplasma presence in carrot seedlings Phytoplasma presence in carrot seedlings Eleonora Satta^{1*} el<mark>eonora.satta2@unibo.it</mark>, Gaia Carminati¹, Assunta Bertaccini¹ 1 Department of Agricultural and Food Sciences Alma Mater Studiorum - Università di Bologna 40126 Bologna BO, Italy

Abstract

Phytoplasmas are cell wall lacking bacteria, insect transmitted, and worldwide infecting carrots with severe epidemics. Molecular tests on up to 4four-months-old carrot seedlings from phytoplasma-positive seed batches, carried out under insect proof conditions, indicated that in the early stages of the plant development, the phytoplasma- infected and not water- stressed plants are asymptomatic. The phytoplasma seed transmission could represents a dangerous source of infection.

Keywords

Gg~rowth stage Mmolecular detection PCR/RFLP **Infection percentages**

Phytoplasmas are phloem- limited and insect-transmitted bacteria that lack cell walls and that have been associated in carrot (*Daucus carota*) with redness of leaves, shoot proliferation, and reduced tap roots quality. In Europe, phytoplasmas were often detected in carrot and have been identified as belonging to 16SrI and 16SrXII groups (Font et al. [1999;](#page-4-0) Duduk et al. [2008\)](#page-4-0). Phytoplasmas were often associated with severe epidemics worldwide, and in several cases in carrot they were associated with other pathogens such as spiroplasmas and *'Candidatus Liberibacter solanacearum'* (Cebrián et al. [2010;](#page-3-0) Satta et al. [2016\)](#page-4-0). Symptoms associated with these bacteria are undistinguishable; therefore, the presence of the diverse pathogens must be demonstrated by molecular tools such as specific DNA amplification. Although phytoplasmas are mainly insect vectored, recent studies demonstrated their transmissibility through seed in some plant species such as alfalfa, tomato, pea, corn, and coconut (Khan et al. [2002;](#page-4-0) Zwolinska et al. [2010;](#page-5-0) Calari et al. [2011;](#page-3-0) Oropeza et al. [2017;](#page-4-0) Satta et al. [2017,](#page-4-0) [2019\)](#page-5-0). Symptoms have been observed only in a very low percentage of alfalfa and canola seedlings (Khan et al. [2002;](#page-4-0) Olivier et al. [2010\)](#page-4-0). It has been shown that phytoplasmas can spread to phloem-related organs, such as the cellular parenchyma (Siller et al. [1987\)](#page-5-0) and to companion cells (Sears and Klomparens [1989\)](#page-5-0); however, the way in which it occurs has not been clarified since the dimensions of these bacteria are greater than those of the plasmodesmata, and at the same time, the presence of specific movement proteins has not been detected (Zambryski [2004\)](#page-5-0). Phytoplasma infections, however, are very common in floral structures, fruits, seeds, and

embryos (Bertaccini and Marani [1982;](#page-3-0) Jiang et al. [2004;](#page-4-0) Nipah et al. [2007\)](#page-4-0), while the analyses carried out on pollen did not show their presence (Nečas et al. [2008\)](#page-4-0). The recent success in the cultivation of phytoplasmas isolated from maize seedlings deriving from infected mother plants, demonstrated also their viability in the colonized seedlings (Satta et al. [2019\)](#page-5-0).

Surveys were carried out to verify the presence of phytoplasmas in seed carrots by Carminati et al. [\(2019\)](#page-3-0) in which 29 batches of seeds were tested. Of these, 24 were positive for phytoplasmas. A total of 400 seeds from three lots positive for 16SrI phytoplasmas (lot numbers 12, 15, and 18) and one negative lot (lot number 5) were sown in sterile soil in insect-proof greenhouse under controlled conditions and showed 83.3 to 90% germination irrespectively of the phytoplasma presence.

The DNAs from the same seedlings were extracted at the cotyledon stage and three more times later, up to 4 four-month-old plants with a CTAB method (Angelini et al. [2001\)](#page-3-0) from 1-g of plant tissue. The DNA was used as template after dilution 1:-30 with sterile distilled water (SDW) for the phytoplasma amplification by nested-PCR of the 16S rRNA gene using the R16F2n/R2 (Gundersen and Lee [1996\)](#page-4-0) and R16(I)F1/R1 (Lee et al. [1994\)](#page-4-0). Phytoplasma identification was achieved by RFLP profiles on R16(I)F1/R1 amplicons with *Tru1*I restriction enzyme and direct amplicon sequencing. Phylogenetic analyses were conducted in MEGA6 (Tamura et al. [2013\)](#page-5-0). For results confirmation, amplification was also carried out on the *tuf* gene using the phytoplasma barcode cocktail -primers (Makarova et al. [2012\)](#page-4-0). Phytoplasma strains are maintained in collection (Bertaccini [2014\)](#page-3-0), and SDW were employed as positive and negative controls, respectively.

Among the 340 germinated seeds, 200 seedlings of the four lots were tested in batches (14 seedlings per batch) or as single plants at different times after sprouting. At the cotyledon stage (about three 3 weeks after sowing), they did not show symptoms; however, phytoplasmas belonging to the ribosomal groups 16SrI and 16SrXII, also in double infection (Table [1](#page-5-1)), were detected in 5 batches out of 14 tested (14 cotyledons per batch) from two phytoplasma positive seed lots (numbers 15 and 18). The 2 two-months-old seedlings resulted also symptomless, and 4 batches (of 14 plants each), in all the three phytoplasma infected lots, were positive to 16SrI phytoplasmas (4/[1](#page-5-1)4 batches) (Table -1). Among the 26 4 four-month-old plants singly tested, only one was positive to 16SrI phytoplasma (lot number 12). Further, 140 seedlings, placed for three 3 weeks under water shortage (stressed samples for further phytoplasma seed transmission verification) showed symptoms of root malformation, leaf whitening, and browning (Fig.ure -1 -1), not present in the watered control plants, and were tested after 3 three -months: 5 out of the 14 batches (of 8 seedlings per batch), resulted positive to 16SrI phytoplasmas (Table -[2](#page-6-0)). For the cotyledon and seedlings batches, the estimated proportion of a single sample infection (*p*) was calculated using the maximum likelix hood estimator of *p*: $p = 1 - Q^{1/K}$, where Q is the observed fraction of not infected batches and K is the number of samples per batch (Swallow [1985\)](#page-5-0). The results showed that the *p* index was 0.03 for cotyledons, 0.02 for the 2two-month-old seedlings, and 0.04 for the stressed seedlings. All the tested materials from the phytoplasma negative seed lot number 5 always resulted negative.

The sequences of all detected phytoplasmas enclosed in 16SrI group (*'Candidatus Phytoplasma asteris*"-related strains) resulted 99% identical to each other, and two representative sequences among them were deposited in GenBank (accession numbers MN398390 and MN398391). The sequence of the 16SrXII phytoplasma (*'Ca. P. solani*-related -strain) detected in single infection in the cotyledon was also deposited in GenBank (accession number MN398389) and clustered with a number of strains classified in the 16SrXII-A subgroup ("stolbur"

phytoplasmas). In the phylogenetic tree, the clustering of these sequences confirmed the phytoplasma identification obtained with the RFLP analyses with the R16(I)F1/R1 amplicons (Fig. **ure** -[2](#page-5-3)). The phytoplasma presence and identity was confirmed by the amplification of the *tuf* gene followed by direct sequencing and blasting into the EPPO-Q-bank database (https://www.eppo.int/RESOURCES/eppo_databases/eppo_q_bank). The anomalies and malformations affecting the floral organs and the fruits of plants of several species infected with phytoplasmas allow the production of seeds with reduced vitality (McCoy et al. [1989\)](#page-4-0), quality, and quantity. However, the severity of this phenomenon is related to the time of the infection. Late infections tend to give rise to normal amounts of seed which are however infected (De La Rue et al[. 2002\)](#page-3-0), and will therefore give the pathogen a greater probability of transmission; on the other hand, an early infection will result in such serious alterations of the mother plant that the seed production is reduced. The recent study on phytoplasma transmission to coconut seedlings (Oropeza et al[. 2017\)](#page-4-0) leaves open the possibility that transmission may be less frequent in the natural environment, and suggests that the bacteria may be present in the early stages of life of the seedling and be destroyed in subsequent phases by mechanisms of endogenous defense. The results in this work confirm that the phytoplasma presence in seedlings from infected seed batches is asymptomatic, except in the waterstressed seedlings. The 16SrI and 16SrXII phytoplasma seed transmission in carrot could be relevant in field conditions when the possible insect vectors can further enhance their dispersion using the few seed-infected plants as source of inoculum. The phytoplasma seed transmission appears therefore to be relevant and could represents a dangerous source of inoculum. However, more study should be carried out to evaluate the percentage of seed transmission and the economic impact of phytoplasma seedling infection directly in the carrot fields.

References

- Angelini E, Clair D, Borgo M, Bertaccini A, Boudon-Padieu E (2001) "Flavescence dorée" in France and Italy oOccurrence of closely related phytoplasma isolates and their near relationship to pPalatine grapevine yellows and an alder phytoplasma. Vitis 40:79–-86
- Bertaccini A (2014) http://www.ipwgnet.org/collection. Accessed 27 Oct 2019-
- Bertaccini A, Marani F (1982) Electron microscopy of two viruses and mycoplasma-like organisms in lilies with deformed flowers. Phytopathol Mediterr 21:8–-14
- Calari A, Paltrinieri S, Contaldo N, Sakalieva D, Mori N, Duduk B, Bertaccini A (2011) Molecular evidence of phytoplasmas in winter oilseed rape, tomato and corn seedlings. Bull Insectol 64(S):S157- S158
- Carminati G, Satta E, Paltrinieri S, Bertaccini A (2019) Simultaneous evaluation of "*Candidatus* Phytoplasma" and "*Candidatus* Liberibacter solanacearum" seed transmission in carrot. Phytopath Moll 9(1):141–-142
- Cebrián MC, Villaescusa FJ, Alfaro-Fernández A, Hermoso de Mendoza A, Córdoba-Sellés MC, Jordá C, Ferrándiz JC, Sanjuán S, Font MI (2010) First report of *Spiroplasma citri* in carrot in Europe. Plant Dis 94(10):1264
- De La Rue SJ, Hopkinson R, Gibb KS (2002) Assessment of *Stylosanthes* seed yield reduction caused by phytoplasma-associated diseases. Aust J Exp Agr 42(8):1053–-1056
- Duduk B, Perić P, Marčić D, Drobnjaković T, Picciau L, Alma A, Bertaccini A (2008) Phytoplasmas in carrots: disease and potential vectors in Serbia. Bull Insectol 61:327–-331
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. Evol 39:783–-791
- Font I, Abad P, Albinana M, Espino AI, Dally EL, Davis RE, Jorda C (1999) Amarilleos y enrojecimientos en zanahoria: una enfermedad a diagnostico. Bol San Veg Plagas 25:405–415
- Gundersen DE, Lee I-M (1996) Ultrasensitive detection of phytoplasmas by nested-PCR assays using two universal primer pairs. Phytopathol Mediterr 35:114–-151
- Jiang H, Wei W, Saiki T, Kawakita H, Watanabe K, Sato M (2004) Distribution patterns of mulberry dwarf phytoplasma in reproductive organs, winter buds, and roots of mulberry trees. J Gen Plant Pathol 70:168–173
- Khan AJ, Botti S, Paltrinieri S, Al-Subhi AM, Bertaccini A (2002) Phytoplasmas in alfalfa seedlings: infected or contaminated seeds? Abstracts 14th IOM. Vienna, July 07-12:6
- Lee I-M, Gundersen DE, Hammond RW, Davis RE (1994) Use of mycoplasma-like organism (MLO) groupspecific oligonucleotide primers for nested-PCR assays to detect mixed-MLO infections in a single host plant. Phytopathology 84:559–-566
- Makarova OV, Contaldo N, Paltrinieri S, Kawube G, Bertaccini A, Nicolaisen M (2012) DNA barcoding for universal identification of "*Candidatus* Phytoplasmas" using a fragment of the elongation factor Tu gene. PLoS One 7(12):e52092
- McCoy RE, Caudwell A, Chang CJ, Chen TA, Chiykowskyi LN, Cousin MT, Dale de Leeuw GTN, Golino DA, Hackett KJ, Kirkptrick BC, Marwitz R, Petzold H, Shina RH, Sugiura M, Whitcomb RF, Yang IL, Zhu BM, Seemüller E (1989) Plant diseases associated with mycoplasma-like organisms. In: The mMycoplasmas, vol. 5. (eds. Whitcomb R.F., Tully J.G.). Academic, New York. Pp. 545–-640
- Nečas T, Mašková V, Krška B (2008) The possibility of ESFY phytoplasma transmission through flowers and seeds. Acta Hortic 781:443–-448
- Nipah J, Jones P, Hodgetts J, Dickinson M (2007) Detection of phytoplasma DNA in embryos from coconut palms in Ghana, and kernels from maize in Peru. Bull Insectol 60:385–-386
- Olivier CY, Galka B, Sèguin-Swartz G (2010) Detection of "aster yellows" phytoplasma DNA in seed and seedlings of canola (*Brassica napus* and B. *rapa*) and AY strain identification. Can J Plant Pathol 32:298–-305
- Oropeza C, Cordova I, Puch-Hau C, Castillo R, Chan JL, Sáenz L (2017) Detection of lethal yellowing phytoplasma in coconut plantlets obtained through in vitro germination of zygotic embryos from the seeds of infected palms. Ann Appl Biol 171:28–-36
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol 4:406–-425
- Satta E, Ramírez AS, Paltrinieri, S, Contaldo N, Benito P, Poveda JB, Bertaccini A (2016) Simultaneous detection of mixed "*Candidatus* Phytoplasma asteris" and "*Ca*. Liberibacter solanacearum" infection in carrot. Phytopathol Mediterr 55(3),401–-409
- Satta E, Nanni IM, Contaldo N, Collina M, Poveda JB, Ramirez AS, Bertaccini A (2017) General phytoplasma detection by a q-PCR method using mycoplasma primers. Mol Cell Probes 35:1–-7

Satta E, Paltrinieri S, Bertaccini A (2019) Phytoplasma transmission by seed. In: Phytoplasmas: pPlant pPathogenic Bacteria-II tTransmission and Management of Phytoplasma Associated Diseases. (Eds Bertaccini AA, PG Weintraub, GP Rao, N Mori) CChapter 6. Springer, Singapore. Pp 131–-147

Sears BB, Klomparens KL (1989) Leaf tip cultures of the evening primrose allow stable, aseptic culture of mycoplasma-like organism. Can J Plant Pathol 11:343–-348

- Siller W, Kuhbandner B, Marwitz R, Petzold H, Seemüller E (1987) Occurrence of mycoplasma-like organisms in parenchyma cells of *Cuscuta odorata* (Ruiz et Pav.). J Phytopathol 119:147–-159
- Swallow WH (1985) Group testing for estimating infection rates and probabilities of disease transmission. Phytopathology 7:5882--5889-
- Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. Proc Natl Acad Sci U S AProc Natl Acad Sci USA 101:11030–-11035
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. Mol Biol Evol 30: 2725–-2729
- Zambryski P (2004) Cell-to-cell transport of proteins and fluorescent tracers via plasmodesmata during plant development. J Cell Biol 164:165–168

Zwolinska A, Krawczyk K, Pospieszny H (2010) First report of "stolbur" phytoplasma infecting pea plants. Abstracts 18th IOM. ChiancianoTerme, 11-16 July 145:152

Fig.ure -1 Seedlings placed in water shortage conditions (stressed samples) and showing symptoms of leaf whitening (left) and browning (right) compared to an asymptomatic leaf from a not stressed carrot seedling (in the middle).

Fig.ure -2 The phylogenetic tree was inferred using the nNeighbor-jJoining method (Saitou and Nei [1987\)](#page-4-0) enclosing ribosomal gene 16Sr DNA. In bold sequences of strains obtained from carrot seedlings 11, 13, and cotyledon 12. The sequence of *Acholeplasma laidlawii* strain PG8A was used as out group. On the right GenBank accession numbers, $\frac{1}{r}$ The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test $(1,000$ replicates) is sure shown next to the branches (Felsenstein [1985\)](#page-4-0). The evolutionary distances were computed using the mMaximum cComposite lLikelihood method (Tamura et al[. 2004\)](#page-5-0) and are in the units of the number of base substitutions per site.

Table -1 Results of phytoplasma detection in carrot seedlings.

*Different color indicate different lots.

Table -2 Results of phytoplasma detection in the 3three--month-olds -stressed carrot seedlings.

*Different color indicate different lots.